

ECOLOGY

Adding Biofuels to the Invasive Species Fire?

S. Raghu,^{1*} R. C. Anderson,² C. C. Daehler,³ A. S. Davis,⁴ R. N. Wiedenmann,⁵ D. Simberloff,⁶ R. N. Mack⁷

The U.S. renewable energy initiative (1) announced in the 2006 presidential State of the Union address (2) has given new impetus to the identification of biofuel crops as sources of energy. However, an earlier presidential directive, Executive Order 13112 (3), attempts to protect the United States from invasive species, unless benefits clearly outweigh potential harms. The policies may conflict because traits deemed ideal in a bioenergy crop are also commonly found among invasive species (see figure).

Biofuel crops may have economic benefits, but studies of concomitant environmental risks of movement into novel habitats are seldom conducted. Although anecdotal claims of “low risk” for some species (4) may be valid, many purportedly beneficial introduced species have had long-term economic and environmental costs owing to their invasiveness (5, 6). For example, *Sorghum halepense* is an introduced forage grass that became an invasive weed in 16 of the 48 U.S. states in which it occurs. Even the most conservative estimate of competitive losses for cotton and soybean crops in three states is in excess of \$30 million annually (7).

Several grasses and woody species have been evaluated for biofuel production, with perennial rhizomatous grasses showing the most economic promise (4, 8). *Arundo donax* (giant reed; native to Asia) and *Phalaris arundinacea* (reed canary grass; native to temperate Europe, Asia, and North America) are two C₃ grasses being considered as biofuel species (8) that are invasive in some U.S. ecosystems. The former threatens riparian areas and alters fire cycles (9); the latter invade wet-

lands (10) and affect wildlife habitat.

The hybrid grass *Miscanthus × giganteus* (native to Asia) and *Panicum virgatum* (switchgrass; native to central and eastern United States) are C₄ grasses being considered in Europe and the United States (4, 11). Several *Miscanthus* species are invasive or have invasive potential (12); in particular, the parent species of *M. × giganteus* (13, 14). *Miscanthus × giganteus* is an allopolyploid that does not produce viable seed and reproduces vegetatively. However, allopolyploidy does not guarantee continued sterility (15) and vegetative propagation is often associated with invasiveness (16, 17) or directly contributes to it (18). Several other traits that make *Miscanthus* potentially valuable as a crop could enhance invasiveness (ability to resprout from below ground, efficient photosynthetic mechanisms, and rapid growth rates) (16, 19).

The U.S. native, *P. virgatum*, shares many traits with *Miscanthus* and can also produce seeds, which may give *P. virgatum* even greater invasive potential. Furthermore, plants native in one region can become invasive when established elsewhere (20). Escape from competitors and natural enemies may help explain the weedy nature of *P. virgatum* outside its endemic range (21). Internationally, there has been little success in eradicating or even controlling an invading grass. Herbicides are used to control invasive grasses on croplands, but they are too expensive to use on rangelands, national parks, and reserves. Development of the most economical tool, biological control with a specific natural enemy, has been avoided because of the perceived risk of its expanding its host range to include commercial grasses, such as wheat, corn, barley, or rice (22).

Balancing costs and benefits of species introductions is a key contemporary challenge. Introducing some plant species as bio-

Biofuel crops, particularly using non-native species, must be introduced with an understanding of possible risks to the environment.

fuel sources may be safe, but safety must be established by agronomic and ecological analyses. Such analyses are already mandatory for biological control agents (23) and transgenic plants (24). Experts must assess ecological risks before introducing biofuel crops, to ensure that we do not add biofuels to the already raging invasive species fire.


References and Notes

- National Energy Policy Development Group, *Reliable, Affordable, and Environmentally Sound Energy for America's Future*; www.whitehouse.gov/energy.
- 31 January 2006, www.gpoaccess.gov/sou/06sou.html.
- Exec. Order 13112: *Invasive Species* (3 February 1999); http://ceq.eh.doe.gov/nepa/regs/eos/eo13112.html.
- E. A. Heaton et al., *Mitigat. Adapt. Strateg. Global Change* **9**, 433 (2004).
- J. J. Ewel et al., *Bioscience* **49**, 619 (1999).
- National Research Council, *Predicting Invasions of Nonindigenous Plants and Plant Pests* (National Academies Press, Washington, DC, 2002), p. 194.
- C. G. McWhorter, *Weed Sci* **41**, 669 (1993).
- I. Lewandowski et al., *Biomass Bioenergy* **25**, 335 (2003).
- T. L. Dudley, in *Invasive Plants of California's Wilds*, C. C. Bossard, J. M. Randall, M. C. Hoshovsky, Eds. (Univ. of California Press, Berkeley, 2000), pp. 53–58.
- J. Molofsky et al., *Biol. Invasions* **1**, 181 (1999).
- I. Lewandowski et al., *Biomass Bioenergy* **19**, 209 (2000).
- Plants database profile; http://plants.usda.gov/java/profile?symbol=MISCA.
- C. M. D'Antonio, P. M. Vitousek, *Annu. Rev. Ecol. Syst.* **23**, 63 (1992).
- Minnesota Department of Natural Resources; www.dnr.state.mn.us/invasives/terrestrialplants/grasses/amursilvergrass.html.
- A. J. Gray et al., *Adv. Ecol. Res.* **21**, 1 (1991).
- C. Daehler, *Biol. Conserv.* **84**, 167 (1998).
- C. S. Kolar, D. M. Lodge, *Trends Ecol. Evol.* **16**, 199 (2001).
- M. L. Moody, D. H. Les, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 14867 (2002).
- P. C. Pheulong et al., *J. Environ. Manage.* **57**, 239 (1999).
- J. M. Randall, J. Marinelli, Eds., *Invasive Plants: Weeds of the Global Garden* (Brooklyn Botanic Garden, Brooklyn, NY, 1996), 111 pp.
- J. Forman, in *Plant Invasions: Ecological Threats and Management Solutions*, L. E. Child et al., Eds. (Backhuys Publishers, Leiden, Netherlands, 2003), pp. 17–39.
- R. D. Goeden, L. A. Andres, in *Handbook of Biological Control: Principles and Applications of Biological Control*, T. S. Bellows, T. W. Fisher, Eds. (Academic Press, London, 1999), pp. 871–890.
- A. W. Sheppard et al., *Biocontrol News Inform.* **24**, 91N (2003).
- A. Hilbeck, D. A. Andow, Eds., *Environmental Risk Assessment of Genetically Modified Organisms* (CABI, Wallingford, UK, 2004), 300 pp.
- A bibliography is provided as supporting online material.

Supporting Online Material

www.sciencemag.org/cgi/content/full/313/5794/1742/DC1

10.1126/science.1129313



Ideal ecological traits of biomass energy crops (4). All traits shown other than perennial growth and sterile seeds are known to contribute to invasiveness. See (25).

¹Illinois Natural History Survey, Champaign, IL 61820;

²Illinois State University, Normal, IL 61709; ³University of Hawaii at Manoa, Honolulu, HI 96822; ⁴USDA-ARS Invasive Weed Management Unit, Urbana, IL 61801; ⁵University of Arkansas, Fayetteville, AR 72701; ⁶University of Tennessee, Knoxville, TN 37996; ⁷Washington State University, Pullman, WA 99164, USA.

*Author for correspondence. E-mail: raghu@uiuc.edu