


Texas – 2010 Profile

HIV/AIDS Epidemic


HIV/AIDS has claimed the lives of more than 550,000 Americans. Today, about 1.1 million Americans are living with HIV, the virus that causes AIDS, and one fifth of those infected are unaware of their infection.

Estimated AIDS Diagnoses among Adults and Adolescents, by Transmission Category, Cumulative through 2008, Texas
N = 76,674*



*MSM, men who have sex with men; IDU, injection drug users; MSM/IDU, men who have sex with men who also inject drugs

Estimated AIDS Diagnoses, by Race/Ethnicity, Cumulative through 2008, Texas
N = 77,070*



*Native Hawaiian/Other Pacific Islander, Unknown Race: <0.1%

Texas reported 77,070 AIDS cases to CDC, cumulatively, from the beginning of the epidemic through December 2008. Texas ranked 4th highest among the 50 states in cumulative reported AIDS cases.


Tuberculosis (TB)

Although the overall rate of TB in the United States has declined substantially since 1992, the rate of decrease among foreign-born persons has been much smaller than that for U.S.-born persons.

In 2008, Texas reported:

- The 4th highest rate of TB among states in the United States (6.2 per 100,000 persons).
- 48.4% of TB cases occurred in foreign-born persons.

TB Cases by Race/Ethnicity, 2008, Texas
N = 1,501*



*Unknown or Missing: <0.1%

Hepatitis A, B, and C Virus (HAV, HBV, HCV)

In the United States, incidence of acute HAV and HBV in 2008 was the lowest ever recorded because of the availability of safe and effective vaccines. But there is no vaccine for HCV, and chronic HBV and HCV account for more than 50% of new cases of chronic liver disease, a leading cause of death. About 4.5 million people are estimated to be living with HBV and HCV infection, and of that number, about 50% are unaware of their status.



From 1999 through 2008, Texas:

- Reported rates of acute hepatitis A decreased by 91%.
- Reported rates of acute hepatitis B decreased by 47%.

Sexually Transmitted Diseases (STDs)

Syphilis – Primary and secondary (P&S) syphilis (the stages when syphilis is most infectious) remains a problem in the southern United States and some urban areas.

- Texas ranked 10th among 50 states, with 5.9 cases of P&S syphilis per 100,000 persons.
- The number of congenital syphilis cases increased from 68 in 1999 to 127 in 2008.



Chlamydia and Gonorrhea – Chlamydial and gonorrheal infections in women are usually asymptomatic and often go undiagnosed. Untreated, these infections can lead to pelvic inflammatory disease, which can cause tubal infertility, ectopic pregnancy, and chronic pelvic pain. In 2008, Texas:

- Ranked 17th among 50 states in chlamydial infections (422 per 100,000 persons) and ranked 15th among 50 states in gonorrheal infections (134.7 per 100,000 persons).
- Reported rates of chlamydia among women (659.5 cases per 100,000) were 3.6 times greater than those among men (182.9 cases per 100,000).

Program Initiatives Supported by CDC

HIV/AIDS – CDC uses a comprehensive approach to HIV prevention that includes surveillance, research, interventions, capacity building, and evaluation. In Texas, CDC supports the state and one city health department, 10 community-based organizations, and two capacity building assistance providers to conduct and support HIV prevention programs. Programs are designed to meet the cultural needs, expectations, and values of the populations they serve and involve affected communities in the HIV prevention community planning process to ensure that funding goes to those who need it most. Research, surveillance, and other prevention efforts are also supported.

STDs – In Texas, CDC funds the state and Houston health departments through the comprehensive STD Prevention System (CSPS) grant program. CSPS supports a community-wide, science-based, interdisciplinary approach to STD prevention that includes behavioral interventions, medical and laboratory services, disease surveillance, outbreak response, professional development, and STD awareness and education campaigns. As part of its CSPS grant, the Texas State Department of Health receives funding specifically for syphilis elimination. CDC also supports an STD-HIV Prevention Training Center in Texas and special surveillance.

TB – In Texas, CDC funds the state and Houston health departments for TB prevention and control activities, including surveillance, case management, and directly observed therapy. These funds also support the identification and evaluation of persons exposed to TB, as well as laboratory services. CDC also supports a regional training and medical consultation center for complex TB cases. TB clinical and epidemiological research is also supported.

Viral Hepatitis – In Texas, CDC funds adult hepatitis prevention coordinators in the state and Houston health departments to integrate viral hepatitis prevention activities into existing public health program. CDC also supports integrated surveillance and other extramural programs.

CDC Funding to Texas, 2009	
HIV/AIDS	\$32,831,239
STDs	\$7,104,971*
TB	\$12,205,908*
Viral Hepatitis	\$963,309

*Also includes HIV appropriated funds

For More Information

Texas: <http://www.dshs.state.tx.us/> CDC: <http://www.cdc.gov/nchhstp/>