

Table 42. Tuberculosis Cases and Percentages by Reason Therapy Stopped: Reporting Areas, 2008¹

Reporting Area	Cases with Initial Drug Regimen Prescribed ²		Completed Therapy		Adverse Event		Moved		Lost		Refused		Died ³		Unknown ⁴	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
United States	12,542	10,813 (86.2)	5	(0.0)	284	(2.3)	315	(2.5)	78	(0.6)	837	(6.7)	210	(1.7)		
Alabama	168	(85.1)	0	(0.0)	2	(1.2)	6	(3.6)	1	(0.6)	16	(9.5)	0	(0.0)		
Alaska	48	(91.7)	0	(0.0)	0	(0.0)	0	(0.0)	1	(2.1)	3	(6.3)	0	(0.0)		
Arizona	212	(73.6)	0	(0.0)	2	(0.9)	13	(6.1)	1	(0.5)	15	(7.1)	25	(11.8)		
Arkansas	80	(85.0)	0	(0.0)	3	(3.8)	4	(5.0)	0	(0.0)	3	(3.8)	2	(2.5)		
California	2,635	(85.4)	0	(0.0)	112	(4.3)	35	(1.3)	13	(0.5)	176	(6.7)	49	(1.9)		
Colorado	101	(93.1)	0	(0.0)	1	(1.0)	3	(3.0)	0	(0.0)	3	(3.0)	0	(0.0)		
Connecticut	97	(92.8)	0	(0.0)	1	(1.0)	1	(1.0)	0	(0.0)	5	(5.2)	0	(0.0)		
Delaware	23	(87.0)	0	(0.0)	2	(8.7)	1	(4.3)	0	(0.0)	0	(0.0)	0	(0.0)		
District of Columbia	53	(79.2)	0	(0.0)	7	(13.2)	1	(1.9)	0	(0.0)	3	(5.7)	0	(0.0)		
Florida	912	(87.4)	0	(0.0)	18	(2.0)	20	(2.2)	1	(0.1)	70	(7.7)	6	(0.7)		
Georgia	458	(84.7)	0	(0.0)	10	(2.2)	10	(2.2)	3	(0.7)	30	(6.6)	17	(3.7)		
Hawaii	123	(87.0)	0	(0.0)	6	(4.9)	0	(0.0)	0	(0.0)	8	(6.5)	2	(1.6)		
Idaho	11	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)		
Illinois	459	(87.6)	0	(0.0)	0	(0.0)	17	(3.7)	9	(2.0)	29	(6.3)	2	(0.4)		
Indiana	115	(83.5)	0	(0.0)	1	(0.9)	3	(2.6)	0	(0.0)	12	(10.4)	3	(2.6)		
Iowa	48	(91.7)	0	(0.0)	2	(4.2)	0	(0.0)	1	(2.1)	1	(2.1)	0	(0.0)		
Kansas	57	(94.7)	0	(0.0)	0	(0.0)	1	(1.8)	0	(0.0)	2	(3.5)	0	(0.0)		
Kentucky	96	(81.3)	0	(0.0)	1	(1.0)	6	(6.3)	2	(2.1)	7	(7.3)	2	(2.1)		
Louisiana	219	(89.5)	0	(0.0)	7	(3.2)	2	(0.9)	2	(0.9)	9	(4.1)	3	(1.4)		
Maine	9	(77.8)	0	(0.0)	1	(11.1)	0	(0.0)	0	(0.0)	0	(0.0)	1	(11.1)		
Maryland	272	(88.2)	0	(0.0)	7	(2.6)	5	(1.8)	0	(0.0)	18	(6.6)	2	(0.7)		
Massachusetts	255	(87.5)	0	(0.0)	12	(4.7)	3	(1.2)	0	(0.0)	15	(5.9)	2	(0.8)		
Michigan	174	(83.3)	1	(0.0)	0	(0.0)	2	(1.1)	3	(1.7)	16	(9.2)	7	(4.0)		
Minnesota	209	(92.8)	0	(0.0)	4	(1.9)	1	(0.5)	1	(0.5)	6	(2.9)	3	(1.4)		
Mississippi	111	(87.4)	0	(0.0)	1	(0.9)	0	(0.0)	0	(0.0)	11	(9.9)	2	(1.8)		
Missouri	106	(90.6)	0	(0.0)	6	(5.7)	0	(0.0)	0	(0.0)	4	(3.8)	0	(0.0)		
Montana	8	(62.5)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	3	(37.5)	0	(0.0)		
Nebraska	33	(81.8)	0	(0.0)	2	(6.1)	1	(3.0)	0	(0.0)	3	(9.1)	0	(0.0)		
Nevada	102	(82.4)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	10	(9.8)	8	(7.8)		
New Hampshire	19	(73.7)	0	(0.0)	0	(0.0)	1	(5.3)	1	(5.3)	2	(10.5)	1	(5.3)		
New Jersey	416	(88.0)	0	(0.0)	1	(0.2)	22	(5.3)	1	(0.2)	26	(6.3)	0	(0.0)		
New Mexico	56	(89.3)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	6	(10.7)	0	(0.0)		

Table 42. Tuberculosis Cases and Percentages by Reason Therapy Stopped: Reporting Areas, 2008¹

Reporting Area	Cases with Initial Drug Regimen Prescribed ²		Completed Therapy		Adverse Event		Moved		Lost		Refused		Died ³		Unknown ⁴	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
New York State ⁵	296	(90.5)	268	(90.5)	0	(0.0)	3	(1.0)	9	(3.0)	2	(0.7)	14	(4.7)	0	(0.0)
New York City	882	(88.8)	783	(88.8)	0	(0.0)	12	(1.4)	16	(1.8)	5	(0.6)	54	(6.1)	12	(1.4)
North Carolina	328	(91.8)	301	(91.8)	0	(0.0)	0	(0.0)	7	(2.1)	0	(0.0)	17	(5.2)	3	(0.9)
North Dakota	3	(33.3)	1	(33.3)	0	(0.0)	0	(0.0)	0	(0.0)	1	(33.3)	0	(0.0)	1	(33.3)
Ohio	209	(85.2)	178	(85.2)	0	(0.0)	4	(1.9)	2	(1.0)	6	(2.9)	15	(7.2)	4	(1.9)
Oklahoma	95	(78.9)	75	(78.9)	0	(0.0)	4	(4.2)	0	(0.0)	0	(0.0)	8	(8.4)	8	(8.4)
Oregon	74	(94.6)	70	(94.6)	0	(0.0)	0	(0.0)	0	(0.0)	1	(1.4)	3	(4.1)	0	(0.0)
Pennsylvania	373	(84.2)	314	(84.2)	3	(0.0)	0	(0.0)	10	(2.7)	5	(1.3)	20	(5.4)	21	(5.6)
Rhode Island	36	(94.4)	34	(94.4)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	2	(5.6)	0	(0.0)
South Carolina	179	(84.9)	152	(84.9)	0	(0.0)	3	(1.7)	4	(2.2)	3	(1.7)	17	(9.5)	0	(0.0)
South Dakota	16	(100.0)	16	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Tennessee	274	(86.1)	236	(86.1)	0	(0.0)	9	(3.3)	1	(0.4)	0	(0.0)	23	(8.4)	5	(1.8)
Texas	1,452	(82.9)	1,204	(82.9)	0	(0.0)	26	(1.8)	101	(7.0)	6	(0.4)	106	(7.3)	9	(0.6)
Utah	26	(92.3)	24	(92.3)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(3.8)	1	(3.8)
Vermont	5	(60.0)	3	(60.0)	0	(0.0)	2	(40.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Virginia	288	(86.5)	249	(86.5)	0	(0.0)	9	(3.1)	5	(1.7)	8	(2.8)	16	(5.6)	1	(0.3)
Washington	222	(89.2)	198	(89.2)	0	(0.0)	2	(0.9)	2	(0.9)	1	(0.5)	18	(8.1)	1	(0.5)
West Virginia	26	(84.6)	22	(84.6)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	4	(15.4)	0	(0.0)
Wisconsin	68	(77.9)	53	(77.9)	1	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	7	(10.3)	7	(10.3)
Wyoming	5	(80.0)	4	(80.0)	0	(0.0)	1	(20.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
American Samoa ⁶	3	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	3	(100.0)
Fed. States of Micronesia ⁶	176	(50.6)	89	(50.6)	0	(0.0)	4	(2.3)	3	(1.7)	0	(0.0)	11	(6.3)	69	(39.2)
Guam ⁶	89	(94.4)	84	(94.4)	0	(0.0)	1	(1.1)	0	(0.0)	0	(0.0)	3	(3.4)	1	(1.1)
Marshall Islands ⁶	123	(86.2)	106	(86.2)	0	(0.0)	0	(0.0)	1	(0.8)	7	(5.7)	3	(2.4)	6	(4.9)
N. Mariana Islands ⁶	35	(80.0)	28	(80.0)	0	(0.0)	5	(14.3)	0	(0.0)	0	(0.0)	1	(2.9)	1	(2.9)
Puerto Rico ⁶	85	(80.0)	68	(80.0)	0	(0.0)	0	(0.0)	2	(2.4)	0	(0.0)	15	(17.6)	0	(0.0)
Republic of Palau ⁶	16	(18.8)	3	(18.8)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(6.3)	12	(75.0)
U.S. Virgin Islands ⁶	4	(50.0)	2	(50.0)	0	(0.0)	0	(0.0)	1	(25.0)	1	(25.0)	0	(0.0)	0	(0.0)

¹Most recent year for which data are available.

²Number of cases in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia).

³Died = Died of any cause.

⁴Includes cases reported as Other, Missing, or Unknown.

⁵Excludes New York City.

⁶Not included in U.S. totals.

Note: Ellipses indicate data not available.