

Your Urinary System and How It Works

National Kidney and Urologic Diseases Information Clearinghouse



National
Institute of
Diabetes and
Digestive
and Kidney
Diseases

NATIONAL
INSTITUTES
OF HEALTH


The organs, tubes, muscles, and nerves that work together to create, store, and carry urine are the urinary system. The urinary system includes two kidneys, two ureters, the bladder, two sphincter muscles, and the urethra.

How does the urinary system work?

Your body takes nutrients from food and uses them to maintain all bodily functions including energy and self-repair. After your body has taken what it needs from the food, waste products are left behind in the blood and in the bowel. The urinary system works with the lungs, skin, and intestines—all of which also excrete wastes—to keep the chemicals and water in your body balanced. Adults eliminate about a quart and a half of urine each day. The amount depends on many factors, especially the amounts of fluid and food a person consumes and how much fluid is lost through sweat and breathing. Certain types of medications can also affect the amount of urine eliminated.

The urinary system removes a type of waste called urea from your blood. Urea is produced when foods containing protein, such as meat, poultry, and certain vegetables, are broken down in the body. Urea is carried in the bloodstream to the kidneys.

The kidneys are bean-shaped organs about the size of your fists. They are near the middle of the back, just below the rib cage. The kidneys remove urea from the blood



Front view of urinary tract.

through tiny filtering units called nephrons. Each nephron consists of a ball formed of small blood capillaries, called a glomerulus, and a small tube called a renal tubule. Urea, together with water and other waste substances, forms the urine as it passes through the nephrons and down the renal tubules of the kidney.

From the kidneys, urine travels down two thin tubes called ureters to the bladder. The ureters are about 8 to 10 inches long. Muscles in the ureter walls constantly tighten and relax to force urine downward away from the kidneys. If urine is allowed to stand still, or back up, a kidney infection can develop. Small amounts of urine are emptied into the bladder from the ureters about every 10 to 15 seconds.



U.S. Department
of Health and
Human Services

The bladder is a hollow muscular organ shaped like a balloon. It sits in your pelvis and is held in place by ligaments attached to other organs and the pelvic bones. The bladder stores urine until you are ready to go to the bathroom to empty it. It swells into a round shape when it is full and gets smaller when empty. If the urinary system is healthy, the bladder can hold up to 16 ounces (2 cups) of urine comfortably for 2 to 5 hours.

Circular muscles called sphincters help keep urine from leaking. The sphincter muscles close tightly like a rubber band around the opening of the bladder into the urethra, the tube that allows urine to pass outside the body.

Nerves in the bladder tell you when it is time to urinate, or empty your bladder. As the bladder first fills with urine, you may notice a feeling that you need to urinate. The sensation to urinate becomes stronger as the bladder continues to fill and reaches its limit. At that point, nerves from the bladder send a message to the brain that the bladder is full, and your urge to empty your bladder intensifies.

When you urinate, the brain signals the bladder muscles to tighten, squeezing urine out of the bladder. At the same time, the brain signals the sphincter muscles to relax. As these muscles relax, urine exits the bladder through the urethra. When all the signals occur in the correct order, normal urination occurs.

What causes problems in the urinary system?

Problems in the urinary system can be caused by aging, illness, or injury. As you get older, changes in the kidneys' structure cause them to lose some of their ability to remove wastes from the blood. Also, the

muscles in your ureters, bladder, and urethra tend to lose some of their strength. You may have more urinary infections because the bladder muscles do not tighten enough to empty your bladder completely. A decrease in strength of muscles of the sphincters and the pelvis can also cause incontinence, the unwanted leakage of urine. Illness or injury can also prevent the kidneys from filtering the blood completely or block the passage of urine.

How are problems in the urinary system detected?

Urinalysis is a test that studies the content of urine for abnormal substances such as protein or signs of infection. This test involves urinating into a special container and leaving the sample to be studied.

Urodynamic tests evaluate the storage of urine in the bladder and the flow of urine from the bladder through the urethra. Your doctor may want to do a urodynamic test if you are having symptoms that suggest problems with the muscles or nerves of your lower urinary system and pelvis—ureters, bladder, urethra, and sphincter muscles.

Urodynamic tests measure the contraction of the bladder muscle as it fills and empties. The test is done by inserting a small tube called a catheter through your urethra into your bladder to fill it either with water or a gas. Another small tube is inserted into your rectum or vagina to measure the pressure put on your bladder when you strain or cough. Other bladder tests use x-ray dye instead of water so that x-ray pictures can be taken when the bladder fills and empties to detect any abnormalities in the shape and function of the bladder. These tests take about an hour.

What are some disorders of the urinary system?

Disorders of the urinary system range in severity from easy to treat to life threatening.

Benign prostatic hyperplasia (BPH) is a condition in men that affects the prostate gland, which is part of the male reproductive system. The prostate is located at the bottom of the bladder and surrounds the urethra. BPH is an enlargement of the prostate gland that can interfere with urinary function in older men. It causes blockage by squeezing the urethra, which can make it difficult to urinate. Men with BPH frequently have other bladder symptoms including an increase in frequency of bladder emptying both during the day and at night. Most men over age 60 have some BPH, but not all have problems with blockage. There are many different treatment options for BPH.

Painful bladder syndrome/Interstitial cystitis (PBS/IC) is a chronic bladder disorder also known as frequency-urgency-dysuria syndrome. In this disorder, the bladder wall can become inflamed and irritated. The inflammation can lead to scarring and stiffening of the bladder, decreased bladder capacity, pinpoint bleeding, and, in rare cases, ulcers in the bladder lining. The cause of IC is unknown at this time.

Kidney stones is the term commonly used to refer to stones, or calculi, in the urinary system. Stones form in the kidneys and may be found anywhere in the urinary system. They vary in size. Some stones cause great pain while others cause very little. The aim of treatment is to remove the stones, prevent infection, and prevent recurrence. Both nonsurgical and surgical treatments are used. Kidney stones affect men more often than women.

Prostatitis is inflammation of the prostate gland that results in urinary frequency and urgency, burning or painful urination, a condition called dysuria, and pain in the lower back and genital area, among other symptoms. In some cases, prostatitis is caused by bacterial infection and can be treated with antibiotics. But the more common forms of prostatitis are not associated with any known infecting organism. Antibiotics are often ineffective in treating the nonbacterial forms of prostatitis.

Proteinuria is the presence of abnormal amounts of protein in the urine. Healthy kidneys take wastes out of the blood but leave in protein. Protein in the urine does not cause a problem by itself. But it may be a sign that your kidneys are not working properly.

Renal (kidney) failure results when the kidneys are not able to regulate water and chemicals in the body or remove waste products from your blood. Acute renal failure (ARF) is the sudden onset of kidney failure. This condition can be caused by an accident that injures the kidneys, loss of a lot of blood, or some drugs or poisons. ARF may lead to permanent loss of kidney function. But if the kidneys are not seriously damaged, they may recover. Chronic kidney disease (CKD) is the gradual reduction of kidney function that may lead to permanent kidney failure, or end-stage renal disease (ESRD). You may go several years without knowing you have CKD.

Urinary tract infections (UTIs) are caused by bacteria in the urinary tract. Women get UTIs more often than men. UTIs are treated with antibiotics. Drinking lots of fluids also helps by flushing out the bacteria.

The name of the UTI depends on its location in the urinary tract. An infection in the bladder is called cystitis. If the infection is in one or both of the kidneys, the infection

is called pyelonephritis. This type of UTI can cause serious damage to the kidneys if it is not adequately treated.

Urinary incontinence, loss of bladder control, is the involuntary passage of urine. There are many causes and types of incontinence, and many treatment options. Treatments range from simple exercises to surgery. Women are affected by urinary incontinence more often than men.

Urinary retention, or bladder-emptying problems, is a common urological problem with many possible causes. Normally, urination can be initiated voluntarily and the bladder empties completely. Urinary retention is the abnormal holding of urine in the bladder. Acute urinary retention is the sudden inability to urinate, causing pain and discomfort. Causes can include an obstruction in the urinary system, stress, or neurologic problems. Chronic urinary retention refers to the persistent presence of urine left in the bladder after incomplete emptying. Common causes of chronic urinary retention are bladder muscle failure, nerve damage, or obstructions in the urinary tract. Treatment for urinary retention depends on the cause.

Who can help me with a urinary problem?

Your primary doctor can help you with some urinary problems. Your pediatrician may be able to treat some of your child's urinary problems. But some problems may require the attention of a urologist, a doctor who specializes in treating problems of the urinary system and the male reproductive system. A gynecologist is a doctor who specializes in the female reproductive system and may be able to help with some urinary problems. A urogynecologist is a gynecologist who specializes in the female urinary system. A nephrologist specializes in treating diseases of the kidney.

Points to Remember

- Your urinary system filters waste and extra fluid from your blood.
- Problems in the urinary system include kidney failure, urinary tract infections, kidney stones, prostate enlargement, and bladder control problems.
- Health professionals who treat urinary problems include general practitioners (your primary doctor), pediatricians, urologists, gynecologists, urogynecologists, and nephrologists.

For More Information

American Urological Association Foundation

1000 Corporate Boulevard
Linthicum, MD 21090
Phone: 1-800-828-7866 or
410-689-3700
Fax: 410-689-3998
Email: auafoundation@auafoundation.org
Internet: www.UrologyHealth.org

American Kidney Fund

11921 Rockville Pike
Suite 300
Rockville, MD 20852
Phone: 1-800-638-8299
Internet: www.kidneyfund.org

American Society of Pediatric Nephrology

Northwestern University
Feinberg School of Medicine
Pediatrics W140
303 East Chicago Avenue
Chicago, IL 60611-3008
Phone: 312-503-4000
Fax: 312-503-1181
E-mail: aspn@northwestern.edu
Internet: www.aspneph.com/research.html

American Urogynecologic Society

2025 M Street, NW
Suite 800
Washington, DC 20036
Phone: 202-367-1167
Fax: 202-367-2167
Email: info@aug.org
Internet: www.augs.org

Interstitial Cystitis Association of America, Inc.

110 North Washington Street
Suite 340
Rockville, MD 20850-2223
Phone: 1-800-HELP-ICA (435-7422) or
301-610-5300
Fax: 301-610-5308
Email: icamail@ichelp.org
Internet: www.ichelp.org

National Association for Continence

P.O. Box 1019
Charleston, SC 29402-1019
Phone: 1-800-BLADDER (252-3337) or
843-377-0900
Email: memberservices@nafc.org
Internet: www.nafc.org

National Kidney Foundation

30 East 33rd Street
New York, NY 10016
Phone: 1-800-622-9010 or 212-889-2210
Fax: 212-689-9261
Internet: www.kidney.org

The Prostatitis Foundation

1063 30th Street
Box 8
Smithshire, IL 61478
Phone: 1-888-891-4200
Fax: 309-325-7184
Internet: www.prostatitis.org

The Simon Foundation for Continence

P.O. Box 835
Wilmette, IL 60091
Phone: 1-800-23-SIMON (237-4666) or
847-864-3913
Fax: 847-864-9758
Email: simoninfo@simonfoundation.org
cbgartley@simonfoundation.org
Internet: www.simonfoundation.org

National Kidney and Urologic Diseases Information Clearinghouse

3 Information Way
Bethesda, MD 20892-3580
Phone: 1-800-891-5390
Fax: 703-738-4929
Email: nkudic@info.niddk.nih.gov
Internet: www.kidney.niddk.nih.gov

The National Kidney and Urologic Diseases Information Clearinghouse (NKUDIC) is a service of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The NIDDK is part of the National Institutes of Health of the U.S. Department of Health and Human Services. Established in 1987, the Clearinghouse provides information about diseases of the kidneys and urologic system to people with kidney and urologic disorders and to their families, health care professionals, and the public. The NKUDIC answers inquiries, develops and distributes publications, and works closely with professional and patient organizations and Government agencies to coordinate resources about kidney and urologic diseases.

Publications produced by the Clearinghouse are carefully reviewed by both NIDDK scientists and outside experts.

You may also find additional information about this topic by visiting MedlinePlus at www.medlineplus.gov.

This publication is not copyrighted. The Clearinghouse encourages users of this fact sheet to duplicate and distribute as many copies as desired.

This fact sheet is also available at www.kidney.niddk.nih.gov.



U.S. DEPARTMENT OF HEALTH
AND HUMAN SERVICES
National Institutes of Health

NIH Publication No. 07-3195
August 2007