

Table 27. Extrapulmonary Tuberculosis Cases and Percentages by Site of Disease: Reporting Areas, 2010

Reporting Area	Total Extrapulmonary Cases ¹		Total Extrapulmonary Sites		Site of Disease													
	No.	%	No.	%	Pleural	Lymphatic		Bone and/or Joint		Genitourinary		Meningeal		Peritoneal		Other		
						No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.
United States	2,438	(16.1)	2,522	(40.1)	407	1,012	(40.1)	259	(10.3)	117	(4.6)	138	(5.5)	142	(5.6)	447	(17.7)	
Alabama	29	(24.1)	7	(24.1)	7	(24.1)	6	(20.7)	0	(0.0)	0	(0.0)	2	(6.9)	1	(3.4)	6	(20.7)
Alaska	8	(0.0)	3	(37.5)	1	(12.5)	1	(12.5)	1	(12.5)	1	(12.5)	0	(0.0)	1	(12.5)	2	(25.0)
Arizona	53	(7.4)	25	(46.3)	4	(7.4)	4	(7.4)	6	(11.1)	7	(13.0)	7	(13.0)	1	(1.9)	7	(13.0)
Arkansas	12	(16.7)	4	(33.3)	0	(0.0)	0	(0.0)	0	(0.0)	2	(16.7)	0	(0.0)	0	(0.0)	4	(33.3)
California	495	(14.5)	192	(38.6)	42	(8.4)	42	(8.4)	30	(6.0)	23	(4.6)	23	(4.6)	30	(6.0)	109	(21.9)
Colorado	22	(9.1)	8	(36.4)	7	(31.8)	7	(31.8)	1	(4.5)	2	(9.1)	2	(9.1)	1	(4.5)	1	(4.5)
Connecticut	21	(14.3)	16	(76.2)	1	(4.8)	1	(4.8)	0	(0.0)	0	(0.0)	0	(0.0)	1	(4.8)	0	(0.0)
Delaware	8	(37.5)	1	(12.5)	1	(12.5)	1	(12.5)	1	(12.5)	1	(12.5)	0	(0.0)	0	(0.0)	2	(25.0)
District of Columbia	14	(21.4)	2	(14.3)	3	(21.4)	3	(21.4)	1	(7.1)	0	(0.0)	0	(0.0)	1	(7.1)	4	(28.6)
Florida	156	(18.0)	65	(40.4)	10	(6.2)	10	(6.2)	6	(3.7)	5	(3.1)	5	(3.1)	13	(8.1)	33	(20.5)
Georgia	88	(22.0)	31	(34.1)	6	(6.6)	6	(6.6)	1	(1.1)	11	(12.1)	11	(12.1)	3	(3.3)	19	(20.9)
Hawaii	20	(18.2)	7	(31.8)	2	(9.1)	2	(9.1)	0	(0.0)	0	(0.0)	0	(0.0)	3	(13.6)	6	(27.3)
Idaho	3	(33.3)	2	(66.7)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Illinois	88	(16.7)	35	(36.5)	13	(13.5)	13	(13.5)	4	(4.2)	4	(4.2)	4	(4.2)	4	(4.2)	20	(20.8)
Indiana	15	(20.0)	4	(26.7)	1	(6.7)	1	(6.7)	0	(0.0)	1	(6.7)	1	(6.7)	1	(6.7)	5	(33.3)
Iowa	13	(26.7)	5	(33.3)	4	(26.7)	4	(26.7)	0	(0.0)	0	(0.0)	1	(6.7)	1	(6.7)	0	(0.0)
Kansas	16	(25.0)	7	(43.8)	4	(25.0)	4	(25.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(6.3)
Kentucky	16	(12.5)	6	(37.5)	3	(18.8)	3	(18.8)	0	(0.0)	0	(0.0)	0	(0.0)	1	(6.3)	4	(25.0)
Louisiana	43	(20.9)	11	(25.6)	2	(4.7)	2	(4.7)	5	(11.6)	2	(4.7)	2	(4.7)	1	(2.3)	13	(30.2)
Maine	5	(0.0)	5	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Maryland	62	(9.1)	29	(43.9)	10	(15.2)	10	(15.2)	1	(1.5)	5	(7.6)	5	(7.6)	3	(4.5)	12	(18.2)
Massachusetts	55	(17.9)	28	(50.0)	3	(5.4)	3	(5.4)	4	(7.1)	1	(1.8)	1	(1.8)	2	(3.6)	8	(14.3)
Michigan	53	(20.4)	17	(31.5)	5	(9.3)	5	(9.3)	0	(0.0)	5	(9.3)	5	(9.3)	1	(1.9)	15	(27.8)
Minnesota	60	(4.8)	36	(58.1)	3	(4.8)	3	(4.8)	2	(3.2)	2	(3.2)	2	(3.2)	8	(12.9)	8	(12.9)
Mississippi	16	(18.8)	4	(25.0)	4	(25.0)	4	(25.0)	1	(6.3)	1	(6.3)	1	(6.3)	0	(0.0)	3	(18.8)
Missouri	18	(33.3)	4	(22.2)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(5.6)	7	(38.9)
Nebraska	8	(12.5)	3	(37.5)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(12.5)	0	(0.0)	3	(37.5)
Nevada	28	(0.0)	16	(53.3)	7	(23.3)	7	(23.3)	1	(3.3)	0	(0.0)	0	(0.0)	1	(3.3)	5	(16.7)
New Hampshire	3	(0.0)	3	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
New Jersey	96	(15.8)	46	(45.5)	16	(15.8)	16	(15.8)	5	(5.0)	6	(5.9)	6	(5.9)	4	(4.0)	8	(7.9)
New Mexico	8	(0.0)	3	(37.5)	2	(25.0)	2	(25.0)	1	(12.5)	0	(0.0)	0	(0.0)	1	(12.5)	1	(12.5)

Table 27. (Cont'd) Extrapulmonary Tuberculosis Cases and Percentages by Site of Disease: Reporting Areas, 2010

Reporting Area	Total Extrapulmonary Cases ¹	Total Extrapulmonary Sites	Site of Disease													
			Pleural		Lymphatic		Bone and/or Joint		Genitourinary		Meningeal		Peritoneal		Other	
			No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
New York	213	225	47	(20.9)	96	(42.7)	15	(6.7)	14	(6.2)	14	(6.2)	17	(7.6)	22	(9.8)
North Carolina	68	74	13	(17.6)	25	(33.8)	11	(14.9)	1	(1.4)	6	(8.1)	2	(2.7)	16	(21.6)
North Dakota	5	5	0	(0.0)	4	(80.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(20.0)	0	(0.0)
Ohio	53	55	8	(14.5)	21	(38.2)	7	(12.7)	3	(5.5)	4	(7.3)	3	(5.5)	9	(16.4)
Oklahoma	27	28	4	(14.3)	13	(46.4)	1	(3.6)	1	(3.6)	0	(0.0)	2	(7.1)	7	(25.0)
Oregon	37	37	9	(24.3)	14	(37.8)	2	(5.4)	2	(5.4)	1	(2.7)	2	(5.4)	7	(18.9)
Pennsylvania	63	64	4	(6.3)	25	(39.1)	12	(18.8)	2	(3.1)	4	(6.3)	5	(7.8)	12	(18.8)
Rhode Island	8	10	0	(0.0)	4	(40.0)	1	(10.0)	3	(30.0)	0	(0.0)	0	(0.0)	2	(20.0)
South Carolina	32	35	9	(25.7)	11	(31.4)	4	(11.4)	3	(8.6)	1	(2.9)	3	(8.6)	4	(11.4)
South Dakota	1	1	1	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Tennessee	36	37	6	(16.2)	10	(27.0)	8	(21.6)	0	(0.0)	3	(8.1)	3	(8.1)	7	(18.9)
Texas	206	215	42	(19.5)	85	(39.5)	19	(8.8)	6	(2.8)	21	(9.8)	10	(4.7)	32	(14.9)
Utah	8	8	0	(0.0)	2	(25.0)	4	(50.0)	0	(0.0)	0	(0.0)	1	(12.5)	1	(12.5)
Vermont	2	2	0	(0.0)	2	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Virginia	53	54	5	(9.3)	28	(51.9)	6	(11.1)	3	(5.6)	1	(1.9)	3	(5.6)	8	(14.8)
Washington	73	79	13	(16.5)	38	(48.1)	6	(7.6)	7	(8.9)	1	(1.3)	5	(6.3)	9	(11.4)
West Virginia	4	5	1	(20.0)	4	(80.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Wisconsin	14	14	1	(7.1)	5	(35.7)	3	(21.4)	1	(7.1)	1	(7.1)	1	(7.1)	2	(14.3)
Wyoming	3	3	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	3	(100.0)
American Samoa ²
Fed. States of Micronesia ²	27	28	8	(28.6)	14	(50.0)	2	(7.1)	0	(0.0)	0	(0.0)	3	(10.7)	1	(3.6)
Guam ²	8	8	5	(62.5)	1	(12.5)	1	(12.5)	0	(0.0)	0	(0.0)	1	(12.5)	0	(0.0)
Marshall Islands ²	65	70	33	(47.1)	22	(31.4)	2	(2.9)	0	(0.0)	1	(1.4)	12	(17.1)	0	(0.0)
N. Mariana Islands ²
Puerto Rico ²	4	4	1	(25.0)	0	(0.0)	1	(25.0)	1	(25.0)	1	(25.0)	0	(0.0)	0	(0.0)
Republic of Palau ²	1	1	0	(0.0)	1	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
U.S. Virgin Islands ²

¹Excludes cases with pulmonary site of disease.

²Not included in U.S. totals.

Note: Ellipses indicate data not available. See Technical Notes (page 9).