

NEW YORK

Table 75. Selected Drug Use, Perceptions of Great Risk, Average Annual Marijuana Initiates, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in New York, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2009-2010 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLCIT DRUGS					
Past Month Illicit Drug Use ¹	1,610	164	546	900	1,446
Past Year Marijuana Use	2,132	216	773	1,142	1,915
Past Month Marijuana Use	1,270	130	480	660	1,140
Past Month Use of Illicit Drugs Other Than Marijuana ¹	606	61	189	357	546
Past Year Cocaine Use	369	15	144	211	355
Past Year Nonmedical Pain Reliever Use	729	79	258	391	649
Perception of Great Risk of Smoking Marijuana Once a Month	5,373	401	401	4,571	4,972
Average Annual Number of Marijuana Initiates ²	172	76	85	11	96
ALCOHOL					
Past Month Alcohol Use	9,113	253	1,490	7,371	8,861
Past Month Binge Alcohol Use ³	3,936	145	944	2,847	3,791
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	6,918	625	728	5,565	6,293
Past Month Alcohol Use (Persons Aged 12 to 20)	768 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	466 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	3,970	151	815	3,004	3,819
Past Month Cigarette Use	3,472	118	709	2,645	3,354
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	12,527	1,073	1,558	9,896	11,454
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	364	37	147	179	327
Illicit Drug Dependence or Abuse ¹	490	68	201	222	422
Alcohol Dependence	593	31	167	395	562
Alcohol Dependence or Abuse	1,204	75	371	758	1,129
Alcohol or Illicit Drug Dependence or Abuse ¹	1,498	117	480	901	1,381
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	447	61	186	200	386
Needing But Not Receiving Treatment for Alcohol Use ⁷	1,146	71	357	718	1,075
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	118	169	835	1,004
Serious Mental Illness ^{9,10}	--	--	170	543	713
Any Mental Illness ^{9,11}	--	--	679	2,363	3,041
Had Serious Thoughts of Suicide	--	--	147	424	571

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to get an overall estimate (12 or older).

⁹ For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

¹⁰ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment in carrying out major life activities.

¹¹ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, regardless of the level of impairment in carrying out major life activities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

NEW YORK

Table 76. Selected Drug Use, Perceptions of Great Risk, Average Annual Rates of First Use of Marijuana, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in New York, by Age Group: Percentages, Annual Averages Based on 2009-2010 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLCIT DRUGS					
Past Month Illicit Drug Use ¹	9.82	10.86	24.39	7.12	9.71
Past Year Marijuana Use	13.00	14.32	34.56	9.03	12.87
Past Month Marijuana Use	7.75	8.58	21.46	5.22	7.66
Past Month Use of Illicit Drugs Other Than Marijuana ¹	3.70	4.01	8.44	2.82	3.67
Past Year Cocaine Use	2.25	0.97	6.42	1.67	2.38
Past Year Nonmedical Pain Reliever Use	4.45	5.26	11.55	3.09	4.36
Perception of Great Risk of Smoking Marijuana Once a Month	32.77	26.59	17.92	36.14	33.40
Average Annual Rate of First Use of Marijuana ²	1.82	5.71	7.43	0.16	1.18
ALCOHOL					
Past Month Alcohol Use	55.58	16.75	66.60	58.28	59.53
Past Month Binge Alcohol Use ³	24.01	9.60	42.22	22.51	25.47
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	42.20	41.41	32.55	44.00	42.28
Past Month Alcohol Use (Persons Aged 12 to 20)	31.64 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	19.21 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	24.21	9.98	36.42	23.75	25.66
Past Month Cigarette Use	21.18	7.79	31.70	20.91	22.53
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	76.41	71.09	69.63	78.24	76.95
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	2.22	2.44	6.59	1.42	2.19
Illicit Drug Dependence or Abuse ¹	2.99	4.49	8.97	1.75	2.84
Alcohol Dependence	3.62	2.05	7.48	3.12	3.78
Alcohol Dependence or Abuse	7.35	4.97	16.60	5.99	7.59
Alcohol or Illicit Drug Dependence or Abuse ¹	9.14	7.76	21.45	7.13	9.28
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	2.73	4.07	8.29	1.58	2.59
Needing But Not Receiving Treatment for Alcohol Use ⁷	6.99	4.69	15.97	5.67	7.22
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	7.84	7.56	6.60	6.75
Serious Mental Illness ^{9,10}	--	--	7.58	4.30	4.79
Any Mental Illness ^{9,11}	--	--	30.34	18.68	20.43
Had Serious Thoughts of Suicide	--	--	6.57	3.35	3.84

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual marijuana initiation rate = $100 * \{ [X_1 \div (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of persons who never used marijuana. Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. Note that the age group is based on a respondent's age at the time of the interview, not his or her age at first use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to get an overall estimate (12 or older).

⁹ For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

¹⁰ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment in carrying out major life activities.

¹¹ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, regardless of the level of impairment in carrying out major life activities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).