

863 **ANIMAL TOXICOLOGY**

864 **Corneal Opacities:** Dogs receiving oral sumatriptan developed corneal opacities and defects  
865 in the corneal epithelium. Corneal opacities were seen at the lowest dosage tested, 2 mg/kg/day,  
866 and were present after 1 month of treatment. Defects in the corneal epithelium were noted in a  
867 60-week study. Earlier examinations for these toxicities were not conducted and no-effect doses  
868 were not established; the lowest dose tested is approximately 0.8 times the recommended human  
869 oral daily dose of 85 mg sumatriptan on a mg/m<sup>2</sup> basis. There was evidence of alterations in  
870 corneal appearance on the first day of intranasal dosing to dogs at all doses tested.

871

872 TREXIMET and IMITREX are registered trademarks of GlaxoSmithKline.

873 ANAPROX is a registered trademark of F. Hoffmann-La Roche Ltd.

874

875



GlaxoSmithKline

876

877 GlaxoSmithKline

878 Research Triangle Park, NC 27709

879

880 ©Year, GlaxoSmithKline. All rights reserved.

881

882 Month Year

883 TRX:XPI

884

885

**MEDICATION GUIDE**

886

**TREXIMET<sup>®</sup> [trex' i-met] Tablets**

887

**(sumatriptan and naproxen sodium)**

888

889 **What is the most important information I should know about TREXIMET?**

890 **1. TREXIMET may increase your chance of a heart attack or stroke that can lead to**  
891 **death. TREXIMET contains 2 medicines: sumatriptan and naproxen sodium (a**  
892 **nonsteroidal anti-inflammatory drug [NSAID]).**

893 **Your chance of a heart attack or stroke increases:**

- 894 • with longer use of NSAID medicines
- 895 • if you have heart disease.

896 **2. TREXIMET should never be used right before or after a heart surgery called a**  
897 **coronary artery bypass graft (CABG).**

898 **3. TREXIMET can cause ulcers and bleeding in the stomach and intestines at any time**  
899 **during your treatment.**

- 900 **Ulcers and bleeding:**  
901 • can happen without warning symptoms  
902 • may cause death.

- 903 **Your chance of getting an ulcer or bleeding increases with:**  
904 • the use of medicines called steroid hormones (corticosteroids) and blood thinners  
905 (anticoagulants)  
906 • longer use  
907 • more frequent use  
908 • smoking  
909 • drinking alcohol  
910 • older age  
911 • having poor health.

912 **4. TREXIMET is not recommended for people with risk factors for heart disease unless a**  
913 **heart exam is done and shows no problems.**

- 914 **Risk factors for heart disease include:**  
915 • high blood pressure  
916 • high cholesterol levels  
917 • smoking  
918 • obesity  
919 • diabetes  
920 • family history of heart disease  
921 • female who has gone through menopause  
922 • male over age 40.

923 **5. “Serotonin syndrome” is a serious and life-threatening problem that may occur with**  
924 **TREXIMET, especially if used with antidepressant medicines called selective serotonin**  
925 **reuptake inhibitors (SSRIs) or selective norepinephrine reuptake inhibitors (SNRIs).**

- 926 **Commonly used SSRIs are:**  
927 • CELEXA<sup>®</sup> (citalopram HBr)  
928 • LEXAPRO<sup>®</sup> (escitalopram oxalate)  
929 • PAXIL<sup>®</sup> (paroxetine)  
930 • PROZAC<sup>®</sup>/SARAFEM<sup>®</sup> (fluoxetine)  
931 • SYMBYAX<sup>®</sup> (olanzapine/fluoxetine)  
932 • ZOLOFT<sup>®</sup> (sertraline)  
933 • LUVOX<sup>®</sup> (fluvoxamine).

- 934 **Commonly used SNRIs are:**  
935 • CYMBALTA<sup>®</sup> (duloxetine)  
936 • EFFEXOR<sup>®</sup> (venlafaxine).

937 **Call your healthcare provider if you have symptoms of serotonin syndrome, which**  
938 **include:**

- 939 • mental changes (hallucinations, agitation, coma)
- 940 • fast heartbeat
- 941 • changes in blood pressure
- 942 • high body temperature or sweating
- 943 • tight muscles
- 944 • trouble walking
- 945 • nausea, vomiting, diarrhea.

946 **6. TREXIMET should only be used:**

- 947 • exactly as prescribed
- 948 • at the lowest dose possible for your treatment
- 949 • for the shortest time needed.

950 **7. TREXIMET already contains an NSAID (naproxen). Do not use TREXIMET with**  
951 **other medicines to lessen pain or fever without talking to your healthcare provider first,**  
952 **because they may contain an NSAID also.**

953

954 **What is TREXIMET?**

955 TREXIMET is a prescription medicine used to treat migraine attacks in adults. It does not  
956 prevent or lessen the number of migraines you have, and it is not for other types of headaches.  
957 TREXIMET contains 2 medicines: sumatriptan and naproxen sodium (an NSAID). This  
958 Medication Guide provides important information you need to know before taking TREXIMET.  
959 It does not take the place of talking with your healthcare provider about your medical condition  
960 or your treatment.

961

962 **How should I take TREXIMET?**

- 963 • Take 1 TREXIMET tablet to treat your migraine headache. Do not take more than 2  
964 TREXIMET tablets in 24 hours. Doses should be separated by at least 2 hours.
- 965 • TREXIMET can be taken with or without food.
- 966 • Do not split, crush, or chew TREXIMET tablets.
- 967 • If you take too much TREXIMET, call the Poison Control Center at 1-800-222-1222.

968

969 **Who should not take TREXIMET?**

970 Do not take TREXIMET right before or after heart bypass surgery. Do not take TREXIMET if  
971 you have or have had:

- 972 • uncontrolled high blood pressure
- 973 • hemiplegic or basilar migraine. (Ask your doctor if you are not sure what type of migraine  
974 you have.)

- 975 • liver problems
- 976 • an asthma attack, hives, or other allergic reaction with aspirin or any other NSAID medicine
- 977 • a heart attack or a history or symptoms of heart disease (such as chest pain or angina)
- 978 • a stroke, mini-stroke (transient ischemic attack or TIA), or other stroke-like syndrome
- 979 • problems with blood circulation to parts of your body, such as less blood flow to your
- 980 intestines (ischemic bowel disease)
- 981 • allergic reactions to sumatriptan, naproxen, or other ingredients in TREXIMET.

982 **Do not take TREXIMET if you take or have taken an antidepressant medicine called a**  
983 **monoamine oxidase (MAO) inhibitor within the last 2 weeks.** Common MAO inhibitors are  
984 isocarboxazid (MARPLAN<sup>®</sup>), phenelzine (NARDIL<sup>®</sup>), tranylcypromine (PARNATE<sup>®</sup>), and  
985 selegiline (ELDEPRYL<sup>®</sup>, EMSAM<sup>®</sup>). Ask your healthcare provider if you are not sure if your  
986 medicine is an MAO inhibitor.

987 **Do not take TREXIMET if you have taken other migraine medicines in the last 24 hours**  
988 **such as:**

- 989 • ergotamine-containing medicine or
- 990 • another triptan medicine.

991 **Before starting TREXIMET, tell your healthcare provider about:**

- 992 • all of your medical conditions including kidney or liver problems
- 993 • all allergies to any medicines
- 994 • chest pain, shortness of breath, irregular heartbeats
- 995 • medicines you may take for migraines, depression, or other health problems such as MAO
- 996 inhibitors, SSRIs, or SNRIs
- 997 • all the prescription and non-prescription medicines you take, including vitamins and herbal
- 998 supplements. Some medicines can interact with TREXIMET and cause serious side effects.

999 **Keep a list of your medicines to show to your healthcare provider. Before starting**  
1000 **TREXIMET, tell your healthcare provider if you:**

- 1001 • are pregnant, think you might be pregnant, or are trying to become pregnant. **TREXIMET**
- 1002 **should not be used by pregnant women late in their pregnancy.**
- 1003 • are breastfeeding
- 1004 • have a headache that is different from your usual migraine
- 1005 • have or have had epilepsy or seizures.

1006

1007 **What are the possible side effects of TREXIMET?**

<p><b>Serious side effects include:</b></p> <ul style="list-style-type: none"><li>• heart attack</li><li>• heartbeat problems</li><li>• stroke</li><li>• high blood pressure</li><li>• heart failure from body swelling (fluid retention)</li><li>• kidney problems including kidney failure</li><li>• bleeding and ulcers in the stomach and intestine</li><li>• low red blood cells (anemia)</li><li>• life-threatening skin reactions</li><li>• life-threatening allergic reactions</li><li>• liver problems including liver failure</li><li>• asthma attacks in people who have asthma</li><li>• loss of blood circulation to areas of your body</li><li>• serotonin syndrome (See list of symptoms in “What is the most important information I should know about TREXIMET?”)</li></ul>	<p><b>Other side effects include:</b></p> <ul style="list-style-type: none"><li>• pain, tightness, or pressure in the chest, neck, and throat</li><li>• stomach pain</li><li>• constipation</li><li>• diarrhea</li><li>• gas</li><li>• heartburn</li><li>• nausea</li><li>• vomiting</li><li>• dizziness</li><li>• drowsiness</li><li>• tiredness</li><li>• weakness</li><li>• tingling and numbness</li><li>• unusual body sensations</li><li>• redness of face (flushed)</li></ul>
--	--

1008 **Get emergency help right away if you have any of the following symptoms:**

- 1009 • shortness of breath or trouble breathing
- 1010 • chest pain
- 1011 • swelling of the face or throat
- 1012 • weakness in one part or on one side of your body
- 1013 • slurred speech.

1014 **Stop TREXIMET and call your healthcare provider right away if you have any of the following symptoms:**

- 1015
- 1016 • nausea that seems out of proportion to your migraine
- 1017 • stomach pain
- 1018 • sudden/severe pain in your belly
- 1019 • vomit blood
- 1020 • blood in your bowel movement or it is black and sticky like tar
- 1021 • itching
- 1022 • skin rash or blisters with fever
- 1023 • yellow skin or eyes
- 1024 • swelling of the arms and legs, hands, feet, face, lips, or tongue
- 1025 • unusual weight gain
- 1026 • more tired or weaker than usual
- 1027 • flu-like symptoms

1028 • serotonin syndrome. See list of symptoms in “What is the most important information I  
1029 should know about TREXIMET?”

1030 Tell your healthcare provider if you have any side effects that bother you or do not go away.  
1031 These are not all of the side effects of TREXIMET. For more information ask your healthcare  
1032 provider.

1033 Call your healthcare provider for medical advice about side effects. You may report side effects  
1034 to FDA at 1-800-FDA-1088.

1035

### 1036 **How should I store TREXIMET?**

1037 • Store TREXIMET at room temperature, 59° to 86°F (15° to 30°C).

1038 • Keep TREXIMET and all medicines out of the reach of children.

1039

### 1040 **General information about TREXIMET**

1041 • Medicines are sometimes prescribed for purposes other than those listed in a Medication  
1042 Guide. Do not use TREXIMET for a condition for which it was not prescribed.

1043 • Do not give TREXIMET to other people, even if they have the same problem you have. It  
1044 may harm them.

1045 • This Medication Guide contains the most important information about TREXIMET. If you  
1046 would like more information, talk with your healthcare provider.

1047 • You can ask your healthcare provider for information written for healthcare professionals.

1048 • For more information call 1-888-825-5249 (toll-free) or visit [www.TREXIMET.com](http://www.TREXIMET.com).

1049

### 1050 **What are the ingredients in TREXIMET?**

1051 **Active ingredients:** sumatriptan succinate and naproxen sodium

1052 **Inactive ingredients:** croscarmellose sodium, dextrose monohydrate, dibasic calcium  
1053 phosphate, FD&C Blue No. 2, lecithin, magnesium stearate, maltodextrin, microcrystalline  
1054 cellulose, povidone, sodium bicarbonate, sodium carboxymethylcellulose, talc, and titanium  
1055 dioxide.

1056

### 1057 **This Medication Guide has been approved by the U.S. Food and Drug Administration.**

1058

1059 TREXIMET, PARNATE, and PAXIL are registered trademarks of GlaxoSmithKline.

1060 The other brands listed are trademarks of their respective owners and are not trademarks of

1061 GlaxoSmithKline. The makers of these brands are not affiliated with and do not endorse

1062 GlaxoSmithKline or its products.

1063

1064



1065  
1066 GlaxoSmithKline  
1067 Research Triangle Park, NC 27709  
1068  
1069 ©Year, GlaxoSmithKline. All rights reserved.  
1070  
1071 Month Year  
1072 TRX:XMG