SCREENING TESTS FOR WOMEN

Check the guidelines listed here to find out about important screening tests for women. These guidelines are recommended by the U.S. Preventive Services Task Force. Keep in mind that these are guidelines only. Your doctor or nurse will personalize the timing of the screening tests you need based on many factors. Ask your doctor or nurse if you don't understand why a certain test is recommended for you. Check with your insurance plan to find out which tests are covered.

Screening tests	18-39	40-49	50-64	65 and older
Blood pressure test	Get tested at least every 2 years if you have normal blood pressure (lower than 120/80).	Get tested at least every 2 years if you have normal blood pressure (lower than 120/80).	Get tested at least every 2 years if you have normal blood pressure (lower than 120/80).	Get tested at least every 2 years if you have normal blood pressure (lower than 120/80).
	Get tested once a year if you have blood pressure between 120/80 and 139/89.	Get tested once a year if you have blood pressure between 120/80 and 139/89.	Get tested once a year if you have blood pressure between 120/80 and 139/89.	Get tested once a year if you have blood pressure between 120/80 and 139/89.
	Discuss treatment with your doctor or nurse if you have blood pressure 140/90 or higher.	Discuss treatment with your doctor or nurse if you have blood pressure 140/90 or higher.	Discuss treatment with your doctor or nurse if you have blood pressure 140/90 or higher.	Discuss treatment with your doctor or nurse if you have blood pressure 140/90 or higher.
Bone mineral den- sity test (osteoporosis screening)			Discuss with your doctor or nurse if you are at risk of osteoporosis.	Get this test at least once at age 65 or older. Talk to your doctor or nurse about repeat testing.
Breast cancer screen- ing (mammogram)		Discuss with your doctor or nurse.	Starting at age 50, get screened every 2 years.	Get screened every 2 years through age 74. Age 75 and older, ask your doctor or nurse if you need to be screened.
Cervical cancer screen- ing (Pap test)*	Get a Pap test every 3 years if you are 21 or older, have had vaginal sex, and have a cervix.	Get a Pap test every 3 years if you have had vaginal sex, and have a cervix.	Get a Pap test every 3 years if you have had vaginal sex, and have a cervix.	Ask your doctor or nurse if you need to get a Pap test.
Chlamydia test	Get tested for chlamydia yearly through age 24 if you are sexually active or pregnant. Age 25 and older, get tested for chlamydia if you are at increased risk, pregnant or not pregnant.	Get tested for chlamydia if you are sexually active and at increased risk, pregnant or not pregnant.	Get tested for chlamydia if you are sexually active and at increased risk.	Get tested for chlamydia if you are sexually active and at increased risk.
Cholesterol test	Starting at age 20, get a cholesterol test regularly if you are at increased risk for heart disease.	Get a cholesterol test regularly if you are at increased risk for heart disease.	Get a cholesterol test regularly if you are at increased risk for heart disease.	Get a cholesterol test regularly if you are at increased risk for heart disease.
	Ask your doctor or nurse how often you need your cholesterol tested.	Ask your doctor or nurse how often you need your cholesterol tested.	Ask your doctor or nurse how often you need your cholesterol tested.	Ask your doctor or nurse how often you need your cholesterol tested.

^{*} Final U.S. Preventive Services Task Force guidelines on cervical cancer screening can be found at www.uspreventiveservicestaskforce.org/uspstf/uspscerv.htm





SCREENING TESTS FOR WOMEN

Screening tests	18-39	40-49	50-64	65 and older
Colorectal cancer screening (using fecal occult blood testing, sig-			Starting at age 50, get screened for colorectal cancer.	Get screened for colorectal cancer through age 75.
moidoscopy, or colonos- copy)			Talk to your doctor or nurse about which screening test is best for you and how often you need it.	Talk to your doctor or nurse about which screening test is best for you and how often you need it.
Diabetes screening	Get screened for diabetes if your blood pressure is higher than 135/80 or if you take medicine for high blood pressure.	Get screened for diabetes if your blood pressure is higher than 135/80 or if you take medicine for high blood pressure.	Get screened for diabetes if your blood pressure is higher than 135/80 or if you take medicine for high blood pressure.	Get screened for diabetes if your blood pressure is higher than 135/80 or if you take medicine for high blood pressure.
Gonorrhea test	Get tested for gonorrhea if you are sexually active and at increased risk, pregnant or not pregnant.	Get tested for gonorrhea if you are sexually active and at increased risk, pregnant or not pregnant.	Get tested for gonorrhea if you are sexually active and at increased risk.	Get tested for gonorrhea if you are sexually active and at increased risk.
HIV test	Get tested if you are at increased risk for HIV.	Get tested if you are at increased risk for HIV.	Get tested if you are at increased risk for HIV.	Get tested if you are at increased risk for HIV.
	Discuss your risk with your doctor or nurse.	Discuss your risk with your doctor or nurse.	Discuss your risk with your doctor or nurse.	Discuss your risk with your doctor or nurse.
	All pregnant women need to be tested for HIV.	All pregnant women need to be tested for HIV.		
Syphilis test	Get tested for syphilis if you are at increased risk or pregnant.	Get tested for syphilis if you are at increased risk or pregnant.	Get tested for syphilis if you are at increased risk.	Get tested for syphilis if you are at increased risk.

Vaccines

Vaccines can protect you from harmful infections. Some adults think only children need vaccines. But this is not true. To find out what vaccines you may need, visit http://www.vaccines.gov.

Content last updated: January 19, 2012



