Diabetes	Heart Failure	Coronary Artery Disease	Preventive Care (measured on population with specified chronic diseases)
DM-1 HbA1c	HF-1 Left Ventricular	CAD-1 Antiplatelet	PC-1Blood Pressure
Management	Function Assessment	Therapy	Measurement
DM-2 HbA1c	HF-2 Left Ventricular	CAD-2 Drug Therapy for	PC-5 Breast Cancer
Control	Ejection Fraction Testing	Lowering LDL Cholesterol	Screening
DM-3 Blood	HF-3 Weight	CAD-3 Beta Blocker	PC-6 Colorectal Cancer
Pressure	Measurement	Therapy – Prior MI	Screening
Management			
DM-4 Lipid	HF-5 Patient	CAD-5 Lipid Profile	PC-7 Influenza
Measurement	Education	-	Vaccination
DM-5 LDL	HF-6 Beta Blocker	CAD-6 LDL Cholesterol	PC-8 Pneumonia
Cholesterol	Therapy	Level	Vaccination
Level			
DM-6 Urine	HF-7 ACE	CAD-7 ACE	
Protein Testing	Inhibitor/ARB	Inhibitor/ARB Therapy	
	Therapy		
DM-7 Eye	HF-8 Warfarin		
Exam	Therapy for Patients		
	with AF		
DM-8 Foot			
Exam			

Table 1: Clinical Quality Measures in the MCMP Demonstration¹

¹ Updated 1/8/2007. Measure specifications are subject to change and may be updated during the course of the demonstration.

Diabetes Mellitus

- 1. HbA1c Management: Testing (DM-1): The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had Hemoglobin A1c (HBA1c) testing
- 2. HbA1c Management: Poor Control (DM-2): The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had HbA1c in poor control (>9.0%)
- 3. Blood Pressure Management (DM-3): The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had a BP < 140/80 mmHg
- 4. Lipid Management Testing: (DM-4): The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had LDL-C screening performed
- 5. Lipid Management: Control < 100mg/dl (DM-5): The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had LDL-C testing l <100 mg/dl
- 6. Urine Protein Screening (DM-6): The percentage of patients 18-75 year of age with diabetes (type 1 or type 2) who had medical attention for nephropathy
- 7. Eye Examination (DM-7): The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had an eye exam (retinal) performed
- 8. Foot Examination (DM-8): The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had a foot exam (visual inspection, sensory exam with monofilament, and pulse exam)

Congestive Heart Failure

- 1. LVF Assessment (HF-1): Percentage of patients with quantitative or qualitative results for LVF assessment
- 2. Left Ventricular Function (LVF) Testing (HF-2): Percentage of patients with LVF testing during the current year for patients hospitalized with a principle diagnosis of HF during the current year
- 3. Weight Measurement (HF-3): Percentage of HF patient visits with weight measurement recorded
- 5. Patient Education (HF-5): Percentage of patients who were provided with patient education on disease management and health behavior changes during one or more visit(s)
- 6. Beta-Blocker Therapy (HF-6): Percentage of patients who were prescribed beta-blocker therapy

- 7. ACE Inhibitor/ARB Therapy (HF-7): Percentage of patients who were prescribed ACE inhibitor or ARB therapy
- 8. Warfarin Therapy for Patients with Atrial Fibrillation (HF-8): Percentage of patients with paroxysmal or chronic atrial fibrillation who were prescribed warfarin therapy

Coronary Artery Disease

- 1. Antiplatelet Therapy (CAD-1): Percentage of patients who were prescribed antiplatelet therapy
- 2. Drug Therapy for Lowering LDL Cholesterol (CAD-2): Percentage of patients who were prescribed a lipid-lowering therapy (based on current ACC/AHA guidelines)
- 3. Beta-Blocker Therapy Prior Myocardial Infarction (MI) (CAD-3): Percentage of patients with prior MI at anytime who were prescribed beta-blocker therap.
- 5. Lipid Profile (CAD-5): Percentage of patients who received at least one lipid profile (or ALL components tests)
- 6. LDL Cholesterol Level (CAD-6): Percentage of patients with most recent LDL cholesterol <100 mg/dl
- 7. ACE Inhibitor or ARB Therapy (CAD-7): Percentage of patients who also have diabetes and/or LVSD who were prescribed ACE inhibitor or ARB therapy

Preventive Care

- 1. Blood Pressure Measurement (PC-1): Percentage of patient visits with blood pressure (BP) measurement recorded
- 5. Breast Cancer Screening (PC-5): The percentage of women 40-69 years of age who had a mammogram to screen for breast cancer
- 6. Colorectal Cancer Screening (PC-6): Percentage of patients screened for colorectal cancer during the one-year measurement period
- 7. Influenza Immunization (PC-7): Percentage of patients who received an influenza immunization during the one-year measurement period
- 8. Pneumonia Vaccination (PC-8): The percentage of patients \geq 65 years and older who ever received a pneumococcal vaccination