PREPARATION INSTRUCTIONS FOR COLONOSCOPY with Golytely

Do these things 7 DAYS BEFORE the procedure:

- Arrange a ride: you will be given medicine that makes you relax and be sleepy, so you cannot drive a car or take a bus home. If you arrive without an escort, your procedure may need to be rescheduled.
- Stop taking iron, Vitamin E, ginko, ginger, aspirin and garlic
- If you are taking Coumadin medicine or if you are diabetic, call your doctor or nurse for special instructions.

Do these things **3 DAYS BEFORE** the procedure:

- Confirm your ride.
- If you need to cancel your appointment, call your doctor or nurse
- Review the diet you need to follow for the next 2 days. Plan your meals according to this diet.

Do these things **2 DAYS BEFORE** the procedure:

- Drink at least 8 glasses of water throughout the day.
- Stop taking all anti-inflammatory medicines. These include: Nuprin, Bufferin, Anacin, Advil, Motrin, Indocin, Voltran, Aleve, Ecotrin, Relafen, Alka Seltzer, Naprosyn, Voltaren (Tylenol is okay to take)
- Stop eating seeds, popcorn, nuts and whole grains
- Do not eat any solid food after midnight

Appointment date: _	
Appointment time: _	
Location:	

Do these things 1 DAY BEFORE the procedure:

- •Start a strict, clear liquid diet. If you put a liquid in a clear glass, and you can see through it, it is probably OK to drink. A clear liquid diet can include:
- -Apple, white grape, and cranberry juices;
- -Beef or chicken broths that are clear:
- -Tea and coffee without milk:
- -Soda pop, Gatorade, Kool-Aid and various -Jell-O flavors (any color except red)
- •Avoid: juices with pulp, milk, cream and solid food.
- •Fill the plastic container containing the laxative powder (**Golytely**) with water, shake well until the powder is dissolved and put the container in the refrigerator.
- •1:30pm: Take the first tablet of **Metoclopramide** by mouth with water. The medicine Metoclopramide will help relieve nausea caused by Golytely.
- •2:00pm: Start drinking the laxative (Golytely) liquid. Drink one full glass every 15 minutes until it is gone. Try to drink one full glass at a time instead of sipping.
- •5:00pm: Take the second tablet of Metoclopramide by mouth with water.
- •Continue to drink clear liquids throughout the day but do not eat any solid food.
- •The laxative (Golytely) liquid will cause you to have many bowel movements. This is needed so you have a clean colon for the procedure. If you have problems completing this preparation, call your doctor or nurse.

Do these things **ON THE DAY OF** the procedure:

- Take your normal medications with a sip of water.
- •You may have clear liquids until 3 hours before the procedure.
- Bring your escort/driver with you to your appointment.
- Arrive 30 minutes before your appointment time.

Do you have QUESTIONS or PROBLEMS completing the instructions? Call:_

There are many different kinds of medical tests and they require different kinds of preparation. For example, some tests require fasting while other tests do not. Talk to your doctor about the test preparation that is right for you.