

This announcement signals the release of the 2010 Mortality Multiple Cause Micro-data Files. The file can be accessed at: http://www.cdc.gov/nchs/data_access/Vitalstatsonline.htm

This document contains the Detailed Tables for the National Vital Statistics Report (NVSR) "*Deaths: Final Data for 2010*." The complete version of this report (NVSR Volume 61, Number 4) will be published in October. Most of the tables make reference (in headnotes and footnotes) to the Technical notes section of the report, which will not be available until publication of the complete report. In the interim, the Technical notes section of the previous report, "*Deaths: Final Data for 2009*" is available for your reference at: http://www.cdc.gov/nchs/data/nvsr/nvsr60/nvsr60_03.pdf. The populations used to calculate death rates were produced under a collaborative arrangement with the U.S. Census Bureau. Populations for 2010 and the intercensal period 2001–2009 are consistent with the 2010 census. Death rates for 2001-2009 have been revised using the intercensal populations and may differ from rates previously published. If you have any questions about the Detailed Tables in this document, please send an email to MortalityData@cdc.gov.

List of Detailed Tables

1. Number of deaths, death rates, and age-adjusted death rates, by race and sex: United States, 1940, 1950, 1960, 1970, and 1980–2010
2. Number of deaths, death rates, and age-adjusted death rates, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1997–2010
3. Number of deaths and death rates, by age, race, and sex: United States, 2010
4. Number of deaths and death rates by Hispanic origin, race for non-Hispanic population, age, and sex: United States, 2010
5. Number of deaths and death rates by age, and age-adjusted death rates, by specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2010 (forthcoming)
6. Abridged life table for the total population, 2010
7. Life expectancy at selected ages by race, Hispanic origin, race for non-Hispanic population, and sex: United States, 2010
8. Life expectancy at birth by race, Hispanic origin, race for non-Hispanic population, and sex: United States, 1940, 1950, 1960, 1970 and 1975-2010
9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2010: United States, 1999–2010
10. Number of deaths from 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010
11. Death rates for 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010
12. Number of deaths from 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010
13. Number of deaths from 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010
14. Death rates for 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010
15. Death rates for 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010
17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to Clostridium difficile, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010
18. Number of deaths, death rates, and age-adjusted death rates for injury deaths, by mechanism and intent of death: United States, 2010
19. Number of deaths, death rates, and age-adjusted death rates for major causes of death: United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, 2010
20. Infant, neonatal, and postneonatal mortality rates by race and sex: United States, 1940, 1950, 1960, 1970, and 1975–2010
21. Number of infant deaths and infant mortality rates for 130 selected causes by race: United States, 2010
22. Number of infant and neonatal deaths and mortality rates, by race for the United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, and by sex for the United States, 2010

Table 1. Number of deaths, death rates, and age-adjusted death rates, by race and sex: United States, 1940, 1950, 1960, 1970, and 1980-2010

[Crude rates on an annual basis per 100,000 population in specified age group; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and estimated as of July 1 for all other years; see "Technical Notes." Beginning 1970, excludes deaths of nonresidents of the United States. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards]

Year	All races ¹			White ²			Black ²			American Indian or Alaska Native ^{2,3}			Asian or Pacific Islander ^{2,4}		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
	Number														
2010	2,468,435	1,232,432	1,236,003	2,114,749	1,051,514	1,063,235	286,959	145,802	141,157	15,565	8,516	7,049	51,162	26,600	24,562
2009	2,437,163	1,217,379	1,219,784	2,086,355	1,037,475	1,048,880	286,623	146,239	140,384	14,960	8,105	6,855	49,225	25,560	23,665
2008	2,471,984	1,226,197	1,245,787	2,120,233	1,046,183	1,074,050	289,072	147,143	141,929	14,776	8,163	6,613	47,903	24,708	23,195
2007	2,423,712	1,203,968	1,219,744	2,074,151	1,023,951	1,050,200	289,585	148,309	141,276	14,367	7,885	6,482	45,609	23,823	21,786
2006	2,426,264	1,201,942	1,224,322	2,077,549	1,022,328	1,055,221	289,971	148,602	141,369	14,037	7,630	6,407	44,707	23,382	21,325
2005	2,448,017	1,207,675	1,240,342	2,098,097	1,028,152	1,069,945	292,808	149,108	143,700	13,918	7,607	6,311	43,194	22,808	20,386
2004	2,397,615	1,181,668	1,215,947	2,056,643	1,007,266	1,049,377	287,315	145,970	141,345	13,124	7,134	5,990	40,533	21,298	19,235
2003	2,448,288	1,201,964	1,246,324	2,103,714	1,025,650	1,078,064	291,300	148,022	143,278	13,147	7,106	6,041	40,127	21,186	18,941
2002	2,443,387	1,199,264	1,244,123	2,102,589	1,025,196	1,077,393	290,051	146,835	143,216	12,415	6,750	5,665	38,332	20,483	17,849
2001	2,416,425	1,183,421	1,233,004	2,079,691	1,011,218	1,068,473	287,709	145,908	141,801	11,977	6,466	5,511	37,048	19,829	17,219
2000	2,403,351	1,177,578	1,225,773	2,071,287	1,007,191	1,064,096	285,826	145,184	140,642	11,363	6,185	5,178	34,875	19,018	15,857
1999	2,391,399	1,175,460	1,215,939	2,061,348	1,005,335	1,056,013	285,064	145,703	139,361	11,312	6,092	5,220	33,675	18,330	15,345
1998	2,337,256	1,157,260	1,179,996	2,015,984	990,190	1,025,794	278,440	143,417	135,023	10,845	5,994	4,851	31,987	17,659	14,328
1997	2,314,245	1,154,039	1,160,206	1,996,393	986,884	1,009,509	276,520	144,110	132,410	10,576	5,985	4,591	30,756	17,060	13,696
1996	2,314,690	1,163,569	1,151,121	1,992,966	991,984	1,000,982	282,089	149,472	132,617	10,127	5,563	4,564	29,508	16,550	12,958
1995	2,312,132	1,172,959	1,139,173	1,987,437	997,277	990,160	286,401	154,175	132,226	9,997	5,574	4,423	28,297	15,933	12,364
1994	2,278,994	1,162,747	1,116,247	1,959,875	988,823	971,052	282,379	153,019	129,360	9,637	5,497	4,140	27,103	15,408	11,695
1993	2,268,553	1,161,797	1,106,756	1,951,437	988,329	963,108	282,151	153,502	128,649	9,579	5,434	4,145	25,386	14,532	10,854
1992	2,175,613	1,122,336	1,053,277	1,873,781	956,957	916,824	269,219	146,630	122,589	8,953	5,181	3,772	23,660	13,568	10,092
1991	2,169,518	1,121,665	1,047,853	1,868,904	956,497	912,407	269,525	147,331	122,194	8,621	4,948	3,673	22,173	12,727	9,446
1990	2,148,463	1,113,417	1,035,046	1,853,254	950,812	902,442	265,498	145,359	120,139	8,316	4,877	3,439	21,127	12,211	8,916
1989	2,150,466	1,114,190	1,036,276	1,853,841	950,852	902,989	267,642	146,393	121,249	8,614	5,066	3,548	20,042	11,688	8,354
1988	2,167,999	1,125,540	1,042,459	1,876,906	965,419	911,487	264,019	144,228	119,791	7,917	4,617	3,300	18,963	11,155	7,808
1987	2,123,323	1,107,958	1,015,365	1,843,067	953,382	889,685	254,814	139,551	115,263	7,602	4,432	3,170	17,689	10,496	7,193
1986	2,105,361	1,104,005	1,001,356	1,831,083	952,554	878,529	250,326	137,214	113,112	7,301	4,365	2,936	16,514	9,795	6,719
1985	2,086,440	1,097,758	988,682	1,819,054	950,455	868,599	244,207	133,610	110,597	7,154	4,181	2,973	15,887	9,441	6,446
1984	2,039,369	1,076,514	962,855	1,781,897	934,529	847,368	235,884	129,147	106,737	6,949	4,117	2,832	14,483	8,627	5,856
1983	2,019,201	1,071,923	947,278	1,765,582	931,779	833,803	233,124	127,911	105,213	6,839	4,064	2,775	13,554	8,126	5,428
1982	1,974,797	1,056,440	918,357	1,729,085	919,239	809,846	226,513	125,610	100,903	6,679	3,974	2,705	12,430	7,564	4,866
1981	1,977,981	1,063,772	914,209	1,731,233	925,490	805,743	228,560	127,296	101,264	6,608	4,016	2,592	11,475	6,908	4,567
1980	1,989,841	1,075,078	914,763	1,738,607	933,878	804,729	233,135	130,138	102,997	6,923	4,193	2,730	11,071	6,809	4,262
1970	1,921,031	1,078,478	842,553	1,682,096	942,437	739,659	225,647	127,540	98,107	5,675	3,391	2,284	---	---	---
1960	1,711,982	975,648	736,334	1,505,335	860,857	644,478	196,010	107,701	88,309	4,528	2,658	1,870	---	---	---
1950	1,452,454	827,749	624,705	1,276,085	731,366	544,719	169,606	92,004	77,602	4,440	2,497	1,943	---	---	---
1940	1,417,269	791,003	626,266	1,231,223	690,901	540,322	178,743	95,517	83,226	4,791	2,527	2,264	---	---	---

Table 1. Number of deaths, death rates, and age-adjusted death rates, by race and sex: United States, 1940, 1950, 1960, 1970, and 1980-2010

[Crude rates on an annual basis per 100,000 population in specified age group; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and estimated as of July 1 for all other years; see "Technical Notes." Beginning 1970, excludes deaths of nonresidents of the United States. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards]

Year	All races ¹			White ²			Black ²			American Indian or Alaska Native ^{2,3}			Asian or Pacific Islander ^{2,4}		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
	Age-adjusted death rate ⁵														
2010	747.0	887.1	634.9	741.8	878.5	630.8	898.2	1,104.0	752.5	628.3	730.2	541.7	424.3	512.1	359.0
2009 ⁵	749.6	890.9	636.8	742.8	880.5	631.3	912.7	1,123.1	763.3	618.2	711.5	538.2	424.2	508.6	360.8
2008 ⁵	774.9	918.8	659.9	767.2	907.1	653.7	947.6	1,168.0	792.0	646.4	759.8	550.8	434.7	518.0	372.1
2007 ⁵	775.3	922.9	658.1	764.3	907.1	649.4	972.1	1,205.0	808.2	663.2	782.9	566.8	435.8	525.3	368.9
2006 ⁵	791.8	943.5	672.2	779.3	925.8	662.3	997.9	1,239.6	828.4	678.4	782.0	591.2	450.3	544.4	380.8
2005 ⁵	815.0	971.9	692.3	801.1	952.9	680.9	1,035.1	1,281.4	862.6	703.9	828.0	604.2	459.2	560.2	384.8
2004 ⁵	813.7	973.3	690.5	798.5	953.3	677.7	1,043.8	1,296.8	869.8	694.0	814.3	596.7	460.2	556.8	388.7
2003 ⁵	843.5	1,010.3	715.2	827.1	988.8	701.6	1,080.5	1,343.4	898.3	728.8	854.2	630.0	480.0	583.0	403.7
2002 ⁵	855.9	1,030.6	723.6	839.0	1,009.0	709.3	1,097.3	1,364.9	913.4	715.5	844.5	613.1	486.0	594.7	405.1
2001 ⁵	858.8	1,035.4	725.6	840.7	1,012.1	710.4	1,106.2	1,380.5	917.9	716.6	837.6	619.1	495.1	603.4	413.7
2000	869.0	1,053.8	731.4	849.8	1,029.4	715.3	1,121.4	1,403.5	927.6	709.3	841.5	604.5	506.4	624.2	416.8
1999	875.6	1,067.0	734.0	854.6	1,040.0	716.6	1,135.7	1,432.6	933.6	780.9	925.9	668.2	519.7	641.2	427.5
1998	870.6	1,069.4	724.7	849.3	1,042.0	707.3	1,127.8	1,430.5	921.6	770.4	943.9	640.5	522.4	646.9	426.7
1997	878.1	1,088.1	725.6	855.7	1,059.1	707.8	1,139.8	1,458.8	922.1	774.0	974.8	625.3	531.8	660.2	432.6
1996	894.1	1,115.7	733.0	869.0	1,082.9	713.6	1,178.4	1,524.2	940.3	763.6	924.8	641.7	543.2	676.1	439.6
1995	909.8	1,143.9	739.4	882.3	1,107.5	718.7	1,213.9	1,585.7	955.9	771.2	932.0	643.9	554.8	693.4	446.7
1994	913.5	1,155.5	738.6	885.6	1,118.7	717.5	1,216.9	1,592.8	954.6	764.8	953.3	618.8	562.7	702.5	452.1
1993	926.1	1,177.3	745.9	897.0	1,138.9	724.1	1,241.2	1,632.2	969.5	796.4	1,006.3	641.6	565.8	709.9	450.4
1992	905.6	1,158.3	725.5	877.7	1,122.4	704.1	1,206.7	1,587.8	942.5	759.0	970.4	599.4	558.5	697.3	445.8
1991	922.3	1,180.5	738.2	893.2	1,143.1	716.1	1,235.4	1,626.1	963.3	763.9	970.6	608.3	566.2	703.4	453.2
1990	938.7	1,202.8	750.9	909.8	1,165.9	728.8	1,250.3	1,644.5	975.1	716.3	916.2	561.8	582.0	716.4	469.3
1989	950.5	1,215.0	761.8	920.2	1,176.6	738.8	1,275.5	1,670.1	998.1	761.6	999.8	586.3	581.3	729.6	458.4
1988	975.7	1,250.7	781.0	947.6	1,215.9	759.1	1,284.3	1,677.6	1,006.8	718.6	917.4	563.6	584.2	732.0	451.0
1987	970.0	1,246.1	774.2	943.4	1,213.4	753.3	1,263.1	1,650.3	989.7	719.8	899.3	583.7	577.3	732.4	448.1
1986	978.6	1,261.7	778.7	952.8	1,230.5	758.1	1,266.7	1,650.1	994.4	720.8	926.7	549.3	576.4	730.5	445.4
1985	988.1	1,278.1	784.5	963.6	1,249.8	764.3	1,261.2	1,634.5	994.4	731.7	926.1	577.2	586.5	755.4	456.7
1984	982.5	1,271.4	779.8	959.7	1,245.9	760.7	1,236.7	1,600.8	976.9	761.7	946.0	567.9	574.4	724.7	443.1
1983	990.0	1,284.5	783.3	967.3	1,259.4	763.9	1,240.5	1,600.7	980.7	757.3	945.0	605.5	565.1	718.8	428.8
1982	985.0	1,279.9	776.6	963.6	1,255.9	758.7	1,221.3	1,580.4	960.1	757.0	940.1	604.4	550.4	738.2	410.3
1981	1,007.1	1,308.2	792.7	984.0	1,282.2	773.6	1,258.4	1,626.6	986.6	784.6	1,030.2	588.0	544.7	710.3	405.3
1980	1,039.1	1,348.1	817.9	1,012.7	1,317.6	796.1	1,314.8	1,697.8	1,033.3	867.0	1,111.5	662.4	589.9	786.5	425.9
1970	1,222.6	1,542.1	971.4	1,193.3	1,513.7	944.0	1,518.1	1,873.9	1,228.7	---	---	---	---	---	---
1960	1,339.2	1,609.0	1,105.3	1,311.3	1,586.0	1,074.4	1,577.5	1,811.1	1,369.7	---	---	---	---	---	---
1950	1,446.0	1,674.2	1,236.0	1,410.8	1,642.5	1,198.0	---	---	---	---	---	---	---	---	---
1940	1,785.0	1,976.0	1,599.4	1,735.3	1,925.2	1,550.4	---	---	---	---	---	---	---	---	---

--- Data not available.

¹For 1940-91, data includes deaths among races not shown separately; beginning in 1992, records coded as "other races" and records for which race was unknown, not stated, or not classifiable were assigned to the race of previous record; see "Technical Notes."

²Multiple-race data were reported by 37 states and the District of Columbia in 2010, by 34 states and the District of Columbia in 2008 and 2009, by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2005, by 15 states in 2004, and by 7 states in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

³Includes Aleuts and Eskimos.

⁴Includes Chinese, Filipino, Hawaiian, Japanese, and other Asian or Pacific Islander groups.

⁵Rates are revised and may differ from those previously published; see "Technical Notes."

⁶For method of computation, see "Technical Notes."

Table 2. Number of deaths, death rates, and age-adjusted death rates, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1997-2010

[Crude rates on an annual basis per 100,000 population in specified group; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and are estimated as of July 1 for all other years; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes"]

Year	All origins ¹			Hispanic			Non-Hispanic ²			Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Number															
2010	2,468,435	1,232,432	1,236,003	144,490	79,622	64,868	2,318,218	1,149,438	1,168,780	1,969,916	971,604	998,312	283,438	143,824	139,614
2009	2,437,163	1,217,379	1,219,784	141,576	78,157	63,419	2,289,999	1,135,852	1,154,147	1,944,606	959,014	985,592	282,982	144,197	138,785
2008	2,471,984	1,226,197	1,245,787	139,241	76,861	62,380	2,327,636	1,146,394	1,181,242	1,981,034	969,288	1,011,746	285,522	145,168	140,354
2007	2,423,712	1,203,968	1,219,744	135,519	75,708	59,811	2,284,446	1,125,974	1,158,472	1,939,606	948,662	990,944	286,366	146,474	139,892
2006	2,426,264	1,201,942	1,224,322	133,004	74,250	58,754	2,288,424	1,124,813	1,163,611	1,944,617	947,966	996,651	286,581	146,729	139,852
2005	2,448,017	1,207,675	1,240,342	131,161	73,788	57,373	2,312,028	1,131,013	1,181,015	1,967,142	954,402	1,012,740	289,163	147,010	142,153
2004	2,397,615	1,181,668	1,215,947	122,416	68,544	53,872	2,269,583	1,109,848	1,159,735	1,933,382	938,143	995,239	283,859	144,022	139,837
2003	2,448,288	1,201,964	1,246,324	122,026	68,119	53,907	2,319,476	1,129,927	1,189,549	1,979,465	956,194	1,023,271	287,968	146,136	141,832
2002	2,443,387	1,199,264	1,244,123	117,135	65,703	51,432	2,318,269	1,129,090	1,189,179	1,981,973	957,645	1,024,328	286,573	144,802	141,771
2001	2,416,425	1,183,421	1,233,004	113,413	63,317	50,096	2,295,244	1,115,683	1,179,561	1,962,810	945,967	1,016,843	284,343	143,971	140,372
2000	2,403,351	1,177,578	1,225,773	107,254	60,172	47,082	2,287,846	1,112,704	1,175,142	1,959,919	944,781	1,015,138	282,676	143,297	139,379
1999	2,391,399	1,175,460	1,215,939	103,740	57,991	45,749	2,279,325	1,112,718	1,166,607	1,953,197	944,913	1,008,284	281,979	143,883	138,096
1998	2,337,256	1,157,260	1,179,996	98,406	55,821	42,585	2,230,127	1,096,677	1,133,450	1,912,802	931,844	980,958	275,264	141,627	133,637
1997	2,314,245	1,154,039	1,160,206	95,460	54,348	41,112	2,209,450	1,094,541	1,114,909	1,895,461	929,703	965,758	273,381	142,241	131,140
Death rate															
2010	799.5	812.0	787.4	286.2	310.8	260.9	897.6	911.1	884.7	984.3	987.5	981.2	718.7	764.5	676.9
2009 ⁴	794.5	807.2	782.1	287.0	311.8	261.4	889.5	903.3	876.3	972.3	975.7	969.1	723.5	773.1	678.4
2008 ⁴	812.9	820.3	805.8	291.3	316.0	265.8	908.2	915.9	900.8	991.6	987.5	995.6	738.6	787.7	693.9
2007 ⁴	804.6	813.1	796.4	293.4	321.6	264.0	895.7	904.2	887.6	972.3	968.3	976.1	749.9	804.8	699.9
2006 ⁴	813.1	819.6	806.9	298.2	326.1	269.0	901.8	908.0	895.8	976.2	969.4	982.8	759.7	816.5	708.1
2005 ⁴	828.4	831.7	825.1	304.9	335.6	272.7	915.7	918.0	913.5	989.1	978.1	999.7	775.8	828.4	728.0
2004 ⁴	818.8	821.6	816.2	295.0	322.8	265.8	903.1	905.3	901.0	973.4	963.2	983.2	770.2	821.1	724.0
2003 ⁴	843.9	843.9	843.9	304.7	332.0	276.0	927.6	926.8	928.3	998.3	984.1	1,011.8	790.5	843.6	742.4
2002 ⁴	849.5	849.2	849.8	303.3	331.5	273.6	931.0	930.0	932.0	1,000.5	986.7	1,013.8	794.8	844.3	749.9
2001 ⁴	848.0	846.0	849.9	305.3	331.8	277.4	926.2	923.5	928.7	992.1	976.3	1,007.2	797.8	849.5	750.9
2000	854.0	853.0	855.0	303.8	331.3	274.6	929.6	928.1	931.0	993.2	978.5	1,007.3	805.5	859.5	756.7
1999	857.0	859.2	854.9	305.7	332.6	277.2	929.9	932.2	927.8	990.7	979.6	1,001.3	812.1	872.8	757.3
1998	847.3	856.4	838.5	303.9	336.0	270.0	916.0	925.3	907.1	972.9	969.2	976.5	805.6	873.7	744.1
1997	848.8	864.6	833.6	309.0	343.2	272.9	913.9	930.4	898.3	967.4	970.6	964.3	813.5	892.9	741.9
Age-adjusted death rate ⁵															
2010	747.0	887.1	634.9	558.6	677.7	463.4	762.6	904.6	649.2	755.0	892.5	643.3	920.4	1,131.7	770.8
2009 ⁴	749.6	890.9	636.8	559.7	675.5	466.1	764.7	908.0	650.5	755.1	893.7	643.1	934.4	1,150.5	781.0
2008 ⁴	774.9	918.8	659.9	579.8	695.3	484.7	790.0	935.9	673.7	779.4	920.2	665.4	969.2	1,195.5	809.6
2007 ⁴	775.3	922.9	658.1	586.1	711.4	484.4	789.5	938.7	671.4	775.3	918.4	660.6	994.6	1,233.6	826.5
2006 ⁴	791.8	943.5	672.2	604.0	732.3	500.2	804.9	958.0	684.6	789.1	935.7	672.4	1,019.5	1,267.2	845.7
2005 ⁴	815.0	971.9	692.3	627.6	771.2	513.8	827.3	985.0	704.4	810.1	961.5	690.7	1,055.2	1,306.2	879.4
2004 ⁴	813.7	973.3	690.5	616.8	750.1	509.5	825.9	986.7	702.2	807.6	962.5	687.2	1,062.9	1,321.0	885.5
2003 ⁴	843.5	1,010.3	715.2	645.3	784.0	534.2	854.6	1,022.6	725.8	834.9	996.7	709.9	1,099.1	1,366.8	913.7
2002 ⁴	855.9	1,030.6	723.6	652.2	799.9	535.9	866.4	1,042.1	733.8	846.4	1,016.5	717.1	1,114.2	1,385.3	927.9
2001 ⁴	858.8	1,035.4	725.6	662.6	808.6	547.0	868.4	1,046.1	734.9	847.1	1,018.8	717.3	1,122.4	1,400.5	931.6
2000	869.0	1,053.8	731.4	665.7	818.1	546.0	877.9	1,063.8	740.0	855.5	1,035.4	721.5	1,137.0	1,422.0	941.2
1999	875.6	1,067.0	734.0	676.4	830.5	555.9	883.9	1,076.4	741.9	859.8	1,045.5	722.3	1,150.1	1,449.4	946.0
1998	870.6	1,069.4	724.7	665.4	833.6	536.9	878.4	1,078.2	732.4	854.1	1,046.7	712.8	1,141.8	1,448.2	932.9
1997	878.1	1,088.1	725.6	669.3	840.5	538.8	885.3	1,096.4	732.6	859.7	1,063.2	712.5	1,154.3	1,476.7	934.2

¹Figures for origin not stated are included in "All origins" but are not distributed among specified origins.

²Includes races other than white and black.

³Multiple-race data were reported by 37 states and the District of Columbia in 2010, by 34 states and the District of Columbia in 2008 and 2009, by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2005, by 15 states in 2004, and by 7 states in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴Rates are revised and may differ from those previously published; see "Technical Notes."

⁵For method of computation, see "Technical Notes."

Table 3. Number of deaths and death rates, by age, race, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes"]

Age	All races			White ¹			Black ¹			American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
	Number														
All ages	2,468,435	1,232,432	1,236,003	2,114,749	1,051,514	1,063,235	286,959	145,802	141,157	15,565	8,516	7,049	51,162	26,600	24,562
Under 1 year	24,586	13,702	10,884	15,954	8,871	7,083	7,401	4,116	3,285	354	213	141	877	502	375
1 - 4 years	4,316	2,460	1,856	3,015	1,718	1,297	1,041	595	446	93	55	38	167	92	75
5 - 9 years	2,330	1,325	1,005	1,691	957	734	494	283	211	46	29	17	99	56	43
10 - 14 years	2,949	1,729	1,220	2,150	1,265	885	651	384	267	62	40	22	86	40	46
15 - 19 years	10,887	7,866	3,021	7,847	5,549	2,298	2,531	1,965	566	242	176	66	267	176	91
20 - 24 years	18,664	13,924	4,740	13,662	10,112	3,550	4,144	3,164	980	373	280	93	485	368	117
25 - 29 years	20,263	14,429	5,834	15,097	10,817	4,280	4,246	2,997	1,249	376	246	130	544	369	175
30 - 34 years	21,996	14,763	7,233	16,328	11,066	5,262	4,674	3,074	1,600	419	279	140	575	344	231
35 - 39 years	28,012	17,614	10,398	20,935	13,391	7,544	5,810	3,440	2,370	513	321	192	754	462	292
40 - 44 years	42,021	25,820	16,201	32,125	20,095	12,030	8,206	4,675	3,531	650	428	222	1,040	622	418
45 - 49 years	73,569	44,946	28,623	57,398	35,738	21,660	13,581	7,665	5,916	990	607	383	1,600	936	664
50 - 54 years	109,638	67,072	42,566	85,651	53,279	32,372	20,512	11,728	8,784	1,213	704	509	2,262	1,361	901
55 - 59 years	139,961	86,775	53,186	110,343	69,195	41,148	25,246	14,939	10,307	1,380	837	543	2,992	1,804	1,188
60 - 64 years	170,841	102,520	68,321	139,240	84,101	55,139	26,570	15,519	11,051	1,324	753	571	3,707	2,147	1,560
65 - 69 years	189,962	109,519	80,443	158,882	92,184	66,698	25,781	14,289	11,492	1,349	738	611	3,950	2,308	1,642
70 - 74 years	217,189	120,185	97,004	184,095	102,609	81,486	26,710	14,106	12,604	1,454	769	685	4,930	2,701	2,229
75 - 79 years	273,348	143,006	130,342	237,879	125,427	112,452	28,238	13,898	14,340	1,388	705	683	5,843	2,976	2,867
80 - 84 years	352,303	168,824	183,479	314,629	152,116	162,513	29,305	12,704	16,601	1,297	593	704	7,072	3,411	3,661
85 years and over	765,474	275,866	489,608	697,733	252,958	444,775	51,795	16,243	35,552	2,042	743	1,299	13,904	5,922	7,982
Not stated	126	87	39	95	66	29	23	18	5	-	-	-	8	3	5
	Rate														
All ages ⁴	799.5	812.0	787.4	861.7	866.1	857.3	682.2	725.4	642.7	365.1	397.5	332.4	301.1	327.0	277.3
Under 1 year ⁵	623.4	680.2	564.0	537.2	584.3	488.0	1,102.1	1,206.5	994.4	455.3	542.5	366.4	389.3	434.4	341.8
1 - 4 years	26.5	29.6	23.3	24.6	27.4	21.6	38.1	42.9	33.2	29.4	34.3	24.4	17.9	19.3	16.3
5 - 9 years	11.5	12.8	10.1	10.9	12.1	9.7	15.0	16.9	13.0	12.2	15.2	*	8.5	9.6	7.4
10 - 14 years	14.3	16.3	12.1	13.6	15.6	11.5	19.1	22.2	16.0	16.6	21.1	12.0	7.8	7.2	8.5
15 - 19 years	49.4	69.6	28.1	47.0	64.7	28.3	67.0	102.5	30.5	61.5	87.1	34.5	22.8	29.3	15.9
20 - 24 years	86.5	126.4	44.8	82.7	119.2	44.2	122.4	189.1	57.3	102.8	147.5	53.7	36.9	55.3	18.0
25 - 29 years	96.0	135.7	55.7	92.6	130.0	53.6	140.3	206.0	79.5	110.4	139.6	79.1	37.9	53.7	23.4
30 - 34 years	110.2	147.7	72.6	106.1	141.5	69.5	164.6	228.4	107.1	134.7	174.5	92.6	40.5	51.4	30.8
35 - 39 years	138.8	175.4	102.6	133.8	169.5	97.3	208.7	263.0	160.5	175.4	216.1	133.5	51.9	67.2	38.1
40 - 44 years	201.1	248.4	154.3	194.7	241.8	146.9	291.4	351.2	237.7	232.1	302.3	160.4	80.3	101.8	61.1
45 - 49 years	324.0	401.0	248.9	314.4	392.5	236.7	458.8	549.8	377.8	348.7	431.2	267.6	132.6	164.8	104.0
50 - 54 years	491.7	613.5	374.5	471.9	592.7	353.5	731.5	893.0	589.3	477.8	570.0	390.5	207.2	268.6	154.0
55 - 59 years	711.7	911.2	524.5	678.9	869.4	496.1	1,104.8	1,425.7	833.0	699.4	878.5	532.2	322.1	427.8	234.2
60 - 64 years	1,015.8	1,269.2	781.7	982.4	1,222.1	756.2	1,523.4	1,977.9	1,151.8	892.0	1,047.5	745.9	492.9	632.2	378.1
65 - 69 years	1,527.6	1,871.3	1,222.0	1,495.8	1,825.2	1,197.3	2,148.8	2,745.1	1,691.9	1,377.8	1,591.5	1,185.6	765.6	982.5	584.3
70 - 74 years	2,340.9	2,831.9	1,926.9	2,315.5	2,792.1	1,905.9	3,041.6	3,862.0	2,457.3	2,202.4	2,555.9	1,906.4	1,285.7	1,555.4	1,062.5
75 - 79 years	3,735.4	4,493.7	3,151.9	3,734.9	4,472.1	3,154.8	4,450.5	5,677.3	3,679.8	3,221.2	3,840.9	2,761.3	2,155.2	2,598.0	1,831.3
80 - 84 years	6,134.1	7,358.2	5,319.8	6,171.5	7,379.2	5,351.7	6,710.4	8,414.7	5,809.8	4,811.0	5,489.7	4,357.2	3,895.1	4,791.6	3,316.8
85 years and over	13,934.3	15,414.3	13,219.2	14,147.6	15,640.3	13,419.3	13,187.2	14,715.3	12,589.9	9,615.3	10,268.1	9,277.9	9,418.1	10,824.5	8,590.1

- Quantity zero.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

¹Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2010, multiple-race data were reported by 37 states and the District of Columbia; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and other Asian or Pacific Islander groups.

⁴Figures for age not stated are included in "All ages" but are not distributed among age groups.

⁵Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births); see "Technical Notes."

Table 4. Number of deaths and death rates by Hispanic origin, race for non-Hispanic population, age, and sex: United States, 2010

[Rates per 100,000 population in specified group; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes"]

Age	All origins ¹			Hispanic			Non-Hispanic ²			Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
	Number														
All ages	2,468,435	1,232,432	1,236,003	144,490	79,622	64,868	2,318,218	1,149,438	1,168,780	1,969,916	971,604	998,312	283,438	143,824	139,614
Under 1 year	24,586	13,702	10,884	5,170	2,870	2,300	19,189	10,709	8,480	11,025	6,144	4,881	7,071	3,931	3,140
1 - 4 years	4,316	2,460	1,856	930	524	406	3,373	1,931	1,442	2,139	1,219	920	993	570	423
5 - 9 years	2,330	1,325	1,005	423	245	178	1,901	1,077	824	1,283	722	561	481	275	206
10 - 14 years	2,949	1,729	1,220	528	298	230	2,406	1,420	986	1,627	969	658	637	374	263
15 - 19 years	10,887	7,866	3,021	1,883	1,436	447	8,964	6,400	2,564	6,013	4,152	1,861	2,482	1,929	553
20 - 24 years	18,664	13,924	4,740	2,912	2,253	659	15,696	11,629	4,067	10,834	7,927	2,907	4,064	3,104	960
25 - 29 years	20,263	14,429	5,834	2,940	2,217	723	17,255	12,156	5,099	12,207	8,637	3,570	4,172	2,940	1,232
30 - 34 years	21,996	14,763	7,233	3,082	2,244	838	18,845	12,470	6,375	13,279	8,846	4,433	4,614	3,031	1,583
35 - 39 years	28,012	17,614	10,398	3,497	2,399	1,098	24,435	15,160	9,275	17,508	11,045	6,463	5,710	3,368	2,342
40 - 44 years	42,021	25,820	16,201	4,645	3,061	1,584	37,213	22,649	14,564	27,491	17,025	10,466	8,097	4,607	3,490
45 - 49 years	73,569	44,946	28,623	6,640	4,310	2,330	66,657	40,459	26,198	50,736	31,406	19,330	13,423	7,568	5,855
50 - 54 years	109,638	67,072	42,566	8,275	5,317	2,958	100,918	61,448	39,470	77,298	47,889	29,409	20,275	11,578	8,697
55 - 59 years	139,961	86,775	53,186	9,504	6,100	3,404	129,931	80,304	49,627	100,742	62,997	37,745	24,964	14,768	10,196
60 - 64 years	170,841	102,520	68,321	10,562	6,409	4,153	159,734	95,733	64,001	128,523	77,555	50,968	26,324	15,370	10,954
65 - 69 years	189,962	109,519	80,443	10,765	6,223	4,542	178,654	102,928	75,726	148,016	85,858	62,158	25,491	14,113	11,378
70 - 74 years	217,189	120,185	97,004	12,197	6,817	5,380	204,474	113,049	91,425	171,789	95,709	76,080	26,439	13,947	12,492
75 - 79 years	273,348	143,006	130,342	14,654	7,664	6,990	258,146	135,025	123,121	223,128	117,699	105,429	27,956	13,738	14,218
80 - 84 years	352,303	168,824	183,479	16,710	8,059	8,651	335,038	160,476	174,562	297,855	144,024	153,831	28,986	12,543	16,443
85 years and over	765,474	275,866	489,608	29,166	11,171	17,995	735,305	264,358	470,947	668,361	241,741	426,620	51,242	16,055	35,187
Not stated	126	87	39	7	5	2	84	57	27	62	40	22	17	15	2

Table 4. Number of deaths and death rates by Hispanic origin, race for non-Hispanic population, age, and sex: United States, 2010

[Rates per 100,000 population in specified group; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes"]

Age	All origins ¹			Hispanic			Non-Hispanic ²			Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
	Rate														
All ages ⁴	799.5	812.0	787.4	286.2	310.8	260.9	897.6	911.1	884.7	984.3	987.5	981.2	718.7	764.5	676.9
Under 1 year ⁵	623.4	680.2	564.0	510.7	556.8	462.9	654.5	714.5	591.8	529.3	575.9	480.4	1,170.4	1,281.5	1,055.7
1 - 4 years	26.5	29.6	23.3	22.7	25.0	20.2	27.7	31.1	24.3	24.7	27.5	21.8	40.2	45.4	34.8
5 - 9 years	11.5	12.8	10.1	8.8	10.0	7.6	12.2	13.6	10.8	11.4	12.5	10.2	15.9	17.9	13.9
10 - 14 years	14.3	16.3	12.1	11.7	12.9	10.4	14.9	17.2	12.5	13.8	16.0	11.5	20.2	23.4	17.0
15 - 19 years	49.4	69.6	28.1	41.5	61.2	20.4	51.2	71.4	30.0	47.5	63.9	30.1	70.6	108.0	31.9
20 - 24 years	86.5	126.4	44.8	67.4	97.9	32.6	90.9	133.5	47.6	85.4	123.2	46.5	129.5	200.2	60.5
25 - 29 years	96.0	135.7	55.7	68.2	97.4	35.5	102.8	145.4	60.5	98.0	137.6	57.8	149.2	219.0	84.7
30 - 34 years	110.2	147.7	72.6	74.7	104.7	42.3	119.0	158.8	79.8	113.6	150.1	76.5	175.4	243.4	114.3
35 - 39 years	138.8	175.4	102.6	90.7	121.7	58.2	149.7	187.8	112.4	144.0	180.8	106.9	218.7	274.7	169.2
40 - 44 years	201.1	248.4	154.3	134.9	173.6	94.4	213.3	262.4	165.2	205.5	253.9	156.9	304.2	366.4	248.5
45 - 49 years	324.0	401.0	248.9	219.7	282.6	155.6	338.6	417.8	261.9	327.0	406.9	247.9	475.4	569.2	392.0
50 - 54 years	491.7	613.5	374.5	338.9	439.1	240.4	508.2	632.0	389.5	485.2	607.1	365.7	753.0	918.4	607.4
55 - 59 years	711.7	911.2	524.5	516.1	686.1	357.5	729.0	930.0	540.1	691.4	881.6	508.4	1,132.8	1,462.7	853.9
60 - 64 years	1,015.8	1,269.2	781.7	769.6	992.8	571.4	1,034.2	1,288.1	798.7	995.3	1,233.3	769.4	1,561.9	2,027.9	1,181.1
65 - 69 years	1,527.6	1,871.3	1,222.0	1,134.9	1,450.9	874.0	1,555.3	1,897.8	1,249.0	1,518.9	1,844.8	1,221.1	2,195.3	2,802.4	1,730.3
70 - 74 years	2,340.9	2,831.9	1,926.9	1,742.1	2,229.5	1,364.2	2,383.7	2,870.6	1,970.4	2,353.0	2,822.4	1,945.9	3,110.7	3,947.8	2,515.2
75 - 79 years	3,735.4	4,493.7	3,151.9	2,868.8	3,592.4	2,349.8	3,792.4	4,547.8	3,208.0	3,786.8	4,518.2	3,207.2	4,542.9	5,789.4	3,760.6
80 - 84 years	6,134.1	7,358.2	5,319.8	4,754.1	5,795.9	4,072.2	6,213.8	7,445.5	5,393.5	6,246.1	7,459.8	5,420.4	6,834.3	8,566.4	5,921.1
85 years and over	13,934.3	15,414.3	13,219.2	10,777.9	11,779.8	10,237.3	14,078.7	15,597.8	13,348.9	14,286.1	15,816.6	13,543.5	13,385.7	14,974.2	12,767.7

¹Figures for origin not stated are included in "All origins" but are not distributed among specified origins.

²Includes races other than white and black.

³Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2010, multiple-race data were reported by 37 states and the District of Columbia; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴Figures for age not stated are included in "All ages" but are not distributed among age groups.

⁵Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births); see "Technical Notes."

Table 6. Abridged life table for the total population, 2010

[For explanation of the columns of the life table, see "United States Life Tables, 2008," *National Vital Statistics Reports*, Volume 61, Number 3]

Age	Probability of dying between ages x to $x + n$ ${}_nq_x$	Number surviving to age x l_x	Number dying between ages x to $x + n$ ${}_nd_x$	Person-years lived between ages x to $x + n$ ${}_nL_x$	Total number of person-years lived above age x T_x	Expectancy of life at age x e_x
0-1	0.006123	100,000	612	99,465	7,866,031	78.7
1-5	0.001071	99,388	106	397,294	7,766,566	78.1
5-10	0.000573	99,281	57	496,250	7,369,272	74.2
10-15	0.000708	99,224	70	495,989	6,873,022	69.3
15-20	0.002463	99,154	244	495,240	6,377,033	64.3
20-25	0.004317	98,910	427	493,529	5,881,793	59.5
25-30	0.004791	98,483	472	491,249	5,388,265	54.7
30-35	0.005500	98,011	539	488,743	4,897,016	50.0
35-40	0.006913	97,472	674	485,752	4,408,273	45.2
40-45	0.009979	96,798	966	481,756	3,922,521	40.5
45-50	0.016044	95,832	1,538	475,582	3,440,766	35.9
50-55	0.024343	94,295	2,295	466,064	2,965,184	31.4
55-60	0.035106	91,999	3,230	452,346	2,499,119	27.2
60-65	0.049847	88,770	4,425	433,346	2,046,773	23.1
65-70	0.074412	84,345	6,276	406,910	1,613,427	19.1
70-75	0.112312	78,068	8,768	369,609	1,206,518	15.5
75-80	0.174772	69,300	12,112	317,693	836,909	12.1
80-85	0.274365	57,189	15,691	248,042	519,216	9.1
85-90	0.430887	41,498	17,881	162,716	271,174	6.5
90-95	0.615150	23,617	14,528	79,221	108,458	4.6
95-100	0.783137	9,089	7,118	24,685	29,237	3.2
100 and over	1.000000	1,971	1,971	4,552	4,552	2.3

Table 7. Life expectancy at selected ages by race, Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race categories are consistent with the 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race.; see "Technical Notes"]

Exact age in years	All races and origins ¹			White ²			Black ²			Hispanic			Non-Hispanic white ²			Non-Hispanic black ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
0	78.7	76.2	81.0	78.9	76.5	81.3	75.1	71.8	78.0	81.2	78.5	83.8	78.8	76.4	81.1	74.7	71.4	77.7
1	78.1	75.7	80.5	78.4	76.0	80.7	75.0	71.8	77.8	80.7	78.0	83.2	78.2	75.8	80.5	74.6	71.3	77.5
5	74.2	71.8	76.6	74.4	72.1	76.7	71.1	67.9	73.9	76.7	74.0	79.2	74.3	71.9	76.6	70.7	67.5	73.6
10	69.3	66.8	71.6	69.5	67.1	71.8	66.1	62.9	69.0	71.8	69.1	74.3	69.3	67.0	71.6	65.8	62.5	68.7
15	64.3	61.9	66.6	64.5	62.1	66.8	61.2	58.0	64.0	66.8	64.1	69.3	64.4	62.0	66.6	60.8	57.6	63.7
20	59.5	57.1	61.7	59.7	57.3	61.9	56.4	53.3	59.1	62.0	59.3	64.4	59.5	57.2	61.7	56.0	52.9	58.8
25	54.7	52.4	56.9	54.9	52.7	57.0	51.7	48.8	54.3	57.1	54.6	59.5	54.7	52.5	56.9	51.4	48.4	54.0
30	50.0	47.8	52.0	50.1	48.0	52.2	47.1	44.3	49.5	52.3	49.8	54.6	50.0	47.9	52.0	46.7	43.9	49.2
35	45.2	43.1	47.2	45.4	43.3	47.4	42.4	39.7	44.7	47.5	45.1	49.7	45.3	43.2	47.2	42.1	39.4	44.5
40	40.5	38.5	42.4	40.7	38.7	42.6	37.8	35.2	40.1	42.7	40.4	44.8	40.6	38.6	42.5	37.6	34.9	39.8
45	35.9	33.9	37.7	36.0	34.1	37.9	33.4	30.8	35.5	38.0	35.7	40.0	36.0	34.0	37.8	33.1	30.5	35.3
50	31.4	29.6	33.2	31.6	29.7	33.3	29.1	26.6	31.1	33.5	31.2	35.3	31.5	29.7	33.2	28.8	26.3	31.0
55	27.2	25.4	28.8	27.3	25.5	28.8	25.1	22.7	27.0	29.0	26.9	30.8	27.2	25.5	28.8	24.9	22.5	26.8
60	23.1	21.5	24.5	23.1	21.6	24.5	21.3	19.2	23.0	24.7	22.8	26.3	23.1	21.5	24.4	21.2	19.0	22.9
65	19.1	17.7	20.3	19.2	17.8	20.3	17.8	15.9	19.3	20.6	18.8	22.0	19.1	17.7	20.3	17.7	15.8	19.1
70	15.5	14.2	16.5	15.5	14.2	16.4	14.6	12.9	15.8	16.8	15.1	18.0	15.4	14.2	16.4	14.5	12.8	15.7
75	12.1	11.0	12.9	12.1	11.0	12.8	11.6	10.2	12.5	13.2	11.7	14.1	12.0	11.0	12.8	11.6	10.1	12.5
80	9.1	8.2	9.7	9.0	8.2	9.6	9.0	7.8	9.6	9.9	8.7	10.7	9.0	8.1	9.6	8.9	7.8	9.6
85	6.5	5.8	6.9	6.5	5.8	6.9	6.8	5.9	7.1	7.1	6.1	7.7	6.5	5.8	6.9	6.7	5.9	7.1
90	4.6	4.1	4.8	4.5	4.0	4.8	5.0	4.4	5.2	5.0	4.2	5.4	4.5	4.0	4.8	5.0	4.4	5.2
95	3.2	2.9	3.3	3.2	2.8	3.3	3.7	3.3	3.8	3.5	2.9	3.7	3.2	2.8	3.3	3.8	3.3	3.8
100	2.3	2.1	2.3	2.3	2.0	2.3	2.8	2.5	2.8	2.4	2.1	2.6	2.3	2.1	2.3	2.9	2.6	2.8

¹ Includes races other than white and black.

²Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

Table 8. Life expectancy at birth by race, Hispanic origin, race for non-Hispanic population, and sex: United States, 1940, 1950, 1960, 1970 and 1975-2010

[Race categories are consistent with the 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race; see "Technical Notes"]

Year	All races and origins ¹			White			Black			Hispanic			Non-Hispanic white			Non-Hispanic black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
2010 ^{2,3}	78.7	76.2	81.0	78.9	76.5	81.3	75.1	71.8	78.0	81.2	78.5	83.8	78.8	76.4	81.1	74.7	71.4	77.7
2009 ^{2,3,4}	78.5	76.0	80.9	78.8	76.4	81.2	74.7	71.4	77.7	81.1	78.4	83.5	78.7	76.3	81.1	74.3	70.9	77.4
2008 ^{2,3,4}	78.2	75.6	80.6	78.5	76.1	80.9	74.3	70.9	77.3	80.8	78.0	83.3	78.4	76.0	80.7	73.9	70.5	77.0
2007 ^{2,3,4}	78.1	75.5	80.6	78.5	76.0	80.9	73.8	70.3	77.0	80.7	77.8	83.2	78.4	75.9	80.8	73.5	69.9	76.7
2006 ^{2,3,4}	77.8	75.2	80.3	78.3	75.8	80.7	73.4	69.9	76.7	80.3	77.5	82.9	78.2	75.7	80.6	73.1	69.5	76.4
2005 ^{2,3,4}	77.6	75.0	80.1	78.0	75.5	80.5	73.0	69.5	76.2	---	---	---	---	---	---	---	---	---
2004 ^{2,3,4}	77.6	75.0	80.1	78.1	75.5	80.5	72.9	69.4	76.1	---	---	---	---	---	---	---	---	---
2003 ^{2,3,4}	77.2	74.5	79.7	77.7	75.1	80.2	72.4	68.9	75.7	---	---	---	---	---	---	---	---	---
2002 ^{2,4}	77.0	74.4	79.6	77.5	74.9	80.1	72.2	68.7	75.4	---	---	---	---	---	---	---	---	---
2001 ^{2,4}	77.0	74.3	79.5	77.5	74.9	80.0	72.0	68.5	75.3	---	---	---	---	---	---	---	---	---
2000	76.8	74.1	79.3	77.3	74.7	79.9	71.8	68.2	75.1	---	---	---	---	---	---	---	---	---
1999	76.7	73.9	79.4	77.3	74.6	79.9	71.4	67.8	74.7	---	---	---	---	---	---	---	---	---
1998	76.7	73.8	79.5	77.3	74.5	80.0	71.3	67.6	74.8	---	---	---	---	---	---	---	---	---
1997	76.5	73.6	79.4	77.1	74.3	79.9	71.1	67.2	74.7	---	---	---	---	---	---	---	---	---
1996	76.1	73.1	79.1	76.8	73.9	79.7	70.2	66.1	74.2	---	---	---	---	---	---	---	---	---
1995	75.8	72.5	78.9	76.5	73.4	79.6	69.6	65.2	73.9	---	---	---	---	---	---	---	---	---
1994	75.7	72.4	79.0	76.5	73.3	79.6	69.5	64.9	73.9	---	---	---	---	---	---	---	---	---
1993	75.5	72.2	78.8	76.3	73.1	79.5	69.2	64.6	73.7	---	---	---	---	---	---	---	---	---
1992	75.8	72.3	79.1	76.5	73.2	79.8	69.6	65.0	73.9	---	---	---	---	---	---	---	---	---
1991	75.5	72.0	78.9	76.3	72.9	79.6	69.3	64.6	73.8	---	---	---	---	---	---	---	---	---
1990	75.4	71.8	78.8	76.1	72.7	79.4	69.1	64.5	73.6	---	---	---	---	---	---	---	---	---
1989	75.1	71.7	78.5	75.9	72.5	79.2	68.8	64.3	73.3	---	---	---	---	---	---	---	---	---
1988	74.9	71.4	78.3	75.6	72.2	78.9	68.9	64.4	73.2	---	---	---	---	---	---	---	---	---
1987	74.9	71.4	78.3	75.6	72.1	78.9	69.1	64.7	73.4	---	---	---	---	---	---	---	---	---
1986	74.7	71.2	78.2	75.4	71.9	78.8	69.1	64.8	73.4	---	---	---	---	---	---	---	---	---
1985	74.7	71.1	78.2	75.3	71.8	78.7	69.3	65.0	73.4	---	---	---	---	---	---	---	---	---
1984	74.7	71.1	78.2	75.3	71.8	78.7	69.5	65.3	73.6	---	---	---	---	---	---	---	---	---
1983	74.6	71.0	78.1	75.2	71.6	78.7	69.4	65.2	73.5	---	---	---	---	---	---	---	---	---
1982	74.5	70.8	78.1	75.1	71.5	78.7	69.4	65.1	73.6	---	---	---	---	---	---	---	---	---
1981	74.1	70.4	77.8	74.8	71.1	78.4	68.9	64.5	73.2	---	---	---	---	---	---	---	---	---
1980	73.7	70.0	77.4	74.4	70.7	78.1	68.1	63.8	72.5	---	---	---	---	---	---	---	---	---
1979	73.9	70.0	77.8	74.6	70.8	78.4	68.5	64.0	72.9	---	---	---	---	---	---	---	---	---
1978	73.5	69.6	77.3	74.1	70.4	78.0	68.1	63.7	72.4	---	---	---	---	---	---	---	---	---
1977	73.3	69.5	77.2	74.0	70.2	77.9	67.7	63.4	72.0	---	---	---	---	---	---	---	---	---
1976	72.9	69.1	76.8	73.6	69.9	77.5	67.2	62.9	71.6	---	---	---	---	---	---	---	---	---
1975	72.6	68.8	76.6	73.4	69.5	77.3	66.8	62.4	71.3	---	---	---	---	---	---	---	---	---
1970	70.8	67.1	74.7	71.7	68.0	75.6	64.1	60.0	68.3	---	---	---	---	---	---	---	---	---
1960	69.7	66.6	73.1	70.6	67.4	74.1	---	---	---	---	---	---	---	---	---	---	---	---
1950	68.2	65.6	71.1	69.1	66.5	72.2	---	---	---	---	---	---	---	---	---	---	---	---
1940	62.9	60.8	65.2	64.2	62.1	66.6	---	---	---	---	---	---	---	---	---	---	---	---

--- Data not available.

¹ Includes races other than white and black.

²Life table data for 2001-2010 are based on revised life table methodology; see "Technical Notes."

³Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010, by 34 states and the District of Columbia in 2009 and 2008, by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2005, by 15 states in 2004, and by 7 states in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴Life table data for 2001-2009 have been re-estimated using new 2001-2009 intercensal population estimates and may differ from data previously published; see "Technical Notes."

Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2010: United States, 1999-2010

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and are estimated as of July 1 for all other years; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004) and year	Age												Age-adjusted rate ³
	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	
All causes													
2010	799.5	623.4	26.5	12.9	67.7	102.9	170.5	407.1	851.9	1,875.1	4,790.2	13,934.3	747.0
2009 ⁴	794.5	659.7	27.4	13.8	69.8	104.4	180.0	418.1	856.7	1,888.7	4,820.2	13,660.1	749.6
2008 ⁴	812.9	678.9	29.3	13.9	74.2	105.1	181.0	419.6	867.1	1,958.4	4,998.1	14,332.4	774.9
2007 ⁴	804.6	702.5	29.4	15.2	78.8	107.2	186.0	420.3	866.7	1,976.0	4,987.1	14,160.9	775.3
2006 ⁴	813.1	705.8	29.1	15.2	81.4	109.0	192.0	427.5	881.3	2,031.4	5,096.1	14,426.7	791.8
2005 ⁴	828.4	710.2	29.9	16.3	80.7	106.8	194.9	431.9	898.5	2,109.7	5,251.8	14,982.4	815.0
2004 ⁴	818.8	695.9	30.3	16.7	79.7	104.1	194.9	426.8	903.2	2,141.0	5,267.4	14,777.6	813.7
2003 ⁴	843.9	704.9	31.8	16.9	81.1	105.2	202.6	433.1	937.3	2,235.0	5,451.3	15,401.4	843.5
2002 ⁴	849.5	709.5	31.4	17.4	80.9	105.1	204.2	431.0	948.7	2,300.3	5,543.8	15,589.5	855.9
2001 ⁴	848.0	687.0	33.4	17.2	80.2	105.6	203.5	426.7	972.5	2,344.2	5,573.7	15,432.6	858.8
2000	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4	869.0
1999	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6	875.6
Diseases of heart (I00-I09,I11,I13,I20-I51)													
2010	193.6	8.3	1.0	0.5	2.4	7.8	25.8	81.6	186.6	409.2	1,172.0	4,285.2	179.1
2009 ⁴	195.4	9.6	0.9	0.5	2.4	7.8	26.7	82.3	190.0	422.8	1,210.8	4,316.9	182.8
2008 ⁴	202.8	9.6	1.2	0.6	2.5	8.1	26.9	85.2	195.3	441.4	1,271.7	4,598.4	192.1
2007 ⁴	204.5	10.2	1.1	0.6	2.5	8.1	27.7	85.2	197.8	454.8	1,308.6	4,668.1	196.1
2006 ⁴	211.7	8.6	1.0	0.6	2.5	8.4	28.5	88.0	205.1	483.0	1,378.0	4,877.6	205.5
2005 ⁴	220.7	8.9	0.9	0.6	2.6	8.3	29.2	89.7	212.8	512.3	1,458.5	5,188.3	216.8
2004 ⁴	222.8	10.5	1.2	0.6	2.5	8.1	29.5	90.2	217.1	535.7	1,504.1	5,233.8	221.6
2003 ⁴	236.1	11.0	1.2	0.6	2.7	8.3	30.8	92.4	232.3	579.8	1,607.7	5,570.7	236.3
2002 ⁴	242.3	12.7	1.1	0.6	2.5	8.0	30.7	93.9	240.5	612.0	1,673.2	5,726.3	244.6
2001 ⁴	245.7	11.9	1.5	0.7	2.5	8.0	29.6	92.4	248.9	632.6	1,723.0	5,784.1	249.5
2000	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1	257.6
1999	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0	266.5
Malignant neoplasms (C00-C97)													
2010	186.2	1.6	2.1	2.2	3.7	8.8	28.8	111.6	300.1	666.1	1,202.2	1,729.5	172.8
2009 ⁴	185.0	1.8	2.2	2.2	3.8	9.0	30.2	112.8	301.7	668.2	1,213.0	1,699.3	173.5
2008 ⁴	186.0	1.7	2.4	2.2	3.8	8.8	30.1	113.4	304.7	688.4	1,230.9	1,724.6	176.4
2007 ⁴	186.9	1.7	2.3	2.4	3.8	8.7	31.0	114.2	311.4	702.9	1,250.1	1,739.4	179.3
2006 ⁴	187.6	1.9	2.4	2.2	3.8	9.3	32.2	116.3	317.7	716.3	1,259.2	1,748.3	181.8
2005 ⁴	189.3	1.9	2.4	2.5	4.0	9.2	33.5	118.6	323.9	733.2	1,272.8	1,778.2	185.1
2004 ⁴	189.2	1.8	2.5	2.5	4.1	9.3	33.6	119.0	330.8	746.8	1,278.6	1,767.4	186.8
2003 ⁴	192.0	1.9	2.5	2.6	4.0	9.5	35.1	122.1	341.6	763.5	1,299.7	1,792.3	190.9
2002 ⁴	193.7	1.9	2.6	2.6	4.2	9.8	36.0	124.1	349.7	787.2	1,308.8	1,812.4	194.3
2001 ⁴	194.3	1.6	2.7	2.4	4.2	10.1	36.8	125.8	359.4	799.7	1,313.7	1,802.9	196.5
2000	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4	199.6
1999	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8	200.8

Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2010: United States, 1999-2010

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and are estimated as of July 1 for all other years; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004) and year	Age												Age-adjusted rate ³
	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	
Chronic lower respiratory diseases (J40-J47)													
2010	44.7	0.9	0.3	0.3	0.3	0.7	1.7	9.9	39.0	146.3	369.9	690.7	42.2
2009 ⁴	44.8	0.7	0.4	0.3	0.4	0.7	1.8	10.4	40.0	147.5	376.4	684.9	42.7
2008 ⁴	46.4	0.8	0.3	0.3	0.4	0.6	1.9	9.9	41.1	155.9	395.4	722.7	44.7
2007 ⁴	42.5	1.0	0.4	0.3	0.3	0.7	1.9	9.5	38.6	145.5	367.1	652.0	41.4
2006 ⁴	41.8	0.7	0.3	0.3	0.4	0.6	1.9	9.1	38.8	147.0	362.0	641.3	41.0
2005 ⁴	44.3	0.8	0.4	0.3	0.3	0.7	2.0	9.4	41.6	158.4	385.0	691.9	43.9
2004 ⁴	41.7	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.1	152.1	366.2	643.2	41.6
2003 ⁴	43.6	0.8	0.4	0.3	0.5	0.7	2.2	8.7	43.1	161.7	382.2	670.2	43.7
2002 ⁴	43.4	1.0	0.4	0.3	0.5	0.8	2.3	8.7	42.2	162.0	385.8	670.3	43.9
2001 ⁴	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.4	44.5	167.3	379.3	658.3	43.9
2000	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6	44.2
1999	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0	45.4
Cerebrovascular diseases (I60-I69)													
2010	41.9	3.3	0.3	0.2	0.4	1.3	4.6	13.1	29.3	81.7	288.3	993.8	39.1
2009 ⁴	42.0	3.7	0.3	0.2	0.4	1.3	4.6	13.7	29.7	82.8	294.9	992.2	39.6
2008 ⁴	44.1	3.4	0.4	0.2	0.4	1.3	4.8	13.7	30.6	87.3	313.3	1,071.0	42.1
2007 ⁴	45.1	3.2	0.3	0.2	0.5	1.3	5.0	14.5	31.7	91.4	320.8	1,110.7	43.5
2006 ⁴	46.0	3.5	0.3	0.2	0.5	1.3	5.1	14.6	32.9	94.9	333.9	1,131.7	44.8
2005 ⁴	48.6	3.1	0.4	0.2	0.5	1.4	5.2	15.0	32.7	99.8	358.4	1,239.7	48.0
2004 ⁴	51.3	3.2	0.3	0.2	0.5	1.4	5.4	14.8	34.0	106.6	385.6	1,331.9	51.2
2003 ⁴	54.4	2.5	0.3	0.2	0.5	1.5	5.6	15.0	35.5	111.9	409.8	1,446.0	54.6
2002 ⁴	56.6	3.0	0.3	0.2	0.4	1.4	5.4	15.1	37.1	119.6	430.0	1,520.1	57.2
2001 ⁴	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.0	38.3	122.9	443.3	1,532.0	58.4
2000	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2	60.9
1999	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8	61.6
Accidents (unintentional injuries) (V01-X59, Y85-Y86)													
2010	39.1	28.1	8.6	4.0	28.3	35.5	36.0	43.7	38.4	43.3	106.1	328.4	38.0
2009 ⁴	38.5	29.5	9.0	4.1	28.6	34.5	36.4	44.5	36.5	42.1	103.5	310.9	37.5
2008 ⁴	40.1	31.8	9.1	4.6	32.5	36.3	38.1	45.8	37.4	43.9	105.7	318.3	39.2
2007 ⁴	41.1	31.0	9.9	5.4	36.8	37.7	39.6	46.2	36.8	44.4	105.0	313.6	40.4
2006 ⁴	40.8	28.4	10.1	5.6	37.9	38.0	40.5	45.5	35.8	43.8	104.7	299.2	40.2
2005 ⁴	39.9	27.0	10.5	5.9	37.1	35.7	38.9	43.2	35.4	45.7	106.0	303.5	39.5
2004 ⁴	38.3	26.2	10.4	6.5	36.8	33.2	37.6	40.7	32.9	43.5	103.6	295.8	38.1
2003 ⁴	37.7	23.8	11.0	6.4	36.9	32.0	38.0	38.8	32.7	43.7	101.6	294.3	37.6
2002 ⁴	37.1	23.9	10.6	6.6	37.7	31.9	37.4	36.7	31.3	44.0	101.1	289.6	37.1
2001 ⁴	35.6	24.3	11.2	6.9	35.8	30.0	35.4	33.9	30.5	42.6	100.7	282.2	35.7
2000	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5	34.9
1999	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4	35.3

Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2010: United States, 1999-2010

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and are estimated as of July 1 for all other years; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004) and year	Age												Age-adjusted rate ³
	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	
Alzheimer's disease (G30)													
2010	27.0	*	*	*	*	*	*	0.3	2.1	19.8	184.5	987.1	25.1
2009 ⁴	25.8	*	*	*	*	*	*	0.2	2.0	19.4	179.1	945.3	24.2
2008 ⁴	27.1	*	*	*	*	*	*	0.2	2.2	21.1	192.5	1,002.2	25.8
2007 ⁴	24.8	*	*	*	*	*	*	0.2	2.2	20.2	175.8	928.7	23.8
2006 ⁴	24.3	*	*	*	*	*	*	0.2	2.1	19.9	175.0	923.4	23.7
2005 ⁴	24.2	*	*	*	*	*	*	0.2	2.1	20.2	177.0	935.5	24.0
2004 ⁴	22.5	*	*	*	*	*	*	0.2	1.8	19.5	168.5	875.3	22.6
2003 ⁴	21.9	*	*	*	*	*	*	0.2	2.0	20.7	164.1	846.8	22.1
2002 ⁴	20.5	*	*	*	*	*	*	0.1	1.9	19.6	157.7	790.9	20.8
2001 ⁴	18.9	*	*	*	*	*	*	0.2	2.1	18.6	147.2	725.4	19.3
2000	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7	18.1
1999	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3	16.5
Diabetes mellitus (E10-E14)													
2010	22.4	*	*	0.1	0.4	1.5	4.4	12.5	32.0	67.6	144.1	285.5	20.8
2009 ⁴	22.4	*	*	0.1	0.4	1.5	4.5	12.8	32.1	69.6	145.8	282.6	21.0
2008 ⁴	23.2	*	*	0.1	0.5	1.4	4.4	12.6	33.3	74.7	153.2	298.9	22.0
2007 ⁴	23.7	*	*	0.1	0.4	1.5	4.6	13.1	34.1	76.7	161.9	302.2	22.8
2006 ⁴	24.3	*	*	0.1	0.4	1.7	4.8	13.1	35.8	80.6	166.2	310.4	23.6
2005 ⁴	25.4	*	*	0.1	0.5	1.6	4.7	13.4	36.9	85.7	177.0	338.8	24.9
2004 ⁴	25.0	*	*	0.1	0.4	1.5	4.6	13.4	36.8	86.2	176.6	328.2	24.7
2003 ⁴	25.6	*	*	0.1	0.4	1.7	4.6	13.9	38.3	90.0	180.7	335.1	25.5
2002 ⁴	25.5	*	*	0.1	0.4	1.6	4.8	13.7	37.5	90.9	182.4	337.0	25.6
2001 ⁴	25.0	*	*	0.1	0.4	1.5	4.3	13.6	38.1	91.0	181.1	328.6	25.4
2000	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7	25.0
1999	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2	25.0
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
2010	16.3	2.7	*	0.1	0.2	0.6	1.8	4.9	13.9	39.3	115.7	333.8	15.3
2009 ⁴	16.0	2.8	*	*	0.2	0.7	2.0	5.2	13.5	38.7	115.1	321.4	15.1
2008 ⁴	15.9	3.5	*	*	0.2	0.6	1.8	5.0	14.1	39.9	113.3	325.6	15.1
2007 ⁴	15.4	3.5	0.1	0.1	0.2	0.7	1.8	5.1	13.4	39.4	112.4	317.9	14.9
2006 ⁴	15.2	4.0	*	*	0.2	0.7	1.8	5.2	13.7	38.8	111.0	316.2	14.8
2005 ⁴	14.9	4.0	*	0.1	0.2	0.7	1.7	4.8	13.5	38.8	110.2	313.1	14.7
2004 ⁴	14.5	4.3	*	0.1	0.2	0.6	1.8	5.0	13.5	38.1	108.2	306.4	14.5
2003 ⁴	14.6	4.6	*	0.1	0.2	0.7	1.8	4.9	13.6	39.7	109.3	309.3	14.7
2002 ⁴	14.2	4.4	*	0.1	0.2	0.7	1.7	4.7	12.9	39.0	108.9	303.4	14.4
2001 ⁴	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.1	40.0	104.0	293.8	14.1
2000	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8	13.5
1999	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9	13.0

Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2010: United States, 1999-2010

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and are estimated as of July 1 for all other years; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004) and year	Age												Age-adjusted rate ³
	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	
Influenza and pneumonia (J09-J18)													
2010	16.2	4.9	0.6	0.2	0.4	0.9	1.9	4.3	9.9	27.9	102.4	426.2	15.1
2009 ⁴	17.5	6.3	0.9	0.6	1.0	2.0	3.2	6.5	11.7	29.5	107.0	433.8	16.5
2008 ⁴	18.5	5.5	0.9	0.2	0.5	0.9	2.1	5.1	10.9	30.5	118.6	512.3	17.6
2007 ⁴	17.5	5.4	0.7	0.3	0.4	0.8	1.8	4.3	9.5	28.2	113.5	506.7	16.8
2006 ⁴	18.9	6.5	0.8	0.2	0.4	0.9	1.9	4.6	9.9	31.6	127.3	547.0	18.4
2005 ⁴	21.3	6.6	0.7	0.3	0.4	0.9	2.1	5.1	11.2	35.1	142.0	644.9	21.0
2004 ⁴	20.4	6.8	0.8	0.2	0.4	0.8	2.0	4.6	10.8	34.2	139.1	622.8	20.4
2003 ⁴	22.5	8.1	1.0	0.4	0.5	1.0	2.2	5.2	11.2	36.9	150.8	703.0	22.6
2002 ⁴	22.8	6.7	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.2	156.6	732.4	23.2
2001 ⁴	21.8	7.5	0.7	0.2	0.5	0.9	2.2	4.6	10.8	36.2	148.3	700.1	22.2
2000	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1	23.7
1999	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8	23.5
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
2010	12.4	0.7	10.5	14.0	16.0	19.6	17.5	13.7	15.7	17.6	12.1
2009 ⁴	12.0	0.6	10.0	13.1	16.1	19.2	16.4	13.7	15.8	16.4	11.8
2008 ⁴	11.8	0.5	9.9	13.2	15.9	18.6	16.0	13.6	16.1	16.4	11.6
2007 ⁴	11.5	0.5	9.6	13.3	15.7	17.7	15.3	12.4	16.2	17.0	11.3
2006 ⁴	11.2	0.5	9.8	12.7	15.2	17.2	14.4	12.4	15.8	17.3	11.0
2005 ⁴	11.0	0.7	9.9	12.7	15.1	16.5	13.7	12.4	16.8	18.3	10.9
2004 ⁴	11.1	0.7	10.3	12.9	15.2	16.6	13.7	12.2	16.3	17.6	11.0
2003 ⁴	10.9	0.6	9.6	12.9	15.0	15.9	13.7	12.6	16.4	17.9	10.8
2002 ⁴	11.0	0.6	9.8	12.8	15.3	15.8	13.5	13.4	17.7	18.9	10.9
2001 ^{4,5}	10.7	0.7	9.9	12.8	14.7	15.1	13.2	13.2	17.4	17.8	10.7
2000	10.4	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6	10.4
1999	10.5	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3	10.5
Septicemia (A40-A41)													
2010	11.3	5.5	0.4	0.2	0.3	0.8	1.9	5.2	12.6	30.1	76.0	179.0	10.6
2009 ⁴	11.6	5.5	0.4	0.2	0.3	0.9	2.2	5.4	13.1	31.4	79.2	182.4	11.0
2008 ⁴	11.8	7.0	0.6	0.2	0.3	0.9	2.1	5.7	13.3	31.4	82.0	189.8	11.3
2007 ⁴	11.6	6.8	0.5	0.2	0.4	0.7	2.1	5.5	12.8	32.2	79.5	190.8	11.2
2006 ⁴	11.5	6.7	0.6	0.2	0.3	0.7	2.0	5.2	12.6	31.6	82.1	193.0	11.2
2005 ⁴	11.6	7.5	0.5	0.2	0.3	0.8	1.9	5.2	12.8	32.2	81.3	203.4	11.4
2004 ⁴	11.4	6.8	0.5	0.2	0.3	0.8	1.9	5.4	12.8	32.1	81.5	199.6	11.3
2003 ⁴	11.7	7.0	0.5	0.2	0.4	0.8	2.1	5.3	13.0	32.3	84.8	213.7	11.8
2002 ⁴	11.8	7.5	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.5	86.3	213.4	11.9
2001 ⁴	11.3	7.8	0.7	0.2	0.3	0.7	1.8	5.0	12.4	32.6	82.2	210.3	11.5
2000	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7	11.3
1999	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7	11.3

Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2010: United States, 1999-2010

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and are estimated as of July 1 for all other years; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004) and year	Age												Age-adjusted rate ³
	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	
Chronic liver disease and cirrhosis (K70,K73-K74)													
2010	10.3	*	*	*	0.1	1.2	5.9	19.2	26.8	26.3	27.7	21.8	9.4
2009 ⁴	10.0	*	*	*	0.1	1.1	6.0	18.7	25.9	25.4	27.2	21.1	9.1
2008 ⁴	9.9	*	*	*	0.1	1.1	6.1	18.5	25.0	26.3	28.0	21.9	9.2
2007 ⁴	9.7	*	*	*	0.1	1.0	6.0	18.7	24.2	26.2	28.2	21.7	9.1
2006 ⁴	9.2	*	*	*	0.1	0.8	5.9	17.8	22.6	25.6	28.9	21.1	8.8
2005 ⁴	9.3	*	*	*	0.1	0.8	6.2	17.7	23.3	26.8	28.9	21.3	8.9
2004 ⁴	9.2	*	*	*		0.8	6.4	18.0	22.4	27.4	28.7	21.1	9.0
2003 ⁴	9.5	*	*	*	*	0.9	6.8	18.3	22.9	29.2	29.9	21.2	9.3
2002 ⁴	9.5	*	*	*	0.1	1.0	7.1	18.0	22.8	29.3	31.3	22.5	9.4
2001 ⁴	9.5	*	*	*	0.1	1.0	7.4	18.4	22.9	29.8	30.2	22.7	9.5
2000	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1	9.5
1999	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2	9.6
Essential hypertension and hypertensive renal disease (I10,I12,I15)													
2010	8.6	*	*	*	0.0	0.3	1.0	3.1	7.3	16.7	51.8	212.0	8.0
2009 ⁴	8.4	*	*	*	0.1	0.3	1.0	3.1	7.1	16.3	51.0	208.0	7.8
2008 ⁴	8.5	*	*	*	0.1	0.3	1.0	3.0	7.2	16.5	51.9	215.3	8.0
2007 ⁴	8.0	*	*	*	0.1	0.2	0.9	2.8	6.4	15.9	49.2	209.1	7.6
2006 ⁴	8.0	*	*	*	0.0	0.3	0.9	3.0	6.8	16.5	50.8	206.1	7.7
2005 ⁴	8.4	*	*	*	0.1	0.2	0.9	2.7	6.4	17.5	55.5	228.0	8.3
2004 ⁴	7.9	*	*	*	0.1	0.3	0.8	2.7	6.3	16.9	52.5	212.2	7.9
2003 ⁴	7.6	*	*	*	0.1	0.2	0.8	2.5	6.3	16.8	51.6	199.4	7.6
2002 ⁴	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	15.9	48.1	189.6	7.1
2001 ⁴	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.4	47.6	175.6	6.9
2000	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9	6.5
1999	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1	6.2
Parkinson's disease (G20-G21)													
2010	7.1	*	*	*	*	*	*	0.2	1.3	11.8	74.8	165.9	6.8
2009 ⁴	6.7	*	*	*	*	*	*	0.2	1.3	11.2	70.8	157.0	6.5
2008 ⁴	6.7	*	*	*	*	*	*	0.2	1.2	12.3	71.2	157.4	6.6
2007 ⁴	6.7	*	*	*	*	*	*	0.1	1.2	11.7	71.5	157.0	6.5
2006 ⁴	6.6	*	*	*	*	*	*	0.2	1.2	12.0	69.5	157.6	6.5
2005 ⁴	6.6	*	*	*	*	*	*	0.2	1.4	12.8	71.1	156.0	6.6
2004 ⁴	6.1	*	*	*	*	*	*	0.2	1.2	11.9	67.4	145.1	6.2
2003 ⁴	6.2	*	*	*	*	*	*	0.2	1.3	12.6	67.6	145.8	6.3
2002 ⁴	5.9	*	*	*	*	*	*	0.1	1.2	12.1	63.8	142.2	6.0
2001 ⁴	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.5	137.0	5.9
2000	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9	5.7
1999	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4	5.4

Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2010: United States, 1999-2010

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for census years and are estimated as of July 1 for all other years; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004) and year	Age											Age-adjusted rate ³	
	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years		85 years and over
Pneumonitis due to solids and liquids (J69)													
2010	5.5	*	*	*	0.1	0.2	0.3	1.1	2.8	8.6	38.2	152.3	5.1
2009 ⁴	5.2	*	*	*	0.1	0.2	0.4	1.1	2.8	7.7	35.7	146.7	4.9
2008 ⁴	5.5	*	*	*	0.1	0.2	0.4	1.1	2.7	8.2	38.5	157.8	5.2
2007 ⁴	5.6	*	*	*	0.1	0.2	0.4	1.0	2.7	8.8	39.6	167.7	5.4
2006 ⁴	5.7	*	*	*	0.1	0.2	0.4	1.0	2.7	9.1	40.4	169.6	5.5
2005 ⁴	5.8	*	*	*	0.1	0.2	0.4	1.1	2.7	9.2	42.5	178.0	5.8
2004 ⁴	5.7	*	*	*	0.1	0.2	0.4	0.9	2.5	9.5	42.8	176.3	5.7
2003 ⁴	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.5	44.9	186.0	6.0
2002 ⁴	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.2	195.5	6.2
2001 ⁴	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.7	193.4	6.2
2000	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6	6.1
1999	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6	5.6

* Figure does not meet standards of reliability or precision, see "Technical Notes."

... Category not applicable.

¹Figures for age not stated included in "All ages" but not distributed among age groups.

²Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births); see "Technical Notes."

³For method of computation, see "Technical Notes."

⁴Rates are revised and may differ from those previously published; see "Technical Notes."

⁵Figures include September 11, 2001 related deaths for which death certificates were filed as of October 24, 2002; see "Technical Notes" from *Deaths: Final Data for 2001*.

Table 10. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All causes	2,468,435	24,586	4,316	5,279	29,551	42,259	70,033	183,207	310,802	407,151	625,651	765,474	126
Salmonella infections (A01-A02)	28	1	-	-	-	-	1	4	4	3	10	5	-
Shigellosis and amebiasis (A03,A06)	3	-	-	-	1	1	-	-	-	-	1	-	-
Certain other intestinal infections (A04,A07-A09)	10,276	322	20	12	16	26	61	264	635	1,400	3,267	4,253	-
Tuberculosis (A16-A19)	569	-	1	3	6	14	26	66	107	111	126	108	1
Respiratory tuberculosis (A16)	423	-	-	2	5	10	16	47	80	83	92	88	-
Other tuberculosis (A17-A19)	146	-	1	1	1	4	10	19	27	28	34	20	1
Whooping cough (A37)	26	25	-	-	-	-	-	-	1	-	-	-	-
Scarlet fever and erysipelas (A38,A46)	3	-	-	-	1	-	-	-	1	-	-	1	-
Meningococcal infection (A39)	79	11	10	4	16	6	8	9	7	1	4	3	-
Septicemia (A40-A41)	34,812	215	62	67	141	312	767	2,333	4,604	6,545	9,931	9,834	1
Syphilis (A50-A53)	28	2	-	-	-	-	1	-	7	8	6	4	-
Acute poliomyelitis (A80)	-	-	-	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)	9	-	1	1	-	-	-	3	1	2	-	1	-
Measles (B05)	2	-	-	-	-	1	-	1	-	-	-	-	-
Viral hepatitis (B15-B19)	7,564	1	-	1	8	51	342	2,376	3,218	879	510	178	-
Human immunodeficiency virus (HIV) disease (B20-B24)	8,369	-	2	3	147	741	1,898	3,123	1,822	486	124	23	-
Malaria (B50-B54)	10	-	-	-	-	3	-	-	4	2	1	-	-
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	5,805	119	67	48	69	115	216	460	956	1,149	1,444	1,162	-
Malignant neoplasms (C00-C97)	574,743	62	346	916	1,604	3,619	11,809	50,211	109,501	144,635	157,025	95,010	5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	8,474	-	1	1	31	43	231	1,157	2,214	2,007	1,721	1,068	-
Malignant neoplasm of esophagus (C15)	14,490	-	-	-	2	32	222	1,483	3,683	3,953	3,517	1,598	-
Malignant neoplasm of stomach (C16)	11,390	-	-	1	22	136	433	1,180	2,075	2,590	2,901	2,052	-
Malignant neoplasms of colon, rectum and anus (C18-C21)	52,622	-	-	4	54	312	1,345	5,052	9,453	11,459	13,846	11,096	1
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	20,305	4	19	20	31	90	315	2,493	5,819	4,814	4,569	2,130	1
Malignant neoplasm of pancreas (C25)	36,888	-	-	3	8	64	468	2,869	7,382	9,658	10,452	5,984	-
Malignant neoplasm of larynx (C32)	3,691	-	-	-	1	6	41	413	955	1,068	855	352	-
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	158,318	-	-	7	31	162	1,351	12,093	31,147	48,606	46,658	18,262	1
Malignant melanoma of skin (C43)	9,154	-	-	2	40	169	447	1,115	1,911	2,063	2,165	1,242	-
Malignant neoplasm of breast (C50)	41,435	-	-	1	7	323	2,045	5,915	9,104	8,710	8,385	6,945	-
Malignant neoplasm of cervix uteri (C53)	3,939	-	-	-	12	191	531	989	916	606	438	256	-
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	8,402	-	-	-	3	31	166	679	1,892	2,323	2,035	1,273	-
Malignant neoplasm of ovary (C56)	14,572	-	-	2	22	97	345	1,510	3,104	3,724	3,722	2,046	-
Malignant neoplasm of prostate (C61)	28,561	-	-	-	2	1	20	493	2,555	5,866	10,135	9,488	1
Malignant neoplasms of kidney and renal pelvis (C64-C65)	13,219	-	16	30	30	41	246	1,262	2,778	3,418	3,348	2,050	-
Malignant neoplasm of bladder (C67)	14,731	-	-	-	-	17	90	602	1,731	3,019	5,027	4,244	1

Table 10. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	14,164	13	101	296	206	366	782	1,987	3,347	3,305	2,711	1,050	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	55,590	28	115	286	533	771	1,261	3,439	8,002	12,789	17,069	11,297	-
Hodgkin's disease (C81)	1,231	-	-	3	48	119	122	124	203	220	259	133	-
Non-Hodgkin's lymphoma (C82-C85)	20,294	2	3	30	103	204	429	1,316	2,912	4,621	6,223	4,451	-
Leukemia (C91-C95)	22,569	25	112	253	376	439	608	1,377	2,973	4,914	6,883	4,609	-
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	11,428	-	-	-	4	8	101	619	1,898	3,027	3,686	2,085	-
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	68	1	-	-	2	1	1	3	16	7	18	19	-
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	64,798	17	94	263	569	767	1,470	5,480	11,433	14,657	17,471	12,577	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14,917	48	59	82	93	163	279	751	1,438	2,580	4,710	4,714	-
Anemias (D50-D64)	4,852	15	29	26	109	142	178	264	369	524	1,091	2,104	1
Diabetes mellitus (E10-E14)	69,071	3	4	26	165	606	1,789	5,610	11,677	14,687	18,822	15,682	-
Nutritional deficiencies (E40-E64)	2,948	3	6	2	7	18	37	106	240	380	803	1,346	-
Malnutrition (E40-E46)	2,790	2	3	-	7	17	36	98	231	366	772	1,258	-
Other nutritional deficiencies (E50-E64)	158	1	3	2	-	1	1	8	9	14	31	88	-
Meningitis (G00,G03)	608	58	22	11	22	33	53	97	115	67	83	47	-
Parkinson's disease (G20-G21)	22,032	-	-	-	3	5	7	80	489	2,567	9,769	9,112	-
Alzheimer's disease (G30)	83,494	-	-	-	-	-	7	121	750	4,291	24,099	54,226	-
Major cardiovascular diseases (I00-I78)	780,213	468	213	282	1,286	4,012	13,313	45,149	83,886	114,483	205,180	311,910	31
Diseases of heart (I00-I09,I11,I13,I20-I51)	597,689	329	159	185	1,028	3,222	10,594	36,729	68,077	88,851	153,080	235,407	28
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	2,987	1	2	3	12	25	56	163	342	534	852	997	-
Hypertensive heart disease (I11)	33,678	-	-	2	47	371	1,358	4,097	5,369	4,567	6,330	11,529	8
Hypertensive heart and renal disease (I13)	2,807	-	-	-	1	19	84	193	311	330	702	1,167	-
Ischemic heart diseases (I20-I25)	379,559	20	5	9	113	1,023	5,330	23,008	46,291	60,934	100,056	142,754	16
Acute myocardial infarction (I21-I22)	122,071	14	2	4	61	353	2,053	8,862	17,680	22,219	32,058	38,763	2
Other acute ischemic heart diseases (I24)	4,170	2	-	-	4	21	94	383	657	709	995	1,305	-
Other forms of chronic ischemic heart disease (I20,I25)	253,318	4	3	5	48	649	3,183	13,763	27,954	38,006	67,003	102,686	14
Atherosclerotic cardiovascular disease, so described (I25.0)	57,438	-	-	-	12	257	1,345	5,949	10,711	9,917	12,458	16,781	8
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	195,880	4	3	5	36	392	1,838	7,814	17,243	28,089	54,545	85,905	6
Other heart diseases (I26-I51)	178,658	308	152	171	855	1,784	3,766	9,268	15,764	22,486	45,140	78,960	4
Acute and subacute endocarditis (I33)	1,103	1	2	-	15	37	61	149	217	235	256	130	-
Diseases of pericardium and acute myocarditis (I30-I31,I40)	776	25	18	7	36	48	58	95	120	97	147	125	-
Heart failure (I50)	57,757	26	13	9	42	107	298	1,116	2,912	5,864	15,094	32,274	2

Table 10. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	119,022	256	119	155	762	1,592	3,349	7,908	12,515	16,290	29,643	46,431	2
Essential hypertension and hypertensive renal disease (I10,I12,I15)	26,634	3	1	2	21	110	394	1,384	2,672	3,636	6,763	11,648	-
Cerebrovascular diseases (I60-I69)	129,476	130	50	90	190	517	1,904	5,910	10,693	17,736	37,659	54,595	2
Atherosclerosis (I70)	7,230	2	-	-	2	6	23	126	366	741	1,802	4,162	-
Other diseases of circulatory system (I71-I78)	19,184	4	3	5	45	157	398	1,000	2,078	3,519	5,876	6,098	1
Aortic aneurysm and dissection (I71)	10,431	-	-	3	38	110	311	621	1,219	2,049	3,390	2,690	-
Other diseases of arteries, arterioles and capillaries (I72-I78)	8,753	4	3	2	7	47	87	379	859	1,470	2,486	3,408	1
Other disorders of circulatory system (I80-I99)	4,241	39	3	2	48	140	327	634	687	600	822	938	1
Influenza and pneumonia (J09-J18)	50,097	195	91	71	181	385	773	1,926	3,627	6,066	13,369	23,411	2
Influenza (J09-J11)	500	16	13	12	31	44	53	103	84	39	44	61	-
Pneumonia (J12-J18)	49,597	179	78	59	150	341	720	1,823	3,543	6,027	13,325	23,350	2
Other acute lower respiratory infections (J20-J22,U04)	213	28	12	4	3	2	10	21	17	14	31	71	-
Acute bronchitis and bronchiolitis (J20-J21)	177	27	12	4	3	2	10	19	15	12	21	52	-
Other and unspecified acute lower respiratory infections (J22,U04)	36	1	-	-	-	-	-	2	2	2	10	19	-
Chronic lower respiratory diseases (J40-J47)	138,080	37	51	133	149	272	709	4,452	14,242	31,777	48,309	37,945	4
Bronchitis, chronic and unspecified (J40-J42)	620	25	16	5	6	8	8	22	52	92	127	259	-
Emphysema (J43)	10,034	1	-	1	2	9	44	411	1,264	2,650	3,556	2,095	1
Asthma (J45-J46)	3,404	6	31	119	132	210	303	571	513	374	498	647	-
Other chronic lower respiratory diseases (J44,J47)	124,022	5	4	8	9	45	354	3,448	12,413	28,661	44,128	34,944	3
Pneumoconioses and chemical effects (J60-J66,J68)	845	-	1	-	-	1	2	19	47	155	364	256	-
Pneumonitis due to solids and liquids (J69)	17,011	18	13	9	42	69	142	487	1,005	1,869	4,990	8,367	-
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	31,187	296	114	68	134	239	468	1,455	3,268	6,267	10,067	8,811	-
Peptic ulcer (K25-K28)	2,977	-	-	1	4	23	58	241	420	495	809	925	1
Diseases of appendix (K35-K38)	415	7	2	6	9	8	13	35	57	68	101	109	-
Hernia (K40-K46)	1,832	27	5	3	7	4	30	112	191	274	496	683	-
Chronic liver disease and cirrhosis (K70,K73-K74)	31,903	3	2	3	34	487	2,423	8,651	9,764	5,720	3,620	1,195	1
Alcoholic liver disease (K70)	15,990	-	-	-	24	378	1,769	5,465	5,257	2,209	762	125	1
Other chronic liver disease and cirrhosis (K73-K74)	15,913	3	2	3	10	109	654	3,186	4,507	3,511	2,858	1,070	-
Cholelithiasis and other disorders of gallbladder (K80-K82)	3,332	-	-	-	7	15	51	102	288	505	942	1,421	1
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,476	105	13	22	68	243	726	2,222	5,082	8,541	15,118	18,335	1
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	203	5	3	2	-	5	8	10	13	32	67	58	-
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	5,894	-	-	4	3	17	48	154	367	681	1,652	2,968	-
Renal failure (N17-N19)	44,362	100	10	16	65	220	668	2,057	4,698	7,824	13,395	15,308	1
Other disorders of kidney (N25,N27)	17	-	-	-	-	1	2	1	4	4	4	1	-
Infections of kidney (N10-N12,N13.6,N15.1)	608	7	2	2	9	6	20	60	63	97	145	197	-

Table 10. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hyperplasia of prostate (N40)	489	-	-	-	-	-	1	3	13	44	140	288	-
Inflammatory diseases of female pelvic organs (N70-N76)	137	-	-	-	3	2	6	10	17	26	40	33	-
Pregnancy, childbirth and the puerperium (O00-O99)	825	1	163	367	223	64	6	1	-	-	-
Pregnancy with abortive outcome (O00-O07)	37	-	7	16	13	1	-	-	-	-	-
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	788	1	156	351	210	63	6	1	-	-	-
Certain conditions originating in the perinatal period (P00-P96)	12,128	12,008	52	25	13	9	11	4	2	2	1	-	1
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9,673	5,107	507	298	412	397	449	680	774	396	351	302	-
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	38,360	3,052	243	100	510	888	1,225	2,204	2,548	3,105	6,827	17,631	27
All other diseases (Residual)	269,844	753	482	779	1,872	3,297	6,576	16,727	26,186	32,793	65,182	115,192	5
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120,859	1,110	1,394	1,643	12,341	14,573	14,792	19,667	14,023	9,407	13,853	18,040	16
Transport accidents (V01-V99,Y85)	37,961	81	469	956	7,549	6,090	5,124	5,955	4,806	2,942	2,620	1,366	3
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)	35,332	79	449	890	7,250	5,746	4,745	5,392	4,335	2,676	2,458	1,309	3
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)	1,029	2	17	30	154	134	149	208	156	89	63	27	-
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	1,600	-	3	36	145	210	230	355	315	177	99	30	-
Nontransport accidents (W00-X59,Y86)	82,898	1,029	925	687	4,792	8,483	9,668	13,712	9,217	6,465	11,233	16,674	13
Falls (W00-W19)	26,009	10	24	28	211	299	493	1,283	2,011	2,988	7,249	11,412	1
Accidental discharge of firearms (W32-W34)	606	-	25	37	145	107	91	89	50	38	19	5	-
Accidental drowning and submersion (W65-W74)	3,782	39	436	251	656	476	409	578	417	276	171	70	3
Accidental exposure to smoke, fire and flames (X00-X09)	2,782	21	147	135	127	155	231	446	468	398	400	253	1
Accidental poisoning and exposure to noxious substances (X40-X49)	33,041	6	34	54	3,183	6,767	7,476	9,662	4,451	837	338	227	6
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)	16,678	953	259	182	470	679	968	1,654	1,820	1,928	3,056	4,707	2
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,364	274	4,600	5,735	6,571	8,799	6,384	2,974	2,052	968	7
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	19,392	81	2,046	2,594	2,914	4,092	3,387	2,053	1,544	679	2
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)	18,972	193	2,554	3,141	3,657	4,707	2,997	921	508	289	5
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16,259	311	385	261	4,678	4,258	2,473	1,997	1,065	452	250	112	17

Table 10. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	11,078	11	43	165	3,889	3,331	1,673	1,097	533	207	90	34	5
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	5,181	300	342	96	789	927	800	900	532	245	160	78	12
Legal intervention (Y35,Y89.0)	412	-	1	1	74	102	108	70	46	9	1	-	-
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	4,908	108	82	67	456	806	932	1,295	726	195	136	102	3
Discharge of firearms, undetermined intent (Y22-Y24)	252	-	3	14	53	54	26	44	33	10	10	5	-
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	4,656	108	79	53	403	752	906	1,251	693	185	126	97	3
Operations of war and their sequelae (Y36,Y89.1)	9	-	-	-	-	2	-	-	4	-	-	3	-
Complications of medical and surgical care (Y40-Y84,Y88)	2,490	22	19	22	40	61	122	242	418	499	629	416	-
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ¹	7,298	2	-	4	6	6	27	149	426	1,063	2,462	3,153	-
Drug-induced deaths ^{2,3}	40,393	23	43	63	3,667	7,864	8,923	11,935	5,911	1,137	520	303	4
Alcohol-induced deaths ^{2,4}	25,692	1	-	-	152	899	3,076	8,612	7,986	3,420	1,250	291	5
Injury by firearms ^{2,5}	31,672	11	71	298	6,201	6,172	4,790	5,380	4,039	2,316	1,664	723	7

- Quantity zero.

... Category not applicable.

¹Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

²Included in selected categories above.

³Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁴Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁵Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 11. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes	799.5	623.4	26.5	12.9	67.7	102.9	170.5	407.1	851.9	1,875.1	4,790.2	13,934.3
Salmonella infections (A01-A02)	0.0	*	*	*	*	*	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	3.3	8.2	0.1	*	*	0.1	0.1	0.6	1.7	6.4	25.0	77.4
Tuberculosis (A16-A19)	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.0	2.0
Respiratory tuberculosis (A16)	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6
Other tuberculosis (A17-A19)	0.0	*	*	*	*	*	*	*	0.1	0.1	0.3	0.4
Whooping cough (A37)	0.0	0.6	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	*	*	*	*	*	*	*	*	*	*	*
Septicemia (A40-A41)	11.3	5.5	0.4	0.2	0.3	0.8	1.9	5.2	12.6	30.1	76.0	179.0
Syphilis (A50-A53)	0.0	*	*	*	*	*	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.4	*	*	*	*	0.1	0.8	5.3	8.8	4.0	3.9	3.2
Human immunodeficiency virus (HIV) disease (B20-B24)	2.7	*	*	*	0.3	1.8	4.6	6.9	5.0	2.2	0.9	0.4
Malaria (B50-B54)	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.9	3.0	0.4	0.1	0.2	0.3	0.5	1.0	2.6	5.3	11.1	21.2
Malignant neoplasms (C00-C97)	186.2	1.6	2.1	2.2	3.7	8.8	28.8	111.6	300.1	666.1	1,202.2	1,729.5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	2.7	*	*	*	0.1	0.1	0.6	2.6	6.1	9.2	13.2	19.4
Malignant neoplasm of esophagus (C15)	4.7	*	*	*	*	0.1	0.5	3.3	10.1	18.2	26.9	29.1
Malignant neoplasm of stomach (C16)	3.7	*	*	*	0.1	0.3	1.1	2.6	5.7	11.9	22.2	37.4
Malignant neoplasms of colon, rectum and anus (C18-C21)	17.0	*	*	*	0.1	0.8	3.3	11.2	25.9	52.8	106.0	202.0
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	6.6	*	*	0.0	0.1	0.2	0.8	5.5	16.0	22.2	35.0	38.8
Malignant neoplasm of pancreas (C25)	11.9	*	*	*	*	0.2	1.1	6.4	20.2	44.5	80.0	108.9
Malignant neoplasm of larynx (C32)	1.2	*	*	*	*	*	0.1	0.9	2.6	4.9	6.5	6.4
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	51.3	*	*	*	0.1	0.4	3.3	26.9	85.4	223.9	357.2	332.4
Malignant melanoma of skin (C43)	3.0	*	*	*	0.1	0.4	1.1	2.5	5.2	9.5	16.6	22.6
Malignant neoplasm of breast (C50)	13.4	*	*	*	*	0.8	5.0	13.1	25.0	40.1	64.2	126.4
Malignant neoplasm of cervix uteri (C53)	1.3	*	*	*	*	0.5	1.3	2.2	2.5	2.8	3.4	4.7
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	2.7	*	*	*	*	0.1	0.4	1.5	5.2	10.7	15.6	23.2
Malignant neoplasm of ovary (C56)	4.7	*	*	*	0.1	0.2	0.8	3.4	8.5	17.2	28.5	37.2
Malignant neoplasm of prostate (C61)	9.3	*	*	*	*	*	0.0	1.1	7.0	27.0	77.6	172.7
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.3	*	*	0.1	0.1	0.1	0.6	2.8	7.6	15.7	25.6	37.3

Table 11. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)	4.8	*	*	*	*	*	0.2	1.3	4.7	13.9	38.5	77.3
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	4.6	*	0.6	0.7	0.5	0.9	1.9	4.4	9.2	15.2	20.8	19.1
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	18.0	0.7	0.7	0.7	1.2	1.9	3.1	7.6	21.9	58.9	130.7	205.6
Hodgkin's disease (C81)	0.4	*	*	*	0.1	0.3	0.3	0.3	0.6	1.0	2.0	2.4
Non-Hodgkin's lymphoma (C82-C85)	6.6	*	*	0.1	0.2	0.5	1.0	2.9	8.0	21.3	47.6	81.0
Leukemia (C91-C95)	7.3	0.6	0.7	0.6	0.9	1.1	1.5	3.1	8.1	22.6	52.7	83.9
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.7	*	*	*	*	*	0.2	1.4	5.2	13.9	28.2	38.0
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	*	*	*	*	*	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	21.0	*	0.6	0.6	1.3	1.9	3.6	12.2	31.3	67.5	133.8	228.9
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.8	1.2	0.4	0.2	0.2	0.4	0.7	1.7	3.9	11.9	36.1	85.8
Anemias (D50-D64)	1.6	*	0.2	0.1	0.2	0.3	0.4	0.6	1.0	2.4	8.4	38.3
Diabetes mellitus (E10-E14)	22.4	*	*	0.1	0.4	1.5	4.4	12.5	32.0	67.6	144.1	285.5
Nutritional deficiencies (E40-E64)	1.0	*	*	*	*	*	0.1	0.2	0.7	1.8	6.1	24.5
Malnutrition (E40-E46)	0.9	*	*	*	*	*	0.1	0.2	0.6	1.7	5.9	22.9
Other nutritional deficiencies (E50-E64)	0.1	*	*	*	*	*	*	*	*	*	0.2	1.6
Meningitis (G00,G03)	0.2	1.5	0.1	*	0.1	0.1	0.1	0.2	0.3	0.3	0.6	0.9
Parkinson's disease (G20-G21)	7.1	*	*	*	*	*	*	0.2	1.3	11.8	74.8	165.9
Alzheimer's disease (G30)	27.0	*	*	*	*	*	*	0.3	2.1	19.8	184.5	987.1
Major cardiovascular diseases (I00-I78)	252.7	11.9	1.3	0.7	2.9	9.8	32.4	100.3	229.9	527.2	1,570.9	5,677.9
Diseases of heart (I00-I09,I11,I13,I20-I51)	193.6	8.3	1.0	0.5	2.4	7.8	25.8	81.6	186.6	409.2	1,172.0	4,285.2
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	1.0	*	*	*	*	0.1	0.1	0.4	0.9	2.5	6.5	18.1
Hypertensive heart disease (I11)	10.9	*	*	*	0.1	0.9	3.3	9.1	14.7	21.0	48.5	209.9
Hypertensive heart and renal disease (I13)	0.9	*	*	*	*	*	0.2	0.4	0.9	1.5	5.4	21.2
Ischemic heart diseases (I20-I25)	122.9	0.5	*	*	0.3	2.5	13.0	51.1	126.9	280.6	766.1	2,598.6
Acute myocardial infarction (I21-I22)	39.5	*	*	*	0.1	0.9	5.0	19.7	48.5	102.3	245.4	705.6
Other acute ischemic heart diseases (I24)	1.4	*	*	*	*	0.1	0.2	0.9	1.8	3.3	7.6	23.8
Other forms of chronic ischemic heart disease (I20,I25)	82.0	*	*	*	0.1	1.6	7.8	30.6	76.6	175.0	513.0	1,869.3
Atherosclerotic cardiovascular disease, so described (I25.0)	18.6	*	*	*	*	0.6	3.3	13.2	29.4	45.7	95.4	305.5
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	63.4	*	*	*	0.1	1.0	4.5	17.4	47.3	129.4	417.6	1,563.8
Other heart diseases (I26-I51)	57.9	7.8	0.9	0.4	2.0	4.3	9.2	20.6	43.2	103.6	345.6	1,437.4
Acute and subacute endocarditis (I33)	0.4	*	*	*	*	0.1	0.1	0.3	0.6	1.1	2.0	2.4
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.3	0.6	*	*	0.1	0.1	0.1	0.2	0.3	0.4	1.1	2.3
Heart failure (I50)	18.7	0.7	*	*	0.1	0.3	0.7	2.5	8.0	27.0	115.6	587.5

Table 11. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	38.6	6.5	0.7	0.4	1.7	3.9	8.2	17.6	34.3	75.0	227.0	845.2
Essential hypertension and hypertensive renal disease (I10,I12,I15)	8.6	*	*	*	0.0	0.3	1.0	3.1	7.3	16.7	51.8	212.0
Cerebrovascular diseases (I60-I69)	41.9	3.3	0.3	0.2	0.4	1.3	4.6	13.1	29.3	81.7	288.3	993.8
Atherosclerosis (I70)	2.3	*	*	*	*	*	0.1	0.3	1.0	3.4	13.8	75.8
Other diseases of circulatory system (I71-I78)	6.2	*	*	*	0.1	0.4	1.0	2.2	5.7	16.2	45.0	111.0
Aortic aneurysm and dissection (I71)	3.4	*	*	*	0.1	0.3	0.8	1.4	3.3	9.4	26.0	49.0
Other diseases of arteries, arterioles and capillaries (I72-I78)	2.8	*	*	*	*	0.1	0.2	0.8	2.4	6.8	19.0	62.0
Other disorders of circulatory system (I80-I99)	1.4	1.0	*	*	0.1	0.3	0.8	1.4	1.9	2.8	6.3	17.1
Influenza and pneumonia (J09-J18)	16.2	4.9	0.6	0.2	0.4	0.9	1.9	4.3	9.9	27.9	102.4	426.2
Influenza (J09-J11)	0.2	*	*	*	0.1	0.1	0.1	0.2	0.2	0.2	0.3	1.1
Pneumonia (J12-J18)	16.1	4.5	0.5	0.1	0.3	0.8	1.8	4.1	9.7	27.8	102.0	425.1
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.7	*	*	*	*	*	0.0	*	*	0.2	1.3
Acute bronchitis and bronchiolitis (J20-J21)	0.1	0.7	*	*	*	*	*	*	*	*	0.2	0.9
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	*	*	*	*	*	*	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	44.7	0.9	0.3	0.3	0.3	0.7	1.7	9.9	39.0	146.3	369.9	690.7
Bronchitis, chronic and unspecified (J40-J42)	0.2	0.6	*	*	*	*	*	0.0	0.1	0.4	1.0	4.7
Emphysema (J43)	3.2	*	*	*	*	*	0.1	0.9	3.5	12.2	27.2	38.1
Asthma (J45-J46)	1.1	*	0.2	0.3	0.3	0.5	0.7	1.3	1.4	1.7	3.8	11.8
Other chronic lower respiratory diseases (J44,J47)	40.2	*	*	*	*	0.1	0.9	7.7	34.0	132.0	337.9	636.1
Pneumoconioses and chemical effects (J60-J66,J68)	0.3	*	*	*	*	*	*	*	0.1	0.7	2.8	4.7
Pneumonitis due to solids and liquids (J69)	5.5	*	*	*	0.1	0.2	0.3	1.1	2.8	8.6	38.2	152.3
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	10.1	7.5	0.7	0.2	0.3	0.6	1.1	3.2	9.0	28.9	77.1	160.4
Peptic ulcer (K25-K28)	1.0	*	*	*	*	0.1	0.1	0.5	1.2	2.3	6.2	16.8
Diseases of appendix (K35-K38)	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.8	2.0
Hernia (K40-K46)	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.3	3.8	12.4
Chronic liver disease and cirrhosis (K70,K73-K74)	10.3	*	*	*	0.1	1.2	5.9	19.2	26.8	26.3	27.7	21.8
Alcoholic liver disease (K70)	5.2	*	*	*	0.1	0.9	4.3	12.1	14.4	10.2	5.8	2.3
Other chronic liver disease and cirrhosis (K73-K74)	5.2	*	*	*	*	0.3	1.6	7.1	12.4	16.2	21.9	19.5
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.1	*	*	*	*	*	0.1	0.2	0.8	2.3	7.2	25.9
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16.3	2.7	*	0.1	0.2	0.6	1.8	4.9	13.9	39.3	115.7	333.8
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.1	*	*	*	*	*	*	*	*	0.1	0.5	1.1
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.9	*	*	*	*	*	0.1	0.3	1.0	3.1	12.6	54.0
Renal failure (N17-N19)	14.4	2.5	*	*	0.1	0.5	1.6	4.6	12.9	36.0	102.6	278.7
Other disorders of kidney (N25,N27)	*	*	*	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	*	*	*	*	*	0.0	0.1	0.2	0.4	1.1	3.6

Table 11. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.2
Inflammatory diseases of female pelvic organs (N70-N76)	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.6
Pregnancy, childbirth and the puerperium (O00-O99)	0.3	*	0.4	0.9	0.5	0.1	*	*	*	*
Pregnancy with abortive outcome (O00-O07)	0.0	*	*	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.3	*	0.4	0.9	0.5	0.1	*	*	*	*
Certain conditions originating in the perinatal period (P00-P96)	3.9	304.5	0.3	0.1	*	*	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.1	129.5	3.1	0.7	0.9	1.0	1.1	1.5	2.1	1.8	2.7	5.5
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	12.4	77.4	1.5	0.2	1.2	2.2	3.0	4.9	7.0	14.3	52.3	320.9
All other diseases (Residual)	87.4	19.1	3.0	1.9	4.3	8.0	16.0	37.2	71.8	151.0	499.1	2,096.9
Accidents (unintentional injuries) (V01-X59, Y85-Y86)	39.1	28.1	8.6	4.0	28.3	35.5	36.0	43.7	38.4	43.3	106.1	328.4
Transport accidents (V01-V99, Y85)	12.3	2.1	2.9	2.3	17.3	14.8	12.5	13.2	13.2	13.5	20.1	24.9
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	11.4	2.0	2.8	2.2	16.6	14.0	11.6	12.0	11.9	12.3	18.8	23.8
Other land transport accidents (V01, V05-V06, V09.1, V09.3-V09.9, V10-V11, V15-V18, V19.3, V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9)	0.3	*	*	0.1	0.4	0.3	0.4	0.5	0.4	0.4	0.5	0.5
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99, Y85)	0.5	*	*	0.1	0.3	0.5	0.6	0.8	0.9	0.8	0.8	0.5
Nontransport accidents (W00-X59, Y86)	26.8	26.1	5.7	1.7	11.0	20.7	23.5	30.5	25.3	29.8	86.0	303.5
Falls (W00-W19)	8.4	*	0.1	0.1	0.5	0.7	1.2	2.9	5.5	13.8	55.5	207.7
Accidental discharge of firearms (W32-W34)	0.2	*	0.2	0.1	0.3	0.3	0.2	0.2	0.1	0.2	*	*
Accidental drowning and submersion (W65-W74)	1.2	1.0	2.7	0.6	1.5	1.2	1.0	1.3	1.1	1.3	1.3	1.3
Accidental exposure to smoke, fire and flames (X00-X09)	0.9	0.5	0.9	0.3	0.3	0.4	0.6	1.0	1.3	1.8	3.1	4.6
Accidental poisoning and exposure to noxious substances (X40-X49)	10.7	*	0.2	0.1	7.3	16.5	18.2	21.5	12.2	3.9	2.6	4.1
Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64, W75-W99, X10-X39, X50-X59, Y86)	5.4	24.2	1.6	0.4	1.1	1.7	2.4	3.7	5.0	8.9	23.4	85.7
Intentional self-harm (suicide) (*U03, X60-X84, Y87.0)	12.4	0.7	10.5	14.0	16.0	19.6	17.5	13.7	15.7	17.6
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	6.3	0.2	4.7	6.3	7.1	9.1	9.3	9.5	11.8	12.4
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60-X71, X75-X84, Y87.0)	6.1	0.5	5.9	7.6	8.9	10.5	8.2	4.2	3.9	5.3

Table 11. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by age: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5.3	7.9	2.4	0.6	10.7	10.4	6.0	4.4	2.9	2.1	1.9	2.0
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	3.6	*	0.3	0.4	8.9	8.1	4.1	2.4	1.5	1.0	0.7	0.6
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	1.7	7.6	2.1	0.2	1.8	2.3	1.9	2.0	1.5	1.1	1.2	1.4
Legal intervention (Y35,Y89.0)	0.1	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.6	2.7	0.5	0.2	1.0	2.0	2.3	2.9	2.0	0.9	1.0	1.9
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.5	2.7	0.5	0.1	0.9	1.8	2.2	2.8	1.9	0.9	1.0	1.8
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.8	0.6	*	0.1	0.1	0.1	0.3	0.5	1.1	2.3	4.8	7.6
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ³	2.4	*	*	*	*	*	0.1	0.3	1.2	4.9	18.8	57.4
Drug-induced deaths ^{4,5}	13.1	0.6	0.3	0.2	8.4	19.2	21.7	26.5	16.2	5.2	4.0	5.5
Alcohol-induced deaths ^{4,6}	8.3	*	*	*	0.3	2.2	7.5	19.1	21.9	15.8	9.6	5.3
Injury by firearms ^{4,7}	10.3	*	0.4	0.7	14.2	15.0	11.7	12.0	11.1	10.7	12.7	13.2

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Figures for age not stated included in "All ages" but not distributed among age groups.

²Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births); see "Technical Notes."

³Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁴Included in selected categories above.

⁵Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁶Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	2,468,435	1,232,432	1,236,003	2,114,749	1,051,514	1,063,235	286,959	145,802	141,157
Salmonella infections (A01-A02)	28	17	11	19	14	5	5	1	4
Shigellosis and amebiasis (A03,A06)	3	1	2	1	1	-	1	-	1
Certain other intestinal infections (A04,A07-A09)	10,276	4,046	6,230	9,248	3,582	5,666	827	370	457
Tuberculosis (A16-A19)	569	351	218	338	200	138	120	86	34
Respiratory tuberculosis (A16)	423	276	147	239	146	93	94	74	20
Other tuberculosis (A17-A19)	146	75	71	99	54	45	26	12	14
Whooping cough (A37)	26	13	13	24	11	13	2	2	-
Scarlet fever and erysipelas (A38,A46)	3	1	2	2	1	1	-	-	-
Meningococcal infection (A39)	79	40	39	67	34	33	9	5	4
Septicemia (A40-A41)	34,812	16,069	18,743	27,985	12,976	15,009	6,001	2,691	3,310
Syphilis (A50-A53)	28	20	8	15	12	3	12	8	4
Acute poliomyelitis (A80)	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)	9	4	5	8	3	5	-	-	-
Measles (B05)	2	2	-	1	1	-	-	-	-
Viral hepatitis (B15-B19)	7,564	5,038	2,526	6,061	4,097	1,964	1,101	707	394
Human immunodeficiency virus (HIV) disease (B20-B24)	8,369	6,099	2,270	3,575	2,951	624	4,662	3,047	1,615
Malaria (B50-B54)	10	7	3	2	1	1	6	5	1
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	5,805	2,970	2,835	4,802	2,439	2,363	777	407	370
Malignant neoplasms (C00-C97)	574,743	301,037	273,706	491,686	258,272	233,414	65,930	33,967	31,963
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	8,474	5,815	2,659	7,151	4,885	2,266	1,006	727	279
Malignant neoplasm of esophagus (C15)	14,490	11,416	3,074	12,761	10,148	2,613	1,437	1,044	393
Malignant neoplasm of stomach (C16)	11,390	6,703	4,687	8,524	5,023	3,501	2,000	1,180	820
Malignant neoplasms of colon, rectum and anus (C18-C21)	52,622	27,284	25,338	43,854	22,765	21,089	7,005	3,618	3,387
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	20,305	13,658	6,647	15,949	10,627	5,322	2,826	1,986	840
Malignant neoplasm of pancreas (C25)	36,888	18,699	18,189	31,413	16,064	15,349	4,327	2,096	2,231
Malignant neoplasm of larynx (C32)	3,691	2,951	740	3,007	2,410	597	613	480	133
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	158,318	87,740	70,578	137,698	75,675	62,023	16,688	9,793	6,895
Malignant melanoma of skin (C43)	9,154	6,002	3,152	8,944	5,905	3,039	135	62	73

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of breast (C50)	41,435	439	40,996	34,183	349	33,834	6,109	78	6,031
Malignant neoplasm of cervix uteri (C53)	3,939	...	3,939	2,970	...	2,970	790	...	790
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	8,402	...	8,402	6,688	...	6,688	1,437	...	1,437
Malignant neoplasm of ovary (C56)	14,572	...	14,572	12,841	...	12,841	1,279	...	1,279
Malignant neoplasm of prostate (C61)	28,561	28,561	...	23,172	23,172	...	4,854	4,854	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	13,219	8,436	4,783	11,659	7,474	4,185	1,185	708	477
Malignant neoplasm of bladder (C67)	14,731	10,429	4,302	13,404	9,647	3,757	1,073	615	458
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	14,164	7,977	6,187	12,876	7,305	5,571	896	466	430
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	55,590	30,777	24,813	48,584	27,084	21,500	5,450	2,824	2,626
Hodgkin's disease (C81)	1,231	714	517	1,087	633	454	114	60	54
Non-Hodgkin's lymphoma (C82-C85)	20,294	11,047	9,247	18,198	9,907	8,291	1,509	825	684
Leukemia (C91-C95)	22,569	12,851	9,718	20,088	11,524	8,564	1,816	957	859
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	11,428	6,117	5,311	9,154	4,979	4,175	2,003	978	1,025
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	68	48	20	57	41	16	8	4	4
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	64,798	34,150	30,648	56,008	29,739	26,269	6,820	3,436	3,384
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14,917	7,721	7,196	13,287	6,966	6,321	1,220	550	670
Anemias (D50-D64)	4,852	2,017	2,835	3,741	1,525	2,216	1,007	446	561
Diabetes mellitus (E10-E14)	69,071	35,490	33,581	54,250	28,486	25,764	12,126	5,640	6,486
Nutritional deficiencies (E40-E64)	2,948	1,141	1,807	2,503	974	1,529	367	134	233
Malnutrition (E40-E46)	2,790	1,079	1,711	2,358	919	1,439	357	129	228
Other nutritional deficiencies (E50-E64)	158	62	96	145	55	90	10	5	5
Meningitis (G00,G03)	608	303	305	442	221	221	138	64	74
Parkinson's disease (G20-G21)	22,032	12,871	9,161	20,792	12,173	8,619	804	452	352
Alzheimer's disease (G30)	83,494	25,364	58,130	76,928	23,442	53,486	5,220	1,488	3,732
Major cardiovascular diseases (I00-I78)	780,213	383,547	396,666	667,081	327,381	339,700	92,898	45,572	47,326
Diseases of heart (I00-I09,I11,I13,I20-I51)	597,689	307,384	290,305	514,323	264,425	249,898	69,083	35,089	33,994
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	2,987	996	1,991	2,644	885	1,759	232	79	153

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart disease (I11)	33,678	16,242	17,436	25,097	11,726	13,371	7,729	4,075	3,654
Hypertensive heart and renal disease (I13)	2,807	1,285	1,522	1,960	864	1,096	748	377	371
Ischemic heart diseases (I20-I25)	379,559	207,580	171,979	330,277	181,386	148,891	39,630	20,615	19,015
Acute myocardial infarction (I21-I22)	122,071	67,435	54,636	106,204	59,181	47,023	12,743	6,445	6,298
Other acute ischemic heart diseases (I24)	4,170	2,156	2,014	3,519	1,798	1,721	557	303	254
Other forms of chronic ischemic heart disease (I20,I25)	253,318	137,989	115,329	220,554	120,407	100,147	26,330	13,867	12,463
Atherosclerotic cardiovascular disease, so described (I25.0)	57,438	33,703	23,735	47,470	27,790	19,680	8,361	4,930	3,431
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	195,880	104,286	91,594	173,084	92,617	80,467	17,969	8,937	9,032
Other heart diseases (I26-I51)	178,658	81,281	97,377	154,345	69,564	84,781	20,744	9,943	10,801
Acute and subacute endocarditis (I33)	1,103	627	476	889	507	382	183	100	83
Diseases of pericardium and acute myocarditis (I30-I31,I40)	776	405	371	617	320	297	126	64	62
Heart failure (I50)	57,757	24,385	33,372	51,290	21,540	29,750	5,528	2,444	3,084
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	119,022	55,864	63,158	101,549	47,197	54,352	14,907	7,335	7,572
Essential hypertension and hypertensive renal disease (I10,I12,I15)	26,634	10,846	15,788	20,560	8,229	12,331	5,116	2,218	2,898
Cerebrovascular diseases (I60-I69)	129,476	52,367	77,109	109,119	43,424	65,695	15,965	6,938	9,027
Atherosclerosis (I70)	7,230	2,933	4,297	6,513	2,618	3,895	594	254	340
Other diseases of circulatory system (I71-I78)	19,184	10,017	9,167	16,566	8,685	7,881	2,140	1,073	1,067
Aortic aneurysm and dissection (I71)	10,431	6,096	4,335	9,134	5,359	3,775	972	552	420
Other diseases of arteries, arterioles and capillaries (I72-I78)	8,753	3,921	4,832	7,432	3,326	4,106	1,168	521	647
Other disorders of circulatory system (I80-I99)	4,241	2,053	2,188	3,377	1,616	1,761	763	383	380
Influenza and pneumonia (J09-J18)	50,097	23,615	26,482	43,296	20,238	23,058	4,936	2,380	2,556
Influenza (J09-J11)	500	250	250	409	205	204	68	32	36
Pneumonia (J12-J18)	49,597	23,365	26,232	42,887	20,033	22,854	4,868	2,348	2,520
Other acute lower respiratory infections (J20-J22,U04)	213	94	119	182	80	102	22	10	12
Acute bronchitis and bronchiolitis (J20-J21)	177	74	103	147	61	86	21	9	12
Other and unspecified acute lower respiratory infections (J22,U04)	36	20	16	35	19	16	1	1	-
Chronic lower respiratory diseases (J40-J47)	138,080	65,423	72,657	127,176	59,632	67,544	8,715	4,532	4,183
Bronchitis, chronic and unspecified (J40-J42)	620	267	353	557	232	325	48	26	22
Emphysema (J43)	10,034	5,227	4,807	9,294	4,764	4,530	596	363	233
Asthma (J45-J46)	3,404	1,283	2,121	2,366	790	1,576	879	408	471

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other chronic lower respiratory diseases (J44,J47)	124,022	58,646	65,376	114,959	53,846	61,113	7,192	3,735	3,457
Pneumoconioses and chemical effects (J60-J66,J68)	845	812	33	803	773	30	36	33	3
Pneumonitis due to solids and liquids (J69)	17,011	9,208	7,803	15,235	8,264	6,971	1,380	724	656
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	31,187	15,791	15,396	27,376	13,968	13,408	2,926	1,378	1,548
Peptic ulcer (K25-K28)	2,977	1,460	1,517	2,572	1,233	1,339	296	172	124
Diseases of appendix (K35-K38)	415	258	157	355	216	139	44	30	14
Hernia (K40-K46)	1,832	814	1,018	1,635	714	921	166	82	84
Chronic liver disease and cirrhosis (K70,K73-K74)	31,903	20,798	11,105	28,014	18,352	9,662	2,635	1,715	920
Alcoholic liver disease (K70)	15,990	11,441	4,549	13,997	10,125	3,872	1,226	822	404
Other chronic liver disease and cirrhosis (K73-K74)	15,913	9,357	6,556	14,017	8,227	5,790	1,409	893	516
Cholelithiasis and other disorders of gallbladder (K80-K82)	3,332	1,552	1,780	2,912	1,368	1,544	300	121	179
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,476	24,865	25,611	40,205	20,172	20,033	8,841	4,016	4,825
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	203	97	106	161	75	86	32	18	14
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	5,894	2,775	3,119	4,770	2,270	2,500	948	432	516
Renal failure (N17-N19)	44,362	21,989	22,373	35,261	17,824	17,437	7,857	3,565	4,292
Other disorders of kidney (N25,N27)	17	4	13	13	3	10	4	1	3
Infections of kidney (N10-N12,N13.6,N15.1)	608	187	421	526	158	368	65	26	39
Hyperplasia of prostate (N40)	489	489	...	444	444	...	37	37	...
Inflammatory diseases of female pelvic organs (N70-N76)	137	...	137	113	...	113	21	...	21
Pregnancy, childbirth and the puerperium (O00-O99)	825	...	825	504	...	504	264	...	264
Pregnancy with abortive outcome (O00-O07)	37	...	37	19	...	19	18	...	18
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	788	...	788	485	...	485	246	...	246
Certain conditions originating in the perinatal period (P00-P96)	12,128	6,803	5,325	7,513	4,233	3,280	4,031	2,216	1,815
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9,673	4,960	4,713	7,566	3,860	3,706	1,656	872	784
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	38,360	16,057	22,303	32,668	13,321	19,347	4,935	2,351	2,584
All other diseases (Residual)	269,844	109,547	160,297	235,449	94,966	140,483	28,532	11,902	16,630

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120,859	75,921	44,938	104,945	65,360	39,585	12,069	8,074	3,995
Transport accidents (V01-V99,Y85)	37,961	26,783	11,178	31,637	22,348	9,289	4,751	3,421	1,330
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)	35,332	24,723	10,609	29,392	20,589	8,803	4,477	3,206	1,271
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)	1,029	794	235	840	647	193	136	106	30
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	1,600	1,266	334	1,405	1,112	293	138	109	29
Nontransport accidents (W00-X59,Y86)	82,898	49,138	33,760	73,308	43,012	30,296	7,318	4,653	2,665
Falls (W00-W19)	26,009	13,049	12,960	24,133	11,967	12,166	1,164	669	495
Accidental discharge of firearms (W32-W34)	606	515	91	474	396	78	113	101	12
Accidental drowning and submersion (W65-W74)	3,782	2,936	846	2,970	2,267	703	571	478	93
Accidental exposure to smoke, fire and flames (X00-X09)	2,782	1,624	1,158	2,135	1,251	884	577	335	242
Accidental poisoning and exposure to noxious substances (X40-X49)	33,041	21,117	11,924	29,283	18,717	10,566	2,988	1,896	1,092
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)	16,678	9,897	6,781	14,313	8,414	5,899	1,905	1,174	731
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,364	30,277	8,087	34,690	27,422	7,268	2,144	1,755	389
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	19,392	16,962	2,430	17,909	15,648	2,261	1,079	965	114
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)	18,972	13,315	5,657	16,781	11,774	5,007	1,065	790	275
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16,259	12,774	3,485	7,863	5,648	2,215	7,818	6,704	1,114
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	11,078	9,340	1,738	4,647	3,555	1,092	6,151	5,553	598
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	5,181	3,434	1,747	3,216	2,093	1,123	1,667	1,151	516
Legal intervention (Y35,Y89.0)	412	399	13	295	287	8	98	96	2

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Events of undetermined intent (Y10-Y34, Y87.2, Y89.9)	4,908	2,930	1,978	4,147	2,448	1,699	594	381	213
Discharge of firearms, undetermined intent (Y22-Y24)	252	203	49	210	171	39	32	25	7
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21, Y25-Y34, Y87.2, Y89.9)	4,656	2,727	1,929	3,937	2,277	1,660	562	356	206
Operations of war and their sequelae (Y36, Y89.1)	9	9	-	7	7	-	1	1	-
Complications of medical and surgical care (Y40-Y84, Y88)	2,490	1,167	1,323	2,023	971	1,052	401	169	232
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	7,298	2,889	4,409	6,671	2,623	4,048	488	203	285
Drug-induced deaths ^{5,6}	40,393	24,376	16,017	36,020	21,697	14,323	3,561	2,210	1,351
Alcohol-induced deaths ^{5,7}	25,692	19,038	6,654	22,167	16,527	5,640	2,330	1,686	644
Injury by firearms ^{5,8}	31,672	27,356	4,316	23,490	20,014	3,476	7,454	6,721	733

- Quantity zero.

... Category not applicable.

¹Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and other Asian or Pacific Islander groups.

⁴Included in "Certain other intestinal infections (A04, A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, F11.1-F11.5, F11.7-F11.9, F12.1-F12.5, F12.7-F12.9, F13.1-F13.5, F13.7-F13.9, F14.1-F14.5, F14.7-F14.9, F15.1-F15.5, F15.7-F15.9, F16.1-F16.5, F16.7-F16.9, F17.3-F17.5, F17.7-F17.9, F18.1-F18.5, F18.7-F18.9, F19.1-F19.5, F19.7-F19.9, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2-J70.4, K85.3, L10.5, L27.0-L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R50.2, R78.1-R78.5, X40-X44, X60-X64, X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, R78.0, X45, X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4, W32-W34, X72-X74, X93-X95, Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
All causes	15,565	8,516	7,049	51,162	26,600	24,562
Salmonella infections (A01-A02)	-	-	-	4	2	2
Shigellosis and amebiasis (A03,A06)	-	-	-	1	-	1
Certain other intestinal infections (A04,A07-A09)	62	30	32	139	64	75
Tuberculosis (A16-A19)	15	6	9	96	59	37
Respiratory tuberculosis (A16)	12	5	7	78	51	27
Other tuberculosis (A17-A19)	3	1	2	18	8	10
Whooping cough (A37)	-	-	-	-	-	-
Scarlet fever and erysipelas (A38,A46)	-	-	-	1	-	1
Meningococcal infection (A39)	-	-	-	3	1	2
Septicemia (A40-A41)	244	106	138	582	296	286
Syphilis (A50-A53)	-	-	-	1	-	1
Acute poliomyelitis (A80)	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)	-	-	-	1	1	-
Measles (B05)	1	1	-	-	-	-
Viral hepatitis (B15-B19)	99	63	36	303	171	132
Human immunodeficiency virus (HIV) disease (B20-B24)	61	48	13	71	53	18
Malaria (B50-B54)	-	-	-	2	1	1
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	53	28	25	173	96	77
Malignant neoplasms (C00-C97)	2,962	1,588	1,374	14,165	7,210	6,955
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	41	25	16	276	178	98
Malignant neoplasm of esophagus (C15)	63	49	14	229	175	54
Malignant neoplasm of stomach (C16)	87	57	30	779	443	336
Malignant neoplasms of colon, rectum and anus (C18-C21)	287	152	135	1,476	749	727
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	196	143	53	1,334	902	432
Malignant neoplasm of pancreas (C25)	177	97	80	971	442	529
Malignant neoplasm of larynx (C32)	25	20	5	46	41	5
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	785	445	340	3,147	1,827	1,320
Malignant melanoma of skin (C43)	22	12	10	53	23	30

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of breast (C50)	172	2	170	971	10	961
Malignant neoplasm of cervix uteri (C53)	36	...	36	143	...	143
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	36	...	36	241	...	241
Malignant neoplasm of ovary (C56)	73	...	73	379	...	379
Malignant neoplasm of prostate (C61)	117	117	...	418	418	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	110	74	36	265	180	85
Malignant neoplasm of bladder (C67)	47	31	16	207	136	71
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	73	38	35	319	168	151
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	230	129	101	1,326	740	586
Hodgkin's disease (C81)	5	3	2	25	18	7
Non-Hodgkin's lymphoma (C82-C85)	70	41	29	517	274	243
Leukemia (C91-C95)	107	59	48	558	311	247
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	48	26	22	223	134	89
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	-	-	-	3	3	-
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	385	197	188	1,585	778	807
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	33	27	350	172	178
Anemias (D50-D64)	18	5	13	86	41	45
Diabetes mellitus (E10-E14)	857	432	425	1,838	932	906
Nutritional deficiencies (E40-E64)	29	9	20	49	24	25
Malnutrition (E40-E46)	28	9	19	47	22	25
Other nutritional deficiencies (E50-E64)	1	-	1	2	2	-
Meningitis (G00,G03)	7	4	3	21	14	7
Parkinson's disease (G20-G21)	62	40	22	374	206	168
Alzheimer's disease (G30)	264	89	175	1,082	345	737
Major cardiovascular diseases (I00-I78)	3,603	1,972	1,631	16,631	8,622	8,009
Diseases of heart (I00-I09,I11,I13,I20-I51)	2,793	1,608	1,185	11,490	6,262	5,228
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	14	6	8	97	26	71

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart disease (I11)	169	100	69	683	341	342
Hypertensive heart and renal disease (I13)	18	10	8	81	34	47
Ischemic heart diseases (I20-I25)	1,831	1,098	733	7,821	4,481	3,340
Acute myocardial infarction (I21-I22)	594	367	227	2,530	1,442	1,088
Other acute ischemic heart diseases (I24)	39	24	15	55	31	24
Other forms of chronic ischemic heart disease (I20,I25)	1,198	707	491	5,236	3,008	2,228
Atherosclerotic cardiovascular disease, so described (I25.0)	372	233	139	1,235	750	485
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	826	474	352	4,001	2,258	1,743
Other heart diseases (I26-I51)	761	394	367	2,808	1,380	1,428
Acute and subacute endocarditis (I33)	11	5	6	20	15	5
Diseases of pericardium and acute myocarditis (I30-I31,I40)	10	7	3	23	14	9
Heart failure (I50)	225	91	134	714	310	404
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	515	291	224	2,051	1,041	1,010
Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	62	82	814	337	477
Cerebrovascular diseases (I60-I69)	559	258	301	3,833	1,747	2,086
Atherosclerosis (I70)	19	5	14	104	56	48
Other diseases of circulatory system (I71-I78)	88	39	49	390	220	170
Aortic aneurysm and dissection (I71)	48	26	22	277	159	118
Other diseases of arteries, arterioles and capillaries (I72-I78)	40	13	27	113	61	52
Other disorders of circulatory system (I80-I99)	23	9	14	78	45	33
Influenza and pneumonia (J09-J18)	326	172	154	1,539	825	714
Influenza (J09-J11)	4	2	2	19	11	8
Pneumonia (J12-J18)	322	170	152	1,520	814	706
Other acute lower respiratory infections (J20-J22,U04)	2	1	1	7	3	4
Acute bronchitis and bronchiolitis (J20-J21)	2	1	1	7	3	4
Other and unspecified acute lower respiratory infections (J22,U04)	-	-	-	-	-	-
Chronic lower respiratory diseases (J40-J47)	702	349	353	1,487	910	577
Bronchitis, chronic and unspecified (J40-J42)	4	4	-	11	5	6
Emphysema (J43)	44	21	23	100	79	21
Asthma (J45-J46)	23	12	11	136	73	63

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Other chronic lower respiratory diseases (J44,J47)	631	312	319	1,240	753	487
Pneumoconioses and chemical effects (J60-J66,J68)	4	4	-	2	2	-
Pneumonitis due to solids and liquids (J69)	83	40	43	313	180	133
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	238	113	125	647	332	315
Peptic ulcer (K25-K28)	22	8	14	87	47	40
Diseases of appendix (K35-K38)	5	3	2	11	9	2
Hernia (K40-K46)	14	8	6	17	10	7
Chronic liver disease and cirrhosis (K70,K73-K74)	787	429	358	467	302	165
Alcoholic liver disease (K70)	592	351	241	175	143	32
Other chronic liver disease and cirrhosis (K73-K74)	195	78	117	292	159	133
Cholelithiasis and other disorders of gallbladder (K80-K82)	20	11	9	100	52	48
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	153	186	1,091	524	567
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	1	-	1	9	4	5
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	41	20	21	135	53	82
Renal failure (N17-N19)	297	133	164	947	467	480
Other disorders of kidney (N25,N27)	-	-	-	-	-	-
Infections of kidney (N10-N12,N13.6,N15.1)	5	1	4	12	2	10
Hyperplasia of prostate (N40)	-	-	...	8	8	...
Inflammatory diseases of female pelvic organs (N70-N76)	3	...	3	-	...	-
Pregnancy, childbirth and the puerperium (O00-O99)	15	...	15	42	...	42
Pregnancy with abortive outcome (O00-O07)	-	...	-	-	...	-
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	15	...	15	42	...	42
Certain conditions originating in the perinatal period (P00-P96)	117	76	41	467	278	189
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	71	59	321	157	164
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	240	129	111	517	256	261
All other diseases (Residual)	1,549	725	824	4,314	1,954	2,360

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,701	1,150	551	2,144	1,337	807
Transport accidents (V01-V99,Y85)	684	462	222	889	552	337
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)	638	425	213	825	503	322
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)	22	19	3	31	22	9
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	24	18	6	33	27	6
Nontransport accidents (W00-X59,Y86)	1,017	688	329	1,255	785	470
Falls (W00-W19)	161	97	64	551	316	235
Accidental discharge of firearms (W32-W34)	14	14	-	5	4	1
Accidental drowning and submersion (W65-W74)	68	60	8	173	131	42
Accidental exposure to smoke, fire and flames (X00-X09)	45	26	19	25	12	13
Accidental poisoning and exposure to noxious substances (X40-X49)	521	332	189	249	172	77
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)	208	159	49	252	150	102
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	344	125	1,061	756	305
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	178	146	32	226	203	23
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)	291	198	93	835	553	282
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	204	53	321	218	103
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	113	98	15	167	134	33
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	144	106	38	154	84	70
Legal intervention (Y35,Y89.0)	9	6	3	10	10	-

Table 12. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	87	46	41	80	55	25
Discharge of firearms, undetermined intent (Y22-Y24)	7	4	3	3	3	-
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34, Y87.2,Y89.9)	80	42	38	77	52	25
Operations of war and their sequelae (Y36,Y89.1)	-	-	-	1	1	-
Complications of medical and surgical care (Y40-Y84,Y88)	21	10	11	45	17	28
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	39	20	19	100	43	57
Drug-induced deaths ^{5,6}	458	250	208	354	219	135
Alcohol-induced deaths ^{5,7}	931	611	320	264	214	50
Injury by firearms ^{5,8}	317	267	50	411	354	57

- Quantity zero.

... Category not applicable.

¹Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and other Asian or Pacific Islander groups.

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins			Hispanic			Non-Hispanic ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	2,468,435	1,232,432	1,236,003	144,490	79,622	64,868	2,318,218	1,149,438	1,168,780
Salmonella infections (A01-A02)	28	17	11	3	-	3	25	17	8
Shigellosis and amebiasis (A03,A06)	3	1	2	-	-	-	3	1	2
Certain other intestinal infections (A04,A07-A09)	10,276	4,046	6,230	603	269	334	9,657	3,770	5,887
Tuberculosis (A16-A19)	569	351	218	98	63	35	467	285	182
Respiratory tuberculosis (A16)	423	276	147	73	52	21	347	222	125
Other tuberculosis (A17-A19)	146	75	71	25	11	14	120	63	57
Whooping cough (A37)	26	13	13	11	5	6	15	8	7
Scarlet fever and erysipelas (A38,A46)	3	1	2	1	1	-	2	-	2
Meningococcal infection (A39)	79	40	39	22	10	12	57	30	27
Septicemia (A40-A41)	34,812	16,069	18,743	2,035	1,012	1,023	32,716	15,028	17,688
Syphilis (A50-A53)	28	20	8	3	3	-	25	17	8
Acute poliomyelitis (A80)	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)	9	4	5	-	-	-	9	4	5
Measles (B05)	2	2	-	1	1	-	1	1	-
Viral hepatitis (B15-B19)	7,564	5,038	2,526	1,118	766	352	6,414	4,248	2,166
Human immunodeficiency virus (HIV) disease (B20-B24)	8,369	6,099	2,270	1,134	910	224	7,175	5,140	2,035
Malaria (B50-B54)	10	7	3	-	-	-	10	7	3
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	5,805	2,970	2,835	441	233	208	5,353	2,732	2,621
Malignant neoplasms (C00-C97)	574,743	301,037	273,706	31,119	16,450	14,669	542,581	284,018	258,563
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	8,474	5,815	2,659	412	298	114	8,035	5,497	2,538
Malignant neoplasm of esophagus (C15)	14,490	11,416	3,074	592	475	117	13,870	10,919	2,951
Malignant neoplasm of stomach (C16)	11,390	6,703	4,687	1,554	873	681	9,814	5,815	3,999
Malignant neoplasms of colon, rectum and anus (C18-C21)	52,622	27,284	25,338	3,179	1,782	1,397	49,358	25,455	23,903
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	20,305	13,658	6,647	2,331	1,584	747	17,929	12,045	5,884
Malignant neoplasm of pancreas (C25)	36,888	18,699	18,189	2,142	1,046	1,096	34,694	17,620	17,074
Malignant neoplasm of larynx (C32)	3,691	2,951	740	211	188	23	3,469	2,753	716

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins			Hispanic			Non-Hispanic ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	158,318	87,740	70,578	4,953	3,046	1,907	153,042	84,519	68,523
Malignant melanoma of skin (C43)	9,154	6,002	3,152	216	125	91	8,926	5,870	3,056
Malignant neoplasm of breast (C50)	41,435	439	40,996	2,312	17	2,295	39,044	421	38,623
Malignant neoplasm of cervix uteri (C53)	3,939	...	3,939	476	...	476	3,456	...	3,456
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	8,402	...	8,402	528	...	528	7,853	...	7,853
Malignant neoplasm of ovary (C56)	14,572	...	14,572	848	...	848	13,691	...	13,691
Malignant neoplasm of prostate (C61)	28,561	28,561	...	1,535	1,535	...	26,973	26,973	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	13,219	8,436	4,783	944	588	356	12,255	7,832	4,423
Malignant neoplasm of bladder (C67)	14,731	10,429	4,302	519	352	167	14,187	10,063	4,124
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	14,164	7,977	6,187	884	514	370	13,269	7,456	5,813
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	55,590	30,777	24,813	3,634	1,968	1,666	51,881	28,760	23,121
Hodgkin's disease (C81)	1,231	714	517	133	82	51	1,095	629	466
Non-Hodgkin's lymphoma (C82-C85)	20,294	11,047	9,247	1,327	711	616	18,937	10,314	8,623
Leukemia (C91-C95)	22,569	12,851	9,718	1,444	785	659	21,096	12,051	9,045
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	11,428	6,117	5,311	728	388	340	10,687	5,720	4,967
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	68	48	20	2	2	-	66	46	20
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	64,798	34,150	30,648	3,849	2,059	1,790	60,835	32,020	28,815
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14,917	7,721	7,196	784	404	380	14,107	7,302	6,805
Anemias (D50-D64)	4,852	2,017	2,835	236	117	119	4,602	1,892	2,710
Diabetes mellitus (E10-E14)	69,071	35,490	33,581	6,556	3,372	3,184	62,357	32,027	30,330
Nutritional deficiencies (E40-E64)	2,948	1,141	1,807	175	81	94	2,768	1,057	1,711
Malnutrition (E40-E46)	2,790	1,079	1,711	166	78	88	2,620	999	1,621

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins			Hispanic			Non-Hispanic ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other nutritional deficiencies (E50-E64)	158	62	96	9	3	6	148	58	90
Meningitis (G00,G03)	608	303	305	71	33	38	533	266	267
Parkinson's disease (G20-G21)	22,032	12,871	9,161	982	555	427	21,035	12,308	8,727
Alzheimer's disease (G30)	83,494	25,364	58,130	3,427	1,171	2,256	79,984	24,169	55,815
Major cardiovascular diseases (I00-I78)	780,213	383,547	396,666	40,154	21,220	18,934	738,164	361,226	376,938
Diseases of heart									
(I00-I09,I11,I13,I20-I51)	597,689	307,384	290,305	30,006	16,421	13,585	566,098	290,006	276,092
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	2,987	996	1,991	154	52	102	2,830	942	1,888
Hypertensive heart disease (I11)	33,678	16,242	17,436	2,094	1,169	925	31,406	14,967	16,439
Hypertensive heart and renal disease (I13)	2,807	1,285	1,522	215	108	107	2,586	1,174	1,412
Ischemic heart diseases (I20-I25)	379,559	207,580	171,979	20,494	11,546	8,948	357,969	195,358	162,611
Acute myocardial infarction (I21-I22)	122,071	67,435	54,636	6,554	3,726	2,828	115,254	63,543	51,711
Other acute ischemic heart diseases (I24)	4,170	2,156	2,014	134	60	74	4,029	2,090	1,939
Other forms of chronic ischemic heart disease (I20,I25)	253,318	137,989	115,329	13,806	7,760	6,046	238,686	129,725	108,961
Atherosclerotic cardiovascular disease, so described (I25.0)	57,438	33,703	23,735	3,419	2,242	1,177	53,660	31,208	22,452
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	195,880	104,286	91,594	10,387	5,518	4,869	185,026	98,517	86,509
Other heart diseases (I26-I51)	178,658	81,281	97,377	7,049	3,546	3,503	171,307	77,565	93,742
Acute and subacute endocarditis (I33)	1,103	627	476	78	47	31	1,023	579	444
Diseases of pericardium and acute myocarditis (I30-I31,I40)	776	405	371	57	31	26	716	372	344
Heart failure (I50)	57,757	24,385	33,372	2,024	936	1,088	55,661	23,409	32,252
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	119,022	55,864	63,158	4,890	2,532	2,358	113,907	53,205	60,702
Essential hypertension and hypertensive renal disease (I10,I12,I15)	26,634	10,846	15,788	1,712	764	948	24,872	10,058	14,814
Cerebrovascular diseases (I60-I69)	129,476	52,367	77,109	7,274	3,382	3,892	122,000	48,894	73,106
Atherosclerosis (I70)	7,230	2,933	4,297	269	127	142	6,952	2,803	4,149
Other diseases of circulatory system (I71-I78)	19,184	10,017	9,167	893	526	367	18,242	9,465	8,777
Aortic aneurysm and dissection (I71)	10,431	6,096	4,335	438	295	143	9,966	5,781	4,185

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins			Hispanic			Non-Hispanic ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of arteries, arterioles and capillaries (I72-I78)	8,753	3,921	4,832	455	231	224	8,276	3,684	4,592
Other disorders of circulatory system (I80-I99)	4,241	2,053	2,188	259	135	124	3,964	1,908	2,056
Influenza and pneumonia (J09-J18)	50,097	23,615	26,482	3,025	1,565	1,460	46,937	21,981	24,956
Influenza (J09-J11)	500	250	250	92	51	41	408	199	209
Pneumonia (J12-J18)	49,597	23,365	26,232	2,933	1,514	1,419	46,529	21,782	24,747
Other acute lower respiratory infections (J20-J22,U04)	213	94	119	13	8	5	200	86	114
Acute bronchitis and bronchiolitis (J20-J21)	177	74	103	12	7	5	165	67	98
Other and unspecified acute lower respiratory infections (J22,U04)	36	20	16	1	1	-	35	19	16
Chronic lower respiratory diseases (J40-J47)	138,080	65,423	72,657	4,172	2,174	1,998	133,638	63,100	70,538
Bronchitis, chronic and unspecified (J40-J42)	620	267	353	45	21	24	574	245	329
Emphysema (J43)	10,034	5,227	4,807	250	152	98	9,749	5,051	4,698
Asthma (J45-J46)	3,404	1,283	2,121	292	120	172	3,099	1,157	1,942
Other chronic lower respiratory diseases (J44,J47)	124,022	58,646	65,376	3,585	1,881	1,704	120,216	56,647	63,569
Pneumoconioses and chemical effects (J60-J66,J68)	845	812	33	23	23	-	820	787	33
Pneumonitis due to solids and liquids (J69)	17,011	9,208	7,803	691	396	295	16,300	8,796	7,504
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	31,187	15,791	15,396	2,021	1,013	1,008	29,107	14,737	14,370
Peptic ulcer (K25-K28)	2,977	1,460	1,517	140	76	64	2,826	1,376	1,450
Diseases of appendix (K35-K38)	415	258	157	28	22	6	385	234	151
Hernia (K40-K46)	1,832	814	1,018	120	44	76	1,707	766	941
Chronic liver disease and cirrhosis (K70,K73-K74)	31,903	20,798	11,105	4,348	3,067	1,281	27,455	17,657	9,798
Alcoholic liver disease (K70)	15,990	11,441	4,549	2,389	1,937	452	13,540	9,458	4,082
Other chronic liver disease and cirrhosis (K73-K74)	15,913	9,357	6,556	1,959	1,130	829	13,915	8,199	5,716
Cholelithiasis and other disorders of gallbladder (K80-K82)	3,332	1,552	1,780	247	130	117	3,075	1,419	1,656
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,476	24,865	25,611	3,252	1,670	1,582	47,120	23,140	23,980

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins			Hispanic			Non-Hispanic ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	203	97	106	12	6	6	191	91	100
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	5,894	2,775	3,119	326	162	164	5,552	2,603	2,949
Renal failure (N17-N19)	44,362	21,989	22,373	2,913	1,502	1,411	41,361	20,442	20,919
Other disorders of kidney (N25,N27)	17	4	13	1	-	1	16	4	12
Infections of kidney (N10-N12,N13.6,N15.1)	608	187	421	44	12	32	562	175	387
Hyperplasia of prostate (N40)	489	489	...	28	28	...	459	459	...
Inflammatory diseases of female pelvic organs (N70-N76)	137	...	137	12	...	12	125	...	125
Pregnancy, childbirth and the puerperium (O00-O99)	825	...	825	146	...	146	670	...	670
Pregnancy with abortive outcome (O00-O07)	37	...	37	8	...	8	28	...	28
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	788	...	788	138	...	138	642	...	642
Certain conditions originating in the perinatal period (P00-P96)	12,128	6,803	5,325	2,529	1,456	1,073	9,446	5,269	4,177
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9,673	4,960	4,713	1,786	903	883	7,845	4,032	3,813
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	38,360	16,057	22,303	2,069	1,065	1,004	36,151	14,900	21,251
All other diseases (Residual)	269,844	109,547	160,297	13,964	6,565	7,399	255,372	102,717	152,655
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120,859	75,921	44,938	10,476	7,594	2,882	109,968	68,032	41,936
Transport accidents (V01-V99,Y85)	37,961	26,783	11,178	4,778	3,518	1,260	33,045	23,165	9,880
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)	35,332	24,723	10,609	4,509	3,291	1,218	30,700	21,345	9,355

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins			Hispanic			Non-Hispanic ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)	1,029	794	235	169	146	23	854	642	212
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	1,600	1,266	334	100	81	19	1,491	1,178	313
Nontransport accidents (W00-X59,Y86)	82,898	49,138	33,760	5,698	4,076	1,622	76,923	44,867	32,056
Falls (W00-W19)	26,009	13,049	12,960	1,275	791	484	24,683	12,234	12,449
Accidental discharge of firearms (W32-W34)	606	515	91	37	34	3	568	480	88
Accidental drowning and submersion (W65-W74)	3,782	2,936	846	527	447	80	3,241	2,476	765
Accidental exposure to smoke, fire and flames (X00-X09)	2,782	1,624	1,158	171	113	58	2,595	1,499	1,096
Accidental poisoning and exposure to noxious substances (X40-X49)	33,041	21,117	11,924	2,601	1,918	683	30,292	19,086	11,206
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)	16,678	9,897	6,781	1,087	773	314	15,544	9,092	6,452
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,364	30,277	8,087	2,661	2,168	493	35,562	27,994	7,568
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	19,392	16,962	2,430	962	868	94	18,365	16,036	2,329
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)	18,972	13,315	5,657	1,699	1,300	399	17,197	11,958	5,239
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16,259	12,774	3,485	2,890	2,435	455	13,252	10,238	3,014
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	11,078	9,340	1,738	1,919	1,706	213	9,082	7,565	1,517

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins			Hispanic			Non-Hispanic ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	5,181	3,434	1,747	971	729	242	4,170	2,673	1,497
Legal intervention (Y35,Y89.0)	412	399	13	75	75	-	335	322	13
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	4,908	2,930	1,978	349	242	107	4,531	2,667	1,864
Discharge of firearms, undetermined intent (Y22-Y24)	252	203	49	25	21	4	227	182	45
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	4,656	2,727	1,929	324	221	103	4,304	2,485	1,819
Operations of war and their sequelae (Y36,Y89.1)	9	9	-	-	-	-	9	9	-
Complications of medical and surgical care (Y40-Y84,Y88)	2,490	1,167	1,323	148	80	68	2,337	1,084	1,253
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	7,298	2,889	4,409	380	158	222	6,908	2,726	4,182
Drug-induced deaths ^{5,6}	40,393	24,376	16,017	2,788	1,944	844	37,420	22,303	15,117
Alcohol-induced deaths ^{5,7}	25,692	19,038	6,654	3,326	2,759	567	22,242	16,178	6,064
Injury by firearms ^{5,8}	31,672	27,356	4,316	3,008	2,694	314	28,519	24,532	3,987

- Quantity zero.

... Category not applicable.

¹Includes races other than white and black.

²Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

³Includes deaths for which Hispanic origin was not reported on the death certificate.

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	1,969,916	971,604	998,312	283,438	143,824	139,614	5,727	3,372	2,355
Salmonella infections (A01-A02)	16	14	2	5	1	4	-	-	-
Shigellosis and amebiasis (A03,A06)	1	1	-	1	-	1	-	-	-
Certain other intestinal infections (A04,A07-A09)	8,661	3,323	5,338	801	357	444	16	7	9
Tuberculosis (A16-A19)	238	135	103	119	86	33	4	3	1
Respiratory tuberculosis (A16)	165	93	72	93	74	19	3	2	1
Other tuberculosis (A17-A19)	73	42	31	26	12	14	1	1	-
Whooping cough (A37)	13	6	7	2	2	-	-	-	-
Scarlet fever and erysipelas (A38,A46)	1	-	1	-	-	-	-	-	-
Meningococcal infection (A39)	46	25	21	8	4	4	-	-	-
Septicemia (A40-A41)	25,951	11,963	13,988	5,958	2,673	3,285	61	29	32
Syphilis (A50-A53)	12	9	3	12	8	4	-	-	-
Acute poliomyelitis (A80)	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)	8	3	5	-	-	-	-	-	-
Measles (B05)	-	-	-	-	-	-	-	-	-
Viral hepatitis (B15-B19)	4,942	3,331	1,611	1,089	697	392	32	24	8
Human immunodeficiency virus (HIV) disease (B20-B24)	2,454	2,046	408	4,598	3,000	1,598	60	49	11
Malaria (B50-B54)	2	1	1	6	5	1	-	-	-
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	4,359	2,205	2,154	771	404	367	11	5	6
Malignant neoplasms (C00-C97)	460,567	241,816	218,751	65,276	33,607	31,669	1,043	569	474
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	6,728	4,580	2,148	996	719	277	27	20	7
Malignant neoplasm of esophagus (C15)	12,161	9,665	2,496	1,424	1,037	387	28	22	6
Malignant neoplasm of stomach (C16)	6,983	4,157	2,826	1,974	1,162	812	22	15	7
Malignant neoplasms of colon, rectum and anus (C18-C21)	40,686	20,990	19,696	6,946	3,584	3,362	85	47	38
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	13,643	9,069	4,574	2,788	1,958	830	45	29	16
Malignant neoplasm of pancreas (C25)	29,278	15,013	14,265	4,292	2,079	2,213	52	33	19
Malignant neoplasm of larynx (C32)	2,793	2,219	574	605	473	132	11	10	1

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	132,658	72,584	60,074	16,536	9,711	6,825	323	175	148
Malignant melanoma of skin (C43)	8,721	5,775	2,946	134	61	73	12	7	5
Malignant neoplasm of breast (C50)	31,868	333	31,535	6,063	77	5,986	79	1	78
Malignant neoplasm of cervix uteri (C53)	2,506	...	2,506	776	...	776	7	...	7
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	6,162	...	6,162	1,421	...	1,421	21	...	21
Malignant neoplasm of ovary (C56)	11,984	...	11,984	1,264	...	1,264	33	...	33
Malignant neoplasm of prostate (C61)	21,657	21,657	...	4,801	4,801	...	53	53	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	10,713	6,882	3,831	1,175	702	473	20	16	4
Malignant neoplasm of bladder (C67)	12,877	9,289	3,588	1,065	610	455	25	14	11
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	12,002	6,801	5,201	881	453	428	11	7	4
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	44,975	25,120	19,855	5,390	2,792	2,598	75	49	26
Hodgkin's disease (C81)	952	549	403	114	60	54	3	3	-
Non-Hodgkin's lymphoma (C82-C85)	16,873	9,192	7,681	1,488	813	675	30	22	8
Leukemia (C91-C95)	18,654	10,744	7,910	1,794	945	849	29	15	14
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	8,441	4,596	3,845	1,986	970	1,016	13	9	4
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	55	39	16	8	4	4	-	-	-
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	52,172	27,682	24,490	6,745	3,388	3,357	114	71	43
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12,494	6,557	5,937	1,211	543	668	26	15	11
Anemias (D50-D64)	3,503	1,404	2,099	995	442	553	14	8	6
Diabetes mellitus (E10-E14)	47,746	25,134	22,612	11,996	5,572	6,424	158	91	67
Nutritional deficiencies (E40-E64)	2,325	891	1,434	366	134	232	5	3	2

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malnutrition (E40-E46)	2,190	840	1,350	356	129	227	4	2	2
Other nutritional deficiencies (E50-E64)	135	51	84	10	5	5	1	1	-
Meningitis (G00,G03)	369	185	184	137	63	74	4	4	-
Parkinson's disease (G20-G21)	19,818	11,627	8,191	795	445	350	15	8	7
Alzheimer's disease (G30)	73,502	22,278	51,224	5,169	1,471	3,698	83	24	59
Major cardiovascular diseases (I00-I78)	626,610	305,928	320,682	91,807	44,993	46,814	1,895	1,101	794
Diseases of heart (I00-I09,I11,I13,I20-I51)	483,973	247,765	236,208	68,215	34,597	33,618	1,585	957	628
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	2,493	831	1,662	228	79	149	3	2	1
Hypertensive heart disease (I11)	22,966	10,533	12,433	7,611	4,007	3,604	178	106	72
Hypertensive heart and renal disease (I13)	1,744	757	987	747	377	370	6	3	3
Ischemic heart diseases (I20-I25)	309,492	169,637	139,855	39,047	20,282	18,765	1,096	676	420
Acute myocardial infarction (I21-I22)	99,600	55,411	44,189	12,597	6,366	6,231	263	166	97
Other acute ischemic heart diseases (I24)	3,388	1,739	1,649	549	298	251	7	6	1
Other forms of chronic ischemic heart disease (I20,I25)	206,504	112,487	94,017	25,901	13,618	12,283	826	504	322
Atherosclerotic cardiovascular disease, so described (I25.0)	43,882	25,424	18,458	8,216	4,832	3,384	359	253	106
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	162,622	87,063	75,559	17,685	8,786	8,899	467	251	216
Other heart diseases (I26-I51)	147,278	66,007	81,271	20,582	9,852	10,730	302	170	132
Acute and subacute endocarditis (I33)	810	460	350	183	100	83	2	1	1
Diseases of pericardium and acute myocarditis (I30-I31,I40)	563	290	273	124	64	60	3	2	1
Heart failure (I50)	49,253	20,596	28,657	5,497	2,428	3,069	72	40	32
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	96,652	44,661	51,991	14,778	7,260	7,518	225	127	98
Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,878	7,476	11,402	5,059	2,196	2,863	50	24	26
Cerebrovascular diseases (I60-I69)	101,849	40,040	61,809	15,833	6,890	8,943	202	91	111
Atherosclerosis (I70)	6,242	2,490	3,752	588	252	336	9	3	6

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of circulatory system (I71-I78)	15,668	8,157	7,511	2,112	1,058	1,054	49	26	23
Aortic aneurysm and dissection (I71)	8,697	5,063	3,634	956	541	415	27	20	7
Other diseases of arteries, arterioles and capillaries (I72-I78)	6,971	3,094	3,877	1,156	517	639	22	6	16
Other disorders of circulatory system (I80-I99)	3,118	1,483	1,635	749	374	375	18	10	8
Influenza and pneumonia (J09-J18)	40,244	18,657	21,587	4,854	2,340	2,514	135	69	66
Influenza (J09-J11)	319	155	164	66	31	35	-	-	-
Pneumonia (J12-J18)	39,925	18,502	21,423	4,788	2,309	2,479	135	69	66
Other acute lower respiratory infections (J20-J22,U04)	169	72	97	22	10	12	-	-	-
Acute bronchitis and bronchiolitis (J20-J21)	135	54	81	21	9	12	-	-	-
Other and unspecified acute lower respiratory infections (J22,U04)	34	18	16	1	1	-	-	-	-
Chronic lower respiratory diseases (J40-J47)	122,887	57,390	65,497	8,614	4,480	4,134	270	149	121
Bronchitis, chronic and unspecified (J40-J42)	514	212	302	48	26	22	1	1	-
Emphysema (J43)	9,021	4,594	4,427	586	358	228	35	24	11
Asthma (J45-J46)	2,082	676	1,406	862	399	463	13	6	7
Other chronic lower respiratory diseases (J44,J47)	111,270	51,908	59,362	7,118	3,697	3,421	221	118	103
Pneumoconioses and chemical effects (J60-J66,J68)	778	748	30	36	33	3	2	2	-
Pneumonitis due to solids and liquids (J69)	14,539	7,862	6,677	1,369	716	653	20	16	4
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	25,350	12,949	12,401	2,900	1,360	1,540	59	41	18
Peptic ulcer (K25-K28)	2,428	1,153	1,275	290	168	122	11	8	3
Diseases of appendix (K35-K38)	328	195	133	42	28	14	2	2	-
Hernia (K40-K46)	1,515	671	844	162	78	84	5	4	1
Chronic liver disease and cirrhosis (K70,K73-K74)	23,655	15,270	8,385	2,601	1,694	907	100	74	26
Alcoholic liver disease (K70)	11,607	8,185	3,422	1,206	808	398	61	46	15
Other chronic liver disease and cirrhosis (K73-K74)	12,048	7,085	4,963	1,395	886	509	39	28	11

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Cholelithiasis and other disorders of gallbladder (K80-K82)	2,663	1,236	1,427	296	120	176	10	3	7
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,963	18,503	18,460	8,769	3,984	4,785	104	55	49
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	150	70	80	32	18	14	-	-	-
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	4,445	2,108	2,337	937	427	510	16	10	6
Renal failure (N17-N19)	32,356	16,322	16,034	7,796	3,538	4,258	88	45	43
Other disorders of kidney (N25,N27)	12	3	9	4	1	3	-	-	-
Infections of kidney (N10-N12,N13.6,N15.1)	481	147	334	65	26	39	2	-	2
Hyperplasia of prostate (N40)	416	416	...	35	35	...	2	2	...
Inflammatory diseases of female pelvic organs (N70-N76)	101	...	101	21	...	21	-	...	-
Pregnancy, childbirth and the puerperium (O00-O99)	357	...	357	261	...	261	9	...	9
Pregnancy with abortive outcome (O00-O07)	11	...	11	17	...	17	1	...	1
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	346	...	346	244	...	244	8	...	8
Certain conditions originating in the perinatal period (P00-P96)	5,065	2,830	2,235	3,846	2,116	1,730	153	78	75
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5,832	2,981	2,851	1,608	850	758	42	25	17
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	30,602	12,246	18,356	4,840	2,293	2,547	140	92	48
All other diseases (Residual)	221,437	88,360	133,077	28,230	11,758	16,472	508	265	243
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,420	57,747	36,673	11,853	7,920	3,933	415	295	120
Transport accidents (V01-V99,Y85)	26,868	18,845	8,023	4,668	3,358	1,310	138	100	38

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)	24,898	17,318	7,580	4,398	3,146	1,252	123	87	36
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9, V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	669	499	170	135	105	30	6	6	-
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	1,301	1,028	273	135	107	28	9	7	2
Nontransport accidents (W00-X59,Y86)	67,552	38,902	28,650	7,185	4,562	2,623	277	195	82
Falls (W00-W19)	22,842	11,178	11,664	1,147	660	487	51	24	27
Accidental discharge of firearms (W32-W34)	439	364	75	112	100	12	1	1	-
Accidental drowning and submersion (W65-W74)	2,462	1,838	624	551	459	92	14	13	1
Accidental exposure to smoke, fire and flames (X00-X09)	1,956	1,130	826	569	331	238	16	12	4
Accidental poisoning and exposure to noxious substances (X40-X49)	26,620	16,746	9,874	2,938	1,861	1,077	148	113	35
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39, X50-X59,Y86)	13,233	7,646	5,587	1,868	1,151	717	47	32	15
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,010	25,238	6,772	2,091	1,712	379	141	115	26
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	16,928	14,762	2,166	1,057	946	111	65	58	7
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)	15,082	10,476	4,606	1,034	766	268	76	57	19

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide)									
(*U01-*U02,X85-Y09,Y87.1)	5,035	3,267	1,768	7,679	6,580	1,099	117	101	16
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	2,775	1,894	881	6,051	5,460	591	77	69	8
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	2,260	1,373	887	1,628	1,120	508	40	32	8
Legal intervention (Y35,Y89.0)	220	212	8	97	95	2	2	2	-
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	3,783	2,194	1,589	587	377	210	28	21	7
Discharge of firearms, undetermined intent (Y22-Y24)	186	151	35	32	25	7	-	-	-
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	3,597	2,043	1,554	555	352	203	28	21	7
Operations of war and their sequelae (Y36,Y89.1)	7	7	-	1	1	-	-	-	-
Complications of medical and surgical care (Y40-Y84,Y88)	1,875	888	987	398	169	229	5	3	2
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	6,298	2,471	3,827	476	196	280	10	5	5
Drug-induced deaths ^{5,6}	33,145	19,689	13,456	3,502	2,170	1,332	185	129	56
Alcohol-induced deaths ^{5,7}	18,806	13,735	5,071	2,296	1,661	635	124	101	23
Injury by firearms ^{5,8}	20,513	17,350	3,163	7,330	6,607	723	145	130	15

- Quantity zero.

... Category not applicable.

¹Includes races other than white and black.

²Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

³Includes deaths for which Hispanic origin was not reported on the death certificate.

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

Table 13. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearm, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ²			Non-Hispanic black ²			Origin not stated ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	799.5	812.0	787.4	861.7	866.1	857.3	682.2	725.4	642.7
Salmonella infections (A01-A02)	0.0	*	*	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	3.3	2.7	4.0	3.8	3.0	4.6	2.0	1.8	2.1
Tuberculosis (A16-A19)	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
Respiratory tuberculosis (A16)	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.4	0.1
Other tuberculosis (A17-A19)	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Whooping cough (A37)	0.0	*	*	0.0	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Septicemia (A40-A41)	11.3	10.6	11.9	11.4	10.7	12.1	14.3	13.4	15.1
Syphilis (A50-A53)	0.0	0.0	*	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.4	3.3	1.6	2.5	3.4	1.6	2.6	3.5	1.8
Human immunodeficiency virus (HIV) disease (B20-B24)	2.7	4.0	1.4	1.5	2.4	0.5	11.1	15.2	7.4
Malaria (B50-B54)	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.9	2.0	1.8	2.0	2.0	1.9	1.8	2.0	1.7
Malignant neoplasms (C00-C97)	186.2	198.3	174.4	200.3	212.7	188.2	156.7	169.0	145.5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	2.7	3.8	1.7	2.9	4.0	1.8	2.4	3.6	1.3
Malignant neoplasm of esophagus (C15)	4.7	7.5	2.0	5.2	8.4	2.1	3.4	5.2	1.8
Malignant neoplasm of stomach (C16)	3.7	4.4	3.0	3.5	4.1	2.8	4.8	5.9	3.7
Malignant neoplasms of colon, rectum and anus (C18-C21)	17.0	18.0	16.1	17.9	18.8	17.0	16.7	18.0	15.4
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	6.6	9.0	4.2	6.5	8.8	4.3	6.7	9.9	3.8
Malignant neoplasm of pancreas (C25)	11.9	12.3	11.6	12.8	13.2	12.4	10.3	10.4	10.2
Malignant neoplasm of larynx (C32)	1.2	1.9	0.5	1.2	2.0	0.5	1.5	2.4	0.6
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	51.3	57.8	45.0	56.1	62.3	50.0	39.7	48.7	31.4
Malignant melanoma of skin (C43)	3.0	4.0	2.0	3.6	4.9	2.5	0.3	0.3	0.3

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of breast (C50)	13.4	0.3	26.1	13.9	0.3	27.3	14.5	0.4	27.5
Malignant neoplasm of cervix uteri (C53)	1.3	...	2.5	1.2	...	2.4	1.9	...	3.6
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	2.7	...	5.4	2.7	...	5.4	3.4	...	6.5
Malignant neoplasm of ovary (C56)	4.7	...	9.3	5.2	...	10.4	3.0	...	5.8
Malignant neoplasm of prostate (C61)	9.3	18.8	...	9.4	19.1	...	11.5	24.1	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.3	5.6	3.0	4.8	6.2	3.4	2.8	3.5	2.2
Malignant neoplasm of bladder (C67)	4.8	6.9	2.7	5.5	7.9	3.0	2.6	3.1	2.1
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	4.6	5.3	3.9	5.2	6.0	4.5	2.1	2.3	2.0
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	18.0	20.3	15.8	19.8	22.3	17.3	13.0	14.0	12.0
Hodgkin's disease (C81)	0.4	0.5	0.3	0.4	0.5	0.4	0.3	0.3	0.2
Non-Hodgkin's lymphoma (C82-C85)	6.6	7.3	5.9	7.4	8.2	6.7	3.6	4.1	3.1
Leukemia (C91-C95)	7.3	8.5	6.2	8.2	9.5	6.9	4.3	4.8	3.9
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.7	4.0	3.4	3.7	4.1	3.4	4.8	4.9	4.7
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.0	0.0	0.0	0.0	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	21.0	22.5	19.5	22.8	24.5	21.2	16.2	17.1	15.4
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.8	5.1	4.6	5.4	5.7	5.1	2.9	2.7	3.1
Anemias (D50-D64)	1.6	1.3	1.8	1.5	1.3	1.8	2.4	2.2	2.6
Diabetes mellitus (E10-E14)	22.4	23.4	21.4	22.1	23.5	20.8	28.8	28.1	29.5
Nutritional deficiencies (E40-E64)	1.0	0.8	1.2	1.0	0.8	1.2	0.9	0.7	1.1
Malnutrition (E40-E46)	0.9	0.7	1.1	1.0	0.8	1.2	0.8	0.6	1.0
Other nutritional deficiencies (E50-E64)	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Meningitis (G00,G03)	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
Parkinson's disease (G20-G21)	7.1	8.5	5.8	8.5	10.0	6.9	1.9	2.2	1.6
Alzheimer's disease (G30)	27.0	16.7	37.0	31.3	19.3	43.1	12.4	7.4	17.0
Major cardiovascular diseases (I00-I78)	252.7	252.7	252.7	271.8	269.7	273.9	220.8	226.7	215.5
Diseases of heart (I00-I09,I11,I13,I20-I51)	193.6	202.5	184.9	209.6	217.8	201.5	164.2	174.6	154.8
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	1.0	0.7	1.3	1.1	0.7	1.4	0.6	0.4	0.7

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart disease (I11)	10.9	10.7	11.1	10.2	9.7	10.8	18.4	20.3	16.6
Hypertensive heart and renal disease (I13)	0.9	0.8	1.0	0.8	0.7	0.9	1.8	1.9	1.7
Ischemic heart diseases (I20-I25)	122.9	136.8	109.6	134.6	149.4	120.1	94.2	102.6	86.6
Acute myocardial infarction (I21-I22)	39.5	44.4	34.8	43.3	48.7	37.9	30.3	32.1	28.7
Other acute ischemic heart diseases (I24)	1.4	1.4	1.3	1.4	1.5	1.4	1.3	1.5	1.2
Other forms of chronic ischemic heart disease (I20,I25)	82.0	90.9	73.5	89.9	99.2	80.8	62.6	69.0	56.7
Atherosclerotic cardiovascular disease, so described (I25.0)	18.6	22.2	15.1	19.3	22.9	15.9	19.9	24.5	15.6
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	63.4	68.7	58.4	70.5	76.3	64.9	42.7	44.5	41.1
Other heart diseases (I26-I51)	57.9	53.6	62.0	62.9	57.3	68.4	49.3	49.5	49.2
Acute and subacute endocarditis (I33)	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.4
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3
Heart failure (I50)	18.7	16.1	21.3	20.9	17.7	24.0	13.1	12.2	14.0
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	38.6	36.8	40.2	41.4	38.9	43.8	35.4	36.5	34.5
Essential hypertension and hypertensive renal disease (I10,I12,I15)	8.6	7.1	10.1	8.4	6.8	9.9	12.2	11.0	13.2
Cerebrovascular diseases (I60-I69)	41.9	34.5	49.1	44.5	35.8	53.0	38.0	34.5	41.1
Atherosclerosis (I70)	2.3	1.9	2.7	2.7	2.2	3.1	1.4	1.3	1.5
Other diseases of circulatory system (I71-I78)	6.2	6.6	5.8	6.7	7.2	6.4	5.1	5.3	4.9
Aortic aneurysm and dissection (I71)	3.4	4.0	2.8	3.7	4.4	3.0	2.3	2.7	1.9
Other diseases of arteries, arterioles and capillaries (I72-I78)	2.8	2.6	3.1	3.0	2.7	3.3	2.8	2.6	2.9
Other disorders of circulatory system (I80-I99)	1.4	1.4	1.4	1.4	1.3	1.4	1.8	1.9	1.7
Influenza and pneumonia (J09-J18)	16.2	15.6	16.9	17.6	16.7	18.6	11.7	11.8	11.6
Influenza (J09-J11)	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Pneumonia (J12-J18)	16.1	15.4	16.7	17.5	16.5	18.4	11.6	11.7	11.5
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Acute bronchitis and bronchiolitis (J20-J21)	0.1	0.0	0.1	0.1	0.1	0.1	0.0	*	*
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	0.0	*	0.0	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	44.7	43.1	46.3	51.8	49.1	54.5	20.7	22.5	19.0
Bronchitis, chronic and unspecified (J40-J42)	0.2	0.2	0.2	0.2	0.2	0.3	0.1	0.1	0.1
Emphysema (J43)	3.2	3.4	3.1	3.8	3.9	3.7	1.4	1.8	1.1
Asthma (J45-J46)	1.1	0.8	1.4	1.0	0.7	1.3	2.1	2.0	2.1

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other chronic lower respiratory diseases (J44,J47)	40.2	38.6	41.7	46.8	44.4	49.3	17.1	18.6	15.7
Pneumoconioses and chemical effects (J60-J66,J68)	0.3	0.5	0.0	0.3	0.6	0.0	0.1	0.2	*
Pneumonitis due to solids and liquids (J69)	5.5	6.1	5.0	6.2	6.8	5.6	3.3	3.6	3.0
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	10.1	10.4	9.8	11.2	11.5	10.8	7.0	6.9	7.0
Peptic ulcer (K25-K28)	1.0	1.0	1.0	1.0	1.0	1.1	0.7	0.9	0.6
Diseases of appendix (K35-K38)	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	*
Hernia (K40-K46)	0.6	0.5	0.6	0.7	0.6	0.7	0.4	0.4	0.4
Chronic liver disease and cirrhosis (K70,K73-K74)	10.3	13.7	7.1	11.4	15.1	7.8	6.3	8.5	4.2
Alcoholic liver disease (K70)	5.2	7.5	2.9	5.7	8.3	3.1	2.9	4.1	1.8
Other chronic liver disease and cirrhosis (K73-K74)	5.2	6.2	4.2	5.7	6.8	4.7	3.3	4.4	2.3
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.1	1.0	1.1	1.2	1.1	1.2	0.7	0.6	0.8
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16.3	16.4	16.3	16.4	16.6	16.2	21.0	20.0	22.0
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.9	1.8	2.0	1.9	1.9	2.0	2.3	2.1	2.3
Renal failure (N17-N19)	14.4	14.5	14.3	14.4	14.7	14.1	18.7	17.7	19.5
Other disorders of kidney (N25,N27)	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	0.1	0.3	0.2	0.1	0.3	0.2	0.1	0.2
Hyperplasia of prostate (N40)	0.2	0.3	...	0.2	0.4	...	0.1	0.2	...
Inflammatory diseases of female pelvic organs (N70-N76)	0.0	...	0.1	0.0	...	0.1	0.0	...	0.1
Pregnancy, childbirth and the puerperium (O00-O99)	0.3	...	0.5	0.2	...	0.4	0.6	...	1.2
Pregnancy with abortive outcome (O00-O07)	0.0	...	0.0	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.3	...	0.5	0.2	...	0.4	0.6	...	1.1
Certain conditions originating in the perinatal period (P00-P96)	3.9	4.5	3.4	3.1	3.5	2.6	9.6	11.0	8.3
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.1	3.3	3.0	3.1	3.2	3.0	3.9	4.3	3.6
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	12.4	10.6	14.2	13.3	11.0	15.6	11.7	11.7	11.8
All other diseases (Residual)	87.4	72.2	102.1	95.9	78.2	113.3	67.8	59.2	75.7

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidents (unintentional injuries) (V01-X59, Y85-Y86)	39.1	50.0	28.6	42.8	53.8	31.9	28.7	40.2	18.2
Transport accidents (V01-V99, Y85)	12.3	17.6	7.1	12.9	18.4	7.5	11.3	17.0	6.1
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	11.4	16.3	6.8	12.0	17.0	7.1	10.6	15.9	5.8
Other land transport accidents (V01, V05-V06, V09.1, V09.3-V09.9, V10-V11, V15-V18, V19.3, V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9)	0.3	0.5	0.1	0.3	0.5	0.2	0.3	0.5	0.1
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99, Y85)	0.5	0.8	0.2	0.6	0.9	0.2	0.3	0.5	0.1
Nontransport accidents (W00-X59, Y86)	26.8	32.4	21.5	29.9	35.4	24.4	17.4	23.1	12.1
Falls (W00-W19)	8.4	8.6	8.3	9.8	9.9	9.8	2.8	3.3	2.3
Accidental discharge of firearms (W32-W34)	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	*
Accidental drowning and submersion (W65-W74)	1.2	1.9	0.5	1.2	1.9	0.6	1.4	2.4	0.4
Accidental exposure to smoke, fire and flames (X00-X09)	0.9	1.1	0.7	0.9	1.0	0.7	1.4	1.7	1.1
Accidental poisoning and exposure to noxious substances (X40-X49)	10.7	13.9	7.6	11.9	15.4	8.5	7.1	9.4	5.0
Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64, W75-W99, X10-X39, X50-X59, Y86)	5.4	6.5	4.3	5.8	6.9	4.8	4.5	5.8	3.3
Intentional self-harm (suicide) (*U03, X60-X84, Y87.0)	12.4	19.9	5.2	14.1	22.6	5.9	5.1	8.7	1.8
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	6.3	11.2	1.5	7.3	12.9	1.8	2.6	4.8	0.5
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60-X71, X75-X84, Y87.0)	6.1	8.8	3.6	6.8	9.7	4.0	2.5	3.9	1.3
Assault (homicide) (*U01-*U02, X85-Y09, Y87.1)	5.3	8.4	2.2	3.2	4.7	1.8	18.6	33.4	5.1
Assault (homicide) by discharge of firearms (*U01.4, X93-X95)	3.6	6.2	1.1	1.9	2.9	0.9	14.6	27.6	2.7
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3, *U01.5-*U01.9, *U02, X85-X92, X96-Y09, Y87.1)	1.7	2.3	1.1	1.3	1.7	0.9	4.0	5.7	2.3
Legal intervention (Y35, Y89.0)	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.6	1.9	1.3	1.7	2.0	1.4	1.4	1.9	1.0
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34, Y87.2,Y89.9)	1.5	1.8	1.2	1.6	1.9	1.3	1.3	1.8	0.9
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.8	0.8	0.8	0.8	0.8	0.8	1.0	0.8	1.1
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	2.4	1.9	2.8	2.7	2.2	3.3	1.2	1.0	1.3
Drug-induced deaths ^{5,6}	13.1	16.1	10.2	14.7	17.9	11.5	8.5	11.0	6.2
Alcohol-induced deaths ^{5,7}	8.3	12.5	4.2	9.0	13.6	4.5	5.5	8.4	2.9
Injury by firearms ^{5,8}	10.3	18.0	2.7	9.6	16.5	2.8	17.7	33.4	3.3

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
All causes	365.1	397.5	332.4	301.1	327.0	277.3
Salmonella infections (A01-A02)	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	1.5	1.4	1.5	0.8	0.8	0.8
Tuberculosis (A16-A19)	*	*	*	0.6	0.7	0.4
Respiratory tuberculosis (A16)	*	*	*	0.5	0.6	0.3
Other tuberculosis (A17-A19)	*	*	*	*	*	*
Whooping cough (A37)	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*
Meningococcal infection (A39)	*	*	*	*	*	*
Septicemia (A40-A41)	5.7	4.9	6.5	3.4	3.6	3.2
Syphilis (A50-A53)	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.3	2.9	1.7	1.8	2.1	1.5
Human immunodeficiency virus (HIV) disease (B20-B24)	1.4	2.2	*	0.4	0.7	*
Malaria (B50-B54)	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.2	1.3	1.2	1.0	1.2	0.9
Malignant neoplasms (C00-C97)	69.5	74.1	64.8	83.4	88.6	78.5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	1.0	1.2	*	1.6	2.2	1.1
Malignant neoplasm of esophagus (C15)	1.5	2.3	*	1.3	2.2	0.6
Malignant neoplasm of stomach (C16)	2.0	2.7	1.4	4.6	5.4	3.8
Malignant neoplasms of colon, rectum and anus (C18-C21)	6.7	7.1	6.4	8.7	9.2	8.2
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	4.6	6.7	2.5	7.9	11.1	4.9
Malignant neoplasm of pancreas (C25)	4.2	4.5	3.8	5.7	5.4	6.0
Malignant neoplasm of larynx (C32)	0.6	0.9	*	0.3	0.5	*
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	18.4	20.8	16.0	18.5	22.5	14.9
Malignant melanoma of skin (C43)	0.5	*	*	0.3	0.3	0.3

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of breast (C50)	4.0	*	8.0	5.7	*	10.8
Malignant neoplasm of cervix uteri (C53)	0.8	...	1.7	0.8	...	1.6
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	0.8	...	1.7	1.4	...	2.7
Malignant neoplasm of ovary (C56)	1.7	...	3.4	2.2	...	4.3
Malignant neoplasm of prostate (C61)	2.7	5.5	...	2.5	5.1	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	2.6	3.5	1.7	1.6	2.2	1.0
Malignant neoplasm of bladder (C67)	1.1	1.4	*	1.2	1.7	0.8
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	1.7	1.8	1.7	1.9	2.1	1.7
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	5.4	6.0	4.8	7.8	9.1	6.6
Hodgkin's disease (C81)	*	*	*	0.1	*	*
Non-Hodgkin's lymphoma (C82-C85)	1.6	1.9	1.4	3.0	3.4	2.7
Leukemia (C91-C95)	2.5	2.8	2.3	3.3	3.8	2.8
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	1.1	1.2	1.0	1.3	1.6	1.0
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	9.0	9.2	8.9	9.3	9.6	9.1
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1.4	1.5	1.3	2.1	2.1	2.0
Anemias (D50-D64)	*	*	*	0.5	0.5	0.5
Diabetes mellitus (E10-E14)	20.1	20.2	20.0	10.8	11.5	10.2
Nutritional deficiencies (E40-E64)	0.7	*	0.9	0.3	0.3	0.3
Malnutrition (E40-E46)	0.7	*	*	0.3	0.3	0.3
Other nutritional deficiencies (E50-E64)	*	*	*	*	*	*
Meningitis (G00,G03)	*	*	*	0.1	*	*
Parkinson's disease (G20-G21)	1.5	1.9	1.0	2.2	2.5	1.9
Alzheimer's disease (G30)	6.2	4.2	8.3	6.4	4.2	8.3
Major cardiovascular diseases (I00-I78)	84.5	92.0	76.9	97.9	106.0	90.4
Diseases of heart (I00-I09,I11,I13,I20-I51)	65.5	75.0	55.9	67.6	77.0	59.0
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	*	*	*	0.6	0.3	0.8

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart disease (I11)	4.0	4.7	3.3	4.0	4.2	3.9
Hypertensive heart and renal disease (I13)	*	*	*	0.5	0.4	0.5
Ischemic heart diseases (I20-I25)	42.9	51.2	34.6	46.0	55.1	37.7
Acute myocardial infarction (I21-I22)	13.9	17.1	10.7	14.9	17.7	12.3
Other acute ischemic heart diseases (I24)	0.9	1.1	*	0.3	0.4	0.3
Other forms of chronic ischemic heart disease (I20,I25)	28.1	33.0	23.2	30.8	37.0	25.2
Atherosclerotic cardiovascular disease, so described (I25.0)	8.7	10.9	6.6	7.3	9.2	5.5
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	19.4	22.1	16.6	23.5	27.8	19.7
Other heart diseases (I26-I51)	17.8	18.4	17.3	16.5	17.0	16.1
Acute and subacute endocarditis (I33)	*	*	*	0.1	*	*
Diseases of pericardium and acute myocarditis (I30-I31,I40)	*	*	*	0.1	*	*
Heart failure (I50)	5.3	4.2	6.3	4.2	3.8	4.6
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	12.1	13.6	10.6	12.1	12.8	11.4
Essential hypertension and hypertensive renal disease (I10,I12,I15)	3.4	2.9	3.9	4.8	4.1	5.4
Cerebrovascular diseases (I60-I69)	13.1	12.0	14.2	22.6	21.5	23.5
Atherosclerosis (I70)	*	*	*	0.6	0.7	0.5
Other diseases of circulatory system (I71-I78)	2.1	1.8	2.3	2.3	2.7	1.9
Aortic aneurysm and dissection (I71)	1.1	1.2	1.0	1.6	2.0	1.3
Other diseases of arteries, arterioles and capillaries (I72-I78)	0.9	*	1.3	0.7	0.7	0.6
Other disorders of circulatory system (I80-I99)	0.5	*	*	0.5	0.6	0.4
Influenza and pneumonia (J09-J18)	7.6	8.0	7.3	9.1	10.1	8.1
Influenza (J09-J11)	*	*	*	*	*	*
Pneumonia (J12-J18)	7.6	7.9	7.2	8.9	10.0	8.0
Other acute lower respiratory infections (J20-J22,U04)	*	*	*	*	*	*
Acute bronchitis and bronchiolitis (J20-J21)	*	*	*	*	*	*
Other and unspecified acute lower respiratory infections (J22,U04)	*	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	16.5	16.3	16.6	8.8	11.2	6.5
Bronchitis, chronic and unspecified (J40-J42)	*	*	*	*	*	*
Emphysema (J43)	1.0	1.0	1.1	0.6	1.0	0.2

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Asthma (J45-J46)	0.5	*	*	0.8	0.9	0.7
Other chronic lower respiratory diseases (J44,J47)	14.8	14.6	15.0	7.3	9.3	5.5
Pneumoconioses and chemical effects (J60-J66,J68)	*	*	*	*	*	*
Pneumonitis due to solids and liquids (J69)	1.9	1.9	2.0	1.8	2.2	1.5
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	5.6	5.3	5.9	3.8	4.1	3.6
Peptic ulcer (K25-K28)	0.5	*	*	0.5	0.6	0.5
Diseases of appendix (K35-K38)	*	*	*	*	*	*
Hernia (K40-K46)	*	*	*	*	*	*
Chronic liver disease and cirrhosis (K70,K73-K74)	18.5	20.0	16.9	2.7	3.7	1.9
Alcoholic liver disease (K70)	13.9	16.4	11.4	1.0	1.8	0.4
Other chronic liver disease and cirrhosis (K73-K74)	4.6	3.6	5.5	1.7	2.0	1.5
Cholelithiasis and other disorders of gallbladder (K80-K82)	0.5	*	*	0.6	0.6	0.5
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8.0	7.1	8.8	6.4	6.4	6.4
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	*	*	*	*	*	*
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.0	0.9	1.0	0.8	0.7	0.9
Renal failure (N17-N19)	7.0	6.2	7.7	5.6	5.7	5.4
Other disorders of kidney (N25,N27)	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	*	*	*	*	*	*
Hyperplasia of prostate (N40)	*	*	...	*	*	...
Inflammatory diseases of female pelvic organs (N70-N76)	*	...	*	*	...	*
Pregnancy, childbirth and the puerperium (O00-O99)	*	...	*	0.2	...	0.5
Pregnancy with abortive outcome (O00-O07)	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	*	...	*	0.2	...	0.5
Certain conditions originating in the perinatal period (P00-P96)	2.7	3.5	1.9	2.7	3.4	2.1
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.0	3.3	2.8	1.9	1.9	1.9
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	5.6	6.0	5.2	3.0	3.1	2.9

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
All other diseases (Residual)	36.3	33.8	38.9	25.4	24.0	26.6
Accidents (unintentional injuries) (V01-X59, Y85-Y86)	39.9	53.7	26.0	12.6	16.4	9.1
Transport accidents (V01-V99, Y85)	16.0	21.6	10.5	5.2	6.8	3.8
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	15.0	19.8	10.0	4.9	6.2	3.6
Other land transport accidents (V01, V05-V06, V09.1, V09.3-V09.9, V10-V11, V15-V18, V19.3, V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9)	0.5	*	*	0.2	0.3	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99, Y85)	0.6	*	*	0.2	0.3	*
Nontransport accidents (W00-X59, Y86)	23.9	32.1	15.5	7.4	9.7	5.3
Falls (W00-W19)	3.8	4.5	3.0	3.2	3.9	2.7
Accidental discharge of firearms (W32-W34)	*	*	*	*	*	*
Accidental drowning and submersion (W65-W74)	1.6	2.8	*	1.0	1.6	0.5
Accidental exposure to smoke, fire and flames (X00-X09)	1.1	1.2	*	0.1	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)	12.2	15.5	8.9	1.5	2.1	0.9
Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64, W75-W99, X10-X39, X50-X59, Y86)	4.9	7.4	2.3	1.5	1.8	1.2
Intentional self-harm (suicide) (*U03, X60-X84, Y87.0)	11.0	16.1	5.9	6.2	9.3	3.4
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	4.2	6.8	1.5	1.3	2.5	0.3
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60-X71, X75-X84, Y87.0)	6.8	9.2	4.4	4.9	6.8	3.2
Assault (homicide) (*U01-*U02, X85-Y09, Y87.1)	6.0	9.5	2.5	1.9	2.7	1.2
Assault (homicide) by discharge of firearms (*U01.4, X93-X95)	2.7	4.6	*	1.0	1.6	0.4
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3, *U01.5-*U01.9, *U02, X85-X92, X96-Y09, Y87.1)	3.4	4.9	1.8	0.9	1.0	0.8

Table 14. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Legal intervention (Y35,Y89.0)	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	2.0	2.1	1.9	0.5	0.7	0.3
Discharge of firearms, undetermined intent (Y22-Y24)	*	*	*	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.9	2.0	1.8	0.5	0.6	0.3
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.5	*	*	0.3	*	0.3
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	0.9	0.9	*	0.6	0.5	0.6
Drug-induced deaths ^{5,6}	10.7	11.7	9.8	2.1	2.7	1.5
Alcohol-induced deaths ^{5,7}	21.8	28.5	15.1	1.6	2.6	0.6
Injury by firearms ^{5,8}	7.4	12.5	2.4	2.4	4.4	0.6

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	799.5	812.0	787.4	286.2	310.8	260.9	897.6	911.1	884.7
Salmonella infections (A01-A02)	0.0	*	*	*	*	*	0.0	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	3.3	2.7	4.0	1.2	1.1	1.3	3.7	3.0	4.5
Tuberculosis (A16-A19)	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
Respiratory tuberculosis (A16)	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1
Other tuberculosis (A17-A19)	0.0	0.0	0.0	0.0	*	*	0.0	0.0	0.0
Whooping cough (A37)	0.0	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	0.0	0.0	0.0	*	*	0.0	0.0	0.0
Septicemia (A40-A41)	11.3	10.6	11.9	4.0	4.0	4.1	12.7	11.9	13.4
Syphilis (A50-A53)	0.0	0.0	*	*	*	*	0.0	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.4	3.3	1.6	2.2	3.0	1.4	2.5	3.4	1.6
Human immunodeficiency virus (HIV) disease (B20-B24)	2.7	4.0	1.4	2.2	3.6	0.9	2.8	4.1	1.5
Malaria (B50-B54)	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.9	2.0	1.8	0.9	0.9	0.8	2.1	2.2	2.0
Malignant neoplasms (C00-C97)	186.2	198.3	174.4	61.6	64.2	59.0	210.1	225.1	195.7
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	2.7	3.8	1.7	0.8	1.2	0.5	3.1	4.4	1.9
Malignant neoplasm of esophagus (C15)	4.7	7.5	2.0	1.2	1.9	0.5	5.4	8.7	2.2
Malignant neoplasm of stomach (C16)	3.7	4.4	3.0	3.1	3.4	2.7	3.8	4.6	3.0
Malignant neoplasms of colon, rectum and anus (C18-C21)	17.0	18.0	16.1	6.3	7.0	5.6	19.1	20.2	18.1
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	6.6	9.0	4.2	4.6	6.2	3.0	6.9	9.5	4.5
Malignant neoplasm of pancreas (C25)	11.9	12.3	11.6	4.2	4.1	4.4	13.4	14.0	12.9
Malignant neoplasm of larynx (C32)	1.2	1.9	0.5	0.4	0.7	0.1	1.3	2.2	0.5

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	51.3	57.8	45.0	9.8	11.9	7.7	59.3	67.0	51.9
Malignant melanoma of skin (C43)	3.0	4.0	2.0	0.4	0.5	0.4	3.5	4.7	2.3
Malignant neoplasm of breast (C50)	13.4	0.3	26.1	4.6	*	9.2	15.1	0.3	29.2
Malignant neoplasm of cervix uteri (C53)	1.3	...	2.5	0.9	...	1.9	1.3	...	2.6
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	2.7	...	5.4	1.0	...	2.1	3.0	...	5.9
Malignant neoplasm of ovary (C56)	4.7	...	9.3	1.7	...	3.4	5.3	...	10.4
Malignant neoplasm of prostate (C61)	9.3	18.8	...	3.0	6.0	...	10.4	21.4	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.3	5.6	3.0	1.9	2.3	1.4	4.7	6.2	3.3
Malignant neoplasm of bladder (C67)	4.8	6.9	2.7	1.0	1.4	0.7	5.5	8.0	3.1
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	4.6	5.3	3.9	1.8	2.0	1.5	5.1	5.9	4.4
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	18.0	20.3	15.8	7.2	7.7	6.7	20.1	22.8	17.5
Hodgkin's disease (C81)	0.4	0.5	0.3	0.3	0.3	0.2	0.4	0.5	0.4
Non-Hodgkin's lymphoma (C82-C85)	6.6	7.3	5.9	2.6	2.8	2.5	7.3	8.2	6.5
Leukemia (C91-C95)	7.3	8.5	6.2	2.9	3.1	2.7	8.2	9.6	6.8
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.7	4.0	3.4	1.4	1.5	1.4	4.1	4.5	3.8
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52,C57-C60, C62-C63,C66,C68-C69,C73-C80,C97)	21.0	22.5	19.5	7.6	8.0	7.2	23.6	25.4	21.8
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.8	5.1	4.6	1.6	1.6	1.5	5.5	5.8	5.2
Anemias (D50-D64)	1.6	1.3	1.8	0.5	0.5	0.5	1.8	1.5	2.1
Diabetes mellitus (E10-E14)	22.4	23.4	21.4	13.0	13.2	12.8	24.1	25.4	23.0
Nutritional deficiencies (E40-E64)	1.0	0.8	1.2	0.3	0.3	0.4	1.1	0.8	1.3
Malnutrition (E40-E46)	0.9	0.7	1.1	0.3	0.3	0.4	1.0	0.8	1.2

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other nutritional deficiencies (E50-E64)	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1
Meningitis (G00,G03)	0.2	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.2
Parkinson's disease (G20-G21)	7.1	8.5	5.8	1.9	2.2	1.7	8.1	9.8	6.6
Alzheimer's disease (G30)	27.0	16.7	37.0	6.8	4.6	9.1	31.0	19.2	42.3
Major cardiovascular diseases (I00-I78)	252.7	252.7	252.7	79.5	82.8	76.2	285.8	286.3	285.3
Diseases of heart									
(I00-I09,I11,I13,I20-I51)	193.6	202.5	184.9	59.4	64.1	54.6	219.2	229.9	209.0
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	1.0	0.7	1.3	0.3	0.2	0.4	1.1	0.7	1.4
Hypertensive heart disease (I11)	10.9	10.7	11.1	4.1	4.6	3.7	12.2	11.9	12.4
Hypertensive heart and renal disease (I13)	0.9	0.8	1.0	0.4	0.4	0.4	1.0	0.9	1.1
Ischemic heart diseases (I20-I25)	122.9	136.8	109.6	40.6	45.1	36.0	138.6	154.8	123.1
Acute myocardial infarction (I21-I22)	39.5	44.4	34.8	13.0	14.5	11.4	44.6	50.4	39.1
Other acute ischemic heart diseases (I24)	1.4	1.4	1.3	0.3	0.2	0.3	1.6	1.7	1.5
Other forms of chronic ischemic heart disease (I20,I25)	82.0	90.9	73.5	27.4	30.3	24.3	92.4	102.8	82.5
Atherosclerotic cardiovascular disease, so described (I25.0)	18.6	22.2	15.1	6.8	8.8	4.7	20.8	24.7	17.0
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	63.4	68.7	58.4	20.6	21.5	19.6	71.6	78.1	65.5
Other heart diseases (I26-I51)	57.9	53.6	62.0	14.0	13.8	14.1	66.3	61.5	71.0
Acute and subacute endocarditis (I33)	0.4	0.4	0.3	0.2	0.2	0.1	0.4	0.5	0.3
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.3	0.3	0.2	0.1	0.1	0.1	0.3	0.3	0.3
Heart failure (I50)	18.7	16.1	21.3	4.0	3.7	4.4	21.6	18.6	24.4
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	38.6	36.8	40.2	9.7	9.9	9.5	44.1	42.2	45.9
Essential hypertension and hypertensive renal disease (I10,I12,I15)	8.6	7.1	10.1	3.4	3.0	3.8	9.6	8.0	11.2
Cerebrovascular diseases (I60-I69)	41.9	34.5	49.1	14.4	13.2	15.7	47.2	38.8	55.3
Atherosclerosis (I70)	2.3	1.9	2.7	0.5	0.5	0.6	2.7	2.2	3.1
Other diseases of circulatory system (I71-I78)	6.2	6.6	5.8	1.8	2.1	1.5	7.1	7.5	6.6
Aortic aneurysm and dissection (I71)	3.4	4.0	2.8	0.9	1.2	0.6	3.9	4.6	3.2

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of arteries, arterioles and capillaries (I72-I78)	2.8	2.6	3.1	0.9	0.9	0.9	3.2	2.9	3.5
Other disorders of circulatory system (I80-I99)	1.4	1.4	1.4	0.5	0.5	0.5	1.5	1.5	1.6
Influenza and pneumonia (J09-J18)	16.2	15.6	16.9	6.0	6.1	5.9	18.2	17.4	18.9
Influenza (J09-J11)	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Pneumonia (J12-J18)	16.1	15.4	16.7	5.8	5.9	5.7	18.0	17.3	18.7
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1
Acute bronchitis and bronchiolitis (J20-J21)	0.1	0.0	0.1	*	*	*	0.1	0.1	0.1
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	0.0	*	*	*	*	0.0	*	*
Chronic lower respiratory diseases (J40-J47)	44.7	43.1	46.3	8.3	8.5	8.0	51.7	50.0	53.4
Bronchitis, chronic and unspecified (J40-J42)	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.2	0.2
Emphysema (J43)	3.2	3.4	3.1	0.5	0.6	0.4	3.8	4.0	3.6
Asthma (J45-J46)	1.1	0.8	1.4	0.6	0.5	0.7	1.2	0.9	1.5
Other chronic lower respiratory diseases (J44,J47)	40.2	38.6	41.7	7.1	7.3	6.9	46.5	44.9	48.1
Pneumoconioses and chemical effects (J60-J66,J68)	0.3	0.5	0.0	0.0	0.1	*	0.3	0.6	0.0
Pneumonitis due to solids and liquids (J69)	5.5	6.1	5.0	1.4	1.5	1.2	6.3	7.0	5.7
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	10.1	10.4	9.8	4.0	4.0	4.1	11.3	11.7	10.9
Peptic ulcer (K25-K28)	1.0	1.0	1.0	0.3	0.3	0.3	1.1	1.1	1.1
Diseases of appendix (K35-K38)	0.1	0.2	0.1	0.1	0.1	*	0.1	0.2	0.1
Hernia (K40-K46)	0.6	0.5	0.6	0.2	0.2	0.3	0.7	0.6	0.7
Chronic liver disease and cirrhosis (K70,K73-K74)	10.3	13.7	7.1	8.6	12.0	5.2	10.6	14.0	7.4
Alcoholic liver disease (K70)	5.2	7.5	2.9	4.7	7.6	1.8	5.2	7.5	3.1
Other chronic liver disease and cirrhosis (K73-K74)	5.2	6.2	4.2	3.9	4.4	3.3	5.4	6.5	4.3
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.1	1.0	1.1	0.5	0.5	0.5	1.2	1.1	1.3
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16.3	16.4	16.3	6.4	6.5	6.4	18.2	18.3	18.2

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.9	1.8	2.0	0.6	0.6	0.7	2.1	2.1	2.2
Renal failure (N17-N19)	14.4	14.5	14.3	5.8	5.9	5.7	16.0	16.2	15.8
Other disorders of kidney (N25,N27)	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	0.1	0.3	0.1	*	0.1	0.2	0.1	0.3
Hyperplasia of prostate (N40)	0.2	0.3	...	0.1	0.1	...	0.2	0.4	...
Inflammatory diseases of female pelvic organs (N70-N76)	0.0	...	0.1	*	...	*	0.0	...	0.1
Pregnancy, childbirth and the puerperium (O00-O99)	0.3	...	0.5	0.3	...	0.6	0.3	...	0.5
Pregnancy with abortive outcome (O00-O07)	0.0	...	0.0	*	...	*	0.0	...	0.0
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.3	...	0.5	0.3	...	0.6	0.2	...	0.5
Certain conditions originating in the perinatal period (P00-P96)	3.9	4.5	3.4	5.0	5.7	4.3	3.7	4.2	3.2
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.1	3.3	3.0	3.5	3.5	3.6	3.0	3.2	2.9
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	12.4	10.6	14.2	4.1	4.2	4.0	14.0	11.8	16.1
All other diseases (Residual)	87.4	72.2	102.1	27.7	25.6	29.8	98.9	81.4	115.6
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39.1	50.0	28.6	20.8	29.6	11.6	42.6	53.9	31.7
Transport accidents (V01-V99,Y85)	12.3	17.6	7.1	9.5	13.7	5.1	12.8	18.4	7.5
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)	11.4	16.3	6.8	8.9	12.8	4.9	11.9	16.9	7.1

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9, V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	0.3	0.5	0.1	0.3	0.6	0.1	0.3	0.5	0.2
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.5	0.8	0.2	0.2	0.3	*	0.6	0.9	0.2
Nontransport accidents (W00-X59,Y86)	26.8	32.4	21.5	11.3	15.9	6.5	29.8	35.6	24.3
Falls (W00-W19)	8.4	8.6	8.3	2.5	3.1	1.9	9.6	9.7	9.4
Accidental discharge of firearms (W32-W34)	0.2	0.3	0.1	0.1	0.1	*	0.2	0.4	0.1
Accidental drowning and submersion (W65-W74)	1.2	1.9	0.5	1.0	1.7	0.3	1.3	2.0	0.6
Accidental exposure to smoke, fire and flames (X00-X09)	0.9	1.1	0.7	0.3	0.4	0.2	1.0	1.2	0.8
Accidental poisoning and exposure to noxious substances (X40-X49)	10.7	13.9	7.6	5.2	7.5	2.7	11.7	15.1	8.5
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39, X50-X59,Y86)	5.4	6.5	4.3	2.2	3.0	1.3	6.0	7.2	4.9
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12.4	19.9	5.2	5.3	8.5	2.0	13.8	22.2	5.7
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	6.3	11.2	1.5	1.9	3.4	0.4	7.1	12.7	1.8
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)	6.1	8.8	3.6	3.4	5.1	1.6	6.7	9.5	4.0
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5.3	8.4	2.2	5.7	9.5	1.8	5.1	8.1	2.3
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	3.6	6.2	1.1	3.8	6.7	0.9	3.5	6.0	1.1

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	1.7	2.3	1.1	1.9	2.8	1.0	1.6	2.1	1.1
Legal intervention (Y35,Y89.0)	0.1	0.3	*	0.1	0.3	*	0.1	0.3	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.6	1.9	1.3	0.7	0.9	0.4	1.8	2.1	1.4
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	0.1	0.0	0.0	0.1	*	0.1	0.1	0.0
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.5	1.8	1.2	0.6	0.9	0.4	1.7	2.0	1.4
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.8	0.8	0.8	0.3	0.3	0.3	0.9	0.9	0.9
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	2.4	1.9	2.8	0.8	0.6	0.9	2.7	2.2	3.2
Drug-induced deaths ^{5,6}	13.1	16.1	10.2	5.5	7.6	3.4	14.5	17.7	11.4
Alcohol-induced deaths ^{5,7}	8.3	12.5	4.2	6.6	10.8	2.3	8.6	12.8	4.6
Injury by firearms ^{5,8}	10.3	18.0	2.7	6.0	10.5	1.3	11.0	19.4	3.0

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Figures for origin not stated are included in "All origins" but not distributed among specified origins.

²Includes races other than white and black.

³Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
All causes	984.3	987.5	981.2	718.7	764.5	676.9
Salmonella infections (A01-A02)	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	4.3	3.4	5.2	2.0	1.9	2.2
Tuberculosis (A16-A19)	0.1	0.1	0.1	0.3	0.5	0.2
Respiratory tuberculosis (A16)	0.1	0.1	0.1	0.2	0.4	*
Other tuberculosis (A17-A19)	0.0	0.0	0.0	0.1	*	*
Whooping cough (A37)	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	0.0	0.0	*	*	*
Septicemia (A40-A41)	13.0	12.2	13.7	15.1	14.2	15.9
Syphilis (A50-A53)	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.5	3.4	1.6	2.8	3.7	1.9
Human immunodeficiency virus (HIV) disease (B20-B24)	1.2	2.1	0.4	11.7	15.9	7.7
Malaria (B50-B54)	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	2.2	2.2	2.1	2.0	2.1	1.8
Malignant neoplasms (C00-C97)	230.1	245.8	215.0	165.5	178.6	153.5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	3.4	4.7	2.1	2.5	3.8	1.3
Malignant neoplasm of esophagus (C15)	6.1	9.8	2.5	3.6	5.5	1.9
Malignant neoplasm of stomach (C16)	3.5	4.2	2.8	5.0	6.2	3.9
Malignant neoplasms of colon, rectum and anus (C18-C21)	20.3	21.3	19.4	17.6	19.1	16.3
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	6.8	9.2	4.5	7.1	10.4	4.0
Malignant neoplasm of pancreas (C25)	14.6	15.3	14.0	10.9	11.1	10.7
Malignant neoplasm of larynx (C32)	1.4	2.3	0.6	1.5	2.5	0.6

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	66.3	73.8	59.0	41.9	51.6	33.1
Malignant melanoma of skin (C43)	4.4	5.9	2.9	0.3	0.3	0.4
Malignant neoplasm of breast (C50)	15.9	0.3	31.0	15.4	0.4	29.0
Malignant neoplasm of cervix uteri (C53)	1.3	...	2.5	2.0	...	3.8
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	3.1	...	6.1	3.6	...	6.9
Malignant neoplasm of ovary (C56)	6.0	...	11.8	3.2	...	6.1
Malignant neoplasm of prostate (C61)	10.8	22.0	...	12.2	25.5	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	5.4	7.0	3.8	3.0	3.7	2.3
Malignant neoplasm of bladder (C67)	6.4	9.4	3.5	2.7	3.2	2.2
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	6.0	6.9	5.1	2.2	2.4	2.1
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	22.5	25.5	19.5	13.7	14.8	12.6
Hodgkin's disease (C81)	0.5	0.6	0.4	0.3	0.3	0.3
Non-Hodgkin's lymphoma (C82-C85)	8.4	9.3	7.5	3.8	4.3	3.3
Leukemia (C91-C95)	9.3	10.9	7.8	4.5	5.0	4.1
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	4.2	4.7	3.8	5.0	5.2	4.9
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.0	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52,C57-C60, C62-C63,C66,C68-C69,C73-C80,C97)	26.1	28.1	24.1	17.1	18.0	16.3
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6.2	6.7	5.8	3.1	2.9	3.2
Anemias (D50-D64)	1.8	1.4	2.1	2.5	2.3	2.7
Diabetes mellitus (E10-E14)	23.9	25.5	22.2	30.4	29.6	31.1
Nutritional deficiencies (E40-E64)	1.2	0.9	1.4	0.9	0.7	1.1
Malnutrition (E40-E46)	1.1	0.9	1.3	0.9	0.7	1.1
Other nutritional deficiencies (E50-E64)	0.1	0.1	0.1	*	*	*

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Meningitis (G00,G03)	0.2	0.2	0.2	0.3	0.3	0.4
Parkinson's disease (G20-G21)	9.9	11.8	8.1	2.0	2.4	1.7
Alzheimer's disease (G30)	36.7	22.6	50.3	13.1	7.8	17.9
Major cardiovascular diseases (I00-I78)	313.1	310.9	315.2	232.8	239.2	227.0
Diseases of heart (I00-I09,I11,I13,I20-I51)	241.8	251.8	232.2	173.0	183.9	163.0
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	1.2	0.8	1.6	0.6	0.4	0.7
Hypertensive heart disease (I11)	11.5	10.7	12.2	19.3	21.3	17.5
Hypertensive heart and renal disease (I13)	0.9	0.8	1.0	1.9	2.0	1.8
Ischemic heart diseases (I20-I25)	154.6	172.4	137.5	99.0	107.8	91.0
Acute myocardial infarction (I21-I22)	49.8	56.3	43.4	31.9	33.8	30.2
Other acute ischemic heart diseases (I24)	1.7	1.8	1.6	1.4	1.6	1.2
Other forms of chronic ischemic heart disease (I20,I25)	103.2	114.3	92.4	65.7	72.4	59.6
Atherosclerotic cardiovascular disease, so described (I25.0)	21.9	25.8	18.1	20.8	25.7	16.4
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	81.3	88.5	74.3	44.8	46.7	43.1
Other heart diseases (I26-I51)	73.6	67.1	79.9	52.2	52.4	52.0
Acute and subacute endocarditis (I33)	0.4	0.5	0.3	0.5	0.5	0.4
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.3	0.3	0.3	0.3	0.3	0.3
Heart failure (I50)	24.6	20.9	28.2	13.9	12.9	14.9
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	48.3	45.4	51.1	37.5	38.6	36.5
Essential hypertension and hypertensive renal disease (I10,I12,I15)	9.4	7.6	11.2	12.8	11.7	13.9
Cerebrovascular diseases (I60-I69)	50.9	40.7	60.8	40.1	36.6	43.4
Atherosclerosis (I70)	3.1	2.5	3.7	1.5	1.3	1.6
Other diseases of circulatory system (I71-I78)	7.8	8.3	7.4	5.4	5.6	5.1
Aortic aneurysm and dissection (I71)	4.3	5.1	3.6	2.4	2.9	2.0

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of arteries, arterioles and capillaries (I72-I78)	3.5	3.1	3.8	2.9	2.7	3.1
Other disorders of circulatory system (I80-I99)	1.6	1.5	1.6	1.9	2.0	1.8
Influenza and pneumonia (J09-J18)	20.1	19.0	21.2	12.3	12.4	12.2
Influenza (J09-J11)	0.2	0.2	0.2	0.2	0.2	0.2
Pneumonia (J12-J18)	19.9	18.8	21.1	12.1	12.3	12.0
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.1	0.1	0.1	*	*
Acute bronchitis and bronchiolitis (J20-J21)	0.1	0.1	0.1	0.1	*	*
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	61.4	58.3	64.4	21.8	23.8	20.0
Bronchitis, chronic and unspecified (J40-J42)	0.3	0.2	0.3	0.1	0.1	0.1
Emphysema (J43)	4.5	4.7	4.4	1.5	1.9	1.1
Asthma (J45-J46)	1.0	0.7	1.4	2.2	2.1	2.2
Other chronic lower respiratory diseases (J44,J47)	55.6	52.8	58.3	18.0	19.7	16.6
Pneumoconioses and chemical effects (J60-J66,J68)	0.4	0.8	0.0	0.1	0.2	*
Pneumonitis due to solids and liquids (J69)	7.3	8.0	6.6	3.5	3.8	3.2
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	12.7	13.2	12.2	7.4	7.2	7.5
Peptic ulcer (K25-K28)	1.2	1.2	1.3	0.7	0.9	0.6
Diseases of appendix (K35-K38)	0.2	0.2	0.1	0.1	0.1	*
Hernia (K40-K46)	0.8	0.7	0.8	0.4	0.4	0.4
Chronic liver disease and cirrhosis (K70,K73-K74)	11.8	15.5	8.2	6.6	9.0	4.4
Alcoholic liver disease (K70)	5.8	8.3	3.4	3.1	4.3	1.9
Other chronic liver disease and cirrhosis (K73-K74)	6.0	7.2	4.9	3.5	4.7	2.5
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.3	1.3	1.4	0.8	0.6	0.9
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18.5	18.8	18.1	22.2	21.2	23.2

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.1	0.1	0.1	0.1	*	*
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	2.2	2.1	2.3	2.4	2.3	2.5
Renal failure (N17-N19)	16.2	16.6	15.8	19.8	18.8	20.6
Other disorders of kidney (N25,N27)	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	0.1	0.3	0.2	0.1	0.2
Hyperplasia of prostate (N40)	0.2	0.4	...	0.1	0.2	...
Inflammatory diseases of female pelvic organs (N70-N76)	0.1	...	0.1	0.1	...	0.1
Pregnancy, childbirth and the puerperium (O00-O99)	0.2	...	0.4	0.7	...	1.3
Pregnancy with abortive outcome (O00-O07)	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.2	...	0.3	0.6	...	1.2
Certain conditions originating in the perinatal period (P00-P96)	2.5	2.9	2.2	9.8	11.2	8.4
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2.9	3.0	2.8	4.1	4.5	3.7
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	15.3	12.4	18.0	12.3	12.2	12.3
All other diseases (Residual)	110.6	89.8	130.8	71.6	62.5	79.9
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47.2	58.7	36.0	30.1	42.1	19.1
Transport accidents (V01-V99,Y85)	13.4	19.2	7.9	11.8	17.9	6.4
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)	12.4	17.6	7.5	11.2	16.7	6.1

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9, V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	0.3	0.5	0.2	0.3	0.6	0.1
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.7	1.0	0.3	0.3	0.6	0.1
Nontransport accidents (W00-X59,Y86)	33.8	39.5	28.2	18.2	24.3	12.7
Falls (W00-W19)	11.4	11.4	11.5	2.9	3.5	2.4
Accidental discharge of firearms (W32-W34)	0.2	0.4	0.1	0.3	0.5	*
Accidental drowning and submersion (W65-W74)	1.2	1.9	0.6	1.4	2.4	0.4
Accidental exposure to smoke, fire and flames (X00-X09)	1.0	1.1	0.8	1.4	1.8	1.2
Accidental poisoning and exposure to noxious substances (X40-X49)	13.3	17.0	9.7	7.4	9.9	5.2
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39, X50-X59,Y86)	6.6	7.8	5.5	4.7	6.1	3.5
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16.0	25.7	6.7	5.3	9.1	1.8
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	8.5	15.0	2.1	2.7	5.0	0.5
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)	7.5	10.6	4.5	2.6	4.1	1.3
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2.5	3.3	1.7	19.5	35.0	5.3
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	1.4	1.9	0.9	15.3	29.0	2.9
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9, *U02,X85-X92,X96-Y09,Y87.1)	1.1	1.4	0.9	4.1	6.0	2.5

Table 15. Death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Legal intervention (Y35,Y89.0)	0.1	0.2	*	0.2	0.5	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.9	2.2	1.6	1.5	2.0	1.0
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	0.2	0.0	0.1	0.1	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.8	2.1	1.5	1.4	1.9	1.0
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.9	0.9	1.0	1.0	0.9	1.1
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	3.1	2.5	3.8	1.2	1.0	1.4
Drug-induced deaths ^{5,6}	16.6	20.0	13.2	8.9	11.5	6.5
Alcohol-induced deaths ^{5,7}	9.4	14.0	5.0	5.8	8.8	3.1
Injury by firearms ^{5,8}	10.2	17.6	3.1	18.6	35.1	3.5

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Figures for origin not stated are included in "All origins" but not distributed among specified origins.

²Includes races other than white and black.

³Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9, F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1, M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	747.0	887.1	634.9	741.8	878.5	630.8	898.2	1,104.0	752.5
Salmonella infections (A01-A02)	0.0	*	*	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	3.1	3.1	3.2	3.2	3.1	3.3	2.6	2.9	2.5
Tuberculosis (A16-A19)	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.6	0.2
Respiratory tuberculosis (A16)	0.1	0.2	0.1	0.1	0.1	0.0	0.3	0.5	0.1
Other tuberculosis (A17-A19)	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Whooping cough (A37)	0.0	*	*	0.0	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Septicemia (A40-A41)	10.6	11.7	9.7	9.8	10.9	9.0	19.4	22.0	17.9
Syphilis (A50-A53)	0.0	0.0	*	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.1	2.9	1.4	2.1	2.9	1.3	2.8	3.9	1.9
Human immunodeficiency virus (HIV) disease (B20-B24)	2.6	3.8	1.4	1.4	2.3	0.5	11.6	16.5	7.5
Malaria (B50-B54)	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.8	2.1	1.5	1.7	2.0	1.5	2.3	2.8	1.9
Malignant neoplasms (C00-C97)	172.8	209.9	146.7	172.4	208.2	146.9	203.8	264.8	167.1
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	2.5	3.8	1.4	2.5	3.7	1.4	2.8	4.8	1.4
Malignant neoplasm of esophagus (C15)	4.3	7.6	1.6	4.4	7.8	1.6	4.2	7.4	2.0
Malignant neoplasm of stomach (C16)	3.4	4.6	2.5	3.0	4.0	2.2	6.3	9.3	4.4
Malignant neoplasms of colon, rectum and anus (C18-C21)	15.8	19.0	13.3	15.3	18.4	12.9	21.8	27.8	17.9
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	6.0	8.8	3.6	5.5	8.1	3.3	7.8	12.4	4.3
Malignant neoplasm of pancreas (C25)	11.0	12.8	9.6	10.9	12.7	9.5	13.7	15.8	12.1
Malignant neoplasm of larynx (C32)	1.1	2.0	0.4	1.1	1.9	0.4	1.8	3.4	0.7
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	47.6	60.3	38.1	48.3	60.1	39.3	51.4	73.7	36.5

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant melanoma of skin (C43)	2.8	4.1	1.7	3.2	4.7	2.0	0.4	0.5	0.4
Malignant neoplasm of breast (C50)	12.4	0.3	22.1	12.0	0.3	21.5	17.8	0.6	30.3
Malignant neoplasm of cervix uteri (C53)	1.2	...	2.3	1.1	...	2.1	2.2	...	3.9
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	2.5	...	4.5	2.3	...	4.2	4.5	...	7.6
Malignant neoplasm of ovary (C56)	4.4	...	7.9	4.5	...	8.2	3.9	...	6.7
Malignant neoplasm of prostate (C61)	8.7	21.9	...	8.1	20.2	...	17.2	48.0	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	3.9	5.7	2.5	4.1	5.9	2.6	3.6	5.2	2.6
Malignant neoplasm of bladder (C67)	4.5	7.8	2.2	4.7	8.2	2.2	3.7	5.6	2.5
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	4.3	5.2	3.4	4.6	5.6	3.7	2.5	3.1	2.2
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	17.0	22.0	13.2	17.2	22.4	13.3	17.1	21.6	14.1
Hodgkin's disease (C81)	0.4	0.5	0.3	0.4	0.5	0.3	0.3	0.4	0.3
Non-Hodgkin's lymphoma (C82-C85)	6.2	7.9	4.9	6.4	8.1	5.1	4.6	5.9	3.7
Leukemia (C91-C95)	6.9	9.2	5.2	7.2	9.6	5.4	5.7	7.3	4.6
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.4	4.3	2.8	3.2	4.1	2.6	6.5	8.0	5.6
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.0	0.0	0.0	0.0	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	19.5	23.9	16.3	19.7	24.1	16.3	20.8	25.5	17.7
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.6	5.8	3.7	4.7	6.0	3.8	4.0	4.5	3.6
Anemias (D50-D64)	1.5	1.5	1.4	1.3	1.3	1.2	3.0	3.0	2.9
Diabetes mellitus (E10-E14)	20.8	24.9	17.6	19.0	23.1	15.6	38.7	43.6	35.1
Nutritional deficiencies (E40-E64)	0.9	0.9	0.9	0.9	0.8	0.9	1.3	1.2	1.3
Malnutrition (E40-E46)	0.8	0.8	0.8	0.8	0.8	0.8	1.2	1.2	1.3
Other nutritional deficiencies (E50-E64)	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Meningitis (G00,G03)	0.2	0.2	0.2	0.2	0.2	0.1	0.4	0.4	0.4
Parkinson's disease (G20-G21)	6.8	10.4	4.6	7.3	11.0	4.9	3.0	4.9	2.1
Alzheimer's disease (G30)	25.1	21.0	27.3	26.0	21.7	28.4	20.6	17.9	21.7
Major cardiovascular diseases (I00-I78)	234.2	282.0	196.1	229.6	277.1	191.1	304.1	366.6	258.5
Diseases of heart (I00-I09,I11,I13,I20-I51)	179.1	225.1	143.3	176.9	222.9	140.4	224.9	280.6	185.3

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	0.9	0.7	1.0	0.9	0.8	1.0	0.7	0.5	0.8
Hypertensive heart disease (I11)	10.0	11.2	8.7	8.6	9.5	7.5	23.6	29.0	19.3
Hypertensive heart and renal disease (I13)	0.8	0.9	0.7	0.7	0.7	0.6	2.4	2.8	2.0
Ischemic heart diseases (I20-I25)	113.6	151.3	84.9	113.5	151.9	83.8	131.2	169.0	104.9
Acute myocardial infarction (I21-I22)	36.5	48.0	27.5	36.6	48.5	27.1	41.8	51.8	34.7
Other acute ischemic heart diseases (I24)	1.3	1.5	1.0	1.2	1.5	1.0	1.7	2.2	1.4
Other forms of chronic ischemic heart disease (I20,I25)	75.9	101.8	56.4	75.7	101.9	55.7	87.6	115.0	68.8
Atherosclerotic cardiovascular disease, so described (I25.0)	17.0	23.1	11.8	16.3	22.2	11.3	26.0	36.6	18.4
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	58.9	78.7	44.5	59.4	79.8	44.4	61.6	78.4	50.4
Other heart diseases (I26-I51)	53.7	60.9	48.0	53.1	60.1	47.5	67.1	79.2	58.4
Acute and subacute endocarditis (I33)	0.3	0.4	0.3	0.3	0.4	0.2	0.5	0.6	0.4
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.3
Heart failure (I50)	17.3	19.2	15.9	17.4	19.2	16.0	19.3	22.3	17.3
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	35.9	41.0	31.6	35.2	40.1	31.0	46.9	55.9	40.4
Essential hypertension and hypertensive renal disease (I10,I12,I15)	8.0	8.0	7.8	7.0	7.0	6.9	17.0	18.3	15.8
Cerebrovascular diseases (I60-I69)	39.1	39.3	38.3	37.7	37.6	37.2	53.0	56.6	49.6
Atherosclerosis (I70)	2.2	2.3	2.0	2.2	2.3	2.1	2.1	2.4	1.9
Other diseases of circulatory system (I71-I78)	5.8	7.3	4.7	5.8	7.3	4.6	7.1	8.7	5.9
Aortic aneurysm and dissection (I71)	3.2	4.4	2.3	3.2	4.5	2.3	3.0	4.0	2.3
Other diseases of arteries, arterioles and capillaries (I72-I78)	2.6	2.9	2.4	2.6	2.8	2.3	4.0	4.7	3.6
Other disorders of circulatory system (I80-I99)	1.3	1.4	1.2	1.2	1.3	1.1	2.2	2.6	2.0
Influenza and pneumonia (J09-J18)	15.1	18.2	13.1	14.9	17.8	13.0	16.8	21.3	14.1
Influenza (J09-J11)	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2
Pneumonia (J12-J18)	14.9	18.0	12.9	14.7	17.6	12.8	16.7	21.1	14.0
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	*
Acute bronchitis and bronchiolitis (J20-J21)	0.0	0.0	0.0	0.0	0.0	0.1	0.1	*	*
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	0.0	*	0.0	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	42.2	48.7	38.0	44.6	50.6	40.8	29.0	39.8	22.8

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Bronchitis, chronic and unspecified (J40-J42)	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.1
Emphysema (J43)	3.1	3.8	2.6	3.3	4.0	2.8	2.0	3.1	1.3
Asthma (J45-J46)	1.0	0.9	1.2	0.9	0.7	1.0	2.3	2.2	2.3
Other chronic lower respiratory diseases (J44,J47)	37.9	43.8	34.1	40.3	45.8	36.8	24.5	34.3	19.1
Pneumoconioses and chemical effects (J60-J66,J68)	0.3	0.6	0.0	0.3	0.7	0.0	0.1	0.4	*
Pneumonitis due to solids and liquids (J69)	5.1	7.3	3.8	5.2	7.4	3.9	4.9	6.9	3.7
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	9.5	11.7	8.0	9.7	11.9	8.1	9.2	11.0	8.2
Peptic ulcer (K25-K28)	0.9	1.0	0.8	0.9	1.0	0.8	0.9	1.3	0.7
Diseases of appendix (K35-K38)	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	*
Hernia (K40-K46)	0.5	0.6	0.5	0.6	0.6	0.5	0.6	0.6	0.5
Chronic liver disease and cirrhosis (K70,K73-K74)	9.4	12.9	6.2	9.9	13.6	6.5	6.7	9.8	4.3
Alcoholic liver disease (K70)	4.7	7.0	2.6	5.0	7.4	2.8	3.1	4.5	1.9
Other chronic liver disease and cirrhosis (K73-K74)	4.7	5.9	3.6	4.9	6.2	3.8	3.7	5.2	2.5
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.2	1.0
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15.3	18.7	13.0	14.0	17.5	11.6	29.3	34.0	26.4
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.0	0.1	0.0	0.0	0.0	0.0	0.1	*	*
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.8	2.2	1.5	1.6	2.0	1.4	3.3	4.0	2.9
Renal failure (N17-N19)	13.4	16.5	11.5	12.2	15.4	10.2	25.9	29.8	23.5
Other disorders of kidney (N25,N27)	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2
Hyperplasia of prostate (N40)	0.1	0.4	...	0.1	0.4	...	0.1	0.4	...
Inflammatory diseases of female pelvic organs (N70-N76)	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
Pregnancy, childbirth and the puerperium (O00-O99)	0.3	...	0.6	0.2	...	0.5	0.6	...	1.2
Pregnancy with abortive outcome (O00-O07)	0.0	...	0.0	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.3	...	0.5	0.2	...	0.4	0.6	...	1.2
Certain conditions originating in the perinatal period (P00-P96)	4.2	4.7	3.8	3.5	3.8	3.1	8.3	9.0	7.6

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.2	3.3	3.1	3.2	3.3	3.1	3.6	3.9	3.4
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	11.7	11.9	11.1	11.5	11.6	11.0	14.9	16.5	13.4
All other diseases (Residual)	81.1	80.9	79.6	81.3	80.8	79.8	94.3	98.9	90.2
Accidents (unintentional injuries) (V01-X59, Y85-Y86)	38.0	51.5	25.6	40.3	54.0	27.3	31.3	46.0	19.3
Transport accidents (V01-V99, Y85)	12.1	17.6	6.9	12.5	18.1	7.2	11.6	17.9	6.1
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	11.3	16.2	6.5	11.7	16.7	6.8	10.9	16.7	5.9
Other land transport accidents (V01, V05-V06, V09.1, V09.3-V09.9, V10-V11, V15-V18, V19.3, V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9)	0.3	0.5	0.1	0.3	0.5	0.2	0.3	0.6	0.2
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99, Y85)	0.5	0.8	0.2	0.5	0.9	0.2	0.3	0.6	0.1
Nontransport accidents (W00-X59, Y86)	25.9	33.9	18.7	27.7	35.9	20.1	19.7	28.1	13.2
Falls (W00-W19)	7.9	9.9	6.4	8.4	10.4	6.8	3.8	5.5	2.7
Accidental discharge of firearms (W32-W34)	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.5	*
Accidental drowning and submersion (W65-W74)	1.2	1.9	0.5	1.2	1.9	0.6	1.3	2.3	0.4
Accidental exposure to smoke, fire and flames (X00-X09)	0.9	1.1	0.7	0.8	1.0	0.6	1.6	2.1	1.2
Accidental poisoning and exposure to noxious substances (X40-X49)	10.6	13.8	7.5	11.9	15.3	8.5	7.3	10.0	5.0
Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64, W75-W99, X10-X39, X50-X59, Y86)	5.2	7.0	3.6	5.2	7.1	3.6	5.5	7.7	3.7
Intentional self-harm (suicide) (*U03, X60-X84, Y87.0)	12.1	19.8	5.0	13.6	22.0	5.6	5.2	9.1	1.8
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	6.1	11.2	1.5	6.9	12.5	1.8	2.7	5.0	0.5
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60-X71, X75-X84, Y87.0)	6.0	8.7	3.5	6.7	9.5	3.9	2.6	4.0	1.3
Assault (homicide) (*U01-*U02, X85-Y09, Y87.1)	5.3	8.4	2.3	3.3	4.7	1.8	17.7	31.5	5.0

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All races			White ¹			Black ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	3.6	6.1	1.1	1.9	2.9	0.9	13.8	25.7	2.6
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	1.7	2.2	1.1	1.3	1.7	0.9	3.9	5.7	2.3
Legal intervention (Y35,Y89.0)	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.6	1.9	1.2	1.7	2.0	1.3	1.4	2.0	1.0
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.5	1.8	1.2	1.6	1.9	1.3	1.4	1.9	0.9
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.8	0.8	0.7	0.7	0.8	0.7	1.2	1.2	1.2
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	2.2	2.3	2.2	2.3	2.3	2.3	1.7	1.9	1.6
Drug-induced deaths ^{5,6}	12.9	15.9	10.0	14.6	17.7	11.4	8.7	11.6	6.2
Alcohol-induced deaths ^{5,7}	7.6	11.7	3.9	8.0	12.2	4.0	5.9	9.6	3.0
Injury by firearms ^{5,8}	10.1	17.9	2.7	9.2	16.1	2.7	16.9	31.8	3.3

¹Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
All causes	628.3	730.2	541.7	424.3	512.1	359.0
Salmonella infections (A01-A02)	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	3.0	4.1	2.5	1.3	1.4	1.2
Tuberculosis (A16-A19)	*	*	*	0.8	1.1	0.5
Respiratory tuberculosis (A16)	*	*	*	0.6	1.0	0.4
Other tuberculosis (A17-A19)	*	*	*	*	*	*
Whooping cough (A37)	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*
Meningococcal infection (A39)	*	*	*	*	*	*
Septicemia (A40-A41)	10.5	9.8	10.7	5.0	6.2	4.2
Syphilis (A50-A53)	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.8	3.6	1.9	2.1	2.5	1.8
Human immunodeficiency virus (HIV) disease (B20-B24)	1.6	2.6	*	0.4	0.7	*
Malaria (B50-B54)	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.9	2.1	1.6	1.3	1.6	1.1
Malignant neoplasms (C00-C97)	122.4	151.0	102.0	108.9	131.0	93.5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	1.5	2.1	*	2.0	2.9	1.3
Malignant neoplasm of esophagus (C15)	2.5	4.2	*	1.7	3.0	0.7
Malignant neoplasm of stomach (C16)	3.5	5.1	2.3	6.1	8.3	4.6
Malignant neoplasms of colon, rectum and anus (C18-C21)	11.7	14.2	9.8	11.4	13.3	9.9
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	7.8	12.1	4.2	10.0	14.8	6.1
Malignant neoplasm of pancreas (C25)	7.1	8.1	6.1	7.8	8.1	7.5
Malignant neoplasm of larynx (C32)	1.1	2.1	*	0.4	0.8	*
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	33.1	41.6	26.3	24.8	33.8	18.3
Malignant melanoma of skin (C43)	0.8	*	*	0.4	0.4	0.4
Malignant neoplasm of breast (C50)	6.4	*	11.5	6.7	*	11.9

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of cervix uteri (C53)	1.2	...	2.2	1.0	...	1.7
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	1.5	...	2.7	1.7	...	3.0
Malignant neoplasm of ovary (C56)	2.9	...	5.2	2.6	...	4.8
Malignant neoplasm of prostate (C61)	6.1	15.3	...	3.8	9.6	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.7	6.9	3.0	1.9	3.0	1.2
Malignant neoplasm of bladder (C67)	2.2	3.6	*	1.8	2.9	1.1
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	2.3	2.6	2.1	2.2	2.6	1.9
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	9.9	13.2	7.8	10.3	13.5	8.0
Hodgkin's disease (C81)	*	*	*	0.2	*	*
Non-Hodgkin's lymphoma (C82-C85)	3.1	3.6	2.6	4.1	5.1	3.4
Leukemia (C91-C95)	4.4	6.4	3.3	4.2	5.5	3.3
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	2.2	2.8	1.8	1.8	2.6	1.3
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	16.1	18.5	14.4	12.2	13.8	11.0
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2.7	3.4	2.0	3.0	3.6	2.6
Anemias (D50-D64)	*	*	*	0.7	0.8	0.6
Diabetes mellitus (E10-E14)	36.4	39.2	34.1	15.5	18.0	13.6
Nutritional deficiencies (E40-E64)	1.6	*	1.8	0.5	0.5	0.4
Malnutrition (E40-E46)	1.5	*	*	0.4	0.5	0.4
Other nutritional deficiencies (E50-E64)	*	*	*	*	*	*
Meningitis (G00,G03)	*	*	*	0.1	*	*
Parkinson's disease (G20-G21)	3.8	6.1	2.2	3.6	4.9	2.7
Alzheimer's disease (G30)	17.2	14.4	18.6	10.9	9.0	12.2
Major cardiovascular diseases (I00-I78)	169.1	200.3	142.8	145.9	175.6	123.4
Diseases of heart (I00-I09,I11,I13,I20-I51)	128.6	158.7	103.5	100.9	127.2	81.2
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	*	*	*	0.8	0.5	1.0
Hypertensive heart disease (I11)	6.6	8.3	5.2	5.8	6.6	5.2

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart and renal disease (I13)	*	*	*	0.7	0.7	0.7
Ischemic heart diseases (I20-I25)	84.9	109.6	64.9	68.7	91.0	52.2
Acute myocardial infarction (I21-I22)	26.8	35.3	20.0	22.1	28.9	16.9
Other acute ischemic heart diseases (I24)	1.6	1.9	*	0.5	0.7	0.3
Other forms of chronic ischemic heart disease (I20,I25)	56.5	72.5	43.6	46.2	61.4	34.9
Atherosclerotic cardiovascular disease, so described (I25.0)	15.7	21.2	11.2	10.0	13.4	7.3
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	40.8	51.3	32.4	36.1	48.1	27.5
Other heart diseases (I26-I51)	35.4	38.9	32.0	24.9	28.4	22.1
Acute and subacute endocarditis (I33)	*	*	*	0.1	*	*
Diseases of pericardium and acute myocarditis (I30-I31,I40)	*	*	*	0.1	*	*
Heart failure (I50)	12.7	11.8	13.1	6.7	7.1	6.4
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	21.9	26.2	18.3	17.8	20.9	15.4
Essential hypertension and hypertensive renal disease (I10,I12,I15)	6.9	6.6	7.1	7.5	7.6	7.5
Cerebrovascular diseases (I60-I69)	28.1	29.8	26.5	33.2	35.2	31.4
Atherosclerosis (I70)	*	*	*	1.0	1.3	0.8
Other diseases of circulatory system (I71-I78)	4.4	4.5	4.3	3.3	4.3	2.5
Aortic aneurysm and dissection (I71)	2.5	3.0	2.1	2.3	3.1	1.7
Other diseases of arteries, arterioles and capillaries (I72-I78)	1.9	*	2.2	1.0	1.2	0.8
Other disorders of circulatory system (I80-I99)	0.8	*	*	0.6	0.9	0.5
Influenza and pneumonia (J09-J18)	15.9	19.2	13.7	14.4	19.1	11.3
Influenza (J09-J11)	*	*	*	*	*	*
Pneumonia (J12-J18)	15.8	19.1	13.6	14.3	18.9	11.2
Other acute lower respiratory infections (J20-J22,U04)	*	*	*	*	*	*
Acute bronchitis and bronchiolitis (J20-J21)	*	*	*	*	*	*
Other and unspecified acute lower respiratory infections (J22,U04)	*	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	33.8	38.8	30.3	13.9	21.0	9.1
Bronchitis, chronic and unspecified (J40-J42)	*	*	*	*	*	*
Emphysema (J43)	1.9	2.1	1.7	0.9	1.8	0.3
Asthma (J45-J46)	0.8	*	*	1.0	1.3	0.9
Other chronic lower respiratory diseases (J44,J47)	31.0	35.8	27.7	11.8	17.8	7.8

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Pneumoconioses and chemical effects (J60-J66,J68)	*	*	*	*	*	*
Pneumonitis due to solids and liquids (J69)	4.0	4.2	3.7	3.0	4.4	2.1
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	11.4	12.9	10.4	5.6	6.8	4.7
Peptic ulcer (K25-K28)	1.0	*	*	0.8	1.1	0.6
Diseases of appendix (K35-K38)	*	*	*	*	*	*
Hernia (K40-K46)	*	*	*	*	*	*
Chronic liver disease and cirrhosis (K70,K73-K74)	22.8	25.5	20.3	3.2	4.4	2.2
Alcoholic liver disease (K70)	16.2	20.1	12.7	1.0	1.9	0.3
Other chronic liver disease and cirrhosis (K73-K74)	6.6	5.5	7.6	2.2	2.5	1.9
Cholelithiasis and other disorders of gallbladder (K80-K82)	0.8	*	*	0.9	1.1	0.7
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16.4	16.7	16.2	9.6	11.0	8.6
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	*	*	*	*	*	*
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	2.3	2.7	2.1	1.2	1.2	1.3
Renal failure (N17-N19)	14.0	14.0	14.0	8.3	9.8	7.2
Other disorders of kidney (N25,N27)	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	*	*	*	*	*	*
Hyperplasia of prostate (N40)	*	*	...	*	*	...
Inflammatory diseases of female pelvic organs (N70-N76)	*	...	*	*	...	*
Pregnancy, childbirth and the puerperium (O00-O99)	*	...	*	0.2	...	0.4
Pregnancy with abortive outcome (O00-O07)	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	*	...	*	0.2	...	0.4
Certain conditions originating in the perinatal period (P00-P96)	2.1	2.7	1.5	2.9	3.3	2.4
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2.5	2.6	2.3	2.0	1.9	2.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	8.1	8.6	7.4	4.3	4.8	3.9
All other diseases (Residual)	65.8	63.1	66.1	38.2	40.7	36.1
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46.9	64.5	30.7	15.0	20.3	10.7

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Transport accidents (V01-V99, Y85)	16.9	23.0	11.0	5.5	7.2	4.0
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	15.7	21.1	10.6	5.1	6.5	3.9
Other land transport accidents (V01, V05-V06, V09.1, V09.3-V09.9, V10-V11, V15-V18, V19.3, V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9)	0.5	*	*	0.2	0.3	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99, Y85)	0.6	*	*	0.2	0.4	*
Nontransport accidents (W00-X59, Y86)	30.1	41.5	19.6	9.5	13.0	6.7
Falls (W00-W19)	7.4	9.5	5.7	4.9	6.7	3.6
Accidental discharge of firearms (W32-W34)	*	*	*	*	*	*
Accidental drowning and submersion (W65-W74)	1.6	2.8	*	1.1	1.6	0.5
Accidental exposure to smoke, fire and flames (X00-X09)	1.3	1.5	*	0.2	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)	13.0	16.8	9.3	1.4	2.1	0.8
Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64, W75-W99, X10-X39, X50-X59, Y86)	6.5	10.3	3.2	1.9	2.5	1.5
Intentional self-harm (suicide) (*U03, X60-X84, Y87.0)	10.8	15.5	6.1	6.2	9.5	3.4
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	4.3	6.7	1.8	1.3	2.5	0.2
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60-X71, X75-X84, Y87.0)	6.6	8.8	4.3	4.9	7.1	3.2
Assault (homicide) (*U01-*U02, X85-Y09, Y87.1)	5.7	8.8	2.5	1.8	2.6	1.2
Assault (homicide) by discharge of firearms (*U01.4, X93-X95)	2.5	4.1	*	0.9	1.5	0.3
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3, *U01.5-*U01.9, *U02, X85-X92, X96-Y09, Y87.1)	3.2	4.7	1.8	0.9	1.0	0.8
Legal intervention (Y35, Y89.0)	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34, Y87.2, Y89.9)	2.0	2.3	1.9	0.5	0.7	0.3

Table 16. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by race and sex: United States, 2010

[Age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female
Discharge of firearms, undetermined intent (Y22-Y24)	*	*	*	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21, Y25-Y34, Y87.2, Y89.9)	1.9	2.1	1.8	0.5	0.7	0.3
Operations of war and their sequelae (Y36, Y89.1)	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84, Y88)	0.8	*	*	0.3	*	0.4
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	2.0	2.8	*	0.9	1.0	0.9
Drug-induced deaths ^{5,6}	11.4	12.3	10.4	2.0	2.7	1.5
Alcohol-induced deaths ^{5,7}	25.4	34.9	16.7	1.6	2.8	0.5
Injury by firearms ^{5,8}	7.3	11.7	2.6	2.3	4.2	0.6

¹Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

⁴Included in "Certain other intestinal infections (A04, A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, F11.1-F11.5, F11.7-F11.9, F12.1-F12.5, F12.7-F12.9, F13.1-F13.5, F13.7-F13.9, F14.1-F14.5, F14.7-F14.9, F15.1-F15.5, F15.7-F15.9, F16.1-F16.5, F16.7-F16.9, F17.3-F17.5, F17.7-F17.9, F18.1-F18.5, F18.7-F18.9, F19.1-F19.5, F19.7-F19.9, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2-J70.4, K85.3, L10.5, L27.0-L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R50.2, R78.1-R78.5, X40-X44, X60-X64, X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, R78.0, X45, X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4, W32-W34, X72-X74, X93-X95, Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	747.0	887.1	634.9	558.6	677.7	463.4	762.6	904.6	649.2
Salmonella infections (A01-A02)	0.0	*	*	*	*	*	0.0	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	3.1	3.1	3.2	2.5	2.6	2.5	3.2	3.1	3.2
Tuberculosis (A16-A19)	0.2	0.2	0.1	0.3	0.5	0.2	0.2	0.2	0.1
Respiratory tuberculosis (A16)	0.1	0.2	0.1	0.3	0.4	0.1	0.1	0.2	0.1
Other tuberculosis (A17-A19)	0.0	0.0	0.0	0.1	*	*	0.0	0.0	0.0
Whooping cough (A37)	0.0	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	0.0	0.0	0.0	*	*	0.0	0.0	0.0
Septicemia (A40-A41)	10.6	11.7	9.7	8.3	9.3	7.4	10.7	11.8	10.0
Syphilis (A50-A53)	0.0	0.0	*	*	*	*	0.0	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.1	2.9	1.4	3.3	4.6	2.1	2.0	2.8	1.3
Human immunodeficiency virus (HIV) disease (B20-B24)	2.6	3.8	1.4	2.8	4.6	1.1	2.6	3.8	1.5
Malaria (B50-B54)	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.8	2.1	1.5	1.5	1.6	1.3	1.8	2.1	1.6
Malignant neoplasms (C00-C97)	172.8	209.9	146.7	119.7	149.4	99.4	177.1	214.6	150.7
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	2.5	3.8	1.4	1.5	2.4	0.8	2.6	3.9	1.5
Malignant neoplasm of esophagus (C15)	4.3	7.6	1.6	2.3	4.1	0.8	4.5	7.9	1.7
Malignant neoplasm of stomach (C16)	3.4	4.6	2.5	5.8	7.6	4.4	3.2	4.4	2.3
Malignant neoplasms of colon, rectum and anus (C18-C21)	15.8	19.0	13.3	12.3	15.9	9.7	16.1	19.2	13.6
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	6.0	8.8	3.6	8.8	12.9	5.4	5.7	8.5	3.4
Malignant neoplasm of pancreas (C25)	11.0	12.8	9.6	8.6	9.4	8.0	11.2	13.0	9.8
Malignant neoplasm of larynx (C32)	1.1	2.0	0.4	0.8	1.8	0.2	1.1	2.0	0.4

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	47.6	60.3	38.1	20.4	29.6	13.8	49.9	62.8	40.2
Malignant melanoma of skin (C43)	2.8	4.1	1.7	0.8	1.0	0.6	2.9	4.4	1.8
Malignant neoplasm of breast (C50)	12.4	0.3	22.1	8.0	*	14.4	12.7	0.3	22.8
Malignant neoplasm of cervix uteri (C53)	1.2	...	2.3	1.4	...	2.6	1.2	...	2.2
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	2.5	...	4.5	1.9	...	3.4	2.5	...	4.5
Malignant neoplasm of ovary (C56)	4.4	...	7.9	3.1	...	5.6	4.4	...	8.0
Malignant neoplasm of prostate (C61)	8.7	21.9	...	7.3	18.4	...	8.8	22.1	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	3.9	5.7	2.5	3.6	5.0	2.5	4.0	5.8	2.6
Malignant neoplasm of bladder (C67)	4.5	7.8	2.2	2.3	3.8	1.3	4.6	8.1	2.3
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	4.3	5.2	3.4	2.8	3.5	2.3	4.4	5.4	3.6
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	17.0	22.0	13.2	13.5	16.5	11.2	17.2	22.3	13.3
Hodgkin's disease (C81)	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3
Non-Hodgkin's lymphoma (C82-C85)	6.2	7.9	4.9	5.2	6.2	4.5	6.2	8.0	4.9
Leukemia (C91-C95)	6.9	9.2	5.2	4.9	6.0	4.0	7.0	9.4	5.3
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.4	4.3	2.8	3.0	3.7	2.4	3.5	4.4	2.8
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	19.5	23.9	16.3	14.5	17.3	12.3	20.0	24.4	16.6
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.6	5.8	3.7	3.2	4.0	2.7	4.7	5.9	3.8
Anemias (D50-D64)	1.5	1.5	1.4	1.0	1.1	0.8	1.5	1.6	1.5
Diabetes mellitus (E10-E14)	20.8	24.9	17.6	27.1	31.2	23.7	20.4	24.4	17.2
Nutritional deficiencies (E40-E64)	0.9	0.9	0.9	0.8	0.9	0.8	0.9	0.9	0.9
Malnutrition (E40-E46)	0.8	0.8	0.8	0.8	0.9	0.7	0.8	0.8	0.8

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other nutritional deficiencies (E50-E64)	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0
Meningitis (G00,G03)	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Parkinson's disease (G20-G21)	6.8	10.4	4.6	5.1	7.2	3.7	7.0	10.6	4.7
Alzheimer's disease (G30)	25.1	21.0	27.3	18.5	16.6	19.5	25.5	21.3	27.8
Major cardiovascular diseases (I00-I78)	234.2	282.0	196.1	177.9	213.8	149.6	238.3	287.1	199.3
Diseases of heart (I00-I09,I11,I13,I20-I51)	179.1	225.1	143.3	132.8	165.1	107.8	182.6	229.6	145.8
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	0.9	0.7	1.0	0.6	0.4	0.8	0.9	0.7	1.1
Hypertensive heart disease (I11)	10.0	11.2	8.7	8.5	10.1	7.1	10.1	11.3	8.8
Hypertensive heart and renal disease (I13)	0.8	0.9	0.7	0.9	1.0	0.8	0.8	0.9	0.7
Ischemic heart diseases (I20-I25)	113.6	151.3	84.9	92.3	119.0	72.0	115.2	153.7	85.8
Acute myocardial infarction (I21-I22)	36.5	48.0	27.5	28.8	37.2	22.3	37.2	48.9	27.9
Other acute ischemic heart diseases (I24)	1.3	1.5	1.0	0.6	0.5	0.6	1.3	1.6	1.0
Other forms of chronic ischemic heart disease (I20,I25)	75.9	101.8	56.4	62.9	81.2	49.1	76.7	103.2	56.8
Atherosclerotic cardiovascular disease, so described (I25.0)	17.0	23.1	11.8	13.8	19.6	9.1	17.2	23.3	12.0
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	58.9	78.7	44.5	49.1	61.6	39.9	59.5	79.9	44.8
Other heart diseases (I26-I51)	53.7	60.9	48.0	30.5	34.6	27.1	55.5	62.9	49.5
Acute and subacute endocarditis (I33)	0.3	0.4	0.3	0.3	0.3	0.2	0.4	0.4	0.3
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.2	0.3	0.2	0.1	0.2	0.1	0.2	0.3	0.2
Heart failure (I50)	17.3	19.2	15.9	10.0	11.4	9.0	17.7	19.7	16.3
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	35.9	41.0	31.6	20.1	22.7	17.8	37.1	42.5	32.7
Essential hypertension and hypertensive renal disease (I10,I12,I15)	8.0	8.0	7.8	7.8	8.1	7.5	8.0	8.0	7.8
Cerebrovascular diseases (I60-I69)	39.1	39.3	38.3	32.1	33.9	30.2	39.6	39.6	38.9
Atherosclerosis (I70)	2.2	2.3	2.0	1.3	1.6	1.2	2.2	2.4	2.1
Other diseases of circulatory system (I71-I78)	5.8	7.3	4.7	3.8	5.1	2.9	6.0	7.5	4.8
Aortic aneurysm and dissection (I71)	3.2	4.4	2.3	1.8	2.6	1.1	3.3	4.5	2.3

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of arteries, arterioles and capillaries (I72-I78)	2.6	2.9	2.4	2.1	2.5	1.7	2.7	2.9	2.5
Other disorders of circulatory system (I80-I99)	1.3	1.4	1.2	0.9	1.0	0.8	1.3	1.4	1.2
Influenza and pneumonia (J09-J18)	15.1	18.2	13.1	13.7	17.0	11.4	15.2	18.3	13.2
Influenza (J09-J11)	0.1	0.2	0.2	0.2	0.3	0.2	0.1	0.1	0.1
Pneumonia (J12-J18)	14.9	18.0	12.9	13.4	16.7	11.2	15.1	18.1	13.0
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1
Acute bronchitis and bronchiolitis (J20-J21)	0.0	0.0	0.0	*	*	*	0.0	0.0	0.1
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	0.0	*	*	*	*	0.0	*	*
Chronic lower respiratory diseases (J40-J47)	42.2	48.7	38.0	19.6	25.2	15.9	43.9	50.4	39.7
Bronchitis, chronic and unspecified (J40-J42)	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2
Emphysema (J43)	3.1	3.8	2.6	1.1	1.7	0.8	3.2	3.9	2.7
Asthma (J45-J46)	1.0	0.9	1.2	0.9	0.7	1.1	1.1	0.9	1.2
Other chronic lower respiratory diseases (J44,J47)	37.9	43.8	34.1	17.3	22.5	13.9	39.4	45.3	35.6
Pneumoconioses and chemical effects (J60-J66,J68)	0.3	0.6	0.0	0.1	0.3	*	0.3	0.7	0.0
Pneumonitis due to solids and liquids (J69)	5.1	7.3	3.8	3.3	4.7	2.4	5.3	7.4	3.9
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	9.5	11.7	8.0	8.5	9.8	7.5	9.6	11.8	8.1
Peptic ulcer (K25-K28)	0.9	1.0	0.8	0.6	0.7	0.5	0.9	1.1	0.8
Diseases of appendix (K35-K38)	0.1	0.2	0.1	0.1	0.1	*	0.1	0.2	0.1
Hernia (K40-K46)	0.5	0.6	0.5	0.5	0.4	0.6	0.6	0.6	0.5
Chronic liver disease and cirrhosis (K70,K73-K74)	9.4	12.9	6.2	13.7	19.8	8.2	9.0	12.2	6.1
Alcoholic liver disease (K70)	4.7	7.0	2.6	6.8	11.7	2.4	4.5	6.5	2.7
Other chronic liver disease and cirrhosis (K73-K74)	4.7	5.9	3.6	6.9	8.1	5.7	4.5	5.7	3.4
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.0	1.2	0.9	1.1	1.4	0.9	1.0	1.2	0.9
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15.3	18.7	13.0	14.1	17.3	11.9	15.3	18.8	13.1

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.0	0.1	0.0	*	*	*	0.0	0.1	0.0
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.8	2.2	1.5	1.5	1.8	1.3	1.8	2.2	1.5
Renal failure (N17-N19)	13.4	16.5	11.5	12.5	15.5	10.6	13.5	16.5	11.5
Other disorders of kidney (N25,N27)	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	0.1	0.2	0.2	*	0.2	0.2	0.1	0.2
Hyperplasia of prostate (N40)	0.1	0.4	...	0.1	0.4	...	0.1	0.4	...
Inflammatory diseases of female pelvic organs (N70-N76)	0.0	...	0.1	*	...	*	0.0	...	0.1
Pregnancy, childbirth and the puerperium (O00-O99)	0.3	...	0.6	0.3	...	0.6	0.3	...	0.6
Pregnancy with abortive outcome (O00-O07)	0.0	...	0.0	*	...	*	0.0	...	0.0
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.3	...	0.5	0.3	...	0.5	0.3	...	0.5
Certain conditions originating in the perinatal period (P00-P96)	4.2	4.7	3.8	3.4	3.9	3.0	4.4	4.8	4.0
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.2	3.3	3.1	2.7	2.8	2.7	3.3	3.4	3.2
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	11.7	11.9	11.1	7.0	7.2	6.6	12.1	12.3	11.5
All other diseases (Residual)	81.1	80.9	79.6	58.1	60.3	55.3	82.8	82.5	81.3
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38.0	51.5	25.6	25.8	37.2	14.9	39.7	53.6	27.1
Transport accidents (V01-V99,Y85)	12.1	17.6	6.9	10.2	15.0	5.5	12.4	18.0	7.1
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)	11.3	16.2	6.5	9.6	14.0	5.3	11.5	16.6	6.8

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3,V19.8- V19.9,V80.0-V80.2,V80.6-V80.9, V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	0.3	0.5	0.1	0.4	0.6	0.1	0.3	0.5	0.2
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.5	0.8	0.2	0.2	0.4	*	0.5	0.9	0.2
Nontransport accidents (W00-X59,Y86)	25.9	33.9	18.7	15.6	22.2	9.4	27.4	35.7	19.9
Falls (W00-W19)	7.9	9.9	6.4	5.2	7.0	3.8	8.0	10.0	6.5
Accidental discharge of firearms (W32-W34)	0.2	0.3	0.1	0.1	0.1	*	0.2	0.4	0.1
Accidental drowning and submersion (W65-W74)	1.2	1.9	0.5	1.0	1.7	0.3	1.3	1.9	0.6
Accidental exposure to smoke, fire and flames (X00-X09)	0.9	1.1	0.7	0.5	0.7	0.3	0.9	1.2	0.7
Accidental poisoning and exposure to noxious substances (X40-X49)	10.6	13.8	7.5	5.6	8.1	3.1	11.6	14.9	8.3
Other and unspecified nontransport accidents and their sequelae (W20- W31,W35-W64,W75-W99,X10-X39, X50-X59,Y86)	5.2	7.0	3.6	3.1	4.6	1.8	5.3	7.2	3.7
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12.1	19.8	5.0	5.9	9.9	2.1	13.1	21.4	5.4
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	6.1	11.2	1.5	2.2	4.2	0.4	6.6	12.2	1.6
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)	6.0	8.7	3.5	3.6	5.6	1.7	6.4	9.2	3.8
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5.3	8.4	2.3	5.3	8.7	1.8	5.3	8.2	2.3
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	3.6	6.1	1.1	3.4	5.8	0.9	3.7	6.1	1.2

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, drug-induced causes, alcohol-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All origins ¹			Hispanic			Non-Hispanic ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	1.7	2.2	1.1	1.9	2.9	0.9	1.6	2.1	1.2
Legal intervention (Y35,Y89.0)	0.1	0.3	*	0.2	0.3	*	0.1	0.3	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.6	1.9	1.2	0.7	1.0	0.5	1.7	2.1	1.4
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	0.1	0.0	0.0	0.1	*	0.1	0.1	0.0
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.5	1.8	1.2	0.7	0.9	0.4	1.6	1.9	1.3
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.8	0.8	0.7	0.5	0.6	0.4	0.8	0.8	0.8
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	2.2	2.3	2.2	1.8	1.8	1.8	2.2	2.3	2.2
Drug-induced deaths ^{5,6}	12.9	15.9	10.0	6.1	8.4	3.8	14.2	17.3	11.1
Alcohol-induced deaths ^{5,7}	7.6	11.7	3.9	9.1	16.0	3.0	7.4	11.2	4.0
Injury by firearms ^{5,8}	10.1	17.9	2.7	5.9	10.5	1.3	10.7	19.1	2.9

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Figures for origin not stated are included in "All origins" but not distributed among specified origins.

²Includes races other than white and black.

³Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
All causes	755.0	892.5	643.3	920.4	1,131.7	770.8
Salmonella infections (A01-A02)	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	3.2	3.2	3.3	2.7	3.0	2.5
Tuberculosis (A16-A19)	0.1	0.1	0.1	0.3	0.6	0.2
Respiratory tuberculosis (A16)	0.1	0.1	0.0	0.3	0.5	*
Other tuberculosis (A17-A19)	0.0	0.0	0.0	0.1	*	*
Whooping cough (A37)	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	0.0	0.0	*	*	*
Septicemia (A40-A41)	9.9	11.0	9.1	20.0	22.6	18.3
Syphilis (A50-A53)	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*
Viral hepatitis (B15-B19)	1.9	2.6	1.2	2.8	4.0	1.9
Human immunodeficiency virus (HIV) disease (B20-B24)	1.1	1.8	0.4	12.0	17.0	7.9
Malaria (B50-B54)	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.7	2.0	1.5	2.3	2.9	2.0
Malignant neoplasms (C00-C97)	176.5	212.6	150.6	208.8	271.1	171.4
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	2.6	3.8	1.4	2.9	4.9	1.4
Malignant neoplasm of esophagus (C15)	4.6	8.2	1.7	4.3	7.7	2.0
Malignant neoplasm of stomach (C16)	2.7	3.6	1.9	6.5	9.5	4.5
Malignant neoplasms of colon, rectum and anus (C18-C21)	15.5	18.5	13.1	22.4	28.6	18.4
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	5.2	7.5	3.2	8.0	12.7	4.4
Malignant neoplasm of pancreas (C25)	11.1	12.9	9.6	14.0	16.2	12.4
Malignant neoplasm of larynx (C32)	1.1	1.9	0.4	1.8	3.5	0.7
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	50.8	62.7	41.7	52.6	75.6	37.3

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10), Second Edition*; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Malignant melanoma of skin (C43)	3.4	5.1	2.1	0.4	0.5	0.4
Malignant neoplasm of breast (C50)	12.3	0.3	22.1	18.4	0.6	31.3
Malignant neoplasm of cervix uteri (C53)	1.0	...	2.0	2.3	...	4.0
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	2.3	...	4.2	4.6	...	7.7
Malignant neoplasm of ovary (C56)	4.6	...	8.3	4.0	...	6.8
Malignant neoplasm of prostate (C61)	8.1	20.3	...	17.5	49.0	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.1	6.0	2.6	3.8	5.4	2.6
Malignant neoplasm of bladder (C67)	4.9	8.5	2.3	3.8	5.7	2.6
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	4.8	5.9	3.9	2.6	3.1	2.2
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	17.4	22.6	13.4	17.5	22.1	14.4
Hodgkin's disease (C81)	0.4	0.5	0.3	0.3	0.4	0.3
Non-Hodgkin's lymphoma (C82-C85)	6.4	8.2	5.1	4.7	6.0	3.7
Leukemia (C91-C95)	7.3	9.8	5.4	5.8	7.5	4.7
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.2	4.1	2.6	6.6	8.2	5.7
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.0	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	20.1	24.6	16.6	21.3	26.0	18.2
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.8	6.1	3.9	4.1	4.6	3.7
Anemias (D50-D64)	1.3	1.3	1.3	3.1	3.1	3.0
Diabetes mellitus (E10-E14)	18.2	22.3	14.9	39.6	44.6	35.9
Nutritional deficiencies (E40-E64)	0.8	0.8	0.9	1.3	1.3	1.3
Malnutrition (E40-E46)	0.8	0.8	0.8	1.3	1.2	1.3
Other nutritional deficiencies (E50-E64)	0.0	0.0	0.0	*	*	*
Meningitis (G00,G03)	0.2	0.2	0.2	0.4	0.4	0.4
Parkinson's disease (G20-G21)	7.5	11.2	5.0	3.1	5.0	2.1

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10), Second Edition*; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Alzheimer's disease (G30)	26.4	22.0	28.9	20.9	18.2	22.0
Major cardiovascular diseases (I00-I78)	232.8	281.1	193.5	310.5	374.5	263.9
Diseases of heart (I00-I09,I11,I13,I20-I51)	179.9	226.9	142.5	229.5	286.3	189.1
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	0.9	0.8	1.1	0.7	0.5	0.8
Hypertensive heart disease (I11)	8.6	9.4	7.5	24.0	29.6	19.6
Hypertensive heart and renal disease (I13)	0.6	0.7	0.6	2.4	2.9	2.1
Ischemic heart diseases (I20-I25)	115.0	154.2	84.4	133.4	171.9	106.7
Acute myocardial infarction (I21-I22)	37.3	49.4	27.5	42.7	52.9	35.4
Other acute ischemic heart diseases (I24)	1.3	1.5	1.0	1.8	2.3	1.4
Other forms of chronic ischemic heart disease (I20,I25)	76.4	103.2	56.0	89.0	116.7	69.8
Atherosclerotic cardiovascular disease, so described (I25.0)	16.4	22.3	11.4	26.4	37.1	18.7
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	60.0	81.0	44.6	62.5	79.6	51.1
Other heart diseases (I26-I51)	54.7	61.9	48.9	68.9	81.4	60.0
Acute and subacute endocarditis (I33)	0.3	0.4	0.3	0.5	0.7	0.4
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.2	0.2	0.2	0.3	0.4	0.3
Heart failure (I50)	17.8	19.7	16.4	19.8	22.8	17.7
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	36.3	41.4	32.0	48.2	57.5	41.5
Essential hypertension and hypertensive renal disease (I10,I12,I15)	6.9	6.9	6.7	17.4	18.8	16.1
Cerebrovascular diseases (I60-I69)	37.8	37.5	37.4	54.3	58.1	50.7
Atherosclerosis (I70)	2.2	2.4	2.1	2.2	2.5	2.0
Other diseases of circulatory system (I71-I78)	5.9	7.5	4.7	7.2	8.9	6.0
Aortic aneurysm and dissection (I71)	3.3	4.6	2.4	3.1	4.1	2.3
Other diseases of arteries, arterioles and capillaries (I72-I78)	2.6	2.8	2.4	4.1	4.9	3.7

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10), Second Edition*; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Other disorders of circulatory system (I80-I99)	1.2	1.3	1.1	2.3	2.6	2.0
Influenza and pneumonia (J09-J18)	14.9	17.7	13.1	17.1	21.7	14.3
Influenza (J09-J11)	0.1	0.1	0.1	0.2	0.2	0.2
Pneumonia (J12-J18)	14.8	17.6	12.9	16.9	21.5	14.2
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.1	0.1	0.1	*	*
Acute bronchitis and bronchiolitis (J20-J21)	0.1	0.0	0.0	0.1	*	*
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	46.6	52.5	42.9	29.6	40.7	23.3
Bronchitis, chronic and unspecified (J40-J42)	0.2	0.2	0.2	0.1	0.2	0.1
Emphysema (J43)	3.4	4.1	3.0	2.0	3.1	1.3
Asthma (J45-J46)	0.9	0.6	1.0	2.4	2.3	2.4
Other chronic lower respiratory diseases (J44,J47)	42.1	47.5	38.7	25.1	35.1	19.5
Pneumoconioses and chemical effects (J60-J66,J68)	0.3	0.7	0.0	0.1	0.4	*
Pneumonitis due to solids and liquids (J69)	5.4	7.6	4.0	5.0	7.1	3.8
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	9.7	11.9	8.1	9.5	11.3	8.5
Peptic ulcer (K25-K28)	0.9	1.1	0.8	0.9	1.3	0.7
Diseases of appendix (K35-K38)	0.1	0.2	0.1	0.1	0.2	*
Hernia (K40-K46)	0.6	0.6	0.5	0.6	0.6	0.5
Chronic liver disease and cirrhosis (K70,K73-K74)	9.4	12.7	6.4	6.9	10.0	4.5
Alcoholic liver disease (K70)	4.7	6.8	2.8	3.1	4.6	1.9
Other chronic liver disease and cirrhosis (K73-K74)	4.7	5.9	3.6	3.8	5.4	2.6
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.0	1.2	0.9	1.1	1.2	1.0
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13.8	17.4	11.5	30.1	34.9	27.0
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.0	0.1	0.0	0.1	*	*

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.6	2.0	1.4	3.4	4.1	2.9
Renal failure (N17-N19)	12.1	15.3	10.1	26.6	30.6	24.0
Other disorders of kidney (N25,N27)	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	0.1	0.2	0.2	0.2	0.2
Hyperplasia of prostate (N40)	0.1	0.4	...	0.1	0.4	...
Inflammatory diseases of female pelvic organs (N70-N76)	0.0	...	0.1	0.1	...	0.1
Pregnancy, childbirth and the puerperium (O00-O99)	0.2	...	0.4	0.7	...	1.3
Pregnancy with abortive outcome (O00-O07)	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.2	...	0.4	0.6	...	1.2
Certain conditions originating in the perinatal period (P00-P96)	3.3	3.6	3.0	8.8	9.5	8.0
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.2	3.3	3.1	3.8	4.2	3.6
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	11.9	12.0	11.4	15.3	17.0	13.8
All other diseases (Residual)	82.9	82.2	81.5	96.5	101.3	92.3
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42.4	56.6	29.2	32.4	47.5	20.0
Transport accidents (V01-V99,Y85)	12.9	18.5	7.5	12.1	18.6	6.4
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)	11.9	17.1	7.0	11.4	17.4	6.1

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10), Second Edition*; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9, V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	0.3	0.5	0.2	0.4	0.6	0.2
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.6	0.9	0.3	0.3	0.6	0.1
Nontransport accidents (W00-X59,Y86)	29.6	38.1	21.7	20.3	28.9	13.6
Falls (W00-W19)	8.6	10.6	7.0	3.9	5.7	2.8
Accidental discharge of firearms (W32-W34)	0.2	0.4	0.1	0.3	0.5	*
Accidental drowning and submersion (W65-W74)	1.2	1.8	0.6	1.3	2.3	0.5
Accidental exposure to smoke, fire and flames (X00-X09)	0.9	1.1	0.7	1.6	2.2	1.2
Accidental poisoning and exposure to noxious substances (X40-X49)	13.3	16.9	9.6	7.5	10.3	5.2
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39, X50-X59,Y86)	5.4	7.3	3.8	5.6	8.0	3.8
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15.0	24.2	6.2	5.4	9.4	1.9
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	7.7	14.0	2.0	2.7	5.3	0.6
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)	7.2	10.3	4.3	2.7	4.2	1.3
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2.5	3.3	1.8	18.6	33.1	5.2
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	1.4	2.0	0.9	14.6	27.2	2.8
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9, *U02,X85-X92,X96-Y09,Y87.1)	1.1	1.4	0.9	4.1	6.0	2.4

Table 17. Age-adjusted death rates for 113 selected causes, Enterocolitis due to *Clostridium difficile*, Drug-induced causes, Alcohol-induced causes, and Injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 2010

[Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10), Second Edition*; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female
Legal intervention (Y35,Y89.0)	0.1	0.2	*	0.3	0.5	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.8	2.2	1.5	1.5	2.1	1.0
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	0.2	0.0	0.1	0.1	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.8	2.0	1.5	1.4	2.0	1.0
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.8	0.8	0.7	1.2	1.3	1.2
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	2.3	2.3	2.3	1.7	1.9	1.6
Drug-induced deaths ^{5,6}	16.4	19.8	12.9	9.0	11.9	6.5
Alcohol-induced deaths ^{5,7}	7.8	11.6	4.2	6.1	9.8	3.1
Injury by firearms ^{5,8}	9.5	16.6	3.0	17.8	33.4	3.4

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Figures for origin not stated are included in "All origins" but not distributed among specified origins.

²Includes races other than white and black.

³Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 37 states and the District of Columbia in 2010; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

⁵Included in selected categories above.

⁶Includes ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. Trend data for Drug-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁷Includes ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. Trend data for Alcohol-induced deaths, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

⁸Includes ICD-10 codes U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0. Trend data for Injury by firearms, previously shown in this report, can be found through a link from the online version of this report, available from <http://www.cdc.gov/nchs/deaths.htm>.

Table 18. Number of deaths, death rates, and age-adjusted death rates for injury deaths, by mechanism and intent of death: United States, 2010

[Totals for selected causes of death differ from those shown in other tables that utilize standard mortality tabulation lists, see "Technical Notes." Rates are per 100,000 population; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Figure(s) in brackets [] applies to the code or range of codes preceding it. For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Mechanism and intent of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Number	Rate	Age-adjusted rate ¹
All injury (*U01-*U03,V01-Y36,Y85-Y87,Y89)	180,811	58.6	57.1
Unintentional (V01-X59,Y85-Y86)	120,859	39.1	38.0
Suicide (*U03,X60-X84,Y87.0)	38,364	12.4	12.1
Homicide (*U01-*U02,X85-Y09,Y87.1)	16,259	5.3	5.3
Undetermined (Y10-Y34,Y87.2,Y89.9)	4,908	1.6	1.6
Legal intervention/war (Y35-Y36,Y89[.0,.1])	421	0.1	0.2
Cut/pierce (W25-W29,W45-W46,X78,X99,Y28,Y35.4)	2,598	0.8	0.8
Unintentional (W25-W29,W45-W46)	105	0.0	0.0
Suicide (X78)	673	0.2	0.2
Homicide (X99)	1,799	0.6	0.6
Undetermined (Y28)	21	0.0	0.0
Legal intervention/war (Y35.4)	-	*	*
Drowning (W65-W74,X71,X92,Y21)	4,521	1.5	1.4
Unintentional (W65-W74)	3,782	1.2	1.2
Suicide (X71)	409	0.1	0.1
Homicide (X92)	52	0.0	0.0
Undetermined (Y21)	278	0.1	0.1
Fall (W00-W19,X80,Y01,Y30)	26,852	8.7	8.1
Unintentional (W00-W19)	26,009	8.4	7.9
Suicide (X80)	781	0.3	0.3
Homicide (Y01)	12	*	*
Undetermined (Y30)	50	0.0	0.0
Fire/hot object or substance (*U01.3,X00-X19,X76-X77,X97-X98,Y26-Y27,Y36.3) ²	3,194	1.0	1.0
Unintentional (X00-X19)	2,845	0.9	0.9
Suicide (X76-X77)	131	0.0	0.0
Homicide (*U01.3,X97-X98)	92	0.0	0.0
Undetermined (Y26-Y27)	126	0.0	0.0
Legal intervention/war (Y36.3)	-	*	*
Fire/flame (X00-X09,X76,X97,Y26)	3,127	1.0	1.0
Unintentional (X00-X09)	2,782	0.9	0.9
Suicide (X76)	131	0.0	0.0
Homicide (X97)	89	0.0	0.0
Undetermined (Y26)	125	0.0	0.0
Hot object/substance (X10-X19,X77,X98,Y27)	67	0.0	0.0
Unintentional (X10-X19)	63	0.0	0.0
Suicide (X77)	-	*	*
Homicide (X98)	3	*	*
Undetermined (Y27)	1	*	*
Firearm (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0)	31,672	10.3	10.1

Table 18. Number of deaths, death rates, and age-adjusted death rates for injury deaths, by mechanism and intent of death: United States, 2010

[Totals for selected causes of death differ from those shown in other tables that utilize standard mortality tabulation lists, see "Technical Notes." Rates are per 100,000 population; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Figure(s) in brackets [] applies to the code or range of codes preceding it. For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Mechanism and intent of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Number	Rate	Age-adjusted rate ¹
Unintentional (W32-W34)	606	0.2	0.2
Suicide (X72-X74)	19,392	6.3	6.1
Homicide (*U01.4,X93-X95)	11,078	3.6	3.6
Undetermined (Y22-Y24)	252	0.1	0.1
Legal intervention/war (Y35.0)	344	0.1	0.1
Machinery (W24,W30-W31) ³	590	0.2	0.2
All transport (*U01.1,V01-V99,X82,Y03,Y32,Y36.1)	37,402	12.1	11.9
Unintentional (V01-V99)	37,236	12.1	11.8
Suicide (X82)	114	0.0	0.0
Homicide (*U01.1,Y03)	39	0.0	0.0
Undetermined (Y32)	13	*	*
Legal intervention/war (Y36.1)	-	*	*
Motor vehicle traffic (V02-V04[.1,.9],V09.2,V12-V14[.3-.9],V19[.4-.6],V20-V28[.3-.9],V29-V79[.4-.9],V80[.3-.5],V81.1,V82.1,V83-V86[.0-.3],V87[.0-.8],V89.2) ³	33,687	10.9	10.7
Occupant (V30-V79[.4-.9],V83-V86[.0-.3]) ³	10,246	3.3	3.3
Motorcyclist (V20-V28[.3-.9],V29[.4-.9]) ³	4,177	1.4	1.3
Pedal cyclist (V12-V14[.3-.9],V19[.4-.6]) ³	551	0.2	0.2
Pedestrian (V02-V04[.1,.9],V09.2) ³	4,383	1.4	1.4
Other (V80[.3-.5],V81.1,V82.1) ³	10	*	*
Unspecified (V87[.0-.8],V89.2) ³	14,320	4.6	4.6
Pedal cyclist, other (V10-V11,V12-V14[.0-.2],V15-V18,V19[.0-.3,.8,.9]) ³	242	0.1	0.1
Pedestrian, other (V01,V02-V04[.0],V05,V06,V09[.0,.1,.3,.9]) ³	1,074	0.3	0.3
Other land transport (V20-V28[.0-.2],V29-V79[.0-.3],V80[.0-.2,.6-.9],V81-V82[.0,.2-.9],V83-V86[.4-.9],V87.9,V88[.0-.9],V89[.0,.1,.3,.9],X82,Y03,Y32)	1,524	0.5	0.5
Unintentional (V20-V28[.0-.2],V29-V79[.0-.3],V80[.0-.2,.6-.9],V81-V82[.0,.2-.9],V83-V86[.4-.9],V87.9,V88[.0-.9],V89[.0,.1,.3,.9])	1,358	0.4	0.4
Suicide (X82)	114	0.0	0.0
Homicide (Y03)	39	0.0	0.0
Undetermined (Y32)	13	*	*
Other transport (*U01.1,V90-V99,Y36.1)	875	0.3	0.3
Unintentional (V90-V99)	875	0.3	0.3
Homicide (*U01.1)	-	*	*
Legal intervention/war (Y36.1)	-	*	*
Natural/environmental (W42-W43,W53-W64,W92-W99,X20-X39,X51-X57) ³	1,576	0.5	0.5
Overexertion (X50) ³	10	*	*
Poisoning (*U01[.6-.7],X40-X49,X60-X69,X85-X90,Y10-Y19,Y35.2)	42,917	13.9	13.7
Unintentional (X40-X49)	33,041	10.7	10.6
Suicide (X60-X69)	6,599	2.1	2.1

Table 18. Number of deaths, death rates, and age-adjusted death rates for injury deaths, by mechanism and intent of death: United States, 2010

[Totals for selected causes of death differ from those shown in other tables that utilize standard mortality tabulation lists, see "Technical Notes." Rates are per 100,000 population; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Figure(s) in brackets [] applies to the code or range of codes preceding it. For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Mechanism and intent of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Number	Rate	Age-adjusted rate ¹
Homicide (*U01[.6-.7],X85-X90)	79	0.0	0.0
Undetermined (Y10-Y19)	3,197	1.0	1.0
Legal intervention/war (Y35.2)	1	*	*
Struck by or against (W20-W22,W50-W52,X79,Y00,Y04,Y29,Y35.3)	912	0.3	0.3
Unintentional (W20-W22,W50-W52)	788	0.3	0.2
Suicide (X79)	-	*	*
Homicide (Y00,Y04)	123	0.0	0.0
Undetermined (Y29)	1	*	*
Legal intervention/war (Y35.3)	-	*	*
Suffocation (W75-W84,X70,X91,Y20)	16,362	5.3	5.2
Unintentional (W75-W84)	6,165	2.0	1.9
Suicide (X70)	9,493	3.1	3.1
Homicide (X91)	544	0.2	0.2
Undetermined (Y20)	160	0.1	0.0
Other specified, classifiable (*U01[.0,.2,.5],*U03.0,W23,W35-W41,W44,W49,W85-W91,X75,X81,X96,Y02,Y05-Y07,Y25,Y31,Y35[.1,.5],Y36[.0,.2,.4-.8],Y85)	2,002	0.6	0.7
Unintentional (W23,W35-W41,W44,W49,W85-W91,Y85)	1,395	0.5	0.5
Suicide (*U03.0,X75,X81)	337	0.1	0.1
Homicide (*U01[.0,.2,.5],X96,Y02,Y05-Y07)	211	0.1	0.1
Undetermined (Y25,Y31)	15	*	*
Legal intervention/war (Y35[.1,.5],Y36[.0,.2,.4-.8])	44	0.0	0.0
Other specified, not elsewhere classified (*U01.8,*U02,X58,X83,Y08,Y33,Y35.6,Y86-Y87,Y89[.0-.1])	1,973	0.6	0.6
Unintentional (X58,Y86)	1,023	0.3	0.3
Suicide (X83,Y87.0)	245	0.1	0.1
Homicide (*U01.8,*U02,Y08,Y87.1)	495	0.2	0.2
Undetermined (Y33,Y87.2)	184	0.1	0.1
Legal intervention/war (Y35.6,Y89[.0-.1])	26	0.0	0.0
Unspecified (*U01.9,*U03.9,X59,X84,Y09,Y34,Y35.7,Y36.9,Y89.9)	8,230	2.7	2.5
Unintentional (X59)	5,688	1.8	1.7
Suicide (*U03.9,X84)	190	0.1	0.1
Homicide (*U01.9,Y09)	1,735	0.6	0.6
Undetermined (Y34,Y89.9)	611	0.2	0.2
Legal intervention/war (Y35.7,Y36.9)	6	*	*

0.0 Quantity more than zero but less than 0.05.

- Quantity zero.

* Figure does not meet standard of reliability or precision; see "Technical Notes."

¹ For method of computation, see "Technical Notes."

² Codes *U01.3 and Y36.3 cannot be divided separately into the subcategories shown below; therefore, subcategories may not add to the total.

³ Intent of death is unintentional.

Table 19. Number of deaths, death rates, and age-adjusted death rates for major causes of death: United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, 2010

[Rates per 100,000 population; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates for the United States and each state are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Rates for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are based on the 2010 census, estimated as of July 1, 2010; see "Technical Notes." Numbers after causes of death are categories of the International Classification of Diseases, Tenth Revision (ICD-10). The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the ICD-10; see "Technical Notes." For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Area	All causes		Human immunodeficiency virus (HIV) disease (B20-B24)			Malignant neoplasms (C00-C97)			Diabetes mellitus (E10-E14)			
	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹
United States ²	2,468,435	799.5	747.0	8,369	2.7	2.6	574,743	186.2	172.8	69,071	22.4	20.8
Alabama	48,038	1,005.0	939.7	153	3.2	3.2	10,196	213.3	191.7	1,302	27.2	25.0
Alaska	3,728	524.9	771.5	6	*	*	884	124.5	176.9	86	12.1	19.6
Arizona	46,762	731.6	693.1	102	1.6	1.6	10,678	167.1	154.2	1,389	21.7	20.3
Arkansas	28,916	991.7	892.7	64	2.2	2.2	6,475	222.1	194.7	850	29.2	25.8
California	234,012	628.2	646.7	742	2.0	1.9	56,453	151.5	156.9	7,061	19.0	19.7
Colorado	31,465	625.6	682.7	67	1.3	1.2	7,035	139.9	149.5	721	14.3	15.2
Connecticut	28,692	802.8	652.9	83	2.3	2.0	6,954	194.6	163.4	662	18.5	15.4
Delaware	7,706	858.2	769.9	46	5.1	4.9	1,909	212.6	185.7	196	21.8	19.2
District of Columbia	4,672	776.4	792.4	121	20.1	20.4	1,041	173.0	178.3	145	24.1	24.9
Florida	173,791	924.4	701.1	1,068	5.7	5.4	41,467	220.6	165.6	5,024	26.7	20.1
Georgia	71,263	735.6	845.4	495	5.1	5.0	15,435	159.3	174.8	1,996	20.6	23.1
Hawaii	9,617	707.0	589.6	13	*	*	2,266	166.6	140.9	272	20.0	16.6
Idaho	11,429	729.1	731.6	9	*	*	2,530	161.4	159.9	353	22.5	22.5
Illinois	99,931	778.8	736.9	300	2.3	2.3	24,070	187.6	178.6	2,507	19.5	18.5
Indiana	56,743	875.2	820.6	91	1.4	1.5	13,164	203.0	188.6	1,587	24.5	22.7
Iowa	27,745	910.8	721.7	15	*	*	6,358	208.7	171.9	733	24.1	19.3
Kansas	24,502	858.8	762.2	27	0.9	0.9	5,377	188.5	171.3	655	23.0	20.5
Kentucky	41,983	967.5	915.0	53	1.2	1.2	9,930	228.8	208.3	1,212	27.9	26.0
Louisiana	40,667	897.1	903.8	225	5.0	5.0	9,203	203.0	197.6	1,205	26.6	26.4
Maine	12,750	959.8	749.6	9	*	*	3,248	244.5	187.9	376	28.3	21.8
Maryland	43,325	750.4	728.6	316	5.5	5.0	10,269	177.9	171.2	1,191	20.6	19.9
Massachusetts	52,583	803.1	675.0	120	1.8	1.6	12,993	198.4	171.3	1,028	15.7	13.3
Michigan	88,021	890.6	786.2	150	1.5	1.5	20,620	208.6	182.9	2,697	27.3	24.0
Minnesota	38,972	734.8	661.5	48	0.9	0.9	9,612	181.2	167.2	1,037	19.6	17.7
Mississippi	28,965	976.1	962.0	118	4.0	4.2	6,271	211.3	201.4	926	31.2	30.3
Missouri	55,281	923.1	819.5	105	1.8	1.8	12,626	210.8	185.6	1,425	23.8	21.2
Montana	8,827	892.1	754.7	7	*	*	1,923	194.4	161.0	227	22.9	19.1
Nebraska	15,171	830.7	717.8	12	*	*	3,438	188.2	167.4	452	24.7	21.7
Nevada	19,623	726.6	795.4	60	2.2	2.2	4,529	167.7	174.2	350	13.0	13.8
New Hampshire	10,201	774.9	690.4	9	*	*	2,525	191.8	167.9	236	17.9	16.1
New Jersey	69,495	790.4	691.1	378	4.3	3.9	16,815	191.3	169.5	2,098	23.9	21.0
New Mexico	15,931	773.7	749.0	33	1.6	1.6	3,358	163.1	152.4	643	31.2	29.8
New York	146,432	755.7	665.5	990	5.1	4.7	35,431	182.8	163.1	3,642	18.8	16.7
North Carolina	78,773	826.1	804.9	328	3.4	3.3	18,061	189.4	179.0	2,042	21.4	20.5
North Dakota	5,944	883.7	704.3	2	*	*	1,269	188.7	157.1	192	28.5	22.2
Ohio	108,711	942.3	815.7	138	1.2	1.1	25,083	217.4	187.7	3,470	30.1	25.8
Oklahoma	36,529	973.8	915.5	50	1.3	1.4	7,831	208.8	191.3	1,089	29.0	26.9
Oregon	31,890	832.4	723.1	47	1.2	1.2	7,638	199.4	173.9	1,052	27.5	23.7
Pennsylvania	124,596	980.9	765.9	266	2.1	1.9	29,055	228.7	181.6	3,242	25.5	20.0
Rhode Island	9,579	910.1	721.7	20	1.9	1.8	2,266	215.3	178.3	211	20.0	16.0
South Carolina	41,614	899.7	854.8	184	4.0	3.9	9,356	202.3	183.6	1,128	24.4	22.6
South Dakota	7,100	872.0	715.1	10	*	*	1,655	203.3	171.0	240	29.5	24.5
Tennessee	59,578	938.8	890.8	207	3.3	3.2	13,593	214.2	195.7	1,687	26.6	24.7
Texas	166,527	662.3	772.3	773	3.1	3.2	36,717	146.0	165.9	4,744	18.9	21.6
Utah	14,776	534.6	703.2	13	*	*	2,810	101.7	133.7	471	17.0	22.7
Vermont	5,380	859.8	718.7	7	*	*	1,392	222.5	183.2	151	24.1	20.7
Virginia	59,032	737.8	741.6	142	1.8	1.7	14,080	176.0	172.4	1,530	19.1	18.8
Washington	48,146	716.0	692.3	76	1.1	1.0	11,874	176.6	170.5	1,499	22.3	21.6
West Virginia	21,275	1,148.1	933.6	14	*	*	4,685	252.8	198.0	776	41.9	32.9
Wisconsin	47,308	831.9	719.0	51	0.9	0.8	11,279	198.3	174.5	1,158	20.4	17.7
Wyoming	4,438	787.4	778.8	6	*	*	1,016	180.3	172.6	105	18.6	18.5
Puerto Rico	29,153	783.3	712.8	334	9.0	9.0	5,174	139.0	123.3	2,951	79.3	70.2
Virgin Islands	715	672.8	663.2	7	*	*	131	123.3	109.3	45	42.3	42.3
Guam	857	537.5	810.6	2	*	*	141	88.4	133.6	41	25.7	37.1
American Samoa	224	403.8	932.9	1	*	*	29	52.3	152.5	23	41.5	104.2
Northern Marianas	174	325.1	863.3	-	*	*	26	48.6	123.0	10	*	*

Table 19. Number of deaths, death rates, and age-adjusted death rates for major causes of death: United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, 2010

[Rates per 100,000 population; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates for the United States and each state are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Rates for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are based on the 2010 census, estimated as of July 1, 2010; see "Technical Notes." Numbers after causes of death are categories of the International Classification of Diseases, Tenth Revision (ICD-10). The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the ICD-10; see "Technical Notes." For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Area	Parkinson's disease (G20-G21)			Alzheimer's disease (G30)			Diseases of heart (I00-I09,I11,I13,I20-I51)			Essential hypertension and hypertensive renal disease (I10,I12,I15)		
	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹
United States ²	22,032	7.1	6.8	83,494	27.0	25.1	597,689	193.6	179.1	26,634	8.6	8.0
Alabama	342	7.2	6.9	1,523	31.9	31.2	12,083	252.8	236.0	549	11.5	10.6
Alaska	32	4.5	9.2	85	12.0	25.9	707	99.5	151.5	25	3.5	6.4
Arizona	489	7.7	7.4	2,327	36.4	35.3	9,954	155.7	146.7	508	7.9	7.6
Arkansas	206	7.1	6.4	955	32.8	29.6	7,274	249.5	222.5	221	7.6	6.8
California	2,238	6.0	6.5	10,856	29.1	30.1	58,641	157.4	161.9	3,733	10.0	10.4
Colorado	305	6.1	7.2	1,334	26.5	31.1	6,038	120.1	132.8	261	5.2	5.9
Connecticut	237	6.6	5.3	820	22.9	17.1	7,127	199.4	155.7	321	9.0	7.0
Delaware	57	6.3	5.8	215	23.9	21.9	1,776	197.8	175.7	62	6.9	6.2
District of Columbia	26	4.3	4.6	114	18.9	18.7	1,306	217.0	222.4	57	9.5	9.6
Florida	1,755	9.3	6.7	4,831	25.7	18.1	41,737	222.0	162.3	1,821	9.7	7.1
Georgia	472	4.9	6.3	2,080	21.5	28.3	15,987	165.0	192.6	1,076	11.1	12.9
Hawaii	84	6.2	5.0	189	13.9	10.5	2,239	164.6	134.7	98	7.2	5.9
Idaho	123	7.8	8.3	410	26.2	26.8	2,495	159.2	159.3	121	7.7	7.6
Illinois	909	7.1	6.9	2,927	22.8	20.9	24,959	194.5	181.7	871	6.8	6.3
Indiana	485	7.5	7.2	1,940	29.9	27.8	13,388	206.5	191.8	454	7.0	6.5
Iowa	320	10.5	8.1	1,411	46.3	32.9	6,880	225.8	173.3	318	10.4	7.8
Kansas	256	9.0	7.9	825	28.9	24.2	5,433	190.4	164.9	157	5.5	4.6
Kentucky	296	6.8	6.8	1,464	33.7	33.5	9,662	222.7	210.1	295	6.8	6.5
Louisiana	265	5.8	6.2	1,295	28.6	30.7	10,282	226.8	229.4	390	8.6	8.9
Maine	137	10.3	8.2	502	37.8	28.5	2,628	197.8	151.1	87	6.5	5.0
Maryland	389	6.7	6.9	986	17.1	16.8	10,915	189.1	182.2	433	7.5	7.3
Massachusetts	459	7.0	5.9	1,773	27.1	21.2	12,043	183.9	150.0	433	6.6	5.3
Michigan	816	8.3	7.4	2,736	27.7	24.0	23,326	236.0	204.2	815	8.2	7.1
Minnesota	512	9.7	8.9	1,451	27.4	23.4	7,185	135.5	119.4	439	8.3	7.3
Mississippi	174	5.9	6.1	927	31.2	32.6	7,542	254.2	251.1	531	17.9	17.6
Missouri	477	8.0	7.2	1,986	33.2	28.8	13,840	231.1	201.8	437	7.3	6.3
Montana	94	9.5	8.1	302	30.5	25.2	1,849	186.9	154.2	76	7.7	6.3
Nebraska	183	10.0	8.7	565	30.9	24.9	3,355	183.7	154.2	176	9.6	8.0
Nevada	149	5.5	6.6	296	11.0	14.2	4,811	178.1	197.3	128	4.7	5.7
New Hampshire	116	8.8	8.0	396	30.1	26.9	2,290	174.0	152.7	80	6.1	5.4
New Jersey	645	7.3	6.5	1,878	21.4	17.7	18,730	213.0	182.0	750	8.5	7.2
New Mexico	171	8.3	8.3	343	16.7	16.8	3,224	156.6	151.2	122	5.9	5.7
New York	972	5.0	4.5	2,616	13.5	11.3	44,981	232.1	199.9	2,048	10.6	9.1
North Carolina	636	6.7	6.8	2,817	29.5	30.3	17,154	179.9	174.9	844	8.9	8.7
North Dakota	61	9.1	7.0	361	53.7	37.2	1,395	207.4	158.0	64	9.5	7.0
Ohio	920	8.0	6.9	4,109	35.6	29.7	26,164	226.8	192.4	1,220	10.6	8.9
Oklahoma	249	6.6	6.4	1,015	27.1	26.1	9,426	251.3	235.2	343	9.1	8.6
Oregon	356	9.3	8.3	1,300	33.9	28.5	6,198	161.8	137.9	444	11.6	9.8
Pennsylvania	1,184	9.3	7.1	3,591	28.3	20.0	31,556	248.4	187.0	977	7.7	5.7
Rhode Island	96	9.1	7.1	338	32.1	22.6	2,322	220.6	167.1	70	6.7	4.9
South Carolina	381	8.2	8.3	1,570	33.9	34.7	9,295	201.0	189.9	417	9.0	8.5
South Dakota	86	10.6	8.3	398	48.9	35.9	1,616	198.5	155.2	96	11.8	9.1
Tennessee	435	6.9	6.9	2,440	38.4	38.5	14,582	229.8	217.4	594	9.4	9.0
Texas	1,492	5.9	7.6	5,209	20.7	26.8	38,253	152.1	181.1	1,722	6.8	8.3
Utah	170	6.2	8.9	375	13.6	19.3	2,889	104.5	143.2	125	4.5	6.1
Vermont	70	11.2	9.6	238	38.0	30.8	1,172	187.3	153.6	49	7.8	6.5
Virginia	520	6.5	7.0	1,848	23.1	24.4	13,404	167.5	168.5	588	7.3	7.5
Washington	514	7.6	7.9	3,025	45.0	43.6	10,602	157.7	151.5	506	7.5	7.1
West Virginia	142	7.7	6.2	594	32.1	26.0	4,897	264.3	211.2	264	14.2	11.3
Wisconsin	492	8.7	7.4	1,762	31.0	25.3	11,115	195.4	165.1	399	7.0	5.9
Wyoming	37	6.6	6.8	146	25.9	27.2	962	170.7	169.8	19	*	*
Puerto Rico	137	3.7	3.4	1,860	50.0	46.0	5,176	139.1	124.9	496	13.3	12.0
Virgin Islands	4	*	*	18	*	*	174	163.7	149.7	20	18.8	21.5
Guam	2	*	*	3	*	*	245	153.7	254.9	20	12.5	16.6
American Samoa	-	*	*	-	*	*	42	75.7	168.9	7	*	*
Northern Marianas	1	*	*	-	*	*	37	69.1	167.0	3	*	*

Table 19. Number of deaths, death rates, and age-adjusted death rates for major causes of death: United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, 2010

[Rates per 100,000 population; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates for the United States and each state are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Rates for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are based on the 2010 census, estimated as of July 1, 2010; see "Technical Notes." Numbers after causes of death are categories of the International Classification of Diseases, Tenth Revision (ICD-10). The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the ICD-10; see "Technical Notes." For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Area	Cerebrovascular diseases (I60-I69)			Influenza and pneumonia (J09-J18)			Chronic lower respiratory diseases (J40-J47)			Chronic liver disease and cirrhosis (K70,K73-K74)		
	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹
United States ²	129,476	41.9	39.1	50,097	16.2	15.1	138,080	44.7	42.2	31,903	10.3	9.4
Alabama	2,619	54.8	51.6	942	19.7	18.7	2,866	60.0	55.4	504	10.5	9.4
Alaska	167	23.5	40.9	64	9.0	15.8	176	24.8	41.5	70	9.9	9.8
Arizona	2,138	33.4	31.9	768	12.0	11.4	2,926	45.8	43.1	870	13.6	12.7
Arkansas	1,741	59.7	53.8	647	22.2	19.9	1,773	60.8	53.5	327	11.2	9.9
California	13,662	36.7	38.1	5,882	15.8	16.4	12,987	34.9	37.0	4,287	11.5	11.3
Colorado	1,607	32.0	36.1	550	10.9	12.3	2,199	43.7	49.7	599	11.9	11.2
Connecticut	1,349	37.7	29.4	568	15.9	12.1	1,289	36.1	29.4	303	8.5	7.3
Delaware	407	45.3	40.7	138	15.4	13.8	441	49.1	44.0	93	10.4	8.8
District of Columbia	196	32.6	32.7	80	13.3	13.5	147	24.4	25.7	54	9.0	8.9
Florida	8,432	44.8	32.8	2,259	12.0	8.9	10,337	55.0	40.1	2,480	13.2	10.6
Georgia	3,762	38.8	46.3	1,456	15.0	18.3	3,816	39.4	46.7	743	7.7	7.5
Hawaii	605	44.5	35.8	289	21.2	17.0	296	21.8	18.0	102	7.5	6.5
Idaho	642	41.0	42.0	208	13.3	13.5	727	46.4	47.0	150	9.6	9.3
Illinois	5,349	41.7	39.2	2,212	17.2	16.1	5,231	40.8	39.3	1,123	8.8	8.2
Indiana	3,082	47.5	44.5	1,175	18.1	16.8	3,794	58.5	55.3	634	9.8	9.0
Iowa	1,537	50.5	38.0	560	18.4	13.7	1,698	55.7	44.7	277	9.1	7.9
Kansas	1,370	48.0	41.2	550	19.3	16.4	1,580	55.4	50.0	244	8.6	7.8
Kentucky	1,992	45.9	44.1	943	21.7	21.0	2,779	64.0	60.0	463	10.7	9.4
Louisiana	1,977	43.6	44.9	878	19.4	20.2	1,939	42.8	43.6	390	8.6	7.9
Maine	602	45.3	34.5	234	17.6	13.4	809	60.9	47.4	154	11.6	8.8
Maryland	2,279	39.5	38.8	925	16.0	15.7	2,039	35.3	35.0	453	7.8	7.0
Massachusetts	2,516	38.4	31.3	1,291	19.7	16.0	2,383	36.4	31.0	584	8.9	7.7
Michigan	4,474	45.3	39.5	1,529	15.5	13.6	5,079	51.4	45.6	1,130	11.4	10.0
Minnesota	2,167	40.9	36.1	595	11.2	9.7	2,016	38.0	35.1	414	7.8	7.0
Mississippi	1,520	51.2	51.2	567	19.1	19.2	1,661	56.0	55.2	291	9.8	9.0
Missouri	3,001	50.1	44.0	1,188	19.8	17.4	3,557	59.4	52.6	530	8.8	7.8
Montana	494	49.9	41.8	168	17.0	13.6	602	60.8	51.3	124	12.5	10.7
Nebraska	876	48.0	40.5	266	14.6	11.9	1,008	55.2	48.8	160	8.8	8.1
Nevada	796	29.5	33.3	471	17.4	19.8	1,186	43.9	49.5	316	11.7	11.1
New Hampshire	500	38.0	33.5	191	14.5	12.6	609	46.3	41.7	103	7.8	6.4
New Jersey	3,402	38.7	33.3	1,128	12.8	11.0	3,106	35.3	31.4	732	8.3	7.3
New Mexico	806	39.1	38.4	294	14.3	14.2	1,022	49.6	47.7	377	18.3	17.0
New York	6,213	32.1	27.9	4,642	24.0	20.6	6,847	35.3	31.4	1,434	7.4	6.6
North Carolina	4,298	45.1	44.7	1,700	17.8	17.7	4,495	47.1	46.1	932	9.8	8.9
North Dakota	382	56.8	42.9	128	19.0	13.9	356	52.9	43.1	70	10.4	10.1
Ohio	5,755	49.9	42.6	1,975	17.1	14.6	6,717	58.2	50.5	1,244	10.8	9.4
Oklahoma	1,980	52.8	50.0	778	20.7	19.7	2,720	72.5	67.4	501	13.4	12.5
Oregon	1,793	46.8	40.1	417	10.9	9.2	1,971	51.4	45.3	509	13.3	11.3
Pennsylvania	6,701	52.8	39.3	2,324	18.3	13.6	6,202	48.8	37.9	1,200	9.4	7.8
Rhode Island	431	40.9	31.4	197	18.7	13.8	507	48.2	38.5	123	11.7	9.7
South Carolina	2,293	49.6	47.9	747	16.2	15.9	2,264	48.9	46.3	571	12.3	10.8
South Dakota	416	51.1	39.9	169	20.8	15.8	451	55.4	46.0	85	10.4	10.1
Tennessee	3,205	50.5	48.7	1,352	21.3	20.5	3,551	56.0	52.7	735	11.6	10.2
Texas	9,180	36.5	44.4	3,022	12.0	14.6	8,921	35.5	43.0	2,873	11.4	11.7
Utah	739	26.7	37.1	348	12.6	17.1	671	24.3	33.1	167	6.0	7.3
Vermont	265	42.3	35.3	60	9.6	7.9	335	53.5	45.3	55	8.8	7.2
Virginia	3,293	41.2	42.1	1,195	14.9	15.5	2,969	37.1	38.1	693	8.7	7.9
Washington	2,548	37.9	37.0	578	8.6	8.3	2,737	40.7	40.4	783	11.6	10.5
West Virginia	1,104	59.6	47.8	436	23.5	19.0	1,482	80.0	62.7	244	13.2	10.7
Wisconsin	2,609	45.9	38.7	904	15.9	13.3	2,474	43.5	38.0	528	9.3	8.2
Wyoming	204	36.2	37.2	109	19.3	19.5	332	58.9	59.5	75	13.3	12.5
Puerto Rico	1,500	40.3	36.4	818	22.0	19.9	1,089	29.3	26.5	201	5.4	4.6
Virgin Islands	34	32.0	33.4	2	*	*	5	*	*	12	*	*
Guam	66	41.4	71.6	11	*	*	24	15.1	24.5	19	*	*
American Samoa	21	37.9	94.5	7	*	*	10	*	*	-	*	*
Northern Marianas	17	*	*	5	*	*	6	*	*	1	*	*

Table 19. Number of deaths, death rates, and age-adjusted death rates for major causes of death: United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, 2010

[Rates per 100,000 population; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates for the United States and each state are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Rates for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are based on the 2010 census, estimated as of July 1, 2010; see "Technical Notes." Numbers after causes of death are categories of the International Classification of Diseases, Tenth Revision (ICD-10). The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the ICD-10; see "Technical Notes." For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Area	Nephritis, nephrotic syndrome, and nephrosis (N00-N07,N17-N19,N25-N27)			Accidents (V01-X59,Y85-Y86)			Motor vehicle accidents ³			Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)		
	Number	Rate	Age- adjusted rate ¹	Number	Rate	Age- adjusted rate ¹	Number	Rate	Age- adjusted rate ¹	Number	Rate	Age- adjusted rate ¹
United States ²	50,476	16.3	15.3	120,859	39.1	38.0	35,332	11.4	11.3	38,364	12.4	12.1
Alabama	1,184	24.8	23.1	2,394	50.1	49.6	931	19.5	19.4	679	14.2	14.0
Alaska	56	7.9	13.5	366	51.5	58.7	71	10.0	10.4	164	23.1	22.8
Arizona	537	8.4	7.9	3,018	47.2	46.7	792	12.4	12.3	1,093	17.1	17.0
Arkansas	737	25.3	22.8	1,461	50.1	49.4	608	20.9	20.7	447	15.3	15.5
California	3,112	8.4	8.7	10,435	28.0	27.8	2,922	7.8	7.7	3,913	10.5	10.3
Colorado	426	8.5	9.6	2,106	41.9	43.5	483	9.6	9.5	865	17.2	16.8
Connecticut	599	16.8	13.4	1,337	37.4	33.6	331	9.3	9.1	353	9.9	9.4
Delaware	158	17.6	15.4	357	39.8	39.2	111	12.4	12.5	106	11.8	11.3
District of Columbia	84	14.0	14.4	212	35.2	35.1	38	6.3	6.0	41	6.8	6.9
Florida	3,297	17.5	12.9	8,875	47.2	43.1	2,536	13.5	13.0	2,789	14.8	13.7
Georgia	1,753	18.1	21.3	3,745	38.7	40.7	1,324	13.7	13.9	1,133	11.7	11.7
Hawaii	208	15.3	12.8	432	31.8	29.6	124	9.1	9.1	207	15.2	15.0
Idaho	186	11.9	12.1	654	41.7	42.1	214	13.7	13.8	290	18.5	18.8
Illinois	2,612	20.4	19.3	3,997	31.2	30.4	1,033	8.1	7.9	1,178	9.2	9.0
Indiana	1,516	23.4	22.0	2,534	39.1	38.5	768	11.8	11.8	864	13.3	13.1
Iowa	323	10.6	8.1	1,273	41.8	37.2	400	13.1	12.7	372	12.2	12.1
Kansas	577	20.2	17.7	1,317	46.2	44.0	482	16.9	16.6	401	14.1	13.9
Kentucky	1,079	24.9	23.8	2,632	60.7	60.5	821	18.9	18.8	631	14.5	14.2
Louisiana	1,218	26.9	27.3	1,999	44.1	44.4	718	15.8	15.8	557	12.3	12.3
Maine	272	20.5	15.8	540	40.7	36.5	171	12.9	12.2	186	14.0	13.2
Maryland	803	13.9	13.6	1,446	25.0	24.7	513	8.9	8.8	502	8.7	8.3
Massachusetts	1,381	21.1	17.5	2,060	31.5	28.5	387	5.9	5.5	598	9.1	8.8
Michigan	1,720	17.4	15.3	3,770	38.1	36.2	1,051	10.6	10.3	1,263	12.8	12.5
Minnesota	897	16.9	15.0	2,103	39.6	36.7	515	9.7	9.5	606	11.4	11.2
Mississippi	738	24.9	24.7	1,685	56.8	56.8	682	23.0	22.9	388	13.1	13.0
Missouri	1,305	21.8	19.2	2,975	49.7	47.9	871	14.5	14.4	856	14.3	14.0
Montana	133	13.4	11.3	548	55.4	53.2	192	19.4	19.6	227	22.9	21.8
Nebraska	290	15.9	13.4	700	38.3	35.8	211	11.6	11.3	193	10.6	10.4
Nevada	471	17.4	19.5	1,088	40.3	41.3	286	10.6	10.7	547	20.3	19.8
New Hampshire	206	15.6	14.2	517	39.3	37.2	135	10.3	10.1	196	14.9	14.1
New Jersey	1,580	18.0	15.6	2,486	28.3	26.7	579	6.6	6.5	719	8.2	7.7
New Mexico	276	13.4	13.1	1,233	59.9	60.7	331	16.1	16.4	413	20.1	20.1
New York	2,439	12.6	11.0	5,004	25.8	24.2	1,323	6.8	6.6	1,547	8.0	7.7
North Carolina	1,892	19.8	19.5	4,144	43.5	43.2	1,383	14.5	14.5	1,174	12.3	12.0
North Dakota	104	15.5	11.5	285	42.4	38.8	97	14.4	14.5	106	15.8	15.6
Ohio	2,066	17.9	15.4	5,124	44.4	42.4	1,251	10.8	10.6	1,439	12.5	12.2
Oklahoma	595	15.9	15.0	2,288	61.0	60.3	716	19.1	19.0	618	16.5	16.5
Oregon	400	10.4	8.9	1,566	40.9	37.8	324	8.5	8.1	685	17.9	17.1
Pennsylvania	2,982	23.5	17.8	5,751	45.3	41.1	1,441	11.3	11.0	1,576	12.4	11.9
Rhode Island	199	18.9	14.5	475	45.1	40.0	91	8.6	8.2	129	12.3	12.3
South Carolina	963	20.8	19.9	2,274	49.2	48.9	812	17.6	17.5	637	13.8	13.5
South Dakota	74	9.1	7.3	393	48.3	44.5	144	17.7	17.3	140	17.2	17.5
Tennessee	983	15.5	14.8	3,539	55.8	54.9	1,099	17.3	17.1	943	14.9	14.6
Texas	3,878	15.4	18.4	9,212	36.6	39.0	3,331	13.2	13.4	2,891	11.5	11.7
Utah	263	9.5	13.1	970	35.1	40.6	272	9.8	10.6	473	17.1	18.3
Vermont	50	8.0	6.7	298	47.6	42.7	76	12.1	11.8	106	16.9	15.7
Virginia	1,595	19.9	20.2	2,527	31.6	31.6	724	9.0	9.0	963	12.0	11.7
Washington	552	8.2	8.0	2,609	38.8	37.6	549	8.2	7.9	957	14.2	13.9
West Virginia	475	25.6	20.3	1,234	66.6	63.7	310	16.7	16.2	279	15.1	14.1
Wisconsin	1,163	20.5	17.5	2,525	44.4	40.8	620	10.9	10.6	793	13.9	13.4
Wyoming	69	12.2	12.5	346	61.4	59.8	138	24.5	23.1	131	23.2	22.4
Puerto Rico	987	26.5	23.7	1,018	27.4	25.9	352	9.5	9.1	286	7.7	7.6
Virgin Islands	14	*	*	25	23.5	22.5	7	*	*	8	*	*
Guam	20	12.5	19.0	45	28.2	36.7	17	*	*	31	19.4	19.2
American Samoa	4	*	*	14	*	*	2	*	*	-	*	*
Northern Marianas	7	*	*	10	*	*	4	*	*	3	*	*

Table 19. Number of deaths, death rates, and age-adjusted death rates for major causes of death: United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, 2010

[Rates per 100,000 population; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates for the United States and each state are based on populations enumerated in the 2010 census as of April 1; see "Technical Notes." Rates for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are based on the 2010 census, estimated as of July 1, 2010; see "Technical Notes." Numbers after causes of death are categories of the International Classification of Diseases, Tenth Revision (ICD-10). The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the ICD-10; see "Technical Notes." For explanation of asterisks preceding cause-of-death codes, see "Technical Notes"]

Area	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)			Alcohol-induced causes ⁴			Drug-induced causes ⁵			Injury by firearms ⁶		
	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹	Number	Rate	Age-adjusted rate ¹
United States ²	16,259	5.3	5.3	25,692	8.3	7.6	40,393	13.1	12.9	31,672	10.3	10.1
Alabama	391	8.2	8.3	267	5.6	5.0	585	12.2	12.6	782	16.4	16.2
Alaska	44	6.2	6.1	122	17.2	16.3	84	11.8	11.9	144	20.3	20.4
Arizona	418	6.5	6.7	987	15.4	14.7	1,141	17.9	18.2	931	14.6	14.6
Arkansas	188	6.4	6.6	179	6.1	5.7	374	12.8	13.3	419	14.4	14.4
California	1,954	5.2	5.1	4,292	11.5	11.2	4,258	11.4	11.1	2,935	7.9	7.7
Colorado	172	3.4	3.4	733	14.6	13.5	676	13.4	13.1	555	11.0	10.8
Connecticut	144	4.0	4.2	192	5.4	4.8	372	10.4	10.3	209	5.8	5.9
Delaware	61	6.8	7.0	74	8.2	7.3	147	16.4	16.8	88	9.8	9.9
District of Columbia	119	19.8	17.2	87	14.5	14.1	90	15.0	15.0	99	16.5	14.2
Florida	1,114	5.9	6.2	1,751	9.3	7.8	3,181	16.9	17.0	2,268	12.1	11.5
Georgia	640	6.6	6.6	630	6.5	6.2	1,124	11.6	11.4	1,223	12.6	12.6
Hawaii	24	1.8	1.7	87	6.4	5.5	154	11.3	11.1	45	3.3	3.2
Idaho	23	1.5	1.5	162	10.3	9.9	184	11.7	12.2	198	12.6	12.8
Illinois	783	6.1	6.1	690	5.4	5.0	1,344	10.5	10.4	1,064	8.3	8.2
Indiana	315	4.9	4.9	435	6.7	6.3	964	14.9	14.9	709	10.9	10.8
Iowa	55	1.8	1.9	233	7.6	6.7	258	8.5	8.8	213	7.0	6.8
Kansas	105	3.7	3.8	201	7.0	6.5	288	10.1	10.2	300	10.5	10.5
Kentucky	199	4.6	4.7	302	7.0	6.2	1,036	23.9	24.2	555	12.8	12.4
Louisiana	541	11.9	12.0	235	5.2	4.8	616	13.6	13.7	864	19.1	19.2
Maine	26	2.0	2.0	132	9.9	7.6	140	10.5	10.7	113	8.5	7.9
Maryland	438	7.6	7.7	291	5.0	4.5	674	11.7	11.4	538	9.3	9.3
Massachusetts	204	3.1	3.2	485	7.4	6.5	836	12.8	12.5	270	4.1	4.1
Michigan	610	6.2	6.4	928	9.4	8.3	1,723	17.4	17.3	1,076	10.9	11.0
Minnesota	112	2.1	2.1	429	8.1	7.3	427	8.1	7.9	365	6.9	6.8
Mississippi	284	9.6	9.8	167	5.6	5.3	353	11.9	12.0	475	16.0	16.1
Missouri	438	7.3	7.5	399	6.7	6.0	1,024	17.1	17.4	846	14.1	14.0
Montana	30	3.0	3.0	147	14.9	13.0	123	12.4	13.3	164	16.6	15.4
Nebraska	60	3.3	3.4	145	7.9	7.4	130	7.1	7.3	152	8.3	8.2
Nevada	164	6.1	6.1	336	12.4	11.4	581	21.5	20.8	395	14.6	14.5
New Hampshire	19	*	*	127	9.6	7.9	164	12.5	12.1	118	9.0	8.2
New Jersey	388	4.4	4.6	548	6.2	5.6	903	10.3	10.1	456	5.2	5.2
New Mexico	152	7.4	7.6	425	20.6	19.7	487	23.7	24.3	301	14.6	14.9
New York	898	4.6	4.6	1,217	6.3	5.7	1,760	9.1	8.8	1,011	5.2	5.1
North Carolina	539	5.7	5.7	669	7.0	6.3	1,125	11.8	11.7	1,123	11.8	11.6
North Dakota	14	*	*	78	11.6	11.4	26	3.9	3.9	65	9.7	9.5
Ohio	569	4.9	5.1	840	7.3	6.5	1,911	16.6	16.7	1,148	10.0	9.9
Oklahoma	214	5.7	5.7	440	11.7	11.1	728	19.4	19.7	538	14.3	14.4
Oregon	115	3.0	2.9	572	14.9	12.9	576	15.0	14.6	458	12.0	11.4
Pennsylvania	689	5.4	5.6	728	5.7	5.0	1,980	15.6	15.8	1,307	10.3	10.1
Rhode Island	27	2.6	2.6	102	9.7	8.2	176	16.7	16.3	49	4.7	4.6
South Carolina	322	7.0	7.1	394	8.5	7.5	697	15.1	15.0	648	14.0	14.0
South Dakota	17	*	*	89	10.9	10.7	48	5.9	6.6	75	9.2	9.2
Tennessee	410	6.5	6.5	542	8.5	7.5	1,132	17.8	17.7	932	14.7	14.4
Texas	1,363	5.4	5.4	1,579	6.3	6.3	2,492	9.9	10.0	2,714	10.8	11.0
Utah	52	1.9	1.9	182	6.6	7.7	457	16.5	17.9	314	11.4	12.2
Vermont	8	*	*	59	9.4	7.9	68	10.9	10.3	70	11.2	10.3
Virginia	373	4.7	4.7	414	5.2	4.7	571	7.1	7.0	875	10.9	10.8
Washington	180	2.7	2.7	850	12.6	11.4	962	14.3	13.8	609	9.1	8.9
West Virginia	86	4.6	5.0	139	7.5	6.3	520	28.1	29.3	273	14.7	14.1
Wisconsin	159	2.8	2.8	500	8.8	7.8	635	11.2	11.0	501	8.8	8.6
Wyoming	9	*	*	80	14.2	13.5	88	15.6	15.5	92	16.3	15.6
Puerto Rico	959	25.8	26.1	200	5.4	4.7	148	4.0	4.1	923	24.8	25.2
Virgin Islands	56	52.7	63.3	15	*	*	2	*	*	53	49.9	59.7
Guam	4	*	*	1	*	*	2	*	*	3	*	*
American Samoa	5	*	*	-	*	*	-	*	*	1	*	*
Northern Marianas	1	*	*	3	*	*	-	*	*	-	*	*

- Quantity zero.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

¹Death rates are affected by the population composition of the area. Age-adjusted death rates should be used for comparisons between areas; for method of computation, see "Technical Notes."

²Excludes data for Puerto Rico, Virgin Islands, Guam, American Samoa and Northern Marianas.

³ICD-10 codes for Motor vehicle accidents are V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.4,V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0, and V89.2; see "Technical Notes."

⁴Causes of death attributable to alcohol-induced mortality include ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15; see "Technical Notes."

⁵Causes of death attributable to drug-induced mortality include ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.1-F11.5,F11.7-F11.9,F12.1-F12.5,F12.7-F12.9,F13.1-F13.5,F13.7-F13.9,F14.1-F14.5,F14.7-F14.9,F15.1-F15.5,F15.7-F15.9,F16.1-F16.5,F16.7-F16.9,F17.3-F17.5,F17.7-F17.9,F18.1-F18.5,F18.7-F18.9,F19.1-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14; see "Technical Notes."

⁶ICD-10 codes for Injury by firearms are *U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24, and Y35.0; see "Technical Notes."

Table 20. Infant, neonatal, and postneonatal mortality rates by race and sex: United States, 1940, 1950, 1960, 1970, and 1975-2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 1,000 live births in specified group. Beginning in 1980, race for live births is tabulated according to race of mother; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards]

Year	All races						All other ¹					
	Both sexes		Male	Female	White ¹		Total ¹		Black ¹		Male	Female
	Both sexes	Male			Female	Both sexes	Male	Female	Both sexes	Male		
Infant mortality rate												
Race of mother ²												
2010	6.15	7.02	5.32	5.20	5.92	4.51	9.28	10.63	7.99	11.63	13.17	10.14
2009	6.39	7.01	5.75	5.30	5.79	4.78	10.02	11.06	8.94	12.64	14.08	11.15
2008	6.61	7.21	5.97	5.55	6.05	5.02	10.16	11.11	9.18	12.74	13.93	11.50
2007	6.75	7.38	6.09	5.64	6.17	5.08	10.55	11.51	9.54	13.24	14.49	11.94
2006	6.69	7.32	6.03	5.56	6.10	4.99	10.60	11.54	9.61	13.29	14.38	12.16
2005	6.87	7.56	6.15	5.73	6.32	5.11	10.92	11.98	9.82	13.73	15.15	12.27
2004	6.79	7.47	6.09	5.66	6.22	5.07	10.92	12.01	9.77	13.79	15.19	12.33
2003	6.85	7.60	6.07	5.72	6.36	5.05	11.09	12.24	9.90	14.01	15.53	12.43
2002	6.97	7.64	6.27	5.79	6.42	5.13	11.41	12.24	10.55	14.36	15.43	13.25
2001	6.85	7.52	6.14	5.65	6.21	5.06	11.33	12.44	10.18	14.02	15.48	12.52
2000	6.91	7.57	6.21	5.68	6.22	5.11	11.44	12.57	10.26	14.09	15.50	12.63
1999	7.06	7.72	6.36	5.77	6.35	5.15	11.94	12.94	10.90	14.56	15.92	13.16
1998	7.20	7.83	6.54	5.95	6.47	5.41	11.92	13.01	10.79	14.31	15.75	12.82
1997	7.23	7.95	6.47	6.03	6.67	5.36	11.76	12.83	10.65	14.16	15.47	12.82
1996	7.32	8.02	6.59	6.07	6.67	5.44	12.18	13.31	11.01	14.68	16.04	13.27
1995	7.59	8.33	6.81	6.29	6.99	5.55	12.61	13.53	11.65	15.12	16.34	13.86
1994	8.02	8.81	7.20	6.57	7.22	5.89	13.47	14.82	12.08	15.83	17.49	14.12
1993	8.37	9.25	7.43	6.82	7.56	6.05	14.07	15.58	12.52	16.52	18.33	14.67
1992	8.52	9.39	7.61	6.92	7.69	6.12	14.44	15.72	13.10	16.85	18.38	15.26
1991	8.94	10.00	7.84	7.30	8.26	6.30	15.07	16.53	13.57	17.57	19.38	15.71
1990	9.22	10.26	8.13	7.56	8.51	6.56	15.52	16.96	14.03	17.96	19.62	16.25
1989	9.81	10.81	8.77	8.08	9.01	7.10	16.33	17.60	15.02	18.61	20.02	17.15
1988	9.95	10.99	8.86	8.36	9.35	7.31	16.08	17.33	14.79	18.54	20.04	16.99
1987	10.08	11.17	8.94	8.48	9.45	7.45	16.46	18.06	14.80	18.75	20.63	16.83
1986	10.35	11.55	9.10	8.80	9.87	7.67	16.72	18.45	14.91	18.90	20.91	16.81
1985	10.64	11.91	9.32	9.17	10.39	7.88	16.84	18.33	15.28	19.01	20.76	17.22
1984	10.79	11.90	9.62	9.30	10.38	8.17	17.05	18.37	15.69	19.15	20.67	17.58
1983	11.16	12.31	9.96	9.61	10.66	8.49	17.80	19.44	16.11	19.98	21.95	17.96
1982	11.52	12.77	10.21	9.94	11.08	8.73	18.31	20.07	16.49	20.48	22.45	18.44
1981	11.93	13.14	10.66	10.34	11.50	9.12	18.82	20.36	17.24	20.81	22.54	19.03
1980	12.60	13.93	11.21	10.86	12.12	9.52	20.19	21.89	18.43	22.19	24.16	20.15
Race of child ³												
1980	12.60	13.93	11.21	11.00	12.27	9.65	19.12	20.73	17.47	21.37	23.27	19.43
1979	13.07	14.50	11.56	11.42	12.82	9.94	19.81	21.47	18.09	21.78	23.66	19.85
1978	13.78	15.26	12.23	12.01	13.37	10.58	21.06	23.15	18.90	23.11	25.39	20.77
1977	14.12	15.75	12.40	12.34	13.90	10.68	21.68	23.71	19.58	23.64	25.91	21.30
1976	15.24	16.82	13.57	13.31	14.81	11.71	23.50	25.51	21.42	25.54	27.83	23.19
1975	16.07	17.86	14.18	14.17	15.94	12.30	24.23	26.24	22.17	26.21	28.32	24.03
1970	20.01	22.37	17.52	17.75	19.95	15.42	30.92	34.20	27.53	32.65	36.18	29.01
1960	26.04	29.33	22.59	22.91	26.01	19.64	43.21	47.88	38.46	44.32	49.12	39.43
1950	29.21	32.75	25.48	26.77	30.21	23.13	44.46	48.87	39.93	43.91	48.27	39.44
1940	47.02	52.45	41.29	43.23	48.32	37.84	73.78	82.21	65.19	72.94	81.07	64.61

Table 20. Infant, neonatal, and postneonatal mortality rates by race and sex: United States, 1940, 1950, 1960, 1970, and 1975-2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 1,000 live births in specified group. Beginning in 1980, race for live births is tabulated according to race of mother; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards]

Year	All races						All other ¹					
	Both sexes		Male	Female	White ¹		Total ¹		Black ¹			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Neonatal mortality rate												
Race of mother ²												
2010	4.05	4.59	3.53	3.46	3.91	3.03	6.00	6.81	5.21	7.49	8.37	6.65
2009	4.18	4.53	3.81	3.48	3.76	3.19	6.48	7.10	5.83	8.17	9.04	7.28
2008	4.29	4.67	3.89	3.62	3.94	3.28	6.54	7.14	5.92	8.23	8.99	7.45
2007	4.42	4.79	4.02	3.70	4.01	3.37	6.86	7.49	6.22	8.65	9.48	7.78
2006	4.45	4.84	4.05	3.72	4.05	3.37	7.00	7.58	6.40	8.82	9.49	8.12
2005	4.54	4.93	4.12	3.79	4.10	3.46	7.18	7.88	6.47	9.07	9.96	8.14
2004	4.52	4.94	4.09	3.78	4.14	3.41	7.19	7.82	6.54	9.13	9.95	8.27
2003	4.62	5.08	4.14	3.87	4.26	3.46	7.40	8.14	6.64	9.40	10.40	8.37
2002	4.66	5.06	4.25	3.89	4.27	3.50	7.55	8.03	7.05	9.51	10.13	8.87
2001	4.54	4.97	4.08	3.78	4.15	3.39	7.37	8.06	6.65	9.21	10.15	8.25
2000	4.63	5.06	4.17	3.82	4.16	3.46	7.60	8.39	6.79	9.38	10.39	8.35
1999	4.73	5.11	4.33	3.88	4.19	3.56	7.94	8.60	7.25	9.77	10.72	8.79
1998	4.80	5.21	4.37	3.98	4.31	3.63	7.91	8.63	7.17	9.55	10.51	8.56
1997	4.77	5.20	4.32	3.99	4.37	3.59	7.74	8.36	7.09	9.40	10.12	8.65
1996	4.77	5.18	4.34	3.97	4.31	3.62	7.86	8.59	7.12	9.56	10.45	8.65
1995	4.91	5.36	4.44	4.08	4.50	3.64	8.13	8.71	7.53	9.85	10.63	9.05
1994	5.12	5.58	4.64	4.20	4.55	3.83	8.60	9.51	7.65	10.21	11.32	9.07
1993	5.29	5.75	4.81	4.29	4.64	3.92	9.02	9.90	8.11	10.69	11.76	9.59
1992	5.37	5.84	4.89	4.35	4.72	3.96	9.19	10.02	8.32	10.83	11.83	9.79
1991	5.59	6.17	4.98	4.53	5.01	4.04	9.52	10.54	8.47	11.25	12.56	9.89
1990	5.85	6.50	5.16	4.79	5.38	4.17	9.86	10.79	8.89	11.55	12.69	10.38
1989	6.23	6.79	5.63	5.15	5.66	4.60	10.30	11.08	9.49	11.92	12.84	10.97
1988	6.32	6.95	5.65	5.27	5.84	4.67	10.33	11.22	9.42	12.05	13.14	10.93
1987	6.46	7.11	5.79	5.40	5.96	4.82	10.68	11.72	9.61	12.30	13.52	11.05
1986	6.71	7.42	5.97	5.72	6.34	5.05	10.79	11.83	9.70	12.31	13.59	10.98
1985	6.96	7.75	6.13	6.00	6.75	5.21	11.00	12.00	9.95	12.62	13.81	11.39
1984	7.00	7.66	6.31	6.09	6.72	5.41	10.87	11.66	10.06	12.32	13.22	11.40
1983	7.28	8.01	6.52	6.31	6.98	5.61	11.41	12.46	10.33	12.93	14.20	11.63
1982	7.70	8.48	6.88	6.69	7.39	5.94	12.04	13.15	10.88	13.62	14.86	12.34
1981	8.02	8.81	7.20	6.99	7.73	6.20	12.51	13.52	11.48	13.98	15.16	12.77
1980	8.48	9.31	7.60	7.39	8.19	6.54	13.21	14.27	12.13	14.62	15.91	13.29
Race of child ³												
1980	8.48	9.31	7.60	7.48	8.29	6.62	12.52	13.51	11.49	14.08	15.32	12.81
1979	8.87	9.79	7.89	7.88	8.80	6.92	12.89	13.91	11.83	14.31	15.45	13.14
1978	9.49	10.54	8.38	8.39	9.34	7.38	14.01	15.54	12.43	15.47	17.17	13.72
1977	9.88	11.00	8.70	8.75	9.83	7.60	14.66	16.02	13.27	16.08	17.60	14.52
1976	10.92	12.03	9.75	9.66	10.73	8.52	16.31	17.68	14.90	17.92	19.47	16.32
1975	11.58	12.91	10.18	10.38	11.70	8.98	16.78	18.21	15.31	18.32	19.78	16.81
1970	15.08	16.96	13.10	13.77	15.55	11.88	21.43	23.87	18.91	22.76	25.37	20.07
1960	18.73	21.24	16.09	17.24	19.66	14.70	26.86	30.04	23.62	27.83	31.13	24.49
1950	20.50	23.34	17.50	19.37	22.18	16.40	27.54	30.76	24.23	27.80	31.09	24.44
1940	28.75	32.56	24.74	27.20	30.85	23.33	39.71	44.87	34.45	39.90	44.78	34.89

Table 20. Infant, neonatal, and postneonatal mortality rates by race and sex: United States, 1940, 1950, 1960, 1970, and 1975-2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 1,000 live births in specified group. Beginning in 1980, race for live births is tabulated according to race of mother; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards]

Year	All races						All other ¹					
	Both sexes		Male	Female	White ¹		Total ¹		Black ¹		Male	Female
	Both sexes	Male			Female	Both sexes	Male	Female	Both sexes	Male		
Postneonatal mortality rate												
Race of mother ²												
2010	2.10	2.43	1.78	1.74	2.01	1.48	3.29	3.81	2.78	4.14	4.80	3.49
2009	2.22	2.48	1.94	1.82	2.04	1.59	3.55	3.96	3.11	4.47	5.05	3.87
2008	2.32	2.54	2.08	1.93	2.12	1.73	3.62	3.97	3.26	4.50	4.93	4.06
2007	2.34	2.58	2.07	1.94	2.16	1.71	3.68	4.02	3.32	4.59	5.01	4.16
2006	2.24	2.48	1.98	1.84	2.05	1.62	3.60	3.96	3.22	4.47	4.89	4.04
2005	2.34	2.63	2.03	1.94	2.22	1.65	3.73	4.10	3.36	4.67	5.19	4.13
2004	2.27	2.53	2.00	1.87	2.07	1.66	3.72	4.19	3.23	4.66	5.24	4.06
2003	2.23	2.52	1.94	1.84	2.09	1.58	3.69	4.10	3.26	4.60	5.13	4.06
2002	2.31	2.58	2.03	1.89	2.15	1.63	3.86	4.21	3.50	4.85	5.30	4.38
2001	2.31	2.55	2.06	1.87	2.06	1.67	3.96	4.37	3.53	4.81	5.32	4.27
2000	2.28	2.51	2.04	1.86	2.06	1.66	3.83	4.18	3.47	4.70	5.11	4.28
1999	2.33	2.61	2.03	1.88	2.16	1.60	4.00	4.34	3.64	4.79	5.20	4.36
1998	2.40	2.62	2.16	1.97	2.16	1.78	4.01	4.38	3.62	4.76	5.24	4.26
1997	2.45	2.75	2.14	2.04	2.30	1.77	4.02	4.47	3.56	4.77	5.34	4.17
1996	2.55	2.84	2.24	2.09	2.36	1.81	4.32	4.72	3.90	5.11	5.60	4.62
1995	2.67	2.97	2.37	2.21	2.49	1.91	4.47	4.82	4.11	5.27	5.71	4.81
1994	2.90	3.22	2.56	2.37	2.67	2.06	4.88	5.32	4.42	5.61	6.17	5.04
1993	3.07	3.50	2.62	2.54	2.92	2.13	5.06	5.68	4.42	5.83	6.57	5.08
1992	3.14	3.55	2.72	2.58	2.97	2.16	5.25	5.69	4.78	6.02	6.54	5.47
1991	3.35	3.82	2.86	2.76	3.25	2.26	5.55	5.99	5.10	6.32	6.82	5.81
1990	3.38	3.76	2.97	2.78	3.14	2.39	5.66	6.16	5.13	6.41	6.93	5.87
1989	3.59	4.01	3.14	2.93	3.35	2.49	6.03	6.52	5.53	6.69	7.18	6.19
1988	3.64	4.04	3.21	3.09	3.51	2.65	5.75	6.11	5.37	6.49	6.90	6.07
1987	3.62	4.06	3.15	3.08	3.49	2.64	5.77	6.34	5.18	6.45	7.10	5.77
1986	3.64	4.13	3.13	3.08	3.53	2.62	5.93	6.62	5.21	6.59	7.33	5.83
1985	3.68	4.15	3.19	3.17	3.64	2.67	5.84	6.33	5.33	6.40	6.95	5.83
1984	3.79	4.23	3.31	3.22	3.65	2.76	6.18	6.71	5.63	6.83	7.46	6.18
1983	3.88	4.30	3.44	3.29	3.68	2.88	6.39	6.98	5.78	7.05	7.75	6.32
1982	3.82	4.29	3.33	3.25	3.68	2.79	6.28	6.92	5.61	6.86	7.59	6.10
1981	3.91	4.34	3.46	3.35	3.77	2.92	6.31	6.84	5.76	6.83	7.38	6.26
1980	4.13	4.62	3.61	3.47	3.93	2.98	6.97	7.62	6.30	7.57	8.25	6.87
Race of child ³												
1980	4.13	4.62	3.61	3.52	3.98	3.02	6.61	7.22	5.97	7.29	7.95	6.62
1979	4.20	4.71	3.67	3.54	4.02	3.03	6.92	7.57	6.25	7.47	8.21	6.71
1978	4.30	4.72	3.85	3.63	4.03	3.20	7.05	7.60	6.48	7.64	8.22	7.05
1977	4.24	4.75	3.71	3.59	4.07	3.08	7.01	7.69	6.31	7.56	8.32	6.78
1976	4.32	4.79	3.83	3.65	4.08	3.19	7.19	7.83	6.52	7.63	8.36	6.88
1975	4.49	4.95	4.00	3.80	4.24	3.33	7.45	8.03	6.86	7.89	8.54	7.22
1970	4.93	5.41	4.42	3.98	4.40	3.54	9.49	10.33	8.62	9.89	10.81	8.94
1960	7.31	8.10	6.49	5.66	6.35	4.94	16.35	17.84	14.84	16.48	17.99	14.95
1950	8.71	9.41	7.98	7.40	8.04	6.73	16.92	18.11	15.70	16.10	17.18	15.00
1940	18.27	19.89	16.55	16.03	17.47	14.50	34.07	37.35	30.74	33.05	36.29	29.72

¹Multiple-race data were reported for deaths by 37 states and the District of Columbia in 2010, by 34 states and the District of Columbia in 2008 and 2009, by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2005, by 15 states in 2004, and by 7 states in 2003; see "Technical Notes." Multiple-race data were reported for births by 38 states and the District of Columbia in 2010, by 32 states and the District of Columbia in 2009, by 30 areas in 2008, by 27 areas in 2007, by 23 areas in 2006, by 19 areas in 2005, by 15 areas in 2004, and by 6 areas in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Infant deaths are based on race of child as stated on the death certificate; live births are based on race of mother as stated on the birth certificate; see "Technical Notes."

³Infant deaths are based on race of child as stated on the death certificate; live births are based on race of parents as stated on the birth certificate; see "Technical Notes."

Table 21. Number of infant deaths and infant mortality rates for 130 selected causes by race: United States, 2010

[Rates are infant deaths (under 1 year) per 100,000 live births in specified group. Infant deaths are based on race of decedent; live births are based on race of mother. The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Number			Rate		
	All races ¹	White ²	Black ²	All races ¹	White ²	Black ²
All causes	24,586	15,954	7,401	614.7	519.8	1,162.9
Certain infectious and parasitic diseases (A00-B99)	696	401	266	17.4	13.1	41.8
Certain intestinal infectious diseases (A00-A08)	7	4	2	*	*	*
Diarrhea and gastroenteritis of infectious origin (A09)	316	170	133	7.9	5.5	20.9
Tuberculosis (A16-A19)	-	-	-	*	*	*
Tetanus (A33,A35)	-	-	-	*	*	*
Diphtheria (A36)	-	-	-	*	*	*
Whooping cough (A37)	25	23	2	0.6	0.7	*
Meningococcal infection (A39)	11	9	2	*	*	*
Septicemia (A40-A41)	215	114	90	5.4	3.7	14.1
Congenital syphilis (A50)	2	1	1	*	*	*
Gonococcal infection (A54)	-	-	-	*	*	*
Viral diseases (A80-B34)	92	58	30	2.3	1.9	4.7
Acute poliomyelitis (A80)	-	-	-	*	*	*
Varicella (chickenpox) (B01)	-	-	-	*	*	*
Measles (B05)	-	-	-	*	*	*
Human immunodeficiency virus (HIV) disease (B20-B24)	-	-	-	*	*	*
Mumps (B26)	-	-	-	*	*	*
Other and unspecified viral diseases (A81-B00,B02-B04,B06-B19,B25,B27-B34)	92	58	30	2.3	1.9	4.7
Candidiasis (B37)	6	5	1	*	*	*
Malaria (B50-B54)	-	-	-	*	*	*
Pneumocystosis (B59)	-	-	-	*	*	*
All other and unspecified infectious and parasitic diseases (A20-A32,A38,A42-A49,A51-A53,A55-A79,B35-B36,B38-B49,B55-B58,B60-B99)	22	17	5	0.6	*	*
Neoplasms (C00-D48)	110	84	16	2.8	2.7	*
Malignant neoplasms (C00-C97)	62	47	8	1.6	1.5	*
Hodgkin's disease and non-Hodgkin's lymphomas (C81-C85)	2	2	-	*	*	*
Leukemia (C91-C95)	25	19	4	0.6	*	*
Other and unspecified malignant neoplasms (C00-C80,C88,C90,C96-C97)	35	26	4	0.9	0.8	*
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	37	8	1.2	1.2	*
Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	95	65	26	2.4	2.1	4.1
Anemias (D50-D64)	15	5	9	*	*	*
Hemorrhagic conditions and other diseases of blood and blood-forming organs (D65-D76)	60	43	14	1.5	1.4	*
Certain disorders involving the immune mechanism (D80-D89)	20	17	3	0.5	*	*
Endocrine, nutritional and metabolic diseases (E00-E88)	188	135	43	4.7	4.4	6.8
Short stature, not elsewhere classified (E34.3)	2	2	-	*	*	*
Nutritional deficiencies (E40-E64)	3	2	-	*	*	*
Cystic fibrosis (E84)	5	5	-	*	*	*
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	48	27	19	1.2	0.9	*
All other endocrine, nutritional and metabolic diseases (E00-E32,E34.0-E34.2,E34.4-E34.9,E65-E83,E85,E88)	130	99	24	3.3	3.2	3.8
Diseases of the nervous system (G00-G98)	345	244	75	8.6	7.9	11.8
Meningitis (G00,G03)	58	33	19	1.5	1.1	*
Infantile spinal muscular atrophy, type I (Werdnig-Hoffman) (G12.0)	4	4	-	*	*	*
Infantile cerebral palsy (G80)	3	2	1	*	*	*
Anoxic brain damage, not elsewhere classified (G93.1)	39	25	11	1.0	0.8	*
Other diseases of nervous system (G04,G06-G11,G12.1-G12.9,G20-G72,G81-G92,G93.0,G93.2-G93.9,G95-G98)	241	180	44	6.0	5.9	6.9
Diseases of the ear and mastoid process (H60-H93)	3	2	-	*	*	*
Diseases of the circulatory system (I00-I99)	507	326	152	12.7	10.6	23.9
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)	90	52	32	2.3	1.7	5.0
Pericarditis, endocarditis and myocarditis (I30,I33,I40)	14	9	4	*	*	*
Cardiomyopathy (I42)	79	62	12	2.0	2.0	*
Cardiac arrest (I46)	18	11	6	*	*	*
Cerebrovascular diseases (I60-I69)	130	74	48	3.3	2.4	7.5
All other diseases of circulatory system (I00-I25,I31,I34-I38,I44-I45,I47-I51,I70-I99)	176	118	50	4.4	3.8	7.9
Diseases of the respiratory system (J00-J98)	574	306	221	14.4	10.0	34.7
Acute upper respiratory infections (J00-J06)	15	12	2	*	*	*
Influenza and pneumonia (J09-J18)	195	101	73	4.9	3.3	11.5
Influenza (J09-J11)	16	6	6	*	*	*
Pneumonia (J12-J18)	179	95	67	4.5	3.1	10.5
Acute bronchitis and acute bronchiolitis (J20-J21)	27	14	10	0.7	*	*
Bronchitis, chronic and unspecified (J40-J42)	25	13	6	0.6	*	*

Table 21. Number of infant deaths and infant mortality rates for 130 selected causes by race: United States, 2010

[Rates are infant deaths (under 1 year) per 100,000 live births in specified group. Infant deaths are based on race of decedent; live births are based on race of mother. The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Number			Rate		
	All races ¹	White ²	Black ²	All races ¹	White ²	Black ²
Asthma (J45-J46)	6	5	1	*	*	*
Pneumonitis due to solids and liquids (J69)	18	8	9	*	*	*
Other and unspecified diseases of respiratory system (J22,J30-J39,J43-J44,J47-J68,J70-J98,U04)	288	153	120	7.2	5.0	18.9
Diseases of the digestive system (K00-K92)	204	141	53	5.1	4.6	8.3
Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	29	20	8	0.7	0.7	*
Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	51	39	11	1.3	1.3	*
All other and unspecified diseases of digestive system (K00-K28,K30-K38,K57-K92)	124	82	34	3.1	2.7	5.3
Diseases of the genitourinary system (N00-N95)	126	73	42	3.2	2.4	6.6
Renal failure and other disorders of kidney (N17-N19,N25,N27)	100	56	35	2.5	1.8	5.5
Other and unspecified diseases of genitourinary system (N00-N15,N20-N23,N26,N28-N95)	26	17	7	0.7	*	*
Certain conditions originating in the perinatal period (P00-P96)	12,008	7,423	4,005	300.2	241.8	629.3
Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)	2,920	1,842	922	73.0	60.0	144.9
Newborn affected by maternal hypertensive disorders (P00.0)	85	49	34	2.1	1.6	5.3
Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	87	56	27	2.2	1.8	4.2
Newborn affected by maternal complications of pregnancy (P01)	1,561	991	486	39.0	32.3	76.4
Newborn affected by incompetent cervix (P01.0)	431	242	154	10.8	7.9	24.2
Newborn affected by premature rupture of membranes (P01.1)	781	520	225	19.5	16.9	35.4
Newborn affected by multiple pregnancy (P01.5)	163	99	60	4.1	3.2	9.4
Newborn affected by other maternal complications of pregnancy (P01.2-P01.4,P01.6-P01.9)	186	130	47	4.7	4.2	7.4
Newborn affected by complications of placenta, cord and membranes (P02)	1,030	643	327	25.8	20.9	51.4
Newborn affected by complications involving placenta (P02.0-P02.3)	492	334	121	12.3	10.9	19.0
Newborn affected by complications involving cord (P02.4-P02.6)	39	26	10	1.0	0.8	*
Newborn affected by chorioamnionitis (P02.7)	497	281	196	12.4	9.2	30.8
Newborn affected by other and unspecified abnormalities of membranes (P02.8-P02.9)	2	2	-	*	*	*
Newborn affected by other complications of labor and delivery (P03)	110	73	33	2.8	2.4	5.2
Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	47	30	15	1.2	1.0	*
Disorders related to length of gestation and fetal malnutrition (P05-P08)	4,233	2,362	1,669	105.8	77.0	262.2
Slow fetal growth and fetal malnutrition (P05)	85	48	34	2.1	1.6	5.3
Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,148	2,314	1,635	103.7	75.4	256.9
Extremely low birth weight or extreme immaturity (P07.0,P07.2)	3,176	1,770	1,248	79.4	57.7	196.1
Other low birth weight or preterm (P07.1,P07.3)	972	544	387	24.3	17.7	60.8
Disorders related to long gestation and high birth weight (P08)	-	-	-	*	*	*
Birth trauma (P10-P15)	19	12	4	*	*	*
Intrauterine hypoxia and birth asphyxia (P20-P21)	314	214	76	7.9	7.0	11.9
Intrauterine hypoxia (P20)	136	92	35	3.4	3.0	5.5
Birth asphyxia (P21)	178	122	41	4.5	4.0	6.4
Respiratory distress of newborn (P22)	514	318	176	12.9	10.4	27.7
Other respiratory conditions originating in the perinatal period (P23-P28)	812	556	221	20.3	18.1	34.7
Congenital pneumonia (P23)	71	49	22	1.8	1.6	3.5
Neonatal aspiration syndromes (P24)	51	42	6	1.3	1.4	*
Interstitial emphysema and related conditions originating in the perinatal period (P25)	106	83	22	2.7	2.7	3.5
Pulmonary hemorrhage originating in the perinatal period (P26)	167	98	60	4.2	3.2	9.4
Chronic respiratory disease originating in the perinatal period (P27)	106	58	40	2.7	1.9	6.3
Atelectasis (P28.0-P28.1)	248	180	58	6.2	5.9	9.1
All other respiratory conditions originating in the perinatal period (P28.2-P28.9)	63	46	13	1.6	1.5	*
Infections specific to the perinatal period (P35-P39)	745	480	234	18.6	15.6	36.8
Bacterial sepsis of newborn (P36)	583	382	176	14.6	12.4	27.7
Omphalitis of newborn with or without mild hemorrhage (P38)	1	1	-	*	*	*
All other infections specific to the perinatal period (P35,P37,P39)	161	97	58	4.0	3.2	9.1
Hemorrhagic and hematological disorders of newborn (P50-P61)	556	396	140	13.9	12.9	22.0
Neonatal hemorrhage (P50-P52,P54)	469	337	118	11.7	11.0	18.5
Hemorrhagic disease of newborn (P53)	1	1	-	*	*	*
Hemolytic disease of newborn due to isoimmunization and other perinatal jaundice (P55-P59)	7	5	-	*	*	*
Hematological disorders (P60-P61)	79	53	22	2.0	1.7	3.5
Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0-P70.2)	3	2	1	*	*	*
Necrotizing enterocolitis of newborn (P77)	472	282	174	11.8	9.2	27.3
Hydrops fetalis not due to hemolytic disease (P83.2)	150	122	15	3.8	4.0	*
Other perinatal conditions (P29,P70.3-P70.9,P71-P76,P78-P81,P83.0-P83.1,P83.3-P83.9,P90-P96)	1,270	837	373	31.8	27.3	58.6
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5,107	3,822	1,010	127.7	124.5	158.7

Table 21. Number of infant deaths and infant mortality rates for 130 selected causes by race: United States, 2010

[Rates are infant deaths (under 1 year) per 100,000 live births in specified group. Infant deaths are based on race of decedent; live births are based on race of mother. The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	Number			Rate		
	All races ¹	White ²	Black ²	All races ¹	White ²	Black ²
Anencephaly and similar malformations (Q00)	293	246	33	7.3	8.0	5.2
Congenital hydrocephalus (Q03)	105	79	17	2.6	2.6	*
Spina bifida (Q05)	15	11	4	*	*	*
Other congenital malformations of nervous system (Q01-Q02,Q04,Q06-Q07)	318	242	58	8.0	7.9	9.1
Congenital malformations of heart (Q20-Q24)	1,148	861	233	28.7	28.1	36.6
Other congenital malformations of circulatory system (Q25-Q28)	176	120	42	4.4	3.9	6.6
Congenital malformations of respiratory system (Q30-Q34)	399	284	95	10.0	9.3	14.9
Congenital malformations of digestive system (Q35-Q45)	88	58	18	2.2	1.9	*
Congenital malformations of genitourinary system (Q50-Q64)	457	346	88	11.4	11.3	13.8
Congenital malformations and deformations of musculoskeletal system, limbs and integument (Q65-Q85)	577	439	106	14.4	14.3	16.7
Down's syndrome (Q90)	85	60	16	2.1	2.0	*
Edward's syndrome (Q91.0-Q91.3)	470	363	83	11.8	11.8	13.0
Patau's syndrome (Q91.4-Q91.7)	244	170	63	6.1	5.5	9.9
Other congenital malformations and deformations (Q10-Q18,Q86-Q89)	542	397	116	13.6	12.9	18.2
Other chromosomal abnormalities, not elsewhere classified (Q92-Q99)	190	146	38	4.8	4.8	6.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	3,052	1,930	994	76.3	62.9	156.2
Sudden infant death syndrome (R95)	2,063	1,311	668	51.6	42.7	105.0
Other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94,R96-R99)	989	619	326	24.7	20.2	51.2
All other diseases (Residual)	20	12	8	0.5	*	*
External causes of mortality (*U01,V01-Y84)	1,551	990	490	38.8	32.3	77.0
Accidents (unintentional injuries) (V01-X59)	1,110	703	354	27.8	22.9	55.6
Transport accidents (V01-V99)	81	60	16	2.0	2.0	*
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)	79	58	16	2.0	1.9	*
Other and unspecified transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9,V90-V99)	2	2	-	*	*	*
Falls (W00-W19)	10	5	5	*	*	*
Accidental discharge of firearms (W32-W34)	-	-	-	*	*	*
Accidental drowning and submersion (W65-W74)	39	31	8	1.0	1.0	*
Accidental suffocation and strangulation in bed (W75)	629	386	217	15.7	12.6	34.1
Other accidental suffocation and strangulation (W76-W77,W81-W84)	218	134	74	5.5	4.4	11.6
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)	58	33	17	1.5	1.1	*
Accidents caused by exposure to smoke, fire and flames (X00-X09)	21	17	4	0.5	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)	6	4	1	*	*	*
Other and unspecified accidents (W20-W31,W35-W64,W85-W99,X10-X39,X50-X59)	48	33	12	1.2	1.1	*
Assault (homicide) (*U01,X85-Y09)	311	213	88	7.8	6.9	13.8
Assault (homicide) by hanging, strangulation and suffocation (X91)	15	14	1	*	*	*
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	11	8	3	*	*	*
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)	82	47	32	2.1	1.5	5.0
Assault (homicide) by other and unspecified means (*U01.0-*U01.3,*U01.5-*U01.9,X85-X90,X92,X96-X99,Y00-Y05,Y08-Y09)	203	144	52	5.1	4.7	8.2
Complications of medical and surgical care (Y40-Y84)	22	15	5	0.6	*	*
Other external causes (Y10-Y36)	108	59	43	2.7	1.9	6.8

* Figure does not meet standards of reliability or precision; see "Technical Notes."

- Quantity zero.

¹Includes races other than white and black.

²Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported, for deaths, by 37 states and the District of Columbia and, for births, by 38 states and the District of Columbia; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

Table 22. Number of infant and neonatal deaths and mortality rates, by race for the United States, each state, Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas, and by sex for the United States, 2010

[Rates are infant (under 1 year) and neonatal (under 28 days) deaths per 1,000 live births in specified group. Infant deaths are based on race of decedent; live births are based on race of mother. See "Technical Notes"]

Sex and area	Infant deaths						Neonatal deaths					
	All races ¹		White ²		Black ²		All races ¹		White ²		Black ²	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
United States ³	24,586	6.15	15,954	5.20	7,401	11.63	16,188	4.05	10,612	3.46	4,769	7.49
Male	13,702	7.02	8,871	5.92	4,116	13.17	8,953	4.59	5,856	3.91	2,616	8.37
Female	10,884	5.32	7,083	4.51	3,285	10.14	7,235	3.53	4,756	3.03	2,153	6.65
Alabama	523	8.71	269	6.63	250	13.66	326	5.43	159	3.92	164	8.96
Alaska	43	3.75	23	3.26	5	*	22	1.92	13	*	5	*
Arizona	522	5.97	403	5.47	54	12.14	332	3.80	262	3.56	35	7.87
Arkansas	282	7.32	201	6.66	77	10.34	160	4.15	109	3.61	47	6.31
California	2,420	4.74	1,853	4.58	324	9.84	1,682	3.30	1,320	3.26	200	6.07
Colorado	392	5.91	330	5.55	47	13.13	287	4.33	239	4.02	36	10.06
Connecticut	199	5.28	134	4.50	61	11.59	150	3.98	104	3.49	45	8.55
Delaware	87	7.66	49	6.36	37	11.87	58	5.10	30	3.89	27	8.66
District of Columbia	72	7.86	16	*	56	10.48	50	5.46	11	*	39	7.30
Florida	1,403	6.54	773	5.04	610	11.33	934	4.35	514	3.35	406	7.54
Georgia	860	6.42	397	5.05	448	9.11	523	3.90	247	3.14	264	5.37
Hawaii	117	6.16	23	3.92	11	*	76	4.00	14	*	9	*
Idaho	112	4.83	105	4.75	1	*	63	2.72	61	2.76	1	*
Illinois	1,118	6.77	680	5.39	394	13.58	769	4.65	496	3.93	238	8.20
Indiana	640	7.62	476	6.65	153	14.96	417	4.97	321	4.48	87	8.51
Iowa	189	4.88	163	4.61	22	11.18	103	2.66	89	2.52	12	*
Kansas	253	6.22	204	5.71	40	12.35	174	4.28	139	3.89	28	8.65
Kentucky	379	6.79	316	6.43	61	11.12	185	3.32	160	3.25	24	4.38
Louisiana	474	7.60	176	4.86	287	11.75	265	4.25	96	2.65	163	6.67
Maine	70	5.40	66	5.39	3	*	49	3.78	46	3.76	2	*
Maryland	498	6.75	184	4.35	297	11.55	350	4.74	127	3.00	212	8.25
Massachusetts	323	4.43	231	4.03	68	7.22	241	3.31	172	3.00	50	5.31
Michigan	817	7.13	481	5.51	318	14.08	550	4.80	321	3.68	219	9.69
Minnesota	308	4.49	231	4.20	44	6.35	209	3.05	157	2.86	29	4.18
Mississippi	387	9.67	144	6.61	242	13.80	220	5.50	75	3.44	145	8.27
Missouri	507	6.61	362	5.78	139	11.84	318	4.14	220	3.51	93	7.92
Montana	71	5.89	61	5.86	-	*	42	3.48	37	3.55	-	*
Nebraska	136	5.25	107	4.72	26	13.40	96	3.70	72	3.18	21	10.82
Nevada	201	5.59	137	4.76	52	13.83	126	3.51	83	2.88	36	9.57
New Hampshire	51	3.96	47	3.90	3	*	27	2.10	25	2.07	1	*
New Jersey	514	4.81	275	3.64	202	10.39	369	3.45	196	2.60	142	7.30
New Mexico	157	5.64	126	5.48	5	*	95	3.41	79	3.43	2	*
New York	1,243	5.09	731	4.24	439	9.21	866	3.54	527	3.05	285	5.98
North Carolina	858	7.01	461	5.35	367	12.28	603	4.93	320	3.71	265	8.87
North Dakota	62	6.81	42	5.43	3	*	46	5.05	33	4.26	3	*
Ohio	1,072	7.71	695	6.27	360	14.78	728	5.23	486	4.38	230	9.44
Oklahoma	404	7.59	255	6.28	62	12.38	227	4.26	143	3.52	35	6.99
Oregon	225	4.94	192	4.72	14	*	153	3.36	129	3.17	9	*
Pennsylvania	1,039	7.25	696	6.28	325	12.62	736	5.14	503	4.54	218	8.47
Rhode Island	79	7.07	60	6.67	17	*	59	5.28	44	4.89	13	*
South Carolina	430	7.37	199	5.32	221	11.32	266	4.56	120	3.21	138	7.07
South Dakota	82	6.94	57	6.03	1	*	55	4.66	39	4.12	-	*
Tennessee	630	7.93	393	6.55	232	13.28	367	4.62	222	3.70	141	8.07
Texas	2,368	6.13	1,788	5.58	533	11.23	1,508	3.91	1,162	3.62	320	6.74
Utah	254	4.86	223	4.53	8	*	176	3.37	161	3.27	3	*
Vermont	26	4.18	26	4.35	-	*	14	*	14	*	-	*
Virginia	700	6.80	347	4.81	328	14.27	475	4.61	233	3.23	224	9.75
Washington	389	4.50	283	4.02	40	8.22	266	3.07	194	2.76	27	5.55
West Virginia	149	7.28	138	7.08	11	*	81	3.96	74	3.79	7	*
Wisconsin	400	5.84	280	4.88	102	14.61	263	3.84	185	3.22	69	9.88
Wyoming	51	6.75	45	6.37	1	*	31	4.10	29	4.10	-	*
Puerto Rico	341	8.09	330	8.75	10	*	242	5.74	233	6.18	8	*
Virgin Islands	13	*	3	*	10	*	9	*	2	*	7	*
Guam	48	14.05	1	*	-	*	29	8.49	-	*	-	*
American Samoa	14	*	-	*	-	*	8	*	-	*	-	*
Northern Marianas	4	*	-	*	-	*	2	*	-	*	-	*

- Quantity zero.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

¹Includes races other than white and black.

²Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported for deaths by 37 states and the District of Columbia and, for births, by 38 states and the District of Columbia; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

³Excludes data for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas.