



Title	Differentiating Neural Stem Cells into Astrocytes
Date Submitted	May 5, 2012
Submitted by -	Efthymiou, Anastasia - anastasia.efthymiou@nih.gov
Adapted from -	Gibco Protocol
Contributors -	Efthymiou, Anastasia
Affiliation(s) -	NIH CRM - NIAMS – Laboratory of Stem Cell Biology

❖ Introduction:



Human astrocyte, courtesy of Dr. Riccardo Cassiani-Ingoni, NIH NINDS

❖ Protocol:

Neural stem cells (NSCs) will proliferate as progenitors a few times even after the complete growth medium is replaced with the appropriate differentiation

medium. If the cells reach 90% confluency, it might be necessary to split the cells at a 1:2 ratio. However, do not split the cells once they reach day 9-10 of differentiation when they can get damaged during the passaging process.

1. Plate the NSCs on a Geltrex - coated culture dish in complete StemPro NSC SFM at 2.5×10^4 cells/cm².
2. After 2 days, change medium to astrocyte differentiation medium. Change the spent medium every 3 to 4 days.

❖ **Materials:**

Geltrex-coated culture dish		
StemPro NSC SFM complete Medium		
astrocyte differentiation medium		
StemPro NSC SFM Complete Media		
Component	Final concentration	Amount
KnockOut™ D-MEM/F-12	1X	97 mL
GlutaMAX™-I Supplement	2 mM	1 mL
bFGF (prep as 100 µg/mL stock)	20 ng/mL	20 µL
EGF (prep as 100 µg/mL stock)	20 ng/mL	20 µL
StemPro® Neural Supplement	2%	2 mL
Astrocyte Differentiation Medium		
Component	Final concentration	Amount
D-MEM	1X	97 mL
N-2 Supplement	1%	1 mL
GlutaMAX™-I Supplement	2 mM	1 mL
FBS	1%	1 mL

❖ Troubleshooting:

❖ **References:**