## UDS Summary Report - 2011 Grantee - Universal

| GRANTEE  |               |  |  |
|--|---------------|--|--|
| PATIENTS   |               |  |  |
| Total Patients   | 91,187        |  |  |
| Number/Percent of Patients by Services Utilized          |               |  |  |
| Medical Services   | 69,191/75.88% |  |  |
| Dental Services  | 23,329/25.58% |  |  |
| Mental Health Services                                   | 1,054/1.16%   |  |  |
| Substance Abuse Services                                 | -/-           |  |  |
| Vision Services  | -/-           |  |  |
| Enabling Services  | 1,941/2.13%   |  |  |
| Number/Percent of Patients by Special Populations        |               |  |  |
| Homeless Patients  | 1,035/1.14%   |  |  |
| Farmworker Patients                                      | 33/0.04%      |  |  |
| School Based Patients                                    | 1,675/1.84%   |  |  |
| Veteran Patients   | 222/0.24%     |  |  |
| Number/Percent of Patients by Age                        |               |  |  |
| Children (< 18 years old)                                | 32,241/35.36% |  |  |
| Adult (18 - 64)  | 51,520/56.50% |  |  |
| Geriatric (age 65 and over)                              | 7,426/8.14%   |  |  |
| Number/Percent of Patients by Insurance Status           |               |  |  |
| Uninsured  | 9,090/9.97%   |  |  |
| Children Uninsured (age 0-19 years)                      | 2,827/7.97%   |  |  |
| Medicaid/CHIP <sup>1</sup>                               | 32,025/35.12% |  |  |
| Medicare   | 8,538/9.36%   |  |  |
| Other Third Party  | 41,534/45.55% |  |  |
| Patients by Income Status (% Known)                      |               |  |  |
| Patients at or below 200% of poverty                     | 92.17%        |  |  |
| Patients at or below 100% of poverty (included in above) | 69.66%        |  |  |
| Patients by Race & Ethnicity (% Known)                   |               |  |  |
| % Racial and/or Ethnic Minority                          | 57.91%        |  |  |
| % Hispanic/Latino Identity                               | 16.24%        |  |  |
| % African American                                       | 43.98%        |  |  |
| % Asian  | 2.94%         |  |  |
| % American Indian/Alaska Native                          | 0.22%         |  |  |
| % Native Hawaiian/Other Pacific Islander                 | 0.40%         |  |  |
| % More than one race                                     | 1.44%         |  |  |
| % Best Served in another language                        | 13.53%        |  |  |
| L. Date Connet he Color/Jated                            |               |  |  |

'-' - Data Cannot be Calculated

<sup>1</sup> Includes Medicaid, Medicaid CHIP, and Other Public Insurance CHIP

|   | GRANTEE |
|---|---------|
| PATIENTS  |         |
| Percent of Medical Patients with Specific Diagnoses |         |
| Hypertension <sup>2</sup>                           | 15.91%  |
| Diabetes <sup>3</sup>                               | 10.02%  |
| Asthma  | 6.95%   |
| HIV   | 0.22%   |
| Prenatal  |         |
| Number of Prenatal Patients                         | 1,900   |
| Number of Prenatal Patients who delivered           | 1,292   |
|   |         |

GRANTEE

ADJUSTED QUARTILE RANKING <sup>5</sup>

## QUALITY OF CARE/HEALTH OUTCOMES

| % Having First Prenatal Visit in 1st Trimester66 <th>Prenatal</th> <th></th> <th></th>   | Prenatal   |        |   |  |
|--|--|--------|---|--|
| Preventive Screening & Immunizations     % of Women with Pap Tests   75.71%   1     % of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity<br>Documented   80.00%   1     % of Adults who Received Weight Screening and Follow up if Appropriate   48.57%   2     % of Adults Assessed for Tobacco Use   84.29%   3     % of Asthmatics Treated with Appropriate Pharmacological Intervention   75.71%   2     % of Ary oy ear Olds Immunized <sup>4</sup> 1   3     % hypertensive Patients with Blood Pressure < 140/90   | % Having First Prenatal Visit in 1st Trimester                         | 72.89% | 3 |  |
| % of Women with Pap Tests75.71%1% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity<br>Documented80.00%80.00%1% of Adults who Received Weight Screening and Follow up if Appropriate48.57%48.57%2% of Adults Assessed for Tobacco Use84.29%33% of Tobacco Users who Received Cessation Advice and/or Medication75.71%23% of Asthmatics Treated with Appropriate Pharmacological Intervention75.71%33% of Two year Olds Immunized 471.43%11Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90   | % Low and Very Low Birth Weight  | 10.79% | 4 |  |
| No<br>DocumentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNumber<br>CommentedNum | Preventive Screening & Immunizations                                   |        |   |  |
| Documented80.00%80.00%1% of Adults who Received Weight Screening and Follow up if Appropriate48.57%2% of Adults Assessed for Tobacco Use84.29%3% of Tobacco Users who Received Cessation Advice and/or Medication75.71%2% of Asthmatics Treated with Appropriate Pharmacological Intervention75.71%3% of Two year Olds Immunized 471.43%1Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90  | % of Women with Pap Tests  | 75.71% | 1 |  |
| % of Adults Assessed for Tobacco Use84.29%84.29%3% of Tobacco Users who Received Cessation Advice and/or Medication75.71%2% of Asthmatics Treated with Appropriate Pharmacological Intervention75.71%3% of Two year Olds Immunized 471.43%1Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90  | · · · · · · · · · · · · · · · · · · ·                                  | 80.00% | 1 |  |
| % of Tobacco Users who Received Cessation Advice and/or Medication 75.71% 2   % of Asthmatics Treated with Appropriate Pharmacological Intervention 75.71% 3   % of Two year Olds Immunized <sup>4</sup> 71.43% 1   Chronic Disease Management   % Hypertensive Patients with Blood Pressure < 140/90  | % of Adults who Received Weight Screening and Follow up if Appropriate | 48.57% | 2 |  |
| % of Asthmatics Treated with Appropriate Pharmacological Intervention   75.71%   3     % of Two year Olds Immunized <sup>4</sup> 71.43%   1     Chronic Disease Management     % Hypertensive Patients with Blood Pressure < 140/90  | % of Adults Assessed for Tobacco Use                                   | 84.29% | 3 |  |
| % of Two year Olds Immunized 4 71.43% 1   Chronic Disease Management 57.59% 3  | % of Tobacco Users who Received Cessation Advice and/or Medication     | 75.71% | 2 |  |
| Chronic Disease Management   % Hypertensive Patients with Blood Pressure < 140/90  | % of Asthmatics Treated with Appropriate Pharmacological Intervention  | 75.71% | 3 |  |
| % Hypertensive Patients with Blood Pressure < 140/90 57.59% 3  | % of Two year Olds Immunized <sup>4</sup>                              | 71.43% | 1 |  |
|  | Chronic Disease Management   |        |   |  |
| % Diabetic Patients with HbA1c <= 9 72.08% 3   | % Hypertensive Patients with Blood Pressure < 140/90                   | 57.59% | 3 |  |
|  | % Diabetic Patients with HbA1c <= 9                                    | 72.08% | 3 |  |

'-' - Data Cannot be Calculated

<sup>2</sup> Hypertensive adults as a percent of estimated adult medical patients of ages 18-85.

<sup>3</sup> Diabetic adults as a percent of estimated adult medical patients of ages 18-75.

 $^{\rm 4}$  Measure was revised in 2011 and is not comparable to calendar year 2010 and prior.

<sup>5</sup> Grantee adjusted quartile results from a statistical model adjusting for special populations, uninsured and minority patients, and EHR use. Clinical performance for each measure is ranked from Quartile 1, highest 25% of reporting grantees, to Quartile 4, lowest 25% of reporting grantees.