UDS Summary Report - 2011 Grantee - Universal

GRANTEE			
PATIENTS			
Total Patients	46,731		
Number/Percent of Patients by Services Utilized			
Medical Services	38,394/82.16%		
Dental Services	11,309/24.20%		
Mental Health Services	0/0.00%		
Substance Abuse Services	0/0.00%		
Vision Services	0/0.00%		
Enabling Services	1,971/4.22%		
Number/Percent of Patients by Special Populations			
Homeless Patients	1,468/3.14%		
Farmworker Patients	6/0.01%		
School Based Patients	0/0.00%		
Veteran Patients	10/0.02%		
Number/Percent of Patients by Age			
Children (< 18 years old)	13,928/29.80%		
Adult (18 - 64)	30,927/66.18%		
Geriatric (age 65 and over)	1,876/4.01%		
Number/Percent of Patients by Insurance Status			
Uninsured	18,684/39.98%		
Children Uninsured (age 0-19 years)	1,905/12.26%		
Medicaid/CHIP ¹	20,127/43.07%		
Medicare	2,834/6.06%		
Other Third Party	5,086/10.88%		
Patients by Income Status (% Known)			
Patients at or below 200% of poverty	98.85%		
Patients at or below 100% of poverty (included in above)	88.41%		
Patients by Race & Ethnicity (% Known)			
% Racial and/or Ethnic Minority	92.53%		
% Hispanic/Latino Identity	12.21%		
% African American	88.06%		
% Asian	2.02%		
% American Indian/Alaska Native	0.11%		
% Native Hawaiian/Other Pacific Islander	0.65%		
% More than one race	0.32%		
% Best Served in another language	4.99%		

'-' - Data Cannot be Calculated

¹ Includes Medicaid, Medicaid CHIP, and Other Public Insurance CHIP

	GRANTEE
PATIENTS	
Percent of Medical Patients with Specific Diagnoses	
Hypertension ²	20.77%
Diabetes ³	12.30%
Asthma	1.96%
HIV	1.63%
Prenatal	
Number of Prenatal Patients	1,391
Number of Prenatal Patients who delivered	786

GRANTEE

ADJUSTED QUARTILE RANKING ⁵

QUALITY OF CARE/HEALTH OUTCOMES

% Having First Prenatal Visit in 1st Trimester58.66%4% Low and Very Low Birth Weight8.94%3Preventive Screening & Immunizations% of Women with Pap Tests64.29%2% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity Documented58.57%2% of Adults who Received Weight Screening and Follow up if Appropriate37.14%3% of Adults Assessed for Tobacco Use85.71%3% of Tobacco Users who Received Cessation Advice and/or Medication34.29%4% of Asthmatics Treated with Appropriate Pharmacological Intervention87.14%2	Prenatal			
Preventive Screening & Immunizations % of Women with Pap Tests 64.29% 2 % of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity Documented 58.57% 2 % of Adults who Received Weight Screening and Follow up if Appropriate 37.14% 3 % of Adults Assessed for Tobacco Use 85.71% 3 % of Tobacco Users who Received Cessation Advice and/or Medication 34.29% 4	% Having First Prenatal Visit in 1st Trimester	58.66%	4	
% of Women with Pap Tests64.29%2% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity Documented58.57%2% of Adults who Received Weight Screening and Follow up if Appropriate37.14%3% of Adults Assessed for Tobacco Use85.71%3% of Tobacco Users who Received Cessation Advice and/or Medication34.29%4	% Low and Very Low Birth Weight	8.94%	3	
% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity Documented58.57%2% of Adults who Received Weight Screening and Follow up if Appropriate37.14%3% of Adults Assessed for Tobacco Use85.71%3% of Tobacco Users who Received Cessation Advice and/or Medication34.29%4	Preventive Screening & Immunizations			
Documented58.57%2% of Adults who Received Weight Screening and Follow up if Appropriate37.14%3% of Adults Assessed for Tobacco Use85.71%3% of Tobacco Users who Received Cessation Advice and/or Medication34.29%4	% of Women with Pap Tests	64.29%	2	
% of Adults Assessed for Tobacco Use 85.71% 3 % of Tobacco Users who Received Cessation Advice and/or Medication 34.29% 4		58.57%	2	
% of Tobacco Users who Received Cessation Advice and/or Medication 34.29% 4	% of Adults who Received Weight Screening and Follow up if Appropriate	37.14%	3	
	% of Adults Assessed for Tobacco Use	85.71%	3	
% of Asthmatics Treated with Appropriate Pharmacological Intervention 87.14%	% of Tobacco Users who Received Cessation Advice and/or Medication	34.29%	4	
	% of Asthmatics Treated with Appropriate Pharmacological Intervention	87.14%	2	
% of Two year Olds Immunized ⁴ 12.86% 4	% of Two year Olds Immunized ⁴	12.86%	4	
Chronic Disease Management				
% Hypertensive Patients with Blood Pressure < 140/90 45.71%	% Hypertensive Patients with Blood Pressure < 140/90	45.71%	4	
% Diabetic Patients with HbA1c <= 9 75.71% 2	% Diabetic Patients with HbA1c <= 9	75.71%	2	

'-' - Data Cannot be Calculated

² Hypertensive adults as a percent of estimated adult medical patients of ages 18-85.

³ Diabetic adults as a percent of estimated adult medical patients of ages 18-75.

 $^{\rm 4}$ Measure was revised in 2011 and is not comparable to calendar year 2010 and prior.

⁵ Grantee adjusted quartile results from a statistical model adjusting for special populations, uninsured and minority patients, and EHR use. Clinical performance for each measure is ranked from Quartile 1, highest 25% of reporting grantees, to Quartile 4, lowest 25% of reporting grantees.