UDS Summary Report - 2011 Grantee - Universal

GRANTEE			
PATIENTS			
Total Patients	9,598		
Number/Percent of Patients by Services Utilized			
Medical Services	9,559/99.59%		
Dental Services	0/0.00%		
Mental Health Services	213/2.22%		
Substance Abuse Services	0/0.00%		
Vision Services	0/0.00%		
Enabling Services	135/1.41%		
Number/Percent of Patients by Special Populations			
Homeless Patients	-/-		
Farmworker Patients	-/-		
School Based Patients	-/-		
Veteran Patients	-/-		
Number/Percent of Patients by Age			
Children (< 18 years old)	2,587/26.95%		
Adult (18 - 64)	6,634/69.12%		
Geriatric (age 65 and over)	377/3.93%		
Number/Percent of Patients by Insurance Status			
Uninsured	3,753/39.10%		
Children Uninsured (age 0-19 years)	833/28.04%		
Medicaid/CHIP ¹	3,595/37.46%		
Medicare	536/5.58%		
Other Third Party	1,714/17.86%		
Patients by Income Status (% Known)			
Patients at or below 200% of poverty	94.40%		
Patients at or below 100% of poverty (included in above)	66.58%		
Patients by Race & Ethnicity (% Known)			
% Racial and/or Ethnic Minority	66.98%		
% Hispanic/Latino Identity	26.16%		
% African American	44.62%		
% Asian	2.89%		
% American Indian/Alaska Native	4.06%		
% Native Hawaiian/Other Pacific Islander	0.22%		
% More than one race	4.30%		
% Best Served in another language	18.18%		
'-' - Data Cannot be Calculated			

'-' - Data Cannot be Calculated

¹ Includes Medicaid, Medicaid CHIP, and Other Public Insurance CHIP

	GRANTEE
PATIENTS	
Percent of Medical Patients with Specific Diagnoses	
Hypertension ²	14.41%
Diabetes ³	7.75%
Asthma	2.69%
HIV	0.04%
Prenatal	
Number of Prenatal Patients	306
Number of Prenatal Patients who delivered	121

GRANTEE

ADJUSTED QUARTILE RANKING ⁵

QUALITY OF CARE/HEALTH OUTCOMES

% Having First Prenatal Visit in 1st Trimester(16)(16)(16)% Low and Very Low Birth Weight(16)(16)(16)(16)Preventive Screening & Immunizations% of Women with Pap Tests(16)(16)(16)% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity(16)(16)(16)% of Adults Aboescend Weight Screening and Follow up 1 Appropriate(16)(16)(16)(16)% of Adults Assessed for Tobacco Use(16)(16)(16)(16)(16)(16)% of Aboescend Weight Appropriate Pharmacological Intervention(16)	Prenatal			
Preventive Screening & Immunizations % of Women with Pap Tests 40.00% 4 % of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity 2.86% 2.86% % of Adults who Received Weight Screening and Follow up if Appropriate 41.43% 2 % of Adults Assessed for Tobacco Use 100% 1 % of Adults Assessed for Tobacco Use 21.43% 4 % of Asthmatics Treated with Appropriate Pharmacological Intervention 94.29% 1 % of Two year Olds Immunized ⁴ 2.86% 4 Khypertensive Patients with Blood Pressure < 140/90	% Having First Prenatal Visit in 1st Trimester	85.29%	1	
% of Women with Pap Tests40.00%40.00%4% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity Documented2.86%2.86%4% of Adults who Received Weight Screening and Follow up if Appropriate41.43%40.00%2% of Adults Assessed for Tobacco Use41.43%40.00%1% of Tobacco Users who Received Cessation Advice and/or Medication21.43%40.00%4% of Asthmatics Treated with Appropriate Pharmacological Intervention94.29%40.00%4% of Two year Olds Immunized 42.86%40.00%4Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90	% Low and Very Low Birth Weight	5.79%	2	
A constraint of the bill Percentile and Counseling on Nutrition and Physical Activity Documented2.86%2.86%% of Adults who Received Weight Screening and Follow up if Appropriate41.43%4% of Adults Assessed for Tobacco Use41.43%4% of Adults Assessed for Tobacco Use100%100%% of Tobacco Users who Received Cessation Advice and/or Medication61.41.43%4% of Asthmatics Treated with Appropriate Pharmacological Intervention94.29%1% of Two year Olds Immunized ⁴ 244Khypertensive Patients with Blood Pressure < 140/90	Preventive Screening & Immunizations			
Documented2.86%2.86%4% of Adults who Received Weight Screening and Follow up if Appropriate41.43%62% of Adults Assessed for Tobacco Use100%100%1% of Tobacco Users who Received Cessation Advice and/or Medication21.43%44% of Asthmatics Treated with Appropriate Pharmacological Intervention94.29%14% of Two year Olds Immunized 42.86%44Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90	% of Women with Pap Tests	40.00%	4	
% of Adults Assessed for Tobacco Use 100% 1 % of Tobacco Users who Received Cessation Advice and/or Medication 21.43% 4 % of Asthmatics Treated with Appropriate Pharmacological Intervention 94.29% 1 % of Two year Olds Immunized ⁴ 2.86% 4 Chronic Disease Management % Hypertensive Patients with Blood Pressure < 140/90		2.86%	4	
% of Tobacco Users who Received Cessation Advice and/or Medication 21.43% 4 % of Asthmatics Treated with Appropriate Pharmacological Intervention 94.29% 4 % of Two year Olds Immunized ⁴ 2.86% 4 Chronic Disease Management % Hypertensive Patients with Blood Pressure < 140/90	% of Adults who Received Weight Screening and Follow up if Appropriate	41.43%	2	
% of Asthmatics Treated with Appropriate Pharmacological Intervention 94.29% 94.29% 1 % of Two year Olds Immunized ⁴ 2.86% 4 Chronic Disease Management % Hypertensive Patients with Blood Pressure < 140/90	% of Adults Assessed for Tobacco Use	100%	1	
% of Two year Olds Immunized 4 2.86% 4 Chronic Disease Management 2.86% 1 % Hypertensive Patients with Blood Pressure < 140/90	% of Tobacco Users who Received Cessation Advice and/or Medication	21.43%	4	
Chronic Disease Management % Hypertensive Patients with Blood Pressure < 140/90	% of Asthmatics Treated with Appropriate Pharmacological Intervention	94.29%	1	
% Hypertensive Patients with Blood Pressure < 140/90 80.00%	% of Two year Olds Immunized ⁴	2.86%	4	
	Chronic Disease Management			
% Diabetic Patients with HbA1c <= 9 80.00% 1	% Hypertensive Patients with Blood Pressure < 140/90	80.00%	1	
	% Diabetic Patients with HbA1c <= 9	80.00%	1	

'-' - Data Cannot be Calculated

² Hypertensive adults as a percent of estimated adult medical patients of ages 18-85.

³ Diabetic adults as a percent of estimated adult medical patients of ages 18-75.

 $^{\rm 4}$ Measure was revised in 2011 and is not comparable to calendar year 2010 and prior.

⁵ Grantee adjusted quartile results from a statistical model adjusting for special populations, uninsured and minority patients, and EHR use. Clinical performance for each measure is ranked from Quartile 1, highest 25% of reporting grantees, to Quartile 4, lowest 25% of reporting grantees.