UDS Summary Report - 2011 Grantee - Universal

GRANTEE	
PATIENTS	
Total Patients	46,128
Number/Percent of Patients by Services Utilized	
Medical Services	36,785/79.75%
Dental Services	14,066/30.49%
Mental Health Services	1,641/3.56%
Substance Abuse Services	332/0.72%
Vision Services	0/0.00%
Enabling Services	465/1.01%
Number/Percent of Patients by Special Populations	
Homeless Patients	6,101/13.23%
Farmworker Patients	162/0.35%
School Based Patients	0/0.00%
Veteran Patients	83/0.18%
Number/Percent of Patients by Age	
Children (< 18 years old)	9,828/21.31%
Adult (18 - 64)	34,354/74.48%
Geriatric (age 65 and over)	1,946/4.22%
Number/Percent of Patients by Insurance Status	
Uninsured	17,436/37.80%
Children Uninsured (age 0-19 years)	1,217/10.86%
Medicaid/CHIP ¹	19,503/42.28%
Medicare	4,019/8.71%
Other Third Party	5,170/11.21%
Patients by Income Status (% Known)	
Patients at or below 200% of poverty	94.28%
Patients at or below 100% of poverty (included in above)	63.22%
Patients by Race & Ethnicity (% Known)	
% Racial and/or Ethnic Minority	12.87%
% Hispanic/Latino Identity	4.00%
% African American	3.45%
% Asian	3.02%
% American Indian/Alaska Native	1.18%
% Native Hawaiian/Other Pacific Islander	0.83%
% More than one race	1.08%
% Best Served in another language	4.29%
L. Date Connet he Coloridated	

'-' - Data Cannot be Calculated

¹ Includes Medicaid, Medicaid CHIP, and Other Public Insurance CHIP

	GRANTEE	
PATIENTS		
Percent of Medical Patients with Specific Diagnoses		
Hypertension ²	18.30%	
Diabetes ³	9.02%	
Asthma	2.59%	
HIV	0.73%	
Prenatal		
Number of Prenatal Patients	437	
Number of Prenatal Patients who delivered	230	

GRANTEE

ADJUSTED QUARTILE RANKING 5

QUALITY OF CARE/HEALTH OUTCOMES

% Having First Prenatal Visit in 1st Trimester6% Having First Prenatal Visit in 1st Trimester6 Preventive Screening & Immunizations % of Women with Pap Tests6% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity2% of Adults who Received Weight Screening and Follow up if Appropriate2% of Adults Assessed for Tobacco Use6% of Adults Assessed for Tobacco Use6% of Astmatics Treated with Appropriate Pharmacological Intervention6% of Astmatics Treated with Appropriate Pharmacological Intervention6% Having Tablescenting8% Hayertensive Patients with Blood Pressure < 140/906% Hayertensive Hablack < 10006% Diabetic Patients with HbA1c <= 96	Prenatal			
Preventive Screening & Immunizations % of Women with Pap Tests 32.86% 4 % of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity 24.29% 24.29% % of Adults who Received Weight Screening and Follow up if Appropriate 28.93% 26 % of Adults Assessed for Tobacco Use 93.97% 1 % of Adults Assessed for Tobacco Use 98.55% 4 % of Asthmatics Treated with Appropriate Pharmacological Intervention 46.98% 4 % of Two year Olds Immunized ⁴ 31.857% 3 Chronic Disease Management 51.83% 2	% Having First Prenatal Visit in 1st Trimester	84.21%	1	
% of Women with Pap Tests32.86%32.86%% of Children and Adolescents with BMI Percentile and Counseling on Nutrition and Physical Activity Documented24.29%24.29%% of Adults who Received Weight Screening and Follow up if Appropriate28.93%20.2% of Adults Assessed for Tobacco Use33.97%30.01% of Tobacco Users who Received Cessation Advice and/or Medication9.85%30.01% of Asthmatics Treated with Appropriate Pharmacological Intervention46.98%30.31% of Two year Olds Immunized 418.57%30.33Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90	% Low and Very Low Birth Weight	10.30%	4	
A constraint of the constraint o	Preventive Screening & Immunizations			
Documented24.29%24.29%2% of Adults who Received Weight Screening and Follow up if Appropriate28.93%2% of Adults Assessed for Tobacco Use93.97%1% of Tobacco Users who Received Cessation Advice and/or Medication9.85%4% of Asthmatics Treated with Appropriate Pharmacological Intervention46.98%4% of Two year Olds Immunized 418.57%3Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90	% of Women with Pap Tests	32.86%	4	
% of Adults Assessed for Tobacco Use93.97%93.97%1% of Tobacco Users who Received Cessation Advice and/or Medication9.85%9.85%4% of Asthmatics Treated with Appropriate Pharmacological Intervention46.98%4% of Two year Olds Immunized 418.57%3Chronic Disease Management% Hypertensive Patients with Blood Pressure < 140/90		24.29%	2	
% of Tobacco Users who Received Cessation Advice and/or Medication 9.85% 9.85% 4 % of Asthmatics Treated with Appropriate Pharmacological Intervention 46.98% 4 % of Two year Olds Immunized ⁴ 18.57% 3 Chronic Disease Management % Hypertensive Patients with Blood Pressure < 140/90	% of Adults who Received Weight Screening and Follow up if Appropriate	28.93%	2	
% of Asthmatics Treated with Appropriate Pharmacological Intervention 46.98% 4 % of Two year Olds Immunized ⁴ 18.57% 3 Chronic Disease Management % Hypertensive Patients with Blood Pressure < 140/90	% of Adults Assessed for Tobacco Use	93.97%	1	
% of Two year Olds Immunized 4 18.57% 3 Chronic Disease Management 2 % Hypertensive Patients with Blood Pressure < 140/90	% of Tobacco Users who Received Cessation Advice and/or Medication	9.85%	4	
Chronic Disease Management 61.83% 62 % Hypertensive Patients with Blood Pressure < 140/90	% of Asthmatics Treated with Appropriate Pharmacological Intervention	46.98%	4	
% Hypertensive Patients with Blood Pressure < 140/90 61.83% 2	% of Two year Olds Immunized ⁴	18.57%	3	
	Chronic Disease Management			
% Diabetic Patients with HbA1c <= 9 72.86% 3	% Hypertensive Patients with Blood Pressure < 140/90	61.83%	2	
	% Diabetic Patients with HbA1c <= 9	72.86%	3	

'-' - Data Cannot be Calculated

² Hypertensive adults as a percent of estimated adult medical patients of ages 18-85.

³ Diabetic adults as a percent of estimated adult medical patients of ages 18-75.

 $^{\rm 4}$ Measure was revised in 2011 and is not comparable to calendar year 2010 and prior.

⁵ Grantee adjusted quartile results from a statistical model adjusting for special populations, uninsured and minority patients, and EHR use. Clinical performance for each measure is ranked from Quartile 1, highest 25% of reporting grantees, to Quartile 4, lowest 25% of reporting grantees.