

Incidence

Supplemental Materials #2: Rates and Trends by Site and Sex for 1992-2000

Incidence SEER 12 Areas	Both Sexes	Male	Female	Both Sexes	Male	Female
All Races	Rate	Rate	Rate	APC	APC	APC
All Sites	477.7	574.2	414.8	-0.7 *	-1.7 *	0.2
Brain and Other Nervous System	6.4	7.7	5.4	-0.5	-0.6	-0.5
Breast			132.5			0.8 *
Cervix Uteri			10.0			-2.5 *
Colon and Rectum	54.5	65.0	46.8	-0.6	-0.9 *	-0.3
Corpus and Uterus, NOS			24.5			-0.1
Esophagus	4.5	7.5	2.1	0.3	0.4	-0.4
Hodgkin Lymphoma	2.7	3.0	2.4	-0.4	-0.1	-0.6
Kidney and Renal Pelvis	10.8	15.1	7.5	1.3 *	1.1 *	1.3 *
Larynx	4.1	7.4	1.6	-2.6 *	-3.2 *	-1.0
Leukemia	12.4	16.2	9.5	-1.3 *	-1.5 *	-1.0
Liver and Intrahepatic Bile Duct	5.5	8.3	3.2	3.9 *	3.7 *	3.7 *
Lung and Bronchus	64.0	84.5	49.4	-1.2 *	-2.3 *	0.0
Melanoma of the Skin	15.7	19.8	12.8	2.5 *	2.8 *	2.3 *
Myeloma	5.6	7.0	4.6	-0.7	-0.8	-0.9
Non-Hodgkin's Lymphomas	19.1	23.7	15.3	0.0	-0.5	0.9 *
Oral Cavity and Pharynx	11.3	16.9	6.7	-2.0 *	-2.3 *	-1.5 *
Ovary			17.0			-0.8 *
Pancreas	11.1	12.7	9.9	-0.4	-0.4	-0.5
Prostate		180.6			-3.1 *	
Stomach	9.3	13.5	6.3	-1.3 *	-1.9 *	-0.5
Testis		5.0			1.3 *	
Thyroid	6.4	3.5	9.2	3.4 *	1.8 *	4.0 *
Urinary Bladder	20.4	36.1	9.2	-0.1	-0.1	-0.5

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard population by 5-year age groups.

~ Statistic could not be calculated.

^ Statistic not displayed due to less than 25 cases.

* The APC is significantly different from zero (p<0.05).