


Table 2.18  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	178.65	0.11		
High Five States				
Kentucky	211.31 <sup>b</sup>	0.97	(01)	18.28 <sup>c</sup>
West Virginia	206.56 <sup>b</sup>	1.36	(02)	15.62 <sup>c</sup>
Mississippi	204.74 <sup>b</sup>	1.18	(03)	14.60 <sup>c</sup>
Louisiana	204.05 <sup>b</sup>	0.97	(04)	14.21 <sup>c</sup>
Tennessee	200.56 <sup>b</sup>	0.79	(05)	12.26 <sup>c</sup>
Low Five States				
New Mexico	158.17 <sup>b</sup>	1.26	(47)	-11.46 <sup>c</sup>
Colorado	154.59 <sup>b</sup>	0.86	(48)	-13.47 <sup>c</sup>
Arizona	152.84 <sup>b</sup>	0.68	(49)	-14.45 <sup>c</sup>
Hawaii	147.81 <sup>b</sup>	1.43	(50)	-17.26 <sup>c</sup>
Utah	128.35 <sup>b</sup>	1.15	(51)	-28.16 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	198.24 <sup>b</sup>	0.89	(07)	10.97 <sup>c</sup>	Montana	172.53	1.78	(37)	-3.43
Alaska	181.35	3.05	(23)	1.51	Nebraska	174.00	1.35	(35)	-2.60
Arizona	152.84 <sup>b</sup>	0.68	(49)	-14.45 <sup>c</sup>	Nevada	183.03	1.26	(22)	2.45
Arkansas	199.69 <sup>b</sup>	1.12	(06)	11.78 <sup>c</sup>	New Hampshire	179.46	1.60	(27)	0.45
California	163.07 <sup>b</sup>	0.31	(46)	-8.72 <sup>c</sup>	New Jersey	179.05	0.62	(29)	0.22
Colorado	154.59 <sup>b</sup>	0.86	(48)	-13.47 <sup>c</sup>	New Mexico	158.17 <sup>b</sup>	1.26	(47)	-11.46 <sup>c</sup>
Connecticut	173.41 <sup>b</sup>	0.94	(36)	-2.94	New York	166.74 <sup>b</sup>	0.40	(44)	-6.67 <sup>c</sup>
Delaware	190.45 <sup>b</sup>	2.00	(14)	6.60 <sup>c</sup>	North Carolina	186.28 <sup>b</sup>	0.64	(19)	4.27
D.C.	196.47 <sup>b</sup>	2.61	(08)	9.98 <sup>c</sup>	North Dakota	171.11	2.15	(39)	-4.22
Florida	169.96 <sup>b</sup>	0.38	(40)	-4.86	Ohio	194.90 <sup>b</sup>	0.55	(10)	9.09 <sup>c</sup>
Georgia	179.23	0.67	(28)	0.32	Oklahoma	194.77 <sup>b</sup>	1.01	(11)	9.02 <sup>c</sup>
Hawaii	147.81 <sup>b</sup>	1.43	(50)	-17.26 <sup>c</sup>	Oregon	179.83	0.95	(26)	0.66
Idaho	165.84 <sup>b</sup>	1.52	(45)	-7.17 <sup>c</sup>	Pennsylvania	187.50 <sup>b</sup>	0.50	(17)	4.95
Illinois	186.65 <sup>b</sup>	0.54	(18)	4.48	Rhode Island	180.37	1.73	(24)	0.96
Indiana	195.42 <sup>b</sup>	0.77	(09)	9.39 <sup>c</sup>	South Carolina	187.94 <sup>b</sup>	0.90	(15)	5.20 <sup>c</sup>
Iowa	178.34	1.01	(32)	-0.18	South Dakota	168.20 <sup>b</sup>	1.92	(43)	-5.85 <sup>c</sup>
Kansas	178.80	1.10	(30)	0.08	Tennessee	200.56 <sup>b</sup>	0.79	(05)	12.26 <sup>c</sup>
Kentucky	211.31 <sup>b</sup>	0.97	(01)	18.28 <sup>c</sup>	Texas	171.30 <sup>b</sup>	0.41	(38)	-4.12
Louisiana	204.05 <sup>b</sup>	0.97	(04)	14.21 <sup>c</sup>	Utah	128.35 <sup>b</sup>	1.15	(51)	-28.16 <sup>c</sup>
Maine	192.79 <sup>b</sup>	1.55	(12)	7.91 <sup>c</sup>	Vermont	176.21	2.24	(34)	-1.37
Maryland	184.15 <sup>b</sup>	0.82	(20)	3.08	Virginia	183.16 <sup>b</sup>	0.70	(21)	2.52
Massachusetts	180.05	0.71	(25)	0.78	Washington	176.74	0.75	(33)	-1.07
Michigan	187.53 <sup>b</sup>	0.59	(16)	4.97	West Virginia	206.56 <sup>b</sup>	1.36	(02)	15.62 <sup>c</sup>
Minnesota	169.94 <sup>b</sup>	0.80	(41)	-4.87	Wisconsin	178.60	0.77	(31)	-0.03
Mississippi	204.74 <sup>b</sup>	1.18	(03)	14.60 <sup>c</sup>	Wyoming	169.82	2.54	(42)	-4.94
Missouri	191.40 <sup>b</sup>	0.77	(13)	7.13 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.19  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	219.38	0.18		
High Five States				
Mississippi	274.20 <sup>b</sup>	2.16	(01)	24.99 <sup>c</sup>
Kentucky	267.18 <sup>b</sup>	1.72	(02)	21.79 <sup>c</sup>
Louisiana	260.82 <sup>b</sup>	1.72	(03)	18.89 <sup>c</sup>
Alabama	259.03 <sup>b</sup>	1.60	(04)	18.07 <sup>c</sup>
Tennessee	257.92 <sup>b</sup>	1.41	(05)	17.56 <sup>c</sup>
Low Five States				
New Mexico	190.10 <sup>b</sup>	2.08	(47)	-13.35 <sup>c</sup>
Colorado	185.02 <sup>b</sup>	1.47	(48)	-15.66 <sup>c</sup>
Hawaii	184.64 <sup>b</sup>	2.39	(49)	-15.84 <sup>c</sup>
Arizona	182.08 <sup>b</sup>	1.12	(50)	-17.00 <sup>c</sup>
Utah	154.13 <sup>b</sup>	1.91	(51)	-29.74 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	259.03 <sup>b</sup>	1.60	(04)	18.07 <sup>c</sup>	Montana	203.37 <sup>b</sup>	2.91	(42)	-7.30 <sup>c</sup>
Alaska	209.46	5.03	(38)	-4.52	Nebraska	215.19	2.30	(29)	-1.91
Arizona	182.08 <sup>b</sup>	1.12	(50)	-17.00 <sup>c</sup>	Nevada	213.31	2.04	(32)	-2.77
Arkansas	253.69 <sup>b</sup>	1.94	(08)	15.64 <sup>c</sup>	New Hampshire	218.24	2.75	(28)	-0.52
California	194.87 <sup>b</sup>	0.53	(46)	-11.17 <sup>c</sup>	New Jersey	213.85 <sup>b</sup>	1.06	(31)	-2.52
Colorado	185.02 <sup>b</sup>	1.47	(48)	-15.66 <sup>c</sup>	New Mexico	190.10 <sup>b</sup>	2.08	(47)	-13.35 <sup>c</sup>
Connecticut	212.00 <sup>b</sup>	1.63	(34)	-3.36	New York	201.34 <sup>b</sup>	0.69	(43)	-8.22 <sup>c</sup>
Delaware	229.56	3.38	(18)	4.64	North Carolina	236.93 <sup>b</sup>	1.14	(15)	8.00 <sup>c</sup>
D.C.	256.34 <sup>b</sup>	4.84	(06)	16.85 <sup>c</sup>	North Dakota	210.23	3.59	(36)	-4.17
Florida	205.95 <sup>b</sup>	0.62	(41)	-6.12 <sup>c</sup>	Ohio	243.35 <sup>b</sup>	0.97	(10)	10.93 <sup>c</sup>
Georgia	230.77 <sup>b</sup>	1.24	(17)	5.19 <sup>c</sup>	Oklahoma	242.96 <sup>b</sup>	1.73	(11)	10.75 <sup>c</sup>
Hawaii	184.64 <sup>b</sup>	2.39	(49)	-15.84 <sup>c</sup>	Oregon	214.35	1.57	(30)	-2.29
Idaho	195.90 <sup>b</sup>	2.49	(45)	-10.70 <sup>c</sup>	Pennsylvania	232.37 <sup>b</sup>	0.86	(16)	5.92 <sup>c</sup>
Illinois	229.38 <sup>b</sup>	0.94	(19)	4.56	Rhode Island	228.82	3.05	(20)	4.30
Indiana	244.88 <sup>b</sup>	1.36	(09)	11.62 <sup>c</sup>	South Carolina	241.26 <sup>b</sup>	1.59	(12)	9.97 <sup>c</sup>
Iowa	220.08	1.72	(26)	0.32	South Dakota	206.04 <sup>b</sup>	3.21	(40)	-6.08 <sup>c</sup>
Kansas	221.55	1.89	(25)	0.99	Tennessee	257.92 <sup>b</sup>	1.41	(05)	17.56 <sup>c</sup>
Kentucky	267.18 <sup>b</sup>	1.72	(02)	21.79 <sup>c</sup>	Texas	212.50 <sup>b</sup>	0.72	(33)	-3.14
Louisiana	260.82 <sup>b</sup>	1.72	(03)	18.89 <sup>c</sup>	Utah	154.13 <sup>b</sup>	1.91	(51)	-29.74 <sup>c</sup>
Maine	240.04 <sup>b</sup>	2.69	(13)	9.42 <sup>c</sup>	Vermont	211.92	3.82	(35)	-3.40
Maryland	226.47 <sup>b</sup>	1.44	(23)	3.23	Virginia	228.46 <sup>b</sup>	1.24	(21)	4.14
Massachusetts	222.63	1.24	(24)	1.48	Washington	209.64 <sup>b</sup>	1.25	(37)	-4.44
Michigan	228.13 <sup>b</sup>	1.02	(22)	3.99	West Virginia	254.76 <sup>b</sup>	2.33	(07)	16.13 <sup>c</sup>
Minnesota	206.75 <sup>b</sup>	1.36	(39)	-5.76 <sup>c</sup>	Wisconsin	218.85	1.31	(27)	-0.24
Mississippi	274.20 <sup>b</sup>	2.16	(01)	24.99 <sup>c</sup>	Wyoming	199.45 <sup>b</sup>	4.17	(44)	-9.08 <sup>c</sup>
Missouri	237.57 <sup>b</sup>	1.33	(14)	8.29 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.20  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	151.06	0.13		
High Five States				
Kentucky	173.63 <sup>b</sup>	1.17	(01)	14.94 <sup>c</sup>
West Virginia	173.15 <sup>b</sup>	1.68	(02)	14.62 <sup>c</sup>
Louisiana	165.83 <sup>b</sup>	1.16	(03)	9.78 <sup>c</sup>
Ohio	163.39 <sup>b</sup>	0.68	(04)	8.16 <sup>c</sup>
Indiana	163.17 <sup>b</sup>	0.94	(05)	8.02 <sup>c</sup>
Low Five States				
Colorado	134.38 <sup>b</sup>	1.07	(47)	-11.04 <sup>c</sup>
New Mexico	134.32 <sup>b</sup>	1.56	(48)	-11.09 <sup>c</sup>
Arizona	129.98 <sup>b</sup>	0.86	(49)	-13.96 <sup>c</sup>
Hawaii	119.64 <sup>b</sup>	1.75	(50)	-20.80 <sup>c</sup>
Utah	109.61 <sup>b</sup>	1.43	(51)	-27.44 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	157.38 <sup>b</sup>	1.05	(20)	4.18	Montana	150.54	2.27	(33)	-0.34
Alaska	159.58	3.85	(15)	5.64 <sup>c</sup>	Nebraska	145.72	1.66	(39)	-3.53
Arizona	129.98 <sup>b</sup>	0.86	(49)	-13.96 <sup>c</sup>	Nevada	158.37 <sup>b</sup>	1.60	(18)	4.83
Arkansas	161.16 <sup>b</sup>	1.36	(10)	6.68 <sup>c</sup>	New Hampshire	154.69	1.99	(23)	2.40
California	141.71 <sup>b</sup>	0.39	(45)	-6.19 <sup>c</sup>	New Jersey	157.68 <sup>b</sup>	0.77	(19)	4.38
Colorado	134.38 <sup>b</sup>	1.07	(47)	-11.04 <sup>c</sup>	New Mexico	134.32 <sup>b</sup>	1.56	(48)	-11.09 <sup>c</sup>
Connecticut	149.61	1.16	(35)	-0.96	New York	145.18 <sup>b</sup>	0.49	(40)	-3.89
Delaware	162.79 <sup>b</sup>	2.48	(06)	7.76 <sup>c</sup>	North Carolina	152.72	0.76	(28)	1.10
D.C.	160.42	3.08	(12)	6.20 <sup>c</sup>	North Dakota	144.15	2.70	(41)	-4.58
Florida	141.93 <sup>b</sup>	0.48	(44)	-6.04 <sup>c</sup>	Ohio	163.39 <sup>b</sup>	0.68	(04)	8.16 <sup>c</sup>
Georgia	146.81 <sup>b</sup>	0.79	(37)	-2.81	Oklahoma	161.17 <sup>b</sup>	1.23	(09)	6.69 <sup>c</sup>
Hawaii	119.64 <sup>b</sup>	1.75	(50)	-20.80 <sup>c</sup>	Oregon	155.48 <sup>b</sup>	1.19	(22)	2.92
Idaho	143.54 <sup>b</sup>	1.93	(42)	-4.98	Pennsylvania	158.49 <sup>b</sup>	0.61	(17)	4.91
Illinois	160.11 <sup>b</sup>	0.66	(14)	5.99 <sup>c</sup>	Rhode Island	151.29	2.10	(30)	0.15
Indiana	163.17 <sup>b</sup>	0.94	(05)	8.02 <sup>c</sup>	South Carolina	151.03	1.07	(31)	-0.02
Iowa	150.98	1.26	(32)	-0.06	South Dakota	141.51 <sup>b</sup>	2.41	(46)	-6.33 <sup>c</sup>
Kansas	149.92	1.36	(34)	-0.76	Tennessee	161.96 <sup>b</sup>	0.94	(07)	7.22 <sup>c</sup>
Kentucky	173.63 <sup>b</sup>	1.17	(01)	14.94 <sup>c</sup>	Texas	142.76 <sup>b</sup>	0.50	(43)	-5.50 <sup>c</sup>
Louisiana	165.83 <sup>b</sup>	1.16	(03)	9.78 <sup>c</sup>	Utah	109.61 <sup>b</sup>	1.43	(51)	-27.44 <sup>c</sup>
Maine	161.61 <sup>b</sup>	1.91	(08)	6.98 <sup>c</sup>	Vermont	152.83	2.80	(27)	1.17
Maryland	157.30 <sup>b</sup>	0.99	(21)	4.13	Virginia	153.87	0.85	(26)	1.86
Massachusetts	153.99	0.87	(24)	1.94	Washington	153.91	0.94	(25)	1.88
Michigan	160.91 <sup>b</sup>	0.74	(11)	6.52 <sup>c</sup>	West Virginia	173.15 <sup>b</sup>	1.68	(02)	14.62 <sup>c</sup>
Minnesota	146.04 <sup>b</sup>	1.00	(38)	-3.32	Wisconsin	152.00	0.95	(29)	0.62
Mississippi	158.85 <sup>b</sup>	1.37	(16)	5.15 <sup>c</sup>	Wyoming	148.32	3.23	(36)	-1.82
Missouri	160.37 <sup>b</sup>	0.95	(13)	6.16 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.18  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.28	0.02		
High Five States				
Iowa	5.53 <sup>b</sup>	0.18	(01)	29.17 <sup>c</sup>
Oregon	5.19 <sup>b</sup>	0.16	(02)	21.37 <sup>c</sup>
Idaho	5.14	0.27	(03)	20.13 <sup>c</sup>
Nebraska	5.12 <sup>b</sup>	0.24	(04)	19.77 <sup>c</sup>
Washington	5.12 <sup>b</sup>	0.13	(05)	19.74 <sup>c</sup>
Low Five States				
New York	3.70 <sup>b</sup>	0.06	(47)	-13.53
New Jersey	3.69 <sup>b</sup>	0.09	(48)	-13.83
New Mexico	3.66	0.19	(49)	-14.47
D.C.	3.47	0.35	(50)	-18.91 <sup>c</sup>
Hawaii	2.39 <sup>b</sup>	0.19	(51)	-44.16 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.58	0.14	(19)	6.98	Montana	4.53	0.29	(23)	5.76
Alaska	4.20	0.43	(36)	-1.77	Nebraska	5.12 <sup>b</sup>	0.24	(04)	19.77 <sup>c</sup>
Arizona	4.11	0.11	(38)	-3.84	Nevada	4.10	0.18	(39)	-4.10
Arkansas	4.95 <sup>b</sup>	0.18	(10)	15.62 <sup>c</sup>	New Hampshire	4.76	0.26	(14)	11.24
California	4.31	0.05	(32)	0.71	New Jersey	3.69 <sup>b</sup>	0.09	(48)	-13.83
Colorado	4.40	0.14	(29)	2.72	New Mexico	3.66	0.19	(49)	-14.47
Connecticut	4.22	0.15	(35)	-1.47	New York	3.70 <sup>b</sup>	0.06	(47)	-13.53
Delaware	4.54	0.31	(21)	6.15	North Carolina	4.05	0.09	(41)	-5.36
D.C.	3.47	0.35	(50)	-18.91 <sup>c</sup>	North Dakota	5.01	0.39	(08)	17.20 <sup>c</sup>
Florida	3.95 <sup>b</sup>	0.06	(42)	-7.79	Ohio	4.44	0.08	(27)	3.66
Georgia	3.87 <sup>b</sup>	0.10	(45)	-9.58	Oklahoma	4.68	0.16	(15)	9.36
Hawaii	2.39 <sup>b</sup>	0.19	(51)	-44.16 <sup>c</sup>	Oregon	5.19 <sup>b</sup>	0.16	(02)	21.37 <sup>c</sup>
Idaho	5.14	0.27	(03)	20.13 <sup>c</sup>	Pennsylvania	4.10	0.08	(40)	-4.23
Illinois	3.89 <sup>b</sup>	0.08	(43)	-9.11	Rhode Island	4.24	0.27	(34)	-0.96
Indiana	4.62	0.12	(17)	7.89	South Carolina	4.33	0.14	(31)	1.27
Iowa	5.53 <sup>b</sup>	0.18	(01)	29.17 <sup>c</sup>	South Dakota	4.93	0.35	(11)	15.29 <sup>c</sup>
Kansas	5.07 <sup>b</sup>	0.19	(07)	18.39 <sup>c</sup>	Tennessee	4.99 <sup>b</sup>	0.12	(09)	16.67 <sup>c</sup>
Kentucky	4.49	0.14	(24)	4.88	Texas	4.18	0.06	(37)	-2.26
Louisiana	4.53	0.14	(22)	5.85	Utah	4.46	0.21	(26)	4.32
Maine	5.12	0.26	(06)	19.74 <sup>c</sup>	Vermont	4.47	0.36	(25)	4.54
Maryland	3.78 <sup>b</sup>	0.12	(46)	-11.57	Virginia	3.87 <sup>b</sup>	0.10	(44)	-9.48
Massachusetts	4.24	0.11	(33)	-0.80	Washington	5.12 <sup>b</sup>	0.13	(05)	19.74 <sup>c</sup>
Michigan	4.86 <sup>b</sup>	0.10	(13)	13.61	West Virginia	4.57	0.21	(20)	6.91
Minnesota	4.33	0.13	(30)	1.28	Wisconsin	4.90 <sup>b</sup>	0.13	(12)	14.44
Mississippi	4.67	0.18	(16)	9.04	Wyoming	4.40	0.41	(28)	2.86
Missouri	4.58	0.12	(18)	6.99					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.19  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.23	0.03		
High Five States				
Iowa	6.95 <sup>b</sup>	0.31	(01)	32.78 <sup>c</sup>
Oregon	6.56 <sup>b</sup>	0.27	(02)	25.35 <sup>c</sup>
Washington	6.48 <sup>b</sup>	0.21	(03)	23.81 <sup>c</sup>
Maine	6.47	0.43	(04)	23.66 <sup>c</sup>
Kansas	6.39 <sup>b</sup>	0.31	(05)	22.06 <sup>c</sup>
Low Five States				
New Jersey	4.65 <sup>b</sup>	0.15	(47)	-11.08
Maryland	4.57	0.19	(48)	-12.60
New Mexico	4.48	0.31	(49)	-14.38
D.C.	4.44	0.60	(50)	-15.21 <sup>c</sup>
Hawaii	2.92 <sup>b</sup>	0.30	(51)	-44.21 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.55	0.23	(18)	6.04	Montana	5.53	0.46	(19)	5.67
Alaska	4.96	0.71	(36)	-5.27	Nebraska	6.01	0.38	(09)	14.89
Arizona	4.87	0.18	(42)	-6.98	Nevada	5.21	0.30	(32)	-0.38
Arkansas	5.90	0.29	(11)	12.79	New Hampshire	6.11	0.44	(06)	16.75 <sup>c</sup>
California	5.32	0.08	(29)	1.70	New Jersey	4.65 <sup>b</sup>	0.15	(47)	-11.08
Colorado	5.45	0.23	(23)	4.10	New Mexico	4.48	0.31	(49)	-14.38
Connecticut	5.07	0.25	(34)	-3.06	New York	4.74 <sup>b</sup>	0.10	(45)	-9.35
Delaware	5.75	0.53	(12)	9.84	North Carolina	5.01	0.16	(35)	-4.21
D.C.	4.44	0.60	(50)	-15.21 <sup>c</sup>	North Dakota	5.63	0.60	(15)	7.67
Florida	4.81 <sup>b</sup>	0.10	(43)	-8.07	Ohio	5.36	0.14	(27)	2.35
Georgia	4.67	0.16	(46)	-10.72	Oklahoma	5.60	0.26	(17)	7.06
Hawaii	2.92 <sup>b</sup>	0.30	(51)	-44.21 <sup>c</sup>	Oregon	6.56 <sup>b</sup>	0.27	(02)	25.35 <sup>c</sup>
Idaho	5.63	0.40	(14)	7.68	Pennsylvania	4.95	0.12	(37)	-5.46
Illinois	4.74 <sup>b</sup>	0.13	(44)	-9.34	Rhode Island	5.24	0.45	(31)	0.07
Indiana	5.46	0.19	(22)	4.31	South Carolina	5.42	0.23	(25)	3.52
Iowa	6.95 <sup>b</sup>	0.31	(01)	32.78 <sup>c</sup>	South Dakota	5.66	0.54	(13)	8.17
Kansas	6.39 <sup>b</sup>	0.31	(05)	22.06 <sup>c</sup>	Tennessee	6.03 <sup>b</sup>	0.21	(08)	15.31 <sup>c</sup>
Kentucky	5.46	0.23	(21)	4.42	Texas	4.89	0.10	(40)	-6.46
Louisiana	5.63	0.24	(16)	7.66	Utah	5.32	0.33	(28)	1.76
Maine	6.47	0.43	(04)	23.66 <sup>c</sup>	Vermont	4.93	0.56	(39)	-5.73
Maryland	4.57	0.19	(48)	-12.60	Virginia	4.87	0.17	(41)	-6.89
Massachusetts	5.38	0.19	(26)	2.79	Washington	6.48 <sup>b</sup>	0.21	(03)	23.81 <sup>c</sup>
Michigan	5.99 <sup>b</sup>	0.16	(10)	14.42	West Virginia	5.31	0.33	(30)	1.42
Minnesota	5.18	0.21	(33)	-1.01	Wisconsin	6.06 <sup>b</sup>	0.21	(07)	15.78 <sup>c</sup>
Mississippi	5.49	0.29	(20)	5.03	Wyoming	4.94	0.62	(38)	-5.66
Missouri	5.45	0.20	(24)	4.09					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.20  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.48	0.02		
High Five States				
Idaho	4.62	0.35	(01)	32.68 <sup>c</sup>
North Dakota	4.42	0.49	(02)	26.87 <sup>c</sup>
Iowa	4.33 <sup>b</sup>	0.23	(03)	24.32 <sup>c</sup>
South Dakota	4.32	0.45	(04)	23.95 <sup>c</sup>
Nebraska	4.30	0.30	(05)	23.52 <sup>c</sup>
Low Five States				
New Mexico	2.94	0.24	(47)	-15.48 <sup>c</sup>
New Jersey	2.90 <sup>b</sup>	0.11	(48)	-16.69 <sup>c</sup>
New York	2.87 <sup>b</sup>	0.07	(49)	-17.51 <sup>c</sup>
D.C.	2.67	0.41	(50)	-23.47 <sup>c</sup>
Hawaii	1.95 <sup>b</sup>	0.23	(51)	-43.93 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.76	0.17	(21)	8.05	Montana	3.63	0.37	(25)	4.09
Alaska	3.56	0.56	(31)	2.15	Nebraska	4.30	0.30	(05)	23.52 <sup>c</sup>
Arizona	3.46	0.14	(35)	-0.66	Nevada	3.09	0.22	(45)	-11.44
Arkansas	4.15	0.22	(06)	19.21 <sup>c</sup>	New Hampshire	3.62	0.32	(26)	4.01
California	3.45	0.06	(36)	-0.83	New Jersey	2.90 <sup>b</sup>	0.11	(48)	-16.69 <sup>c</sup>
Colorado	3.51	0.17	(33)	0.75	New Mexico	2.94	0.24	(47)	-15.48 <sup>c</sup>
Connecticut	3.52	0.19	(32)	0.97	New York	2.87 <sup>b</sup>	0.07	(49)	-17.51 <sup>c</sup>
Delaware	3.60	0.38	(28)	3.45	North Carolina	3.28	0.11	(40)	-5.83
D.C.	2.67	0.41	(50)	-23.47 <sup>c</sup>	North Dakota	4.42	0.49	(02)	26.87 <sup>c</sup>
Florida	3.19 <sup>b</sup>	0.07	(43)	-8.38	Ohio	3.66	0.10	(22)	5.17
Georgia	3.22	0.12	(41)	-7.53	Oklahoma	3.84	0.19	(20)	10.36
Hawaii	1.95 <sup>b</sup>	0.23	(51)	-43.93 <sup>c</sup>	Oregon	4.05	0.20	(08)	16.16 <sup>c</sup>
Idaho	4.62	0.35	(01)	32.68 <sup>c</sup>	Pennsylvania	3.41	0.09	(38)	-2.18
Illinois	3.19	0.10	(42)	-8.32	Rhode Island	3.43	0.33	(37)	-1.60
Indiana	3.93	0.15	(16)	12.67	South Carolina	3.47	0.16	(34)	-0.28
Iowa	4.33 <sup>b</sup>	0.23	(03)	24.32 <sup>c</sup>	South Dakota	4.32	0.45	(04)	23.95 <sup>c</sup>
Kansas	3.93	0.23	(15)	12.77	Tennessee	4.14 <sup>b</sup>	0.15	(07)	18.86 <sup>c</sup>
Kentucky	3.66	0.17	(23)	5.09	Texas	3.60	0.08	(30)	3.31
Louisiana	3.61	0.17	(27)	3.54	Utah	3.65	0.26	(24)	4.84
Maine	4.00	0.31	(09)	14.69	Vermont	3.99	0.47	(10)	14.63
Maryland	3.15	0.14	(44)	-9.56	Virginia	3.07	0.12	(46)	-11.93
Massachusetts	3.35	0.13	(39)	-3.81	Washington	3.94	0.15	(14)	12.99
Michigan	3.95 <sup>b</sup>	0.12	(13)	13.27	West Virginia	3.99	0.27	(12)	14.42
Minnesota	3.60	0.16	(29)	3.41	Wisconsin	3.91	0.16	(18)	12.13
Mississippi	3.99	0.22	(11)	14.48	Wyoming	3.91	0.53	(17)	12.15
Missouri	3.86	0.15	(19)	10.93					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 4.24  
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	23.02	0.05		
High Five States				
D.C.	27.96 <sup>b</sup>	1.29	(01)	21.46 <sup>c</sup>
Louisiana	26.32 <sup>b</sup>	0.46	(02)	14.35 <sup>c</sup>
New Jersey	26.06 <sup>b</sup>	0.31	(03)	13.20 <sup>c</sup>
Ohio	25.22 <sup>b</sup>	0.27	(04)	9.57
Mississippi	24.93	0.55	(05)	8.28
Low Five States				
Vermont	20.70	1.04	(47)	-10.09 <sup>c</sup>
Arizona	20.48 <sup>b</sup>	0.34	(48)	-11.02 <sup>c</sup>
Montana	20.48	0.84	(49)	-11.05 <sup>c</sup>
Colorado	19.92 <sup>b</sup>	0.41	(50)	-13.45 <sup>c</sup>
Hawaii	17.78 <sup>b</sup>	0.69	(51)	-22.79 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	23.98	0.41	(12)	4.18	Montana	20.48	0.84	(49)	-11.05 <sup>c</sup>
Alaska	23.51	1.41	(20)	2.11	Nebraska	21.20	0.64	(44)	-7.91
Arizona	20.48 <sup>b</sup>	0.34	(48)	-11.02 <sup>c</sup>	Nevada	23.32	0.61	(22)	1.31
Arkansas	23.62	0.53	(17)	2.59	New Hampshire	21.42	0.74	(40)	-6.95
California	22.32 <sup>b</sup>	0.15	(28)	-3.04	New Jersey	26.06 <sup>b</sup>	0.31	(03)	13.20 <sup>c</sup>
Colorado	19.92 <sup>b</sup>	0.41	(50)	-13.45 <sup>c</sup>	New Mexico	21.14	0.62	(45)	-8.16
Connecticut	22.54	0.45	(26)	-2.08	New York	22.53	0.20	(27)	-2.14
Delaware	23.05	0.94	(23)	0.11	North Carolina	23.54	0.30	(19)	2.27
D.C.	27.96 <sup>b</sup>	1.29	(01)	21.46 <sup>c</sup>	North Dakota	21.95	1.07	(30)	-4.65
Florida	21.51 <sup>b</sup>	0.19	(37)	-6.55	Ohio	25.22 <sup>b</sup>	0.27	(04)	9.57
Georgia	22.98	0.31	(24)	-0.17	Oklahoma	23.82	0.48	(16)	3.47
Hawaii	17.78 <sup>b</sup>	0.69	(51)	-22.79 <sup>c</sup>	Oregon	21.48	0.44	(38)	-6.72
Idaho	21.27	0.74	(43)	-7.61	Pennsylvania	24.09 <sup>b</sup>	0.24	(10)	4.64
Illinois	24.17 <sup>b</sup>	0.26	(09)	4.97	Rhode Island	21.93	0.81	(31)	-4.76
Indiana	23.87	0.36	(15)	3.68	South Carolina	23.97	0.43	(14)	4.11
Iowa	21.83	0.48	(34)	-5.17	South Dakota	20.87	0.93	(46)	-9.33
Kansas	22.87	0.53	(25)	-0.64	Tennessee	23.98	0.36	(13)	4.17
Kentucky	23.36	0.43	(21)	1.47	Texas	22.20 <sup>b</sup>	0.20	(29)	-3.56
Louisiana	26.32 <sup>b</sup>	0.46	(02)	14.35 <sup>c</sup>	Utah	21.54	0.63	(36)	-6.41
Maine	21.45	0.70	(39)	-6.84	Vermont	20.70	1.04	(47)	-10.09 <sup>c</sup>
Maryland	24.85 <sup>b</sup>	0.39	(07)	7.96	Virginia	24.79 <sup>b</sup>	0.34	(08)	7.69
Massachusetts	21.87	0.33	(33)	-4.99	Washington	21.90	0.35	(32)	-4.89
Michigan	24.02 <sup>b</sup>	0.29	(11)	4.32	West Virginia	23.59	0.63	(18)	2.48
Minnesota	21.34 <sup>b</sup>	0.38	(42)	-7.32	Wisconsin	21.60 <sup>b</sup>	0.36	(35)	-6.16
Mississippi	24.93	0.55	(05)	8.28	Wyoming	21.39	1.23	(41)	-7.07
Missouri	24.92 <sup>b</sup>	0.38	(06)	8.24					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 5.14  
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.39	0.02		
High Five States				
Mississippi	3.88 <sup>b</sup>	0.22	(01)	62.23 <sup>c</sup>
Louisiana	3.35 <sup>b</sup>	0.17	(02)	40.17 <sup>c</sup>
Arkansas	3.32 <sup>b</sup>	0.21	(03)	39.10 <sup>c</sup>
Alabama	3.04 <sup>b</sup>	0.15	(04)	27.32 <sup>c</sup>
D.C.	2.94	0.43	(05)	23.07 <sup>c</sup>
Low Five States				
Minnesota	1.49 <sup>b</sup>	0.10	(47)	-37.51 <sup>c</sup>
Rhode Island	1.45 <sup>b</sup>	0.21	(48)	-39.20 <sup>c</sup>
Massachusetts	1.39 <sup>b</sup>	0.09	(49)	-41.90 <sup>c</sup>
Utah	1.30 <sup>b</sup>	0.16	(50)	-45.74 <sup>c</sup>
Vermont	1.14 <sup>b</sup>	0.25	(51)	-52.17 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.04 <sup>b</sup>	0.15	(04)	27.32 <sup>c</sup>	Montana	1.94	0.28	(36)	-18.64 <sup>c</sup>
Alaska	1.99	0.39	(34)	-16.66 <sup>c</sup>	Nebraska	1.59 <sup>b</sup>	0.18	(45)	-33.64 <sup>c</sup>
Arizona	1.90 <sup>b</sup>	0.11	(41)	-20.44 <sup>c</sup>	Nevada	2.69	0.21	(12)	12.61
Arkansas	3.32 <sup>b</sup>	0.21	(03)	39.10 <sup>c</sup>	New Hampshire	2.09	0.24	(32)	-12.71
California	2.35	0.05	(23)	-1.68	New Jersey	2.44	0.10	(20)	2.12
Colorado	1.74 <sup>b</sup>	0.12	(42)	-27.26 <sup>c</sup>	New Mexico	2.65	0.23	(15)	10.97
Connecticut	1.64 <sup>b</sup>	0.13	(44)	-31.34 <sup>c</sup>	New York	2.38	0.07	(22)	-0.34
Delaware	2.18	0.30	(27)	-8.67	North Carolina	2.35	0.10	(24)	-1.88
D.C.	2.94	0.43	(05)	23.07 <sup>c</sup>	North Dakota	1.54	0.30	(46)	-35.65 <sup>c</sup>
Florida	2.68 <sup>b</sup>	0.07	(13)	12.17	Ohio	2.59	0.09	(17)	8.37
Georgia	2.68	0.11	(14)	12.16	Oklahoma	2.91	0.18	(07)	21.63 <sup>c</sup>
Hawaii	1.93	0.24	(37)	-19.29 <sup>c</sup>	Oregon	1.92	0.14	(38)	-19.78 <sup>c</sup>
Idaho	2.14	0.24	(30)	-10.46	Pennsylvania	2.25	0.08	(26)	-5.72
Illinois	2.63	0.09	(16)	9.83	Rhode Island	1.45 <sup>b</sup>	0.21	(48)	-39.20 <sup>c</sup>
Indiana	2.38	0.12	(21)	-0.31	South Carolina	2.85	0.15	(10)	19.03 <sup>c</sup>
Iowa	2.11	0.16	(31)	-11.87	South Dakota	1.91	0.30	(39)	-20.18 <sup>c</sup>
Kansas	2.18	0.18	(28)	-8.81	Tennessee	2.79	0.13	(11)	16.79 <sup>c</sup>
Kentucky	2.91	0.16	(06)	21.92 <sup>c</sup>	Texas	2.91 <sup>b</sup>	0.07	(08)	21.59 <sup>c</sup>
Louisiana	3.35 <sup>b</sup>	0.17	(02)	40.17 <sup>c</sup>	Utah	1.30 <sup>b</sup>	0.16	(50)	-45.74 <sup>c</sup>
Maine	1.90	0.22	(40)	-20.34 <sup>c</sup>	Vermont	1.14 <sup>b</sup>	0.25	(51)	-52.17 <sup>c</sup>
Maryland	2.28	0.12	(25)	-4.77	Virginia	2.18	0.10	(29)	-8.89
Massachusetts	1.39 <sup>b</sup>	0.09	(49)	-41.90 <sup>c</sup>	Washington	1.96 <sup>b</sup>	0.11	(35)	-18.16 <sup>c</sup>
Michigan	2.00 <sup>b</sup>	0.09	(33)	-16.17 <sup>c</sup>	West Virginia	2.90	0.24	(09)	21.51 <sup>c</sup>
Minnesota	1.49 <sup>b</sup>	0.10	(47)	-37.51 <sup>c</sup>	Wisconsin	1.70 <sup>b</sup>	0.11	(43)	-28.98 <sup>c</sup>
Mississippi	3.88 <sup>b</sup>	0.22	(01)	62.23 <sup>c</sup>	Wyoming	2.53	0.45	(19)	6.01
Missouri	2.57	0.13	(18)	7.71					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 6.25  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	16.69	0.03		
High Five States				
West Virginia	19.96 <sup>b</sup>	0.42	(01)	19.59 <sup>c</sup>
Mississippi	19.92 <sup>b</sup>	0.37	(02)	19.35 <sup>c</sup>
D.C.	19.89 <sup>b</sup>	0.83	(03)	19.17 <sup>c</sup>
Kentucky	19.85 <sup>b</sup>	0.30	(04)	18.91 <sup>c</sup>
Louisiana	19.70 <sup>b</sup>	0.30	(05)	18.04 <sup>c</sup>
Low Five States				
Connecticut	14.82 <sup>b</sup>	0.27	(47)	-11.20 <sup>c</sup>
Idaho	14.52 <sup>b</sup>	0.45	(48)	-13.00 <sup>c</sup>
Hawaii	14.37 <sup>b</sup>	0.44	(49)	-13.92 <sup>c</sup>
Arizona	13.99 <sup>b</sup>	0.21	(50)	-16.17 <sup>c</sup>
Utah	12.23 <sup>b</sup>	0.35	(51)	-26.73 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	18.36 <sup>b</sup>	0.27	(09)	10.02 <sup>c</sup>	Montana	16.11	0.54	(36)	-3.49
Alaska	17.02	0.95	(24)	1.99	Nebraska	18.36 <sup>b</sup>	0.43	(10)	9.97
Arizona	13.99 <sup>b</sup>	0.21	(50)	-16.17 <sup>c</sup>	Nevada	17.89	0.40	(16)	7.17
Arkansas	18.40 <sup>b</sup>	0.34	(08)	10.24 <sup>c</sup>	New Hampshire	15.83	0.47	(38)	-5.18
California	15.17 <sup>b</sup>	0.10	(43)	-9.12	New Jersey	18.16 <sup>b</sup>	0.20	(13)	8.78
Colorado	14.92 <sup>b</sup>	0.27	(46)	-10.61 <sup>c</sup>	New Mexico	15.82	0.40	(39)	-5.21
Connecticut	14.82 <sup>b</sup>	0.27	(47)	-11.20 <sup>c</sup>	New York	16.23 <sup>b</sup>	0.12	(33)	-2.79
Delaware	17.17	0.60	(21)	2.86	North Carolina	16.28	0.19	(30)	-2.49
D.C.	19.89 <sup>b</sup>	0.83	(03)	19.17 <sup>c</sup>	North Dakota	17.70	0.68	(18)	6.01
Florida	15.43 <sup>b</sup>	0.11	(42)	-7.56	Ohio	18.41 <sup>b</sup>	0.17	(07)	10.30 <sup>c</sup>
Georgia	16.36	0.20	(29)	-1.98	Oklahoma	18.34 <sup>b</sup>	0.31	(11)	9.88
Hawaii	14.37 <sup>b</sup>	0.44	(49)	-13.92 <sup>c</sup>	Oregon	15.99	0.28	(37)	-4.18
Idaho	14.52 <sup>b</sup>	0.45	(48)	-13.00 <sup>c</sup>	Pennsylvania	18.13 <sup>b</sup>	0.15	(14)	8.61
Illinois	18.42 <sup>b</sup>	0.17	(06)	10.34 <sup>c</sup>	Rhode Island	15.82	0.51	(40)	-5.22
Indiana	18.10 <sup>b</sup>	0.23	(15)	8.46	South Carolina	16.88	0.27	(27)	1.11
Iowa	17.52	0.31	(20)	4.97	South Dakota	16.96	0.61	(25)	1.59
Kansas	17.11	0.34	(22)	2.53	Tennessee	18.21 <sup>b</sup>	0.24	(12)	9.07
Kentucky	19.85 <sup>b</sup>	0.30	(04)	18.91 <sup>c</sup>	Texas	16.23 <sup>b</sup>	0.13	(31)	-2.78
Louisiana	19.70 <sup>b</sup>	0.30	(05)	18.04 <sup>c</sup>	Utah	12.23 <sup>b</sup>	0.35	(51)	-26.73 <sup>c</sup>
Maine	16.92	0.46	(26)	1.34	Vermont	16.15	0.68	(35)	-3.22
Maryland	17.70 <sup>b</sup>	0.25	(17)	6.04	Virginia	16.62	0.21	(28)	-0.42
Massachusetts	16.17	0.21	(34)	-3.14	Washington	14.93 <sup>b</sup>	0.22	(45)	-10.59 <sup>c</sup>
Michigan	17.05	0.18	(23)	2.12	West Virginia	19.96 <sup>b</sup>	0.42	(01)	19.59 <sup>c</sup>
Minnesota	14.97 <sup>b</sup>	0.24	(44)	-10.32 <sup>c</sup>	Wisconsin	15.51 <sup>b</sup>	0.22	(41)	-7.07
Mississippi	19.92 <sup>b</sup>	0.37	(02)	19.35 <sup>c</sup>	Wyoming	16.23	0.79	(32)	-2.78
Missouri	17.61 <sup>b</sup>	0.23	(19)	5.47					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.26  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	20.15	0.06		
High Five States				
Louisiana	25.06 <sup>b</sup>	0.53	(01)	24.34 <sup>c</sup>
Mississippi	24.95 <sup>b</sup>	0.65	(02)	23.79 <sup>c</sup>
Kentucky	24.30 <sup>b</sup>	0.52	(03)	20.58 <sup>c</sup>
West Virginia	24.21 <sup>b</sup>	0.72	(04)	20.13 <sup>c</sup>
D.C.	23.07	1.44	(05)	14.47 <sup>c</sup>
Low Five States				
Colorado	17.41 <sup>b</sup>	0.45	(47)	-13.61 <sup>c</sup>
Connecticut	17.27 <sup>b</sup>	0.47	(48)	-14.28 <sup>c</sup>
Arizona	16.79 <sup>b</sup>	0.34	(49)	-16.67 <sup>c</sup>
Idaho	15.88 <sup>b</sup>	0.71	(50)	-21.19 <sup>c</sup>
Utah	14.33 <sup>b</sup>	0.58	(51)	-28.88 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	22.91 <sup>b</sup>	0.48	(06)	13.71 <sup>c</sup>	Montana	17.76	0.86	(45)	-11.87 <sup>c</sup>
Alaska	20.04	1.55	(29)	-0.54	Nebraska	22.46	0.74	(12)	11.47 <sup>c</sup>
Arizona	16.79 <sup>b</sup>	0.34	(49)	-16.67 <sup>c</sup>	Nevada	20.71	0.63	(20)	2.80
Arkansas	22.54 <sup>b</sup>	0.58	(08)	11.87 <sup>c</sup>	New Hampshire	19.33	0.82	(35)	-4.07
California	18.10 <sup>b</sup>	0.16	(43)	-10.17 <sup>c</sup>	New Jersey	21.99 <sup>b</sup>	0.34	(15)	9.12
Colorado	17.41 <sup>b</sup>	0.45	(47)	-13.61 <sup>c</sup>	New Mexico	18.73	0.66	(38)	-7.07
Connecticut	17.27 <sup>b</sup>	0.47	(48)	-14.28 <sup>c</sup>	New York	19.43	0.22	(34)	-3.57
Delaware	20.32	1.00	(24)	0.84	North Carolina	19.85	0.33	(31)	-1.50
D.C.	23.07	1.44	(05)	14.47 <sup>c</sup>	North Dakota	21.63	1.15	(17)	7.35
Florida	18.35 <sup>b</sup>	0.19	(42)	-8.95	Ohio	22.50 <sup>b</sup>	0.30	(09)	11.66 <sup>c</sup>
Georgia	20.18	0.36	(26)	0.12	Oklahoma	22.91 <sup>b</sup>	0.53	(07)	13.69 <sup>c</sup>
Hawaii	18.69	0.76	(40)	-7.27	Oregon	18.50 <sup>b</sup>	0.46	(41)	-8.20
Idaho	15.88 <sup>b</sup>	0.71	(50)	-21.19 <sup>c</sup>	Pennsylvania	22.29 <sup>b</sup>	0.27	(14)	10.60 <sup>c</sup>
Illinois	22.48 <sup>b</sup>	0.30	(11)	11.57 <sup>c</sup>	Rhode Island	19.60	0.90	(33)	-2.74
Indiana	22.49 <sup>b</sup>	0.41	(10)	11.60 <sup>c</sup>	South Carolina	20.48	0.46	(23)	1.63
Iowa	20.62	0.53	(21)	2.31	South Dakota	20.12	1.00	(28)	-0.15
Kansas	21.17	0.58	(19)	5.08	Tennessee	22.36 <sup>b</sup>	0.42	(13)	10.94 <sup>c</sup>
Kentucky	24.30 <sup>b</sup>	0.52	(03)	20.58 <sup>c</sup>	Texas	20.22	0.22	(25)	0.32
Louisiana	25.06 <sup>b</sup>	0.53	(01)	24.34 <sup>c</sup>	Utah	14.33 <sup>b</sup>	0.58	(51)	-28.88 <sup>c</sup>
Maine	20.51	0.79	(22)	1.76	Vermont	18.81	1.15	(37)	-6.63
Maryland	21.97 <sup>b</sup>	0.45	(16)	9.04	Virginia	19.95	0.37	(30)	-1.01
Massachusetts	19.62	0.37	(32)	-2.63	Washington	17.66 <sup>b</sup>	0.36	(46)	-12.35 <sup>c</sup>
Michigan	20.17	0.31	(27)	0.07	West Virginia	24.21 <sup>b</sup>	0.72	(04)	20.13 <sup>c</sup>
Minnesota	18.01 <sup>b</sup>	0.40	(44)	-10.63 <sup>c</sup>	Wisconsin	18.69 <sup>b</sup>	0.39	(39)	-7.25
Mississippi	24.95 <sup>b</sup>	0.65	(02)	23.79 <sup>c</sup>	Wyoming	18.94	1.29	(36)	-6.02
Missouri	21.59 <sup>b</sup>	0.40	(18)	7.16					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.27  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	14.08	0.04		
High Five States				
D.C.	17.74 <sup>b</sup>	1.02	(01)	26.00 <sup>c</sup>
West Virginia	16.85 <sup>b</sup>	0.52	(02)	19.68 <sup>c</sup>
Kentucky	16.60 <sup>b</sup>	0.36	(03)	17.89 <sup>c</sup>
Mississippi	16.23 <sup>b</sup>	0.44	(04)	15.27 <sup>c</sup>
Louisiana	15.70 <sup>b</sup>	0.35	(05)	11.55 <sup>c</sup>
Low Five States				
Washington	12.69 <sup>b</sup>	0.27	(47)	-9.88
Minnesota	12.65 <sup>b</sup>	0.29	(48)	-10.16 <sup>c</sup>
Arizona	11.64 <sup>b</sup>	0.25	(49)	-17.29 <sup>c</sup>
Hawaii	10.78 <sup>b</sup>	0.52	(50)	-23.40 <sup>c</sup>
Utah	10.44 <sup>b</sup>	0.44	(51)	-25.83 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	15.09	0.32	(14)	7.23	Montana	14.73	0.70	(19)	4.62
Alaska	14.13	1.18	(29)	0.40	Nebraska	15.07	0.52	(15)	7.05
Arizona	11.64 <sup>b</sup>	0.25	(49)	-17.29 <sup>c</sup>	Nevada	15.31	0.50	(09)	8.73
Arkansas	15.18	0.41	(12)	7.85	New Hampshire	13.21	0.57	(40)	-6.14
California	12.92 <sup>b</sup>	0.12	(46)	-8.19	New Jersey	15.48 <sup>b</sup>	0.24	(08)	9.98
Colorado	13.01	0.33	(44)	-7.56	New Mexico	13.54	0.50	(37)	-3.82
Connecticut	13.04	0.34	(43)	-7.35	New York	14.00	0.15	(31)	-0.52
Delaware	14.34	0.72	(24)	1.85	North Carolina	13.63	0.23	(36)	-3.14
D.C.	17.74 <sup>b</sup>	1.02	(01)	26.00 <sup>c</sup>	North Dakota	14.79	0.84	(18)	5.06
Florida	12.98 <sup>b</sup>	0.14	(45)	-7.80	Ohio	15.49 <sup>b</sup>	0.20	(07)	10.06 <sup>c</sup>
Georgia	13.75	0.24	(35)	-2.29	Oklahoma	14.80	0.37	(17)	5.13
Hawaii	10.78 <sup>b</sup>	0.52	(50)	-23.40 <sup>c</sup>	Oregon	13.92	0.35	(33)	-1.14
Idaho	13.42	0.59	(38)	-4.69	Pennsylvania	15.21 <sup>b</sup>	0.18	(11)	8.04
Illinois	15.60 <sup>b</sup>	0.20	(06)	10.83 <sup>c</sup>	Rhode Island	13.28	0.61	(39)	-5.63
Indiana	14.96	0.28	(16)	6.27	South Carolina	14.06	0.32	(30)	-0.09
Iowa	15.23	0.39	(10)	8.22	South Dakota	14.23	0.74	(25)	1.09
Kansas	14.00	0.41	(32)	-0.53	Tennessee	15.15 <sup>b</sup>	0.29	(13)	7.61
Kentucky	16.60 <sup>b</sup>	0.36	(03)	17.89 <sup>c</sup>	Texas	13.14 <sup>b</sup>	0.15	(41)	-6.63
Louisiana	15.70 <sup>b</sup>	0.35	(05)	11.55 <sup>c</sup>	Utah	10.44 <sup>b</sup>	0.44	(51)	-25.83 <sup>c</sup>
Maine	14.38	0.56	(23)	2.15	Vermont	14.23	0.84	(26)	1.09
Maryland	14.56	0.30	(22)	3.46	Virginia	14.19	0.26	(27)	0.80
Massachusetts	13.81	0.25	(34)	-1.88	Washington	12.69 <sup>b</sup>	0.27	(47)	-9.88
Michigan	14.72	0.22	(20)	4.54	West Virginia	16.85 <sup>b</sup>	0.52	(02)	19.68 <sup>c</sup>
Minnesota	12.65 <sup>b</sup>	0.29	(48)	-10.16 <sup>c</sup>	Wisconsin	13.13	0.27	(42)	-6.70
Mississippi	16.23 <sup>b</sup>	0.44	(04)	15.27 <sup>c</sup>	Wyoming	14.18	0.99	(28)	0.71
Missouri	14.62	0.28	(21)	3.89					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 7.14  
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.18	0.02		
High Five States				
D.C.	6.35 <sup>b</sup>	0.61	(01)	51.94 <sup>c</sup>
New Jersey	5.17 <sup>b</sup>	0.14	(02)	23.77 <sup>c</sup>
Delaware	5.16	0.44	(03)	23.55 <sup>c</sup>
New York	5.09 <sup>b</sup>	0.09	(04)	21.93 <sup>c</sup>
Pennsylvania	4.93 <sup>b</sup>	0.11	(05)	18.13 <sup>c</sup>
Low Five States				
Wyoming	3.17	0.48	(47)	-23.98 <sup>c</sup>
Arizona	2.96 <sup>b</sup>	0.13	(48)	-29.15 <sup>c</sup>
Utah	2.85 <sup>b</sup>	0.23	(49)	-31.68 <sup>c</sup>
Arkansas	2.80 <sup>b</sup>	0.18	(50)	-32.99 <sup>c</sup>
Alaska	1.69 <sup>b</sup>	0.39	(51)	-59.59 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.37 <sup>b</sup>	0.15	(45)	-19.30 <sup>c</sup>	Montana	4.29	0.38	(18)	2.82
Alaska	1.69 <sup>b</sup>	0.39	(51)	-59.59 <sup>c</sup>	Nebraska	4.72	0.30	(08)	13.05
Arizona	2.96 <sup>b</sup>	0.13	(48)	-29.15 <sup>c</sup>	Nevada	3.32 <sup>b</sup>	0.23	(46)	-20.38 <sup>c</sup>
Arkansas	2.80 <sup>b</sup>	0.18	(50)	-32.99 <sup>c</sup>	New Hampshire	4.39	0.33	(15)	5.20
California	4.10	0.07	(26)	-1.86	New Jersey	5.17 <sup>b</sup>	0.14	(02)	23.77 <sup>c</sup>
Colorado	3.39 <sup>b</sup>	0.17	(44)	-18.79 <sup>c</sup>	New Mexico	3.42	0.25	(43)	-18.09 <sup>c</sup>
Connecticut	4.54	0.20	(14)	8.67	New York	5.09 <sup>b</sup>	0.09	(04)	21.93 <sup>c</sup>
Delaware	5.16	0.44	(03)	23.55 <sup>c</sup>	North Carolina	4.00	0.12	(30)	-4.27
D.C.	6.35 <sup>b</sup>	0.61	(01)	51.94 <sup>c</sup>	North Dakota	4.70	0.49	(10)	12.57
Florida	3.74 <sup>b</sup>	0.08	(37)	-10.41	Ohio	4.61 <sup>b</sup>	0.11	(13)	10.49
Georgia	3.63 <sup>b</sup>	0.12	(39)	-12.97	Oklahoma	3.91	0.19	(31)	-6.39
Hawaii	3.76	0.31	(36)	-9.88	Oregon	3.81	0.18	(33)	-8.78
Idaho	4.25	0.33	(19)	1.81	Pennsylvania	4.93 <sup>b</sup>	0.11	(05)	18.13 <sup>c</sup>
Illinois	4.63 <sup>b</sup>	0.11	(12)	10.75	Rhode Island	3.77	0.33	(34)	-9.69
Indiana	4.38	0.15	(17)	4.82	South Carolina	4.08	0.17	(28)	-2.26
Iowa	4.90	0.23	(06)	17.24 <sup>c</sup>	South Dakota	4.14	0.42	(25)	-0.94
Kansas	3.53	0.21	(41)	-15.37 <sup>c</sup>	Tennessee	3.77	0.14	(35)	-9.84
Kentucky	3.51 <sup>b</sup>	0.17	(42)	-15.90 <sup>c</sup>	Texas	3.60 <sup>b</sup>	0.08	(40)	-13.74
Louisiana	3.68	0.17	(38)	-11.98	Utah	2.85 <sup>b</sup>	0.23	(49)	-31.68 <sup>c</sup>
Maine	4.22	0.31	(20)	1.11	Vermont	4.38	0.48	(16)	4.93
Maryland	4.64	0.17	(11)	11.14	Virginia	4.19	0.14	(24)	0.29
Massachusetts	4.19	0.14	(23)	0.39	Washington	4.04	0.15	(29)	-3.23
Michigan	4.82 <sup>b</sup>	0.13	(07)	15.44 <sup>c</sup>	West Virginia	4.21	0.26	(21)	0.75
Minnesota	4.09	0.17	(27)	-2.04	Wisconsin	4.71	0.17	(09)	12.74
Mississippi	3.83	0.21	(32)	-8.30	Wyoming	3.17	0.48	(47)	-23.98 <sup>c</sup>
Missouri	4.20	0.15	(22)	0.52					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.18  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.31	0.02		
High Five States				
Maine	6.13 <sup>b</sup>	0.28	(01)	42.19 <sup>c</sup>
New Hampshire	5.97 <sup>b</sup>	0.29	(02)	38.65 <sup>c</sup>
D.C.	5.41	0.43	(03)	25.61 <sup>c</sup>
Rhode Island	5.36	0.30	(04)	24.32 <sup>c</sup>
Massachusetts	5.27 <sup>b</sup>	0.12	(05)	22.27 <sup>c</sup>
Low Five States				
Texas	3.59 <sup>b</sup>	0.06	(47)	-16.59 <sup>c</sup>
California	3.59 <sup>b</sup>	0.05	(48)	-16.71 <sup>c</sup>
New Mexico	3.35 <sup>b</sup>	0.18	(49)	-22.16 <sup>c</sup>
Hawaii	3.22 <sup>b</sup>	0.21	(50)	-25.30 <sup>c</sup>
Utah	2.92 <sup>b</sup>	0.17	(51)	-32.27 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.91	0.12	(44)	-9.23	Montana	4.34	0.28	(31)	0.75
Alaska	4.43	0.45	(25)	2.82	Nebraska	4.33	0.21	(32)	0.52
Arizona	3.89 <sup>b</sup>	0.11	(46)	-9.74	Nevada	4.65	0.20	(16)	8.00
Arkansas	3.90	0.16	(45)	-9.50	New Hampshire	5.97 <sup>b</sup>	0.29	(02)	38.65 <sup>c</sup>
California	3.59 <sup>b</sup>	0.05	(48)	-16.71 <sup>c</sup>	New Jersey	4.13	0.09	(37)	-4.16
Colorado	4.04	0.14	(39)	-6.32	New Mexico	3.35 <sup>b</sup>	0.18	(49)	-22.16 <sup>c</sup>
Connecticut	4.51	0.15	(20)	4.75	New York	4.24	0.06	(35)	-1.61
Delaware	4.24	0.30	(34)	-1.49	North Carolina	3.94 <sup>b</sup>	0.09	(41)	-8.63
D.C.	5.41	0.43	(03)	25.61 <sup>c</sup>	North Dakota	4.03	0.33	(40)	-6.38
Florida	4.09 <sup>b</sup>	0.06	(38)	-5.07	Ohio	5.18 <sup>b</sup>	0.09	(06)	20.18 <sup>c</sup>
Georgia	3.93 <sup>b</sup>	0.10	(42)	-8.75	Oklahoma	4.62	0.15	(18)	7.29
Hawaii	3.22 <sup>b</sup>	0.21	(50)	-25.30 <sup>c</sup>	Oregon	4.80	0.15	(12)	11.39
Idaho	4.49	0.25	(21)	4.18	Pennsylvania	5.03 <sup>b</sup>	0.08	(07)	16.83 <sup>c</sup>
Illinois	4.63 <sup>b</sup>	0.08	(17)	7.38	Rhode Island	5.36	0.30	(04)	24.32 <sup>c</sup>
Indiana	4.84 <sup>b</sup>	0.12	(09)	12.36	South Carolina	4.43	0.14	(26)	2.73
Iowa	4.75	0.17	(14)	10.37	South Dakota	4.78	0.33	(13)	11.02
Kansas	3.91	0.16	(43)	-9.12	Tennessee	4.18	0.11	(36)	-2.94
Kentucky	4.60	0.14	(19)	6.88	Texas	3.59 <sup>b</sup>	0.06	(47)	-16.59 <sup>c</sup>
Louisiana	4.28	0.14	(33)	-0.69	Utah	2.92 <sup>b</sup>	0.17	(51)	-32.27 <sup>c</sup>
Maine	6.13 <sup>b</sup>	0.28	(01)	42.19 <sup>c</sup>	Vermont	4.47	0.36	(23)	3.68
Maryland	4.48	0.13	(22)	3.89	Virginia	4.35	0.11	(30)	0.88
Massachusetts	5.27 <sup>b</sup>	0.12	(05)	22.27 <sup>c</sup>	Washington	4.84 <sup>b</sup>	0.12	(11)	12.28
Michigan	4.84 <sup>b</sup>	0.10	(10)	12.32	West Virginia	4.73	0.20	(15)	9.78
Minnesota	4.46	0.13	(24)	3.63	Wisconsin	5.02 <sup>b</sup>	0.13	(08)	16.52 <sup>c</sup>
Mississippi	4.38	0.17	(29)	1.62	Wyoming	4.42	0.41	(27)	2.56
Missouri	4.41	0.12	(28)	2.47					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.19  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.73	0.03		
High Five States				
Maine	11.54 <sup>b</sup>	0.58	(01)	49.30 <sup>c</sup>
New Hampshire	10.70 <sup>b</sup>	0.60	(02)	38.47 <sup>c</sup>
D.C.	9.82	0.91	(03)	26.98 <sup>c</sup>
Rhode Island	9.80	0.63	(04)	26.78 <sup>c</sup>
Ohio	9.58 <sup>b</sup>	0.19	(05)	23.98 <sup>c</sup>
Low Five States				
Texas	6.58 <sup>b</sup>	0.12	(47)	-14.94
California	6.24 <sup>b</sup>	0.09	(48)	-19.34 <sup>c</sup>
New Mexico	5.87 <sup>b</sup>	0.36	(49)	-24.02 <sup>c</sup>
Hawaii	5.63 <sup>b</sup>	0.41	(50)	-27.22 <sup>c</sup>
Utah	5.37 <sup>b</sup>	0.35	(51)	-30.54 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.43	0.26	(36)	-3.83	Montana	7.52	0.54	(33)	-2.76
Alaska	7.04	0.83	(43)	-8.99	Nebraska	7.84	0.43	(29)	1.38
Arizona	6.93 <sup>b</sup>	0.22	(45)	-10.30	Nevada	7.95	0.38	(26)	2.85
Arkansas	6.87	0.31	(46)	-11.10	New Hampshire	10.70 <sup>b</sup>	0.60	(02)	38.47 <sup>c</sup>
California	6.24 <sup>b</sup>	0.09	(48)	-19.34 <sup>c</sup>	New Jersey	7.16	0.19	(40)	-7.35
Colorado	7.28	0.28	(38)	-5.83	New Mexico	5.87 <sup>b</sup>	0.36	(49)	-24.02 <sup>c</sup>
Connecticut	8.22	0.32	(19)	6.29	New York	7.51	0.13	(34)	-2.80
Delaware	8.11	0.63	(21)	4.87	North Carolina	7.15	0.19	(41)	-7.52
D.C.	9.82	0.91	(03)	26.98 <sup>c</sup>	North Dakota	7.08	0.65	(42)	-8.45
Florida	7.22 <sup>b</sup>	0.12	(39)	-6.60	Ohio	9.58 <sup>b</sup>	0.19	(05)	23.98 <sup>c</sup>
Georgia	7.01 <sup>b</sup>	0.20	(44)	-9.31	Oklahoma	8.46	0.32	(16)	9.48
Hawaii	5.63 <sup>b</sup>	0.41	(50)	-27.22 <sup>c</sup>	Oregon	8.50	0.31	(15)	9.91
Idaho	7.87	0.49	(27)	1.77	Pennsylvania	9.31 <sup>b</sup>	0.17	(07)	20.47 <sup>c</sup>
Illinois	8.52 <sup>b</sup>	0.18	(14)	10.26	Rhode Island	9.80	0.63	(04)	26.78 <sup>c</sup>
Indiana	8.91 <sup>b</sup>	0.25	(09)	15.30 <sup>c</sup>	South Carolina	8.03	0.28	(22)	3.83
Iowa	8.69	0.34	(11)	12.36	South Dakota	8.73	0.66	(10)	12.89
Kansas	7.43	0.34	(37)	-3.95	Tennessee	7.46	0.23	(35)	-3.49
Kentucky	8.34	0.29	(18)	7.86	Texas	6.58 <sup>b</sup>	0.12	(47)	-14.94
Louisiana	7.65	0.28	(31)	-1.00	Utah	5.37 <sup>b</sup>	0.35	(51)	-30.54 <sup>c</sup>
Maine	11.54 <sup>b</sup>	0.58	(01)	49.30 <sup>c</sup>	Vermont	8.20	0.75	(20)	6.09
Maryland	8.01	0.26	(23)	3.57	Virginia	7.84	0.22	(28)	1.43
Massachusetts	9.45 <sup>b</sup>	0.25	(06)	22.27 <sup>c</sup>	Washington	8.41	0.25	(17)	8.74
Michigan	8.60 <sup>b</sup>	0.19	(13)	11.31	West Virginia	8.66	0.42	(12)	11.99
Minnesota	7.83	0.26	(30)	1.33	Wisconsin	8.94 <sup>b</sup>	0.26	(08)	15.65 <sup>c</sup>
Mississippi	7.96	0.35	(25)	2.95	Wyoming	7.60	0.80	(32)	-1.73
Missouri	8.00	0.24	(24)	3.47					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.





Table 8.20  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.61	0.01		
High Five States				
D.C.	2.22	0.36	(01)	38.54 <sup>c</sup>
New Hampshire	2.21	0.23	(02)	37.33 <sup>c</sup>
Massachusetts	2.07 <sup>b</sup>	0.10	(03)	28.98 <sup>c</sup>
Rhode Island	2.07	0.24	(04)	28.66 <sup>c</sup>
Washington	1.94	0.11	(05)	20.74 <sup>c</sup>
Low Five States				
New Mexico	1.22	0.15	(47)	-23.81 <sup>c</sup>
Hawaii	1.19	0.17	(48)	-26.05 <sup>c</sup>
Alabama	1.14 <sup>b</sup>	0.09	(49)	-29.08 <sup>c</sup>
Kansas	1.14 <sup>b</sup>	0.12	(50)	-29.12 <sup>c</sup>
Utah	0.81 <sup>b</sup>	0.12	(51)	-49.45 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.14 <sup>b</sup>	0.09	(49)	-29.08 <sup>c</sup>	Montana	1.60	0.23	(26)	-0.41
Alaska	1.86	0.42	(06)	16.14 <sup>c</sup>	Nebraska	1.47	0.17	(40)	-8.40
Arizona	1.30 <sup>b</sup>	0.09	(44)	-18.91 <sup>c</sup>	Nevada	1.68	0.16	(21)	4.74
Arkansas	1.48	0.13	(38)	-7.91	New Hampshire	2.21	0.23	(02)	37.33 <sup>c</sup>
California	1.51	0.04	(36)	-6.24	New Jersey	1.76	0.08	(14)	9.58
Colorado	1.40	0.11	(43)	-12.96	New Mexico	1.22	0.15	(47)	-23.81 <sup>c</sup>
Connecticut	1.74	0.13	(16)	8.35	New York	1.80 <sup>b</sup>	0.05	(12)	11.84
Delaware	1.25	0.22	(45)	-22.20 <sup>c</sup>	North Carolina	1.45	0.07	(41)	-9.94
D.C.	2.22	0.36	(01)	38.54 <sup>c</sup>	North Dakota	1.55	0.28	(34)	-3.42
Florida	1.48	0.05	(39)	-7.98	Ohio	1.84	0.07	(08)	14.74
Georgia	1.61	0.08	(25)	0.14	Oklahoma	1.53	0.12	(35)	-4.70
Hawaii	1.19	0.17	(48)	-26.05 <sup>c</sup>	Oregon	1.82	0.13	(10)	13.45
Idaho	1.49	0.20	(37)	-7.32	Pennsylvania	1.73	0.06	(17)	7.94
Illinois	1.69	0.07	(19)	5.35	Rhode Island	2.07	0.24	(04)	28.66 <sup>c</sup>
Indiana	1.66	0.10	(22)	3.36	South Carolina	1.59	0.11	(29)	-0.99
Iowa	1.62	0.13	(24)	0.72	South Dakota	1.43	0.24	(42)	-11.02
Kansas	1.14 <sup>b</sup>	0.12	(50)	-29.12 <sup>c</sup>	Tennessee	1.56	0.09	(32)	-2.86
Kentucky	1.59	0.11	(28)	-0.81	Texas	1.24 <sup>b</sup>	0.05	(46)	-23.04 <sup>c</sup>
Louisiana	1.60	0.11	(27)	-0.41	Utah	0.81 <sup>b</sup>	0.12	(51)	-49.45 <sup>c</sup>
Maine	1.80	0.20	(11)	11.88	Vermont	1.68	0.29	(20)	4.80
Maryland	1.80	0.11	(13)	11.82	Virginia	1.57	0.08	(30)	-2.09
Massachusetts	2.07 <sup>b</sup>	0.10	(03)	28.98 <sup>c</sup>	Washington	1.94	0.11	(05)	20.74 <sup>c</sup>
Michigan	1.84	0.08	(09)	14.61	West Virginia	1.55	0.16	(33)	-3.20
Minnesota	1.74	0.11	(15)	8.62	Wisconsin	1.85	0.10	(07)	15.22 <sup>c</sup>
Mississippi	1.56	0.14	(31)	-2.77	Wyoming	1.71	0.35	(18)	6.52
Missouri	1.64	0.10	(23)	2.44					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.18  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.41	0.01		
High Five States				
Maine	0.54	0.09	(01)	31.45 <sup>c</sup>
Louisiana	0.53	0.05	(02)	31.03 <sup>c</sup>
Vermont	0.52	0.13	(03)	27.60 <sup>c</sup>
Wisconsin	0.47	0.04	(04)	16.23 <sup>c</sup>
Tennessee	0.47	0.04	(05)	16.06 <sup>c</sup>
Low Five States				
Maryland	0.35	0.04	(40)	-14.02
Utah	0.35	0.06	(41)	-14.97
Georgia	0.33	0.03	(42)	-17.95 <sup>c</sup>
Arizona	0.33	0.03	(43)	-20.06 <sup>c</sup>
New Hampshire	0.31	0.07	(44)	-23.20 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.41	0.04	(28)	-0.01	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.41	0.07	(25)	0.30
Arizona	0.33	0.03	(43)	-20.06 <sup>c</sup>	Nevada	0.43	0.06	(18)	4.35
Arkansas	0.44	0.05	(11)	7.97	New Hampshire	0.31	0.07	(44)	-23.20 <sup>c</sup>
California	0.41	0.02	(26)	0.17	New Jersey	0.44	0.03	(12)	7.55
Colorado	0.37	0.04	(35)	-8.61	New Mexico	0.38	0.06	(34)	-7.25
Connecticut	0.43	0.05	(15)	4.68	New York	0.43	0.02	(13)	5.23
Delaware	0.41	0.10	(23)	1.19	North Carolina	0.42	0.03	(19)	3.53
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.40	0.02	(31)	-1.59	Ohio	0.44	0.03	(10)	8.55
Georgia	0.33	0.03	(42)	-17.95 <sup>c</sup>	Oklahoma	0.42	0.05	(22)	2.10
Hawaii	-	-	(-)	-	Oregon	0.46	0.05	(06)	13.77
Idaho	0.41	0.08	(24)	0.40	Pennsylvania	0.45	0.03	(08)	10.81
Illinois	0.40	0.03	(32)	-2.18	Rhode Island	0.38	0.08	(33)	-6.64
Indiana	0.35	0.03	(39)	-13.18	South Carolina	0.40	0.04	(29)	-0.77
Iowa	0.42	0.05	(20)	2.94	South Dakota	-	-	(-)	-
Kansas	0.37	0.05	(37)	-10.02	Tennessee	0.47	0.04	(05)	16.06 <sup>c</sup>
Kentucky	0.42	0.04	(21)	2.14	Texas	0.40	0.02	(30)	-0.78
Louisiana	0.53	0.05	(02)	31.03 <sup>c</sup>	Utah	0.35	0.06	(41)	-14.97
Maine	0.54	0.09	(01)	31.45 <sup>c</sup>	Vermont	0.52	0.13	(03)	27.60 <sup>c</sup>
Maryland	0.35	0.04	(40)	-14.02	Virginia	0.43	0.03	(16)	4.67
Massachusetts	0.35	0.03	(38)	-13.02	Washington	0.37	0.03	(36)	-9.86
Michigan	0.43	0.03	(17)	4.42	West Virginia	0.46	0.07	(07)	12.62
Minnesota	0.43	0.04	(14)	4.90	Wisconsin	0.47	0.04	(04)	16.23 <sup>c</sup>
Mississippi	0.41	0.05	(27)	-0.01	Wyoming	-	-	(-)	-
Missouri	0.44	0.04	(09)	8.98					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.19  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.50	0.01		
High Five States				
Rhode Island	0.72	0.17	(01)	43.26 <sup>c</sup>
Maine	0.71	0.15	(02)	41.15 <sup>c</sup>
Louisiana	0.64	0.08	(03)	27.50 <sup>c</sup>
Wisconsin	0.62	0.07	(04)	23.06 <sup>c</sup>
Nebraska	0.60	0.12	(05)	19.22 <sup>c</sup>
Low Five States				
Arizona	0.42	0.05	(36)	-16.41 <sup>c</sup>
Utah	0.42	0.10	(37)	-16.58 <sup>c</sup>
Iowa	0.41	0.07	(38)	-17.93 <sup>c</sup>
Georgia	0.39	0.05	(39)	-21.10 <sup>c</sup>
Maryland	0.38	0.05	(40)	-23.62 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.49	0.07	(25)	-1.73	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.60	0.12	(05)	19.22 <sup>c</sup>
Arizona	0.42	0.05	(36)	-16.41 <sup>c</sup>	Nevada	0.59	0.11	(06)	17.32 <sup>c</sup>
Arkansas	0.49	0.09	(24)	-1.63	New Hampshire	-	-	(-)	-
California	0.50	0.03	(20)	0.09	New Jersey	0.54	0.05	(11)	7.34
Colorado	0.48	0.07	(27)	-3.58	New Mexico	0.45	0.10	(34)	-10.44
Connecticut	0.50	0.08	(19)	0.63	New York	0.52	0.03	(15)	4.33
Delaware	-	-	(-)	-	North Carolina	0.58	0.05	(07)	15.37 <sup>c</sup>
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.50	0.03	(22)	-0.86	Ohio	0.55	0.05	(09)	9.36
Georgia	0.39	0.05	(39)	-21.10 <sup>c</sup>	Oklahoma	0.50	0.08	(21)	-0.41
Hawaii	-	-	(-)	-	Oregon	0.54	0.08	(10)	8.91
Idaho	-	-	(-)	-	Pennsylvania	0.57	0.04	(08)	14.52
Illinois	0.47	0.04	(30)	-5.76	Rhode Island	0.72	0.17	(01)	43.26 <sup>c</sup>
Indiana	0.48	0.06	(28)	-4.93	South Carolina	0.46	0.07	(32)	-7.80
Iowa	0.41	0.07	(38)	-17.93 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.43	0.08	(35)	-14.20	Tennessee	0.51	0.06	(17)	2.18
Kentucky	0.54	0.07	(12)	7.26	Texas	0.51	0.03	(18)	1.23
Louisiana	0.64	0.08	(03)	27.50 <sup>c</sup>	Utah	0.42	0.10	(37)	-16.58 <sup>c</sup>
Maine	0.71	0.15	(02)	41.15 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.38	0.05	(40)	-23.62 <sup>c</sup>	Virginia	0.53	0.06	(14)	6.69
Massachusetts	0.47	0.06	(29)	-5.48	Washington	0.49	0.06	(23)	-1.09
Michigan	0.54	0.05	(13)	7.07	West Virginia	0.49	0.10	(26)	-2.86
Minnesota	0.52	0.07	(16)	3.24	Wisconsin	0.62	0.07	(04)	23.06 <sup>c</sup>
Mississippi	0.45	0.09	(33)	-9.71	Wyoming	-	-	(-)	-
Missouri	0.47	0.06	(31)	-6.46					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.20  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.33	0.01		
High Five States				
Louisiana	0.45	0.06	(01)	37.48 <sup>c</sup>
Tennessee	0.43	0.05	(02)	30.71 <sup>c</sup>
Iowa	0.42	0.07	(03)	26.90 <sup>c</sup>
West Virginia	0.42	0.09	(04)	26.87 <sup>c</sup>
Missouri	0.41	0.05	(05)	25.25 <sup>c</sup>
Low Five States				
Massachusetts	0.27	0.04	(34)	-19.84 <sup>c</sup>
Washington	0.26	0.04	(35)	-20.62 <sup>c</sup>
Indiana	0.26	0.04	(36)	-21.21 <sup>c</sup>
Colorado	0.26	0.05	(37)	-22.56 <sup>c</sup>
Arizona	0.24	0.04	(38)	-27.52 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.33	0.05	(23)	-0.05	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.24	0.04	(38)	-27.52 <sup>c</sup>	Nevada	0.28	0.07	(31)	-14.09
Arkansas	0.40	0.07	(07)	20.61 <sup>c</sup>	New Hampshire	-	-	(-)	-
California	0.34	0.02	(21)	1.70	New Jersey	0.35	0.04	(14)	7.20
Colorado	0.26	0.05	(37)	-22.56 <sup>c</sup>	New Mexico	0.30	0.07	(29)	-8.98
Connecticut	0.35	0.06	(15)	7.11	New York	0.35	0.03	(16)	6.91
Delaware	-	-	(-)	-	North Carolina	0.30	0.03	(30)	-10.47
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.32	0.02	(26)	-3.80	Ohio	0.36	0.03	(11)	9.01
Georgia	0.28	0.03	(33)	-15.01 <sup>c</sup>	Oklahoma	0.36	0.06	(10)	9.29
Hawaii	-	-	(-)	-	Oregon	0.40	0.06	(06)	22.14 <sup>c</sup>
Idaho	-	-	(-)	-	Pennsylvania	0.36	0.03	(13)	8.62
Illinois	0.34	0.03	(19)	2.37	Rhode Island	-	-	(-)	-
Indiana	0.26	0.04	(36)	-21.21 <sup>c</sup>	South Carolina	0.35	0.05	(18)	6.17
Iowa	0.42	0.07	(03)	26.90 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.33	0.07	(24)	-1.10	Tennessee	0.43	0.05	(02)	30.71 <sup>c</sup>
Kentucky	0.31	0.05	(28)	-6.15	Texas	0.32	0.02	(25)	-3.06
Louisiana	0.45	0.06	(01)	37.48 <sup>c</sup>	Utah	0.28	0.07	(32)	-14.71
Maine	0.40	0.10	(08)	19.56 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.31	0.05	(27)	-6.11	Virginia	0.34	0.04	(22)	1.69
Massachusetts	0.27	0.04	(34)	-19.84 <sup>c</sup>	Washington	0.26	0.04	(35)	-20.62 <sup>c</sup>
Michigan	0.34	0.03	(20)	1.84	West Virginia	0.42	0.09	(04)	26.87 <sup>c</sup>
Minnesota	0.36	0.05	(12)	8.84	Wisconsin	0.35	0.05	(17)	6.60
Mississippi	0.38	0.07	(09)	15.67 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	0.41	0.05	(05)	25.25 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.18  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.01	0.02		
High Five States				
North Dakota	5.36	0.39	(01)	33.80 <sup>c</sup>
Louisiana	5.06 <sup>b</sup>	0.15	(02)	26.32 <sup>c</sup>
Oklahoma	5.05 <sup>b</sup>	0.16	(03)	26.15 <sup>c</sup>
Iowa	4.94 <sup>b</sup>	0.17	(04)	23.35 <sup>c</sup>
West Virginia	4.79 <sup>b</sup>	0.21	(05)	19.60 <sup>c</sup>
Low Five States				
New York	3.39 <sup>b</sup>	0.06	(47)	-15.51 <sup>c</sup>
Connecticut	3.32 <sup>b</sup>	0.13	(48)	-17.17 <sup>c</sup>
Utah	3.16 <sup>b</sup>	0.18	(49)	-21.15 <sup>c</sup>
D.C.	2.88 <sup>b</sup>	0.32	(50)	-28.04 <sup>c</sup>
Hawaii	2.70 <sup>b</sup>	0.19	(51)	-32.69 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.96	0.13	(29)	-1.22	Montana	3.83	0.27	(35)	-4.41
Alaska	4.33	0.47	(18)	7.98	Nebraska	4.39	0.21	(15)	9.52
Arizona	3.69	0.11	(42)	-7.97	Nevada	3.96	0.18	(28)	-1.20
Arkansas	4.32	0.16	(19)	7.91	New Hampshire	3.83	0.23	(34)	-4.30
California	3.60 <sup>b</sup>	0.05	(44)	-10.10	New Jersey	3.53 <sup>b</sup>	0.09	(45)	-11.79
Colorado	3.82	0.13	(36)	-4.77	New Mexico	4.47	0.21	(12)	11.63
Connecticut	3.32 <sup>b</sup>	0.13	(48)	-17.17 <sup>c</sup>	New York	3.39 <sup>b</sup>	0.06	(47)	-15.51 <sup>c</sup>
Delaware	3.81	0.28	(37)	-4.94	North Carolina	4.16	0.10	(25)	3.78
D.C.	2.88 <sup>b</sup>	0.32	(50)	-28.04 <sup>c</sup>	North Dakota	5.36	0.39	(01)	33.80 <sup>c</sup>
Florida	3.49 <sup>b</sup>	0.05	(46)	-12.98	Ohio	4.52 <sup>b</sup>	0.08	(10)	12.90
Georgia	3.78	0.10	(39)	-5.54	Oklahoma	5.05 <sup>b</sup>	0.16	(03)	26.15 <sup>c</sup>
Hawaii	2.70 <sup>b</sup>	0.19	(51)	-32.69 <sup>c</sup>	Oregon	3.76	0.14	(40)	-6.15
Idaho	3.88	0.23	(33)	-3.29	Pennsylvania	4.04	0.07	(26)	0.75
Illinois	4.23	0.08	(23)	5.48	Rhode Island	3.89	0.25	(32)	-2.84
Indiana	4.54 <sup>b</sup>	0.12	(09)	13.43	South Carolina	4.29	0.14	(21)	6.96
Iowa	4.94 <sup>b</sup>	0.17	(04)	23.35 <sup>c</sup>	South Dakota	4.45	0.31	(13)	11.12
Kansas	4.66 <sup>b</sup>	0.18	(06)	16.26 <sup>c</sup>	Tennessee	4.27	0.11	(22)	6.49
Kentucky	4.62 <sup>b</sup>	0.14	(07)	15.26 <sup>c</sup>	Texas	4.56 <sup>b</sup>	0.07	(08)	13.87
Louisiana	5.06 <sup>b</sup>	0.15	(02)	26.32 <sup>c</sup>	Utah	3.16 <sup>b</sup>	0.18	(49)	-21.15 <sup>c</sup>
Maine	4.40	0.23	(14)	9.72	Vermont	3.79	0.33	(38)	-5.35
Maryland	3.90	0.12	(31)	-2.63	Virginia	3.99	0.10	(27)	-0.45
Massachusetts	3.72	0.10	(41)	-7.04	Washington	3.94	0.11	(30)	-1.68
Michigan	4.18	0.09	(24)	4.40	West Virginia	4.79 <sup>b</sup>	0.21	(05)	19.60 <sup>c</sup>
Minnesota	4.36	0.13	(16)	8.92	Wisconsin	4.32	0.12	(20)	7.88
Mississippi	4.49	0.17	(11)	12.01	Wyoming	3.62	0.36	(43)	-9.77
Missouri	4.35	0.12	(17)	8.65					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.19  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.84	0.03		
High Five States				
North Dakota	7.80	0.69	(01)	33.69 <sup>c</sup>
Iowa	7.24 <sup>b</sup>	0.31	(02)	24.04 <sup>c</sup>
Oklahoma	7.20 <sup>b</sup>	0.29	(03)	23.36 <sup>c</sup>
Louisiana	7.13 <sup>b</sup>	0.28	(04)	22.23 <sup>c</sup>
Kansas	6.84	0.33	(05)	17.26 <sup>c</sup>
Low Five States				
Florida	5.11 <sup>b</sup>	0.10	(47)	-12.51
Connecticut	4.75 <sup>b</sup>	0.24	(48)	-18.62 <sup>c</sup>
Utah	4.30 <sup>b</sup>	0.31	(49)	-26.32 <sup>c</sup>
D.C.	4.20	0.59	(50)	-28.09 <sup>c</sup>
Hawaii	4.11 <sup>b</sup>	0.36	(51)	-29.63 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.47	0.23	(38)	-6.34	Montana	5.51	0.48	(36)	-5.54
Alaska	5.76	0.81	(28)	-1.27	Nebraska	6.17	0.39	(22)	5.65
Arizona	5.16 <sup>b</sup>	0.19	(44)	-11.60	Nevada	5.65	0.33	(30)	-3.26
Arkansas	6.33	0.30	(17)	8.39	New Hampshire	5.56	0.43	(35)	-4.80
California	5.29 <sup>b</sup>	0.09	(42)	-9.45	New Jersey	5.30	0.17	(41)	-9.21
Colorado	5.58	0.25	(32)	-4.45	New Mexico	6.18	0.37	(21)	5.87
Connecticut	4.75 <sup>b</sup>	0.24	(48)	-18.62 <sup>c</sup>	New York	5.13 <sup>b</sup>	0.11	(45)	-12.16
Delaware	5.36	0.51	(40)	-8.19	North Carolina	6.47	0.19	(11)	10.90
D.C.	4.20	0.59	(50)	-28.09 <sup>c</sup>	North Dakota	7.80	0.69	(01)	33.69 <sup>c</sup>
Florida	5.11 <sup>b</sup>	0.10	(47)	-12.51	Ohio	6.57 <sup>b</sup>	0.16	(08)	12.60
Georgia	5.61	0.19	(31)	-3.92	Oklahoma	7.20 <sup>b</sup>	0.29	(03)	23.36 <sup>c</sup>
Hawaii	4.11 <sup>b</sup>	0.36	(51)	-29.63 <sup>c</sup>	Oregon	5.49	0.25	(37)	-5.99
Idaho	5.25	0.40	(43)	-10.01	Pennsylvania	5.88	0.14	(26)	0.81
Illinois	6.22	0.15	(20)	6.54	Rhode Island	5.37	0.46	(39)	-8.05
Indiana	6.44	0.22	(14)	10.32	South Carolina	6.45	0.26	(13)	10.56
Iowa	7.24 <sup>b</sup>	0.31	(02)	24.04 <sup>c</sup>	South Dakota	5.99	0.54	(23)	2.59
Kansas	6.84	0.33	(05)	17.26 <sup>c</sup>	Tennessee	6.40	0.22	(15)	9.65
Kentucky	6.69	0.27	(06)	14.58	Texas	6.53 <sup>b</sup>	0.12	(10)	11.81
Louisiana	7.13 <sup>b</sup>	0.28	(04)	22.23 <sup>c</sup>	Utah	4.30 <sup>b</sup>	0.31	(49)	-26.32 <sup>c</sup>
Maine	6.28	0.43	(18)	7.57	Vermont	5.12	0.58	(46)	-12.25
Maryland	5.74	0.22	(29)	-1.61	Virginia	5.99	0.20	(24)	2.54
Massachusetts	5.57	0.20	(33)	-4.53	Washington	5.57	0.20	(34)	-4.61
Michigan	5.82	0.16	(27)	-0.23	West Virginia	6.46	0.37	(12)	10.73
Minnesota	6.39	0.24	(16)	9.53	Wisconsin	6.25	0.22	(19)	7.14
Mississippi	6.57	0.33	(09)	12.58	Wyoming	5.94	0.70	(25)	1.73
Missouri	6.64 <sup>b</sup>	0.22	(07)	13.81					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 11.20  
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.61	0.02		
High Five States				
Louisiana	3.48 <sup>b</sup>	0.17	(01)	33.73 <sup>c</sup>
West Virginia	3.45 <sup>b</sup>	0.23	(02)	32.32 <sup>c</sup>
Oklahoma	3.37 <sup>b</sup>	0.18	(03)	29.54 <sup>c</sup>
North Dakota	3.28	0.41	(04)	26.07 <sup>c</sup>
Iowa	3.23	0.18	(05)	23.89 <sup>c</sup>
Low Five States				
Florida	2.17 <sup>b</sup>	0.06	(47)	-16.77 <sup>c</sup>
New York	2.14 <sup>b</sup>	0.06	(48)	-17.90 <sup>c</sup>
D.C.	1.80	0.33	(49)	-31.07 <sup>c</sup>
Wyoming	1.67	0.34	(50)	-35.90 <sup>c</sup>
Hawaii	1.53 <sup>b</sup>	0.20	(51)	-41.32 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.78	0.14	(20)	6.79	Montana	2.49	0.29	(35)	-4.23
Alaska	2.97	0.54	(14)	14.13	Nebraska	2.99	0.24	(13)	14.61
Arizona	2.44	0.12	(40)	-6.28	Nevada	2.44	0.20	(41)	-6.45
Arkansas	2.74	0.18	(23)	5.16	New Hampshire	2.50	0.25	(34)	-4.06
California	2.30 <sup>b</sup>	0.05	(43)	-11.81	New Jersey	2.25 <sup>b</sup>	0.09	(45)	-13.63
Colorado	2.45	0.14	(38)	-5.89	New Mexico	3.04	0.23	(09)	16.78 <sup>c</sup>
Connecticut	2.28	0.14	(44)	-12.42	New York	2.14 <sup>b</sup>	0.06	(48)	-17.90 <sup>c</sup>
Delaware	2.69	0.32	(26)	3.27	North Carolina	2.46	0.10	(37)	-5.51
D.C.	1.80	0.33	(49)	-31.07 <sup>c</sup>	North Dakota	3.28	0.41	(04)	26.07 <sup>c</sup>
Florida	2.17 <sup>b</sup>	0.06	(47)	-16.77 <sup>c</sup>	Ohio	3.04 <sup>b</sup>	0.09	(11)	16.52 <sup>c</sup>
Georgia	2.47	0.10	(36)	-5.25	Oklahoma	3.37 <sup>b</sup>	0.18	(03)	29.54 <sup>c</sup>
Hawaii	1.53 <sup>b</sup>	0.20	(51)	-41.32 <sup>c</sup>	Oregon	2.37	0.15	(42)	-8.88
Idaho	2.71	0.26	(25)	3.97	Pennsylvania	2.65	0.08	(30)	1.79
Illinois	2.75	0.09	(22)	5.39	Rhode Island	2.84	0.29	(18)	8.97
Indiana	3.08 <sup>b</sup>	0.13	(07)	18.17 <sup>c</sup>	South Carolina	2.67	0.14	(28)	2.57
Iowa	3.23	0.18	(05)	23.89 <sup>c</sup>	South Dakota	3.18	0.36	(06)	21.95 <sup>c</sup>
Kansas	2.97	0.19	(15)	14.06	Tennessee	2.68	0.12	(27)	2.79
Kentucky	3.04	0.15	(10)	16.66 <sup>c</sup>	Texas	3.02 <sup>b</sup>	0.07	(12)	15.78 <sup>c</sup>
Louisiana	3.48 <sup>b</sup>	0.17	(01)	33.73 <sup>c</sup>	Utah	2.17	0.20	(46)	-16.54 <sup>c</sup>
Maine	3.04	0.26	(08)	16.87 <sup>c</sup>	Vermont	2.72	0.38	(24)	4.35
Maryland	2.55	0.13	(32)	-2.11	Virginia	2.51	0.11	(33)	-3.51
Massachusetts	2.45	0.11	(39)	-6.01	Washington	2.64	0.12	(31)	1.31
Michigan	2.96 <sup>b</sup>	0.10	(16)	13.69	West Virginia	3.45 <sup>b</sup>	0.23	(02)	32.32 <sup>c</sup>
Minnesota	2.76	0.14	(21)	5.89	Wisconsin	2.79	0.13	(19)	6.92
Mississippi	2.89	0.18	(17)	10.94	Wyoming	1.67	0.34	(50)	-35.90 <sup>c</sup>
Missouri	2.67	0.12	(29)	2.53					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.18  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.17	0.01		
High Five States				
West Virginia	1.86 <sup>b</sup>	0.13	(01)	58.78 <sup>c</sup>
D.C.	1.73	0.24	(02)	47.77 <sup>c</sup>
Louisiana	1.57 <sup>b</sup>	0.08	(03)	33.89 <sup>c</sup>
Kentucky	1.55 <sup>b</sup>	0.08	(04)	32.11 <sup>c</sup>
Tennessee	1.47 <sup>b</sup>	0.07	(05)	25.55 <sup>c</sup>
Low Five States				
South Dakota	0.76	0.13	(47)	-34.68 <sup>c</sup>
Oregon	0.76 <sup>b</sup>	0.06	(48)	-35.43 <sup>c</sup>
Wyoming	0.73	0.16	(49)	-37.89 <sup>c</sup>
Colorado	0.71 <sup>b</sup>	0.06	(50)	-39.65 <sup>c</sup>
Utah	0.32 <sup>b</sup>	0.06	(51)	-72.33 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.47 <sup>b</sup>	0.08	(06)	25.54 <sup>c</sup>	Montana	0.90	0.13	(41)	-23.21 <sup>c</sup>
Alaska	0.95	0.22	(36)	-19.21 <sup>c</sup>	Nebraska	1.02	0.10	(32)	-12.58
Arizona	0.87 <sup>b</sup>	0.05	(43)	-25.48 <sup>c</sup>	Nevada	1.25	0.10	(19)	7.10
Arkansas	1.21	0.09	(22)	3.81	New Hampshire	1.17	0.13	(25)	-0.43
California	0.91 <sup>b</sup>	0.02	(39)	-22.60 <sup>c</sup>	New Jersey	1.26	0.05	(17)	7.65
Colorado	0.71 <sup>b</sup>	0.06	(50)	-39.65 <sup>c</sup>	New Mexico	0.92	0.10	(38)	-21.33 <sup>c</sup>
Connecticut	1.04	0.07	(31)	-11.06	New York	1.21	0.03	(23)	3.45
Delaware	1.35	0.17	(10)	15.65 <sup>c</sup>	North Carolina	1.33	0.05	(14)	13.87
D.C.	1.73	0.24	(02)	47.77 <sup>c</sup>	North Dakota	0.81	0.15	(46)	-30.37 <sup>c</sup>
Florida	1.35 <sup>b</sup>	0.03	(11)	15.47 <sup>c</sup>	Ohio	1.42 <sup>b</sup>	0.05	(08)	20.96 <sup>c</sup>
Georgia	1.07	0.05	(30)	-8.17	Oklahoma	1.13	0.08	(27)	-3.19
Hawaii	0.97	0.11	(35)	-17.48 <sup>c</sup>	Oregon	0.76 <sup>b</sup>	0.06	(48)	-35.43 <sup>c</sup>
Idaho	0.82	0.11	(45)	-29.99 <sup>c</sup>	Pennsylvania	1.28	0.04	(16)	9.19
Illinois	1.26	0.04	(18)	7.50	Rhode Island	1.22	0.14	(21)	4.15
Indiana	1.21	0.06	(24)	3.31	South Carolina	1.32	0.07	(15)	13.19
Iowa	0.97	0.08	(34)	-16.78 <sup>c</sup>	South Dakota	0.76	0.13	(47)	-34.68 <sup>c</sup>
Kansas	0.85 <sup>b</sup>	0.08	(44)	-27.17 <sup>c</sup>	Tennessee	1.47 <sup>b</sup>	0.07	(05)	25.55 <sup>c</sup>
Kentucky	1.55 <sup>b</sup>	0.08	(04)	32.11 <sup>c</sup>	Texas	1.16	0.03	(26)	-0.76
Louisiana	1.57 <sup>b</sup>	0.08	(03)	33.89 <sup>c</sup>	Utah	0.32 <sup>b</sup>	0.06	(51)	-72.33 <sup>c</sup>
Maine	1.34	0.13	(13)	14.36	Vermont	0.88	0.16	(42)	-24.82 <sup>c</sup>
Maryland	1.36	0.07	(09)	16.16 <sup>c</sup>	Virginia	1.11	0.05	(28)	-5.19
Massachusetts	1.10	0.06	(29)	-5.78	Washington	0.90 <sup>b</sup>	0.05	(40)	-22.90 <sup>c</sup>
Michigan	1.35 <sup>b</sup>	0.05	(12)	15.16 <sup>c</sup>	West Virginia	1.86 <sup>b</sup>	0.13	(01)	58.78 <sup>c</sup>
Minnesota	0.94 <sup>b</sup>	0.06	(37)	-19.64 <sup>c</sup>	Wisconsin	1.02	0.06	(33)	-13.14
Mississippi	1.45	0.10	(07)	23.51 <sup>c</sup>	Wyoming	0.73	0.16	(49)	-37.89 <sup>c</sup>
Missouri	1.23	0.06	(20)	5.44					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.19  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	2.10	0.02		
High Five States				
D.C.	3.56	0.54	(01)	69.30 <sup>c</sup>
West Virginia	3.22 <sup>b</sup>	0.25	(02)	53.09 <sup>c</sup>
Mississippi	2.90 <sup>b</sup>	0.22	(03)	38.20 <sup>c</sup>
Louisiana	2.75 <sup>b</sup>	0.17	(04)	30.91 <sup>c</sup>
Kentucky	2.72 <sup>b</sup>	0.17	(05)	29.28 <sup>c</sup>
Low Five States				
Idaho	1.33 <sup>b</sup>	0.20	(46)	-36.65 <sup>c</sup>
South Dakota	1.33	0.26	(47)	-36.66 <sup>c</sup>
Oregon	1.19 <sup>b</sup>	0.12	(48)	-43.28 <sup>c</sup>
Colorado	1.12 <sup>b</sup>	0.11	(49)	-46.69 <sup>c</sup>
Utah	0.55 <sup>b</sup>	0.11	(50)	-73.94 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.67 <sup>b</sup>	0.16	(07)	27.02 <sup>c</sup>	Montana	1.58	0.25	(41)	-24.77 <sup>c</sup>
Alaska	1.73	0.46	(35)	-17.82 <sup>c</sup>	Nebraska	1.84	0.21	(34)	-12.42
Arizona	1.48 <sup>b</sup>	0.10	(44)	-29.49 <sup>c</sup>	Nevada	1.99	0.19	(28)	-5.25
Arkansas	2.24	0.18	(20)	6.80	New Hampshire	2.15	0.27	(24)	2.17
California	1.67 <sup>b</sup>	0.05	(38)	-20.50 <sup>c</sup>	New Jersey	2.23	0.11	(21)	6.00
Colorado	1.12 <sup>b</sup>	0.11	(49)	-46.69 <sup>c</sup>	New Mexico	1.61	0.19	(40)	-23.24 <sup>c</sup>
Connecticut	1.96	0.16	(31)	-6.76	New York	2.22	0.07	(22)	5.41
Delaware	2.31	0.32	(16)	9.96	North Carolina	2.46	0.11	(10)	16.83 <sup>c</sup>
D.C.	3.56	0.54	(01)	69.30 <sup>c</sup>	North Dakota	1.63	0.31	(39)	-22.59 <sup>c</sup>
Florida	2.36 <sup>b</sup>	0.07	(11)	12.47	Ohio	2.58 <sup>b</sup>	0.10	(08)	22.65 <sup>c</sup>
Georgia	2.02	0.11	(26)	-3.79	Oklahoma	1.97	0.15	(30)	-6.49
Hawaii	1.84	0.24	(33)	-12.35	Oregon	1.19 <sup>b</sup>	0.12	(48)	-43.28 <sup>c</sup>
Idaho	1.33 <sup>b</sup>	0.20	(46)	-36.65 <sup>c</sup>	Pennsylvania	2.35	0.09	(15)	11.79
Illinois	2.30	0.09	(17)	9.50	Rhode Island	2.36	0.31	(14)	12.07
Indiana	2.01	0.12	(27)	-4.42	South Carolina	2.46	0.15	(09)	17.06 <sup>c</sup>
Iowa	1.71	0.15	(36)	-18.62 <sup>c</sup>	South Dakota	1.33	0.26	(47)	-36.66 <sup>c</sup>
Kansas	1.53 <sup>b</sup>	0.15	(42)	-27.26 <sup>c</sup>	Tennessee	2.68 <sup>b</sup>	0.14	(06)	27.48 <sup>c</sup>
Kentucky	2.72 <sup>b</sup>	0.17	(05)	29.28 <sup>c</sup>	Texas	2.17	0.07	(23)	3.21
Louisiana	2.75 <sup>b</sup>	0.17	(04)	30.91 <sup>c</sup>	Utah	0.55 <sup>b</sup>	0.11	(50)	-73.94 <sup>c</sup>
Maine	2.28	0.26	(19)	8.66	Vermont	1.45	0.30	(45)	-31.02 <sup>c</sup>
Maryland	2.36	0.14	(13)	12.35	Virginia	2.08	0.12	(25)	-0.93
Massachusetts	1.99	0.12	(29)	-5.36	Washington	1.50 <sup>b</sup>	0.10	(43)	-28.43 <sup>c</sup>
Michigan	2.36	0.10	(12)	12.45	West Virginia	3.22 <sup>b</sup>	0.25	(02)	53.09 <sup>c</sup>
Minnesota	1.67 <sup>b</sup>	0.12	(37)	-20.45 <sup>c</sup>	Wisconsin	1.87	0.12	(32)	-11.15
Mississippi	2.90 <sup>b</sup>	0.22	(03)	38.20 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	2.29	0.13	(18)	8.78					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.20  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.45	0.01		
High Five States				
West Virginia	0.76	0.11	(01)	68.29 <sup>c</sup>
Kentucky	0.68	0.07	(02)	50.62 <sup>c</sup>
Louisiana	0.68	0.07	(03)	50.61 <sup>c</sup>
Maine	0.63	0.12	(04)	38.90 <sup>c</sup>
Nevada	0.62	0.10	(05)	38.13 <sup>c</sup>
Low Five States				
Wisconsin	0.36	0.05	(35)	-19.55 <sup>c</sup>
Mississippi	0.36	0.07	(36)	-21.06 <sup>c</sup>
California	0.33 <sup>b</sup>	0.02	(37)	-27.14 <sup>c</sup>
New Mexico	0.33	0.08	(38)	-27.49 <sup>c</sup>
Kansas	0.32	0.06	(39)	-29.72 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.53	0.06	(11)	17.47 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.37	0.09	(30)	-17.53 <sup>c</sup>
Arizona	0.37	0.05	(34)	-19.04 <sup>c</sup>	Nevada	0.62	0.10	(05)	38.13 <sup>c</sup>
Arkansas	0.40	0.07	(26)	-12.53	New Hampshire	0.42	0.11	(25)	-6.63
California	0.33 <sup>b</sup>	0.02	(37)	-27.14 <sup>c</sup>	New Jersey	0.55	0.05	(09)	21.32 <sup>c</sup>
Colorado	0.37	0.06	(32)	-18.53 <sup>c</sup>	New Mexico	0.33	0.08	(38)	-27.49 <sup>c</sup>
Connecticut	0.39	0.06	(27)	-13.67	New York	0.47	0.03	(18)	3.99
Delaware	-	-	(-)	-	North Carolina	0.51	0.04	(13)	13.12
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.51	0.03	(14)	11.81	Ohio	0.53	0.04	(12)	17.28 <sup>c</sup>
Georgia	0.37	0.04	(31)	-17.97 <sup>c</sup>	Oklahoma	0.48	0.07	(16)	5.84
Hawaii	-	-	(-)	-	Oregon	0.42	0.06	(24)	-6.04
Idaho	-	-	(-)	-	Pennsylvania	0.49	0.03	(15)	7.44
Illinois	0.48	0.04	(17)	5.41	Rhode Island	-	-	(-)	-
Indiana	0.58	0.06	(07)	29.05 <sup>c</sup>	South Carolina	0.45	0.06	(20)	0.03
Iowa	0.45	0.07	(21)	-0.71	South Dakota	-	-	(-)	-
Kansas	0.32	0.06	(39)	-29.72 <sup>c</sup>	Tennessee	0.53	0.05	(10)	17.66 <sup>c</sup>
Kentucky	0.68	0.07	(02)	50.62 <sup>c</sup>	Texas	0.38	0.03	(29)	-15.96 <sup>c</sup>
Louisiana	0.68	0.07	(03)	50.61 <sup>c</sup>	Utah	-	-	(-)	-
Maine	0.63	0.12	(04)	38.90 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.60	0.06	(06)	32.08 <sup>c</sup>	Virginia	0.39	0.04	(28)	-14.66
Massachusetts	0.46	0.05	(19)	2.17	Washington	0.43	0.05	(23)	-5.41
Michigan	0.56	0.04	(08)	23.76 <sup>c</sup>	West Virginia	0.76	0.11	(01)	68.29 <sup>c</sup>
Minnesota	0.37	0.05	(33)	-19.01 <sup>c</sup>	Wisconsin	0.36	0.05	(35)	-19.55 <sup>c</sup>
Mississippi	0.36	0.07	(36)	-21.06 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	0.44	0.05	(22)	-1.94					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.26  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.10	0.02		
High Five States				
Wisconsin	8.06 <sup>b</sup>	0.16	(01)	13.55
North Dakota	7.97	0.46	(02)	12.27
Wyoming	7.96	0.55	(03)	12.22
Montana	7.93	0.39	(04)	11.73
Michigan	7.90 <sup>b</sup>	0.12	(05)	11.29
Low Five States				
Nevada	6.43	0.24	(47)	-9.45
New Mexico	6.34	0.25	(48)	-10.63
D.C.	6.26	0.47	(49)	-11.85
Arizona	6.18 <sup>b</sup>	0.14	(50)	-12.94
Hawaii	5.02 <sup>b</sup>	0.26	(51)	-29.27 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.47	0.17	(19)	5.29	Montana	7.93	0.39	(04)	11.73
Alaska	7.46	0.63	(20)	5.17	Nebraska	7.09	0.27	(29)	-0.09
Arizona	6.18 <sup>b</sup>	0.14	(50)	-12.94	Nevada	6.43	0.24	(47)	-9.45
Arkansas	7.78	0.22	(09)	9.63	New Hampshire	6.79	0.31	(38)	-4.35
California	6.72 <sup>b</sup>	0.06	(44)	-5.34	New Jersey	6.75	0.12	(41)	-4.93
Colorado	6.78	0.18	(39)	-4.42	New Mexico	6.34	0.25	(48)	-10.63
Connecticut	7.00	0.19	(31)	-1.38	New York	6.72 <sup>b</sup>	0.08	(45)	-5.37
Delaware	6.93	0.38	(33)	-2.38	North Carolina	7.06	0.13	(30)	-0.46
D.C.	6.26	0.47	(49)	-11.85	North Dakota	7.97	0.46	(02)	12.27
Florida	6.75 <sup>b</sup>	0.08	(42)	-4.95	Ohio	7.51 <sup>b</sup>	0.11	(17)	5.86
Georgia	6.81	0.13	(37)	-4.05	Oklahoma	7.60	0.20	(13)	7.11
Hawaii	5.02 <sup>b</sup>	0.26	(51)	-29.27 <sup>c</sup>	Oregon	7.27	0.19	(25)	2.50
Idaho	7.35	0.32	(22)	3.51	Pennsylvania	7.52 <sup>b</sup>	0.10	(15)	5.94
Illinois	7.54 <sup>b</sup>	0.11	(14)	6.24	Rhode Island	7.50	0.35	(18)	5.71
Indiana	7.76 <sup>b</sup>	0.15	(10)	9.38	South Carolina	6.87	0.17	(35)	-3.22
Iowa	7.51	0.21	(16)	5.87	South Dakota	6.98	0.39	(32)	-1.61
Kansas	7.64	0.23	(12)	7.67	Tennessee	7.35	0.15	(21)	3.60
Kentucky	7.31	0.18	(24)	2.95	Texas	6.77 <sup>b</sup>	0.08	(40)	-4.59
Louisiana	7.24	0.18	(27)	1.98	Utah	6.73	0.26	(43)	-5.19
Maine	7.32	0.31	(23)	3.11	Vermont	7.10	0.46	(28)	-0.01
Maryland	6.82	0.16	(36)	-3.89	Virginia	6.89	0.14	(34)	-2.94
Massachusetts	6.69	0.14	(46)	-5.70	Washington	7.26	0.15	(26)	2.26
Michigan	7.90 <sup>b</sup>	0.12	(05)	11.29	West Virginia	7.79	0.27	(08)	9.74
Minnesota	7.76 <sup>b</sup>	0.17	(11)	9.31	Wisconsin	8.06 <sup>b</sup>	0.16	(01)	13.55
Mississippi	7.84	0.23	(06)	10.42	Wyoming	7.96	0.55	(03)	12.22
Missouri	7.83 <sup>b</sup>	0.16	(07)	10.33					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
 SE Standard error of the rate.  
 PD Percent difference between state rate and total U.S. rate.  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.27  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.61	0.04		
High Five States				
Wisconsin	11.36 <sup>b</sup>	0.30	(01)	18.24 <sup>c</sup>
North Dakota	11.07	0.82	(02)	15.26 <sup>c</sup>
Missouri	11.02 <sup>b</sup>	0.29	(03)	14.74
Minnesota	10.80 <sup>b</sup>	0.31	(04)	12.41
Indiana	10.77 <sup>b</sup>	0.29	(05)	12.07
Low Five States				
D.C.	8.70	0.89	(47)	-9.43
New Hampshire	8.31	0.54	(48)	-13.48
New Mexico	8.13	0.43	(49)	-15.36 <sup>c</sup>
Arizona	7.97 <sup>b</sup>	0.23	(50)	-17.05 <sup>c</sup>
Hawaii	6.37 <sup>b</sup>	0.44	(51)	-33.74 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.25	0.32	(20)	6.65	Montana	10.76	0.68	(06)	11.99
Alaska	8.94	1.09	(44)	-6.95	Nebraska	9.97	0.50	(25)	3.79
Arizona	7.97 <sup>b</sup>	0.23	(50)	-17.05 <sup>c</sup>	Nevada	8.74	0.42	(46)	-9.06
Arkansas	10.40	0.40	(13)	8.22	New Hampshire	8.31	0.54	(48)	-13.48
California	8.81 <sup>b</sup>	0.11	(45)	-8.36	New Jersey	9.17	0.22	(39)	-4.60
Colorado	8.96	0.33	(42)	-6.71	New Mexico	8.13	0.43	(49)	-15.36 <sup>c</sup>
Connecticut	9.63	0.35	(29)	0.27	New York	9.16	0.15	(40)	-4.65
Delaware	9.53	0.70	(31)	-0.78	North Carolina	9.99	0.24	(24)	3.92
D.C.	8.70	0.89	(47)	-9.43	North Dakota	11.07	0.82	(02)	15.26 <sup>c</sup>
Florida	8.95 <sup>b</sup>	0.13	(43)	-6.82	Ohio	10.10	0.20	(22)	5.14
Georgia	9.46	0.25	(32)	-1.58	Oklahoma	10.51	0.37	(09)	9.34
Hawaii	6.37 <sup>b</sup>	0.44	(51)	-33.74 <sup>c</sup>	Oregon	9.76	0.34	(28)	1.62
Idaho	9.23	0.54	(37)	-3.98	Pennsylvania	10.12	0.18	(21)	5.37
Illinois	10.28	0.20	(19)	6.98	Rhode Island	10.32	0.65	(17)	7.37
Indiana	10.77 <sup>b</sup>	0.29	(05)	12.07	South Carolina	9.57	0.32	(30)	-0.36
Iowa	10.50	0.38	(10)	9.24	South Dakota	9.23	0.68	(36)	-3.96
Kansas	10.30	0.41	(18)	7.16	Tennessee	10.45	0.29	(11)	8.76
Kentucky	9.95	0.34	(26)	3.58	Texas	9.18	0.15	(38)	-4.50
Louisiana	10.02	0.34	(23)	4.25	Utah	9.38	0.47	(34)	-2.43
Maine	10.40	0.57	(12)	8.24	Vermont	10.38	0.86	(14)	8.07
Maryland	9.29	0.30	(35)	-3.27	Virginia	9.43	0.25	(33)	-1.81
Massachusetts	9.12	0.25	(41)	-5.14	Washington	9.95	0.28	(27)	3.57
Michigan	10.71 <sup>b</sup>	0.22	(07)	11.49	West Virginia	10.37	0.48	(15)	7.91
Minnesota	10.80 <sup>b</sup>	0.31	(04)	12.41	Wisconsin	11.36 <sup>b</sup>	0.30	(01)	18.24 <sup>c</sup>
Mississippi	10.62	0.43	(08)	10.57	Wyoming	10.34	0.96	(16)	7.61
Missouri	11.02 <sup>b</sup>	0.29	(03)	14.74					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 13.28  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.32	0.02		
High Five States				
Alaska	6.40	0.79	(01)	20.32 <sup>c</sup>
Wyoming	6.03	0.65	(02)	13.39
West Virginia	6.02	0.32	(03)	13.18
Mississippi	5.98	0.27	(04)	12.48
Arkansas	5.94	0.26	(05)	11.78
Low Five States				
Arizona	4.72 <sup>b</sup>	0.16	(47)	-11.25
D.C.	4.60	0.52	(48)	-13.48
Nevada	4.57	0.27	(49)	-14.08
Vermont	4.53	0.48	(50)	-14.75
Hawaii	3.89 <sup>b</sup>	0.32	(51)	-26.78 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.65	0.20	(18)	6.24	Montana	5.79	0.45	(11)	8.88
Alaska	6.40	0.79	(01)	20.32 <sup>c</sup>	Nebraska	4.94	0.30	(45)	-6.98
Arizona	4.72 <sup>b</sup>	0.16	(47)	-11.25	Nevada	4.57	0.27	(49)	-14.08
Arkansas	5.94	0.26	(05)	11.78	New Hampshire	5.84	0.39	(07)	9.91
California	5.19	0.07	(33)	-2.28	New Jersey	5.12	0.14	(35)	-3.62
Colorado	5.23	0.21	(29)	-1.59	New Mexico	4.95	0.30	(44)	-6.85
Connecticut	5.23	0.22	(30)	-1.64	New York	5.03	0.09	(40)	-5.31
Delaware	5.01	0.44	(41)	-5.69	North Carolina	5.09	0.14	(37)	-4.29
D.C.	4.60	0.52	(48)	-13.48	North Dakota	5.79	0.54	(10)	8.98
Florida	4.98 <sup>b</sup>	0.09	(42)	-6.25	Ohio	5.74	0.13	(13)	7.95
Georgia	5.06	0.15	(38)	-4.74	Oklahoma	5.54	0.23	(21)	4.26
Hawaii	3.89 <sup>b</sup>	0.32	(51)	-26.78 <sup>c</sup>	Oregon	5.41	0.22	(24)	1.83
Idaho	5.79	0.39	(09)	8.99	Pennsylvania	5.73 <sup>b</sup>	0.12	(14)	7.72
Illinois	5.72	0.13	(15)	7.70	Rhode Island	5.65	0.41	(17)	6.35
Indiana	5.78	0.18	(12)	8.77	South Carolina	4.95	0.19	(43)	-6.82
Iowa	5.53	0.24	(22)	4.02	South Dakota	5.24	0.46	(28)	-1.49
Kansas	5.81	0.26	(08)	9.31	Tennessee	5.32	0.17	(26)	0.14
Kentucky	5.44	0.21	(23)	2.31	Texas	5.04	0.09	(39)	-5.11
Louisiana	5.38	0.21	(25)	1.21	Utah	4.73	0.30	(46)	-10.99
Maine	5.21	0.35	(31)	-1.93	Vermont	4.53	0.48	(50)	-14.75
Maryland	5.15	0.18	(34)	-3.12	Virginia	5.11	0.15	(36)	-3.81
Massachusetts	5.21	0.16	(32)	-1.97	Washington	5.25	0.17	(27)	-1.24
Michigan	5.89 <sup>b</sup>	0.14	(06)	10.74	West Virginia	6.02	0.32	(03)	13.18
Minnesota	5.67	0.20	(16)	6.67	Wisconsin	5.63	0.18	(19)	5.85
Mississippi	5.98	0.27	(04)	12.48	Wyoming	6.03	0.65	(02)	13.39
Missouri	5.58	0.18	(20)	5.07					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.18  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.47	0.02		
High Five States				
Hawaii	7.29 <sup>b</sup>	0.32	(01)	33.26 <sup>c</sup>
D.C.	7.28 <sup>b</sup>	0.50	(02)	33.16 <sup>c</sup>
Louisiana	7.22 <sup>b</sup>	0.18	(03)	32.04 <sup>c</sup>
Texas	7.20 <sup>b</sup>	0.08	(04)	31.59 <sup>c</sup>
California	6.77 <sup>b</sup>	0.06	(05)	23.83 <sup>c</sup>
Low Five States				
South Dakota	4.05 <sup>b</sup>	0.30	(47)	-25.91 <sup>c</sup>
Nebraska	3.90 <sup>b</sup>	0.20	(48)	-28.60 <sup>c</sup>
Idaho	3.89 <sup>b</sup>	0.23	(49)	-28.82 <sup>c</sup>
Montana	3.87 <sup>b</sup>	0.26	(50)	-29.28 <sup>c</sup>
Utah	3.51 <sup>b</sup>	0.19	(51)	-35.74 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.49	0.15	(14)	0.41	Montana	3.87 <sup>b</sup>	0.26	(50)	-29.28 <sup>c</sup>
Alaska	5.53	0.50	(13)	1.08	Nebraska	3.90 <sup>b</sup>	0.20	(48)	-28.60 <sup>c</sup>
Arizona	5.06	0.12	(27)	-7.46	Nevada	5.93	0.22	(08)	8.52
Arkansas	5.77	0.19	(09)	5.45	New Hampshire	4.45 <sup>b</sup>	0.25	(43)	-18.63 <sup>c</sup>
California	6.77 <sup>b</sup>	0.06	(05)	23.83 <sup>c</sup>	New Jersey	5.14	0.10	(24)	-5.96
Colorado	4.67 <sup>b</sup>	0.15	(39)	-14.62	New Mexico	6.40 <sup>b</sup>	0.25	(07)	16.98 <sup>c</sup>
Connecticut	4.95	0.16	(32)	-9.57	New York	5.54	0.07	(12)	1.34
Delaware	5.42	0.33	(16)	-0.84	North Carolina	4.95 <sup>b</sup>	0.10	(31)	-9.55
D.C.	7.28 <sup>b</sup>	0.50	(02)	33.16 <sup>c</sup>	North Dakota	4.11 <sup>b</sup>	0.33	(46)	-24.88 <sup>c</sup>
Florida	5.34	0.07	(19)	-2.33	Ohio	4.71 <sup>b</sup>	0.09	(37)	-13.82
Georgia	4.82 <sup>b</sup>	0.11	(33)	-11.85	Oklahoma	5.43	0.17	(15)	-0.65
Hawaii	7.29 <sup>b</sup>	0.32	(01)	33.26 <sup>c</sup>	Oregon	5.08	0.16	(25)	-7.16
Idaho	3.89 <sup>b</sup>	0.23	(49)	-28.82 <sup>c</sup>	Pennsylvania	5.02 <sup>b</sup>	0.08	(28)	-8.16
Illinois	5.21	0.09	(22)	-4.68	Rhode Island	5.35	0.30	(17)	-2.12
Indiana	4.78 <sup>b</sup>	0.12	(36)	-12.58	South Carolina	5.22	0.15	(21)	-4.50
Iowa	4.25 <sup>b</sup>	0.16	(45)	-22.20 <sup>c</sup>	South Dakota	4.05 <sup>b</sup>	0.30	(47)	-25.91 <sup>c</sup>
Kansas	4.47 <sup>b</sup>	0.17	(42)	-18.18 <sup>c</sup>	Tennessee	5.24	0.13	(20)	-4.17
Kentucky	4.96	0.15	(29)	-9.26	Texas	7.20 <sup>b</sup>	0.08	(04)	31.59 <sup>c</sup>
Louisiana	7.22 <sup>b</sup>	0.18	(03)	32.04 <sup>c</sup>	Utah	3.51 <sup>b</sup>	0.19	(51)	-35.74 <sup>c</sup>
Maine	4.52 <sup>b</sup>	0.24	(41)	-17.31 <sup>c</sup>	Vermont	4.36	0.35	(44)	-20.27 <sup>c</sup>
Maryland	5.35	0.14	(18)	-2.21	Virginia	4.95 <sup>b</sup>	0.11	(30)	-9.44
Massachusetts	5.74	0.13	(11)	4.88	Washington	5.75	0.13	(10)	5.15
Michigan	5.06 <sup>b</sup>	0.10	(26)	-7.46	West Virginia	4.55 <sup>b</sup>	0.20	(40)	-16.88 <sup>c</sup>
Minnesota	4.69 <sup>b</sup>	0.13	(38)	-14.26	Wisconsin	4.82 <sup>b</sup>	0.13	(34)	-11.87
Mississippi	6.46 <sup>b</sup>	0.21	(06)	18.12 <sup>c</sup>	Wyoming	4.79	0.42	(35)	-12.40
Missouri	5.16	0.13	(23)	-5.67					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.19  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.12	0.03		
High Five States				
D.C.	11.78 <sup>b</sup>	0.97	(01)	45.10 <sup>c</sup>
Louisiana	11.05 <sup>b</sup>	0.34	(02)	36.07 <sup>c</sup>
Texas	10.67 <sup>b</sup>	0.15	(03)	31.46 <sup>c</sup>
Hawaii	10.58 <sup>b</sup>	0.57	(04)	30.32 <sup>c</sup>
Mississippi	10.06 <sup>b</sup>	0.40	(05)	23.94 <sup>c</sup>
Low Five States				
North Dakota	5.59 <sup>b</sup>	0.58	(47)	-31.11 <sup>c</sup>
Idaho	5.58 <sup>b</sup>	0.41	(48)	-31.31 <sup>c</sup>
South Dakota	5.16 <sup>b</sup>	0.50	(49)	-36.49 <sup>c</sup>
Montana	5.07 <sup>b</sup>	0.45	(50)	-37.57 <sup>c</sup>
Utah	4.88 <sup>b</sup>	0.33	(51)	-39.85 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.38	0.28	(11)	3.20	Montana	5.07 <sup>b</sup>	0.45	(50)	-37.57 <sup>c</sup>
Alaska	8.19	0.92	(16)	0.90	Nebraska	5.82 <sup>b</sup>	0.37	(46)	-28.37 <sup>c</sup>
Arizona	7.16 <sup>b</sup>	0.22	(33)	-11.84	Nevada	8.25	0.38	(15)	-1.56
Arkansas	8.30	0.34	(13)	2.23	New Hampshire	6.29 <sup>b</sup>	0.45	(42)	-22.49 <sup>c</sup>
California	9.75 <sup>b</sup>	0.11	(06)	20.10 <sup>c</sup>	New Jersey	7.72	0.20	(24)	-4.97
Colorado	6.83 <sup>b</sup>	0.27	(39)	-15.89 <sup>c</sup>	New Mexico	9.37	0.45	(07)	15.43 <sup>c</sup>
Connecticut	7.64	0.30	(26)	-5.95	New York	8.49	0.14	(10)	4.53
Delaware	8.78	0.64	(09)	8.09	North Carolina	7.44 <sup>b</sup>	0.19	(28)	-8.35
D.C.	11.78 <sup>b</sup>	0.97	(01)	45.10 <sup>c</sup>	North Dakota	5.59 <sup>b</sup>	0.58	(47)	-31.11 <sup>c</sup>
Florida	8.10	0.12	(18)	-0.19	Ohio	7.13 <sup>b</sup>	0.16	(34)	-12.15
Georgia	7.22 <sup>b</sup>	0.21	(31)	-11.13	Oklahoma	7.96	0.30	(21)	-1.97
Hawaii	10.58 <sup>b</sup>	0.57	(04)	30.32 <sup>c</sup>	Oregon	7.21	0.27	(32)	-11.21
Idaho	5.58 <sup>b</sup>	0.41	(48)	-31.31 <sup>c</sup>	Pennsylvania	7.68	0.15	(25)	-5.41
Illinois	7.73	0.17	(23)	-4.85	Rhode Island	8.33	0.57	(12)	2.60
Indiana	7.01 <sup>b</sup>	0.22	(35)	-12.71	South Carolina	7.97	0.28	(20)	-1.87
Iowa	6.02 <sup>b</sup>	0.28	(44)	-25.87 <sup>c</sup>	South Dakota	5.16 <sup>b</sup>	0.50	(49)	-36.49 <sup>c</sup>
Kansas	6.54 <sup>b</sup>	0.32	(41)	-19.47 <sup>c</sup>	Tennessee	7.97	0.24	(19)	-1.86
Kentucky	7.25	0.27	(30)	-10.70	Texas	10.67 <sup>b</sup>	0.15	(03)	31.46 <sup>c</sup>
Louisiana	11.05 <sup>b</sup>	0.34	(02)	36.07 <sup>c</sup>	Utah	4.88 <sup>b</sup>	0.33	(51)	-39.85 <sup>c</sup>
Maine	6.62	0.44	(40)	-18.45 <sup>c</sup>	Vermont	5.90 <sup>b</sup>	0.61	(45)	-27.34 <sup>c</sup>
Maryland	8.11	0.26	(17)	-0.12	Virginia	7.80	0.22	(22)	-3.98
Massachusetts	9.34 <sup>b</sup>	0.25	(08)	15.05 <sup>c</sup>	Washington	8.28	0.24	(14)	2.01
Michigan	7.34 <sup>b</sup>	0.18	(29)	-9.56	West Virginia	6.23 <sup>b</sup>	0.36	(43)	-23.32 <sup>c</sup>
Minnesota	6.89 <sup>b</sup>	0.24	(37)	-15.18 <sup>c</sup>	Wisconsin	6.86 <sup>b</sup>	0.23	(38)	-15.56 <sup>c</sup>
Mississippi	10.06 <sup>b</sup>	0.40	(05)	23.94 <sup>c</sup>	Wyoming	6.95	0.77	(36)	-14.37
Missouri	7.56	0.23	(27)	-6.93					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.20  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.27	0.02		
High Five States				
Hawaii	4.33	0.33	(01)	32.26 <sup>c</sup>
Texas	4.29 <sup>b</sup>	0.09	(02)	31.11 <sup>c</sup>
California	4.24 <sup>b</sup>	0.07	(03)	29.47 <sup>c</sup>
Louisiana	4.19 <sup>b</sup>	0.18	(04)	27.99 <sup>c</sup>
New Mexico	3.84	0.26	(05)	17.33 <sup>c</sup>
Low Five States				
Kansas	2.68	0.18	(47)	-18.20 <sup>c</sup>
Delaware	2.58	0.31	(48)	-21.12 <sup>c</sup>
Idaho	2.38 <sup>b</sup>	0.25	(49)	-27.43 <sup>c</sup>
Utah	2.33 <sup>b</sup>	0.21	(50)	-28.71 <sup>c</sup>
Nebraska	2.30 <sup>b</sup>	0.21	(51)	-29.69 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.25	0.15	(12)	-0.79	Montana	2.75	0.31	(45)	-15.87 <sup>c</sup>
Alaska	3.10	0.52	(24)	-5.45	Nebraska	2.30 <sup>b</sup>	0.21	(51)	-29.69 <sup>c</sup>
Arizona	3.19	0.13	(17)	-2.46	Nevada	3.75	0.25	(06)	14.63
Arkansas	3.69	0.21	(09)	12.78	New Hampshire	2.83	0.27	(40)	-13.68
California	4.24 <sup>b</sup>	0.07	(03)	29.47 <sup>c</sup>	New Jersey	3.11	0.11	(22)	-5.16
Colorado	2.84	0.16	(39)	-13.41	New Mexico	3.84	0.26	(05)	17.33 <sup>c</sup>
Connecticut	2.81	0.16	(42)	-14.06	New York	3.20	0.07	(14)	-2.15
Delaware	2.58	0.31	(48)	-21.12 <sup>c</sup>	North Carolina	2.92	0.11	(35)	-10.79
D.C.	3.71	0.47	(08)	13.19	North Dakota	2.82	0.37	(41)	-13.83
Florida	2.95 <sup>b</sup>	0.07	(31)	-9.93	Ohio	2.77 <sup>b</sup>	0.09	(43)	-15.27 <sup>c</sup>
Georgia	2.92	0.11	(34)	-10.73	Oklahoma	3.28	0.17	(11)	0.31
Hawaii	4.33	0.33	(01)	32.26 <sup>c</sup>	Oregon	3.18	0.17	(18)	-2.93
Idaho	2.38 <sup>b</sup>	0.25	(49)	-27.43 <sup>c</sup>	Pennsylvania	2.92 <sup>b</sup>	0.08	(36)	-10.81
Illinois	3.23	0.09	(13)	-1.46	Rhode Island	3.11	0.30	(21)	-4.87
Indiana	3.01	0.13	(29)	-8.20	South Carolina	2.94	0.15	(32)	-10.23
Iowa	2.77	0.17	(44)	-15.38 <sup>c</sup>	South Dakota	3.03	0.35	(28)	-7.53
Kansas	2.68	0.18	(47)	-18.20 <sup>c</sup>	Tennessee	3.04	0.13	(27)	-7.08
Kentucky	3.12	0.16	(20)	-4.63	Texas	4.29 <sup>b</sup>	0.09	(02)	31.11 <sup>c</sup>
Louisiana	4.19 <sup>b</sup>	0.18	(04)	27.99 <sup>c</sup>	Utah	2.33 <sup>b</sup>	0.21	(50)	-28.71 <sup>c</sup>
Maine	2.84	0.25	(38)	-13.21	Vermont	3.05	0.41	(26)	-6.97
Maryland	3.14	0.14	(19)	-3.98	Virginia	2.72 <sup>b</sup>	0.11	(46)	-16.85 <sup>c</sup>
Massachusetts	2.92	0.12	(33)	-10.70	Washington	3.52	0.14	(10)	7.36
Michigan	3.20	0.10	(15)	-2.34	West Virginia	3.10	0.22	(23)	-5.40
Minnesota	2.86	0.14	(37)	-12.70	Wisconsin	3.06	0.13	(25)	-6.61
Mississippi	3.74	0.21	(07)	14.18	Wyoming	3.00	0.46	(30)	-8.49
Missouri	3.19	0.13	(16)	-2.44					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.24  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	50.63	0.06		
High Five States				
Kentucky	74.03 <sup>b</sup>	0.57	(01)	46.20 <sup>c</sup>
West Virginia	66.99 <sup>b</sup>	0.77	(02)	32.30 <sup>c</sup>
Arkansas	66.24 <sup>b</sup>	0.65	(03)	30.83 <sup>c</sup>
Tennessee	65.74 <sup>b</sup>	0.45	(04)	29.83 <sup>c</sup>
Mississippi	65.11 <sup>b</sup>	0.66	(05)	28.59 <sup>c</sup>
Low Five States				
California	39.86 <sup>b</sup>	0.16	(47)	-21.28 <sup>c</sup>
Hawaii	37.66 <sup>b</sup>	0.72	(48)	-25.63 <sup>c</sup>
Colorado	37.36 <sup>b</sup>	0.43	(49)	-26.20 <sup>c</sup>
New Mexico	35.88 <sup>b</sup>	0.60	(50)	-29.14 <sup>c</sup>
Utah	21.45 <sup>b</sup>	0.47	(51)	-57.63 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	61.28 <sup>b</sup>	0.49	(08)	21.04 <sup>c</sup>	Montana	48.15	0.94	(33)	-4.91
Alaska	53.27	1.65	(20)	5.20 <sup>c</sup>	Nebraska	47.27 <sup>b</sup>	0.71	(36)	-6.63 <sup>c</sup>
Arizona	40.85 <sup>b</sup>	0.35	(46)	-19.33 <sup>c</sup>	Nevada	55.06 <sup>b</sup>	0.69	(16)	8.75 <sup>c</sup>
Arkansas	66.24 <sup>b</sup>	0.65	(03)	30.83 <sup>c</sup>	New Hampshire	50.77	0.85	(29)	0.27
California	39.86 <sup>b</sup>	0.16	(47)	-21.28 <sup>c</sup>	New Jersey	46.17 <sup>b</sup>	0.32	(39)	-8.82 <sup>c</sup>
Colorado	37.36 <sup>b</sup>	0.43	(49)	-26.20 <sup>c</sup>	New Mexico	35.88 <sup>b</sup>	0.60	(50)	-29.14 <sup>c</sup>
Connecticut	45.66 <sup>b</sup>	0.49	(40)	-9.83 <sup>c</sup>	New York	43.69 <sup>b</sup>	0.20	(44)	-13.71 <sup>c</sup>
Delaware	57.32 <sup>b</sup>	1.09	(13)	13.21 <sup>c</sup>	North Carolina	57.27 <sup>b</sup>	0.35	(14)	13.10 <sup>c</sup>
D.C.	46.73	1.28	(38)	-7.71 <sup>c</sup>	North Dakota	43.69 <sup>b</sup>	1.10	(43)	-13.71 <sup>c</sup>
Florida	50.16	0.21	(31)	-0.93	Ohio	58.14 <sup>b</sup>	0.30	(11)	14.82 <sup>c</sup>
Georgia	53.90 <sup>b</sup>	0.37	(18)	6.45 <sup>c</sup>	Oklahoma	62.13 <sup>b</sup>	0.57	(06)	22.71 <sup>c</sup>
Hawaii	37.66 <sup>b</sup>	0.72	(48)	-25.63 <sup>c</sup>	Oregon	51.14	0.51	(28)	1.01
Idaho	42.53 <sup>b</sup>	0.77	(45)	-16.00 <sup>c</sup>	Pennsylvania	51.71 <sup>b</sup>	0.26	(24)	2.13
Illinois	52.43 <sup>b</sup>	0.29	(22)	3.55	Rhode Island	52.03	0.93	(23)	2.77
Indiana	61.63 <sup>b</sup>	0.44	(07)	21.72 <sup>c</sup>	South Carolina	56.81 <sup>b</sup>	0.49	(15)	12.20 <sup>c</sup>
Iowa	51.20	0.55	(27)	1.13	South Dakota	46.93 <sup>b</sup>	1.02	(37)	-7.32 <sup>c</sup>
Kansas	53.50 <sup>b</sup>	0.61	(19)	5.66 <sup>c</sup>	Tennessee	65.74 <sup>b</sup>	0.45	(04)	29.83 <sup>c</sup>
Kentucky	74.03 <sup>b</sup>	0.57	(01)	46.20 <sup>c</sup>	Texas	47.66 <sup>b</sup>	0.22	(34)	-5.88 <sup>c</sup>
Louisiana	60.99 <sup>b</sup>	0.53	(09)	20.45 <sup>c</sup>	Utah	21.45 <sup>b</sup>	0.47	(51)	-57.63 <sup>c</sup>
Maine	57.55 <sup>b</sup>	0.85	(12)	13.65 <sup>c</sup>	Vermont	51.55	1.21	(25)	1.82
Maryland	51.53	0.43	(26)	1.77	Virginia	53.10 <sup>b</sup>	0.38	(21)	4.88
Massachusetts	50.47	0.38	(30)	-0.32	Washington	49.25	0.40	(32)	-2.73
Michigan	54.90 <sup>b</sup>	0.32	(17)	8.43 <sup>c</sup>	West Virginia	66.99 <sup>b</sup>	0.77	(02)	32.30 <sup>c</sup>
Minnesota	44.68 <sup>b</sup>	0.41	(42)	-11.77 <sup>c</sup>	Wisconsin	47.61 <sup>b</sup>	0.40	(35)	-5.96 <sup>c</sup>
Mississippi	65.11 <sup>b</sup>	0.66	(05)	28.59 <sup>c</sup>	Wyoming	44.79 <sup>b</sup>	1.31	(41)	-11.55 <sup>c</sup>
Missouri	60.16 <sup>b</sup>	0.43	(10)	18.82 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.25  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	65.65	0.10		
High Five States				
Kentucky	99.67 <sup>b</sup>	1.03	(01)	51.81 <sup>c</sup>
Mississippi	97.29 <sup>b</sup>	1.27	(02)	48.19 <sup>c</sup>
Arkansas	92.54 <sup>b</sup>	1.16	(03)	40.96 <sup>c</sup>
Tennessee	91.49 <sup>b</sup>	0.82	(04)	39.35 <sup>c</sup>
Alabama	89.39 <sup>b</sup>	0.92	(05)	36.16 <sup>c</sup>
Low Five States				
Arizona	50.25 <sup>b</sup>	0.58	(47)	-23.46 <sup>c</sup>
California	49.18 <sup>b</sup>	0.27	(48)	-25.09 <sup>c</sup>
Colorado	45.11 <sup>b</sup>	0.72	(49)	-31.29 <sup>c</sup>
New Mexico	44.42 <sup>b</sup>	1.00	(50)	-32.34 <sup>c</sup>
Utah	28.06 <sup>b</sup>	0.81	(51)	-57.26 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	89.39 <sup>b</sup>	0.92	(05)	36.16 <sup>c</sup>	Montana	57.09 <sup>b</sup>	1.53	(39)	-13.04 <sup>c</sup>
Alaska	62.89	2.72	(28)	-4.21	Nebraska	62.42	1.24	(31)	-4.92
Arizona	50.25 <sup>b</sup>	0.58	(47)	-23.46 <sup>c</sup>	Nevada	62.53	1.09	(30)	-4.75
Arkansas	92.54 <sup>b</sup>	1.16	(03)	40.96 <sup>c</sup>	New Hampshire	61.97	1.44	(33)	-5.60 <sup>c</sup>
California	49.18 <sup>b</sup>	0.27	(48)	-25.09 <sup>c</sup>	New Jersey	57.88 <sup>b</sup>	0.55	(38)	-11.84 <sup>c</sup>
Colorado	45.11 <sup>b</sup>	0.72	(49)	-31.29 <sup>c</sup>	New Mexico	44.42 <sup>b</sup>	1.00	(50)	-32.34 <sup>c</sup>
Connecticut	55.88 <sup>b</sup>	0.84	(41)	-14.88 <sup>c</sup>	New York	55.25 <sup>b</sup>	0.36	(42)	-15.85 <sup>c</sup>
Delaware	69.24	1.83	(19)	5.47 <sup>c</sup>	North Carolina	79.28 <sup>b</sup>	0.65	(12)	20.76 <sup>c</sup>
D.C.	64.74	2.41	(25)	-1.39	North Dakota	56.55 <sup>b</sup>	1.87	(40)	-13.86 <sup>c</sup>
Florida	63.45 <sup>b</sup>	0.35	(26)	-3.35	Ohio	77.38 <sup>b</sup>	0.54	(13)	17.86 <sup>c</sup>
Georgia	75.81 <sup>b</sup>	0.70	(14)	15.47 <sup>c</sup>	Oklahoma	82.67 <sup>b</sup>	1.00	(08)	25.92 <sup>c</sup>
Hawaii	51.17 <sup>b</sup>	1.26	(46)	-22.06 <sup>c</sup>	Oregon	61.17 <sup>b</sup>	0.84	(35)	-6.83 <sup>c</sup>
Idaho	51.32 <sup>b</sup>	1.27	(45)	-21.83 <sup>c</sup>	Pennsylvania	68.47 <sup>b</sup>	0.46	(20)	4.29
Illinois	67.78 <sup>b</sup>	0.51	(21)	3.25	Rhode Island	66.32	1.64	(23)	1.02
Indiana	82.01 <sup>b</sup>	0.78	(09)	24.92 <sup>c</sup>	South Carolina	79.62 <sup>b</sup>	0.90	(11)	21.28 <sup>c</sup>
Iowa	67.46	0.95	(22)	2.75	South Dakota	62.19	1.76	(32)	-5.28 <sup>c</sup>
Kansas	70.61 <sup>b</sup>	1.06	(16)	7.56 <sup>c</sup>	Tennessee	91.49 <sup>b</sup>	0.82	(04)	39.35 <sup>c</sup>
Kentucky	99.67 <sup>b</sup>	1.03	(01)	51.81 <sup>c</sup>	Texas	63.43 <sup>b</sup>	0.39	(27)	-3.38
Louisiana	84.36 <sup>b</sup>	0.96	(07)	28.50 <sup>c</sup>	Utah	28.06 <sup>b</sup>	0.81	(51)	-57.26 <sup>c</sup>
Maine	73.11 <sup>b</sup>	1.46	(15)	11.36 <sup>c</sup>	Vermont	61.60	2.03	(34)	-6.18 <sup>c</sup>
Maryland	65.58	0.77	(24)	-0.11	Virginia	70.55 <sup>b</sup>	0.68	(17)	7.46 <sup>c</sup>
Massachusetts	62.60 <sup>b</sup>	0.66	(29)	-4.65	Washington	58.07 <sup>b</sup>	0.66	(37)	-11.54 <sup>c</sup>
Michigan	70.34 <sup>b</sup>	0.56	(18)	7.15 <sup>c</sup>	West Virginia	87.53 <sup>b</sup>	1.35	(06)	33.32 <sup>c</sup>
Minnesota	55.20 <sup>b</sup>	0.70	(43)	-15.91 <sup>c</sup>	Wisconsin	59.94 <sup>b</sup>	0.68	(36)	-8.69 <sup>c</sup>
Mississippi	97.29 <sup>b</sup>	1.27	(02)	48.19 <sup>c</sup>	Wyoming	52.80 <sup>b</sup>	2.13	(44)	-19.57 <sup>c</sup>
Missouri	79.80 <sup>b</sup>	0.77	(10)	21.56 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 15.26  
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	39.65	0.07		
High Five States				
Kentucky	55.51 <sup>b</sup>	0.66	(01)	40.01 <sup>c</sup>
West Virginia	51.87 <sup>b</sup>	0.92	(02)	30.83 <sup>c</sup>
Nevada	48.81 <sup>b</sup>	0.89	(03)	23.11 <sup>c</sup>
Delaware	48.48 <sup>b</sup>	1.35	(04)	22.29 <sup>c</sup>
Indiana	47.21 <sup>b</sup>	0.51	(05)	19.09 <sup>c</sup>
Low Five States				
California	33.07 <sup>b</sup>	0.19	(47)	-16.60 <sup>c</sup>
Colorado	31.93 <sup>b</sup>	0.53	(48)	-19.47 <sup>c</sup>
New Mexico	29.12 <sup>b</sup>	0.73	(49)	-26.54 <sup>c</sup>
Hawaii	27.02 <sup>b</sup>	0.83	(50)	-31.84 <sup>c</sup>
Utah	16.10 <sup>b</sup>	0.55	(51)	-59.40 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	41.10	0.54	(26)	3.65	Montana	41.33	1.19	(25)	4.23
Alaska	45.53	2.08	(11)	14.83 <sup>c</sup>	Nebraska	36.04 <sup>b</sup>	0.83	(39)	-9.10 <sup>c</sup>
Arizona	33.20 <sup>b</sup>	0.43	(46)	-16.26 <sup>c</sup>	Nevada	48.81 <sup>b</sup>	0.89	(03)	23.11 <sup>c</sup>
Arkansas	46.34 <sup>b</sup>	0.73	(09)	16.89 <sup>c</sup>	New Hampshire	42.97	1.06	(18)	8.38 <sup>c</sup>
California	33.07 <sup>b</sup>	0.19	(47)	-16.60 <sup>c</sup>	New Jersey	38.28 <sup>b</sup>	0.38	(37)	-3.45
Colorado	31.93 <sup>b</sup>	0.53	(48)	-19.47 <sup>c</sup>	New Mexico	29.12 <sup>b</sup>	0.73	(49)	-26.54 <sup>c</sup>
Connecticut	38.82	0.60	(33)	-2.09	New York	35.82 <sup>b</sup>	0.25	(41)	-9.65 <sup>c</sup>
Delaware	48.48 <sup>b</sup>	1.35	(04)	22.29 <sup>c</sup>	North Carolina	41.62 <sup>b</sup>	0.40	(24)	4.97
D.C.	34.82	1.44	(44)	-12.19 <sup>c</sup>	North Dakota	34.33 <sup>b</sup>	1.34	(45)	-13.41 <sup>c</sup>
Florida	39.33	0.25	(32)	-0.79	Ohio	44.51 <sup>b</sup>	0.35	(12)	12.27 <sup>c</sup>
Georgia	38.74	0.41	(35)	-2.29	Oklahoma	46.87 <sup>b</sup>	0.66	(07)	18.21 <sup>c</sup>
Hawaii	27.02 <sup>b</sup>	0.83	(50)	-31.84 <sup>c</sup>	Oregon	43.59 <sup>b</sup>	0.63	(16)	9.94 <sup>c</sup>
Idaho	35.59 <sup>b</sup>	0.97	(42)	-10.24 <sup>c</sup>	Pennsylvania	40.01	0.31	(29)	0.92
Illinois	41.89 <sup>b</sup>	0.34	(22)	5.65 <sup>c</sup>	Rhode Island	43.00	1.13	(17)	8.45 <sup>c</sup>
Indiana	47.21 <sup>b</sup>	0.51	(05)	19.09 <sup>c</sup>	South Carolina	39.99	0.55	(30)	0.86
Iowa	39.41	0.65	(31)	-0.59	South Dakota	35.48	1.21	(43)	-10.50 <sup>c</sup>
Kansas	41.00	0.72	(27)	3.41	Tennessee	47.18 <sup>b</sup>	0.51	(06)	18.99 <sup>c</sup>
Kentucky	55.51 <sup>b</sup>	0.66	(01)	40.01 <sup>c</sup>	Texas	35.94 <sup>b</sup>	0.25	(40)	-9.34 <sup>c</sup>
Louisiana	44.12 <sup>b</sup>	0.60	(14)	11.28 <sup>c</sup>	Utah	16.10 <sup>b</sup>	0.55	(51)	-59.40 <sup>c</sup>
Maine	46.36 <sup>b</sup>	1.02	(08)	16.94 <sup>c</sup>	Vermont	44.32	1.51	(13)	11.78 <sup>c</sup>
Maryland	41.78 <sup>b</sup>	0.52	(23)	5.39 <sup>c</sup>	Virginia	40.72	0.44	(28)	2.70
Massachusetts	42.45 <sup>b</sup>	0.46	(20)	7.08 <sup>c</sup>	Washington	42.83 <sup>b</sup>	0.50	(19)	8.03 <sup>c</sup>
Michigan	43.87 <sup>b</sup>	0.39	(15)	10.64 <sup>c</sup>	West Virginia	51.87 <sup>b</sup>	0.92	(02)	30.83 <sup>c</sup>
Minnesota	37.23 <sup>b</sup>	0.51	(38)	-6.10 <sup>c</sup>	Wisconsin	38.74	0.49	(34)	-2.29
Mississippi	42.33 <sup>b</sup>	0.71	(21)	6.77 <sup>c</sup>	Wyoming	38.48	1.65	(36)	-2.95
Missouri	45.95 <sup>b</sup>	0.51	(10)	15.90 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.18  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.73	0.01		
High Five States				
Oklahoma	3.48 <sup>b</sup>	0.14	(01)	27.26 <sup>c</sup>
Utah	3.41 <sup>b</sup>	0.19	(02)	24.93 <sup>c</sup>
Kentucky	3.34 <sup>b</sup>	0.12	(03)	22.18 <sup>c</sup>
Colorado	3.33 <sup>b</sup>	0.12	(04)	21.97 <sup>c</sup>
Vermont	3.33	0.31	(05)	21.95 <sup>c</sup>
Low Five States				
New York	2.17 <sup>b</sup>	0.05	(47)	-20.43 <sup>c</sup>
Alaska	2.16	0.30	(48)	-21.10 <sup>c</sup>
North Dakota	2.10	0.24	(49)	-23.22 <sup>c</sup>
Hawaii	1.53 <sup>b</sup>	0.15	(50)	-44.02 <sup>c</sup>
D.C.	1.23 <sup>b</sup>	0.20	(51)	-54.79 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.09	0.11	(12)	13.15	Montana	2.84	0.23	(27)	4.00
Alaska	2.16	0.30	(48)	-21.10 <sup>c</sup>	Nebraska	3.07	0.18	(14)	12.37
Arizona	2.87	0.09	(25)	5.05	Nevada	2.85	0.16	(26)	4.29
Arkansas	2.61	0.13	(36)	-4.30	New Hampshire	2.91	0.20	(22)	6.37
California	2.58 <sup>b</sup>	0.04	(37)	-5.59	New Jersey	2.74	0.08	(30)	0.31
Colorado	3.33 <sup>b</sup>	0.12	(04)	21.97 <sup>c</sup>	New Mexico	2.73	0.17	(31)	-0.21
Connecticut	2.65	0.12	(35)	-3.12	New York	2.17 <sup>b</sup>	0.05	(47)	-20.43 <sup>c</sup>
Delaware	2.88	0.25	(24)	5.40	North Carolina	3.09 <sup>b</sup>	0.08	(13)	13.01
D.C.	1.23 <sup>b</sup>	0.20	(51)	-54.79 <sup>c</sup>	North Dakota	2.10	0.24	(49)	-23.22 <sup>c</sup>
Florida	2.93 <sup>b</sup>	0.05	(21)	7.36	Ohio	2.80	0.07	(29)	2.44
Georgia	2.42 <sup>b</sup>	0.08	(43)	-11.37	Oklahoma	3.48 <sup>b</sup>	0.14	(01)	27.26 <sup>c</sup>
Hawaii	1.53 <sup>b</sup>	0.15	(50)	-44.02 <sup>c</sup>	Oregon	3.12	0.12	(11)	14.30
Idaho	3.22	0.21	(08)	17.73 <sup>c</sup>	Pennsylvania	2.90	0.06	(23)	6.04
Illinois	2.56	0.06	(38)	-6.27	Rhode Island	2.69	0.21	(33)	-1.44
Indiana	3.01	0.10	(16)	10.34	South Carolina	2.55	0.10	(39)	-6.48
Iowa	2.72	0.13	(32)	-0.43	South Dakota	2.53	0.24	(40)	-7.37
Kansas	3.12	0.15	(10)	14.36	Tennessee	2.97	0.10	(19)	8.58
Kentucky	3.34 <sup>b</sup>	0.12	(03)	22.18 <sup>c</sup>	Texas	2.48 <sup>b</sup>	0.05	(41)	-9.25
Louisiana	2.21 <sup>b</sup>	0.10	(46)	-19.04 <sup>c</sup>	Utah	3.41 <sup>b</sup>	0.19	(02)	24.93 <sup>c</sup>
Maine	2.97	0.20	(20)	8.55	Vermont	3.33	0.31	(05)	21.95 <sup>c</sup>
Maryland	2.80	0.10	(28)	2.68	Virginia	2.98	0.09	(17)	9.09
Massachusetts	3.02	0.09	(15)	10.64	Washington	2.97	0.10	(18)	8.61
Michigan	2.42 <sup>b</sup>	0.07	(42)	-11.28	West Virginia	3.26	0.18	(07)	19.35 <sup>c</sup>
Minnesota	2.29 <sup>b</sup>	0.09	(45)	-16.10 <sup>c</sup>	Wisconsin	2.65	0.09	(34)	-2.83
Mississippi	2.37	0.13	(44)	-13.29	Wyoming	3.27	0.35	(06)	19.78 <sup>c</sup>
Missouri	3.15 <sup>b</sup>	0.10	(09)	15.29 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.19  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.08	0.02		
High Five States				
Utah	5.26	0.35	(01)	28.88 <sup>c</sup>
Oklahoma	5.15 <sup>b</sup>	0.25	(02)	26.10 <sup>c</sup>
Vermont	5.01	0.59	(03)	22.54 <sup>c</sup>
Colorado	4.99 <sup>b</sup>	0.23	(04)	22.20 <sup>c</sup>
Wyoming	4.92	0.66	(05)	20.33 <sup>c</sup>
Low Five States				
North Dakota	3.30	0.45	(47)	-19.32 <sup>c</sup>
New York	3.26 <sup>b</sup>	0.09	(48)	-20.22 <sup>c</sup>
Alaska	2.53	0.46	(49)	-38.13 <sup>c</sup>
Hawaii	2.19 <sup>b</sup>	0.26	(50)	-46.49 <sup>c</sup>
D.C.	1.94 <sup>b</sup>	0.42	(51)	-52.41 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.75	0.21	(08)	16.38 <sup>c</sup>	Montana	3.59	0.38	(44)	-12.19
Alaska	2.53	0.46	(49)	-38.13 <sup>c</sup>	Nebraska	4.24	0.32	(27)	3.72
Arizona	4.44	0.17	(18)	8.63	Nevada	4.52	0.29	(15)	10.65
Arkansas	3.87	0.24	(35)	-5.33	New Hampshire	4.27	0.37	(25)	4.47
California	3.90	0.07	(33)	-4.48	New Jersey	4.02	0.14	(32)	-1.65
Colorado	4.99 <sup>b</sup>	0.23	(04)	22.20 <sup>c</sup>	New Mexico	3.89	0.30	(34)	-4.78
Connecticut	4.03	0.22	(31)	-1.44	New York	3.26 <sup>b</sup>	0.09	(48)	-20.22 <sup>c</sup>
Delaware	4.64	0.48	(09)	13.54	North Carolina	4.56	0.16	(13)	11.70
D.C.	1.94 <sup>b</sup>	0.42	(51)	-52.41 <sup>c</sup>	North Dakota	3.30	0.45	(47)	-19.32 <sup>c</sup>
Florida	4.46 <sup>b</sup>	0.09	(17)	9.14	Ohio	4.27	0.13	(24)	4.63
Georgia	3.80	0.16	(39)	-6.94	Oklahoma	5.15 <sup>b</sup>	0.25	(02)	26.10 <sup>c</sup>
Hawaii	2.19 <sup>b</sup>	0.26	(50)	-46.49 <sup>c</sup>	Oregon	4.29	0.22	(23)	5.07
Idaho	4.84	0.38	(07)	18.56 <sup>c</sup>	Pennsylvania	4.31	0.12	(21)	5.56
Illinois	3.80	0.12	(40)	-7.04	Rhode Island	4.18	0.41	(29)	2.35
Indiana	4.60	0.18	(10)	12.51	South Carolina	4.10	0.21	(30)	0.34
Iowa	3.86	0.23	(36)	-5.39	South Dakota	3.63	0.43	(42)	-11.07
Kansas	4.50	0.27	(16)	10.28	Tennessee	4.59	0.19	(11)	12.43
Kentucky	4.91 <sup>b</sup>	0.23	(06)	20.13 <sup>c</sup>	Texas	3.72 <sup>b</sup>	0.09	(41)	-9.03
Louisiana	3.37 <sup>b</sup>	0.20	(45)	-17.46 <sup>c</sup>	Utah	5.26	0.35	(01)	28.88 <sup>c</sup>
Maine	4.26	0.36	(26)	4.39	Vermont	5.01	0.59	(03)	22.54 <sup>c</sup>
Maryland	4.40	0.20	(19)	7.60	Virginia	4.35	0.17	(20)	6.46
Massachusetts	4.30	0.17	(22)	5.17	Washington	4.21	0.17	(28)	2.98
Michigan	3.61 <sup>b</sup>	0.13	(43)	-11.71	West Virginia	4.59	0.32	(12)	12.41
Minnesota	3.32 <sup>b</sup>	0.17	(46)	-18.84 <sup>c</sup>	Wisconsin	3.85	0.17	(37)	-5.76
Mississippi	3.85	0.26	(38)	-5.86	Wyoming	4.92	0.66	(05)	20.33 <sup>c</sup>
Missouri	4.55	0.18	(14)	11.47					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.20  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.73	0.01		
High Five States				
West Virginia	2.27	0.20	(01)	31.29 <sup>c</sup>
Montana	2.24	0.29	(02)	29.56 <sup>c</sup>
Nebraska	2.22	0.21	(03)	28.30 <sup>c</sup>
Oregon	2.20	0.14	(04)	27.64 <sup>c</sup>
Oklahoma	2.20	0.15	(05)	27.55 <sup>c</sup>
Low Five States				
Louisiana	1.40	0.11	(46)	-19.12 <sup>c</sup>
Nevada	1.38	0.15	(47)	-20.03 <sup>c</sup>
Mississippi	1.35	0.13	(48)	-21.64 <sup>c</sup>
North Dakota	1.08	0.23	(49)	-37.26 <sup>c</sup>
Hawaii	0.95 <sup>b</sup>	0.16	(50)	-45.10 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.89	0.12	(18)	9.31	Montana	2.24	0.29	(02)	29.56 <sup>c</sup>
Alaska	1.73	0.38	(29)	0.28	Nebraska	2.22	0.21	(03)	28.30 <sup>c</sup>
Arizona	1.56	0.09	(39)	-9.95	Nevada	1.38	0.15	(47)	-20.03 <sup>c</sup>
Arkansas	1.62	0.14	(36)	-6.06	New Hampshire	1.86	0.22	(22)	7.44
California	1.58 <sup>b</sup>	0.04	(38)	-8.70	New Jersey	1.82	0.08	(23)	5.33
Colorado	2.05	0.13	(12)	18.80 <sup>c</sup>	New Mexico	1.79	0.18	(26)	3.69
Connecticut	1.70	0.13	(33)	-1.40	New York	1.42 <sup>b</sup>	0.05	(45)	-17.88 <sup>c</sup>
Delaware	1.59	0.25	(37)	-7.70	North Carolina	2.08 <sup>b</sup>	0.09	(11)	20.16 <sup>c</sup>
D.C.	-	-	(-)	-	North Dakota	1.08	0.23	(49)	-37.26 <sup>c</sup>
Florida	1.70	0.05	(32)	-1.34	Ohio	1.75	0.07	(28)	1.20
Georgia	1.48	0.08	(43)	-14.23	Oklahoma	2.20	0.15	(05)	27.55 <sup>c</sup>
Hawaii	0.95 <sup>b</sup>	0.16	(50)	-45.10 <sup>c</sup>	Oregon	2.20	0.14	(04)	27.64 <sup>c</sup>
Idaho	1.79	0.22	(25)	3.81	Pennsylvania	1.87	0.07	(19)	8.29
Illinois	1.69	0.07	(35)	-2.12	Rhode Island	1.77	0.23	(27)	2.65
Indiana	1.87	0.10	(21)	8.26	South Carolina	1.43	0.11	(44)	-17.17 <sup>c</sup>
Iowa	1.87	0.15	(20)	8.28	South Dakota	1.69	0.27	(34)	-1.97
Kansas	2.09	0.16	(09)	21.09 <sup>c</sup>	Tennessee	1.81	0.10	(24)	4.73
Kentucky	2.14	0.13	(07)	23.83 <sup>c</sup>	Texas	1.55	0.05	(41)	-10.28
Louisiana	1.40	0.11	(46)	-19.12 <sup>c</sup>	Utah	1.92	0.19	(17)	11.37
Maine	2.02	0.22	(13)	16.78 <sup>c</sup>	Vermont	2.08	0.33	(10)	20.68 <sup>c</sup>
Maryland	1.72	0.10	(31)	-0.67	Virginia	1.98	0.10	(15)	14.75
Massachusetts	2.14 <sup>b</sup>	0.10	(06)	23.89 <sup>c</sup>	Washington	2.01	0.11	(14)	16.48 <sup>c</sup>
Michigan	1.55	0.07	(40)	-10.09	West Virginia	2.27	0.20	(01)	31.29 <sup>c</sup>
Minnesota	1.53	0.10	(42)	-11.35	Wisconsin	1.72	0.10	(30)	-0.63
Mississippi	1.35	0.13	(48)	-21.64 <sup>c</sup>	Wyoming	1.98	0.37	(16)	14.49
Missouri	2.09	0.11	(08)	21.23 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.18  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.42	0.01		
High Five States				
D.C.	5.70 <sup>b</sup>	0.45	(01)	66.55 <sup>c</sup>
North Carolina	4.04 <sup>b</sup>	0.09	(02)	18.02 <sup>c</sup>
Alabama	4.00 <sup>b</sup>	0.13	(03)	16.90 <sup>c</sup>
Virginia	3.98 <sup>b</sup>	0.10	(04)	16.27 <sup>c</sup>
Maryland	3.96 <sup>b</sup>	0.12	(05)	15.60 <sup>c</sup>
Low Five States				
Florida	2.95 <sup>b</sup>	0.05	(47)	-13.85
Alaska	2.95	0.40	(48)	-13.88
Rhode Island	2.85	0.22	(49)	-16.73 <sup>c</sup>
Nevada	2.58 <sup>b</sup>	0.15	(50)	-24.70 <sup>c</sup>
Hawaii	2.47 <sup>b</sup>	0.18	(51)	-27.73 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.00 <sup>b</sup>	0.13	(03)	16.90 <sup>c</sup>	Montana	3.24	0.24	(38)	-5.39
Alaska	2.95	0.40	(48)	-13.88	Nebraska	3.23	0.18	(39)	-5.58
Arizona	2.95 <sup>b</sup>	0.09	(46)	-13.74	Nevada	2.58 <sup>b</sup>	0.15	(50)	-24.70 <sup>c</sup>
Arkansas	3.46	0.15	(25)	1.22	New Hampshire	3.57	0.23	(22)	4.22
California	3.20 <sup>b</sup>	0.04	(40)	-6.59	New Jersey	3.41	0.09	(27)	-0.23
Colorado	3.32	0.13	(33)	-3.07	New Mexico	2.97	0.17	(45)	-13.08
Connecticut	3.15	0.13	(42)	-7.94	New York	3.07 <sup>b</sup>	0.05	(44)	-10.20
Delaware	3.60	0.27	(19)	5.22	North Carolina	4.04 <sup>b</sup>	0.09	(02)	18.02 <sup>c</sup>
D.C.	5.70 <sup>b</sup>	0.45	(01)	66.55 <sup>c</sup>	North Dakota	3.70	0.32	(13)	8.26
Florida	2.95 <sup>b</sup>	0.05	(47)	-13.85	Ohio	3.69 <sup>b</sup>	0.08	(14)	7.71
Georgia	3.56	0.10	(23)	4.15	Oklahoma	3.45	0.13	(26)	0.77
Hawaii	2.47 <sup>b</sup>	0.18	(51)	-27.73 <sup>c</sup>	Oregon	3.62	0.13	(17)	5.89
Idaho	3.30	0.21	(36)	-3.55	Pennsylvania	3.36	0.07	(30)	-1.93
Illinois	3.57	0.08	(21)	4.27	Rhode Island	2.85	0.22	(49)	-16.73 <sup>c</sup>
Indiana	3.57	0.10	(20)	4.30	South Carolina	3.87	0.13	(07)	13.07
Iowa	3.31	0.14	(34)	-3.42	South Dakota	3.19	0.27	(41)	-6.72
Kansas	3.81	0.16	(09)	11.25	Tennessee	3.90 <sup>b</sup>	0.11	(06)	14.05
Kentucky	3.51	0.13	(24)	2.70	Texas	3.30	0.06	(35)	-3.49
Louisiana	3.81	0.13	(08)	11.45	Utah	3.39	0.19	(29)	-1.01
Maine	3.09	0.20	(43)	-9.76	Vermont	3.32	0.31	(32)	-2.91
Maryland	3.96 <sup>b</sup>	0.12	(05)	15.60 <sup>c</sup>	Virginia	3.98 <sup>b</sup>	0.10	(04)	16.27 <sup>c</sup>
Massachusetts	3.29	0.10	(37)	-3.91	Washington	3.33	0.10	(31)	-2.63
Michigan	3.80 <sup>b</sup>	0.08	(10)	10.92	West Virginia	3.61	0.18	(18)	5.46
Minnesota	3.76	0.12	(11)	9.79	Wisconsin	3.67	0.11	(15)	7.19
Mississippi	3.71	0.16	(12)	8.53	Wyoming	3.41	0.36	(28)	-0.40
Missouri	3.66	0.11	(16)	6.90					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.19  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.37	0.03		
High Five States				
D.C.	7.63 <sup>b</sup>	0.85	(01)	74.48 <sup>c</sup>
Tennessee	5.23 <sup>b</sup>	0.20	(02)	19.65 <sup>c</sup>
Maryland	5.07	0.22	(03)	16.09 <sup>c</sup>
Virginia	5.03 <sup>b</sup>	0.19	(04)	15.13 <sup>c</sup>
Wyoming	5.00	0.65	(05)	14.39
Low Five States				
Florida	3.68 <sup>b</sup>	0.08	(47)	-15.92 <sup>c</sup>
Alaska	3.41	0.64	(48)	-22.00 <sup>c</sup>
Rhode Island	3.34	0.37	(49)	-23.70 <sup>c</sup>
Nevada	3.22 <sup>b</sup>	0.25	(50)	-26.43 <sup>c</sup>
Hawaii	3.19 <sup>b</sup>	0.31	(51)	-27.08 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.99	0.22	(06)	14.06	Montana	3.93	0.40	(42)	-9.99
Alaska	3.41	0.64	(48)	-22.00 <sup>c</sup>	Nebraska	4.13	0.32	(37)	-5.43
Arizona	3.68 <sup>b</sup>	0.16	(46)	-15.72 <sup>c</sup>	Nevada	3.22 <sup>b</sup>	0.25	(50)	-26.43 <sup>c</sup>
Arkansas	4.46	0.26	(25)	1.94	New Hampshire	4.30	0.39	(32)	-1.62
California	4.10 <sup>b</sup>	0.08	(39)	-6.16	New Jersey	4.36	0.15	(29)	-0.33
Colorado	4.30	0.23	(31)	-1.56	New Mexico	3.77	0.29	(45)	-13.84
Connecticut	4.07	0.23	(41)	-6.79	New York	3.91 <sup>b</sup>	0.10	(43)	-10.54
Delaware	4.69	0.49	(20)	7.20	North Carolina	4.98 <sup>b</sup>	0.17	(07)	13.95
D.C.	7.63 <sup>b</sup>	0.85	(01)	74.48 <sup>c</sup>	North Dakota	4.76	0.54	(18)	8.98
Florida	3.68 <sup>b</sup>	0.08	(47)	-15.92 <sup>c</sup>	Ohio	4.61	0.13	(21)	5.38
Georgia	4.45	0.17	(26)	1.86	Oklahoma	4.60	0.24	(22)	5.19
Hawaii	3.19 <sup>b</sup>	0.31	(51)	-27.08 <sup>c</sup>	Oregon	4.83	0.24	(15)	10.44
Idaho	4.12	0.36	(38)	-5.75	Pennsylvania	4.29	0.12	(33)	-1.75
Illinois	4.57	0.13	(23)	4.45	Rhode Island	3.34	0.37	(49)	-23.70 <sup>c</sup>
Indiana	4.86	0.19	(13)	11.16	South Carolina	4.93	0.23	(10)	12.78
Iowa	4.32	0.24	(30)	-1.27	South Dakota	4.14	0.46	(36)	-5.18
Kansas	4.97	0.28	(08)	13.71	Tennessee	5.23 <sup>b</sup>	0.20	(02)	19.65 <sup>c</sup>
Kentucky	4.88	0.24	(12)	11.74	Texas	4.18	0.10	(35)	-4.32
Louisiana	4.86	0.24	(14)	11.14	Utah	4.21	0.32	(34)	-3.65
Maine	4.08	0.35	(40)	-6.69	Vermont	3.88	0.52	(44)	-11.24
Maryland	5.07	0.22	(03)	16.09 <sup>c</sup>	Virginia	5.03 <sup>b</sup>	0.19	(04)	15.13 <sup>c</sup>
Massachusetts	4.38	0.18	(27)	0.25	Washington	4.53	0.19	(24)	3.69
Michigan	4.91 <sup>b</sup>	0.15	(11)	12.42	West Virginia	4.95	0.33	(09)	13.31
Minnesota	4.74	0.21	(19)	8.49	Wisconsin	4.79	0.20	(17)	9.62
Mississippi	4.38	0.27	(28)	0.24	Wyoming	5.00	0.65	(05)	14.39
Missouri	4.80	0.19	(16)	9.92					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 18.20  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.75	0.02		
High Five States				
D.C.	4.57	0.52	(01)	66.23 <sup>c</sup>
North Carolina	3.39 <sup>b</sup>	0.11	(02)	23.55 <sup>c</sup>
Virginia	3.29 <sup>b</sup>	0.12	(03)	19.64 <sup>c</sup>
Alabama	3.26	0.15	(04)	18.69 <sup>c</sup>
Mississippi	3.23	0.19	(05)	17.71 <sup>c</sup>
Low Five States				
Florida	2.38 <sup>b</sup>	0.06	(47)	-13.51
New Mexico	2.37	0.21	(48)	-13.86
Nevada	2.05 <sup>b</sup>	0.18	(49)	-25.45 <sup>c</sup>
Wyoming	1.99	0.38	(50)	-27.68 <sup>c</sup>
Hawaii	1.92 <sup>b</sup>	0.22	(51)	-29.97 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.26	0.15	(04)	18.69 <sup>c</sup>	Montana	2.67	0.30	(29)	-2.76
Alaska	2.51	0.49	(40)	-8.55	Nebraska	2.58	0.22	(36)	-6.17
Arizona	2.38	0.12	(46)	-13.33	Nevada	2.05 <sup>b</sup>	0.18	(49)	-25.45 <sup>c</sup>
Arkansas	2.70	0.18	(28)	-1.75	New Hampshire	3.07	0.28	(09)	11.92
California	2.54 <sup>b</sup>	0.05	(38)	-7.41	New Jersey	2.75	0.10	(22)	0.26
Colorado	2.60	0.15	(32)	-5.40	New Mexico	2.37	0.21	(48)	-13.86
Connecticut	2.51	0.15	(41)	-8.58	New York	2.50 <sup>b</sup>	0.06	(42)	-9.03
Delaware	2.84	0.32	(21)	3.36	North Carolina	3.39 <sup>b</sup>	0.11	(02)	23.55 <sup>c</sup>
D.C.	4.57	0.52	(01)	66.23 <sup>c</sup>	North Dakota	2.94	0.38	(16)	7.20
Florida	2.38 <sup>b</sup>	0.06	(47)	-13.51	Ohio	3.07 <sup>b</sup>	0.09	(10)	11.83
Georgia	3.00	0.11	(13)	9.25	Oklahoma	2.60	0.15	(34)	-5.52
Hawaii	1.92 <sup>b</sup>	0.22	(51)	-29.97 <sup>c</sup>	Oregon	2.72	0.16	(25)	-0.89
Idaho	2.71	0.26	(26)	-1.28	Pennsylvania	2.73	0.08	(24)	-0.64
Illinois	2.92	0.09	(17)	6.22	Rhode Island	2.60	0.28	(33)	-5.48
Indiana	2.70	0.12	(27)	-1.60	South Carolina	3.14	0.15	(07)	14.42
Iowa	2.55	0.16	(37)	-7.08	South Dakota	2.43	0.31	(44)	-11.64
Kansas	3.01	0.19	(11)	9.51	Tennessee	2.99	0.13	(14)	8.70
Kentucky	2.59	0.14	(35)	-5.87	Texas	2.65	0.07	(30)	-3.41
Louisiana	3.14	0.16	(08)	14.28	Utah	2.74	0.23	(23)	-0.40
Maine	2.41	0.23	(45)	-12.29	Vermont	2.89	0.38	(19)	5.36
Maryland	3.23	0.14	(06)	17.54 <sup>c</sup>	Virginia	3.29 <sup>b</sup>	0.12	(03)	19.64 <sup>c</sup>
Massachusetts	2.54	0.11	(39)	-7.72	Washington	2.44	0.12	(43)	-11.03
Michigan	2.98	0.10	(15)	8.54	West Virginia	2.64	0.20	(31)	-3.96
Minnesota	3.01	0.14	(12)	9.41	Wisconsin	2.90	0.13	(18)	5.60
Mississippi	3.23	0.19	(05)	17.71 <sup>c</sup>	Wyoming	1.99	0.38	(50)	-27.68 <sup>c</sup>
Missouri	2.84	0.12	(20)	3.54					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.22  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	6.55	0.02		
High Five States				
West Virginia	7.60 <sup>b</sup>	0.26	(01)	15.95 <sup>c</sup>
Michigan	7.35 <sup>b</sup>	0.12	(02)	12.10
Kentucky	7.30 <sup>b</sup>	0.18	(03)	11.45
Indiana	7.30 <sup>b</sup>	0.15	(04)	11.34
Wisconsin	7.28 <sup>b</sup>	0.15	(05)	11.02
Low Five States				
D.C.	5.91	0.45	(47)	-9.89
Georgia	5.84 <sup>b</sup>	0.12	(48)	-10.84
Nevada	5.73 <sup>b</sup>	0.23	(49)	-12.56
Hawaii	5.59	0.28	(50)	-14.67
New Mexico	5.46 <sup>b</sup>	0.23	(51)	-16.66 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.53	0.16	(26)	-0.31	Montana	6.64	0.35	(21)	1.30
Alaska	6.62	0.60	(22)	1.01	Nebraska	7.09	0.27	(09)	8.14
Arizona	6.00 <sup>b</sup>	0.14	(44)	-8.42	Nevada	5.73 <sup>b</sup>	0.23	(49)	-12.56
Arkansas	6.74	0.21	(20)	2.82	New Hampshire	6.03	0.30	(43)	-7.95
California	6.30 <sup>b</sup>	0.06	(32)	-3.80	New Jersey	6.55	0.12	(25)	-0.05
Colorado	5.96	0.17	(45)	-9.04	New Mexico	5.46 <sup>b</sup>	0.23	(51)	-16.66 <sup>c</sup>
Connecticut	6.39	0.18	(30)	-2.50	New York	6.16 <sup>b</sup>	0.08	(39)	-6.05
Delaware	6.47	0.37	(27)	-1.30	North Carolina	6.11 <sup>b</sup>	0.12	(42)	-6.78
D.C.	5.91	0.45	(47)	-9.89	North Dakota	6.44	0.41	(28)	-1.72
Florida	6.19 <sup>b</sup>	0.07	(38)	-5.50	Ohio	7.15 <sup>b</sup>	0.11	(07)	9.15
Georgia	5.84 <sup>b</sup>	0.12	(48)	-10.84	Oklahoma	7.20	0.19	(06)	9.90
Hawaii	5.59	0.28	(50)	-14.67	Oregon	6.91	0.18	(16)	5.47
Idaho	6.61	0.30	(23)	0.85	Pennsylvania	7.13 <sup>b</sup>	0.10	(08)	8.74
Illinois	6.83	0.10	(18)	4.25	Rhode Island	6.25	0.32	(35)	-4.68
Indiana	7.30 <sup>b</sup>	0.15	(04)	11.34	South Carolina	6.14	0.16	(41)	-6.36
Iowa	7.03	0.20	(13)	7.33	South Dakota	6.29	0.37	(34)	-4.04
Kansas	7.08	0.22	(10)	8.09	Tennessee	7.08 <sup>b</sup>	0.15	(11)	8.04
Kentucky	7.30 <sup>b</sup>	0.18	(03)	11.45	Texas	6.30	0.08	(33)	-3.88
Louisiana	6.78	0.18	(19)	3.38	Utah	5.95	0.25	(46)	-9.22
Maine	7.02	0.30	(14)	7.15	Vermont	6.21	0.42	(37)	-5.28
Maryland	6.14	0.15	(40)	-6.30	Virginia	6.40	0.13	(29)	-2.33
Massachusetts	6.35	0.13	(31)	-3.05	Washington	6.97	0.15	(15)	6.36
Michigan	7.35 <sup>b</sup>	0.12	(02)	12.10	West Virginia	7.60 <sup>b</sup>	0.26	(01)	15.95 <sup>c</sup>
Minnesota	7.04	0.16	(12)	7.41	Wisconsin	7.28 <sup>b</sup>	0.15	(05)	11.02
Mississippi	6.25	0.21	(36)	-4.70	Wyoming	6.84	0.51	(17)	4.35
Missouri	6.59	0.14	(24)	0.58					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.23  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.40	0.04		
High Five States				
Indiana	9.72 <sup>b</sup>	0.27	(01)	15.76 <sup>c</sup>
Kansas	9.63	0.40	(02)	14.74
Minnesota	9.57 <sup>b</sup>	0.30	(03)	14.03
Ohio	9.43 <sup>b</sup>	0.19	(04)	12.31
Wisconsin	9.41 <sup>b</sup>	0.27	(05)	12.10
Low Five States				
Hawaii	7.47	0.48	(47)	-11.04
Arizona	7.47 <sup>b</sup>	0.23	(48)	-11.05
North Dakota	7.44	0.68	(49)	-11.34
New Mexico	6.74 <sup>b</sup>	0.39	(50)	-19.73 <sup>c</sup>
Nevada	6.67 <sup>b</sup>	0.36	(51)	-20.61 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.28	0.29	(27)	-1.34	Montana	8.14	0.58	(28)	-2.99
Alaska	7.89	1.02	(40)	-6.01	Nebraska	9.02	0.47	(14)	7.38
Arizona	7.47 <sup>b</sup>	0.23	(48)	-11.05	Nevada	6.67 <sup>b</sup>	0.36	(51)	-20.61 <sup>c</sup>
Arkansas	8.77	0.36	(19)	4.47	New Hampshire	7.68	0.52	(43)	-8.58
California	8.10	0.11	(30)	-3.56	New Jersey	8.09	0.21	(31)	-3.59
Colorado	8.00	0.31	(36)	-4.71	New Mexico	6.74 <sup>b</sup>	0.39	(50)	-19.73 <sup>c</sup>
Connecticut	8.07	0.32	(33)	-3.93	New York	7.95	0.14	(38)	-5.25
Delaware	8.42	0.65	(22)	0.34	North Carolina	7.63 <sup>b</sup>	0.21	(45)	-9.16
D.C.	9.40	0.92	(06)	11.95	North Dakota	7.44	0.68	(49)	-11.34
Florida	7.77 <sup>b</sup>	0.12	(42)	-7.43	Ohio	9.43 <sup>b</sup>	0.19	(04)	12.31
Georgia	7.66	0.23	(44)	-8.77	Oklahoma	8.93	0.33	(16)	6.34
Hawaii	7.47	0.48	(47)	-11.04	Oregon	8.59	0.32	(21)	2.35
Idaho	8.07	0.51	(32)	-3.84	Pennsylvania	9.21 <sup>b</sup>	0.17	(09)	9.75
Illinois	8.85	0.19	(17)	5.38	Rhode Island	8.75	0.60	(20)	4.25
Indiana	9.72 <sup>b</sup>	0.27	(01)	15.76 <sup>c</sup>	South Carolina	7.97	0.29	(37)	-5.05
Iowa	9.21	0.35	(10)	9.68	South Dakota	7.80	0.63	(41)	-7.04
Kansas	9.63	0.40	(02)	14.74	Tennessee	9.25	0.27	(07)	10.21
Kentucky	9.22	0.32	(08)	9.79	Texas	8.06	0.14	(34)	-3.96
Louisiana	9.00	0.32	(15)	7.21	Utah	7.48	0.42	(46)	-10.94
Maine	9.15	0.53	(12)	9.00	Vermont	8.06	0.76	(35)	-4.03
Maryland	7.93	0.27	(39)	-5.56	Virginia	8.33	0.24	(24)	-0.78
Massachusetts	8.30	0.24	(25)	-1.16	Washington	8.79	0.26	(18)	4.71
Michigan	9.19 <sup>b</sup>	0.21	(11)	9.46	West Virginia	9.14	0.45	(13)	8.85
Minnesota	9.57 <sup>b</sup>	0.30	(03)	14.03	Wisconsin	9.41 <sup>b</sup>	0.27	(05)	12.10
Mississippi	8.29	0.38	(26)	-1.28	Wyoming	8.11	0.86	(29)	-3.36
Missouri	8.38	0.25	(23)	-0.14					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.24  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.20	0.02		
High Five States				
West Virginia	6.37 <sup>b</sup>	0.32	(01)	22.62 <sup>c</sup>
Michigan	6.06 <sup>b</sup>	0.14	(02)	16.64 <sup>c</sup>
Wyoming	5.95	0.64	(03)	14.45
Oklahoma	5.93	0.23	(04)	14.12
Kentucky	5.91	0.21	(05)	13.75
Low Five States				
Georgia	4.57 <sup>b</sup>	0.14	(47)	-12.09
New Mexico	4.40	0.28	(48)	-15.39 <sup>c</sup>
Colorado	4.39 <sup>b</sup>	0.19	(49)	-15.48 <sup>c</sup>
Hawaii	4.17	0.32	(50)	-19.85 <sup>c</sup>
D.C.	3.48 <sup>b</sup>	0.44	(51)	-33.00 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.32	0.19	(23)	2.34	Montana	5.37	0.42	(21)	3.38
Alaska	5.75	0.76	(06)	10.73	Nebraska	5.71	0.32	(08)	9.90
Arizona	4.78	0.16	(43)	-8.03	Nevada	4.81	0.28	(42)	-7.51
Arkansas	5.21	0.24	(25)	0.31	New Hampshire	4.96	0.36	(36)	-4.57
California	4.97	0.07	(34)	-4.28	New Jersey	5.46	0.14	(18)	5.16
Colorado	4.39 <sup>b</sup>	0.19	(49)	-15.48 <sup>c</sup>	New Mexico	4.40	0.28	(48)	-15.39 <sup>c</sup>
Connecticut	5.18	0.21	(27)	-0.33	New York	4.89	0.09	(38)	-5.87
Delaware	4.96	0.43	(37)	-4.60	North Carolina	5.00	0.14	(31)	-3.70
D.C.	3.48 <sup>b</sup>	0.44	(51)	-33.00 <sup>c</sup>	North Dakota	5.53	0.51	(14)	6.45
Florida	4.89 <sup>b</sup>	0.09	(39)	-5.98	Ohio	5.55	0.12	(12)	6.90
Georgia	4.57 <sup>b</sup>	0.14	(47)	-12.09	Oklahoma	5.93	0.23	(04)	14.12
Hawaii	4.17	0.32	(50)	-19.85 <sup>c</sup>	Oregon	5.65	0.22	(09)	8.82
Idaho	5.44	0.37	(20)	4.75	Pennsylvania	5.63 <sup>b</sup>	0.11	(10)	8.28
Illinois	5.47	0.12	(17)	5.20	Rhode Island	4.58	0.36	(46)	-11.84
Indiana	5.61	0.17	(11)	7.97	South Carolina	4.81	0.19	(41)	-7.35
Iowa	5.48	0.23	(15)	5.48	South Dakota	5.07	0.44	(30)	-2.44
Kansas	5.25	0.25	(24)	1.02	Tennessee	5.47	0.17	(16)	5.24
Kentucky	5.91	0.21	(05)	13.75	Texas	5.00	0.09	(33)	-3.86
Louisiana	5.15	0.20	(28)	-0.87	Utah	4.63	0.29	(45)	-10.86
Maine	5.46	0.35	(19)	5.04	Vermont	4.96	0.50	(35)	-4.53
Maryland	4.85	0.17	(40)	-6.62	Virginia	5.00	0.15	(32)	-3.77
Massachusetts	5.08	0.15	(29)	-2.34	Washington	5.54	0.18	(13)	6.68
Michigan	6.06 <sup>b</sup>	0.14	(02)	16.64 <sup>c</sup>	West Virginia	6.37 <sup>b</sup>	0.32	(01)	22.62 <sup>c</sup>
Minnesota	5.19	0.18	(26)	-0.14	Wisconsin	5.73	0.18	(07)	10.32
Mississippi	4.76	0.24	(44)	-8.43	Wyoming	5.95	0.64	(03)	14.45
Missouri	5.32	0.17	(22)	2.41					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.24  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.47	0.01		
High Five States				
D.C.	3.61	0.35	(01)	46.02 <sup>c</sup>
Mississippi	3.48 <sup>b</sup>	0.15	(02)	40.66 <sup>c</sup>
Louisiana	3.13 <sup>b</sup>	0.12	(03)	26.70 <sup>c</sup>
South Carolina	2.99 <sup>b</sup>	0.11	(04)	20.88 <sup>c</sup>
Washington	2.95 <sup>b</sup>	0.10	(05)	19.57 <sup>c</sup>
Low Five States				
Connecticut	2.00 <sup>b</sup>	0.10	(47)	-19.14 <sup>c</sup>
Kansas	1.93 <sup>b</sup>	0.11	(48)	-21.97 <sup>c</sup>
Alaska	1.93	0.30	(49)	-22.03 <sup>c</sup>
South Dakota	1.92	0.21	(50)	-22.47 <sup>c</sup>
Utah	1.39 <sup>b</sup>	0.12	(51)	-43.56 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.87 <sup>b</sup>	0.11	(08)	15.98 <sup>c</sup>	Montana	2.04	0.19	(44)	-17.61 <sup>c</sup>
Alaska	1.93	0.30	(49)	-22.03 <sup>c</sup>	Nebraska	2.13	0.15	(42)	-13.99
Arizona	2.04 <sup>b</sup>	0.08	(43)	-17.24 <sup>c</sup>	Nevada	2.32	0.14	(32)	-6.22
Arkansas	2.86	0.13	(09)	15.61 <sup>c</sup>	New Hampshire	2.50	0.19	(24)	1.23
California	2.53	0.04	(21)	2.28	New Jersey	2.16 <sup>b</sup>	0.07	(39)	-12.54
Colorado	2.03 <sup>b</sup>	0.10	(45)	-17.65 <sup>c</sup>	New Mexico	2.22	0.15	(37)	-10.33
Connecticut	2.00 <sup>b</sup>	0.10	(47)	-19.14 <sup>c</sup>	New York	2.14 <sup>b</sup>	0.04	(41)	-13.57
Delaware	2.50	0.23	(23)	1.24	North Carolina	2.57	0.07	(19)	4.13
D.C.	3.61	0.35	(01)	46.02 <sup>c</sup>	North Dakota	2.21	0.25	(38)	-10.39
Florida	2.77 <sup>b</sup>	0.05	(10)	12.24	Ohio	2.63	0.06	(15)	6.42
Georgia	2.58	0.08	(18)	4.27	Oklahoma	2.61	0.12	(17)	5.45
Hawaii	2.93	0.20	(06)	18.55 <sup>c</sup>	Oregon	2.43	0.11	(28)	-1.57
Idaho	2.64	0.19	(14)	6.92	Pennsylvania	2.29	0.06	(34)	-7.49
Illinois	2.40	0.06	(29)	-2.84	Rhode Island	2.15	0.19	(40)	-13.17
Indiana	2.30	0.08	(33)	-6.87	South Carolina	2.99 <sup>b</sup>	0.11	(04)	20.88 <sup>c</sup>
Iowa	2.44	0.12	(26)	-1.07	South Dakota	1.92	0.21	(50)	-22.47 <sup>c</sup>
Kansas	1.93 <sup>b</sup>	0.11	(48)	-21.97 <sup>c</sup>	Tennessee	2.92 <sup>b</sup>	0.09	(07)	18.15 <sup>c</sup>
Kentucky	2.72	0.11	(11)	9.94	Texas	2.52	0.05	(22)	2.13
Louisiana	3.13 <sup>b</sup>	0.12	(03)	26.70 <sup>c</sup>	Utah	1.39 <sup>b</sup>	0.12	(51)	-43.56 <sup>c</sup>
Maine	2.48	0.18	(25)	0.29	Vermont	2.69	0.27	(12)	9.02
Maryland	2.61	0.10	(16)	5.50	Virginia	2.27	0.08	(35)	-8.22
Massachusetts	2.37	0.08	(30)	-3.89	Washington	2.95 <sup>b</sup>	0.10	(05)	19.57 <sup>c</sup>
Michigan	2.55	0.07	(20)	3.26	West Virginia	2.66	0.16	(13)	7.60
Minnesota	2.00 <sup>b</sup>	0.09	(46)	-18.95 <sup>c</sup>	Wisconsin	2.44	0.09	(27)	-1.42
Mississippi	3.48 <sup>b</sup>	0.15	(02)	40.66 <sup>c</sup>	Wyoming	2.27	0.29	(36)	-8.26
Missouri	2.34	0.08	(31)	-5.42					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.25  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.79	0.02		
High Five States				
D.C.	5.98	0.71	(01)	57.89 <sup>c</sup>
Mississippi	5.88 <sup>b</sup>	0.30	(02)	55.27 <sup>c</sup>
Louisiana	5.22 <sup>b</sup>	0.23	(03)	37.75 <sup>c</sup>
South Carolina	4.77 <sup>b</sup>	0.21	(04)	25.84 <sup>c</sup>
Tennessee	4.66 <sup>b</sup>	0.18	(05)	22.92 <sup>c</sup>
Low Five States				
South Dakota	2.85	0.37	(47)	-24.79 <sup>c</sup>
Colorado	2.78 <sup>b</sup>	0.17	(48)	-26.52 <sup>c</sup>
Nebraska	2.77 <sup>b</sup>	0.26	(49)	-26.79 <sup>c</sup>
Alaska	2.73	0.53	(50)	-27.85 <sup>c</sup>
Utah	1.73 <sup>b</sup>	0.19	(51)	-54.36 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.65 <sup>b</sup>	0.21	(06)	22.81 <sup>c</sup>	Montana	2.93	0.34	(45)	-22.67 <sup>c</sup>
Alaska	2.73	0.53	(50)	-27.85 <sup>c</sup>	Nebraska	2.77 <sup>b</sup>	0.26	(49)	-26.79 <sup>c</sup>
Arizona	3.05 <sup>b</sup>	0.14	(42)	-19.42 <sup>c</sup>	Nevada	3.30	0.24	(38)	-12.83
Arkansas	4.54	0.26	(07)	19.79 <sup>c</sup>	New Hampshire	3.90	0.36	(20)	2.94
California	3.76	0.07	(25)	-0.80	New Jersey	3.18 <sup>b</sup>	0.13	(41)	-15.95 <sup>c</sup>
Colorado	2.78 <sup>b</sup>	0.17	(48)	-26.52 <sup>c</sup>	New Mexico	3.51	0.27	(33)	-7.48
Connecticut	3.04 <sup>b</sup>	0.19	(43)	-19.69 <sup>c</sup>	New York	3.39 <sup>b</sup>	0.09	(37)	-10.49
Delaware	3.92	0.43	(19)	3.52	North Carolina	4.06	0.14	(16)	7.20
D.C.	5.98	0.71	(01)	57.89 <sup>c</sup>	North Dakota	3.49	0.46	(34)	-7.75
Florida	4.30 <sup>b</sup>	0.09	(10)	13.57	Ohio	4.02	0.12	(17)	6.12
Georgia	4.22	0.16	(12)	11.30	Oklahoma	3.80	0.21	(22)	0.27
Hawaii	4.53	0.37	(08)	19.48 <sup>c</sup>	Oregon	3.69	0.20	(28)	-2.71
Idaho	3.72	0.33	(26)	-1.82	Pennsylvania	3.56	0.10	(30)	-6.08
Illinois	3.51	0.11	(32)	-7.39	Rhode Island	3.25	0.35	(39)	-14.13
Indiana	3.53	0.16	(31)	-6.76	South Carolina	4.77 <sup>b</sup>	0.21	(04)	25.84 <sup>c</sup>
Iowa	3.78	0.22	(24)	-0.35	South Dakota	2.85	0.37	(47)	-24.79 <sup>c</sup>
Kansas	2.87 <sup>b</sup>	0.21	(46)	-24.14 <sup>c</sup>	Tennessee	4.66 <sup>b</sup>	0.18	(05)	22.92 <sup>c</sup>
Kentucky	4.37	0.21	(09)	15.42 <sup>c</sup>	Texas	3.99	0.09	(18)	5.26
Louisiana	5.22 <sup>b</sup>	0.23	(03)	37.75 <sup>c</sup>	Utah	1.73 <sup>b</sup>	0.19	(51)	-54.36 <sup>c</sup>
Maine	3.79	0.33	(23)	0.05	Vermont	4.11	0.52	(15)	8.42
Maryland	4.15	0.19	(13)	9.44	Virginia	3.64	0.15	(29)	-3.82
Massachusetts	3.71	0.16	(27)	-1.99	Washington	4.23	0.17	(11)	11.70
Michigan	3.87	0.13	(21)	2.15	West Virginia	4.14	0.29	(14)	9.16
Minnesota	2.94 <sup>b</sup>	0.16	(44)	-22.44 <sup>c</sup>	Wisconsin	3.43	0.16	(36)	-9.40
Mississippi	5.88 <sup>b</sup>	0.30	(02)	55.27 <sup>c</sup>	Wyoming	3.21	0.52	(40)	-15.25 <sup>c</sup>
Missouri	3.44	0.16	(35)	-9.33					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 20.26  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.38	0.01		
High Five States				
D.C.	1.87	0.34	(01)	34.85 <sup>c</sup>
Washington	1.85 <sup>b</sup>	0.10	(02)	33.47 <sup>c</sup>
Idaho	1.63	0.21	(03)	17.48 <sup>c</sup>
Oklahoma	1.59	0.12	(04)	15.00 <sup>c</sup>
Wisconsin	1.59	0.10	(05)	14.87
Low Five States				
Arizona	1.17	0.08	(46)	-15.76 <sup>c</sup>
New York	1.16 <sup>b</sup>	0.04	(47)	-16.54 <sup>c</sup>
South Dakota	1.08	0.20	(48)	-21.82 <sup>c</sup>
New Mexico	1.07	0.14	(49)	-23.00 <sup>c</sup>
Utah	1.05	0.14	(50)	-24.10 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.45	0.10	(20)	4.41	Montana	1.23	0.20	(41)	-10.88
Alaska	-	-	(-)	-	Nebraska	1.57	0.18	(07)	13.19
Arizona	1.17	0.08	(46)	-15.76 <sup>c</sup>	Nevada	1.41	0.15	(23)	1.75
Arkansas	1.47	0.13	(15)	6.43	New Hampshire	1.34	0.19	(32)	-3.53
California	1.51	0.04	(11)	8.89	New Jersey	1.35	0.07	(29)	-2.67
Colorado	1.41	0.11	(22)	2.04	New Mexico	1.07	0.14	(49)	-23.00 <sup>c</sup>
Connecticut	1.18	0.10	(44)	-14.70	New York	1.16 <sup>b</sup>	0.04	(47)	-16.54 <sup>c</sup>
Delaware	1.34	0.22	(31)	-3.37	North Carolina	1.36	0.07	(28)	-1.84
D.C.	1.87	0.34	(01)	34.85 <sup>c</sup>	North Dakota	1.18	0.26	(43)	-14.46
Florida	1.46	0.05	(17)	5.19	Ohio	1.49	0.06	(12)	7.44
Georgia	1.33	0.08	(34)	-4.11	Oklahoma	1.59	0.12	(04)	15.00 <sup>c</sup>
Hawaii	1.51	0.19	(09)	9.36	Oregon	1.38	0.11	(26)	-0.35
Idaho	1.63	0.21	(03)	17.48 <sup>c</sup>	Pennsylvania	1.28	0.05	(38)	-7.41
Illinois	1.48	0.06	(14)	6.95	Rhode Island	1.30	0.20	(37)	-5.93
Indiana	1.26	0.08	(39)	-8.64	South Carolina	1.51	0.11	(10)	9.24
Iowa	1.33	0.12	(33)	-3.83	South Dakota	1.08	0.20	(48)	-21.82 <sup>c</sup>
Kansas	1.17	0.12	(45)	-15.30 <sup>c</sup>	Tennessee	1.48	0.09	(13)	7.22
Kentucky	1.37	0.10	(27)	-0.91	Texas	1.32	0.05	(35)	-4.29
Louisiana	1.46	0.11	(16)	5.84	Utah	1.05	0.14	(50)	-24.10 <sup>c</sup>
Maine	1.40	0.17	(24)	1.03	Vermont	1.58	0.28	(06)	14.48
Maryland	1.39	0.09	(25)	0.46	Virginia	1.22	0.08	(42)	-12.08
Massachusetts	1.32	0.08	(36)	-4.31	Washington	1.85 <sup>b</sup>	0.10	(02)	33.47 <sup>c</sup>
Michigan	1.45	0.07	(18)	5.00	West Virginia	1.34	0.15	(30)	-2.83
Minnesota	1.24	0.09	(40)	-10.57	Wisconsin	1.59	0.10	(05)	14.87
Mississippi	1.55	0.13	(08)	12.13	Wyoming	1.45	0.32	(19)	4.93
Missouri	1.43	0.09	(21)	3.10					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 21.14  
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	8.23	0.03		
High Five States				
Washington	9.80 <sup>b</sup>	0.24	(01)	18.96 <sup>c</sup>
Wisconsin	9.46 <sup>b</sup>	0.24	(02)	14.82
Oregon	9.30 <sup>b</sup>	0.29	(03)	12.89
Alabama	9.28 <sup>b</sup>	0.25	(04)	12.74
Montana	9.08	0.55	(05)	10.32
Low Five States				
Missouri	7.36 <sup>b</sup>	0.20	(47)	-10.58
Nevada	7.32	0.34	(48)	-11.12
Alaska	7.06	0.79	(49)	-14.28
Vermont	6.97	0.61	(50)	-15.39 <sup>c</sup>
Hawaii	5.99 <sup>b</sup>	0.39	(51)	-27.30 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.28 <sup>b</sup>	0.25	(04)	12.74	Montana	9.08	0.55	(05)	10.32
Alaska	7.06	0.79	(49)	-14.28	Nebraska	8.10	0.40	(29)	-1.61
Arizona	7.88	0.21	(37)	-4.30	Nevada	7.32	0.34	(48)	-11.12
Arkansas	8.37	0.31	(17)	1.60	New Hampshire	7.85	0.45	(39)	-4.64
California	8.21	0.09	(22)	-0.29	New Jersey	8.71	0.18	(09)	5.80
Colorado	8.66	0.27	(11)	5.17	New Mexico	7.92	0.38	(35)	-3.79
Connecticut	8.11	0.27	(27)	-1.51	New York	8.20	0.12	(25)	-0.39
Delaware	8.21	0.56	(24)	-0.31	North Carolina	7.87	0.17	(38)	-4.37
D.C.	7.58	0.68	(44)	-7.97	North Dakota	8.97	0.69	(08)	8.94
Florida	7.57 <sup>b</sup>	0.11	(45)	-8.06	Ohio	8.11	0.15	(26)	-1.50
Georgia	8.34	0.19	(18)	1.31	Oklahoma	8.06	0.27	(31)	-2.12
Hawaii	5.99 <sup>b</sup>	0.39	(51)	-27.30 <sup>c</sup>	Oregon	9.30 <sup>b</sup>	0.29	(03)	12.89
Idaho	8.27	0.47	(21)	0.42	Pennsylvania	9.08 <sup>b</sup>	0.15	(06)	10.22
Illinois	8.32	0.15	(19)	1.04	Rhode Island	7.89	0.48	(36)	-4.22
Indiana	8.60	0.22	(12)	4.40	South Carolina	7.92	0.24	(34)	-3.78
Iowa	8.98	0.31	(07)	9.04	South Dakota	8.45	0.60	(16)	2.63
Kansas	8.69	0.33	(10)	5.51	Tennessee	8.06	0.21	(32)	-2.17
Kentucky	8.06	0.25	(30)	-2.11	Texas	7.69 <sup>b</sup>	0.12	(43)	-6.65
Louisiana	7.85	0.25	(40)	-4.70	Utah	7.70	0.38	(42)	-6.44
Maine	7.82	0.42	(41)	-5.01	Vermont	6.97	0.61	(50)	-15.39 <sup>c</sup>
Maryland	8.10	0.23	(28)	-1.59	Virginia	8.55	0.20	(14)	3.77
Massachusetts	8.29	0.20	(20)	0.64	Washington	9.80 <sup>b</sup>	0.24	(01)	18.96 <sup>c</sup>
Michigan	8.57	0.17	(13)	4.10	West Virginia	8.48	0.37	(15)	2.97
Minnesota	8.21	0.24	(23)	-0.30	Wisconsin	9.46 <sup>b</sup>	0.24	(02)	14.82
Mississippi	7.52	0.30	(46)	-8.68	Wyoming	8.01	0.75	(33)	-2.77
Missouri	7.36 <sup>b</sup>	0.20	(47)	-10.58					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.18  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	10.84	0.03		
High Five States				
D.C.	13.04	0.67	(01)	20.26 <sup>c</sup>
Louisiana	12.27 <sup>b</sup>	0.24	(02)	13.18
Connecticut	12.13 <sup>b</sup>	0.25	(03)	11.90
Michigan	11.80 <sup>b</sup>	0.15	(04)	8.90
New Hampshire	11.78	0.41	(05)	8.70
Low Five States				
South Dakota	9.90	0.46	(47)	-8.63
Colorado	9.84 <sup>b</sup>	0.22	(48)	-9.26
Arizona	9.34 <sup>b</sup>	0.17	(49)	-13.80
West Virginia	9.28 <sup>b</sup>	0.29	(50)	-14.36
Utah	8.69 <sup>b</sup>	0.30	(51)	-19.80 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	11.18	0.21	(19)	3.14	Montana	10.38	0.43	(40)	-4.28
Alaska	11.22	0.76	(16)	3.49	Nebraska	10.68	0.33	(36)	-1.51
Arizona	9.34 <sup>b</sup>	0.17	(49)	-13.80	Nevada	10.97	0.31	(27)	1.17
Arkansas	11.35	0.27	(12)	4.66	New Hampshire	11.78	0.41	(05)	8.70
California	10.46 <sup>b</sup>	0.08	(38)	-3.47	New Jersey	11.44 <sup>b</sup>	0.16	(10)	5.50
Colorado	9.84 <sup>b</sup>	0.22	(48)	-9.26	New Mexico	10.11	0.32	(44)	-6.74
Connecticut	12.13 <sup>b</sup>	0.25	(03)	11.90	New York	10.99	0.10	(25)	1.37
Delaware	10.93	0.48	(29)	0.80	North Carolina	10.81	0.15	(32)	-0.29
D.C.	13.04	0.67	(01)	20.26 <sup>c</sup>	North Dakota	10.71	0.54	(34)	-1.22
Florida	10.20 <sup>b</sup>	0.09	(43)	-5.91	Ohio	11.35 <sup>b</sup>	0.13	(13)	4.66
Georgia	10.39	0.16	(39)	-4.16	Oklahoma	10.24	0.23	(41)	-5.54
Hawaii	11.02	0.39	(20)	1.70	Oregon	11.01	0.23	(21)	1.61
Idaho	10.68	0.39	(35)	-1.49	Pennsylvania	11.49 <sup>b</sup>	0.12	(09)	5.96
Illinois	11.41 <sup>b</sup>	0.13	(11)	5.22	Rhode Island	10.01	0.41	(46)	-7.64
Indiana	10.99	0.18	(23)	1.43	South Carolina	10.95	0.22	(28)	1.01
Iowa	10.23	0.24	(42)	-5.60	South Dakota	9.90	0.46	(47)	-8.63
Kansas	10.81	0.27	(31)	-0.29	Tennessee	10.92	0.18	(30)	0.73
Kentucky	10.74	0.22	(33)	-0.93	Texas	10.03 <sup>b</sup>	0.10	(45)	-7.48
Louisiana	12.27 <sup>b</sup>	0.24	(02)	13.18	Utah	8.69 <sup>b</sup>	0.30	(51)	-19.80 <sup>c</sup>
Maine	10.98	0.37	(26)	1.29	Vermont	11.00	0.56	(22)	1.48
Maryland	11.51	0.20	(07)	6.22	Virginia	11.25	0.17	(15)	3.79
Massachusetts	11.50 <sup>b</sup>	0.18	(08)	6.09	Washington	10.99	0.19	(24)	1.42
Michigan	11.80 <sup>b</sup>	0.15	(04)	8.90	West Virginia	9.28 <sup>b</sup>	0.29	(50)	-14.36
Minnesota	10.54	0.20	(37)	-2.75	Wisconsin	11.20	0.19	(17)	3.34
Mississippi	11.61	0.28	(06)	7.10	Wyoming	11.33	0.66	(14)	4.48
Missouri	11.18	0.19	(18)	3.15					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.19  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	12.51	0.04		
High Five States				
D.C.	16.34	1.21	(01)	30.60 <sup>c</sup>
Connecticut	14.69 <sup>b</sup>	0.43	(02)	17.44 <sup>c</sup>
Michigan	13.92 <sup>b</sup>	0.25	(03)	11.21 <sup>c</sup>
Louisiana	13.82	0.39	(04)	10.48 <sup>c</sup>
Mississippi	13.77	0.48	(05)	10.05 <sup>c</sup>
Low Five States				
West Virginia	11.23	0.48	(47)	-10.25 <sup>c</sup>
South Dakota	11.09	0.74	(48)	-11.34 <sup>c</sup>
Arizona	10.95 <sup>b</sup>	0.27	(49)	-12.49 <sup>c</sup>
Colorado	10.89 <sup>b</sup>	0.35	(50)	-12.94 <sup>c</sup>
Utah	9.48 <sup>b</sup>	0.47	(51)	-24.24 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	13.33	0.36	(09)	6.52	Montana	12.39	0.71	(30)	-0.99
Alaska	12.34	1.17	(32)	-1.40	Nebraska	12.17	0.54	(37)	-2.73
Arizona	10.95 <sup>b</sup>	0.27	(49)	-12.49 <sup>c</sup>	Nevada	12.30	0.48	(34)	-1.69
Arkansas	13.64	0.45	(06)	9.03	New Hampshire	13.41	0.67	(07)	7.19
California	11.80 <sup>b</sup>	0.13	(43)	-5.73	New Jersey	13.29	0.26	(10)	6.23
Colorado	10.89 <sup>b</sup>	0.35	(50)	-12.94 <sup>c</sup>	New Mexico	11.56	0.51	(45)	-7.64
Connecticut	14.69 <sup>b</sup>	0.43	(02)	17.44 <sup>c</sup>	New York	12.64	0.17	(23)	1.06
Delaware	12.31	0.77	(33)	-1.61	North Carolina	12.14	0.25	(38)	-2.97
D.C.	16.34	1.21	(01)	30.60 <sup>c</sup>	North Dakota	12.79	0.88	(22)	2.23
Florida	11.98 <sup>b</sup>	0.15	(41)	-4.23	Ohio	13.12	0.22	(13)	4.87
Georgia	12.42	0.28	(28)	-0.71	Oklahoma	12.03	0.38	(39)	-3.82
Hawaii	12.88	0.63	(20)	2.97	Oregon	12.22	0.37	(35)	-2.36
Idaho	11.51	0.59	(46)	-8.01	Pennsylvania	13.37 <sup>b</sup>	0.20	(08)	6.88
Illinois	13.13	0.22	(12)	4.90	Rhode Island	12.37	0.71	(31)	-1.14
Indiana	13.10	0.31	(14)	4.70	South Carolina	12.52	0.36	(24)	0.08
Iowa	12.01	0.40	(40)	-4.05	South Dakota	11.09	0.74	(48)	-11.34 <sup>c</sup>
Kansas	12.52	0.44	(25)	0.04	Tennessee	12.96	0.31	(18)	3.55
Kentucky	12.48	0.37	(27)	-0.26	Texas	11.74 <sup>b</sup>	0.17	(44)	-6.20
Louisiana	13.82	0.39	(04)	10.48 <sup>c</sup>	Utah	9.48 <sup>b</sup>	0.47	(51)	-24.24 <sup>c</sup>
Maine	12.20	0.59	(36)	-2.47	Vermont	12.51	0.90	(26)	-0.05
Maryland	12.92	0.34	(19)	3.24	Virginia	12.98	0.29	(17)	3.74
Massachusetts	13.08	0.30	(15)	4.50	Washington	12.41	0.30	(29)	-0.82
Michigan	13.92 <sup>b</sup>	0.25	(03)	11.21 <sup>c</sup>	West Virginia	11.23	0.48	(47)	-10.25 <sup>c</sup>
Minnesota	11.81	0.32	(42)	-5.63	Wisconsin	12.88	0.32	(21)	2.92
Mississippi	13.77	0.48	(05)	10.05 <sup>c</sup>	Wyoming	13.17	1.06	(11)	5.22
Missouri	13.06	0.31	(16)	4.41					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.20  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	9.48	0.03		
High Five States				
Louisiana	11.00 <sup>b</sup>	0.30	(01)	15.97 <sup>c</sup>
D.C.	10.66	0.79	(02)	12.42
New Hampshire	10.62	0.52	(03)	12.00
Maryland	10.44 <sup>b</sup>	0.26	(04)	10.09
Massachusetts	10.27 <sup>b</sup>	0.22	(05)	8.32
Low Five States				
Texas	8.66 <sup>b</sup>	0.12	(47)	-8.73
Rhode Island	8.38	0.49	(48)	-11.69
Utah	7.96 <sup>b</sup>	0.39	(49)	-16.13 <sup>c</sup>
Arizona	7.95 <sup>b</sup>	0.21	(50)	-16.20 <sup>c</sup>
West Virginia	7.67 <sup>b</sup>	0.35	(51)	-19.18 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.46	0.25	(29)	-0.30	Montana	8.67	0.54	(46)	-8.57
Alaska	10.04	0.99	(09)	5.85	Nebraska	9.43	0.42	(30)	-0.57
Arizona	7.95 <sup>b</sup>	0.21	(50)	-16.20 <sup>c</sup>	Nevada	9.77	0.40	(20)	3.00
Arkansas	9.41	0.33	(31)	-0.78	New Hampshire	10.62	0.52	(03)	12.00
California	9.38	0.10	(35)	-1.13	New Jersey	10.01	0.19	(10)	5.53
Colorado	8.93	0.28	(39)	-5.87	New Mexico	8.92	0.40	(40)	-5.91
Connecticut	10.19	0.30	(06)	7.45	New York	9.74	0.13	(21)	2.66
Delaware	9.72	0.60	(22)	2.44	North Carolina	9.67	0.19	(26)	2.00
D.C.	10.66	0.79	(02)	12.42	North Dakota	8.71	0.65	(44)	-8.13
Florida	8.68 <sup>b</sup>	0.12	(45)	-8.50	Ohio	9.92	0.16	(13)	4.58
Georgia	8.86	0.20	(41)	-6.62	Oklahoma	8.73	0.28	(43)	-7.93
Hawaii	9.40	0.48	(32)	-0.85	Oregon	10.00	0.30	(11)	5.42
Idaho	9.84	0.50	(17)	3.70	Pennsylvania	9.97	0.15	(12)	5.08
Illinois	10.07 <sup>b</sup>	0.17	(08)	6.16	Rhode Island	8.38	0.49	(48)	-11.69
Indiana	9.40	0.22	(33)	-0.86	South Carolina	9.68	0.27	(25)	2.11
Iowa	8.77	0.30	(42)	-7.53	South Dakota	9.13	0.61	(38)	-3.78
Kansas	9.38	0.34	(34)	-1.11	Tennessee	9.34	0.22	(37)	-1.56
Kentucky	9.35	0.27	(36)	-1.38	Texas	8.66 <sup>b</sup>	0.12	(47)	-8.73
Louisiana	11.00 <sup>b</sup>	0.30	(01)	15.97 <sup>c</sup>	Utah	7.96 <sup>b</sup>	0.39	(49)	-16.13 <sup>c</sup>
Maine	9.80	0.46	(18)	3.37	Vermont	9.63	0.70	(27)	1.52
Maryland	10.44 <sup>b</sup>	0.26	(04)	10.09	Virginia	9.86	0.21	(15)	3.95
Massachusetts	10.27 <sup>b</sup>	0.22	(05)	8.32	Washington	9.79	0.24	(19)	3.24
Michigan	10.13 <sup>b</sup>	0.18	(07)	6.84	West Virginia	7.67 <sup>b</sup>	0.35	(51)	-19.18 <sup>c</sup>
Minnesota	9.48	0.25	(28)	-0.08	Wisconsin	9.84	0.24	(16)	3.77
Mississippi	9.89	0.34	(14)	4.24	Wyoming	9.71	0.82	(23)	2.40
Missouri	9.69	0.23	(24)	2.13					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 23.14  
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	23.58	0.06		
High Five States				
D.C.	41.33 <sup>b</sup>	2.08	(01)	75.28 <sup>c</sup>
Mississippi	30.95 <sup>b</sup>	0.77	(02)	31.26 <sup>c</sup>
Alabama	28.72 <sup>b</sup>	0.57	(03)	21.81 <sup>c</sup>
Georgia	27.45 <sup>b</sup>	0.47	(04)	16.42 <sup>c</sup>
Montana	27.22	1.10	(05)	15.43 <sup>c</sup>
Low Five States				
Kansas	21.38 <sup>b</sup>	0.60	(47)	-9.35
Wyoming	20.90	1.42	(48)	-11.38 <sup>c</sup>
Arizona	19.72 <sup>b</sup>	0.38	(49)	-16.38 <sup>c</sup>
Florida	19.64 <sup>b</sup>	0.19	(50)	-16.69 <sup>c</sup>
Hawaii	16.23 <sup>b</sup>	0.71	(51)	-31.16 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	28.72 <sup>b</sup>	0.57	(03)	21.81 <sup>c</sup>	Montana	27.22	1.10	(05)	15.43 <sup>c</sup>
Alaska	22.07	1.83	(43)	-6.40	Nebraska	24.68	0.80	(22)	4.67
Arizona	19.72 <sup>b</sup>	0.38	(49)	-16.38 <sup>c</sup>	Nevada	23.40	0.72	(34)	-0.75
Arkansas	25.26	0.64	(17)	7.12	New Hampshire	23.23	0.95	(35)	-1.51
California	23.19	0.19	(36)	-1.67	New Jersey	22.41	0.36	(41)	-4.95
Colorado	23.81	0.56	(29)	0.99	New Mexico	24.28	0.78	(27)	2.96
Connecticut	24.79	0.57	(21)	5.14	New York	22.18 <sup>b</sup>	0.24	(42)	-5.95
Delaware	24.85	1.17	(20)	5.40	North Carolina	25.86 <sup>b</sup>	0.40	(11)	9.66
D.C.	41.33 <sup>b</sup>	2.08	(01)	75.28 <sup>c</sup>	North Dakota	25.16	1.24	(18)	6.67
Florida	19.64 <sup>b</sup>	0.19	(50)	-16.69 <sup>c</sup>	Ohio	25.41 <sup>b</sup>	0.33	(15)	7.74
Georgia	27.45 <sup>b</sup>	0.47	(04)	16.42 <sup>c</sup>	Oklahoma	23.57	0.57	(32)	-0.03
Hawaii	16.23 <sup>b</sup>	0.71	(51)	-31.16 <sup>c</sup>	Oregon	25.70 <sup>b</sup>	0.56	(12)	8.98
Idaho	26.74	0.96	(08)	13.39 <sup>c</sup>	Pennsylvania	23.74	0.28	(31)	0.67
Illinois	25.47 <sup>b</sup>	0.33	(14)	8.01	Rhode Island	22.51	0.98	(40)	-4.55
Indiana	23.75	0.45	(30)	0.72	South Carolina	26.92 <sup>b</sup>	0.57	(07)	14.16 <sup>c</sup>
Iowa	23.93	0.58	(28)	1.48	South Dakota	22.94	1.08	(37)	-2.71
Kansas	21.38 <sup>b</sup>	0.60	(47)	-9.35	Tennessee	25.30 <sup>b</sup>	0.47	(16)	7.30
Kentucky	24.56	0.56	(23)	4.15	Texas	21.45 <sup>b</sup>	0.24	(46)	-9.05
Louisiana	27.07 <sup>b</sup>	0.59	(06)	14.79 <sup>c</sup>	Utah	24.51	0.80	(24)	3.93
Maine	24.44	0.90	(25)	3.64	Vermont	21.98	1.30	(44)	-6.78
Maryland	26.67 <sup>b</sup>	0.53	(09)	13.10 <sup>c</sup>	Virginia	25.99 <sup>b</sup>	0.45	(10)	10.23 <sup>c</sup>
Massachusetts	23.44	0.42	(33)	-0.61	Washington	24.89	0.45	(19)	5.53
Michigan	22.62	0.34	(39)	-4.07	West Virginia	21.74	0.72	(45)	-7.83
Minnesota	24.34	0.48	(26)	3.20	Wisconsin	25.60 <sup>b</sup>	0.47	(13)	8.54
Mississippi	30.95 <sup>b</sup>	0.77	(02)	31.26 <sup>c</sup>	Wyoming	20.90	1.42	(48)	-11.38 <sup>c</sup>
Missouri	22.75	0.43	(38)	-3.54					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 24.18  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.61	0.02		
High Five States				
Hawaii	6.39 <sup>b</sup>	0.30	(01)	76.95 <sup>c</sup>
D.C.	4.80	0.41	(02)	33.01 <sup>c</sup>
Louisiana	4.68 <sup>b</sup>	0.15	(03)	29.71 <sup>c</sup>
California	4.49 <sup>b</sup>	0.05	(04)	24.39 <sup>c</sup>
New Jersey	4.40 <sup>b</sup>	0.10	(05)	21.79 <sup>c</sup>
Low Five States				
Nebraska	2.61 <sup>b</sup>	0.16	(47)	-27.62 <sup>c</sup>
Wyoming	2.59	0.31	(48)	-28.29 <sup>c</sup>
Iowa	2.53 <sup>b</sup>	0.12	(49)	-29.96 <sup>c</sup>
Utah	2.32 <sup>b</sup>	0.15	(50)	-35.61 <sup>c</sup>
Vermont	2.23 <sup>b</sup>	0.25	(51)	-38.23 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.70	0.12	(16)	2.44	Montana	2.62 <sup>b</sup>	0.22	(46)	-27.48 <sup>c</sup>
Alaska	4.32	0.46	(07)	19.71 <sup>c</sup>	Nebraska	2.61 <sup>b</sup>	0.16	(47)	-27.62 <sup>c</sup>
Arizona	2.97 <sup>b</sup>	0.10	(34)	-17.82 <sup>c</sup>	Nevada	3.51	0.18	(21)	-2.84
Arkansas	3.34	0.15	(26)	-7.58	New Hampshire	2.88 <sup>b</sup>	0.20	(41)	-20.31 <sup>c</sup>
California	4.49 <sup>b</sup>	0.05	(04)	24.39 <sup>c</sup>	New Jersey	4.40 <sup>b</sup>	0.10	(05)	21.79 <sup>c</sup>
Colorado	2.91 <sup>b</sup>	0.12	(35)	-19.26 <sup>c</sup>	New Mexico	4.38 <sup>b</sup>	0.21	(06)	21.27 <sup>c</sup>
Connecticut	4.24 <sup>b</sup>	0.15	(09)	17.36 <sup>c</sup>	New York	4.30 <sup>b</sup>	0.06	(08)	19.01 <sup>c</sup>
Delaware	3.96	0.29	(13)	9.68	North Carolina	3.33	0.09	(27)	-7.65
D.C.	4.80	0.41	(02)	33.01 <sup>c</sup>	North Dakota	3.39	0.30	(24)	-6.04
Florida	3.16 <sup>b</sup>	0.05	(31)	-12.38	Ohio	3.29 <sup>b</sup>	0.07	(29)	-8.86
Georgia	3.65	0.10	(18)	1.21	Oklahoma	3.00 <sup>b</sup>	0.13	(33)	-17.00 <sup>c</sup>
Hawaii	6.39 <sup>b</sup>	0.30	(01)	76.95 <sup>c</sup>	Oregon	2.62 <sup>b</sup>	0.11	(45)	-27.45 <sup>c</sup>
Idaho	2.90	0.20	(38)	-19.62 <sup>c</sup>	Pennsylvania	3.39	0.07	(25)	-6.15
Illinois	3.97 <sup>b</sup>	0.08	(11)	9.90	Rhode Island	3.66	0.24	(17)	1.32
Indiana	2.84 <sup>b</sup>	0.09	(43)	-21.43 <sup>c</sup>	South Carolina	3.96	0.13	(12)	9.70
Iowa	2.53 <sup>b</sup>	0.12	(49)	-29.96 <sup>c</sup>	South Dakota	3.53	0.28	(20)	-2.34
Kansas	2.65 <sup>b</sup>	0.13	(44)	-26.51 <sup>c</sup>	Tennessee	3.41	0.10	(23)	-5.57
Kentucky	3.15 <sup>b</sup>	0.12	(32)	-12.71	Texas	3.86 <sup>b</sup>	0.06	(15)	6.85
Louisiana	4.68 <sup>b</sup>	0.15	(03)	29.71 <sup>c</sup>	Utah	2.32 <sup>b</sup>	0.15	(50)	-35.61 <sup>c</sup>
Maine	2.89 <sup>b</sup>	0.19	(39)	-19.89 <sup>c</sup>	Vermont	2.23 <sup>b</sup>	0.25	(51)	-38.23 <sup>c</sup>
Maryland	3.87	0.12	(14)	7.07	Virginia	3.32	0.09	(28)	-7.92
Massachusetts	3.60	0.10	(19)	-0.30	Washington	3.25 <sup>b</sup>	0.10	(30)	-9.89
Michigan	3.43	0.08	(22)	-5.04	West Virginia	2.90 <sup>b</sup>	0.16	(37)	-19.56 <sup>c</sup>
Minnesota	2.87 <sup>b</sup>	0.10	(42)	-20.60 <sup>c</sup>	Wisconsin	2.91 <sup>b</sup>	0.10	(36)	-19.50 <sup>c</sup>
Mississippi	4.21 <sup>b</sup>	0.17	(10)	16.61 <sup>c</sup>	Wyoming	2.59	0.31	(48)	-28.29 <sup>c</sup>
Missouri	2.89 <sup>b</sup>	0.09	(40)	-19.99 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.19  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.00	0.03		
High Five States				
Hawaii	8.86 <sup>b</sup>	0.53	(01)	77.21 <sup>c</sup>
Louisiana	6.65 <sup>b</sup>	0.27	(02)	33.05 <sup>c</sup>
D.C.	6.60	0.76	(03)	32.01 <sup>c</sup>
New Mexico	6.21	0.37	(04)	24.13 <sup>c</sup>
Connecticut	6.20 <sup>b</sup>	0.28	(05)	24.03 <sup>c</sup>
Low Five States				
Oregon	3.66 <sup>b</sup>	0.20	(47)	-26.74 <sup>c</sup>
Iowa	3.65 <sup>b</sup>	0.22	(48)	-26.89 <sup>c</sup>
Vermont	3.44	0.49	(49)	-31.28 <sup>c</sup>
Wyoming	3.07 <sup>b</sup>	0.50	(50)	-38.67 <sup>c</sup>
Utah	2.84 <sup>b</sup>	0.26	(51)	-43.20 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.12	0.23	(17)	2.43	Montana	3.67	0.39	(46)	-26.67 <sup>c</sup>
Alaska	5.10	0.73	(18)	1.95	Nebraska	3.77 <sup>b</sup>	0.30	(45)	-24.64 <sup>c</sup>
Arizona	3.84 <sup>b</sup>	0.16	(43)	-23.25 <sup>c</sup>	Nevada	4.50	0.30	(28)	-10.07
Arkansas	4.45	0.26	(29)	-10.90	New Hampshire	4.13	0.37	(38)	-17.36 <sup>c</sup>
California	5.98 <sup>b</sup>	0.09	(09)	19.73 <sup>c</sup>	New Jersey	6.12 <sup>b</sup>	0.18	(06)	22.48 <sup>c</sup>
Colorado	4.04 <sup>b</sup>	0.22	(41)	-19.08 <sup>c</sup>	New Mexico	6.21	0.37	(04)	24.13 <sup>c</sup>
Connecticut	6.20 <sup>b</sup>	0.28	(05)	24.03 <sup>c</sup>	New York	6.02 <sup>b</sup>	0.12	(08)	20.47 <sup>c</sup>
Delaware	5.65	0.52	(12)	12.93	North Carolina	4.79	0.16	(23)	-4.09
D.C.	6.60	0.76	(03)	32.01 <sup>c</sup>	North Dakota	4.71	0.54	(25)	-5.71
Florida	4.24 <sup>b</sup>	0.09	(34)	-15.20 <sup>c</sup>	Ohio	4.72	0.14	(24)	-5.66
Georgia	4.88	0.18	(22)	-2.31	Oklahoma	4.16 <sup>b</sup>	0.23	(36)	-16.79 <sup>c</sup>
Hawaii	8.86 <sup>b</sup>	0.53	(01)	77.21 <sup>c</sup>	Oregon	3.66 <sup>b</sup>	0.20	(47)	-26.74 <sup>c</sup>
Idaho	3.95	0.36	(42)	-21.05 <sup>c</sup>	Pennsylvania	4.91	0.12	(21)	-1.72
Illinois	5.43	0.14	(14)	8.64	Rhode Island	6.06	0.50	(07)	21.22 <sup>c</sup>
Indiana	4.16 <sup>b</sup>	0.18	(35)	-16.74 <sup>c</sup>	South Carolina	5.74	0.24	(11)	14.88
Iowa	3.65 <sup>b</sup>	0.22	(48)	-26.89 <sup>c</sup>	South Dakota	4.58	0.48	(26)	-8.41
Kansas	3.78 <sup>b</sup>	0.24	(44)	-24.38 <sup>c</sup>	Tennessee	5.00	0.20	(19)	0.02
Kentucky	4.44	0.22	(30)	-11.13	Texas	5.15	0.11	(16)	2.97
Louisiana	6.65 <sup>b</sup>	0.27	(02)	33.05 <sup>c</sup>	Utah	2.84 <sup>b</sup>	0.26	(51)	-43.20 <sup>c</sup>
Maine	4.42	0.36	(31)	-11.63	Vermont	3.44	0.49	(49)	-31.28 <sup>c</sup>
Maryland	5.34	0.22	(15)	6.76	Virginia	4.53	0.17	(27)	-9.34
Massachusetts	5.43	0.19	(13)	8.71	Washington	4.31 <sup>b</sup>	0.18	(32)	-13.85
Michigan	4.92	0.15	(20)	-1.48	West Virginia	4.13	0.30	(39)	-17.46 <sup>c</sup>
Minnesota	4.13 <sup>b</sup>	0.19	(40)	-17.47 <sup>c</sup>	Wisconsin	4.28 <sup>b</sup>	0.18	(33)	-14.36
Mississippi	5.81	0.31	(10)	16.31 <sup>c</sup>	Wyoming	3.07 <sup>b</sup>	0.50	(50)	-38.67 <sup>c</sup>
Missouri	4.16 <sup>b</sup>	0.18	(37)	-16.80 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 24.20  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.57	0.02		
High Five States				
Hawaii	4.55 <sup>b</sup>	0.34	(01)	76.84 <sup>c</sup>
Alaska	3.56	0.57	(02)	38.44 <sup>c</sup>
D.C.	3.47	0.46	(03)	34.69 <sup>c</sup>
California	3.37 <sup>b</sup>	0.06	(04)	31.12 <sup>c</sup>
Louisiana	3.19 <sup>b</sup>	0.16	(05)	24.00 <sup>c</sup>
Low Five States				
Montana	1.76	0.24	(47)	-31.77 <sup>c</sup>
Maine	1.73 <sup>b</sup>	0.19	(48)	-32.66 <sup>c</sup>
Nebraska	1.73 <sup>b</sup>	0.18	(49)	-32.66 <sup>c</sup>
Iowa	1.66 <sup>b</sup>	0.13	(50)	-35.49 <sup>c</sup>
Vermont	1.40 <sup>b</sup>	0.26	(51)	-45.62 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.66	0.14	(16)	3.53	Montana	1.76	0.24	(47)	-31.77 <sup>c</sup>
Alaska	3.56	0.57	(02)	38.44 <sup>c</sup>	Nebraska	1.73 <sup>b</sup>	0.18	(49)	-32.66 <sup>c</sup>
Arizona	2.27	0.11	(29)	-11.96	Nevada	2.65	0.21	(17)	3.08
Arkansas	2.48	0.17	(20)	-3.71	New Hampshire	1.86	0.21	(44)	-27.74 <sup>c</sup>
California	3.37 <sup>b</sup>	0.06	(04)	31.12 <sup>c</sup>	New Jersey	3.17 <sup>b</sup>	0.11	(06)	23.22 <sup>c</sup>
Colorado	2.05 <sup>b</sup>	0.13	(36)	-20.24 <sup>c</sup>	New Mexico	2.89	0.23	(11)	12.23
Connecticut	2.79	0.16	(14)	8.24	New York	3.08 <sup>b</sup>	0.07	(07)	19.51 <sup>c</sup>
Delaware	2.55	0.31	(19)	-0.74	North Carolina	2.26	0.09	(30)	-12.11
D.C.	3.47	0.46	(03)	34.69 <sup>c</sup>	North Dakota	2.43	0.34	(22)	-5.59
Florida	2.29 <sup>b</sup>	0.06	(26)	-11.17	Ohio	2.28 <sup>b</sup>	0.08	(27)	-11.30
Georgia	2.79	0.11	(13)	8.62	Oklahoma	2.11	0.14	(34)	-17.91 <sup>c</sup>
Hawaii	4.55 <sup>b</sup>	0.34	(01)	76.84 <sup>c</sup>	Oregon	1.83 <sup>b</sup>	0.13	(45)	-28.79 <sup>c</sup>
Idaho	2.05	0.23	(37)	-20.28 <sup>c</sup>	Pennsylvania	2.26 <sup>b</sup>	0.07	(31)	-12.35
Illinois	2.93 <sup>b</sup>	0.09	(09)	14.03	Rhode Island	2.19	0.25	(33)	-14.95
Indiana	1.89 <sup>b</sup>	0.10	(43)	-26.62 <sup>c</sup>	South Carolina	2.57	0.14	(18)	-0.26
Iowa	1.66 <sup>b</sup>	0.13	(50)	-35.49 <sup>c</sup>	South Dakota	2.76	0.34	(15)	7.18
Kansas	1.81 <sup>b</sup>	0.15	(46)	-29.54 <sup>c</sup>	Tennessee	2.27	0.11	(28)	-11.72
Kentucky	2.20	0.13	(32)	-14.61	Texas	2.85 <sup>b</sup>	0.07	(12)	10.65
Louisiana	3.19 <sup>b</sup>	0.16	(05)	24.00 <sup>c</sup>	Utah	1.90 <sup>b</sup>	0.19	(41)	-26.08 <sup>c</sup>
Maine	1.73 <sup>b</sup>	0.19	(48)	-32.66 <sup>c</sup>	Vermont	1.40 <sup>b</sup>	0.26	(51)	-45.62 <sup>c</sup>
Maryland	2.92	0.14	(10)	13.37	Virginia	2.42	0.11	(23)	-5.87
Massachusetts	2.31	0.10	(24)	-10.18	Washington	2.43	0.12	(21)	-5.45
Michigan	2.30	0.09	(25)	-10.58	West Virginia	2.03	0.18	(38)	-20.99 <sup>c</sup>
Minnesota	1.96 <sup>b</sup>	0.12	(39)	-23.70 <sup>c</sup>	Wisconsin	1.90 <sup>b</sup>	0.11	(42)	-26.22 <sup>c</sup>
Mississippi	3.06	0.19	(08)	19.11 <sup>c</sup>	Wyoming	2.09	0.38	(35)	-18.77 <sup>c</sup>
Missouri	1.93 <sup>b</sup>	0.10	(40)	-24.94 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 25.14  
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.24	0.01		
High Five States				
Missouri	0.34	0.05	(01)	40.32 <sup>c</sup>
California	0.33 <sup>b</sup>	0.02	(02)	37.37 <sup>c</sup>
Arkansas	0.32	0.07	(03)	34.52 <sup>c</sup>
Pennsylvania	0.30	0.03	(04)	24.90 <sup>c</sup>
North Carolina	0.30	0.04	(05)	24.78 <sup>c</sup>
Low Five States				
South Carolina	0.17	0.04	(26)	-31.11 <sup>c</sup>
Georgia	0.15	0.03	(27)	-36.26 <sup>c</sup>
Virginia	0.15	0.03	(28)	-37.29 <sup>c</sup>
Minnesota	0.14	0.03	(29)	-40.08 <sup>c</sup>
Massachusetts	0.13 <sup>b</sup>	0.03	(30)	-45.65 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.29	0.05	(07)	22.12 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.28	0.04	(10)	14.62	Nevada	0.30	0.07	(06)	24.62 <sup>c</sup>
Arkansas	0.32	0.07	(03)	34.52 <sup>c</sup>	New Hampshire	-	-	(-)	-
California	0.33 <sup>b</sup>	0.02	(02)	37.37 <sup>c</sup>	New Jersey	0.18	0.03	(23)	-25.06 <sup>c</sup>
Colorado	0.26	0.05	(12)	8.59	New Mexico	-	-	(-)	-
Connecticut	-	-	(-)	-	New York	0.19	0.02	(21)	-22.14 <sup>c</sup>
Delaware	-	-	(-)	-	North Carolina	0.30	0.04	(05)	24.78 <sup>c</sup>
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.23	0.02	(16)	-4.76	Ohio	0.21	0.03	(19)	-11.64
Georgia	0.15	0.03	(27)	-36.26 <sup>c</sup>	Oklahoma	0.29	0.06	(09)	19.64 <sup>c</sup>
Hawaii	-	-	(-)	-	Oregon	0.18	0.04	(24)	-26.34 <sup>c</sup>
Idaho	-	-	(-)	-	Pennsylvania	0.30	0.03	(04)	24.90 <sup>c</sup>
Illinois	0.25	0.03	(14)	4.82	Rhode Island	-	-	(-)	-
Indiana	0.18	0.03	(25)	-26.44 <sup>c</sup>	South Carolina	0.17	0.04	(26)	-31.11 <sup>c</sup>
Iowa	-	-	(-)	-	South Dakota	-	-	(-)	-
Kansas	-	-	(-)	-	Tennessee	0.29	0.04	(08)	20.78 <sup>c</sup>
Kentucky	0.23	0.05	(17)	-6.10	Texas	0.26	0.02	(13)	6.76
Louisiana	0.22	0.05	(18)	-10.19	Utah	-	-	(-)	-
Maine	-	-	(-)	-	Vermont	-	-	(-)	-
Maryland	0.27	0.04	(11)	11.73	Virginia	0.15	0.03	(28)	-37.29 <sup>c</sup>
Massachusetts	0.13 <sup>b</sup>	0.03	(30)	-45.65 <sup>c</sup>	Washington	0.20	0.04	(20)	-18.75 <sup>c</sup>
Michigan	0.24	0.03	(15)	-1.44	West Virginia	-	-	(-)	-
Minnesota	0.14	0.03	(29)	-40.08 <sup>c</sup>	Wisconsin	0.19	0.04	(22)	-22.91 <sup>c</sup>
Mississippi	-	-	(-)	-	Wyoming	-	-	(-)	-
Missouri	0.34	0.05	(01)	40.32 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.18  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.50	0.01		
High Five States				
Vermont	0.72	0.14	(01)	43.36 <sup>c</sup>
Idaho	0.67	0.10	(02)	34.09 <sup>c</sup>
Colorado	0.63	0.06	(03)	26.40 <sup>c</sup>
Wyoming	0.63	0.15	(04)	25.10 <sup>c</sup>
California	0.60 <sup>b</sup>	0.02	(05)	20.11 <sup>c</sup>
Low Five States				
Florida	0.42 <sup>b</sup>	0.02	(45)	-15.99 <sup>c</sup>
Nebraska	0.40	0.06	(46)	-21.21 <sup>c</sup>
Georgia	0.39	0.03	(47)	-21.84 <sup>c</sup>
South Carolina	0.36	0.04	(48)	-27.39 <sup>c</sup>
Montana	0.33	0.08	(49)	-35.14 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.43	0.04	(43)	-13.49	Montana	0.33	0.08	(49)	-35.14 <sup>c</sup>
Alaska	-	-	(-)	-	Nebraska	0.40	0.06	(46)	-21.21 <sup>c</sup>
Arizona	0.55	0.04	(10)	9.07	Nevada	0.45	0.06	(37)	-10.53
Arkansas	0.45	0.05	(38)	-10.71	New Hampshire	0.54	0.09	(12)	6.69
California	0.60 <sup>b</sup>	0.02	(05)	20.11 <sup>c</sup>	New Jersey	0.49	0.03	(26)	-1.89
Colorado	0.63	0.06	(03)	26.40 <sup>c</sup>	New Mexico	0.55	0.07	(09)	10.28
Connecticut	0.47	0.05	(30)	-5.60	New York	0.53	0.02	(14)	5.26
Delaware	0.49	0.10	(28)	-3.06	North Carolina	0.45	0.03	(39)	-10.78
D.C.	-	-	(-)	-	North Dakota	0.49	0.11	(25)	-1.61
Florida	0.42 <sup>b</sup>	0.02	(45)	-15.99 <sup>c</sup>	Ohio	0.53	0.03	(13)	5.62
Georgia	0.39	0.03	(47)	-21.84 <sup>c</sup>	Oklahoma	0.46	0.05	(34)	-7.65
Hawaii	0.60	0.09	(06)	20.10 <sup>c</sup>	Oregon	0.46	0.05	(33)	-7.59
Idaho	0.67	0.10	(02)	34.09 <sup>c</sup>	Pennsylvania	0.51	0.03	(18)	2.31
Illinois	0.50	0.03	(23)	-0.17	Rhode Island	0.43	0.08	(44)	-13.77
Indiana	0.46	0.04	(35)	-8.48	South Carolina	0.36	0.04	(48)	-27.39 <sup>c</sup>
Iowa	0.47	0.05	(31)	-6.21	South Dakota	0.50	0.11	(22)	0.06
Kansas	0.58	0.06	(07)	16.18 <sup>c</sup>	Tennessee	0.50	0.04	(24)	-0.23
Kentucky	0.51	0.05	(19)	1.81	Texas	0.52	0.02	(16)	3.25
Louisiana	0.46	0.05	(32)	-7.58	Utah	0.52	0.07	(17)	2.98
Maine	0.51	0.08	(20)	1.75	Vermont	0.72	0.14	(01)	43.36 <sup>c</sup>
Maryland	0.49	0.04	(27)	-2.57	Virginia	0.44	0.03	(41)	-11.39
Massachusetts	0.48	0.04	(29)	-4.91	Washington	0.56	0.04	(08)	12.33
Michigan	0.50	0.03	(21)	0.39	West Virginia	0.44	0.06	(42)	-13.14
Minnesota	0.54	0.04	(11)	7.00	Wisconsin	0.52	0.04	(15)	3.40
Mississippi	0.45	0.05	(40)	-10.80	Wyoming	0.63	0.15	(04)	25.10 <sup>c</sup>
Missouri	0.46	0.04	(36)	-8.78					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.19  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.49	0.01		
High Five States				
Kansas	0.71	0.11	(01)	43.42 <sup>c</sup>
Colorado	0.70	0.09	(02)	42.12 <sup>c</sup>
New Hampshire	0.67	0.15	(03)	36.44 <sup>c</sup>
Utah	0.62	0.12	(04)	24.70 <sup>c</sup>
Ohio	0.58	0.05	(05)	17.70 <sup>c</sup>
Low Five States				
South Carolina	0.41	0.07	(37)	-17.70 <sup>c</sup>
Mississippi	0.41	0.08	(38)	-17.75 <sup>c</sup>
Arkansas	0.39	0.07	(39)	-21.18 <sup>c</sup>
Nevada	0.36	0.08	(40)	-26.20 <sup>c</sup>
West Virginia	0.33	0.08	(41)	-32.89 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.50	0.07	(17)	1.22	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.43	0.10	(32)	-13.26
Arizona	0.53	0.06	(12)	6.30	Nevada	0.36	0.08	(40)	-26.20 <sup>c</sup>
Arkansas	0.39	0.07	(39)	-21.18 <sup>c</sup>	New Hampshire	0.67	0.15	(03)	36.44 <sup>c</sup>
California	0.56	0.03	(06)	12.27	New Jersey	0.49	0.05	(24)	-1.38
Colorado	0.70	0.09	(02)	42.12 <sup>c</sup>	New Mexico	0.55	0.11	(07)	11.62
Connecticut	0.49	0.08	(22)	-1.27	New York	0.54	0.04	(09)	8.39
Delaware	-	-	(-)	-	North Carolina	0.45	0.05	(28)	-8.35
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.43	0.03	(33)	-13.99	Ohio	0.58	0.05	(05)	17.70 <sup>c</sup>
Georgia	0.42	0.05	(35)	-15.01 <sup>c</sup>	Oklahoma	0.46	0.08	(27)	-6.36
Hawaii	-	-	(-)	-	Oregon	0.44	0.07	(30)	-10.00
Idaho	0.49	0.12	(23)	-1.29	Pennsylvania	0.47	0.04	(26)	-5.20
Illinois	0.41	0.04	(36)	-16.71 <sup>c</sup>	Rhode Island	-	-	(-)	-
Indiana	0.49	0.06	(21)	-0.36	South Carolina	0.41	0.07	(37)	-17.70 <sup>c</sup>
Iowa	0.54	0.08	(10)	8.23	South Dakota	-	-	(-)	-
Kansas	0.71	0.11	(01)	43.42 <sup>c</sup>	Tennessee	0.50	0.06	(18)	0.19
Kentucky	0.49	0.07	(20)	0.03	Texas	0.50	0.03	(16)	1.27
Louisiana	0.42	0.07	(34)	-14.12	Utah	0.62	0.12	(04)	24.70 <sup>c</sup>
Maine	0.49	0.12	(25)	-1.64	Vermont	-	-	(-)	-
Maryland	0.53	0.07	(11)	7.97	Virginia	0.51	0.06	(13)	3.13
Massachusetts	0.49	0.06	(19)	0.04	Washington	0.55	0.06	(08)	10.64
Michigan	0.51	0.05	(15)	2.70	West Virginia	0.33	0.08	(41)	-32.89 <sup>c</sup>
Minnesota	0.51	0.06	(14)	3.11	Wisconsin	0.43	0.06	(31)	-12.96
Mississippi	0.41	0.08	(38)	-17.75 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	0.45	0.06	(29)	-8.36					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 26.20  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.50	0.01		
High Five States				
Vermont	0.99	0.23	(01)	96.34 <sup>c</sup>
Idaho	0.80	0.14	(02)	58.62 <sup>c</sup>
Hawaii	0.79	0.14	(03)	57.51 <sup>c</sup>
California	0.64 <sup>b</sup>	0.03	(04)	26.94 <sup>c</sup>
Wisconsin	0.58	0.06	(05)	16.29 <sup>c</sup>
Low Five States				
Virginia	0.41	0.04	(40)	-19.26 <sup>c</sup>
Alabama	0.38	0.05	(41)	-24.29 <sup>c</sup>
Georgia	0.37	0.04	(42)	-27.00 <sup>c</sup>
Nebraska	0.33	0.08	(43)	-33.84 <sup>c</sup>
South Carolina	0.33	0.05	(44)	-34.44 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.38	0.05	(41)	-24.29 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.33	0.08	(43)	-33.84 <sup>c</sup>
Arizona	0.56	0.06	(08)	11.05	Nevada	0.52	0.09	(15)	3.00
Arkansas	0.49	0.07	(23)	-2.53	New Hampshire	0.44	0.11	(33)	-12.62
California	0.64 <sup>b</sup>	0.03	(04)	26.94 <sup>c</sup>	New Jersey	0.50	0.04	(20)	-1.10
Colorado	0.55	0.07	(09)	10.03	New Mexico	0.54	0.10	(11)	6.56
Connecticut	0.46	0.06	(30)	-8.75	New York	0.53	0.03	(13)	4.83
Delaware	-	-	(-)	-	North Carolina	0.44	0.04	(34)	-13.04
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.42 <sup>b</sup>	0.03	(38)	-17.44 <sup>c</sup>	Ohio	0.48	0.04	(25)	-3.82
Georgia	0.37	0.04	(42)	-27.00 <sup>c</sup>	Oklahoma	0.48	0.07	(26)	-5.40
Hawaii	0.79	0.14	(03)	57.51 <sup>c</sup>	Oregon	0.47	0.06	(28)	-6.36
Idaho	0.80	0.14	(02)	58.62 <sup>c</sup>	Pennsylvania	0.54	0.04	(10)	8.20
Illinois	0.56	0.04	(07)	12.36	Rhode Island	0.42	0.10	(36)	-16.12 <sup>c</sup>
Indiana	0.41	0.05	(39)	-17.52 <sup>c</sup>	South Carolina	0.33	0.05	(44)	-34.44 <sup>c</sup>
Iowa	0.42	0.06	(37)	-16.51 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.51	0.08	(18)	1.15	Tennessee	0.50	0.05	(21)	-1.17
Kentucky	0.51	0.06	(17)	1.53	Texas	0.53	0.03	(14)	4.68
Louisiana	0.49	0.06	(22)	-2.49	Utah	0.43	0.09	(35)	-14.20
Maine	0.51	0.11	(16)	2.37	Vermont	0.99	0.23	(01)	96.34 <sup>c</sup>
Maryland	0.45	0.05	(32)	-11.09	Virginia	0.41	0.04	(40)	-19.26 <sup>c</sup>
Massachusetts	0.47	0.05	(29)	-7.41	Washington	0.57	0.06	(06)	12.95
Michigan	0.48	0.04	(24)	-3.66	West Virginia	0.50	0.09	(19)	-0.63
Minnesota	0.53	0.06	(12)	5.68	Wisconsin	0.58	0.06	(05)	16.29 <sup>c</sup>
Mississippi	0.47	0.07	(27)	-5.70	Wyoming	-	-	(-)	-
Missouri	0.45	0.05	(31)	-10.23					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.18  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.36	0.02		
High Five States				
Maine	6.02 <sup>b</sup>	0.27	(01)	38.03 <sup>c</sup>
Vermont	5.89 <sup>b</sup>	0.41	(02)	35.03 <sup>c</sup>
Nevada	5.36 <sup>b</sup>	0.22	(03)	22.93 <sup>c</sup>
Rhode Island	5.27	0.29	(04)	20.66 <sup>c</sup>
Delaware	5.10	0.33	(05)	16.96 <sup>c</sup>
Low Five States				
Mississippi	3.61 <sup>b</sup>	0.16	(47)	-17.37 <sup>c</sup>
Alaska	3.45	0.44	(48)	-20.87 <sup>c</sup>
New Mexico	3.39 <sup>b</sup>	0.18	(49)	-22.22 <sup>c</sup>
Utah	3.36 <sup>b</sup>	0.19	(50)	-22.92 <sup>c</sup>
Hawaii	2.74 <sup>b</sup>	0.19	(51)	-37.29 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.81 <sup>b</sup>	0.12	(45)	-12.66	Montana	4.38	0.28	(23)	0.44
Alaska	3.45	0.44	(48)	-20.87 <sup>c</sup>	Nebraska	3.94	0.20	(41)	-9.73
Arizona	3.89 <sup>b</sup>	0.11	(42)	-10.90	Nevada	5.36 <sup>b</sup>	0.22	(03)	22.93 <sup>c</sup>
Arkansas	4.16	0.16	(32)	-4.76	New Hampshire	4.94	0.27	(09)	13.17
California	4.06 <sup>b</sup>	0.05	(37)	-6.97	New Jersey	4.96 <sup>b</sup>	0.10	(08)	13.57
Colorado	3.83 <sup>b</sup>	0.14	(44)	-12.33	New Mexico	3.39 <sup>b</sup>	0.18	(49)	-22.22 <sup>c</sup>
Connecticut	4.92 <sup>b</sup>	0.16	(10)	12.81	New York	4.38	0.06	(24)	0.33
Delaware	5.10	0.33	(05)	16.96 <sup>c</sup>	North Carolina	4.13	0.10	(33)	-5.30
D.C.	4.36	0.39	(26)	-0.20	North Dakota	4.29	0.33	(27)	-1.59
Florida	4.39	0.06	(22)	0.59	Ohio	5.02 <sup>b</sup>	0.09	(07)	15.08 <sup>c</sup>
Georgia	4.03	0.10	(38)	-7.70	Oklahoma	4.06	0.14	(36)	-6.87
Hawaii	2.74 <sup>b</sup>	0.19	(51)	-37.29 <sup>c</sup>	Oregon	4.73	0.15	(13)	8.43
Idaho	4.36	0.25	(25)	-0.12	Pennsylvania	4.71 <sup>b</sup>	0.08	(16)	7.95
Illinois	4.50	0.08	(21)	3.01	Rhode Island	5.27	0.29	(04)	20.66 <sup>c</sup>
Indiana	4.59	0.12	(18)	5.20	South Carolina	3.99	0.13	(40)	-8.68
Iowa	4.55	0.16	(20)	4.30	South Dakota	4.08	0.29	(35)	-6.52
Kansas	3.87	0.16	(43)	-11.40	Tennessee	4.23	0.12	(30)	-3.02
Kentucky	4.57	0.14	(19)	4.63	Texas	3.79 <sup>b</sup>	0.06	(46)	-13.22
Louisiana	4.27	0.14	(29)	-2.23	Utah	3.36 <sup>b</sup>	0.19	(50)	-22.92 <sup>c</sup>
Maine	6.02 <sup>b</sup>	0.27	(01)	38.03 <sup>c</sup>	Vermont	5.89 <sup>b</sup>	0.41	(02)	35.03 <sup>c</sup>
Maryland	4.72	0.13	(15)	8.06	Virginia	4.12	0.11	(34)	-5.55
Massachusetts	5.10 <sup>b</sup>	0.12	(06)	16.85 <sup>c</sup>	Washington	4.73	0.12	(14)	8.37
Michigan	4.82 <sup>b</sup>	0.10	(12)	10.35	West Virginia	4.68	0.20	(17)	7.21
Minnesota	4.02	0.12	(39)	-7.81	Wisconsin	4.83 <sup>b</sup>	0.12	(11)	10.61
Mississippi	3.61 <sup>b</sup>	0.16	(47)	-17.37 <sup>c</sup>	Wyoming	4.29	0.41	(28)	-1.73
Missouri	4.21	0.11	(31)	-3.43					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.19  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.66	0.04		
High Five States				
Vermont	10.94 <sup>b</sup>	0.89	(01)	42.80 <sup>c</sup>
Maine	10.33 <sup>b</sup>	0.57	(02)	34.84 <sup>c</sup>
Rhode Island	9.75	0.64	(03)	27.33 <sup>c</sup>
Massachusetts	9.12 <sup>b</sup>	0.26	(04)	19.12 <sup>c</sup>
Ohio	9.04 <sup>b</sup>	0.19	(05)	18.05 <sup>c</sup>
Low Five States				
Texas	6.65 <sup>b</sup>	0.13	(47)	-13.22
Utah	6.05 <sup>b</sup>	0.39	(48)	-21.05 <sup>c</sup>
Alaska	5.87	0.89	(49)	-23.29 <sup>c</sup>
New Mexico	5.83 <sup>b</sup>	0.37	(50)	-23.93 <sup>c</sup>
Hawaii	4.62 <sup>b</sup>	0.38	(51)	-39.67 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.77	0.27	(43)	-11.65	Montana	7.71	0.58	(24)	0.71
Alaska	5.87	0.89	(49)	-23.29 <sup>c</sup>	Nebraska	6.93	0.42	(41)	-9.55
Arizona	6.72 <sup>b</sup>	0.22	(45)	-12.19	Nevada	8.69	0.43	(09)	13.48
Arkansas	6.94	0.33	(40)	-9.35	New Hampshire	8.80	0.57	(07)	14.97
California	7.03 <sup>b</sup>	0.10	(38)	-8.26	New Jersey	8.82 <sup>b</sup>	0.22	(06)	15.19 <sup>c</sup>
Colorado	6.89	0.30	(42)	-10.09	New Mexico	5.83 <sup>b</sup>	0.37	(50)	-23.93 <sup>c</sup>
Connecticut	8.69	0.34	(10)	13.47	New York	7.83	0.14	(22)	2.27
Delaware	8.80	0.68	(08)	14.96	North Carolina	7.13	0.21	(37)	-6.93
D.C.	7.17	0.84	(35)	-6.36	North Dakota	7.81	0.68	(23)	1.99
Florida	7.35	0.12	(29)	-3.99	Ohio	9.04 <sup>b</sup>	0.19	(05)	18.05 <sup>c</sup>
Georgia	7.18	0.23	(34)	-6.25	Oklahoma	7.35	0.31	(30)	-4.02
Hawaii	4.62 <sup>b</sup>	0.38	(51)	-39.67 <sup>c</sup>	Oregon	8.20	0.31	(16)	7.05
Idaho	7.48	0.50	(28)	-2.29	Pennsylvania	8.45 <sup>b</sup>	0.17	(14)	10.40
Illinois	7.97	0.18	(20)	4.08	Rhode Island	9.75	0.64	(03)	27.33 <sup>c</sup>
Indiana	8.05	0.25	(19)	5.17	South Carolina	6.76	0.28	(44)	-11.74
Iowa	8.48	0.34	(12)	10.75	South Dakota	7.54	0.62	(27)	-1.52
Kansas	6.97	0.34	(39)	-9.00	Tennessee	7.65	0.25	(25)	-0.14
Kentucky	8.13	0.31	(18)	6.19	Texas	6.65 <sup>b</sup>	0.13	(47)	-13.22
Louisiana	7.30	0.30	(32)	-4.71	Utah	6.05 <sup>b</sup>	0.39	(48)	-21.05 <sup>c</sup>
Maine	10.33 <sup>b</sup>	0.57	(02)	34.84 <sup>c</sup>	Vermont	10.94 <sup>b</sup>	0.89	(01)	42.80 <sup>c</sup>
Maryland	8.18	0.29	(17)	6.82	Virginia	7.34	0.23	(31)	-4.16
Massachusetts	9.12 <sup>b</sup>	0.26	(04)	19.12 <sup>c</sup>	Washington	8.21	0.25	(15)	7.17
Michigan	8.61 <sup>b</sup>	0.21	(11)	12.45	West Virginia	7.94	0.42	(21)	3.74
Minnesota	7.24	0.26	(33)	-5.45	Wisconsin	8.47	0.27	(13)	10.62
Mississippi	6.70	0.35	(46)	-12.51	Wyoming	7.15	0.81	(36)	-6.67
Missouri	7.59	0.24	(26)	-0.88					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
 SE Standard error of the rate.  
 PD Percent difference between state rate and total U.S. rate.  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.20  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.19	0.02		
High Five States				
Maine	3.06	0.25	(01)	39.61 <sup>c</sup>
D.C.	2.81	0.40	(02)	27.96 <sup>c</sup>
Nevada	2.69	0.21	(03)	22.63 <sup>c</sup>
Massachusetts	2.64 <sup>b</sup>	0.11	(04)	20.23 <sup>c</sup>
Maryland	2.61	0.13	(05)	18.92 <sup>c</sup>
Low Five States				
South Dakota	1.69	0.25	(47)	-22.79 <sup>c</sup>
New Mexico	1.53 <sup>b</sup>	0.16	(48)	-30.27 <sup>c</sup>
Alaska	1.52	0.39	(49)	-30.59 <sup>c</sup>
Hawaii	1.38 <sup>b</sup>	0.18	(50)	-37.22 <sup>c</sup>
Utah	1.35 <sup>b</sup>	0.16	(51)	-38.37 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.95	0.11	(38)	-11.18	Montana	1.99	0.25	(35)	-9.20
Alaska	1.52	0.39	(49)	-30.59 <sup>c</sup>	Nebraska	1.99	0.18	(36)	-9.51
Arizona	1.69 <sup>b</sup>	0.10	(46)	-22.77 <sup>c</sup>	Nevada	2.69	0.21	(03)	22.63 <sup>c</sup>
Arkansas	2.27	0.16	(22)	3.30	New Hampshire	2.43	0.24	(13)	10.85
California	2.06	0.05	(32)	-6.24	New Jersey	2.56 <sup>b</sup>	0.09	(07)	16.72 <sup>c</sup>
Colorado	1.75 <sup>b</sup>	0.12	(44)	-20.17 <sup>c</sup>	New Mexico	1.53 <sup>b</sup>	0.16	(48)	-30.27 <sup>c</sup>
Connecticut	2.60	0.15	(06)	18.53 <sup>c</sup>	New York	2.24	0.06	(25)	2.02
Delaware	2.43	0.29	(14)	10.56	North Carolina	2.27	0.09	(21)	3.56
D.C.	2.81	0.40	(02)	27.96 <sup>c</sup>	North Dakota	1.80	0.28	(43)	-17.78 <sup>c</sup>
Florida	2.15	0.06	(28)	-1.81	Ohio	2.52 <sup>b</sup>	0.08	(09)	14.65
Georgia	2.12	0.10	(29)	-3.21	Oklahoma	1.84	0.13	(42)	-16.18 <sup>c</sup>
Hawaii	1.38 <sup>b</sup>	0.18	(50)	-37.22 <sup>c</sup>	Oregon	2.26	0.14	(23)	2.90
Idaho	2.01	0.22	(34)	-8.59	Pennsylvania	2.36	0.07	(17)	7.76
Illinois	2.37	0.08	(16)	7.83	Rhode Island	2.53	0.26	(08)	15.36 <sup>c</sup>
Indiana	2.35	0.11	(19)	7.05	South Carolina	2.17	0.13	(27)	-1.19
Iowa	2.06	0.14	(33)	-6.31	South Dakota	1.69	0.25	(47)	-22.79 <sup>c</sup>
Kansas	1.88	0.15	(40)	-14.18	Tennessee	2.06	0.10	(31)	-5.92
Kentucky	2.25	0.13	(24)	2.58	Texas	1.90 <sup>b</sup>	0.06	(39)	-13.57
Louisiana	2.37	0.14	(15)	7.94	Utah	1.35 <sup>b</sup>	0.16	(51)	-38.37 <sup>c</sup>
Maine	3.06	0.25	(01)	39.61 <sup>c</sup>	Vermont	2.44	0.34	(12)	11.31
Maryland	2.61	0.13	(05)	18.92 <sup>c</sup>	Virginia	2.07	0.10	(30)	-5.72
Massachusetts	2.64 <sup>b</sup>	0.11	(04)	20.23 <sup>c</sup>	Washington	2.33	0.11	(20)	6.06
Michigan	2.36	0.09	(18)	7.67	West Virginia	2.48	0.19	(11)	12.88
Minnesota	1.87	0.11	(41)	-14.60	Wisconsin	2.50	0.12	(10)	14.01
Mississippi	1.73	0.14	(45)	-21.17 <sup>c</sup>	Wyoming	2.19	0.39	(26)	-0.03
Missouri	1.98	0.10	(37)	-9.71					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.