

Table VI-23  
COLON AND RECTUM CANCER (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2003  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <13 <sup>e</sup>	0 to <28 <sup>e</sup>	>=28 <sup>g</sup>	Complete <sup>h</sup>
<u>Race/Sex</u>									
All Races <sup>b</sup>	416,089	238,065	165,688	108,697	61,523	762,462	1,012,722	55,476	1,068,198
Males	207,662	117,724	82,105	52,008	25,932	379,236	494,463	20,326	514,789
Females	208,427	120,341	83,583	56,689	35,591	383,226	518,259	35,150	553,409
White <sup>b</sup>	358,288	208,250	146,687	97,237	55,369	662,584	886,271	49,625	935,896
Males	180,179	103,668	73,262	46,809	23,467	331,729	435,544	18,337	453,881
Females	178,109	104,582	73,425	50,428	31,902	330,855	450,727	31,288	482,015
Black <sup>b</sup>	39,757	20,776	13,531	8,332	4,347	69,195	88,244	3,816	92,060
Males	17,689	9,203	5,991	3,451	1,585	30,968	38,475	1,289	39,764
Females	22,068	11,573	7,540	4,881	2,762	38,227	49,769	2,527	52,296
Asian/Pacific Islander <sup>c</sup>	12,355	6,306	+	+	+	21,142	+	+	+
Males	6,533	3,376	+	+	+	11,139	+	+	+
Females	5,822	2,930	+	+	+	10,003	+	+	+
Hispanic <sup>d</sup>	18,538	9,092	+	+	+	31,876	+	+	+
Males	10,067	4,834	+	+	+	17,043	+	+	+
Females	8,471	4,258	+	+	+	14,833	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2003, of the SEER 11 population diagnosed in the previous 10 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races <sup>c</sup>	0.2008%	-	0.0002%	0.0036%	0.0192%	0.0718%	0.2483%	0.6476%	1.2394%	1.6585%	0.2179%
Males	0.2037%	-	0.0002%	0.0036%	0.0201%	0.0748%	0.2828%	0.7802%	1.4976%	1.8458%	0.2544%
Females	0.1980%	-	0.0003%	0.0035%	0.0183%	0.0689%	0.2156%	0.5293%	1.0450%	1.5567%	0.1896%
White <sup>c</sup>	0.2122%	-	0.0003%	0.0033%	0.0186%	0.0684%	0.2366%	0.6446%	1.2561%	1.6855%	0.2174%
Males	0.2140%	-	0.0002%	0.0034%	0.0193%	0.0713%	0.2717%	0.7812%	1.5130%	1.8739%	0.2541%
Females	0.2105%	-	0.0003%	0.0033%	0.0178%	0.0655%	0.2024%	0.5196%	1.0592%	1.5846%	0.1883%
Black <sup>c</sup>	0.1552%	-	-	0.0038%	0.0186%	0.0871%	0.3176%	0.7551%	1.2116%	1.4315%	0.2275%
Males	0.1470%	-	-	0.0031%	0.0180%	0.0895%	0.3580%	0.8357%	1.3377%	1.4488%	0.2473%
Females	0.1626%	-	-	0.0044%	0.0192%	0.0850%	0.2841%	0.6919%	1.1288%	1.4233%	0.2135%
Asian/Pacific Islander <sup>c</sup>	0.1760%	-	-	0.0044%	0.0219%	0.0775%	0.2564%	0.5698%	1.1153%	1.5422%	0.2039%
Males	0.1918%	-	-	0.0054%	0.0259%	0.0835%	0.2851%	0.7035%	1.4523%	1.8013%	0.2480%
Females	0.1611%	-	-	0.0033%	0.0181%	0.0720%	0.2312%	0.4584%	0.8712%	1.3675%	0.1708%
Hispanic <sup>d</sup>	0.0700%	-	-	0.0032%	0.0147%	0.0482%	0.1741%	0.4576%	0.8753%	1.0560%	0.1508%
Males	0.0726%	-	-	0.0035%	0.0149%	0.0468%	0.1971%	0.5863%	1.1661%	1.2298%	0.1856%
Females	0.0673%	-	-	0.0027%	0.0145%	0.0496%	0.1529%	0.3513%	0.6677%	0.9583%	0.1251%

- <sup>a</sup> US 2003 cancer prevalence counts are based on 2003 cancer prevalence proportions from the SEER registries and 1/1/2003 US population estimates based on the average of 2002 and 2003 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- <sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia excluding Hawaii and Seattle.
- <sup>e</sup> Maximum limited-duration prevalence: 28 years for 1975-2003 SEER 9 data; 13 years for 1990-2003 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- <sup>f</sup> Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- <sup>g h</sup> (g) Cases diagnosed more than 28 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <28 and >=28.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.