

Table 1.21  
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2009<sup>a</sup>  
Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			34-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 34 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 34 Years <sup>d</sup>
All Sites	Both Sexes	4,357,341	4,442,261	4,864,827	12,096,978	12,304,118
	Male	2,246,570	2,279,828	2,484,619	5,717,465	5,784,750
	Female	2,110,771	2,162,433	2,380,208	6,379,513	6,519,368
Oral Cavity & Pharynx	Both Sexes	97,542	111,817	115,695	252,887	278,231
	Male	67,675	76,701	79,161	166,069	180,884
	Female	29,867	35,116	36,534	86,818	97,347
Esophagus	Both Sexes	20,356	24,677	24,737	31,710	37,085
	Male	15,922	19,308	19,356	24,602	28,647
	Female	4,434	5,369	5,381	7,108	8,438
Stomach	Both Sexes	34,419	40,930	41,143	68,386	78,221
	Male	20,553	24,756	24,854	38,804	44,783
	Female	13,866	16,174	16,289	29,582	33,438
Colon & Rectum	Both Sexes	405,797	469,194	478,203	1,111,907	1,239,920
	Male	205,558	237,798	242,129	548,761	608,653
	Female	200,239	231,396	236,074	563,146	631,267
Liver & Intrahepatic Bile Duct	Both Sexes	26,017	29,875	29,931	35,119	39,545
	Male	18,881	21,432	21,488	24,549	27,385
	Female	7,136	8,443	8,443	10,570	12,160
Pancreas	Both Sexes	28,875	35,140	35,140	37,971	45,129
	Male	14,376	17,768	17,768	18,446	22,163
	Female	14,499	17,372	17,372	19,525	22,966
Larynx	Both Sexes	30,294	36,780	37,134	86,894	97,654
	Male	24,153	29,235	29,544	69,765	77,999
	Female	6,141	7,545	7,590	17,129	19,655
Lung & Bronchus	Both Sexes	224,508	284,065	291,615	379,223	462,713
	Male	104,519	133,787	136,648	173,619	212,377
	Female	119,989	150,278	154,967	205,604	250,336
Melanoma of the Skin	Both Sexes	263,883	297,561	309,968	837,215	901,518
	Male	140,746	162,207	170,141	415,003	452,508
	Female	123,137	135,354	139,827	422,212	449,010
Breast	Female	818,223	883,935	937,301	2,681,040	2,841,885
Cervix	Female	37,983	39,767	39,882	202,962	208,595
Corpus & Uterus, NOS	Female	160,329	180,638	180,713	553,870	603,503
Ovary <sup>f</sup>	Female	55,497	64,242	64,294	164,877	185,160

<sup>a</sup> U.S. 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER 9 registries and 1/1/2009 U.S. population estimates based on the average of 2008 and 2009 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 34 years (1975-2008)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2004-2008)  
For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2004; Melanoma in 2005.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 34-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2009.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 34-year limited duration prevalence. The 2004 breast cancer is counted for the breast 5-year and 34-year limited duration prevalence.

In method (e) the 2004 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2005 melanoma is counted for 5-year limited duration prevalence for melanoma.

<sup>f</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.21 - continued  
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2009<sup>a</sup>  
 Using Different Tumor Inclusion Criteria<sup>b</sup>

Site	Sex	5-Year Limited Duration			34-year Limited Duration	
		1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 34 Years <sup>d</sup>	1st Per Site in Previous 5 Years <sup>e</sup>	1st Invasive Tumor Ever <sup>c</sup>	1st Per Site in Previous 34 Years <sup>d</sup>
Prostate	Male	991,610	1,069,850	1,069,879	2,494,988	2,659,051
Testis	Male	41,292	42,052	42,698	194,935	197,533
Urinary Bladder	Both Sexes	196,403	243,253	248,175	540,518	626,417
	Male	148,936	185,112	189,064	402,281	466,103
	Female	47,467	58,141	59,111	138,237	160,314
Kidney & Renal Pelvis	Both Sexes	137,327	166,272	168,023	309,168	359,955
	Male	82,513	101,559	102,846	182,285	214,227
	Female	54,814	64,713	65,177	126,883	145,728
Brain & Nervous System	Both Sexes	42,492	44,665	45,128	121,848	125,138
	Male	23,270	24,518	24,750	64,666	66,457
	Female	19,222	20,147	20,378	57,182	58,681
Thyroid	Both Sexes	150,696	166,265	166,781	456,180	484,996
	Male	33,895	39,037	39,179	99,597	108,060
	Female	116,801	127,228	127,602	356,583	376,936
Hodgkin Lymphoma	Both Sexes	37,947	40,392	40,402	161,071	165,355
	Male	20,359	21,735	21,735	83,458	85,731
	Female	17,588	18,657	18,667	77,613	79,624
Non-Hodgkin Lymphoma	Both Sexes	193,568	226,608	228,204	474,455	529,222
	Male	101,809	120,766	121,658	246,926	276,020
	Female	91,759	105,842	106,546	227,529	253,202
Myeloma	Both Sexes	45,747	53,626	53,674	70,950	81,089
	Male	24,230	28,956	28,992	38,319	44,359
	Female	21,517	24,670	24,682	32,631	36,730
Leukemia	Both Sexes	108,472	124,139	124,315	266,026	291,521
	Male	62,188	71,959	72,042	150,062	165,245
	Female	46,284	52,180	52,273	115,964	126,276
Acute Lymphocytic Leukemia	Both Sexes	15,984	16,396	16,396	62,732	63,353
	Male	9,278	9,448	9,448	34,732	34,950
	Female	6,706	6,948	6,948	28,000	28,403
Childhood (Ages 0-19)	Both Sexes	61,498	61,622	62,135	288,648	289,141
	Male	32,966	33,016	33,253	148,843	149,048
	Female	28,532	28,606	28,882	139,805	140,093
Kaposi Sarcoma	Both Sexes	7,129	7,546	7,546	23,623	24,764
	Male	6,715	7,047	7,047	22,336	23,276
	Female	414	499	499	1,287	1,488
Mesothelioma	Both Sexes	3,040	3,957	3,957	4,570	5,673
	Male	2,095	2,759	2,759	2,694	3,484
	Female	945	1,198	1,198	1,876	2,189

<sup>a</sup> U.S. 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER 9 registries and 1/1/2009 U.S. population estimates based on the average of 2008 and 2009 population estimates from the U.S. Bureau of the Census.

<sup>b</sup> Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

<sup>c</sup> (c) First invasive tumor ever

<sup>d</sup> (d) First invasive tumor for each cancer site diagnosed during the previous 34 years (1975-2008)

<sup>e</sup> (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2004-2008)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2004; Melanoma in 2005.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 34-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2009.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 34-year limited duration prevalence. The 2004 breast cancer is counted for the breast 5-year and 34-year limited duration prevalence.

In method (e) the 2004 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2005 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.22  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2009<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	5,809,017	18,156	41,323	80,500	150,710	346,280	815,636	1,465,844	2,890,567
Females	6,740,016	15,649	35,203	83,134	219,550	622,449	1,226,028	1,600,054	2,937,950
Oral Cavity & Pharynx									
Males	172,502	67	418	1,505	3,416	14,153	41,881	49,752	61,309
Females	91,610	101	493	1,453	3,530	8,334	17,445	21,205	39,049
Esophagus									
Males	24,657	0	0	38	186	1,059	4,426	8,360	10,588
Females	7,170	0	0	0	39	314	1,070	1,656	4,091
Stomach									
Males	39,694	0	17	78	688	2,436	6,519	9,680	20,275
Females	30,232	4	11	165	643	2,169	4,233	5,764	17,243
Colon & Rectum									
Males	558,408	11	69	1,116	5,277	24,270	77,071	129,789	320,804
Females	581,302	0	86	1,202	5,090	22,495	64,292	107,339	380,797
Liver & Intrahep									
Males	24,615	438	554	494	438	1,581	8,481	7,111	5,516
Females	10,913	466	435	332	473	786	2,384	2,609	3,430
Pancreas									
Males	18,537	0	23	68	321	1,356	4,100	5,731	6,939
Females	19,649	0	47	135	418	1,377	3,454	5,181	9,036
Larynx									
Males	71,550	0	0	66	221	2,095	10,475	20,267	38,425
Females	17,521	0	0	48	143	1,051	3,163	4,474	8,642
Lung & Bronchus									
Males	178,447	45	80	454	1,086	5,587	24,209	52,348	94,640
Females	209,203	34	47	363	1,291	7,893	27,437	57,808	114,331
Melanoma of the Skin									
Males	427,519	72	764	5,002	17,503	47,272	91,489	111,843	153,573
Females	448,192	95	950	10,770	33,163	72,253	104,576	99,252	127,133

<sup>a</sup> U.S. 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2009 U.S. population estimates based on the average of 2008 and 2009 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.22 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2009<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	14,303	0	0	22	103	498	2,093	3,896	7,691
Females	2,747,209	0	86	2,487	35,584	212,685	522,225	733,979	1,240,161
Cervix									
Females	247,556	0	68	2,050	16,434	42,782	59,350	52,979	73,893
Corpus & Uterus, NOS									
Females	589,789	0	56	412	5,572	24,294	82,675	148,774	328,006
Ovary <sup>d</sup>									
Females	182,710	66	932	3,503	6,560	19,171	39,536	45,524	67,418
Prostate									
Males	2,496,420	34	69	89	247	19,652	216,595	687,644	1,572,090
Urinary Bladder									
Males	411,161	56	110	595	2,405	11,643	42,320	94,909	259,122
Females	143,089	34	56	162	1,087	4,412	13,373	29,661	94,303
Kidney & Renal Pelvis									
Males	188,341	1,507	2,373	2,601	4,936	16,313	37,165	53,360	70,087
Females	131,799	1,408	2,607	2,769	4,642	11,613	22,637	30,925	55,198
Hodgkin Lymphoma									
Males	90,420	168	2,316	9,264	16,301	23,466	20,120	12,217	6,567
Females	84,475	146	1,642	9,767	16,451	22,548	18,024	9,302	6,597
Non-Hodgkin Lymphoma									
Males	252,012	810	3,472	7,162	12,801	28,228	49,236	62,729	87,574
Females	232,051	387	1,793	4,203	8,302	19,927	39,219	53,951	104,268
Myeloma									
Males	38,403	0	6	50	485	2,628	7,535	12,343	15,356
Females	32,754	0	0	22	276	2,112	5,612	9,822	14,911
Leukemia									
Males	152,597	6,389	12,636	11,647	10,425	12,678	20,796	29,694	48,332
Females	119,096	5,232	10,502	10,167	9,042	9,418	13,463	19,494	41,777
Acute Lymphocytic Leuk									
Males	36,576	5,244	10,969	9,000	5,888	3,114	1,140	773	448
Females	29,614	4,381	8,829	7,398	4,995	2,241	767	687	315

<sup>a</sup> U.S. 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2009 U.S. population estimates based on the average of 2008 and 2009 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21  
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	4,357,341	3,022,968	2,016,342	1,278,550	738,833	437,465	10,479,611	12,096,978	456,359	12,553,337
	Males	2,246,570	1,519,767	947,022	551,691	250,351	137,162	5,193,308	5,717,465	93,632	5,811,097
	Females	2,110,771	1,503,201	1,069,320	726,859	488,482	300,303	5,286,303	6,379,513	362,727	6,742,240
White <sup>b</sup>	Both Sexes	3,710,739	2,639,002	1,778,231	1,149,321	671,706	398,911	9,099,659	10,571,450	401,052	10,972,502
	Males	1,905,553	1,317,934	830,086	498,591	230,018	126,935	4,486,216	4,968,978	87,505	5,056,483
	Females	1,805,186	1,321,068	948,145	650,730	441,688	271,976	4,613,443	5,602,472	313,547	5,916,019
Black <sup>b</sup>	Both Sexes	428,651	261,596	160,625	86,590	43,078	25,102	925,760	1,019,358	26,577	1,045,935
	Males	230,263	144,925	83,740	37,304	12,944	6,229	492,390	518,162	2,449	520,611
	Females	198,388	116,671	76,885	49,286	30,134	18,873	433,370	501,196	24,128	525,324
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	125,291	76,487	46,393	+	+	+	270,475	+	+	+
	Males	54,838	31,418	17,398	+	+	+	111,227	+	+	+
	Females	70,453	45,069	28,995	+	+	+	159,248	+	+	+
Hispanic <sup>d</sup>	Both Sexes	282,399	171,977	105,097	+	+	+	613,843	+	+	+
	Males	135,959	80,564	46,453	+	+	+	284,175	+	+	+
	Females	146,440	91,413	58,644	+	+	+	329,668	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	2.7602%	0.0796%	0.1656%	0.2755%	0.6903%	1.6735%	3.8785%	8.5940%	13.7507%	14.2298%	2.7820%
	Males	2.7630%	0.0844%	0.1757%	0.2615%	0.5299%	1.1401%	3.2133%	9.4824%	18.2603%	20.8174%	3.1605%
	Females	2.7573%	0.0745%	0.1550%	0.2906%	0.8557%	2.2104%	4.5168%	7.7888%	10.1655%	10.5548%	2.5603%
White <sup>c</sup>	Both Sexes	2.9986%	0.0856%	0.1803%	0.3057%	0.7649%	1.7672%	4.0010%	8.9139%	14.3847%	14.6321%	2.9022%
	Males	2.9706%	0.0909%	0.1922%	0.2912%	0.5991%	1.2171%	3.2802%	9.6207%	18.7445%	21.0246%	3.2420%
	Females	3.0268%	0.0801%	0.1677%	0.3215%	0.9437%	2.3423%	4.7170%	8.2543%	10.8365%	11.0719%	2.7132%
Black <sup>c</sup>	Both Sexes	2.0532%	0.0580%	0.1143%	0.1710%	0.4680%	1.3534%	3.7855%	9.1017%	13.5146%	13.0556%	2.6524%
	Males	2.3127%	0.0627%	0.1146%	0.1562%	0.3238%	0.9595%	3.8859%	12.6398%	21.1312%	22.4651%	3.6134%
	Females	1.8172%	0.0532%	0.1140%	0.1855%	0.5948%	1.6978%	3.7015%	6.3897%	8.2741%	8.6546%	2.0550%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1.8421%	0.0631%	0.1247%	0.1789%	0.4388%	1.2640%	2.8924%	5.3976%	8.8455%	10.2471%	1.8928%
	Males	1.6288%	0.0670%	0.1341%	0.1627%	0.2739%	0.6629%	1.7853%	5.0698%	11.0754%	14.8580%	1.9222%
	Females	2.0420%	0.0590%	0.1148%	0.1952%	0.5948%	1.8144%	3.8568%	5.6762%	7.1561%	7.3868%	1.9287%
Hispanic <sup>d</sup>	Both Sexes	1.1836%	0.0738%	0.1603%	0.2112%	0.4946%	1.1892%	2.7498%	6.0169%	9.8943%	9.8695%	1.9915%
	Males	1.0643%	0.0793%	0.1709%	0.2076%	0.3635%	0.6894%	1.9666%	6.3427%	13.6086%	14.8680%	2.2274%
	Females	1.3103%	0.0680%	0.1491%	0.2154%	0.6492%	1.7384%	3.5093%	5.7350%	7.1099%	6.9431%	1.8931%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.21  
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	42,492	25,410	18,285	14,945	10,499	6,075	98,707	121,848	13,554	135,402
	Males	23,270	12,699	9,934	8,317	5,250	3,116	52,790	64,666	6,159	70,825
	Females	19,222	12,711	8,351	6,628	5,249	2,959	45,917	57,182	7,395	64,577
White <sup>b</sup>	Both Sexes	37,122	22,504	15,887	13,207	9,405	5,386	86,549	107,288	5,378	112,666
	Males	20,567	11,403	8,670	7,311	4,640	2,749	46,651	57,269	3,089 <sup>i</sup>	60,358
	Females	16,555	11,101	7,217	5,896	4,765	2,637	39,898	50,019	2,289 <sup>i</sup>	52,308
Black <sup>b</sup>	Both Sexes	3,321	1,874	1,505	1,206	746	546	7,703	9,458	+	+
	Males	1,632	842	806	682	405	290	3,862	4,759	+	+
	Females	1,689	1,032	699	524	341	256	3,841	4,699	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,179	677	504	+	+	+	2,649	+	+	+
	Males	594	333	225	+	+	+	1,326	+	+	+
	Females	585	344	279	+	+	+	1,323	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,251	2,773	1,752	+	+	+	9,900	+	+	+
	Males	2,385	1,417	881	+	+	+	5,242	+	+	+
	Females	1,866	1,356	871	+	+	+	4,658	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0261%	0.0134%	0.0299%	0.0240%	0.0247%	0.0319%	0.0322%	0.0319%	0.0257%	0.0155%	0.0260%
	Males	0.0282%	0.0138%	0.0320%	0.0241%	0.0269%	0.0343%	0.0365%	0.0355%	0.0300%	0.0160%	0.0281%
	Females	0.0240%	0.0129%	0.0276%	0.0238%	0.0225%	0.0296%	0.0281%	0.0286%	0.0223%	0.0152%	0.0240%
White <sup>c</sup>	Both Sexes	0.0292%	0.0145%	0.0330%	0.0267%	0.0282%	0.0367%	0.0360%	0.0346%	0.0288%	0.0165%	0.0291%
	Males	0.0317%	0.0151%	0.0359%	0.0268%	0.0309%	0.0394%	0.0405%	0.0384%	0.0345%	0.0167%	0.0315%
	Females	0.0266%	0.0139%	0.0300%	0.0266%	0.0253%	0.0339%	0.0316%	0.0310%	0.0242%	0.0164%	0.0267%
Black <sup>c</sup>	Both Sexes	0.0165%	0.0097%	0.0217%	0.0175%	0.0140%	0.0164%	0.0196%	0.0222%	0.0116%	0.0128%	0.0164%
	Males	0.0166%	0.0100%	0.0215%	0.0162%	0.0142%	0.0159%	0.0229%	0.0227%	0.0104%	-	0.0165%
	Females	0.0165%	0.0094%	0.0219%	0.0187%	0.0138%	0.0169%	0.0168%	0.0218%	0.0125%	0.0140%	0.0164%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0153%	0.0099%	0.0184%	0.0134%	0.0146%	0.0172%	0.0185%	0.0202%	0.0141%	0.0073%	0.0154%
	Males	0.0154%	0.0101%	0.0180%	0.0137%	0.0140%	0.0166%	0.0200%	0.0227%	0.0106%	0.0114%	0.0155%
	Females	0.0152%	0.0096%	0.0189%	0.0131%	0.0151%	0.0177%	0.0173%	0.0180%	0.0168%	-	0.0154%
Hispanic <sup>d</sup>	Both Sexes	0.0186%	0.0100%	0.0229%	0.0145%	0.0164%	0.0260%	0.0275%	0.0255%	0.0240%	0.0115%	0.0199%
	Males	0.0191%	0.0103%	0.0246%	0.0138%	0.0168%	0.0268%	0.0298%	0.0254%	0.0295%	-	0.0207%
	Females	0.0180%	0.0097%	0.0210%	0.0153%	0.0159%	0.0251%	0.0252%	0.0256%	0.0199%	0.0146%	0.0191%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.25  
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	823,927	683,519	495,838	325,573	202,835	105,335	2,275,147	2,695,153	66,596	2,761,749
	Males	5,704	3,906	1,987	1,113	770	450	12,578	14,113	177	14,290
	Females	818,223	679,613	493,851	324,460	202,065	104,885	2,262,569	2,681,040	66,419	2,747,459
White <sup>b</sup>	Both Sexes	702,358	601,788	440,973	292,285	184,628	95,438	1,988,799	2,370,711	58,903	2,429,614
	Males	4,917	3,269	1,779	989	689	417	10,835	12,242	10 <sup>i</sup>	12,252
	Females	697,441	598,519	439,194	291,296	183,939	95,021	1,977,964	2,358,469	58,893	2,417,362
Black <sup>b</sup>	Both Sexes	81,239	54,142	36,708	22,978	12,847	7,074	191,456	218,378	+	+
	Males	630	503	121	98	73	25	1,339	1,449	+	+
	Females	80,609	53,639	36,587	22,880	12,774	7,049	190,117	216,929	4,916	221,845
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	29,486	21,038	13,310	+	+	+	70,442	+	+	+
	Males	121	79	43	+	+	+	270	+	+	+
	Females	29,365	20,959	13,267	+	+	+	70,172	+	+	+
Hispanic <sup>d</sup>	Both Sexes	54,105	36,800	23,230	+	+	+	126,296	+	+	+
	Males	265	130	58	+	+	+	480	+	+	+
	Females	53,840	36,670	23,172	+	+	+	125,816	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.5971%	-	0.0001%	0.0059%	0.0845%	0.4516%	1.1102%	1.9633%	2.4851%	2.8097%	0.5888%
	Males	0.0067%	-	-	-	0.0005%	0.0024%	0.0092%	0.0236%	0.0424%	0.0588%	0.0077%
	Females	1.1782%	-	0.0003%	0.0121%	0.1712%	0.9038%	2.1668%	3.7213%	4.4271%	4.3442%	1.0826%
White <sup>c</sup>	Both Sexes	0.6485%	-	0.0001%	0.0054%	0.0834%	0.4512%	1.1325%	2.0657%	2.6495%	2.9878%	0.6138%
	Males	0.0072%	-	-	-	0.0005%	0.0021%	0.0087%	0.0244%	0.0455%	0.0626%	0.0079%
	Females	1.2926%	-	0.0003%	0.0111%	0.1729%	0.9206%	2.2489%	3.9711%	4.7687%	4.6169%	1.1415%
Black <sup>c</sup>	Both Sexes	0.4237%	-	-	0.0085%	0.0919%	0.4472%	1.0146%	1.7019%	2.0725%	2.4305%	0.5243%
	Males	0.0063%	-	-	-	-	0.0039%	0.0165%	0.0351%	0.0371%	0.0586%	0.0093%
	Females	0.8033%	-	-	0.0166%	0.1724%	0.8348%	1.8486%	2.9796%	3.4730%	3.5399%	0.9013%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.4647%	-	-	0.0067%	0.0827%	0.4508%	1.0427%	1.4793%	1.7363%	1.6825%	0.4642%
	Males	0.0038%	-	-	-	-	0.0022%	0.0067%	0.0091%	0.0262%	0.0281%	0.0044%
	Females	0.8966%	-	-	0.0132%	0.1602%	0.8616%	1.9451%	2.7291%	3.0319%	2.7088%	0.8413%
Hispanic <sup>d</sup>	Both Sexes	0.2444%	-	-	0.0043%	0.0627%	0.3187%	0.8137%	1.3733%	1.6813%	1.6725%	0.4071%
	Males	0.0018%	-	-	-	-	0.0011%	0.0042%	0.0133%	0.0253%	0.0206%	0.0039%
	Females	0.5023%	-	-	0.0091%	0.1360%	0.6677%	1.5987%	2.5499%	2.9228%	2.6397%	0.7519%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.15  
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	37,983	35,018	36,166	32,413	25,994	20,081	135,337	202,962	44,749	247,711
White <sup>b</sup>	29,788	28,023	29,009	27,086	21,874	16,439	108,627	164,735	37,738	202,473
Black <sup>b</sup>	5,475	4,723	4,389	3,495	2,827	2,717	17,446	25,791	+	+
Asian/Pacific Islander <sup>c</sup>	1,997	1,688	1,758	+	+	+	6,454	+	+	+
Hispanic <sup>d</sup>	8,040	7,423	6,068	+	+	+	25,845	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0778%	-	0.0002%	0.0088%	0.0794%	0.1569%	0.1623%	0.1393%	0.1241%	0.0734%	0.0755%
White <sup>c</sup>	0.0791%	-	-	0.0098%	0.0898%	0.1684%	0.1601%	0.1292%	0.1128%	0.0585%	0.0769%
Black <sup>c</sup>	0.0679%	-	-	0.0088%	0.0599%	0.1261%	0.1624%	0.1506%	0.1418%	0.1364%	0.0718%
Asian/Pacific Islander <sup>c</sup>	0.0743%	-	-	0.0022%	0.0377%	0.1088%	0.1670%	0.1839%	0.1752%	0.1399%	0.0697%
Hispanic <sup>d</sup>	0.0956%	-	-	0.0098%	0.0894%	0.2266%	0.2911%	0.2977%	0.2722%	0.1891%	0.1266%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 6.28  
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	405,797	278,550	177,574	117,163	74,383	39,813	959,520	1,111,907	28,254	1,140,161
	Males	205,558	139,486	87,277	57,095	35,600	16,670	480,383	548,761	9,923	558,684
	Females	200,239	139,064	90,297	60,068	38,783	23,143	479,137	563,146	18,331	581,477
White <sup>b</sup>	Both Sexes	338,421	239,116	153,599	102,486	65,848	35,380	816,532	951,501	24,954	976,455
	Males	173,886	121,152	76,050	50,366	31,868	14,969	413,436	474,638	8,886	483,524
	Females	164,535	117,964	77,549	52,120	33,980	20,411	403,096	476,863	16,068	492,931
Black <sup>b</sup>	Both Sexes	44,714	26,111	16,145	10,114	6,032	3,039	95,277	107,424	2,082	109,506
	Males	19,971	11,222	6,952	4,459	2,419	1,012	41,918	46,467	654	47,121
	Females	24,743	14,889	9,193	5,655	3,613	2,027	53,359	60,957	1,428	62,385
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	16,130	9,554	5,932	+	+	+	34,375	+	+	+
	Males	8,366	4,903	3,077	+	+	+	17,691	+	+	+
	Females	7,764	4,651	2,855	+	+	+	16,684	+	+	+
Hispanic <sup>d</sup>	Both Sexes	27,741	14,273	8,149	+	+	+	54,499	+	+	+
	Males	14,957	7,449	4,267	+	+	+	28,852	+	+	+
	Females	12,784	6,824	3,882	+	+	+	25,647	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.2579%	-	0.0003%	0.0049%	0.0236%	0.0976%	0.3206%	0.7365%	1.4325%	2.0351%	0.2611%
	Males	0.2627%	-	0.0002%	0.0049%	0.0240%	0.1026%	0.3564%	0.8718%	1.7062%	2.3123%	0.3023%
	Females	0.2531%	-	0.0003%	0.0049%	0.0231%	0.0926%	0.2862%	0.6138%	1.2149%	1.8804%	0.2286%
White <sup>c</sup>	Both Sexes	0.2675%	-	0.0003%	0.0051%	0.0234%	0.0944%	0.3039%	0.7120%	1.4327%	2.0561%	0.2575%
	Males	0.2725%	-	-	0.0050%	0.0242%	0.0973%	0.3413%	0.8496%	1.7075%	2.3421%	0.2992%
	Females	0.2624%	-	0.0003%	0.0052%	0.0226%	0.0914%	0.2667%	0.5835%	1.2090%	1.8967%	0.2240%
Black <sup>c</sup>	Both Sexes	0.2122%	-	-	0.0035%	0.0222%	0.1070%	0.3910%	0.9132%	1.4723%	2.0218%	0.2860%
	Males	0.2009%	-	-	0.0033%	0.0201%	0.1150%	0.4131%	1.0387%	1.6332%	2.1002%	0.3118%
	Females	0.2224%	-	-	0.0038%	0.0240%	0.1000%	0.3726%	0.8170%	1.3615%	1.9851%	0.2679%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.2429%	-	-	0.0042%	0.0236%	0.1007%	0.3413%	0.7264%	1.3643%	1.8564%	0.2532%
	Males	0.2566%	-	-	0.0049%	0.0252%	0.1162%	0.3763%	0.8578%	1.6824%	2.1566%	0.2990%
	Females	0.2301%	-	-	0.0035%	0.0221%	0.0865%	0.3109%	0.6146%	1.1233%	1.6703%	0.2182%
Hispanic <sup>d</sup>	Both Sexes	0.1059%	-	-	0.0034%	0.0180%	0.0739%	0.2524%	0.6065%	1.0854%	1.3485%	0.1973%
	Males	0.1080%	-	-	0.0037%	0.0191%	0.0715%	0.2745%	0.7299%	1.3883%	1.5992%	0.2358%
	Females	0.1037%	-	-	0.0029%	0.0168%	0.0764%	0.2309%	0.4998%	0.8584%	1.2017%	0.1684%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.15  
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	160,329	121,079	94,055	67,211	47,637	35,784	431,150	553,870	36,017	589,887
White <sup>b</sup>	139,264	108,689	86,061	62,792	44,595	33,829	386,000	501,593	30,986	532,579
Black <sup>b</sup>	12,241	6,720	4,059	2,419	1,685	908	25,012	28,720	949	29,669
Asian/Pacific Islander <sup>c</sup>	5,612	3,620	2,427	+	+	+	12,824	+	+	+
Hispanic <sup>d</sup>	10,768	6,689	4,352	+	+	+	24,394	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.2174%	-	-	0.0026%	0.0277%	0.1077%	0.3531%	0.8100%	0.9335%	0.7695%	0.2001%
White <sup>c</sup>	0.2447%	-	-	0.0023%	0.0271%	0.1081%	0.3703%	0.8821%	1.0384%	0.8450%	0.2159%
Black <sup>c</sup>	0.1069%	-	-	0.0032%	0.0229%	0.0682%	0.1787%	0.5005%	0.6473%	0.4661%	0.1248%
Asian/Pacific Islander <sup>c</sup>	0.1630%	-	-	0.0036%	0.0298%	0.1307%	0.3918%	0.5721%	0.4937%	0.3731%	0.1511%
Hispanic <sup>d</sup>	0.0962%	-	-	0.0039%	0.0346%	0.1115%	0.2843%	0.5731%	0.5612%	0.4089%	0.1436%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.21  
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	20,356	6,602	2,809	1,252	483	146	30,930	31,710	151	31,861
	Males	15,922	5,119	2,150	926	314	123	24,044	24,602	77	24,679
	Females	4,434	1,483	659	326	169	23	6,886	7,108	74	7,182
White <sup>b</sup>	Both Sexes	18,259	5,939	2,457	1,039	418	132	27,621	28,306	123	28,429
	Males	14,415	4,756	1,957	761	274	109	21,828	22,320	71	22,391
	Females	3,844	1,183	500	278	144	23	5,793	5,986	52	6,038
Black <sup>b</sup>	Both Sexes	1,547	478	287	173	57	14	2,473	2,556	34	2,590
	Males	1,046	242	151	134	39	14	1,561	1,627	10	1,637
	Females	501	236	136	39	18	0	912	929	24	953
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	403	108	46	+	+	+	589	+	+	+
	Males	329	70	26	+	+	+	451	+	+	+
	Females	74	38	20	+	+	+	138	+	+	+
Hispanic <sup>d</sup>	Both Sexes	763	205	113	+	+	+	1,160	+	+	+
	Males	628	173	86	+	+	+	961	+	+	+
	Females	135	32	27	+	+	+	199	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0080%	-	-	-	0.0005%	0.0027%	0.0115%	0.0310%	0.0476%	0.0367%	0.0081%
	Males	0.0125%	-	-	-	0.0008%	0.0041%	0.0191%	0.0539%	0.0791%	0.0704%	0.0138%
	Females	0.0036%	-	-	-	0.0002%	0.0012%	0.0043%	0.0102%	0.0225%	0.0179%	0.0034%
White <sup>c</sup>	Both Sexes	0.0091%	-	-	-	0.0007%	0.0029%	0.0125%	0.0338%	0.0519%	0.0389%	0.0088%
	Males	0.0143%	-	-	-	0.0011%	0.0046%	0.0205%	0.0591%	0.0870%	0.0767%	0.0151%
	Females	0.0039%	-	-	-	-	0.0012%	0.0046%	0.0102%	0.0234%	0.0179%	0.0034%
Black <sup>c</sup>	Both Sexes	0.0052%	-	-	-	-	0.0023%	0.0114%	0.0298%	0.0374%	0.0198%	0.0066%
	Males	0.0070%	-	-	-	-	0.0028%	0.0171%	0.0463%	0.0514%	0.0379%	0.0100%
	Females	0.0036%	-	-	-	-	0.0019%	0.0066%	0.0172%	0.0277%	0.0113%	0.0042%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0043%	-	-	-	-	0.0015%	0.0060%	0.0140%	0.0260%	0.0293%	0.0045%
	Males	0.0066%	-	-	-	-	0.0027%	0.0119%	0.0243%	0.0448%	0.0444%	0.0077%
	Females	0.0021%	-	-	-	-	-	0.0051%	0.0118%	0.0200%	0.0200%	0.0020%
Hispanic <sup>d</sup>	Both Sexes	0.0023%	-	-	-	0.0005%	0.0018%	0.0060%	0.0123%	0.0234%	0.0237%	0.0041%
	Males	0.0036%	-	-	-	0.0008%	0.0028%	0.0102%	0.0240%	0.0398%	0.0515%	0.0075%
	Females	0.0008%	-	-	-	-	-	0.0020%	-	0.0111%	0.0074%	0.0014%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.21  
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	37,947	30,453	27,098	24,139	19,975	13,751	115,151	161,071	13,837	174,908
	Males	20,359	15,887	14,007	12,141	9,981	7,455	60,143	83,458	6,967	90,425
	Females	17,588	14,566	13,091	11,998	9,994	6,296	55,008	77,613	6,870	84,483
White <sup>b</sup>	Both Sexes	31,782	26,338	24,141	21,810	18,539	12,827	99,899	142,784	15,415	158,199
	Males	17,098	13,759	12,552	11,025	9,200	6,802	52,332	73,838	7,993	81,831
	Females	14,684	12,579	11,589	10,785	9,339	6,025	47,567	68,946	7,422	76,368
Black <sup>b</sup>	Both Sexes	4,762	3,055	2,412	1,923	1,149	738	11,932	14,292	1,047	15,339
	Males	2,470	1,514	1,201	928	641	549	6,007	7,461	719	8,180
	Females	2,292	1,541	1,211	995	508	189	5,925	6,831	328	7,159
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	865	647	311	+	+	+	2,023	+	+	+
	Males	465	342	156	+	+	+	1,049	+	+	+
	Females	400	305	155	+	+	+	974	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,640	2,685	1,950	+	+	+	9,597	+	+	+
	Males	1,987	1,607	1,019	+	+	+	5,325	+	+	+
	Females	1,653	1,078	931	+	+	+	4,272	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0287%	0.0007%	0.0100%	0.0380%	0.0520%	0.0436%	0.0305%	0.0278%	0.0247%	0.0194%	0.0286%
	Males	0.0306%	0.0007%	0.0117%	0.0361%	0.0502%	0.0475%	0.0368%	0.0341%	0.0277%	0.0248%	0.0305%
	Females	0.0269%	0.0006%	0.0082%	0.0400%	0.0537%	0.0397%	0.0245%	0.0220%	0.0224%	0.0164%	0.0270%
White <sup>c</sup>	Both Sexes	0.0321%	0.0007%	0.0109%	0.0422%	0.0591%	0.0490%	0.0335%	0.0304%	0.0283%	0.0218%	0.0321%
	Males	0.0340%	0.0007%	0.0127%	0.0390%	0.0571%	0.0523%	0.0402%	0.0370%	0.0312%	0.0278%	0.0337%
	Females	0.0301%	0.0007%	0.0089%	0.0457%	0.0613%	0.0454%	0.0269%	0.0242%	0.0259%	0.0184%	0.0306%
Black <sup>c</sup>	Both Sexes	0.0257%	-	0.0089%	0.0300%	0.0493%	0.0408%	0.0317%	0.0276%	0.0184%	0.0132%	0.0261%
	Males	0.0271%	-	0.0083%	0.0310%	0.0452%	0.0470%	0.0399%	0.0333%	0.0215%	0.0207%	0.0284%
	Females	0.0244%	-	0.0095%	0.0290%	0.0529%	0.0354%	0.0248%	0.0232%	0.0163%	0.0097%	0.0243%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0117%	-	0.0052%	0.0210%	0.0193%	0.0145%	0.0104%	0.0097%	0.0079%	0.0057%	0.0113%
	Males	0.0128%	-	0.0075%	0.0236%	0.0178%	0.0162%	0.0100%	0.0141%	0.0091%	-	0.0124%
	Females	0.0108%	-	0.0027%	0.0184%	0.0207%	0.0130%	0.0108%	0.0058%	0.0070%	0.0053%	0.0101%
Hispanic <sup>d</sup>	Both Sexes	0.0177%	0.0006%	0.0102%	0.0240%	0.0269%	0.0253%	0.0214%	0.0288%	0.0278%	0.0229%	0.0196%
	Males	0.0190%	0.0005%	0.0128%	0.0239%	0.0255%	0.0275%	0.0281%	0.0295%	0.0317%	0.0404%	0.0217%
	Females	0.0163%	0.0008%	0.0074%	0.0242%	0.0285%	0.0228%	0.0148%	0.0282%	0.0249%	0.0126%	0.0178%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.15  
Kaposi SarcomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	7,129	5,158	7,229	2,809	880	262	21,959	23,623	+	+
	Males	6,715	4,810	7,001	2,599	869	216	20,770	22,336	+	+
	Females	414	348	228	210	11	46	1,189	1,287	+	+
White <sup>b</sup>	Both Sexes	4,951	3,763	5,833	2,373	747	190	16,615	17,989	+	+
	Males	4,662	3,462	5,680	2,221	736	155	15,731	17,018	+	+
	Females	289	301	153	152	11	35	884	971	+	+
Black <sup>b</sup>	Both Sexes	1,839	1,172	940	276	56	38	4,193	4,333	+	+
	Males	1,732	1,137	898	241	56	38	3,974	4,114	+	+
	Females	107	35	42	35	0	0	219	219	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	106	88	88	+	+	+	291	+	+	+
	Males	101	88	88	+	+	+	286	+	+	+
	Females	5	0	0	+	+	+	5	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,236	908	1,261	+	+	+	3,785	+	+	+
	Males	1,174	862	1,221	+	+	+	3,612	+	+	+
	Females	62	46	40	+	+	+	173	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0072%	-	-	0.0009%	0.0064%	0.0198%	0.0141%	0.0082%	0.0071%	0.0128%	0.0072%
	Males	0.0140%	-	-	0.0017%	0.0125%	0.0389%	0.0285%	0.0163%	0.0142%	0.0245%	0.0141%
	Females	0.0006%	-	-	-	0.0002%	0.0006%	0.0003%	0.0010%	0.0015%	0.0062%	0.0005%
White <sup>c</sup>	Both Sexes	0.0072%	-	-	0.0007%	0.0056%	0.0194%	0.0144%	0.0084%	0.0080%	0.0131%	0.0070%
	Males	0.0138%	-	-	0.0014%	0.0107%	0.0374%	0.0287%	0.0164%	0.0154%	0.0250%	0.0137%
	Females	0.0006%	-	-	-	-	0.0005%	-	0.0009%	0.0020%	0.0064%	0.0005%
Black <sup>c</sup>	Both Sexes	0.0103%	-	-	0.0025%	0.0154%	0.0313%	0.0182%	0.0110%	0.0030%	0.0107%	0.0109%
	Males	0.0209%	-	-	0.0048%	0.0327%	0.0661%	0.0387%	0.0223%	0.0074%	-	0.0227%
	Females	0.0007%	-	-	-	-	-	-	-	-	0.0109%	0.0008%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0019%	-	-	-	0.0014%	0.0048%	0.0039%	0.0022%	0.0033%	0.0037%	0.0019%
	Males	0.0039%	-	-	-	0.0028%	0.0097%	0.0084%	0.0048%	0.0076%	0.0081%	0.0040%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic <sup>d</sup>	Both Sexes	0.0072%	-	-	0.0014%	0.0077%	0.0237%	0.0174%	0.0102%	0.0138%	0.0305%	0.0096%
	Males	0.0134%	-	-	0.0025%	0.0141%	0.0444%	0.0351%	0.0201%	0.0265%	0.0578%	0.0183%
	Females	0.0007%	-	-	-	-	0.0009%	-	-	0.0044%	0.0145%	0.0011%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.21  
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	137,327	74,063	43,448	27,201	14,677	8,542	277,556	309,168	11,014	320,182
	Males	82,513	43,626	25,550	15,571	8,248	4,782	164,872	182,285	6,074	188,359
	Females	54,814	30,437	17,898	11,630	6,429	3,760	112,684	126,883	4,940	131,823
White <sup>b</sup>	Both Sexes	115,768	63,624	37,373	24,143	13,158	7,432	236,800	265,057	12,524	277,581
	Males	69,632	37,790	22,451	13,998	7,576	4,291	141,644	157,560	5,557	163,117
	Females	46,136	25,834	14,922	10,145	5,582	3,141	95,156	107,497	6,967	114,464
Black <sup>b</sup>	Both Sexes	14,888	7,524	4,574	2,391	1,258	897	29,056	31,804	1,958	33,762
	Males	8,761	4,084	2,254	1,185	530	390	16,155	17,321	673	17,994
	Females	6,127	3,440	2,320	1,206	728	507	12,901	14,483	1,285	15,768
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,256	1,608	809	+	+	+	5,974	+	+	+
	Males	2,039	966	503	+	+	+	3,688	+	+	+
	Females	1,217	642	306	+	+	+	2,286	+	+	+
Hispanic <sup>d</sup>	Both Sexes	11,974	6,084	2,983	+	+	+	22,560	+	+	+
	Males	7,056	3,490	1,581	+	+	+	12,961	+	+	+
	Females	4,918	2,594	1,402	+	+	+	9,599	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0742%	0.0065%	0.0089%	0.0027%	0.0129%	0.0474%	0.1237%	0.2512%	0.3446%	0.2930%	0.0741%
	Males	0.0902%	0.0064%	0.0077%	0.0025%	0.0138%	0.0579%	0.1613%	0.3423%	0.4596%	0.4217%	0.0975%
	Females	0.0584%	0.0066%	0.0103%	0.0030%	0.0120%	0.0369%	0.0876%	0.1686%	0.2531%	0.2212%	0.0547%
White <sup>c</sup>	Both Sexes	0.0806%	0.0067%	0.0096%	0.0026%	0.0131%	0.0501%	0.1287%	0.2626%	0.3625%	0.3051%	0.0774%
	Males	0.0977%	0.0065%	0.0088%	0.0024%	0.0138%	0.0601%	0.1659%	0.3529%	0.4833%	0.4442%	0.1014%
	Females	0.0635%	0.0069%	0.0104%	0.0029%	0.0123%	0.0396%	0.0917%	0.1784%	0.2642%	0.2275%	0.0572%
Black <sup>c</sup>	Both Sexes	0.0631%	0.0084%	0.0098%	0.0035%	0.0157%	0.0478%	0.1406%	0.2687%	0.3401%	0.2784%	0.0776%
	Males	0.0752%	0.0093%	0.0053%	0.0032%	0.0165%	0.0610%	0.1904%	0.3922%	0.4355%	0.3500%	0.1019%
	Females	0.0522%	0.0074%	0.0143%	0.0038%	0.0149%	0.0363%	0.0989%	0.1741%	0.2745%	0.2449%	0.0597%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0419%	0.0030%	0.0031%	0.0019%	0.0069%	0.0255%	0.0677%	0.1431%	0.2122%	0.1909%	0.0430%
	Males	0.0533%	0.0025%	0.0024%	0.0022%	0.0069%	0.0341%	0.0932%	0.2068%	0.2915%	0.2833%	0.0594%
	Females	0.0312%	0.0035%	0.0039%	0.0016%	0.0070%	0.0177%	0.0455%	0.0890%	0.1522%	0.1337%	0.0301%
Hispanic <sup>d</sup>	Both Sexes	0.0444%	0.0055%	0.0075%	0.0018%	0.0128%	0.0431%	0.1186%	0.2477%	0.3628%	0.3233%	0.0746%
	Males	0.0492%	0.0048%	0.0059%	0.0013%	0.0124%	0.0490%	0.1441%	0.3268%	0.4974%	0.4047%	0.0946%
	Females	0.0393%	0.0061%	0.0092%	0.0025%	0.0132%	0.0366%	0.0938%	0.1793%	0.2619%	0.2757%	0.0587%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.21  
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	30,294	20,347	14,369	9,734	6,297	4,132	73,107	86,894	2,248	89,142
	Males	24,153	16,488	11,459	7,754	5,178	3,313	58,562	69,765	1,843	71,608
	Females	6,141	3,859	2,910	1,980	1,119	819	14,545	17,129	405	17,534
White <sup>b</sup>	Both Sexes	25,283	17,333	12,467	8,332	5,459	3,743	62,070	74,141	1,781	75,922
	Males	20,268	14,095	9,929	6,700	4,550	3,044	49,889	59,854	1,457	61,311
	Females	5,015	3,238	2,538	1,632	909	699	12,181	14,287	324	14,611
Black <sup>b</sup>	Both Sexes	4,231	2,428	1,569	1,139	720	301	9,143	10,539	159	10,698
	Males	3,258	1,877	1,257	869	553	201	7,101	8,130	135	8,265
	Females	973	551	312	270	167	100	2,042	2,409	24	2,433
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	447	352	222	+	+	+	1,142	+	+	+
	Males	383	307	196	+	+	+	990	+	+	+
	Females	64	45	26	+	+	+	152	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,660	1,035	665	+	+	+	3,739	+	+	+
	Males	1,403	849	549	+	+	+	3,125	+	+	+
	Females	257	186	116	+	+	+	614	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0181%	-	-	0.0003%	0.0008%	0.0056%	0.0265%	0.0706%	0.1091%	0.0805%	0.0183%
	Males	0.0296%	-	-	0.0003%	0.0009%	0.0076%	0.0427%	0.1246%	0.1996%	0.1877%	0.0333%
	Females	0.0068%	-	-	0.0002%	0.0007%	0.0036%	0.0110%	0.0217%	0.0370%	0.0207%	0.0064%
White <sup>c</sup>	Both Sexes	0.0197%	-	-	0.0003%	0.0009%	0.0061%	0.0280%	0.0719%	0.1136%	0.0827%	0.0190%
	Males	0.0318%	-	-	0.0003%	0.0009%	0.0083%	0.0450%	0.1245%	0.2047%	0.1912%	0.0340%
	Females	0.0075%	-	-	0.0003%	0.0009%	0.0038%	0.0111%	0.0228%	0.0394%	0.0222%	0.0067%
Black <sup>c</sup>	Both Sexes	0.0195%	-	-	-	0.0009%	0.0063%	0.0378%	0.1150%	0.1423%	0.0968%	0.0255%
	Males	0.0322%	-	-	-	0.0017%	0.0071%	0.0602%	0.2203%	0.2775%	0.2555%	0.0496%
	Females	0.0080%	-	-	-	-	0.0056%	0.0191%	0.0344%	0.0494%	0.0226%	0.0093%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0079%	-	-	-	-	0.0022%	0.0081%	0.0271%	0.0544%	0.0545%	0.0083%
	Males	0.0141%	-	-	-	-	0.0041%	0.0135%	0.0536%	0.1074%	0.1290%	0.0168%
	Females	0.0020%	-	-	-	-	-	0.0033%	0.0046%	0.0142%	0.0083%	0.0019%
Hispanic <sup>d</sup>	Both Sexes	0.0070%	-	-	-	0.0006%	0.0033%	0.0144%	0.0483%	0.0852%	0.0757%	0.0134%
	Males	0.0113%	-	-	-	0.0006%	0.0036%	0.0238%	0.0930%	0.1651%	0.1783%	0.0260%
	Females	0.0025%	-	-	-	-	0.0029%	0.0054%	0.0097%	0.0252%	0.0157%	0.0039%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.29  
LeukemiaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	108,472	64,337	39,582	24,594	15,440	8,839	232,921	266,026	5,854	271,880
	Males	62,188	36,459	22,201	13,656	8,601	4,678	132,236	150,062	2,636	152,698
	Females	46,284	27,878	17,381	10,938	6,839	4,161	100,685	115,964	3,218	119,182
White <sup>b</sup>	Both Sexes	94,458	57,374	35,500	22,166	13,964	8,132	205,822	235,994	4,934	240,928
	Males	54,452	32,647	19,822	12,539	7,873	4,321	117,388	133,809	1,893	135,702
	Females	40,006	24,727	15,678	9,627	6,091	3,811	88,434	102,185	3,041	105,226
Black <sup>b</sup>	Both Sexes	8,023	3,883	2,143	1,312	756	399	15,193	16,754	706	17,460
	Males	4,420	1,946	1,115	618	344	207	8,010	8,735	326	9,061
	Females	3,603	1,937	1,028	694	412	192	7,183	8,019	380	8,399
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,692	1,495	1,096	+	+	+	5,876	+	+	+
	Males	1,479	864	604	+	+	+	3,265	+	+	+
	Females	1,213	631	492	+	+	+	2,611	+	+	+
Hispanic <sup>d</sup>	Both Sexes	10,100	5,778	3,983	+	+	+	22,075	+	+	+
	Males	5,812	3,192	2,260	+	+	+	12,537	+	+	+
	Females	4,288	2,586	1,723	+	+	+	9,538	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0634%	0.0291%	0.0545%	0.0229%	0.0222%	0.0350%	0.0677%	0.1442%	0.2317%	0.2592%	0.0644%
	Males	0.0731%	0.0314%	0.0594%	0.0252%	0.0246%	0.0411%	0.0846%	0.1868%	0.3052%	0.3435%	0.0792%
	Females	0.0539%	0.0267%	0.0493%	0.0205%	0.0197%	0.0288%	0.0516%	0.1056%	0.1734%	0.2122%	0.0519%
White <sup>c</sup>	Both Sexes	0.0712%	0.0321%	0.0611%	0.0251%	0.0244%	0.0361%	0.0731%	0.1575%	0.2560%	0.2839%	0.0704%
	Males	0.0817%	0.0348%	0.0660%	0.0274%	0.0267%	0.0427%	0.0904%	0.2022%	0.3373%	0.3763%	0.0863%
	Females	0.0607%	0.0293%	0.0560%	0.0225%	0.0220%	0.0293%	0.0559%	0.1158%	0.1898%	0.2325%	0.0568%
Black <sup>c</sup>	Both Sexes	0.0338%	0.0142%	0.0240%	0.0155%	0.0135%	0.0268%	0.0458%	0.0923%	0.1526%	0.1512%	0.0397%
	Males	0.0381%	0.0162%	0.0260%	0.0166%	0.0153%	0.0320%	0.0595%	0.1224%	0.1793%	0.1918%	0.0483%
	Females	0.0298%	0.0120%	0.0220%	0.0145%	0.0120%	0.0221%	0.0343%	0.0692%	0.1343%	0.1322%	0.0333%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0356%	0.0266%	0.0478%	0.0171%	0.0180%	0.0295%	0.0375%	0.0619%	0.0838%	0.0889%	0.0371%
	Males	0.0410%	0.0277%	0.0561%	0.0190%	0.0208%	0.0333%	0.0453%	0.0731%	0.1076%	0.1246%	0.0441%
	Females	0.0305%	0.0255%	0.0390%	0.0151%	0.0153%	0.0260%	0.0308%	0.0524%	0.0658%	0.0668%	0.0313%
Hispanic <sup>d</sup>	Both Sexes	0.0418%	0.0317%	0.0652%	0.0269%	0.0233%	0.0288%	0.0495%	0.0851%	0.1123%	0.1225%	0.0483%
	Males	0.0458%	0.0345%	0.0707%	0.0304%	0.0254%	0.0333%	0.0574%	0.1000%	0.1271%	0.1762%	0.0556%
	Females	0.0375%	0.0287%	0.0593%	0.0229%	0.0209%	0.0237%	0.0419%	0.0723%	0.1013%	0.0911%	0.0419%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 13.30  
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	15,984	12,034	10,764	8,909	6,862	5,223	45,921	62,732	3,462	66,194
	Males	9,278	6,641	6,045	4,815	3,747	2,713	25,796	34,732	1,847	36,579
	Females	6,706	5,393	4,719	4,094	3,115	2,510	20,125	28,000	1,615	29,615
White <sup>b</sup>	Both Sexes	13,498	10,518	9,397	7,811	6,055	4,804	39,671	54,838	+	+
	Males	7,778	5,861	5,225	4,341	3,345	2,501	22,344	30,480	+	+
	Females	5,720	4,657	4,172	3,470	2,710	2,303	17,327	24,358	+	+
Black <sup>b</sup>	Both Sexes	1,323	684	704	585	371	201	3,177	3,999	+	+
	Males	880	298	404	255	174	103	1,760	2,166	+	+
	Females	443	386	300	330	197	98	1,417	1,833	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	774	521	478	+	+	+	2,069	+	+	+
	Males	419	313	289	+	+	+	1,160	+	+	+
	Females	355	208	189	+	+	+	909	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,649	2,848	2,401	+	+	+	11,256	+	+	+
	Males	2,743	1,519	1,360	+	+	+	6,409	+	+	+
	Females	1,906	1,329	1,041	+	+	+	4,847	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0137%	0.0244%	0.0468%	0.0133%	0.0050%	0.0036%	0.0029%	0.0038%	0.0023%	0.0016%	0.0143%
	Males	0.0157%	0.0261%	0.0512%	0.0158%	0.0062%	0.0041%	0.0034%	0.0045%	0.0029%	0.0022%	0.0159%
	Females	0.0118%	0.0226%	0.0422%	0.0108%	0.0037%	0.0032%	0.0025%	0.0032%	0.0018%	0.0013%	0.0126%
White <sup>c</sup>	Both Sexes	0.0153%	0.0273%	0.0533%	0.0154%	0.0059%	0.0038%	0.0032%	0.0040%	0.0019%	0.0016%	0.0161%
	Males	0.0173%	0.0293%	0.0575%	0.0180%	0.0071%	0.0044%	0.0037%	0.0047%	0.0026%	0.0019%	0.0177%
	Females	0.0133%	0.0252%	0.0490%	0.0125%	0.0046%	0.0033%	0.0027%	0.0033%	0.0014%	0.0015%	0.0143%
Black <sup>c</sup>	Both Sexes	0.0065%	0.0104%	0.0185%	0.0073%	0.0014%	0.0014%	0.0013%	0.0019%	-	-	0.0060%
	Males	0.0079%	0.0128%	0.0207%	0.0084%	0.0015%	-	-	-	-	-	0.0069%
	Females	0.0053%	0.0079%	0.0161%	0.0062%	0.0013%	0.0016%	-	-	-	-	0.0051%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0109%	0.0220%	0.0391%	0.0080%	0.0036%	0.0042%	0.0020%	0.0040%	0.0045%	-	0.0121%
	Males	0.0131%	0.0227%	0.0464%	0.0098%	0.0050%	0.0047%	0.0027%	0.0038%	-	-	0.0139%
	Females	0.0089%	0.0214%	0.0313%	0.0062%	0.0022%	0.0038%	0.0015%	0.0042%	0.0050%	-	0.0104%
Hispanic <sup>d</sup>	Both Sexes	0.0207%	0.0271%	0.0577%	0.0176%	0.0070%	0.0048%	0.0051%	0.0059%	0.0031%	-	0.0178%
	Males	0.0228%	0.0291%	0.0618%	0.0211%	0.0080%	0.0061%	0.0050%	0.0043%	-	-	0.0194%
	Females	0.0185%	0.0250%	0.0533%	0.0134%	0.0059%	0.0033%	0.0053%	0.0073%	-	-	0.0160%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.21  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	26,017	5,399	1,820	823	653	276	33,953	35,119	438	35,557
	Males	18,881	3,689	1,066	475	246	134	24,087	24,549	73	24,622
	Females	7,136	1,710	754	348	407	142	9,866	10,570	365	10,935
White <sup>b</sup>	Both Sexes	18,734	3,962	1,373	620	504	224	24,595	25,507	381	25,888
	Males	13,710	2,733	786	358	155	105	17,564	17,891	195 <sup>i</sup>	18,086
	Females	5,024	1,229	587	262	349	119	7,031	7,616	186 <sup>i</sup>	7,802
Black <sup>b</sup>	Both Sexes	3,793	453	113	106	71	37	4,453	4,606	+	+
	Males	2,708	257	64	48	37	24	3,077	3,149	+	+
	Females	1,085	196	49	58	34	13	1,376	1,457	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,020	868	279	+	+	+	4,253	+	+	+
	Males	2,171	585	193	+	+	+	3,011	+	+	+
	Females	849	283	86	+	+	+	1,242	+	+	+
Hispanic <sup>d</sup>	Both Sexes	3,764	677	268	+	+	+	4,825	+	+	+
	Males	2,713	438	181	+	+	+	3,413	+	+	+
	Females	1,051	239	87	+	+	+	1,412	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted <sup>f</sup>
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0129%	0.0025%	0.0018%	0.0007%	0.0014%	0.0063%	0.0295%	0.0450%	0.0526%	0.0308%	0.0125%
	Males	0.0184%	0.0027%	0.0021%	0.0009%	0.0015%	0.0094%	0.0480%	0.0690%	0.0766%	0.0484%	0.0188%
	Females	0.0074%	0.0024%	0.0015%	0.0006%	0.0013%	0.0032%	0.0118%	0.0232%	0.0334%	0.0210%	0.0070%
White <sup>c</sup>	Both Sexes	0.0102%	0.0028%	0.0017%	0.0008%	0.0012%	0.0047%	0.0246%	0.0334%	0.0343%	0.0223%	0.0095%
	Males	0.0146%	0.0028%	0.0019%	0.0010%	0.0011%	0.0067%	0.0405%	0.0510%	0.0501%	0.0349%	0.0142%
	Females	0.0058%	0.0027%	0.0015%	0.0005%	0.0012%	0.0026%	0.0088%	0.0171%	0.0215%	0.0153%	0.0053%
Black <sup>c</sup>	Both Sexes	0.0104%	-	0.0013%	0.0009%	0.0014%	0.0036%	0.0351%	0.0514%	0.0406%	0.0294%	0.0120%
	Males	0.0150%	-	0.0023%	-	0.0020%	0.0047%	0.0524%	0.0900%	0.0621%	0.0448%	0.0187%
	Females	0.0063%	-	-	0.0013%	-	0.0025%	0.0206%	0.0218%	0.0259%	0.0222%	0.0070%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0305%	0.0029%	0.0028%	-	0.0024%	0.0183%	0.0530%	0.1143%	0.1721%	0.0865%	0.0313%
	Males	0.0442%	0.0034%	0.0022%	-	0.0034%	0.0307%	0.0895%	0.1771%	0.2573%	0.1310%	0.0485%
	Females	0.0176%	0.0024%	0.0033%	-	0.0014%	0.0070%	0.0213%	0.0608%	0.1075%	0.0590%	0.0173%
Hispanic <sup>d</sup>	Both Sexes	0.0099%	0.0028%	0.0017%	0.0006%	0.0007%	0.0067%	0.0354%	0.0630%	0.0661%	0.0460%	0.0159%
	Males	0.0135%	0.0031%	0.0021%	0.0007%	0.0010%	0.0103%	0.0591%	0.0974%	0.0910%	0.0543%	0.0236%
	Females	0.0060%	0.0024%	0.0013%	-	-	0.0028%	0.0124%	0.0332%	0.0474%	0.0412%	0.0091%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.27  
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	224,508	73,096	40,029	21,696	11,093	5,706	355,972	379,223	8,539	387,762
	Males	104,519	32,530	17,506	10,024	4,915	2,722	162,893	173,619	4,871	178,490
	Females	119,989	40,566	22,523	11,672	6,178	2,984	193,079	205,604	3,668	209,272
White <sup>b</sup>	Both Sexes	191,522	64,113	35,002	19,420	9,843	5,042	307,035	327,592	3,735	331,327
	Males	88,141	28,202	14,846	8,931	4,264	2,410	138,623	147,973	1,789	149,762
	Females	103,381	35,911	20,156	10,489	5,579	2,632	168,412	179,619	1,946	181,565
Black <sup>b</sup>	Both Sexes	23,966	6,652	3,723	1,787	967	503	35,864	37,934	351	38,285
	Males	11,723	3,226	1,905	853	496	240	17,555	18,621	186	18,807
	Females	12,243	3,426	1,818	934	471	263	18,309	19,313	165	19,478
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	6,612	1,700	916	+	+	+	9,545	+	+	+
	Males	3,473	809	521	+	+	+	4,980	+	+	+
	Females	3,139	891	395	+	+	+	4,565	+	+	+
Hispanic <sup>d</sup>	Both Sexes	7,956	2,206	1,171	+	+	+	11,845	+	+	+
	Males	3,986	1,069	578	+	+	+	5,883	+	+	+
	Females	3,970	1,137	593	+	+	+	5,962	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0935%	0.0002%	0.0002%	0.0017%	0.0051%	0.0244%	0.1067%	0.3345%	0.6208%	0.5351%	0.0961%
	Males	0.0874%	-	0.0003%	0.0016%	0.0047%	0.0213%	0.1042%	0.3414%	0.6513%	0.6330%	0.1007%
	Females	0.0995%	-	-	0.0017%	0.0056%	0.0276%	0.1090%	0.3283%	0.5966%	0.4805%	0.0933%
White <sup>c</sup>	Both Sexes	0.1001%	0.0002%	0.0003%	0.0018%	0.0051%	0.0247%	0.1039%	0.3399%	0.6450%	0.5473%	0.0982%
	Males	0.0905%	-	0.0003%	0.0017%	0.0048%	0.0211%	0.0973%	0.3340%	0.6572%	0.6278%	0.0996%
	Females	0.1096%	-	-	0.0019%	0.0055%	0.0283%	0.1104%	0.3455%	0.6351%	0.5025%	0.0980%
Black <sup>c</sup>	Both Sexes	0.0803%	-	-	0.0014%	0.0054%	0.0305%	0.1567%	0.4186%	0.6023%	0.4870%	0.1058%
	Males	0.0832%	-	-	0.0014%	0.0056%	0.0274%	0.1818%	0.4882%	0.7220%	0.6712%	0.1262%
	Females	0.0777%	-	-	0.0013%	0.0053%	0.0333%	0.1358%	0.3653%	0.5200%	0.4008%	0.0924%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0711%	-	-	0.0011%	0.0048%	0.0179%	0.0828%	0.2402%	0.4912%	0.4691%	0.0754%
	Males	0.0761%	-	-	-	0.0037%	0.0178%	0.0885%	0.2903%	0.5797%	0.6339%	0.0902%
	Females	0.0663%	-	-	0.0014%	0.0059%	0.0179%	0.0777%	0.1976%	0.4241%	0.3668%	0.0645%
Hispanic <sup>d</sup>	Both Sexes	0.0238%	-	-	0.0011%	0.0031%	0.0122%	0.0435%	0.1517%	0.3030%	0.2768%	0.0460%
	Males	0.0227%	-	-	0.0007%	0.0035%	0.0119%	0.0417%	0.1614%	0.3657%	0.3306%	0.0521%
	Females	0.0250%	-	-	0.0016%	0.0027%	0.0126%	0.0453%	0.1432%	0.2561%	0.2454%	0.0418%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.21  
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	263,883	191,244	140,062	98,190	72,228	48,232	675,956	837,215	39,129	876,344
	Males	140,746	97,383	70,090	46,927	30,976	19,650	347,086	415,003	12,807	427,810
	Females	123,137	93,861	69,972	51,263	41,252	28,582	328,870	422,212	26,322	448,534
White <sup>b</sup>	Both Sexes	254,590	184,997	134,614	94,402	68,857	46,812	651,810	806,429	36,949	843,378
	Males	136,325	94,247	67,392	45,397	29,701	19,065	335,483	400,868	12,222	413,090
	Females	118,265	90,750	67,222	49,005	39,156	27,747	316,327	405,561	24,727	430,288
Black <sup>b</sup>	Both Sexes	1,042	611	564	355	175	109	2,482	2,957	112	3,069
	Males	417	296	287	150	24	35	1,119	1,237	44	1,281
	Females	625	315	277	205	151	74	1,363	1,720	68	1,788
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	682	481	288	+	+	+	1,580	+	+	+
	Males	345	203	127	+	+	+	717	+	+	+
	Females	337	278	161	+	+	+	863	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,723	3,480	2,337	+	+	+	11,964	+	+	+
	Males	1,922	1,159	835	+	+	+	4,429	+	+	+
	Females	2,801	2,321	1,502	+	+	+	7,535	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1615%	0.0004%	0.0035%	0.0289%	0.0903%	0.1826%	0.2901%	0.4395%	0.5233%	0.5543%	0.1596%
	Males	0.1707%	0.0003%	0.0031%	0.0184%	0.0636%	0.1548%	0.3043%	0.5494%	0.7473%	0.8839%	0.1838%
	Females	0.1524%	0.0005%	0.0039%	0.0401%	0.1179%	0.2106%	0.2765%	0.3399%	0.3452%	0.3705%	0.1443%
White <sup>c</sup>	Both Sexes	0.2058%	0.0004%	0.0043%	0.0370%	0.1171%	0.2309%	0.3616%	0.5360%	0.6417%	0.6538%	0.1975%
	Males	0.2160%	0.0003%	0.0039%	0.0233%	0.0805%	0.1917%	0.3732%	0.6616%	0.9072%	1.0450%	0.2234%
	Females	0.1956%	0.0005%	0.0048%	0.0520%	0.1566%	0.2718%	0.3502%	0.4187%	0.4256%	0.4358%	0.1821%
Black <sup>c</sup>	Both Sexes	0.0053%	-	-	0.0013%	0.0034%	0.0060%	0.0065%	0.0200%	0.0299%	0.0308%	0.0067%
	Males	0.0050%	-	-	-	0.0018%	0.0058%	0.0059%	0.0220%	0.0407%	0.0482%	0.0077%
	Females	0.0056%	-	-	0.0019%	0.0048%	0.0062%	0.0069%	0.0186%	0.0224%	0.0227%	0.0063%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0103%	-	0.0011%	0.0020%	0.0068%	0.0123%	0.0159%	0.0287%	0.0258%	0.0434%	0.0103%
	Males	0.0100%	-	0.0016%	0.0018%	0.0054%	0.0090%	0.0163%	0.0309%	0.0329%	0.0617%	0.0108%
	Females	0.0106%	-	-	0.0022%	0.0081%	0.0152%	0.0155%	0.0268%	0.0204%	0.0321%	0.0100%
Hispanic <sup>d</sup>	Both Sexes	0.0225%	-	0.0011%	0.0048%	0.0168%	0.0396%	0.0607%	0.0872%	0.1154%	0.1353%	0.0340%
	Males	0.0160%	-	0.0010%	0.0025%	0.0087%	0.0232%	0.0413%	0.0720%	0.1403%	0.1833%	0.0299%
	Females	0.0294%	-	0.0012%	0.0076%	0.0263%	0.0576%	0.0794%	0.1004%	0.0968%	0.1071%	0.0395%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.13  
MesotheliomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	3,040	486	385	249	189	107	4,122	4,570	+	+
	Males	2,095	256	116	86	60	58	2,526	2,694	+	+
	Females	945	230	269	163	129	49	1,596	1,876	+	+
White <sup>b</sup>	Both Sexes	2,826	397	350	221	177	95	3,758	4,155	+	+
	Males	1,935	217	104	70	48	46	2,303	2,431	+	+
	Females	891	180	246	151	129	49	1,455	1,724	+	+
Black <sup>b</sup>	Both Sexes	144	56	26	26	12	12	251	298	+	+
	Males	109	34	13	13	12	12	169	205	+	+
	Females	35	22	13	13	0	0	82	93	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	30	16	4	+	+	+	50	+	+	+
	Males	9	10	4	+	+	+	23	+	+	+
	Females	21	6	0	+	+	+	27	+	+	+
Hispanic <sup>d</sup>	Both Sexes	266	49	26	+	+	+	358	+	+	+
	Males	142	38	21	+	+	+	201	+	+	+
	Females	124	11	5	+	+	+	157	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted <sup>f</sup> All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0011%	-	-	-	0.0003%	0.0005%	0.0013%	0.0034%	0.0062%	0.0072%	0.0011%
	Males	0.0014%	-	-	-	0.0002%	0.0004%	0.0016%	0.0044%	0.0099%	0.0147%	0.0017%
	Females	0.0008%	-	-	-	0.0004%	0.0007%	0.0010%	0.0025%	0.0033%	0.0030%	0.0008%
White <sup>c</sup>	Both Sexes	0.0013%	-	-	-	0.0003%	0.0006%	0.0015%	0.0039%	0.0070%	0.0078%	0.0013%
	Males	0.0017%	-	-	-	0.0002%	0.0004%	0.0019%	0.0053%	0.0108%	0.0154%	0.0019%
	Females	0.0009%	-	-	-	0.0004%	0.0008%	0.0011%	0.0027%	0.0039%	0.0035%	0.0009%
Black <sup>c</sup>	Both Sexes	0.0005%	-	-	-	-	-	0.0010%	0.0020%	0.0042%	-	0.0007%
	Males	0.0007%	-	-	-	-	-	-	-	0.0089%	-	0.0013%
	Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0003%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0004%	-	-	-	-	-	-	0.0012%	-	0.0043%	0.0004%
	Males	0.0004%	-	-	-	-	-	-	-	-	0.0097%	0.0005%
	Females	0.0004%	-	-	-	-	-	-	-	-	-	0.0003%
Hispanic <sup>d</sup>	Both Sexes	0.0007%	-	-	-	-	0.0006%	0.0022%	0.0049%	0.0059%	0.0061%	0.0013%
	Males	0.0008%	-	-	-	-	-	0.0034%	0.0062%	0.0064%	0.0124%	0.0017%
	Females	0.0006%	-	-	-	-	-	-	0.0037%	0.0055%	-	0.0010%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.21  
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	45,747	15,680	5,928	2,305	905	279	69,411	70,950	263	71,213	
	Males	24,230	8,744	3,306	1,343	513	160	37,464	38,319	95	38,414	
	Females	21,517	6,936	2,622	962	392	119	31,947	32,631	168	32,799	
White <sup>b</sup>	Both Sexes	34,935	12,486	4,640	1,739	649	188	53,654	54,719	101	54,820	
	Males	19,405	7,200	2,680	990	372	117	30,182	30,788	0 <sup>i</sup>	30,788	
	Females	15,530	5,286	1,960	749	277	71	23,472	23,931	101 <sup>i</sup>	24,032	
Black <sup>b</sup>	Both Sexes	9,098	2,589	1,058	465	203	82	13,126	13,521	+	+	
	Males	4,024	1,254	499	301	127	39	6,018	6,245	+	+	
	Females	5,074	1,335	559	164	76	43	7,108	7,276	+	+	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	1,016	354	117	+	+	+	1,537	+	+	+	
	Males	506	173	67	+	+	+	771	+	+	+	
	Females	510	181	50	+	+	+	766	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	3,195	1,233	373	+	+	+	4,921	+	+	+	
	Males	1,761	679	219	+	+	+	2,739	+	+	+	
	Females	1,434	554	154	+	+	+	2,182	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted <sup>f</sup>		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0199%	-	-	0.0002%	0.0021%	0.0099%	0.0304%	0.0745%	0.1031%	0.0936%	0.0199%
	Males	0.0217%	-	-	0.0003%	0.0027%	0.0112%	0.0350%	0.0881%	0.1242%	0.1188%	0.0238%
	Females	0.0181%	-	-	-	0.0014%	0.0086%	0.0259%	0.0621%	0.0863%	0.0795%	0.0167%
White <sup>c</sup>	Both Sexes	0.0194%	-	-	0.0002%	0.0020%	0.0090%	0.0282%	0.0691%	0.0969%	0.0883%	0.0186%
	Males	0.0219%	-	-	0.0003%	0.0025%	0.0109%	0.0332%	0.0840%	0.1218%	0.1182%	0.0230%
	Females	0.0169%	-	-	-	0.0013%	0.0071%	0.0233%	0.0551%	0.0767%	0.0716%	0.0148%
Black <sup>c</sup>	Both Sexes	0.0310%	-	-	-	0.0037%	0.0214%	0.0612%	0.1513%	0.2049%	0.1845%	0.0400%
	Males	0.0302%	-	-	-	0.0045%	0.0200%	0.0690%	0.1766%	0.2127%	0.1929%	0.0437%
	Females	0.0318%	-	-	-	0.0029%	0.0227%	0.0547%	0.1319%	0.1996%	0.1806%	0.0374%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0113%	-	-	-	0.0010%	0.0045%	0.0149%	0.0453%	0.0598%	0.0666%	0.0116%
	Males	0.0115%	-	-	-	0.0017%	0.0053%	0.0157%	0.0508%	0.0693%	0.0697%	0.0130%
	Females	0.0110%	-	-	-	-	0.0038%	0.0142%	0.0406%	0.0527%	0.0647%	0.0105%
Hispanic <sup>d</sup>	Both Sexes	0.0101%	-	-	0.0004%	0.0023%	0.0083%	0.0264%	0.0645%	0.1015%	0.0798%	0.0181%
	Males	0.0108%	-	-	0.0007%	0.0032%	0.0086%	0.0307%	0.0792%	0.1145%	0.1065%	0.0216%
	Females	0.0094%	-	-	-	0.0012%	0.0079%	0.0223%	0.0517%	0.0918%	0.0642%	0.0154%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.25  
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	193,568	124,952	73,118	40,328	23,808	13,206	425,784	474,455	9,881	484,336
	Males	101,809	64,597	37,822	20,781	12,282	6,865	221,874	246,926	5,185	252,111
	Females	91,759	60,355	35,296	19,547	11,526	6,341	203,910	227,529	4,696	232,225
White <sup>b</sup>	Both Sexes	169,344	111,105	64,658	36,119	21,607	12,076	375,558	419,930	10,218	430,148
	Males	89,651	57,853	33,336	18,717	11,169	6,255	196,674	219,545	5,745	225,290
	Females	79,693	53,252	31,322	17,402	10,438	5,821	178,884	200,385	4,473	204,858
Black <sup>b</sup>	Both Sexes	15,176	8,849	5,327	2,662	1,414	748	31,735	34,493	788	35,281
	Males	7,256	4,279	2,863	1,262	730	392	15,520	16,913	326	17,239
	Females	7,920	4,570	2,464	1,400	684	356	16,215	17,580	462	18,042
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,134	3,241	1,818	+	+	+	11,000	+	+	+
	Males	2,740	1,632	930	+	+	+	5,705	+	+	+
	Females	2,394	1,609	888	+	+	+	5,295	+	+	+
Hispanic <sup>d</sup>	Both Sexes	13,379	8,080	4,315	+	+	+	27,683	+	+	+
	Males	7,003	4,207	2,320	+	+	+	14,463	+	+	+
	Females	6,376	3,873	1,995	+	+	+	13,220	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted <sup>f</sup>
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1153%	0.0026%	0.0115%	0.0208%	0.0396%	0.0862%	0.1728%	0.3318%	0.5039%	0.5486%	0.1155%
	Males	0.1221%	0.0033%	0.0150%	0.0252%	0.0458%	0.1019%	0.1966%	0.3775%	0.5621%	0.6296%	0.1323%
	Females	0.1087%	0.0018%	0.0077%	0.0162%	0.0332%	0.0705%	0.1499%	0.2904%	0.4576%	0.5035%	0.1010%
White <sup>c</sup>	Both Sexes	0.1277%	0.0027%	0.0117%	0.0212%	0.0409%	0.0901%	0.1830%	0.3570%	0.5473%	0.5864%	0.1232%
	Males	0.1347%	0.0036%	0.0154%	0.0261%	0.0476%	0.1061%	0.2070%	0.4044%	0.6074%	0.6742%	0.1407%
	Females	0.1207%	0.0018%	0.0077%	0.0159%	0.0337%	0.0735%	0.1592%	0.3127%	0.4983%	0.5374%	0.1077%
Black <sup>c</sup>	Both Sexes	0.0713%	0.0024%	0.0110%	0.0205%	0.0422%	0.0831%	0.1485%	0.2386%	0.2937%	0.2691%	0.0840%
	Males	0.0753%	0.0029%	0.0142%	0.0240%	0.0511%	0.0989%	0.1722%	0.2720%	0.2816%	0.3000%	0.0941%
	Females	0.0677%	0.0019%	0.0077%	0.0172%	0.0344%	0.0693%	0.1286%	0.2129%	0.3021%	0.2547%	0.0757%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0746%	0.0017%	0.0097%	0.0181%	0.0262%	0.0574%	0.1129%	0.1978%	0.3356%	0.3810%	0.0763%
	Males	0.0793%	0.0022%	0.0119%	0.0208%	0.0274%	0.0669%	0.1273%	0.2261%	0.3958%	0.4352%	0.0879%
	Females	0.0702%	-	0.0073%	0.0154%	0.0250%	0.0486%	0.1004%	0.1737%	0.2900%	0.3474%	0.0670%
Hispanic <sup>d</sup>	Both Sexes	0.0547%	0.0023%	0.0088%	0.0160%	0.0299%	0.0652%	0.1331%	0.2453%	0.3831%	0.3971%	0.0874%
	Males	0.0552%	0.0030%	0.0115%	0.0196%	0.0350%	0.0733%	0.1456%	0.2524%	0.3854%	0.4209%	0.0934%
	Females	0.0541%	0.0015%	0.0060%	0.0118%	0.0239%	0.0562%	0.1210%	0.2392%	0.3813%	0.3832%	0.0816%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.27  
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	97,542	58,825	39,428	24,908	16,305	10,015	216,832	252,887	11,555	264,442
	Males	67,675	39,111	25,048	15,519	9,587	5,769	145,098	166,069	6,639	172,708
	Females	29,867	19,714	14,380	9,389	6,718	4,246	71,734	86,818	4,916	91,734
White <sup>b</sup>	Both Sexes	84,225	51,096	34,480	21,818	14,412	8,856	188,211	220,015	9,834	229,849
	Males	59,083	34,536	22,154	13,870	8,646	5,274	127,653	146,590	5,682	152,272
	Females	25,142	16,560	12,326	7,948	5,766	3,582	60,558	73,425	4,152	77,577
Black <sup>b</sup>	Both Sexes	7,297	4,522	2,580	1,678	1,168	597	15,869	18,257	589	18,846
	Males	4,760	2,613	1,450	819	568	167	9,506	10,536	223	10,759
	Females	2,537	1,909	1,130	859	600	430	6,363	7,721	366	8,087
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,677	2,170	1,584	+	+	+	8,161	+	+	+
	Males	2,309	1,311	955	+	+	+	5,024	+	+	+
	Females	1,368	859	629	+	+	+	3,137	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,957	2,644	1,786	+	+	+	10,280	+	+	+
	Males	3,112	1,669	1,044	+	+	+	6,262	+	+	+
	Females	1,845	975	742	+	+	+	4,018	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0584%	0.0005%	0.0019%	0.0056%	0.0145%	0.0430%	0.1195%	0.1973%	0.2200%	0.2022%	0.0571%
	Males	0.0788%	0.0004%	0.0018%	0.0054%	0.0140%	0.0557%	0.1767%	0.3019%	0.3242%	0.3028%	0.0824%
	Females	0.0383%	0.0005%	0.0020%	0.0058%	0.0150%	0.0301%	0.0645%	0.1026%	0.1372%	0.1461%	0.0356%
White <sup>c</sup>	Both Sexes	0.0623%	0.0005%	0.0016%	0.0054%	0.0130%	0.0404%	0.1225%	0.2074%	0.2339%	0.2130%	0.0587%
	Males	0.0848%	0.0004%	0.0012%	0.0048%	0.0130%	0.0536%	0.1837%	0.3153%	0.3458%	0.3201%	0.0853%
	Females	0.0396%	0.0006%	0.0021%	0.0061%	0.0129%	0.0267%	0.0617%	0.1066%	0.1428%	0.1533%	0.0354%
Black <sup>c</sup>	Both Sexes	0.0364%	0.0007%	0.0028%	0.0055%	0.0113%	0.0354%	0.0975%	0.1504%	0.1466%	0.1163%	0.0427%
	Males	0.0477%	-	0.0040%	0.0078%	0.0069%	0.0419%	0.1428%	0.2443%	0.2199%	0.1710%	0.0612%
	Females	0.0261%	-	0.0016%	0.0032%	0.0152%	0.0298%	0.0597%	0.0784%	0.0962%	0.0907%	0.0287%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0529%	-	0.0024%	0.0058%	0.0233%	0.0602%	0.1124%	0.1508%	0.1690%	0.1541%	0.0524%
	Males	0.0671%	-	0.0031%	0.0057%	0.0245%	0.0761%	0.1515%	0.2288%	0.2259%	0.2240%	0.0709%
	Females	0.0395%	-	0.0017%	0.0060%	0.0221%	0.0455%	0.0785%	0.0845%	0.1258%	0.1108%	0.0374%
Hispanic <sup>d</sup>	Both Sexes	0.0198%	0.0004%	0.0012%	0.0036%	0.0085%	0.0216%	0.0582%	0.1100%	0.1379%	0.1238%	0.0322%
	Males	0.0237%	-	0.0008%	0.0037%	0.0076%	0.0257%	0.0774%	0.1556%	0.2027%	0.1730%	0.0438%
	Females	0.0157%	-	0.0016%	0.0035%	0.0096%	0.0171%	0.0395%	0.0705%	0.0892%	0.0951%	0.0228%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 21.15  
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	55,497	31,173	23,579	19,390	15,986	12,050	126,262	164,877	17,881	182,758
White <sup>b</sup>	48,202	27,307	20,665	17,431	14,416	10,736	110,554	145,192	15,290	160,482
Black <sup>b</sup>	4,120	1,938	1,676	1,179	1,042	808	8,717	11,336	1,719	13,055
Asian/Pacific Islander <sup>c</sup>	2,365	1,425	870	+	+	+	5,183	+	+	+
Hispanic <sup>d</sup>	4,541	2,518	1,662	+	+	+	9,748	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races <sup>c</sup>	0.0682%	0.0003%	0.0056%	0.0168%	0.0269%	0.0628%	0.1320%	0.2019%	0.2076%	0.1606%	0.0637%	
White <sup>c</sup>	0.0741%	0.0003%	0.0060%	0.0175%	0.0264%	0.0641%	0.1379%	0.2174%	0.2266%	0.1690%	0.0671%	
Black <sup>c</sup>	0.0374%	-	0.0041%	0.0113%	0.0225%	0.0421%	0.0761%	0.1112%	0.1372%	0.1305%	0.0411%	
Asian/Pacific Islander <sup>c</sup>	0.0633%	-	0.0046%	0.0177%	0.0318%	0.0714%	0.1418%	0.1745%	0.1366%	0.1135%	0.0591%	
Hispanic <sup>d</sup>	0.0386%	-	0.0080%	0.0186%	0.0245%	0.0507%	0.1005%	0.1595%	0.1734%	0.1341%	0.0527%	

- <sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.  
<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census.  
<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.  
<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
 + Not available.

Table 22.21  
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
Race	Sex										
All Races <sup>b</sup>	Both Sexes	28,875	4,434	2,188	1,249	635	454	36,583	37,971	337	38,308
	Males	14,376	2,055	1,015	535	233	141	17,907	18,446	143	18,589
	Females	14,499	2,379	1,173	714	402	313	18,676	19,525	194	19,719
White <sup>b</sup>	Both Sexes	24,202	3,764	1,720	976	464	383	30,578	31,609	278	31,887
	Males	12,300	1,842	860	430	193	130	15,395	15,812	128	15,940
	Females	11,902	1,922	860	546	271	253	15,183	15,797	150	15,947
Black <sup>b</sup>	Both Sexes	3,325	437	266	177	133	44	4,133	4,414	135	4,549
	Males	1,442	138	69	61	40	0	1,674	1,780	56	1,836
	Females	1,883	299	197	116	93	44	2,459	2,634	79	2,713
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	958	181	93	+	+	+	1,258	+	+	+
	Males	473	62	37	+	+	+	580	+	+	+
	Females	485	119	56	+	+	+	678	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,807	257	190	+	+	+	2,348	+	+	+
	Males	811	123	49	+	+	+	1,021	+	+	+
	Females	996	134	141	+	+	+	1,327	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted <sup>f</sup>		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0105%	-	0.0001%	0.0006%	0.0015%	0.0056%	0.0170%	0.0362%	0.0534%	0.0494%	0.0105%
	Males	0.0104%	-	-	0.0003%	0.0013%	0.0057%	0.0187%	0.0404%	0.0554%	0.0559%	0.0113%
	Females	0.0107%	-	0.0002%	0.0008%	0.0018%	0.0056%	0.0154%	0.0323%	0.0518%	0.0458%	0.0099%
White <sup>c</sup>	Both Sexes	0.0109%	-	0.0001%	0.0006%	0.0015%	0.0053%	0.0169%	0.0364%	0.0530%	0.0482%	0.0104%
	Males	0.0110%	-	-	0.0004%	0.0012%	0.0055%	0.0189%	0.0410%	0.0568%	0.0564%	0.0114%
	Females	0.0108%	-	-	0.0008%	0.0018%	0.0050%	0.0149%	0.0322%	0.0499%	0.0437%	0.0095%
Black <sup>c</sup>	Both Sexes	0.0097%	-	-	-	0.0021%	0.0076%	0.0211%	0.0440%	0.0558%	0.0517%	0.0122%
	Males	0.0082%	-	-	-	0.0021%	0.0068%	0.0237%	0.0404%	0.0434%	0.0482%	0.0113%
	Females	0.0111%	-	-	-	0.0021%	0.0083%	0.0190%	0.0467%	0.0643%	0.0533%	0.0128%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0093%	-	-	-	0.0012%	0.0055%	0.0138%	0.0279%	0.0522%	0.0564%	0.0097%
	Males	0.0088%	-	-	-	-	0.0051%	0.0125%	0.0370%	0.0528%	0.0571%	0.0101%
	Females	0.0097%	-	-	-	0.0015%	0.0059%	0.0149%	0.0202%	0.0518%	0.0560%	0.0094%
Hispanic <sup>d</sup>	Both Sexes	0.0048%	-	-	0.0007%	0.0010%	0.0045%	0.0106%	0.0316%	0.0395%	0.0473%	0.0084%
	Males	0.0040%	-	-	-	-	0.0047%	0.0105%	0.0268%	0.0425%	0.0439%	0.0080%
	Females	0.0056%	-	-	0.0010%	0.0015%	0.0043%	0.0107%	0.0357%	0.0372%	0.0493%	0.0087%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.  
<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.  
<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).  
<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.  
<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.  
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.  
+ Not available.

Table 23.15  
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	991,610	772,659	454,494	225,090	39,253	9,569	2,426,184	2,494,988	1,796	2,496,784
White <sup>b</sup>	805,947	649,042	387,249	200,794	35,855	8,795	2,026,925	2,089,734	2,032	2,091,766
Black <sup>b</sup>	135,003	97,839	54,635	19,776	2,784	661	306,001	310,925	168	311,093
Asian/Pacific Islander <sup>c</sup>	19,903	13,544	6,303	+	+	+	41,987	+	+	+
Hispanic <sup>d</sup>	55,469	37,852	20,199	+	+	+	120,434	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup> All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races <sup>c</sup>	1.2781%	-	0.0002%	0.0002%	0.0010%	0.0801%	1.0020%	4.9316%	10.8539%	11.8049%	1.5268%
White <sup>c</sup>	1.3283%	-	0.0002%	-	0.0008%	0.0700%	0.9509%	4.8456%	10.8782%	11.5288%	1.5048%
Black <sup>c</sup>	1.4257%	-	-	-	-	0.1951%	1.9940%	8.6540%	15.7798%	16.4060%	2.3775%
Asian/Pacific Islander <sup>c</sup>	0.6582%	-	-	-	0.0014%	0.0239%	0.3441%	2.0771%	5.8657%	8.3851%	0.8269%
Hispanic <sup>d</sup>	0.4559%	-	-	-	0.0007%	0.0362%	0.5915%	3.4192%	8.6859%	9.4350%	1.1623%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.21  
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	34,419	14,791	8,365	4,910	3,629	1,649	61,601	68,386	1,600	69,986
	Males	20,553	8,158	4,718	2,255	1,834	935	35,303	38,804	903	39,707
	Females	13,866	6,633	3,647	2,655	1,795	714	26,298	29,582	697	30,279
White <sup>b</sup>	Both Sexes	25,785	10,840	6,017	3,678	2,726	1,321	45,621	50,847	1,280	52,127
	Males	16,065	6,143	3,402	1,620	1,418	809	26,912	29,720	698	30,418
	Females	9,720	4,697	2,615	2,058	1,308	512	18,709	21,127	582	21,709
Black <sup>b</sup>	Both Sexes	5,393	2,325	1,354	541	567	164	9,536	10,423	225	10,648
	Males	2,629	1,142	683	286	252	89	4,726	5,134	125	5,259
	Females	2,764	1,183	671	255	315	75	4,810	5,289	100	5,389
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,053	1,484	930	+	+	+	5,974	+	+	+
	Males	1,736	781	544	+	+	+	3,345	+	+	+
	Females	1,317	703	386	+	+	+	2,629	+	+	+
Hispanic <sup>d</sup>	Both Sexes	4,878	1,795	1,071	+	+	+	8,258	+	+	+
	Males	2,709	978	514	+	+	+	4,412	+	+	+
	Females	2,169	817	557	+	+	+	3,846	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0212%	-	-	0.0007%	0.0032%	0.0116%	0.0272%	0.0589%	0.1139%	0.1502%	0.0214%
	Males	0.0243%	-	-	0.0005%	0.0032%	0.0126%	0.0336%	0.0779%	0.1542%	0.2016%	0.0278%
	Females	0.0180%	-	-	0.0011%	0.0032%	0.0106%	0.0209%	0.0417%	0.0818%	0.1215%	0.0165%
White <sup>c</sup>	Both Sexes	0.0176%	-	-	0.0007%	0.0029%	0.0095%	0.0227%	0.0482%	0.0896%	0.1103%	0.0170%
	Males	0.0208%	-	-	0.0005%	0.0031%	0.0102%	0.0284%	0.0665%	0.1244%	0.1482%	0.0224%
	Females	0.0145%	-	-	0.0010%	0.0028%	0.0088%	0.0169%	0.0311%	0.0613%	0.0891%	0.0126%
Black <sup>c</sup>	Both Sexes	0.0210%	-	-	-	0.0043%	0.0164%	0.0384%	0.0784%	0.1336%	0.2031%	0.0279%
	Males	0.0216%	-	-	-	0.0042%	0.0205%	0.0413%	0.0933%	0.1690%	0.2421%	0.0336%
	Females	0.0204%	-	-	-	0.0043%	0.0127%	0.0359%	0.0669%	0.1091%	0.1848%	0.0241%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0427%	-	-	0.0012%	0.0034%	0.0193%	0.0448%	0.1121%	0.2509%	0.4031%	0.0452%
	Males	0.0491%	-	-	-	0.0029%	0.0198%	0.0606%	0.1424%	0.3353%	0.5365%	0.0587%
	Females	0.0368%	-	-	0.0019%	0.0039%	0.0189%	0.0311%	0.0864%	0.1870%	0.3204%	0.0352%
Hispanic <sup>d</sup>	Both Sexes	0.0164%	-	-	0.0012%	0.0045%	0.0161%	0.0375%	0.0817%	0.1593%	0.1975%	0.0297%
	Males	0.0170%	-	-	0.0007%	0.0048%	0.0157%	0.0424%	0.1070%	0.1993%	0.2396%	0.0360%
	Females	0.0157%	-	-	0.0017%	0.0042%	0.0165%	0.0329%	0.0598%	0.1292%	0.1728%	0.0249%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.15  
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	41,292	37,674	33,588	30,617	24,508	18,217	136,985	194,935	16,274	211,209
White <sup>b</sup>	38,482	35,360	31,772	29,384	23,734	17,605	129,064	184,996	15,600	200,596
Black <sup>b</sup>	1,032	1,002	760	416	309	299	3,110	4,012	157	4,169
Asian/Pacific Islander <sup>c</sup>	755	600	491	+	+	+	2,195	+	+	+
Hispanic <sup>d</sup>	5,435	3,985	2,910	+	+	+	14,281	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.0709%	0.0021%	0.0063%	0.0656%	0.1448%	0.1560%	0.0929%	0.0382%	0.0149%	0.0076%	0.0702%
White <sup>c</sup>	0.0858%	0.0020%	0.0074%	0.0801%	0.1794%	0.1891%	0.1089%	0.0444%	0.0172%	0.0088%	0.0850%
Black <sup>c</sup>	0.0152%	0.0016%	0.0032%	0.0100%	0.0281%	0.0345%	0.0265%	0.0128%	-	-	0.0157%
Asian/Pacific Islander <sup>c</sup>	0.0245%	0.0034%	0.0037%	0.0248%	0.0426%	0.0465%	0.0386%	0.0119%	0.0057%	-	0.0235%
Hispanic <sup>d</sup>	0.0511%	0.0023%	0.0082%	0.0665%	0.1172%	0.0921%	0.0494%	0.0216%	0.0063%	-	0.0496%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 26.21  
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	150,696	99,270	68,603	49,444	37,734	28,415	359,113	456,180	40,721	496,901
	Males	33,895	21,428	14,497	11,046	7,911	5,907	78,991	99,597	9,323	108,920
	Females	116,801	77,842	54,106	38,398	29,823	22,508	280,122	356,583	31,398	387,981
White <sup>b</sup>	Both Sexes	131,211	87,013	60,604	44,265	33,836	25,840	315,040	402,611	36,877	439,488
	Males	30,240	19,306	13,177	10,107	7,225	5,348	71,088	89,957	7,849	97,806
	Females	100,971	67,707	47,427	34,158	26,611	20,492	243,952	312,654	29,028	341,682
Black <sup>b</sup>	Both Sexes	9,269	5,798	3,617	2,529	1,657	1,294	20,839	25,217	1,711	26,928
	Males	1,672	1,019	565	531	302	243	3,710	4,428	192	4,620
	Females	7,597	4,779	3,052	1,998	1,355	1,051	17,129	20,789	1,519	22,308
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	6,845	4,582	3,026	+	+	+	16,148	+	+	+
	Males	1,255	839	495	+	+	+	2,881	+	+	+
	Females	5,590	3,743	2,531	+	+	+	13,267	+	+	+
Hispanic <sup>d</sup>	Both Sexes	14,617	9,084	5,853	+	+	+	32,831	+	+	+
	Males	2,507	1,597	953	+	+	+	5,599	+	+	+
	Females	12,110	7,487	4,900	+	+	+	27,232	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.0991%	0.0001%	0.0047%	0.0347%	0.1085%	0.1689%	0.1949%	0.1989%	0.1695%	0.1118%	0.0976%
	Males	0.0430%	-	0.0020%	0.0096%	0.0347%	0.0645%	0.0882%	0.1124%	0.1109%	0.0777%	0.0438%
	Females	0.1544%	-	0.0076%	0.0615%	0.1847%	0.2739%	0.2973%	0.2774%	0.2161%	0.1308%	0.1497%
White <sup>c</sup>	Both Sexes	0.1069%	0.0001%	0.0052%	0.0382%	0.1205%	0.1840%	0.2054%	0.2043%	0.1719%	0.1099%	0.1039%
	Males	0.0477%	-	0.0022%	0.0106%	0.0382%	0.0713%	0.0964%	0.1191%	0.1177%	0.0766%	0.0473%
	Females	0.1663%	-	0.0084%	0.0683%	0.2091%	0.3018%	0.3136%	0.2838%	0.2161%	0.1285%	0.1610%
Black <sup>c</sup>	Both Sexes	0.0462%	-	0.0014%	0.0116%	0.0436%	0.0805%	0.1039%	0.1294%	0.1209%	0.0837%	0.0516%
	Males	0.0170%	-	0.0013%	0.0027%	0.0103%	0.0304%	0.0367%	0.0605%	0.0686%	0.0685%	0.0214%
	Females	0.0729%	-	0.0015%	0.0204%	0.0729%	0.1244%	0.1601%	0.1821%	0.1569%	0.0909%	0.0764%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0987%	-	0.0045%	0.0327%	0.0973%	0.1532%	0.1990%	0.2086%	0.1819%	0.1351%	0.0958%
	Males	0.0368%	-	-	0.0079%	0.0343%	0.0505%	0.0737%	0.0994%	0.0915%	0.0834%	0.0379%
	Females	0.1567%	-	0.0083%	0.0577%	0.1570%	0.2473%	0.3081%	0.3013%	0.2505%	0.1671%	0.1470%
Hispanic <sup>d</sup>	Both Sexes	0.0635%	-	0.0043%	0.0280%	0.0789%	0.1291%	0.1769%	0.1823%	0.1633%	0.1110%	0.0825%
	Males	0.0207%	-	0.0018%	0.0054%	0.0209%	0.0359%	0.0663%	0.0847%	0.0967%	0.0582%	0.0308%
	Females	0.1089%	-	0.0069%	0.0543%	0.1473%	0.2314%	0.2843%	0.2667%	0.2133%	0.1419%	0.1345%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.21  
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	196,403	132,873	88,851	57,162	35,819	20,518	465,947	540,518	13,829	554,347
	Males	148,936	98,526	65,546	42,220	26,648	14,501	348,486	402,281	8,953	411,234
	Females	47,467	34,347	23,305	14,942	9,171	6,017	117,461	138,237	4,876	143,113
White <sup>b</sup>	Both Sexes	180,436	124,001	83,738	54,223	34,150	19,647	433,385	504,580	10,793	515,373
	Males	137,650	92,174	61,856	39,956	25,367	13,884	325,159	376,457	6,861	383,318
	Females	42,786	31,827	21,882	14,267	8,783	5,763	108,226	128,123	3,932	132,055
Black <sup>b</sup>	Both Sexes	8,238	5,262	2,879	1,780	1,144	489	18,001	20,027	224	20,251
	Males	5,449	3,513	2,084	1,339	923	307	12,250	13,774	182	13,956
	Females	2,789	1,749	795	441	221	182	5,751	6,253	42	6,295
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	3,499	1,906	1,135	+	+	+	7,122	+	+	+
	Males	2,693	1,457	802	+	+	+	5,404	+	+	+
	Females	806	449	333	+	+	+	1,718	+	+	+
Hispanic <sup>d</sup>	Both Sexes	7,102	4,189	2,431	+	+	+	15,175	+	+	+
	Males	5,266	3,148	1,855	+	+	+	11,304	+	+	+
	Females	1,836	1,041	576	+	+	+	3,871	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races <sup>c</sup>	Both Sexes	0.1160%	0.0002%	0.0004%	0.0013%	0.0068%	0.0278%	0.1066%	0.3424%	0.7309%	0.9975%	0.1191%
	Males	0.1763%	0.0002%	0.0005%	0.0019%	0.0094%	0.0416%	0.1673%	0.5565%	1.2688%	2.0059%	0.2103%
	Females	0.0566%	-	0.0003%	0.0006%	0.0041%	0.0140%	0.0483%	0.1483%	0.3033%	0.4349%	0.0515%
White <sup>c</sup>	Both Sexes	0.1377%	0.0002%	0.0003%	0.0015%	0.0080%	0.0316%	0.1198%	0.3909%	0.8344%	1.0870%	0.1340%
	Males	0.2079%	-	0.0006%	0.0022%	0.0108%	0.0459%	0.1842%	0.6269%	1.4359%	2.2072%	0.2348%
	Females	0.0671%	-	-	0.0007%	0.0051%	0.0166%	0.0558%	0.1707%	0.3449%	0.4632%	0.0579%
Black <sup>c</sup>	Both Sexes	0.0413%	-	0.0006%	-	0.0038%	0.0148%	0.0610%	0.1707%	0.3220%	0.4910%	0.0578%
	Males	0.0598%	-	-	-	0.0061%	0.0239%	0.1037%	0.2896%	0.5381%	0.8828%	0.0991%
	Females	0.0245%	-	0.0013%	-	0.0019%	0.0070%	0.0254%	0.0795%	0.1734%	0.3077%	0.0308%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0522%	-	-	0.0007%	0.0023%	0.0135%	0.0468%	0.1322%	0.3198%	0.5769%	0.0556%
	Males	0.0811%	-	-	-	0.0029%	0.0231%	0.0779%	0.2249%	0.5727%	1.0636%	0.0992%
	Females	0.0251%	-	-	-	0.0018%	0.0048%	0.0197%	0.0534%	0.1281%	0.2750%	0.0237%
Hispanic <sup>d</sup>	Both Sexes	0.0288%	-	-	0.0009%	0.0034%	0.0131%	0.0485%	0.1619%	0.3513%	0.4761%	0.0571%
	Males	0.0415%	-	-	0.0012%	0.0045%	0.0171%	0.0740%	0.2667%	0.6208%	0.9220%	0.0999%
	Females	0.0153%	-	-	-	0.0021%	0.0087%	0.0238%	0.0712%	0.1493%	0.2150%	0.0258%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 28.12  
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>f</sup>	Complete <sup>g</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	61,498	53,571	46,779	41,740	35,040	29,378	195,592	288,648	+	+
	Males	32,966	27,910	24,725	21,880	17,388	14,625	103,297	148,843	+	+
	Females	28,532	25,661	22,054	19,860	17,652	14,753	92,295	139,805	+	+
White <sup>b</sup>	Both Sexes	49,984	45,279	39,437	35,403	30,616	25,968	163,294	244,939	+	+
	Males	26,941	23,719	20,732	18,754	15,222	13,059	86,564	126,824	+	+
	Females	23,043	21,560	18,705	16,649	15,394	12,909	76,730	118,115	+	+
Black <sup>b</sup>	Both Sexes	7,323	5,314	4,626	4,044	2,867	2,398	20,502	28,256	+	+
	Males	3,865	2,628	2,435	2,032	1,342	1,092	10,540	14,036	+	+
	Females	3,458	2,686	2,191	2,012	1,525	1,306	9,962	14,220	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,435	1,862	1,643	+	+	+	7,069	+	+	+
	Males	1,249	1,023	897	+	+	+	3,709	+	+	+
	Females	1,186	839	746	+	+	+	3,360	+	+	+
Hispanic <sup>d</sup>	Both Sexes	11,982	8,894	6,969	+	+	+	32,515	+	+	+
	Males	6,784	4,826	3,616	+	+	+	17,865	+	+	+
	Females	5,198	4,068	3,353	+	+	+	14,650	+	+	+

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census.

<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <34 and >=34.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.



Table 29.7  
U.S. Childhood Cancer Survivors at January 1, 2009  
Number of People Previously Diagnosed with Cancer as Children (Ages 0-19 Years) in the United States and Alive January 1, 2009  
By International Classification of Childhood Cancer (ICCC)<sup>a</sup> Selected Group and Subgroup, Age Group, and Sex

Site	Sex	Complete Prevalence Counts by Age at Prevalence							All Ages		
		0-19	20-29	30-39	40-49	50-59	60+	All Ages	Diagnosed between 1975-2008	Diagnosed before 1975	% <sup>b</sup>
									Count	Count	
All Sites	Both	111,023	89,411	59,878	53,611	33,505	15,703	363,131	288,469	74,662	21
	Male	59,903	46,813	29,211	23,582	14,395	8,427	182,331	148,974	33,357	18
	Female <sup>c</sup>	51,120	42,598	30,667	30,029	19,110	7,276	180,800	139,495	41,305	23
I Leukemia	Both	34,866	18,665	10,312	4,111	1,220	122	69,296	65,041	4,255	6
	Male	19,076	10,046	5,153	2,139	595	53	37,062	35,117	1,945	5
	Female	15,790	8,619	5,159	1,972	625	69	32,234	29,924	2,310	7
Acute lymphocytic leukemia <sup>d</sup>	Both	29,489	15,460	8,888	3,368	284	1	57,490	54,252	3,238	6
	Male	16,246	8,401	4,572	1,961	179	1	31,360	29,636	1,724	5
	Female	13,243	7,059	4,316	1,407	105	0	26,130	24,616	1,514	6
I(b) Acute myeloid leukemia	Both	3,883	2,094	1,063	539	182	1	7,762	7,272	490	6
	Male	1,985	1,046	427	262	99	1	3,820	3,562	258	7
	Female	1,898	1,048	636	277	83	0	3,942	3,710	232	6
II(a) Hodgkin lymphoma	Both	4,286	9,000	7,873	7,531	4,296	936	33,922	28,323	5,599	17
	Male	2,492	4,438	3,934	3,715	2,105	496	17,180	14,176	3,004	17
	Female	1,794	4,562	3,939	3,816	2,191	440	16,742	14,147	2,595	15
II(b,c,e) Non-Hodgkin lymphoma	Both	6,537	6,581	4,250	2,878	1,614	1,007	22,867	19,231	3,636	16
	Male	4,328	4,344	2,957	1,681	818	595	14,723	12,759	1,964	13
	Female	2,209	2,237	1,293	1,197	796	412	8,144	6,472	1,672	21
III CNS and misc intracranial & intraspinal neoplasms	Both	19,994	14,276	8,534	8,257	4,381	1,222	56,664	45,355	11,309	20
	Male	10,961	7,834	4,644	3,927	1,971	525	29,862	24,704	5,158	17
	Female	9,033	6,442	3,890	4,330	2,410	697	26,802	20,651	6,151	23
IV Neuroblastoma & other peripheral nervous cell tumor	Both	9,536	3,359	2,067	1,491	1,359	683	18,495	14,559	3,936	21
	Male	4,872	1,724	869	526	406	168	8,565	7,354	1,211	14
	Female	4,664	1,635	1,198	965	953	515	9,930	7,205	2,725	27
VI Renal tumors	Both	7,751	4,538	3,662	3,999	2,077	615	22,642	14,989	7,653	34
	Male	3,823	2,145	1,698	2,173	1,142	363	11,344	7,089	4,255	38
	Female	3,928	2,393	1,964	1,826	935	252	11,298	7,900	3,398	30
VIII Malignant bone tumors	Both	3,552	4,167	2,903	2,462	2,175	683	15,942	12,296	3,646	23
	Male	2,029	2,360	1,646	1,143	966	366	8,510	6,842	1,668	20
	Female	1,523	1,807	1,257	1,319	1,209	317	7,432	5,454	1,978	27
VIII(a) Osteosarcoma	Both	1,733	2,249	1,655	1,247	1,123	680	8,687	6,526	2,161	25
	Male	961	1,243	946	573	425	368	4,516	3,559	957	21
	Female	772	1,006	709	674	698	312	4,171	2,967	1,204	29
VIII(c) Ewing tumor & related sarcomas of bone	Both	1,356	1,358	828	640	348	88	4,618	3,987	631	14
	Male	795	777	412	291	167	26	2,468	2,210	258	10
	Female	561	581	416	349	181	62	2,150	1,777	373	17
IX Soft tissue & other extraosseous sarcomas	Both	7,088	6,589	4,004	4,808	3,910	4,122	30,521	20,064	10,457	34
	Male	3,926	3,589	2,069	2,252	1,802	2,285	15,923	10,710	5,213	33
	Female	3,162	3,000	1,935	2,556	2,108	1,837	14,598	9,354	5,244	36
X Germ cell & trophoblastic tumors & neoplasms of gonads	Both	4,868	7,533	5,355	6,390	6,298	6,083	36,527	22,357	14,170	39
	Male	2,584	4,881	3,211	3,314	3,183	2,716	19,889	13,223	6,666	34
	Female	2,284	2,652	2,144	3,076	3,115	3,367	16,638	9,134	7,504	45

<sup>a</sup> International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.

<sup>b</sup> Percent diagnosed before 1975 of all ages complete prevalence counts.

<sup>c</sup> All sites females excludes breast cancer.

<sup>d</sup> SEER site recode Acute Lymphocytic Leukemia was used instead of the ICCC classification lymphocytic leukemia (Ia) because of statistical modeling requirements.

For more information regarding the modeling of the prevalence estimates, see Mariotto et al. Long-Term Survivors of Childhood Cancers in the United States. Cancer Epidemiol Biomarkers Prev 2009, page 1034.