

Table 34.1
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2002-2008
Both Sexes by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	1,746,347	66.3	0.0	220,866	60.1	0.1	124,744	65.2	0.2	8,014	58.4	0.6	195,999	66.2	0.1
Oral Cavity & Pharynx	39,649	67.4	0.3	4,794	47.3	0.8	3,260	68.5	1.0	193	63.5	4.2	3,439	63.9	1.0
Esophagus	16,427	19.8	0.4	2,413	14.1	0.9	858	17.6	1.7	89	14.3	4.3	1,343	17.2	1.3
Stomach	24,160	27.2	0.3	4,760	29.6	0.8	4,786	37.1	0.8	251	24.1	3.0	5,700	27.9	0.7
Colon & Rectum	170,881	64.9	0.1	24,384	58.1	0.4	15,190	69.0	0.5	989	62.7	1.8	18,887	63.6	0.4
Liver & Intrahepatic Bile Duct	22,367	17.9	0.3	3,917	14.5	0.8	5,775	23.9	0.7	299	16.8	2.7	5,674	18.8	0.7
Pancreas	41,763	6.0	0.1	6,258	5.8	0.4	3,312	7.8	0.6	242	5.9	1.8	4,831	6.7	0.5
Larynx	13,051	67.1	0.5	2,474	59.4	1.2	479	68.5	2.5	45	51.4	9.2	1,210	64.6	1.7
Lung & Bronchus	221,654	18.2	0.1	29,178	14.9	0.3	13,789	19.2	0.4	941	12.2	1.3	13,897	17.1	0.4
Melanoma of the Skin	83,903	89.1	0.1	425	71.8	2.5	551	77.4	2.0	99	78.6	4.9	2,807	83.8	0.8
Breast	258,126	88.3	0.1	32,251	78.4	0.3	21,677	90.9	0.2	1,141	84.9	1.3	29,540	86.5	0.3
Cervix uteri	16,904	71.8	0.4	3,169	62.5	1.0	1,864	73.0	1.2	133	62.4	5.1	5,058	74.6	0.7
Corpus & Uterus, NOS	50,745	83.1	0.2	5,508	62.4	0.8	4,190	84.2	0.7	261	84.3	2.6	6,023	81.8	0.6
Ovary ^d	28,032	43.6	0.4	2,678	37.5	1.1	2,303	56.3	1.3	144	46.3	4.8	3,697	48.4	1.0
Prostate	279,977	93.8	0.1	49,114	91.8	0.2	15,892	94.2	0.2	837	89.4	1.3	29,704	92.9	0.2
Testis	13,627	95.8	0.2	420	89.5	1.6	457	91.4	1.6	99	96.5	2.0	2,845	93.0	0.5
Urinary Bladder	76,071	79.0	0.2	4,408	66.5	0.8	3,020	79.9	0.9	161	71.3	4.1	4,646	77.0	0.7
Kidney & Renal Pelvis	51,953	72.1	0.2	6,593	72.9	0.6	2,846	72.0	1.0	443	69.4	2.5	7,535	72.3	0.6
Brain & Nervous System	27,826	34.0	0.3	2,170	42.5	1.2	1,529	42.5	1.5	111	48.2	5.7	4,156	45.7	0.9
Thyroid	42,463	96.6	0.1	3,297	96.1	0.4	4,868	96.3	0.3	216	95.3	1.5	7,261	96.9	0.2
Hodgkin Lymphoma	12,139	87.0	0.3	1,686	83.8	1.0	641	86.6	1.5	40	88.5	5.5	2,169	85.1	0.9
Non-Hodgkin Lymphoma	75,199	69.6	0.2	7,037	64.6	0.6	5,224	67.7	0.7	271	68.1	3.2	9,634	66.9	0.6
Myeloma	19,486	45.9	0.5	4,995	48.0	0.9	1,294	46.7	1.8	116	45.7	5.7	2,754	46.5	1.3
Leukemia	48,487	58.2	0.3	4,568	52.7	0.9	3,031	54.5	1.0	225	60.4	3.6	7,440	60.2	0.7
Mesothelioma	3,954	7.6	0.6	202	15.0	3.4	120	11.5	3.3	-	-	-	433	11.1	2.0

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.2
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2002-2008
Males by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	905,720	66.3	0.1	118,336	62.3	0.2	59,424	61.0	0.2	3,829	56.0	0.9	96,697	65.1	0.2
Oral Cavity & Pharynx	27,928	67.3	0.3	3,326	42.3	1.0	2,091	66.5	1.2	135	61.5	5.2	2,310	60.6	1.2
Esophagus	12,916	20.0	0.4	1,688	12.8	1.1	645	16.9	1.9	60	-	-	1,070	16.7	1.4
Stomach	14,865	25.3	0.4	2,664	25.9	1.0	2,700	37.3	1.1	155	20.4	3.7	3,181	27.5	1.0
Colon & Rectum	87,823	65.4	0.2	11,417	57.0	0.6	7,896	68.7	0.6	487	61.9	2.6	10,137	63.7	0.6
Liver & Intrahepatic Bile Duct	16,084	18.4	0.4	2,866	12.3	0.9	4,025	24.4	0.9	212	20.6	3.4	4,044	18.8	0.8
Pancreas	20,709	5.7	0.2	2,817	5.1	0.6	1,633	7.0	0.8	122	5.2	2.5	2,277	5.4	0.6
Larynx	10,493	67.6	0.5	1,978	59.5	1.3	416	69.1	2.6	33	51.7	11.3	1,050	63.0	1.9
Lung & Bronchus	117,452	15.9	0.1	16,736	13.4	0.3	8,143	16.6	0.5	487	12.0	1.9	7,687	14.4	0.5
Melanoma of the Skin	46,863	86.7	0.2	178	62.0	4.2	274	73.4	3.1	51	68.3	7.6	1,125	76.6	1.6
Prostate	279,977	93.8	0.1	49,114	91.8	0.2	15,892	94.2	0.2	837	89.4	1.3	29,704	92.9	0.2
Testis	13,627	95.8	0.2	420	89.5	1.6	457	91.4	1.6	99	96.5	2.0	2,845	93.0	0.5
Urinary Bladder	57,474	80.6	0.2	2,856	70.6	1.0	2,303	81.8	0.9	127	74.1	4.6	3,400	79.6	0.8
Kidney & Renal Pelvis	31,969	71.6	0.3	3,922	72.2	0.8	1,787	71.7	1.3	256	68.7	3.4	4,508	70.5	0.8
Brain & Nervous System	15,546	32.6	0.4	1,107	36.5	1.7	813	39.3	2.0	59	51.9	7.0	2,246	44.3	1.2
Thyroid	10,052	93.6	0.3	601	93.2	1.3	947	93.9	0.9	40	84.9	5.7	1,300	93.4	0.8
Hodgkin Lymphoma	6,581	86.4	0.5	883	81.1	1.6	361	84.4	2.2	-	-	-	1,221	84.4	1.2
Non-Hodgkin Lymphoma	40,330	68.6	0.3	3,794	61.2	0.9	2,777	65.9	1.0	145	70.3	4.2	5,184	64.3	0.8
Myeloma	10,823	47.7	0.6	2,438	46.5	1.4	664	48.2	2.6	51	48.1	8.2	1,457	49.7	1.8
Leukemia	27,832	58.5	0.3	2,487	53.8	1.2	1,682	55.2	1.4	121	62.2	4.8	4,155	60.9	0.9
Mesothelioma	3,043	5.8	0.6	142	14.4	4.0	86	7.2	3.0	-	-	-	309	8.6	2.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.3
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2002-2008
Females by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	840,627	66.2	0.1	102,530	57.6	0.2	65,320	69.1	0.2	4,185	60.6	0.9	99,302	67.2	0.2
Oral Cavity & Pharynx	11,721	67.8	0.5	1,468	58.4	1.5	1,169	72.2	1.5	58	67.8	6.8	1,129	70.7	1.6
Esophagus	3,511	19.2	0.8	725	16.9	1.7	213	19.9	3.3	29	21.7	8.0	273	19.3	2.8
Stomach	9,295	30.2	0.6	2,096	34.1	1.2	2,086	37.0	1.2	96	30.1	5.2	2,519	28.4	1.1
Colon & Rectum	83,058	64.3	0.2	12,967	59.0	0.5	7,294	69.4	0.6	502	63.4	2.5	8,750	63.5	0.6
Liver & Intrahepatic Bile Duct	6,283	16.7	0.6	1,051	19.6	1.5	1,750	22.8	1.3	87	8.9	3.7	1,630	18.6	1.2
Pancreas	21,054	6.3	0.2	3,441	6.3	0.5	1,679	8.7	0.8	120	6.7	2.6	2,554	7.9	0.7
Larynx	2,558	64.6	1.1	496	59.0	2.8	63	63.8	8.4	-	-	-	160	74.4	4.1
Lung & Bronchus	104,202	20.7	0.2	12,442	16.8	0.4	5,646	22.9	0.7	454	12.5	1.9	6,210	20.4	0.6
Melanoma of the Skin	37,040	92.1	0.2	247	78.7	3.0	277	81.2	2.7	48	90.3	4.7	1,682	88.3	0.9
Breast	256,313	88.4	0.1	31,964	78.4	0.3	21,570	90.8	0.2	1,135	84.9	1.3	29,395	86.4	0.3
Cervix uteri	16,904	71.8	0.4	3,169	62.5	1.0	1,864	73.0	1.2	133	62.4	5.1	5,058	74.6	0.7
Corpus & Uterus, NOS	50,745	83.1	0.2	5,508	62.4	0.8	4,190	84.2	0.7	261	84.3	2.6	6,023	81.8	0.6
Ovary ^d	28,032	43.6	0.4	2,678	37.5	1.1	2,303	56.3	1.3	144	46.3	4.8	3,697	48.4	1.0
Urinary Bladder	18,597	74.3	0.4	1,552	58.7	1.4	717	73.4	1.9	34	60.8	8.6	1,246	69.8	1.5
Kidney & Renal Pelvis	19,984	72.8	0.4	2,671	74.1	1.0	1,059	72.4	1.6	187	70.3	3.6	3,027	74.9	0.9
Brain & Nervous System	12,280	35.6	0.5	1,063	48.8	1.7	716	46.0	2.1	52	45.1	8.9	1,910	47.2	1.3
Thyroid	32,411	97.6	0.1	2,696	96.7	0.4	3,921	96.8	0.3	176	97.7	1.1	5,961	97.6	0.2
Hodgkin Lymphoma	5,558	87.6	0.5	803	86.7	1.4	280	89.3	2.0	-	-	-	948	86.1	1.3
Non-Hodgkin Lymphoma	34,869	70.7	0.3	3,243	68.7	0.9	2,447	69.7	1.0	126	66.1	4.7	4,450	69.9	0.8
Myeloma	8,663	43.8	0.7	2,557	49.4	1.3	630	45.3	2.7	65	43.6	7.9	1,297	43.0	1.9
Leukemia	20,655	57.7	0.4	2,081	51.4	1.3	1,349	53.7	1.5	104	58.3	5.2	3,285	59.3	1.0
Mesothelioma	911	13.9	1.5	60	17.1	5.9	34	25.0	8.8	-	-	-	124	16.6	4.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 34.4

Cancer of the Female Breast

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2002-2008

Race/Ethnicity	All Ages				Ages <50				Ages 50+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	313,482	100.0	87.5	0.1	78,978	100.0	89.0	0.1	234,504	100.0	87.1	0.1
White	256,313	81.8	88.4	0.1	60,268	76.3	90.3	0.1	196,045	83.6	87.8	0.1
White Non-Hispanic	228,006	72.7	88.6	0.1	49,692	62.9	91.0	0.2	178,314	76.0	87.9	0.1
White Hispanic	28,307	9.0	86.5	0.3	10,576	13.4	86.6	0.4	17,731	7.6	86.4	0.3
Black	31,964	10.2	78.4	0.3	10,198	12.9	79.1	0.5	21,766	9.3	78.2	0.3
American Indian/Alaska Native ^c	1,135	0.4	84.9	1.3	384	0.5	85.7	2.2	751	0.3	84.5	1.7
Asian	19,989	6.4	91.2	0.3	6,891	8.7	92.3	0.4	13,098	5.6	90.5	0.3
Asian Indian, Pakistani	1,573	0.5	89.0	1.0	614	0.8	91.8	1.4	959	0.4	87.2	1.4
Chinese	4,017	1.3	91.2	0.6	1,474	1.9	92.8	0.9	2,543	1.1	90.3	0.7
Filipino	5,519	1.8	89.5	0.5	1,693	2.1	90.5	0.9	3,826	1.6	89.0	0.6
Japanese	3,180	1.0	93.2	0.5	705	0.9	94.6	1.0	2,475	1.1	92.8	0.6
Korean	1,438	0.5	91.1	0.9	632	0.8	91.1	1.4	806	0.3	91.2	1.2
Vietnamese	1,249	0.4	91.3	1.0	540	0.7	92.4	1.5	709	0.3	90.5	1.4
Other Asian ^d	3,013	1.0	93.0	0.6	1,233	1.6	94.1	0.8	1,780	0.8	92.2	0.8
Pacific Islander	1,581	0.5	86.7	1.1	481	0.6	84.3	2.1	1,100	0.5	87.8	1.2
Hawaiian	1,041	0.3	88.5	1.2	287	0.4	86.6	2.6	754	0.3	89.3	1.3
Other Pacific Islander ^e	540	0.2	82.9	2.1	194	0.2	81.1	3.5	346	0.1	83.9	2.6
Hispanic ^f	29,395	9.4	86.4	0.3	10,959	13.9	86.6	0.4	18,436	7.9	86.3	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.5

Cancer of the Colon and Rectum

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2002-2008

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	213,274	100.0	64.6	0.1	108,621	100.0	65.0	0.2	104,653	100.0	64.1	0.2
White	170,881	80.1	64.9	0.1	87,823	80.9	65.4	0.2	83,058	79.4	64.3	0.2
White Non-Hispanic	152,689	71.6	65.0	0.1	78,025	71.8	65.7	0.2	74,664	71.3	64.3	0.2
White Hispanic	18,192	8.5	63.5	0.4	9,798	9.0	63.5	0.6	8,394	8.0	63.6	0.6
Black	24,384	11.4	58.1	0.4	11,417	10.5	57.0	0.6	12,967	12.4	59.0	0.5
American Indian/Alaska Native ^c	989	0.5	62.7	1.8	487	0.4	61.9	2.6	502	0.5	63.4	2.5
Asian	14,323	6.7	69.5	0.5	7,397	6.8	69.3	0.6	6,926	6.6	69.7	0.7
Asian Indian, Pakistani	653	0.3	70.6	2.2	377	0.3	70.0	2.8	276	0.3	71.4	3.4
Chinese	3,375	1.6	71.2	0.9	1,713	1.6	71.6	1.3	1,662	1.6	70.9	1.3
Filipino	3,063	1.4	66.4	1.1	1,623	1.5	67.0	1.4	1,440	1.4	65.8	1.5
Japanese	3,014	1.4	69.4	1.0	1,519	1.4	67.6	1.5	1,495	1.4	71.1	1.4
Korean	1,368	0.6	67.6	1.5	690	0.6	66.4	2.2	678	0.6	68.8	2.1
Vietnamese	1,135	0.5	68.3	1.7	612	0.6	67.7	2.4	523	0.5	68.9	2.4
Other Asian ^d	1,715	0.8	73.5	1.3	863	0.8	75.3	1.7	852	0.8	71.6	1.8
Pacific Islander	867	0.4	61.3	2.0	499	0.5	59.7	2.7	368	0.4	63.5	3.1
Hawaiian	589	0.3	61.6	2.4	348	0.3	58.6	3.2	241	0.2	65.8	3.8
Other Pacific Islander ^e	278	0.1	60.7	3.6	151	0.1	62.4	4.9	127	0.1	59.2	5.2
Hispanic ^f	18,887	8.9	63.6	0.4	10,137	9.3	63.7	0.6	8,750	8.4	63.5	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.6

Cancer of the Liver and Intrahepatic Bile Duct

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2002-2008

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	32,571	100.0	18.6	0.3	23,341	100.0	18.8	0.3	9,230	100.0	18.2	0.5
White	22,367	68.7	17.9	0.3	16,084	68.9	18.4	0.4	6,283	68.1	16.7	0.6
White Non-Hispanic	16,866	51.8	17.7	0.4	12,153	52.1	18.2	0.5	4,713	51.1	16.4	0.7
White Hispanic	5,501	16.9	18.5	0.7	3,931	16.8	18.8	0.8	1,570	17.0	17.7	1.3
Black	3,917	12.0	14.5	0.8	2,866	12.3	12.3	0.9	1,051	11.4	19.6	1.5
American Indian/Alaska Native ^c	299	0.9	16.8	2.7	212	0.9	20.6	3.4	87	0.9	8.9	3.7
Asian	5,504	16.9	24.4	0.7	3,815	16.3	25.0	0.9	1,689	18.3	23.0	1.3
Asian Indian, Pakistani	155	0.5	21.7	4.4	110	0.5	22.0	5.5	45	0.5	20.7	7.0
Chinese	1,414	4.3	27.1	1.5	1,005	4.3	27.9	1.8	409	4.4	25.0	2.7
Filipino	807	2.5	21.9	1.9	581	2.5	22.6	2.2	226	2.4	20.2	3.3
Japanese	550	1.7	21.8	2.3	257	1.1	29.0	3.4	293	3.2	17.0	2.9
Korean	698	2.1	25.5	2.1	467	2.0	22.6	2.5	231	2.5	30.9	3.5
Vietnamese	1,160	3.6	25.9	1.7	885	3.8	26.4	1.9	275	3.0	24.3	3.5
Other Asian ^d	720	2.2	21.2	1.9	510	2.2	21.2	2.2	210	2.3	21.3	3.8
Pacific Islander	271	0.8	14.8	2.8	210	0.9	13.5	3.1	61	0.7	20.0	5.8
Hawaiian	153	0.5	12.4	3.6	114	0.5	12.5	4.2	39	0.4	12.2	6.6
Other Pacific Islander ^e	118	0.4	18.4	4.1	96	0.4	14.9	4.2	-	-	-	-
Hispanic ^f	5,674	17.4	18.8	0.7	4,044	17.3	18.8	0.8	1,630	17.7	18.6	1.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.7

Cancer of the Lung and Bronchus

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2002-2008

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	266,439	100.0	17.9	0.1	143,273	100.0	15.6	0.1	123,166	100.0	20.4	0.1
White	221,654	83.2	18.2	0.1	117,452	82.0	15.9	0.1	104,202	84.6	20.7	0.2
White Non-Hispanic	208,247	78.2	18.2	0.1	110,032	76.8	16.0	0.1	98,215	79.7	20.7	0.2
White Hispanic	13,407	5.0	17.1	0.4	7,420	5.2	14.3	0.5	5,987	4.9	20.4	0.6
Black	29,178	11.0	14.9	0.3	16,736	11.7	13.4	0.3	12,442	10.1	16.8	0.4
American Indian/Alaska Native ^c	941	0.4	12.2	1.3	487	0.3	12.0	1.9	454	0.4	12.5	1.9
Asian	12,613	4.7	19.4	0.5	7,458	5.2	16.7	0.6	5,155	4.2	23.2	0.8
Asian Indian, Pakistani	481	0.2	25.4	2.5	331	0.2	21.5	2.8	150	0.1	35.1	4.9
Chinese	3,228	1.2	19.5	0.9	1,877	1.3	16.2	1.1	1,351	1.1	24.0	1.5
Filipino	3,282	1.2	19.5	0.9	2,032	1.4	16.3	1.1	1,250	1.0	24.6	1.6
Japanese	2,068	0.8	17.1	1.0	1,107	0.8	14.9	1.3	961	0.8	19.4	1.6
Korean	996	0.4	18.0	1.5	560	0.4	16.5	1.9	436	0.4	20.1	2.4
Vietnamese	1,369	0.5	19.2	1.4	891	0.6	17.5	1.6	478	0.4	22.1	2.5
Other Asian ^d	1,189	0.4	21.9	1.6	660	0.5	19.2	2.1	529	0.4	25.2	2.4
Pacific Islander	1,176	0.4	17.0	1.3	685	0.5	15.0	1.7	491	0.4	19.6	2.1
Hawaiian	819	0.3	17.0	1.6	457	0.3	14.5	2.1	362	0.3	19.7	2.3
Other Pacific Islander ^e	357	0.1	16.7	2.6	228	0.2	15.6	3.2	129	0.1	18.6	4.3
Hispanic ^f	13,897	5.2	17.1	0.4	7,687	5.4	14.4	0.5	6,210	5.0	20.4	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 34.8

Cancer of the Prostate

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2002-2008

Race/Ethnicity	All Ages				Ages <65				Ages 65+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	357,078	100.0	93.6	0.0	148,007	100.0	96.7	0.1	209,071	100.0	91.4	0.1
White	279,977	78.4	93.8	0.1	113,513	76.7	97.0	0.1	166,464	79.6	91.6	0.1
White Non-Hispanic	251,872	70.5	93.9	0.1	102,220	69.1	97.2	0.1	149,652	71.6	91.7	0.1
White Hispanic	28,105	7.9	92.9	0.2	11,293	7.6	95.1	0.3	16,812	8.0	91.3	0.3
Black	49,114	13.8	91.8	0.2	25,338	17.1	95.1	0.2	23,776	11.4	88.2	0.3
American Indian/Alaska Native ^c	837	0.2	89.4	1.3	352	0.2	92.5	1.7	485	0.2	87.1	1.9
Asian	14,836	4.2	94.5	0.2	4,571	3.1	97.3	0.3	10,265	4.9	93.2	0.3
Asian Indian, Pakistani	1,311	0.4	94.9	0.8	534	0.4	97.8	0.8	777	0.4	92.9	1.1
Chinese	3,254	0.9	94.8	0.5	815	0.6	98.4	0.6	2,439	1.2	93.6	0.6
Filipino	4,173	1.2	93.5	0.5	1,426	1.0	95.7	0.7	2,747	1.3	92.2	0.6
Japanese	2,737	0.8	94.6	0.5	630	0.4	98.0	0.7	2,107	1.0	93.6	0.6
Korean	708	0.2	95.2	1.0	183	0.1	97.0	1.5	525	0.3	94.4	1.3
Vietnamese	797	0.2	93.8	1.0	278	0.2	96.6	1.3	519	0.2	92.2	1.4
Other Asian ^d	1,856	0.5	96.0	0.6	705	0.5	98.8	0.5	1,151	0.6	94.3	0.8
Pacific Islander	1,056	0.3	89.6	1.2	358	0.2	92.1	1.7	698	0.3	88.2	1.5
Hawaiian	645	0.2	88.6	1.5	215	0.1	91.0	2.3	430	0.2	87.5	2.0
Other Pacific Islander ^e	411	0.1	91.3	1.7	143	0.1	93.7	2.4	268	0.1	89.9	2.3
Hispanic ^f	29,704	8.3	92.9	0.2	11,857	8.0	95.1	0.2	17,847	8.5	91.3	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2002-2008 with follow-up through 2009.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.