

PENNSYLVANIA

Table 87. Selected Drug Use, Perceptions of Great Risk, Average Annual Marijuana Initiates, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in Pennsylvania, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2009-2010 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ¹	846	91	295	459	755
Past Year Marijuana Use	1,162	131	428	603	1,031
Past Month Marijuana Use	653	71	261	321	582
Past Month Use of Illicit Drugs Other Than Marijuana ¹	340	39	109	192	301
Past Year Cocaine Use	207	8	70	128	199
Past Year Nonmedical Pain Reliever Use	466	55	157	254	411
Perception of Great Risk of Smoking Marijuana Once a Month	3,490	283	226	2,980	3,207
Average Annual Number of Marijuana Initiates ²	114	50	58	6	64
ALCOHOL					
Past Month Alcohol Use	5,852	152	902	4,798	5,700
Past Month Binge Alcohol Use ³	2,648	87	628	1,933	2,561
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	4,160	367	407	3,386	3,793
Past Month Alcohol Use (Persons Aged 12 to 20)	455 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	307 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	3,251	125	613	2,514	3,127
Past Month Cigarette Use	2,654	99	520	2,035	2,555
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	7,288	611	842	5,836	6,677
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	220	25	84	111	195
Illicit Drug Dependence or Abuse ¹	300	43	112	145	257
Alcohol Dependence	329	15	88	226	314
Alcohol Dependence or Abuse	725	40	226	459	685
Alcohol or Illicit Drug Dependence or Abuse ¹	926	68	280	578	858
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	255	41	97	117	215
Needing But Not Receiving Treatment for Alcohol Use ⁷	689	39	215	435	650
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	76	118	519	637
Serious Mental Illness ^{9,10}	--	--	100	338	438
Any Mental Illness ^{9,11}	--	--	419	1,463	1,882
Had Serious Thoughts of Suicide	--	--	90	265	355

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to get an overall estimate (12 or older).

⁹ For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

¹⁰ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment in carrying out major life activities.

¹¹ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, regardless of the level of impairment in carrying out major life activities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).

PENNSYLVANIA

Table 88. Selected Drug Use, Perceptions of Great Risk, Average Annual Rates of First Use of Marijuana, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in Pennsylvania, by Age Group: Percentages, Annual Averages Based on 2009-2010 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLCIT DRUGS					
Past Month Illicit Drug Use ¹	7.99	9.51	21.70	5.55	7.84
Past Year Marijuana Use	10.97	13.62	31.43	7.29	10.70
Past Month Marijuana Use	6.17	7.40	19.18	3.88	6.04
Past Month Use of Illicit Drugs Other Than Marijuana ¹	3.21	4.07	7.99	2.33	3.13
Past Year Cocaine Use	1.96	0.88	5.17	1.55	2.06
Past Year Nonmedical Pain Reliever Use	4.40	5.75	11.55	3.07	4.27
Perception of Great Risk of Smoking Marijuana Once a Month	32.93	29.41	16.62	36.03	33.28
Average Annual Rate of First Use of Marijuana ²	1.79	5.81	8.27	0.13	1.16
ALCOHOL					
Past Month Alcohol Use	55.23	15.78	66.28	58.01	59.18
Past Month Binge Alcohol Use ³	24.99	9.04	46.13	23.37	26.59
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	39.26	38.13	29.89	40.94	39.38
Past Month Alcohol Use (Persons Aged 12 to 20)	29.47 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	19.88 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	30.69	12.94	45.03	30.39	32.46
Past Month Cigarette Use	25.05	10.28	38.18	24.60	26.53
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	68.78	63.44	61.85	70.55	69.32
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	2.08	2.59	6.19	1.34	2.02
Illicit Drug Dependence or Abuse ¹	2.83	4.50	8.24	1.75	2.67
Alcohol Dependence	3.10	1.56	6.46	2.73	3.26
Alcohol Dependence or Abuse	6.84	4.16	16.58	5.55	7.11
Alcohol or Illicit Drug Dependence or Abuse ¹	8.74	7.02	20.58	6.99	8.91
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	2.41	4.21	7.16	1.42	2.23
Needing But Not Receiving Treatment for Alcohol Use ⁷	6.51	4.07	15.80	5.26	6.75
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	7.91	8.67	6.28	6.62
Serious Mental Illness ^{9,10}	--	--	7.33	4.09	4.55
Any Mental Illness ^{9,11}	--	--	30.82	17.68	19.54
Had Serious Thoughts of Suicide	--	--	6.60	3.20	3.68

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual marijuana initiation rate = $100 * \{ [X_1 \div (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of persons who never used marijuana. Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. Note that the age group is based on a respondent's age at the time of the interview, not his or her age at first use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to get an overall estimate (12 or older).

⁹ For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

¹⁰ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment in carrying out major life activities.

¹¹ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, regardless of the level of impairment in carrying out major life activities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2009 and 2010 (Revised March 2012).