

REFERENCE LIST

1. Kuczmarski RJ, Carrol MD, Flegal KM, Troiano RP. Varying body mass index cut-off points to describe overweight prevalence among U.S. adults: NHANES III (1988 to 1994). *Obes Res.* 1997;5:542-548.
2. Brown CD, Donato KA, Obarzanek, E, et al. Body mass index and prevalence of risk factors for cardiovascular disease. Forthcoming.
3. Association of Life Insurance Medical Directors of America. Society of Actuaries. Blood Pressure Study, 1979. Chicago: The Society; 1980.
4. Stamler R, Stamler J, Riedlinger WF, Algera G, Roberts RH. Weight and blood pressure. Findings in hypertension screening of 1 million Americans. *JAMA.* 1978;240:1607-1610.
5. Criqui MH, Mebane I, Wallace RB, Heiss G, Holdbrook MJ. Multivariate correlates of adult blood pressures in nine North American populations: The Lipid Research Clinics Prevalence Study. *Prev Med.* 1982;11:391-402.
6. Dyer AR, Elliott P. The INTERSALT study: relations of body mass index to blood pressure. INTERSALT Co-operative Research Group. *J Hum Hypertens.* 1989;3:299-308.
7. Westlund K, Nicolaysen R. Ten-year mortality and morbidity related to serum cholesterol. A follow-up of 3,751 men aged 40-49. *Scand J Clin Lab Invest Suppl.* 1972;127:1-24.
8. Lew EA, Garfinkel L. Variations in mortality by weight among 750,000 men and women. *J Chronic Dis.* 1979;32:563-576.
9. Larsson B, Bjorntorp P, Tibblin G. The health consequences of moderate obesity. *Int J Obes.* 1981;5:97-116.
10. Medalie JH, Papier C, Herman JB, et al. Diabetes mellitus among 10,000 adult men. I. 5-year incidence and associated variables. *Isr J Med Sci.* 1974;10:681-697.
11. Hubert HB, Feinleib M, McNamara PM, Castelli WP. Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study. *Circulation.* 1983;67:968-977.
12. Rexrode KM, Hennekens CH, Willett WC, et al. A prospective study of body mass index, weight change, and risk of stroke in women. *JAMA.* 1997;277:1539-1545.
13. Walker SP, Rimm EB, Ascherio A, Kawachi I, Stampfer MJ, Willett WC. Body size and fat distribution as predictors of stroke among US men. *Am J Epidemiol.* 1996;144:1143-1150.
14. Stampfer MJ, Maclure KM, Colditz GA, Manson JE, Willett WC. Risk of sympto-

- matic gallstones in women with severe obesity. *Am J Clin Nutr.* 1992;55:652-658.
15. Khare M, Everhart JE, Maurer KR, Hill MC. Association of ethnicity and body mass index (BMI) with gallstone disease in the United States. *Am J Epidemiol.* 1995;141:S69.
 16. Cicuttini FM, Baker JR, Spector TD. The association of obesity with osteoarthritis of the hand and knee in women: a twin study. *J Rheumatol.* 1996;23:1221-1226.
 17. Hart DJ, Spector TD. The relationship of obesity, fat distribution, and osteoarthritis in women in the general population: the Chingford Study. *J Rheumatol.* 1993;20:331-335.
 18. Hochberg MC, Lethbridge-Cejku M, Scott WW Jr, Reichle R, Plato CC, Tobin JD. The association of body weight, body fatness and body fat distribution with osteoarthritis of the knee: data from the Baltimore Longitudinal Study of Aging. *J Rheumatol.* 1995;22:488- 493.
 19. Millman RP, Carlisle CC, McGarvey ST, Eveloff SE, Levinson PD. Body fat distribution and sleep apnea severity in women. *Chest.* 1995;107:362-366.
 20. Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med.* 1993;328:1230-1235.
 21. Shepard JW Jr. Hypertension, cardiac arrhythmias, myocardial infarction, and stroke in relation to obstructive sleep apnea. *Clin Chest Med.* 1992;13:437-458.
 22. Bostick RM, Potter JD, Kushi LH, et al. Sugar, meat, and fat intake, and nondietary risk factors for colon cancer incidence in Iowa women (United States). *Cancer Causes Control.* 1994;5:38-52.
 23. Chute CG, Willett WC, Colditz GA, et al. A prospective study of body mass, height, and smoking on the risk of colorectal cancer in women. *Cancer Causes Control.* 1991;2:117- 124.
 24. Garland C, Shekelle RB, Barrett-Connor E, Criqui MH, Ross AH, Paul O. Dietary vitamin D and calcium and risk of colorectal cancer: a 19-year prospective study in men. *Lancet.* 1985;1:307-309.
 25. McGinnis JM, Foege WH. Actual causes of death in the United States. *JAMA.* 1993;270:2207-2212.
 26. Broussard BA, Sugarman JR, Bachman-Carter K, et al. Toward comprehensive obesity prevention programs in Native American communities. *Obes Res.* 1995;3:2895-2975.
 27. Najjar MF, Kuczmarski RJ. Anthropometric data and prevalence of overweight for Hispanics: 1982-1984. National Center for Health Statistics. *Vital Health Stat [11].* 1989;239:1-106.
 28. World Health Organization. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organ Tech Rep Ser.* 1995;854:1- 452.
 29. VanItallie TB, Lew EA. Overweight and Underweight. In: Lew EA, Gajewski J, eds. *Medical Risks: Trends in Mortality by Age and Time Elapsed.* Vol 1. New York: Praeger; 1990: Chapter 13.
 30. VanItallie TB. Health implications of overweight and obesity in the United States. *Ann Intern Med.* 1985;103:983-988.
 31. Manson JE, Stampfer MJ, Hennekens CH, Willett WC. Body weight and longevity. A reassessment. *JAMA.* 1987;257:353-358.

32. Troiano RP, Frongillo EA Jr, Sobal J, Levitsky DA. The relationship between body weight and mortality: a quantitative analysis of combined information from existing studies. *Int J Obes Relat Metab Disord.* 1996;20:63-75.
33. National Research Council. Committee on Diet and health. Implications for Reducing Chronic Disease Risk. Washington, DC: National Academy Press; 1989.
34. Forster JL, Jeffery RW, Schmid TL, Kramer FM. Preventing weight gain in adults: a pound of prevention. *Health Psychol.* 1988;7:515-525.
35. Lenfant C. NHLBI Clinical Guidelines: another look. *Circulation.* 1995;91:617-618.
36. Dickersin K, Min YI, Meinert CL. Factors influencing publication of research results. Follow-up of applications submitted to two institutional review boards. *JAMA.* 1992;267:374-378.
37. Dickersin K. The existence of publication bias and risk factors for its occurrence. *JAMA.* 1990;263:1385-1389.
38. Dickersin K, Min YI. Publication bias: the problem that won't go away. *Ann N Y Acad Sci.* 1993;703:135-146.
39. Allison DB, Faith MS, Gorman BS. Publication bias in obesity treatment trials? *Int J Obes Relat Metab Disord.* 1996;20:931-937.
40. Bray GA. Obesity in Perspective: a Conference. John E. Fogarty International Center for Advanced Study in the Health Sciences. Washington, DC: Govt Print Office; 1975. DHEW publication no. (NIH) 75-708.
41. Bray GA. Obesity in America. An overview of the Second Fogarty International Center conference on obesity. *Int J Obes.* 1979;3:363-375.
42. Health implications of obesity. National Institutes of Health Consensus Development Conference Statement. *Ann Intern Med.* 1985;103:1073-1077.
43. U.S. Department of Health and Human Services. Healthy people 2000: National health promotion and disease prevention objectives: Washington, DC: Public Health Service; 1991. DHHS publication no. (PHS) 91-50213.
44. Thomas PR, ed. Weighing the Options. Criteria for Evaluating Weight-Management Programs. Washington, DC: National Academy Press; 1995.
45. Najjar MF, Rowland M. Anthropometric reference data and prevalence of overweight, United States, 1976-80. *Vital Health Stat [11].* 1987;238:1-73.
46. Wolf AM, Colditz GA. Social and economic effects of body weight in the United States. *Am J Clin Nutr.* 1996;63:466S-469S.
47. Flynn MA, Nolph GB, Baker AS, Krause G. Aging in humans: a continuous 20-year study of physiologic and dietary parameters. *J Am Coll Nutr.* 1992;11:660-672.
48. Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *Int J Obes.* 1998;22:39-47.
49. Centers for Disease Control. Prevalence of overweight among adolescents—United States, 1988-91. *MMWR Morb Mortal Wkly Rep.* 1994;43:818-821.
50. McGarvey ST. Obesity in Samoans and a perspective on its etiology in Polynesians. *Am J Clin Nutr.* 1991;53:1586S-1594S.

51. Klatsky AL, Armstrong MA. Cardiovascular risk factors among Asian Americans living in northern California. *Am J Public Health.* 1991;81:1423-1428.
52. Welty TK, Lee ET, Yeh J, et al. Cardiovascular disease risk factors among American Indians. The Strong Heart Study. *Am J Epidemiol.* 1995;142:269-287.
53. Leigh JP, Fries JF, Hubert HB. Gender and race differences in the correlation between body mass and education in the 1971-1975 NHANES I. *J Epidemiol Community Health.* 1992;46:191-196.
54. Kuczmarski RJ. Prevalence of overweight and weight gain in the United States. *Am J Clin Nutr.* 1992;55:495S-502S.
55. Kuczmarski RJ, Flegal KM, Campbell SM, Johnson CL. Increasing prevalence of overweight among US adults. The National Health and Nutrition Examination Surveys, 1960 to 1991. *JAMA.* 1994;272:205-211.
56. Pi-Sunyer FX. Medical hazards of obesity. *Ann Intern Med.* 1993;119:655-660.
57. Seidell JC. The impact of obesity on health status: some implications for health care costs. *Int J Obes Relat Metab Disord.* 1995;19 (Suppl 6):S13-S16.
58. Wolf AM, Colditz GA. Current estimates of the economic costs of obesity in the United States. *Obes Res.* 1998;6:97-106.
59. Wolf AM, Colditz GA. The cost of obesity: the U.S. perspective. *Pharmacoeconomics.* 1994;5:34-37.
60. Colditz GA. Economic costs of obesity. *Am J Clin Nutr.* 1992;55:503S-507S.
61. Levy E, Levy P, Le Pen C, Basdevant A. The economic cost of obesity: the French situation. *Int J Obes Relat Metab Disord.* 1995;19:788-792.
62. Segal L, Carter R, Zimmet P. The cost of obesity - the Australian perspective. *Pharmacoeconomics.* 1994;5:45-52.
63. Seidell JC. Obesity in Europe. *Obes Res.* 1995;3:89S-93S.
64. National Task Force on Prevention and Treatment of Obesity. Towards Prevention of Obesity: Research Directions. *Obes Res.* 1994;2:571-584.
65. World Health Organization. Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation of Obesity. Geneva, 3-5 June 1997.
66. Winkleby MA, Feldman HA, Murray DM. Joint analysis of three U.S. community intervention trials for reduction of cardiovascular disease risk. *J Clin Epidemiol.* 1997;50:645-658.
67. Carleton RA, Lasater TM, Assaf AR, Feldman HA, McKinlay S. The Pawtucket Heart Health Program: community changes in cardiovascular risk factors and projected disease risk. *Am J Public Health.* 1995;85:777-785.
68. Davis SM. Cardiovascular Health Promotion Project. American Indian Program Recaptures Past to Improve Children's Future (Pathways). *Heartmemo (Special Edition).* 1996:25-27.
69. Epstein LH, Valoski A, Wing RR, McCurley J. Ten-year followup of behavioral, family-based treatment for obese children. *JAMA.* 1990;264:2519-2523.
70. Pan XR, Li GW, Hu YH, et al. Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care.* 1997;20:537-544.

71. James WP. A public health approach to the problem of obesity. *Int J Obes Relat Metab Disord.* 1995;19 (Suppl 3):S37-S45.
72. Jeffery RW. Population perspectives on the prevention and treatment of obesity in minority populations. *Am J Clin Nutr.* 1991;53 (6 Suppl):1621S-1624S.
73. Lewin-Epstein N. Determinants of regular source of health care in Black, Mexican, Puerto Rican, and non-Hispanic White population. *Med Care.* 1991;29(6):543-557.
74. O'Hare WP. America's Minorities—The demographics of diversity. *Population Bulletin.* Washington, DC: Population Reference Bureau, Inc.; 1992;47:4.
75. Link BG, Phelan JC. Understanding sociodemographic differences in health—the role of fundamental social causes. *Am J Public Health.* 1996;86:471-473.
76. Ballantyne D, Devine BL, Fife R. Interrelation of age, obesity, cigarette smoking, and blood pressure in hypertensive patients. *BMJ.* 1978;1:880-881.
77. Brennan PJ, Simpson JM, Blacket RB, McGilchrist CA. The effects of body weight on serum cholesterol, serum triglycerides, serum urate, and systolic blood pressure. *Aust N Z J Med.* 1980;10:15-20.
78. Havlik RJ, Hubert HB, Fabsitz RR, Feinleib M. Weight and hypertension. *Ann Intern Med.* 1983;98:855-859.
79. Loggie JM, Horan MJ, Hohn AR, Gruskin AB, Dunbar JB, Havlik RJ. Juvenile hypertension: highlights of a workshop. *J Pediatr.* 1984;104:657-663.
80. MacMahon SW, Blacket RB, Macdonald GJ, Hall W. Obesity, alcohol consumption and blood pressure in Australian men and women. The National Heart Foundation of Australia Risk Factor Prevalence Study. *J Hypertens.* 1984;2:85-91.
81. Colditz GA, Willett WC, Stampfer MJ, et al. Weight as a risk factor for clinical diabetes in women. *Am J Epidemiol.* 1990;132:501-513.
82. Chan JM, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes Care.* 1994;17:961-969.
83. Colditz GA, Willett WC, Rotnitzky A, Manson JE. Weight gain as a risk factor for clinical diabetes mellitus in women. *Ann Intern Med.* 1995;122:481-486.
84. Ford ES, Williamson DF, Liu S. Weight change and diabetes incidence: findings from a national cohort of US adults. *Am J Epidemiol.* 1997;146:214-222.
85. Despres JP, Nadeau A, Tremblay A, et al. Role of deep abdominal fat in the association between regional adipose tissue distribution and glucose tolerance in obese women. *Diabetes.* 1989;38:304-309.
86. Haffner SM, Mitchell BD, Hazuda HP, Stern MP. Greater influence of central distribution of adipose tissue on incidence of non-insulin-dependent diabetes in women than men. *Am J Clin Nutr.* 1991;53:1312-1317.
87. Sparrow D, Borkan GA, Gerzof SG, Wisniewski C, Silbert CK. Relationship of fat distribution to glucose tolerance. Results of computed tomography in male participants of the Normative Aging Study. *Diabetes.* 1986;35:411-415.
88. Lundgren H, Bengtsson C, Blohme G, Lapidus L, Sjostrom L. Adiposity and adipose tissue distribution in relation to incidence of diabetes in women: results from a

- prospective population study in Gothenburg, Sweden. *Int J Obes.* 1989;13:413-423.
89. Ohlson LO, Larsson B, Svardsudd K, et al. The influence of body fat distribution on the incidence of diabetes mellitus. 13.5 years of follow-up of the participants in the study of men born in 1913. *Diabetes.* 1985;34:1055-1058.
 90. Willett WC, Manson JE, Stampfer MJ, et al. Weight, weight change, and coronary heart disease in women. Risk within the 'normal' weight range. *JAMA.* 1995;273:461-465.
 91. Carman WJ, Sowers M, Hawthorne VM, Weissfeld LA. Obesity as a risk factor for osteoarthritis of the hand and wrist: a prospective study. *Am J Epidemiol.* 1994;139:119-129.
 92. Felson DT, Anderson JJ, Naimark A, Walker AM, Meenan RF. Obesity and knee osteoarthritis. The Framingham Study. *Ann Intern Med.* 1988;109:18-24.
 93. Felson DT, Zhang Y, Anthony JM, Naimark A, Anderson JJ. Weight loss reduces the risk for symptomatic knee osteoarthritis in women. The Framingham Study. *Ann Intern Med.* 1992;116:535-539.
 94. Felson DT. Weight and osteoarthritis. *J Rheumatol Suppl.* 1995;43:7-9.
 95. McGoey BV, Deitel M, Saplys RJ, Kliman ME. Effect of weight loss on musculoskeletal pain in the morbidly obese. *J Bone Joint Surg [Br].* 1990;72-B:322-323.
 96. Chua W, Chediak AD. Obstructive sleep apnea. Treatment improves quality of life—and may prevent death. *Postgrad Med.* 1994;95:123-126, 131, 135-138.
 97. Loubé DI, Loubé AA, Mitler MM. Weight loss for obstructive sleep apnea: the optimal therapy for obese patients. *J Am Diet Assoc.* 1994;94:1291-1295.
 98. Davies RJ, Stradling JR. The relationship between neck circumference, radiographic pharyngeal anatomy, and the obstructive sleep apnoea syndrome. *Eur Respir J.* 1990;3:509-514.
 99. Giovannucci E, Ascherio A, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. Physical activity, obesity, and risk for colon cancer and adenoma in men. *Ann Intern Med.* 1995;122:327-334.
 100. Graham S, Marshall J, Haughey B, et al. Dietary epidemiology of cancer of the colon in western New York. *Am J Epidemiol.* 1988;128:490-503.
 101. Klatsky AL, Armstrong MA, Friedman GD, Hiatt RA. The relations of alcoholic beverage use to colon and rectal cancer. *Am J Epidemiol.* 1988;128:1007-1015.
 102. Lee IM, Paffenbarger RS Jr. Quetelet's index and risk of colon cancer in college alumni. *J Natl Cancer Inst.* 1992;84:1326-1331.
 103. Le Marchand L, Wilkens LR, Mi MP. Obesity in youth and middle age and risk of colorectal cancer in men. *Cancer Causes Control.* 1992;3:349-354.
 104. Martinez ME, Giovannucci E, Spiegelman D, et al. Physical activity, body size, and colorectal cancer in women. *Am J Epidemiol.* 1996;146:S73.
 105. Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH. Long-term morbidity and mortality of overweight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935. *N Engl J Med.* 1992;327:1350-1355.
 106. Phillips RL, Snowdon DA. Dietary rela-

- tionships with fatal colorectal cancer among Seventh-Day Adventists. *J Natl Cancer Inst.* 1985;74:307-317.
107. Giovannucci E, Colditz GA, Stampfer MJ, Willett WC. Physical activity, obesity, and risk of colorectal adenoma in women (United States). *Cancer Causes Control.* 1996;7:253-263.
 108. Giovannucci E. Insulin and colon cancer. *Cancer Causes Control.* 1995;6:164-179.
 109. Chu SY, Lee NC, Wingo PA, Senie RT, Greenberg RS, Peterson HB. The relationship between body mass and breast cancer among women enrolled in the Cancer and Steroid Hormone Study. *J Clin Epidemiol.* 1991;44:1197-1206.
 110. Helmrich SP, Shapiro S, Rosenberg L, et al. Risk factors for breast cancer. *Am J Epidemiol.* 1983;117:35-45.
 111. Rosenberg L, Palmer JR, Miller DR, Clarke EA, Shapiro S. A case-control study of alcoholic beverage consumption and breast cancer. *Am J Epidemiol.* 1990;131:6-14.
 112. Willett WC, Browne ML, Bain C, et al. Relative weight and risk of breast cancer among premenopausal women. *Am J Epidemiol.* 1985;122:731-740.
 113. Hunter DJ, Willett WC. Diet, body size, and breast cancer. *Epidemiol Rev.* 1993;15:110-132.
 114. Huang Z, Hankinson SE, Colditz GA, et al. Dual effects of weight and weight gain on breast cancer risk. *JAMA.* 1997;278:1407-1411.
 115. Schottenfeld D, Fraumeni JF. *Cancer Epidemiology and Prevention.* New York: Oxford University Press, 1996.
 116. Hartz AJ, Barboriak PN, Wong A, Katayaa KP, Rimm AA. The association of obesity with infertility and related menstrual abnormalities in women. *Int J Obes.* 1979;3:57-73.
 117. Rich-Edwards JW, Goldman MB, Willett WC, et al. Adolescent body mass index and infertility caused by ovulatory disorder. *Am J Obstet Gynecol.* 1994;171:171-177.
 118. Dunaif A. *Polycystic Ovary Syndrome.* Boston: Blackwell Scientific Publications; 1992.
 119. Garbacia JA Jr., Richter M, Miller S, Barton JJ. Maternal weight and pregnancy complications. *Am J Obstet Gynecol.* 1985;152:238-245.
 120. Keppel KG, Taffel SM. Pregnancy-related weight gain and retention: implications of the 1990 Institute of Medicine guidelines. *Am J Public Health.* 1993;83:1100-1103.
 121. Smith DE, Lewis CE, Caveny JL, Perkins LL, Burke GL, Bild DE. Longitudinal changes in adiposity associated with pregnancy. The CARDIA Study. Coronary Artery Risk Development in Young Adults Study. *JAMA.* 1994;271:1747-1751.
 122. Johnson SR, Kolberg BH, Varner MW, Railsback LD. Maternal obesity and pregnancy. *Surg Gynecol Obstet.* 1987;164:431-437.
 123. Prentice A, Goldberg G. Maternal obesity increases congenital malformations. *Nutr Rev.* 1996;54:146-152.
 124. Stunkard AJ, Wadden TA. Psychological aspects of human obesity. In: Bjorntorp P, Brodoff BN, eds. *Obesity.* Philadelphia: Lippincott; 1992:352-360.
 125. Brownell KD, Fairburn C. Psychosocial consequences of obesity. In: Stunkard AJ, Sobal JG, eds. *Eating Disorders and Obesity: A Comprehensive Handbook.*

- New York: Guilford Press; 1995:417-421.
126. Wadden TA, Stunkard AJ. Psychosocial consequences of obesity and dieting-research and clinical findings. In: Stunkard AJ, Wadden TA, eds. *Obesity Theory and Therapy*. New York: Raven Press; 1993:163-177 .
 127. Faith MS, Allison DB. Assessment of psychological status among obese persons. In: Thompson JK, ed. *Body Image, Eating Disorders, and Obesity: An Integrative Guide For Assessment and Treatment*. Washington, DC: American Psychological Association; 1996:365-387.
 128. Allon N. The stigma of overweight in everyday life. In: Wolman BB, DeBerry S, eds. *Psychological Aspects of Obesity: A Handbook*. New York: Van Nostrand Reinhold; 1982:130-174.
 129. Feinleib M. Epidemiology of obesity in relation to health hazards. *Ann Intern Med*. 1985;103:1019-1024.
 130. Garrison RJ, Castelli WP. Weight and 30-year mortality of men in the Framingham Study. *Ann Intern Med*. 1985;103:1006-1009.
 131. Rabkin SW, Mathewson FA, Hsu PH. Relation of body weight to development of ischemic heart disease in a cohort of young North American men after a 26 year observation period: the Manitoba Study. *Am J Cardiol*. 1977;39:452-458.
 132. Cutler JA, Psaty BM, MacMahon S, Furberg CD. Public health issues in hypertension control: what has been learned from clinical trials. In: Laragh JH, Brenner BM, eds. *Hypertension: Pathophysiology, Diagnosis, and Management*. New York: Raven Press; 1995:253-270.
 133. Denke MA, Sempos CT, Grundy SM. Excess body weight. An underrecognized contributor to high blood cholesterol levels in white American men. *Arch Intern Med*. 1993;153:1093-1103.
 134. Denke MA, Sempos CT, Grundy SM. Excess body weight. An underrecognized contributor to dyslipidemia in white American women. *Arch Intern Med*. 1994;154:401- 410.
 135. Ashley FW Jr, Kannel WB. Relation of weight change to changes in atherogenic traits: the Framingham Study. *J Chronic Dis*. 1974;27:103-114.
 136. Hershcopf RJ, Elahi D, Andres R, et al. Longitudinal changes in serum cholesterol in man: an epidemiologic search for an etiology. *J Chronic Dis*. 1982;35:101-114.
 137. Shekelle RB, Shryock AM, Paul O, et al. Diet, serum cholesterol, and death from coronary heart disease. The Western Electric study. *N Engl J Med*. 1981;304:65-70.
 138. Manson JE, Colditz GA, Stampfer MJ, et al. A prospective study of obesity and risk of coronary heart disease in women. *N Engl J Med*. 1990;322:882-889.
 139. Reeder BA, Angel A, Ledoux M, Rabkin SW, Young TK, Sweet LE. Obesity and its relation to cardiovascular disease risk factors in Canadian adults. Canadian Heart Health Surveys Research Group. *Can Med Assoc J*. 1992;146:2009-2019.
 140. Carlson LA, Lindstedt S. The Stockholm prospective study. 1. The initial values for plasma lipids. *Acta Med Scand Suppl*. 1968;493:1-135.
 141. Mann JI, Lewis B, Shepherd J, et al. Blood lipid concentrations and other cardiovascular risk factors: distribution, prevalence, and detection in Britain. *BMJ*.

- 1988;296:1702-1706.
142. National Cholesterol Education Program. Second Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II). *Circulation*. 1994;89:1333-1445.
 143. Glueck CJ, Taylor HL, Jacobs D, Morrison JA, Beaglehole R, Williams OD. Plasma high-density lipoprotein cholesterol: association with measurements of body mass. The Lipid Research Clinics Program Prevalence Study. *Circulation*. 1980;62:IV-62-69.
 144. Garrison RJ, Wilson PW, Castelli WP, Feinleib M, Kannel WB, McNamara PM. Obesity and lipoprotein cholesterol in the Framingham offspring study. *Metabolism*. 1980;29:1053-1060.
 145. Anderson KM, Wilson PWF, Garrison RJ, Castelli WP. Longitudinal and secular trends in lipoprotein cholesterol measurements in a general population sample: The Framingham offspring study. *Atherosclerosis*. 1987;68:59-66.
 146. Law MR, Wald NJ, Thompson SG. By how much and how quickly does reduction in serum cholesterol concentration lower risk of ischaemic heart disease? *BMJ*. 1994;308:367-773.
 147. Rydker PM, Hennekens CH, Stampfer MJ. A prospective study of Liprotein (a) and the risk for myocardial infarction. *JAMA*. 1993;270:2195-2199.
 148. Miller BD, Alderman EL, Haskell WL, Fair JM, Krauss RM. Predominance of dense low-density lipoprotein particles predicts angiographic benefit of therapy in the Stanford coronary risk intervention study. *Circulation*. 1996;94:2146-2153.
 149. Lamarche B, Tchernof A, Moorjani S, et al. Small, dense low-density lipoprotein particles as a predictor of the risk of ischemic heart disease in men. Prospective results from the Quebec Cardiovascular Study. *Circulation*. 1997;95:69-75.
 150. Reaven GM, Chen YD, Jeppesen J, Maheux P, Krauss RM. Insulin resistance and hyperinsulinemia in individuals with small, dense low density lipoprotein particles. *J Clin Invest*. 1993;92:141-146.
 151. Selby JV, Austin MA, Newman B, et al. LDL subclass phenotypes and the insulin resistance syndrome in women. *Circulation*. 1993;88:381-387.
 152. Tchernof A, Lamarche B, Prud'Homme D, et al. The dense LDL phenotype: association with plasma lipoprotein levels, visceral obesity, and hyperinsulinemia in men. *Diabetes Care*. 1996;19(6):629-637.
 153. Knowler WC, Pettitt DJ, Savage PJ, Bennett PH. Diabetes incidence in Pima Indians: contributions of obesity and parental diabetes. *Am J Epidemiol*. 1981;113:144-156.
 154. Lee ET, Howard BV, Savage PJ, et al. Diabetes and impaired glucose tolerance in three American Indian populations aged 45-74 years. The Strong Heart Study. *Diabetes Care*. 1995;18:599-610.
 155. Higgins M, Kannel W, Garrison R, Pinsky J, Stokes J 3rd. Hazards of obesity—the Framingham experience. *Acta Med Scand Suppl*. 1988;723:23-36.
 156. Donahue RP, Abbott RD. Central obesity and coronary heart disease in men. *Lancet*. 1987;2:1215.
 157. Ducimetiere P, Richard JL. The relationship between subsets of anthropometric upper versus lower body measurements and

- coronary heart disease risk in middle-aged men. The Paris Prospective Study I. *Int J Obes*. 1989;13:111-121.
158. Lapidus L, Bengtsson C, Larsson B, Pennert K, Rybo E, Sjoström L. Distribution of adipose tissue and risk of cardiovascular disease and death: a 12 year follow up of participants in the population study of women in Gothenburg, Sweden. *BMJ*. 1984;289:1257-1261.
 159. Larsson B, Svardsudd K, Welin L, Wilhelmsen L, Björntorp P, Tibblin G. Abdominal adipose tissue distribution, obesity, and risk of cardiovascular disease and death: 13 year follow up of participants in the study of men born in 1913. *BMJ*. 1984;288:1401-1404.
 160. Björntorp P. The associations between obesity, adipose tissue distribution, and disease. *Acta Med Scand Suppl*. 1988;723:121-134.
 161. Fujioka S, Matsuzawa Y, Tokunaga K, Tarui S. Contribution of intra-abdominal fat accumulation to the impairment of glucose and lipid metabolism in human obesity. *Metabolism*. 1987;36:54-59.
 162. Shaper AG, Wannamethee SG, Walker M. Body weight: implications for the prevention of coronary heart disease, stroke, and diabetes mellitus in a cohort study of middle aged men. *BMJ*. 1997;314:1311-1317.
 163. Tokunaga K, Matsuzawa Y, Kotani K, et al. Ideal body weight estimated from the body mass index with the lowest morbidity. *Int J Obes*. 1991;15:1-5.
 164. Jousilahti P, Tuomilehto J, Vartiainen E, Pekkanen J, Puska P. Body weight, cardiovascular risk factors, and coronary mortality. 15-year follow-up of middle-aged men and women in eastern Finland. *Circulation*. 1996;93:1372-1379.
 165. DiBianco R. The changing syndrome of heart failure: an annotated review as we approach the 21st century. *J Hypertens Suppl*. 1994;12:S73-S87.
 166. Kannel WB, Cupples A. Epidemiology and risk profile of cardiac failure. *Cardiovasc Drugs Ther*. 1988;2 (Suppl 1):387-395.
 167. Savage MP, Krolewski AS, Kenien GG, Lebeis MP, Christlieb AR, Lewis SM. Acute myocardial infarction in diabetes mellitus and significance of congestive heart failure as a prognostic factor. *Am J Cardiol*. 1988;62:665-669.
 168. Eriksson H, Svardsudd K, Larsson B, et al. Risk factors for heart failure in the general population: the study of men born in 1913. *Eur Heart J*. 1989;10:647-656.
 169. Eriksson H, Wilhelmsen L, Caidahl K, Svardsudd K. Epidemiology and prognosis of heart failure. *Z Kardiol*. 1991;80 (Suppl 8):1-6.
 170. Shimizu M, Isogai Y. Heart failure due to metabolic heart disorders. *Nippon Rinsho*. 1993;51:1362-1366.
 171. Urbina EM, Gidding SS, Bao W, Pickoff AS, Berdusis K, Berenson GS. Effect of body size, ponderosity, and blood pressure on left ventricular growth in children and young adults in the Bogalusa Heart Study. *Circulation*. 1995;91:2400-2406.
 172. Alpert MA, Hashimi MW. Obesity and the heart. *Am J Med Sci*. 1993;306:117-123.
 173. Garavaglia GE, Messerli FH, Nunez BD, Schmieder RE, Grossman E. Myocardial contractility and left ventricular function in obese patients with essential hypertension. *Am J Cardiol*. 1988;62:594-597.
 174. Ritter MM, Schraudolph M, Richter WO, Herbert M, Wiebecke B, Schwandt P. Obesity, heart failure and pulmonary insuf-

- iciency in a 26-year-old female. *Med Klin*. 1990;85:371-375.
175. Huang C, Ross PD, Lydick E, Wasnich RD. Factors associated with joint pain among postmenopausal women. *Int J Obes Relat Metab Disord*. 1997;21:349-354.
 176. Williams RA, Foulsham BM. Weight reduction in osteoarthritis using phentermine. *Practitioner*. 1981;225:231-232.
 177. Garfinkel L. Overweight and mortality. *Cancer*. 1986;58:1826-1829.
 178. Williamson DF, Madans J, Pamuk E, Flegal KM, Kendrick JS, Serdula MK. A prospective study of childbearing and 10-year weight gain in US white women 25 to 45 years of age. *Int J Obes Relat Metab Disord*. 1994;18:561-569.
 179. Cnattingius S, Bergstrom R, Lipworth L, Kramer MS. Prepregnancy weight and the risk of adverse pregnancy outcomes. *N Eng J Med*. 1998;338:147-152.
 180. Shils ME, Olson JA, Shike M, eds. *Modern Nutrition in Health and Disease*. Philadelphia: Lea & Febiger; 1994.
 181. Cogswell ME, Serdula MK, Hungerford DW, Yip R. Gestational weight gain among average-weight and overweight women—what is excessive? *Am J Obstet Gynecol*. 1995;172:705-712.
 182. Institute of Medicine. Committee on Nutritional Status During Pregnancy and Lactation. *Nutrition During Pregnancy: Part I, Weight Gain; Part II, Nutrient Supplements*. Washington, DC: National Academy Press, 1990.
 183. O'Neil PM, Jarrell MP. Psychological aspects of obesity and dieting. In: Wadden TA, Van Itallie TB, eds. *Treatment Of The Seriously Obese Patient*. New York: Guilford Press; 1992:252-270.
 184. Jarvie GJ, Lahey BB, Graziano W, Frammer E. Childhood obesity and social stigma: what we know and what we don't know. *Dev Rev*. 1983;3:237-273.
 185. DeJong W, Kleck RE. The Social Psychological Effects of Overweight. In: Herman CP, Zanna MP, Higgins ET, eds. *Physical Appearance, Stigma, and Social Behavior*. Hillsdale, NJ: L. Erlbaum; 1986:65-88.
 186. Goodman N, Dornbusch SM, Richardson SA, Hastorf AH. Variant reactions to physical disabilities. *Amer Sociolo Rev*. 1963;28:429-435.
 187. Lerner RM, Gellert E. Body build identification, preference, and aversion in children. *Dev Psychol*. 1969;5:456-462.
 188. Richardson SA, Hastorf AH, Goodman N, Dornbusch SM. Cultural uniformity in reaction to physical disabilities. *Am Soc Rev*. 1961;26:241-247.
 189. Staffieri JR. Body build and behavioral expectancies in young females. *Dev Psychol*. 1972;6:125-127.
 190. Staffieri JR. A study of social stereotype of body image in children. *J Pers Soc Psychol*. 1967;7:101-104.
 191. Lerner RM, Korn SJ. The development of body build stereotypes in males. *Child Dev*. 1972;43:908-920.
 192. Blumberg P, Mellis LP. Medical students' attitudes toward the obese and the morbidly obese. *Int J Eat Disord*. 1985;4:169-175.
 193. Maddox GL, Liederman V. Overweight as a social disability with medical implications. *J Med Educ*. 1969;44:214-220.
 194. Rand CS, Macgregor AM. Morbidly obese patients' perceptions of social discrimina-

- tion before and after surgery for obesity. *South Med J.* 1990;83:1390-1395.
195. Crandall CS, Biernat M. The ideology of anti-fat attitudes. *J Appl Soc Psychol.* 1990;20:227-243.
 196. Maddox GL, Back KW, Liederman WR. Overweight as social deviance and disability. *J Health Soc Behav.* 1968;9:287-298.
 197. Larkin JC, Pines HA. No fat persons need apply: experimental studies of the overweight stereotype and hiring preference. *Sociology of Work and Occupations.* 6:312-327.
 198. Pingitore R, Dugoni BL, Tindale RS, Spring B. Bias against overweight job applicants in a simulated employment interview. *J Appl Psychol.* 1994;79:909-917.
 199. Roe DA, Eickwort KR. Relationships between obesity and associated health factors with unemployment among low income women. *J Am Med Womens Assoc.* 1976;31:193-194, 198-199, 203-204.
 200. Canning H, Mayer J. Obesity: its possible effect on college acceptance. *N Eng J Med.* 1966;275:1172-1174.
 201. Crandall CS. Do heavy-weight students have more difficulty paying for college? *Personality Soc Psychol Bull.* 1991;17:606-611.
 202. Sargent JD, Blanchflower DG. Obesity and stature in adolescence and earnings in young adulthood. Analysis of a British birth cohort. *Arch Pediatr Adolesc Med.* 1994;148:681-687.
 203. Karris L. Prejudice against obese renters. *J Soc Psych.* 1977;101:159-160.
 204. Gortmaker SL, Must A, Perrin JM, Sobol AM, Dietz WH. Social and economic consequences of overweight in adolescence and young adulthood. *N Engl J Med.* 1993;329:1008-1012.
 205. Massara EB, Stunkard AJ. A method of quantifying cultural ideals of beauty and the obese. *Int J Obes.* 1979;3:149-152.
 206. Crandall CS, Martinez R. Culture, ideology, and antifat attitudes. *Personality Soc Psychol Bull.* 1996;22:1165-1176.
 207. Striegel-Moore R, Rodin J. The influences of psychological variables in obesity. In: Brownell KD, Foreyt JP, eds. *Handbook of Eating Disorders: Physiology, Psychology, and Treatment of Obesity, Anorexia, and Bulimia.* New York: Basic Books; 1986:99-121.
 208. Wadden TA, Stunkard AJ. Psychopathology and obesity. *Ann N Y Acad Sci.* 1987;499:55-65.
 209. Lapidus L, Bengtsson C, Hallstrom T, Bjorntorp P. Obesity, adipose tissue distribution and health in women—results from a population study in Gothenburg, Sweden. *Appetite.* 1989;12:25-35.
 210. Lissau I, Sorensen TI. Parental neglect during childhood and increased risk of obesity in young adulthood. *Lancet.* 1994;343:324-327.
 211. Sullivan M, Karlsson J, Sjostrom L, et al. Swedish obese subjects (SOS)—an intervention study of obesity. Baseline evaluation of health and psychosocial functioning in the first 1743 subjects examined. *Int J Obes Relat Metab Disord.* 1993;17:503-512.
 212. Friedman MA, Brownell KD. Psychological correlates of obesity: moving to the next research generation. *Psychol Bull.* 1995;117:3-20.

213. Stunkard AJ, Rush J. Dieting and depression reexamined: a critical review of reports of untoward responses during weight reduction for obesity. *Annals Intern Med.* 1974;81:526- 533.
214. Prather RC, Williamson DA. Psychopathology associated with bulimia, binge eating, and obesity. *Int J Eat Disord.* 1988;7:177-184.
215. Fitzgibbon ML, Stolley MR, Kirschenbaum DS. Obese people who seek treatment have different characteristics than those who do not seek treatment. *Health Psychol.* 1993;12:342-345.
216. Gormally J, Black S, Daston S, Rardin D. The assessment of binge eating severity among obese persons. *Addict Behav.* 1982;7:47-55.
217. Loro AD Jr, Orleans CS. Binge eating in obesity: preliminary findings and guidelines for behavioral analysis and treatment. *Addict Behav.* 1981;6:155-166.
218. Marcus MD, Wing RR, Fairburn CG. Cognitive treatment of binge eating versus behavioral weight control in the treatment of binge eating disorder. *Annals of Behavioral Medicine.* 1995;17:S090.
219. Spitzer RL, Devlin M, Walsh BT, et al. Binge eating disorder: a multisite field trial of the diagnostic criteria. *Int J Eat Disord.* 1992;11:191-203.
220. Yanovski SZ, Gormally JF, Leser MS, Gwirtsman HE, Yanovski JA. Binge eating disorder affects outcome of comprehensive very-low-calorie diet treatment. *Obes Res.* 1994;2:205- 212.
221. Spitzer RL, Yanovski S, Wadden T, et al. Binge eating disorder: its further validation in a multisite study. *Int J Eat Disord.* 1993;13:137-153.
222. Wilfley DE, Agras WS, Telch CF, et al. Group cognitive-behavioral therapy and group interpersonal psychotherapy for the nonpurging bulimic individual: a controlled comparison. *J Consult Clin Psychol.* 1993;61:296-305.
223. Wilfley DE, Cohen LR. Psychological treatment of bulimia nervosa and binge eating disorder. *Psychopharmacol Bull.* 1997;33:437-453.
224. Yanovski SZ. Binge eating disorder: Current knowledge and future directions. *Obes Res.* 1993;1:306-324.
225. Yanovski SZ, Nelson JE, Dubbert BK, Spitzer RL. Association of binge eating disorder and psychiatric comorbidity in obese subjects. *Am J Psychiatry.* 1993;150:1472-1479.
226. Wilfley DE, Pike KM, Striegel-Moore R. Toward an integrated model of risk for binge eating disorder. *J Gender Cult Health.* 1997;2:1-32.
227. Fairburn CG, Wilson GT. *Binge Eating: Nature, Assessment, and Treatment.* New York: Guilford Press; 1993.
228. Castonguay LG, Eldredge KL, Agras WS. Binge eating disorder: current state and future directions. *Clini Psychol Rev.* 1995;15:865-890.
229. Striegel-Moore RH, Wilson GT, Wilfley DE, Elder KA, Brownell KD. Binge eating in an obese community sample. *Int J Eat Disord.* 1998;23:27-37.
230. Telch CF, Agras WS, Rossiter EM. Binge eating increases with increasing adiposity. *Int J Eat Disord.* 1988;7:115-119.
231. Kenardy J, Arnow B, Agras WS. The aversiveness of specific emotional states associated with binge-eating in obese subjects. *Aust N Z J Psychiatry.* 1996;30:839-844.

232. Marcus MD, Wing RR, Ewing L, Kern E, Gooding W, McDermott M. Psychiatric disorders among obese binge eaters. *Int J Eat Disord.* 1990;9:69-77.
233. Marcus MD, Wing RR, Hopkins J. Obese binge eaters: affect, cognitions, and response to behavioural weight control. *J Consult Clin Psychol.* 1988;56:433-439.
234. Molinari E, Ragazzoni P, Morosin A. Psychopathology in obese subjects with and without binge-eating disorder and in bulimic subjects. *Psychol Rep.* 1997;80:1327-1335.
235. Mussell MP, Mitchell JE, de Zwaan M, Crosby RD, Seim HC, Crow SJ. Clinical characteristics associated with binge eating in obese females: a descriptive study. *Int J Obes Relat Metab Disord.* 1996;20:324-331.
236. Specker S, de Zwaan M, Raymond N, Mitchell J. Psychopathology in subgroups of obese women with and without binge eating disorder. *Compr Psychiatry.* 1994;35:185-190.
237. Brody ML, Walsh BT, Devlin MJ. Binge eating disorder: reliability and validity of a new diagnostic category. *J Consult Clin Psychol.* 1994;62:381-386.
238. de Zwaan M, Nutzinger DO, Schoenbeck G. Binge eating in overweight women. *Compr Psychiatry.* 1992;33:256-261.
239. Kuehnel RH, Wadden TA. Binge eating disorder, weight cycling, and psychopathology. *Int J Eat Disord.* 1994;15:321-329.
240. Keefe PH, Wyshogrod D, Weinberger E, Agras WS. Binge eating and outcome of behavioral treatment of obesity: a preliminary report. *Behav Res Ther.* 1984;22:319-321.
241. Garner DM, Wooley SC. Confronting the failure of behavioral and dietary treatments for obesity. *Clin Psychol Rev.* 1991;11:729-780.
242. Telch CF, Agras WS. The effects of a very low-calorie diet on binge eating. *Behav Therapy.* 1993;24:177-194.
243. Yanovski SZ, Sebring NG. Recorded food intake of obese women with binge eating disorder before and after weight loss. *Int J Eat Disord.* 1994;15:135-150.
244. Cash TF, Hicks KL. Being fat versus thinking fat: relationships with body image, eating behaviors, and well-being. *Cogni Ther Res.* 1990;14:327-341.
245. Collins JK. Methodology for the objective measurement of body image. *Int J Eat Disord.* 1987;6:393-399.
246. Collins JK, Beumont PJ, Touyz SW, Krass J, et al. Variability in body shape, perception in anorexic, bulimic, obese, and control subjects. *Int J Eat Disord.* 1987;6:633-638.
247. Gardner RM, Martinez R, Sandoval Y. Obesity and body image: an evaluation of sensory and non-sensory components. *Psychol Med.* 1987;17:927-932.
248. Glucksman ML, Hirsch J. The response of obese patients to weight reduction. 3. The perception of body size. *Psychosom Med.* 1969;31:1-7.
249. Garner DM, Garfinkel PE, Stancer HC, Moldofsky H. Body image disturbances in anorexia nervosa and obesity. *Psychosom Med.* 1976;38:327-336.
250. Grilo CM, Wilfley DE, Brownell KD, Rodin J. Teasing, body image, and self-esteem in a clinical sample of obese women. *Addict Behav.* 1994;19:443-450.
251. Mussell MP, Peterson CB, Weller CL,

- Crosby RD, de Zwaan M, Mitchell JE. Differences in body image and depression among obese women with and without binge eating disorder. *Obes Res.* 1996;4:431-439.
252. Stunkard A, Mendelson M. Obesity and the body image. 1. Characteristics of disturbances in the body image of some obese persons. *Am J Psychiatry.* 1967;123:1296-1300.
253. Stunkard A, Burt V. Obesity and the body image. II. Age at onset of disturbances in the body image. *Am J Psychiatry.* 1967;123:1443-1447.
254. Cash TF. The psychology of physical appearance: aesthetics, attributes, and images. In: Cash TF, Pruzinsky T, eds. *Body Images: Development, Deviance, and Change.* New York: Guilford Press, 1990:51-79.
255. Tiggemann M, Rothblum ED. Gender differences in social consequences of perceived overweight in the United States and Australia. *Sex Roles.* 1988;18:75-86.
256. Rosen JC. Improving body image in obesity. In: Thompson JK, ed. *Body Image, Eating Disorders, and Obesity.* Washington, DC: American Psychological Association; 1996:425- 440.
257. Kumanyika SK, Morssink CB. Cultural appropriateness of weight management programs. In: Dalton S, ed. *Overweight and Weight Management: The Health Professional's Guide To Understanding and Practice.* Gaithersburg, MD: Aspen Publishers; 1997:69-103.
258. Childress AC, Brewerton TD, Hodges EL, Jarrell MP. The Kids' Eating Disorders Survey (KEDS): a study of middle school students. *J Am Acad Child Adolesc Psychiatry.* 1993;32:843-850.
259. Kumanyika S, Wilson JF, Guilford-Davenport M. Weight-related attitudes and behaviors of black women. *J Am Diet Assoc.* 1993;93:416-422.
260. Powell AD, Kahn AS. Racial differences in women's desires to be thin. *Int J Eat Disord.* 1995;17:191-195.
261. Striegel-Moore RH, Schreiber GB, Pike KM, Wilfley DE, Rodin J. Drive for thinness in black and white preadolescent girls. *Int J Eat Disord.* 1995;18:59-69.
262. Desmond SM, Price JH, Hallinan C, Smith D. Black and white adolescents' perceptions of their weight. *J Sch Health.* 1989;59:353-358.
263. Kemper KA, Sargent RG, Drane JW, Valois RF, Hussey JW. Black and white females' perceptions of ideal body size and social norms. *Obes Res.* 1994;2:117-126.
264. Rand CS, Kulda JM. Morbid obesity: a comparison between a general population and obesity surgery patients. *Int J Obes Relat Metab Disord.* 1993;17:657-661.
265. Stevens J, Kumanyika SK, Keil JE. Attitudes toward body size and dieting: differences between elderly black and white women. *Am J Public Health.* 1994;84:1322-1325.
266. Wilfley DE, Schreiber GB, Pike KM, Striegel-Moore RH, Wright DJ, Rodin J. Eating disturbance and body image: a comparison of a community sample of adult black and white women. *Int J Eat Disord.* 1996;20:377-387.
267. Rucker CE, Cash TF. Body images, body-size perceptions, and eating behaviors among African-American and white college women. *Int J Eating Disord.* 1992;12:291-300.
268. Schreiber GG, Robins M, Striegel-Moore

- R, Obarzanek E, Morrison JA, Wright DJ. Weight modification efforts reported by black and white preadolescent girls: National Heart, Lung, and Blood Institute Growth and Health Study. *Pediatrics*. 1996;98:63-70.
269. Rosen JC, Gross J. Prevalence of weight reducing and weight gaining in adolescent girls and boys. *Health Psychol*. 1987;6:131-147.
270. Gittelsohn J, Harris SB, Thorne-Lyman AL, Hanley AJ, Barnie A, Zinman B. Body image concepts differ by age and sex in an Ojibway-Cree community in Canada. *J Nutr*. 1996;126:2990-3000.
271. Cooper RS, Ford E. Comparability of risk factors for coronary heart disease among blacks and whites in the NHANES-I Epidemiologic Follow-up Study. *Ann Epidemiol*. 1992;2:637-645.
272. Folsom AR, Burke GL, Byers CL, et al. Implications of obesity for cardiovascular disease in blacks: the CARDIA and ARIC studies. *Am J Clin Nutr*. 1991;53:1604S-1611S.
273. Fujimoto WY, Newell-Morris LL, Grote M, Bergstrom RW, Shuman WP. Visceral fat obesity and morbidity: NIDDM and atherogenic risk in Japanese-American men and women. *Int J Obes*. 1991;15 (Suppl 2):41-44.
274. Galanis DJ, McGarvey ST, Sobal J, Bausserman L, Levinson PD. Relations of body fat and fat distribution to the serum lipid, apolipoprotein and insulin concentrations of Samoan men and women. *Int J Obes Relat Metab Disord*. 1995;19:731-738.
275. Haffner SM, Diehl AK, Stern MP, Hazuda HP. Central adiposity and gallbladder disease in Mexican Americans. *Am J Epidemiol*. 1989;129:587-595.
276. Havas S, Fujimoto W, Close N, McCarter R, Keller J, Sherwin R. The NHLBI workshop on Hypertension in Hispanic Americans, Native Americans, and Asian/Pacific Islander Americans. *Public Health Rep*. 1996;111:451-458.
277. Howard BV, Lee ET, Cowan LD, et al. Coronary heart disease prevalence and its relation to risk factors in American Indians. The Strong Heart Study. *Am J Epidemiol*. 1995;142:254-268.
278. Lipton RB, Liao Y, Cao G, Cooper RS, McGee D. Determinants of incident non-insulin-dependent diabetes mellitus among blacks and whites in a national sample. The NHANES I Epidemiologic Follow-up Study. *Am J Epidemiol*. 1993;138:826-839.
279. Federation of American Societies for Experimental Biology. Life Sciences Research. Third Report on Nutrition Monitoring in the United States. Washington, DC: Interagency Board for Nutrition Monitoring and Related Research; 1995.
280. Ettinger WH, Wahl PW, Kuller LH, et al. Lipoprotein lipids in older people. Results from the Cardiovascular Health Study. The CHS Collaborative Research Group. *Circulation*. 1992;86:858-869.
281. Naser WB, Thomas J, Semanya K, Thomas DJ, Gillum RF. Obesity and hypertension in a longitudinal study of black physicians: the Meharry Cohort Study. *J Chronic Dis*. 1986;39:105-113.
282. Schmidt-Nowara WW, Coultas DB, Wiggins C, Skipper BE, Samet JM. Snoring in a Hispanic-American population. Risk factors and association with hypertension and other morbidity. *Arch*

- Intern Med. 1990;150:597-601.
283. Warne DK, Charles MA, Hanson RL, et al. Comparison of body size measurements as predictors of NIDDM in Pima Indians. *Diabetes Care*. 1995;18:435-439.
 284. Wei M, Gaskill SP, Haffner SM, Stern MP. Waist circumference as the best predictor of noninsulin dependent diabetes mellitus (NIDDM) compared to body mass index, waist/hip ratio, and other anthropometric measurements in Mexican Americans—a 7-year prospective study. *Obes Res*. 1997;5:16-23.
 285. Lee IM, Manson JE, Hennekens CH, Paffenbarger RS Jr. Body weight and mortality. A 27-year follow-up of middle-aged men. *JAMA*. 1993;270:2823-2828.
 286. Allison DB, Faith MS, Heo M, Kotler DP. Hypothesis concerning the U-shaped relation between body mass index and mortality. *Am J Epidemiol*. 1997;146:339-349.
 287. Stevens J, Cai J, Pamuk ER, Williamson DF, Thun MJ, Wood JL. The effect of age on the association between body-mass index and mortality. *N Engl J Med*. 1998;338:1-7.
 288. Diehr P, Bild DE, Harris TB, Duxbury A, Siscovick D, Rossi M. Body mass index and mortality in nonsmoking older adults: The cardiovascular health study. *Am J Public Health*. 1998;88:623-629.
 289. Cornoni-Huntley JC, Harris TB, Everett DF, et al. An overview of body weight of older persons, including the impact on mortality. The National Health and Nutrition Examination Survey I—Epidemiologic Follow-up Study. *J Clin Epidemiol*. 1991;44:743-753.
 290. Harris T, Cook EF, Garrison R, Higgins M, Kannel W, Goldman L. Body mass index and mortality among nonsmoking older persons. The Framingham Heart Study. *JAMA*. 1988;259:1520-1524.
 291. Baumgartner RN, Heymsfield SB, Roche AF. Human body composition and the epidemiology of chronic disease. *Obes Res*. 1995;3:73-95.
 292. Losonczy KG, Harris TB, Cornoni-Huntley J, et al. Does weight loss from middle age to old age explain the inverse weight mortality relation in old age? *Am J Epidemiol*. 1995;141:312-321.
 293. Fried LP, Kronmal RA, Newman AB, et al. Risk factors for 5-year mortality in older adults (for the Cardiovascular Health Study Collaborative Research Group). *JAMA*. 1998;279:585-592.
 294. Folsom AR, Kaye SA, Sellers TA, et al. Body fat distribution and 5-year risk of death in older women. *JAMA*. 1993;269:483-487.
 295. Willett WC, Stampfer M, Manson J, VanItallie T. New weight guidelines for Americans: justified or injudicious? *Am J Clin Nutr*. 1991;53:1102-1103.
 296. Roche AF. Sarcopenia: a critical review of its measurements and health-related significance in the middle-aged and elderly. *Am J Hum Biol*. 1994;6:33-42.
 297. Durazo-Arvizu RA, McGee DL, Cooper RS, Liao Y, Luke A. Mortality and optimal body mass index in a sample of the U.S. population. *Am J Epidemiol*. 1998;147:739-749.
 298. Sorokin JD, Zonderman AB, Costa PT Jr., Andres R. Twenty year follow-up of the NHANES I cohort: tests of methodologic hypotheses. *Obes Res*. 1996;4 Suppl (1): M46.
 299. Stern MP, Patterson JK, Mitchell BD,

- Haffner SM, Hazuda HP. Overweight and mortality in Mexican Americans. *Int J Obes*. 1990;14:623-629.
300. Johnson JL, Heineman EF, Heiss G, Hames CG, Tyroler HA. Cardiovascular disease risk factors and mortality among black women and white women aged 40-64 years in Evans County, Georgia. *Am J Epidemiol*. 1986;123:209-220.
301. Stevens J, Keil JE, Rust PF, Tyroler HA, Davis CE, Gazes PC. Body mass index and body girths as predictors of mortality in black and white women. *Arch Intern Med*. 1992;152:1257-1262.
302. Tyroler HA, Knowles MG, Wing SB, et al. Ischemic heart disease risk factors and 20-year mortality in middle-age Evans County black males. *Am Heart J*. 1984;108:738-746.
303. Wienpahl J, Ragland DR, Sidney S. Body mass index and 15-year mortality in a cohort of black men and women. *J Clin Epidemiol*. 1990;43:949-960.
304. Durazo-Arvizu R, Cooper RS, Luke A, Prewitt TE, Liao Y, McGee DL. Relative weight and mortality in U.S. blacks and whites: findings from representative national population samples. *Ann Epidemiol*. 1997;7:383-395.
305. Hodge AM, Dowse GK, Collins VR, Zimmet PZ. Mortality in Micronesian Nauruans and Melanesian and Indian Fijians is not associated with obesity. *Am J Epidemiol*. 1996;143:442-455.
306. Hanson RL, McCance DR, Jacobsson LT, et al. The U-shaped association between body mass index and mortality: relationship with weight gain in a Native-American population. *J Clin Epidemiol*. 1995;48:903-916.
307. Pettitt DJ, Lisse JR, Knowler WC, Bennett PH. Mortality as a function of obesity and diabetes mellitus. *Am J Epidemiol*. 1982;115:359-366.
308. Andres R, Muller DC, Sorkin JD. Long-term effects of change in body weight on all-cause mortality. A review. *Ann Intern Med*. 1993;119:737-743.
309. Weight cycling. National Task Force on the Prevention and Treatment of Obesity. *JAMA*. 1994;272:1196-1202.
310. Williamson DF, Pamuk ER. The association between weight loss and increased longevity. A review of the evidence. *Ann Intern Med*. 1993;119:731-736.
311. Williamson DF. "Weight cycling" and mortality: how do the epidemiologists explain the role of intentional weight loss? *J Am Coll Nutr*. 1996;15:6-13.
312. Williamson DF. Intentional weight loss: patterns in the general population and its association with morbidity and mortality. *Int J Obes Relat Metab Disord*. 1997;21 (Suppl 1):S14-S19.
313. French SA, Jeffery RW, Folsom AR, Williamson DF, Byers T. Weight variability in a population-based sample of older women: reliability and intercorrelation of measures. *Int J Obes Relat Metab Disord*. 1995;19:22-29.
314. French SA, Jeffery RW, Folsom AR, Williamson DF, Byers T. History of intentional and unintentional weight loss in a population-based sample of women aged 55 to 69 years. *Obes Res*. 1995;3:163-170.
315. French SA, Jeffery RW, Folsom AR, Williamson DF, Byers T. Relation of weight variability and intentionality of weight loss to disease history and health-related variables in a population-based

- sample of women aged 55-69 years. *Am J Epidemiol.* 1995;142:1306-1314.
316. Meltzer AA, Everhart JE. Unintentional weight loss in the United States. *Am J Epidemiol.* 1995;142:1039-1046.
317. Singh RB, Rastogi SS, Verma R, et al. Randomised controlled trial of cardioprotective diet in patients with recent acute myocardial infarction: results of 1-year follow up. *BMJ.* 1992;304:1015-1019.
318. Williamson DF, Pamuk E, Thun M, Flanders D, Byers T, Heath C. Prospective study of intentional weight loss and mortality in never-smoking overweight US white women aged 40-64 years. *Am J Epidemiol.* 1995;141:1128-1141.
319. Williamson DF, Pamuk E, Thun M, et al. Prospective study of international weight loss and mortality in overweight men aged 40-64 years. *Obes Res.* 1997;5:94S. [abstract].
320. Sjostrom L, Larsson B, Backman L, et al. Swedish obese subjects (SOS). Recruitment for an intervention study and a selected description of the obese state. *Int J Obes Relat Metab Disord.* 1992;16:465-479.
321. Sjostrom CD, Lissner L, Sjostrom L. Relationships between changes in body composition and changes in cardiovascular risk factors: the SOS intervention study. *Obes Res.* 1997;5:519-530.
322. Sjostrom L. Natural history of massive obesity. *Obes Res.* 1995;3:317S. [abstract].
323. Sobal J, Stunkard AJ. Socioeconomic status and obesity: A review of the literature. *Psychol Bull.* 1989;105:260-275.
324. Kumanyika SK. Obesity in minority populations: An epidemiologic assessment. *Obes Res.* 1994;2:166-182.
325. Ravussin E, Valencia ME, Esparza J, Bennett PH, Schultz LO. Effects of a traditional lifestyle on obesity in Pima Indians. *Diabetes Care.* 1994;17(9):1067-1074.
326. Kawate R, Yamakido M, Nishimoto Y. Migrant studies among the Japanese in Hiroshima and Hawaii. *Proc 10th Congr Int Diabetes Federation.* 1980:526-531.
327. Kawate R, Yamakido M, Nishimoto Y. Diabetes mellitus and its vascular complications in Japanese migrants on the Island of Hawaii. *Diabetes Care.* 1979;29:161-170.
328. Taylor R, Bennett P, Uili R, Joffres N, Levey S, Zimmet P. Diabetes in Wallis Polynesians: a comparison of residents of Wallis Island and first generation migrants of Noumea. *Diabetes Res Clin Pract.* 1985;1:169-178.
329. Bouchard C, Perusse L, Leblanc C, Tremblay A, Theriault G. Inheritance of the amount and distribution of human body fat. *Int J Obes.* 1988;12:205-215.
330. Tambs K, Moum T, Eaves L, et al. Genetics and environmental contributions to the variance of the body mass index in a Norwegian sample of first-degree and second-degree relatives. *Am J Human Biology.* 1991;3:257-268.
331. Vogler GP, Sorensen TI, Stunkard AJ, Srinivasan MR, Rao DC. Influences of genes and shared family environment on adult body mass index assessed in an adoption study by a comprehensive path model. *Int J Obes Relat Metab Disord.* 1995;19:40-45.
332. Stunkard AJ, Harris JR, Pedersen NL, McClearn GE. The body-mass index of twins who have been reared apart. *N Engl J Med.* 1990;322:1483-1487.

333. Allison DB, Kaprio J, Korkeila M, Koskenvuo M, Neale MC, Hayakawa K. The heritability of body mass index among an international sample of monozygotic twins reared apart. *Int J Obes Relat Metab Disord.* 1996;20:501-506.
334. Maes HHM, Neale MC, Eaves LJ. Genetic and environmental factors in relative body weight and human adiposity. *Behav Gene.* 1997;27:325-351.
335. Allison DB. Methodological issues in obesity research: examples from new directions in assessment and management. In: VanItallie TB, Simopoulos AP, eds. *Obesity.* Philadelphia: Charles Press; 1995:129-132.
336. Allison DB, Pi-Sunyer FX. *Obesity Treatment: Establishing Goals, Improving Outcomes, and Reviewing the Research Agenda.* New York: Plenum Press; 1995.
337. Allison DB, Faith MS, Nathan JS. Risch's lambda values for human obesity. *Int J Obes.* 1996;20:990-999.
338. Lee JH, Reed DR, Price RA. Familial risk ratios for extreme obesity: implications for mapping human obesity genes. *Int J Obes Relat Metab Disord.* 1997;21:935-940.
339. Perusse L, Chagnon YC, Dionne FT, Bouchard C. The human obesity gene map: the 1996 update. *Obes Res.* 1997;5:49-61.
340. Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue. *Nature.* 1994;372:425-432.
341. Tartaglia LA, Dembski M, Weng X, et al. Identification and expression cloning of leptin receptor, OB-R. *Cell.* 1995;83:1263-1271.
342. Montague CT, Farooqi IS, Whitehead JP, et al. Congenital leptin deficiency is associated with severe early-onset obesity in humans. *Nature.* 1997;387:903-908.
343. Bouchard C, Perusse L, Rice T, Rao DC. The genetics of human obesity. In: Bray GA, Bouchard C, James WPT, eds. *Handbook of Obesity.* New York: M. Dekker; 1998: Chapter 10.
344. Bouchard C, Tremblay A. Genetic effects in human energy expenditure components. *Int J Obes.* 1990;14 (Suppl 1):49-55.
345. Bouchard C, Tremblay A, Despres JP, et al. The response to long-term overfeeding in identical twins. *N Engl J Med.* 1990;322:1477-1482.
346. Anderssen S, Holme I, Urdal P, Hjerermann I. Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press.* 1995;4:343-349.
347. Croft PR, Brigg D, Smith S, Harrison CB, Branthwaite A, Collins MF. How useful is weight reduction in the management of hypertension? *J R Coll Gen Pract.* 1986;36:445- 448.
348. Davis BR, Blafox MD, Oberman A, et al. Reduction in long-term antihypertensive medication requirements. Effects of weight reduction by dietary intervention in overweight persons with mild hypertension. *Arch Intern Med.* 1993;153:1773-1782.
349. Fagerberg B, Andersson OK, Isaksson B, Bjorntorp P. Blood pressure control during weight reduction in obese hypertensive men: separate effects of sodium and energy restriction. *BMJ.* 1984;288:11-14.
350. Grimm RH Jr, Cohen JD, Smith WM, Falvo-Gerard L, Neaton JD. Hypertension

- management in the Multiple Risk Factor Intervention Trial (MRFIT). 6-year intervention results for men in special intervention and usual care groups. *Arch Intern Med.* 1985;145:1191-1199.
351. Haynes RB, Harper AC, Costley SR, et al. Failure of weight reduction to reduce mildly elevated blood pressure: a randomized trial. *J Hypertens.* 1984;2:535-539.
 352. Heyden S. The workingman's diet. II. Effect of weight reduction in obese patients with hypertension, diabetes, hyperuricemia and hyperlipidemia. *Nutr Metab.* 1978;22:141-159.
 353. Kumanyika SK, Bahnson J, Bottom J, et al. *Can J Cardiol.* 1997;13 (Suppl B):369B [abstract].
 354. Langford HG, Blaufox MD, Oberman A, et al. Dietary therapy slows the return of hypertension after stopping prolonged medication. *JAMA.* 1985;253:657-664.
 355. Langford HG, Davis BR, Blaufox D, et al. Effect of drug and diet treatment of mild hypertension on diastolic blood pressure. The TAIM Research Group. *Hypertension.* 1991;17:210-217.
 356. MacMahon SW, Macdonald GJ, Bernstein L, Andrews G, Blacket RB. A randomized controlled trial of weight reduction and metoprolol in the treatment of hypertension in young overweight patients. *Clin Exp Pharmacol Physiol.* 1985;12:267-271.
 357. Ramsay LE, Ramsay MH, Hettiarachchi J, Davies DL, Winchester J. Weight reduction in a blood pressure clinic. *BMJ.* 1978;2:244-245.
 358. Reisin E, Abel R, Modan M, Silverberg DS, Eliahou HE, Modan B. Effect of weight loss without salt restriction on the reduction of blood pressure in overweight hypertensive patients. *N Engl J Med.* 1978;298:1-6.
 359. Stamler R, Stamler J, Grimm R, et al. Nutritional therapy for high blood pressure. Final report of a 4-year randomized controlled trial—the Hypertension Control Program. *JAMA.* 1987;257:1484-1491.
 360. Wassertheil-Smoller S, Langford HG, Blaufox MD, et al. Effective dietary intervention in hypertensives: sodium restriction and weight reduction. *J Am Diet Assoc.* 1985;85:423-430.
 361. Whelton PK, Applegate WB, Ettiger WH, et al. Efficacy of weight loss and reduced sodium intake in the Trial of Nonpharmacologic Interventions in the Elderly (TONE). *Circulation.* 1996;94 (Suppl I):I-178 [abstract].
 362. Agurs-Collins TD, Kumanyika SK, Have TR, Adams-Campbell LL. A randomized controlled trial of weight reduction and exercise for diabetes management in older African-American subjects. *Diabetes Care.* 1997;20:1503-1511.
 363. Fortmann SP, Haskell WL, Wood PD. Effects of weight loss on clinic and ambulatory blood pressure in normotensive men. *Am J Cardiol.* 1988;62:89-93.
 364. Hakala P, Karvetti RL. Weight reduction on lactovegetarian and mixed diets. Changes in weight, nutrient intake, skin-fold thicknesses and blood pressure. *Eur J Clin Nutr.* 1989;43:421-430.
 365. Hellenius ML, de Faire U, Berglund B, Hamsten A, Krakau I. Diet and exercise are equally effective in reducing risk for cardiovascular disease. Results of a randomized controlled study in men with slightly to moderately raised cardiovascular risk factors. *Atherosclerosis.* 1993;103:81-91.

366. The Hypertension Prevention Trial: three-year effects of dietary changes on blood pressure. Hypertension Prevention Trial Research Group. *Arch Intern Med.* 1990;150:153-162.
367. Jeffery RW, Wing RR, Thorson C, et al. Strengthening behavioral interventions for weight loss: a randomized trial of food provision and monetary incentives. *J Consult Clin Psychol.* 1993;61:1038-1045.
368. Karvetti RL, Hakala P. A 7-year follow-up of a weight reduction programme in Finnish primary health care. *Eur J Clin Nutr.* 1992;46:743-752.
369. Katzell LI, Bleecker ER, Colman EG, Rogus EM, Sorkin JD, Goldberg AP. Effects of weight loss vs aerobic exercise training on risk factors for coronary disease in healthy, obese, middle-aged and older men. A randomized controlled trial. *JAMA.* 1995;274:1915- 1921.
370. Puddey IB, Parker M, Beilin LJ, Vandongen R, Masarei JR. Effects of alcohol and caloric restrictions on blood pressure and serum lipids in overweight men. *Hypertension.* 1992;20:533-541.
371. Reid CM, Dart AM, Dewar EM, Jennings GL. Interactions between the effects of exercise and weight loss on risk factors, cardiovascular haemodynamics and left ventricular structure in overweight subjects. *J Hypertens.* 1994;12:291-301.
372. Rocchini AP, Katch V, Schork A, Kelch RP. Insulin and blood pressure during weight loss in obese adolescents. *Hypertension.* 1987;10:267-273.
373. Simkin-Silverman L, Wing RR, Hansen DH, et al. Prevention of cardiovascular risk factor elevations in healthy premenopausal women. *Prev Med.* 1995;24:509-517.
374. Stamler R, Stamler J, Gosch FC, et al. Primary prevention of hypertension by nutritional-hygienic means. Final report of a randomized, controlled trial. *JAMA.* 1989;262:1801- 1807.
375. Stefanick ML, Mackey S, Sheehan M, Ellsworth N, Haskell WL, Wood PD. Effects of the NCEP Step 2 diet and exercise on lipoprotein in postmenopausal women and men with low HDL-cholesterol and high LDL-cholesterol. *New Engl J Med* (in press).
376. Stevens VJ, Corrigan SA, Obarzanek E, et al. Weight loss intervention in phase 1 of the Trials of Hypertension Prevention. The TOHP Collaborative Research Group. *Arch Intern Med.* 1993;153:849-858.
377. Svendsen OL, Hassager C, Christiansen C. Effect of an energy-restrictive diet, with or without exercise, on lean tissue mass, resting metabolic rate, cardiovascular risk factors, and bone in overweight postmenopausal women. *Am J Med.* 1993;95:131-140.
378. The effects of nonpharmacologic interventions on blood pressure of persons with high normal levels. Results of the Trials of Hypertension Prevention, Phase I. *JAMA.* 1992;267:1213-1220.
379. Effects of weight loss and sodium reduction intervention on blood pressure and hypertension incidence in overweight people with high-normal blood pressure. The Trials of Hypertension Prevention, phase II. The Trials of Hypertension Prevention Collaborative Research Group. *Arch Intern Med.* 1997;157:657-667.
380. Wood PD, Stefanick ML, Williams PT, Haskell WL. The effects on plasma lipoproteins of a prudent weight-reducing diet, with or without exercise, in over-

- weight men and women. *N Engl J Med*. 1991;325:461-466.
381. MacMahon S, Cutler J, Brittain E, Higgins M. Obesity and hypertension: epidemiological and clinical issues. *Eur Heart J*. 1987;8 (Suppl B):57-70.
382. Whelton PK, Appel LJ, Espeland MA, et al. Sodium reduction and weight loss in the treatment of hypertension in older persons: A randomized controlled trial of non-pharmacologic interventions in the elderly (TONE). *JAMA*. 1998;279:839-846.
383. Cutler JA. Randomized clinical trials of weight reduction in nonhypertensive persons. *Ann Epidemiol*. 1991;1:363-370.
384. Svendsen OL, Hassager C, Christiansen C. Six months' follow-up on exercise added to a short-term diet in overweight postmenopausal women—effects on body composition, resting metabolic rate, cardiovascular risk factors and bone. *Int J Obes Relat Metab Disord*. 1994;18:692-698.
385. Steven VJ. Weight loss and blood pressure changes over three years: results of the Trials of Hypertension Prevention (TOHP) Phase II. In: *Cardiovascular Health: Coming Together for the 21st Century*. A National Conference. 1998 Feb 19-21; San Francisco, CA: National Heart, Lung, and Blood Institute; 1998. Abstract R.6F.
386. Bremer JM, Scott RS, Lintott CJ. Dexfenfluramine reduces cardiovascular risk factors. *Int J Obes Relat Metab Disord*. 1994;18:199-205.
387. Ditschuneit HH, Flechtner-Mors M, Adler G. The effects of dexfenfluramine on weight loss and cardiovascular risk factors in female patients with upper and lower body obesity. *J Cardiovasc Risk*. 1996;3:397-403.
388. Bone HG, Reedy KR, Sherwin RS. FDA Orlistat Transcript. Endocrinologic and Metabolic Drugs Advisory Committee Meeting # 67. 1997.
389. Bone HG, Reedy KR, Colley CA. FDA Sibutramine Transcript. Endocrinologic and Metabolic Drugs Advisory Committee Meeting # 64. 1996.
390. Mathus-Vliegen EM, van de Voorde K, Kok AM, Res AM. Dexfenfluramine in the treatment of severe obesity: a placebo-controlled investigation of the effects on weight loss, cardiovascular risk factors, food intake and eating behaviour. *J Intern Med*. 1992;232:119-127.
391. Mathus-Vliegen EM. Prolonged surveillance of dexfenfluramine in severe obesity. *Neth J Med*. 1993;43:246-253.
392. O'Connor HT, Richman RM, Steinbeck KS, Caterson ID. Dexfenfluramine treatment of obesity: a double blind trial with posttrial followup. *Int J Obes Relat Metab Disord*. 1995;19:181-189.
393. Pfohl M, Luft D, Blomberg I, Schmulling RM. Long-term changes of body weight and cardiovascular risk factors after weight reduction with group therapy and dexfenfluramine. *Int J Obes Relat Metab Disord*. 1994;18:391-395.
394. Swinburn BA, Carmichael HE, Wilson MR. Dexfenfluramine as an adjunct to a reduced-fat, ad libitum diet: effects on body composition, nutrient intake and cardiovascular risk factors. *Int J Obes Relat Metab Disord*. 1996;20:1033-1040.
395. Weintraub M, Sundaresan PR, Madan M, et al. Long-term weight control study. I (weeks 0 to 34). The enhancement of behavior modification, caloric restriction, and exercise by fenfluramine plus phentermine versus placebo. *Clin Pharmacol Ther*.

- 1992;51:586-594.
396. Arroll B, Beaglehole R. Exercise for hypertension. *Lancet*. 1993;341:1248-1249.
397. Fagard RH. Physical fitness and blood pressure. *J Hyper*. 1993;11:S47-S52.
398. Kelley G, Tran ZV. Aerobic exercise and normotensive adults: a meta-analysis. *Med Sci Sports Exerc*. 1995;27:1371-1377.
399. Dengel JL, Katzel LI, Goldberg AP. Effect of an American Heart Association diet, with or without weight loss, on lipids in obese middle-aged and older men. *Am J Clin Nutr*. 1995;62:715-721.
400. Jalkanen L. The effect of a weight reduction program on cardiovascular risk factors among overweight hypertensives in primary health care. *Scand J Soc Med*. 1991;19:66-71.
401. King AC, Haskell WL, Taylor CB, Kraemer HC, DeBusk RF. Group- vs home-based exercise training in healthy older men and women. A community-based clinical trial. *JAMA*. 1991;266:1535-1542.
402. Marniemi J, Seppanen A, Hakala P. Long-term effects on lipid metabolism of weight reduction on lactovegetarian and mixed diet. *Int J Obes*. 1990;14:113-125.
403. Nilsson PM, Lindholm LH, Schersten BF. Lifestyle changes improve insulin resistance in hyperinsulinaemic subjects: a 1-year intervention study of hypertensives and normotensives in Dalby. *J Hypertens*. 1992;10:1071-1078.
404. Ronnema T, Marniemi J, Puukka P, Kuusi T. Effects of long-term physical exercise on serum lipids, lipoproteins and lipid metabolizing enzymes in type 2 (non-insulin-dependent) diabetic patients. *Diabetes Res*. 1988;7:79-84.
405. Schuler G, Hambrecht R, Schlierf G, et al. Regular physical exercise and low-fat diet. Effects on progression of coronary artery disease. *Circulation*. 1992;86:1-11.
406. Wood PD, Stefanick ML, Dreon DM, et al. Changes in plasma lipids and lipoproteins in overweight men during weight loss through dieting as compared with exercise. *N Engl J Med*. 1988;319:1173-1179.
407. Drent ML, van der Veen EA. Lipase inhibition: a novel concept in the treatment of obesity. *Int J Obes Relat Metab Disord*. 1993;17:241-244.
408. O'Kane M, Wiles PG, Wales JK. Fluoxetine in the treatment of obese type 2 diabetic patients. *Diabet Med*. 1994;11:105-110.
409. Emery EM, Schmid TL, Kahn HS, Filozof PP. A review of the association between abdominal fat distribution, health outcome measures, and modifiable risk factors. *Am J Health Promot*. 1993;7(5):342-353.
410. Bjorntorp P. Abdominal fat distribution and disease: an overview of epidemiological data. *Ann Med*. 1992;24:15-18.
411. Leon AS. Physical Activity and Cardiovascular Health: A National Consensus. IL: Human Kinetics; 1997.
412. Hjermann I, Leren P, Norman N, Helgeland A, Holme I. Serum insulin response to oral glucose load during a dietary intervention trial in healthy coronary high risk men: the Oslo study. *Scand J Clin Lab Invest*. 1980;40:89-94.
413. Heller SR, Clarke P, Daly H, et al. Group education for obese patients with type 2 diabetes: greater success at less cost. *Diabet Med*. 1988;5:552-556.
414. Manning RM, Jung RT, Leese GP, Newton RW. The comparison of four weight reduc-

- tion strategies aimed at overweight diabetic patients. *Diabet Med.* 1995;12:409-415.
415. Colman E, Katzel LI, Rogus E, Coon P, Muller D, Goldberg AP. Weight loss reduces abdominal fat and improves insulin action in middle-aged and older men with impaired glucose tolerance. *Metabolism.* 1995;44:1502-1508.
 416. Fujioka S, Matsuzawa Y, Tokunaga K, et al. Improvement of glucose and lipid metabolism associated with selective reduction of intra-abdominal visceral fat in premenopausal women with visceral fat obesity. *Int J Obes.* 1991;15:853-859.
 417. Marks SJ, Moore NR, Clark ML, Strauss BJ, Hockaday TD. Reduction of visceral adipose tissue and improvement of metabolic indices: effect of dexfenfluramine in NIDDM. *Obes Res.* 1996;4:1-7.
 418. van der Merwe MT, Wing JR, Celgou LH, et al. Metabolic indices in relation to body composition changes during weight loss on Dexfenfluramine in obese women from two South African ethnic groups. *Int J Obes Relat Metab Disord.* 1996;20:768-776.
 419. Ronnema T, Mattila K, Lehtonen A, Kallio V. A controlled randomized study on the effect of long-term physical exercise on the metabolic control in type 2 diabetic patients. *Acta Med Scand.* 1986;220:219-224.
 420. Han TS, van Leer EM, Seidell JC, Lean ME. Waist circumference action levels in the identification of cardiovascular risk factors: prevalence study in a random sample. *BMJ.* 1995;311:1401-1405.
 421. Pouliot MC, Despres JP, Lemieux S, et al. Waist circumference and abdominal sagittal diameter: best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. *Am J Cardiol.* 1994;73:460-468.
 422. van der Kooy K, Leenen R, Seidell JC, Deurenberg P, Droop A, Bakker CJ. Waist-hip ratio is a poor predictor of changes in visceral fat. *Am J Clin Nutr.* 1993;57:327-333.
 423. Bouchard C, Tremblay A, Despres JP, et al. The response to exercise with constant energy intake in identical twins. *Obes Res.* 1994;2:400-410.
 424. Leenen R, van der Kooy K, Deurenberg P, et al. Visceral fat accumulation in obese subjects: relation to energy expenditure and response to weight loss. *Am J Physiol.* 1992;263:E913-E919.
 425. Stallone DD, Stunkard AJ, Wadden TA, Foster GD, Boorstein J, Arger P. Weight loss and body fat distribution: a feasibility study using computed tomography. *Int J Obes.* 1991;15:775-780.
 426. Ross R, Rissanen J, Pedwell H, Clifford J, Shragge P. Influence of diet and exercise on skeletal muscle and visceral adipose tissue in men. *J Appl Physiol.* 1996;81:2445-2455.
 427. Ross R, Rissanen J. Mobilization of visceral and subcutaneous adipose tissue in response to energy restriction and exercise. *Am J Clin Nutr.* 1994;60:695-703.
 428. van der Kooy K, Leenen R, Seidell JC, Deurenberg P, Hautvast JG. Effect of a weight cycle on visceral fat accumulation. *Am J Clin Nutr.* 1993;58:853-857.
 429. Dowling HJ, Pi-Sunyer FX. Race-dependent health risks of upper body obesity. *Diabetes.* 1993;42:537-543.
 430. Kissebah AH, Vydellingum N, Murray R, et al. Relation of body fat distribution to

- metabolic complications of obesity. *J Clin Endocrinol Metab.* 1982;54:254-260.
431. Berglund A, Andersson OK, Berglund G, Fagerberg B. Antihypertensive effect of diet compared with drug treatment in obese men with mild hypertension. *BMJ.* 1989;299:480-485.
432. Frey-Hewitt B, Vranizan KM, Dreon DM, Wood PD. The effect of weight loss by dieting or exercise on resting metabolic rate in overweight men. *Int J Obes.* 1990;14:327-334.
433. Oberman A, Wassertheil-Smoller S, Langford HG, et al. Pharmacologic and nutritional treatment of mild hypertension: changes in cardiovascular risk status. *Ann Intern Med.* 1990;112:89-95.
434. Hammer RL, Barrier CA, Roundy ES, Bradford JM, Fisher AG. Calorie-restricted low-fat diet and exercise in obese women. *Am J Clin Nutr.* 1989;49:77-85.
435. Marks BL, Ward A, Morris DH, Castellani J, Rippe JM. Fat-free mass is maintained in women following a moderate diet and exercise program. *Med Sci Sports Exerc.* 1995;27:1243-1251.
436. Wadden TA, Sternberg JA, Letizia KA, Stunkard AJ, Foster GD. Treatment of obesity by very low calorie diet, behavior therapy, and their combination: a five-year perspective. *Int J Obes.* 1989;13 (Suppl 2):39-46.
437. Wadden TA, Foster GD, Letizia KA. One-year behavioral treatment of obesity: comparison of moderate and severe caloric restriction and the effects of weight maintenance therapy. *J Consult Clin Psychol.* 1994;62:165-171.
438. Wadden TA, Foster GD, Letizia KA, Mullen JL. Long-term effects of dieting on resting metabolic rate in obese outpatients. *JAMA.* 1990;264:707-711.
439. Wing RR, Blair EH, Bononi P, Marcus MD, Watanabe R, Bergman RN. Caloric restriction per se is a significant factor in improvements in glycemic control and insulin sensitivity during weight loss in obese NIDDM patients. *Diabetes Care.* 1994;17:30-36.
440. de Bont AJ, Baker IA, St. Leger AS, et al. A randomised controlled trial of the effect of low-fat diet advice on dietary response in insulin-independent diabetic women. *Diabetologia.* 1981;21:529-533.
441. Jeffery RW, Hellerstedt WL, French SA, Baxter JE. A randomized trial of counseling for fat restriction versus calorie restriction in the treatment of obesity. *Int J Obes Relat Metab Disord.* 1995;19:132-137.
442. Schlundt DG, Hill JO, Pope-Cordle J, Arnold D, Virts KL, Katahn M. Randomized evaluation of a low fat ad libitum carbohydrate diet for weight reduction. *Int J Obes Relat Metab Disord.* 1993;17:623-629.
443. Shah M, McGovern P, French S, Baxter J. Comparison of a low-fat, ad libitum complex-carbohydrate diet with a low-energy diet in moderately obese women. *Am J Clin Nutr.* 1994;59:980-984.
444. Sheppard L, Kristal AR, Kushi LH. Weight loss in women participating in a randomized trial of low-fat diets. *Am J Clin Nutr.* 1991;54:821-828.
445. Bertram SR, Venter I, Stewart RI. Weight loss in obese women—exercise vs. dietary education. *S Afr Med J.* 1990;78:15-18.
446. Verity LS, Ismail AH. Effects of exercise on cardiovascular disease risk in women with NIDDM. *Diabetes Res Clin Pract.*

- 1989;6:27-35.
447. King AC, Haskell WL, Young DR, Oka RK, Stefanick ML. Long-term effects of varying intensities and formats of physical activity on participation rates, fitness, and lipoproteins in men and women aged 50 to 65 years. *Circulation*. 1995;91:2596-2604.
 448. Gordon NF, Scott CB, Levine BD. Comparison of single versus multiple lifestyle interventions: are the antihypertensive effects of exercise training and diet-induced weight loss additive? *Am J Cardiol*. 1997;79:763-767.
 449. Garrow JS, Summerbell CD. Meta-analysis: effect of exercise, with or without dieting, on the body composition of overweight subjects. *Eur J Clin Nutr*. 1995;49:1-10.
 450. Miller WC, Lindeman AK, Wallace J, Niederpruem M. Diet composition, energy intake, and exercise in relation to body fat in men and women. *Am J Clin Nutr*. 1990;52:426-430.
 451. Fitzgerald SJ, Kriska AM, Pereira MA, de Courten MP. Associations among physical activity, television watching, and obesity in adult Pima Indians. *Med Sci Sports Exerc*. 1997;29:910-915.
 452. Klesges RC, Eck LH, Isbell TR, Fulliton W, Hanson CL. Physical activity, body composition, and blood pressure: a multi-method approach. *Med Sci Sports Exerc*. 1991;23:759-765.
 453. Tremblay A, Despres JP, Leblanc C, et al. Effect of intensity of physical activity on body fatness and fat distribution. *Am J Clin Nutr*. 1990;51:153-157.
 454. French SA, Jeffery RW, Forster JL, McGovern PG, Kelder SH, Baxter JE. Predictors of weight change over two years among a population of working adults: the Healthy Worker Project. *Int J Obes Relat Metab Disord*. 1994;18:145-154.
 455. Williams PT. Evidence for the incompatibility of age-neutral overweight and age-neutral physical activity standards from runners. *Am J Clin Nutr*. 1997;65:1391-1396.
 456. Ewbank PP, Darga LL, Lucas CP. Physical activity as a predictor of weight maintenance in previously obese subjects. *Obes Res*. 1995;3:257-263.
 457. Klesges RC, Klesges LM, Haddock CK, Eck LH. A longitudinal analysis of the impact of dietary intake and physical activity on weight change in adults. *Am J Clin Nutr*. 1992;55:818-822.
 458. Ching PL, Willett WC, Rimm EB, Colditz GA, Gortmaker SL, Stampfer MJ. Activity level and risk of overweight in male health professionals. *Am J Public Health*. 1996;86:25-30.
 459. Williamson DF, Madans J, Anda RF, Kleinman JC, Kahn HS, Byers T. Recreational physical activity and ten-year weight change in a US national cohort. *Int J Obes Relat Metab Disord*. 1993;17:279-286.
 460. Kawachi I, Troisi RJ, Rotnitzky AG, Coakley EH, Colditz GA. Can physical activity minimize weight gain in women after smoking cessation? *Am J Public Health*. 1996;86:999-1004.
 461. Bild DE, Sholinsky P, Smith DE, Lewis CE, Hardin JM, Burke GL. Correlates and predictors of weight loss in young adults: the CARDIA study. *Int J Obes Relat Metab Disord*. 1996;20:47-55.
 462. Kayman S, Bruvold W, Stern JS. Maintenance and relapse after weight loss in women: behavioral aspects. *Am J Clin*

- Nutr. 1990;52:800-807.
463. Grodstein F, Levine R, Troy L, Spencer T, Colditz GA, Stampfer MJ. Three-year follow-up of participants in a commercial weight loss program. Can you keep it off? *Arch Intern Med.* 1996;156:1302-1306.
 464. Seidell JC. Environmental influences on regional fat distribution. *Int J Obes.* 1991;15:31-35.
 465. Kaye SA, Folsom AR, Prineas RJ, Potter JD, Gapstur SM. The association of body fat distribution with lifestyle and reproductive factors in a population study of postmenopausal women. *Int J Obes.* 1990;14:583-591.
 466. Slattery ML, McDonald A, Bild DE, et al. Associations of body fat and its distribution with dietary intake, physical activity, alcohol, and smoking in blacks and whites. *Am J Clin Nutr.* 1992;55:943-949.
 467. Troisi RJ, Heinold JW, Vokonas PS, Weiss ST. Cigarette smoking, dietary intake, and physical activity: effects on body fat distribution—the Normative Aging Study. *Am J Clin Nutr.* 1991;53:1104-1111.
 468. Wing RR, Matthews KA, Kuller LH, Meilahn EN, Plantinga P. Waist to hip ratio in middle-aged women. Associations with behavioral and psychological factors and with changes in cardiovascular risk factors. *Arterioscler Throm.* 1991;11:1250-1257.
 469. Andersen RE, Wadden TA, Bartlett SJ, Vogt RA, Weinstock RS. Relation of weight loss to changes in serum lipids and lipoproteins in obese women. *Am J Clin Nutr.* 1995;62:350-357.
 470. Blonk MC, Jacobs MA, Biesheuvel EH, Weeda-Mannak WL, Heine RJ. Influences on weight loss in type 2 diabetic patients: little long-term benefit from group behaviour therapy and exercise training. *Diabet Med.* 1994;11:449-457.
 471. Neumark-Sztainer D, Kaufmann NA, Berry EM. Physical activity within a community-based weight control program: program evaluation and predictors of success. *Public Health Rev.* 1995;23:237-251.
 472. Sweeney ME, Hill JO, Heller PA, Baney R, DiGirolamo M. Severe vs moderate energy restriction with and without exercise in the treatment of obesity: efficiency of weight loss. *Am J Clin Nutr.* 1993;57:127-134.
 473. Wing RR, Epstein LH, Paternostro-Bayles M, Kriska A, Nowalk MP, Gooding W. Exercise in a behavioural weight control programme for obese patients with Type 2 (non-insulin-dependent) diabetes. *Diabetologia.* 1988;31:902-909.
 474. Kaplan RM, Hartwell SL, Wilson DK, Wallace JP. Effects of diet and exercise interventions on control and quality of life in non-insulin-dependent diabetes mellitus. *J Gen Intern Med.* 1987;2:220-228.
 475. Leighton RF, Repka FJ, Birk TJ, et al. The Toledo Exercise and Diet Study. Results at 26 weeks. *Arch Intern Med.* 1990;150:1016-1020.
 476. Long CG, Simpson CM, Allott EA. Psychological and dietetic counselling combined in the treatment of obesity: a comparative study in a hospital outpatient clinic. *Hum Nutr Appl Nutr.* 1983;37:94-102.
 477. Wadden TA, Stunkard AJ. Controlled trial of very low calorie diet, behavior therapy, and their combination in the treatment of obesity. *J Consult Clin Psychol.* 1986;54:482-488.
 478. Craighead LW, Stunkard AJ, O'Brien RM. Behavior therapy and pharmacotherapy for obesity. *Arch Gen Psychiatry.* 1981;38:763-

- 768.
479. Bennett GA. An evaluation of self-instructional training in the treatment of obesity. *Addict Behav.* 1986;11:125-134.
480. Bennett GA. Cognitive rehearsal in the treatment of obesity: a comparison against cue avoidance and social pressure. *Addict Behav.* 1986;11:225-237.
481. Forster JL, Jeffery RW, Sullivan S, Snell MK. A work-site weight control program using financial incentives collected through payroll deduction. *J Occup Med.* 1985;27:804-808.
482. Glasgow RE, Toobert DJ, Hampson SE, Brown JE, Lewinsohn PM, Donnelly J. Improving self-care among older patients with type II diabetes: the Sixty Something... Study. *Patient Educ Couns.* 1992;19:61-74.
483. Jeffery RW, Hellerstedt WL, Schmid TL. Correspondence programs for smoking cessation and weight control: a comparison of two strategies in the Minnesota Heart Health Program. *Health Psychol.* 1990;9:585-598.
484. Lovibond SH, Birrell PC, Langeluddecke P. Changing coronary heart disease risk-factor status: the effects of three behavioral programs. *J Behav Med.* 1986;9:415-437.
485. Muchmore DB, Springer J, Miller M. Self-monitoring of blood glucose in overweight type 2 diabetic patients. *Acta Diabetol.* 1994;31:215-219.
486. Perri MG, Nezu AM, Patti ET, McCann KL. Effect of length of treatment on weight loss. *J Consult Clin Psychol.* 1989;57:450-452.
487. Perri MG, McAllister DA, Gange JJ, Jordan RC, McAdoo G, Nezu AM. Effects of four maintenance programs on the long-term management of obesity. *J Consult Clin Psychol.* 1988;56:529-534.
488. Rosenthal B, Allen GJ, Winter C. Husband involvement in the behavioral treatment of overweight women: initial effects and long-term follow-up. *Int J Obes.* 1980;4:165-173.
489. Smith DE, Heckemeyer CM, Kratt PP, Mason DA. Motivational interviewing to improve adherence to a behavioral weight-control program for older obese women with NIDDM. A pilot study. *Diabetes Care.* 1997;20:52-54.
490. Uusitupa M, Laitinen J, Siitonen O, Vanninen E, Pyorala K. The maintenance of improved metabolic control after intensified diet therapy in recent type 2 diabetes. *Diabetes Res Clin Pract.* 1993;19:227-238.
491. Wadden TA, Berkowitz RI, Vogt RA, Steen SN, Stunkard AJ, Foster GD. Lifestyle modification in the pharmacologic treatment of obesity: a pilot investigation of a potential primary care approach. *Obes Res.* 1997;5:218-226.
492. Wing RR, Marcus MD, Epstein LH, Jawad A. A "family-based" approach to the treatment of obese Type II diabetic patients. *J Consult Clin Psychol.* 1991;59:156-162.
493. Wing RR, Epstein LH, Nowalk MP, Scott N, Koeske R, Hagg S. Does self-monitoring of blood glucose levels improve dietary compliance for obese patients with Type II diabetes? *Am J Med.* 1986;81:830-836.
494. Eldredge KL, Agras WS, Arnow B, et al. The effects of extending cognitive-behavioral therapy for binge eating disorder among initial treatment nonresponders. *Int J Eat Disord.* 1997;21:347-352.
495. Agras WS, Berkowitz RI, Arnow BA, et al.

- Maintenance following a very-low-calorie diet. *J Consult Clin Psychol.* 1996;64:610-613.
496. Sobal J, Devin C. Social aspects of obesity. In: Dalton S, ed. *Overweight and Weight Management: The Health Professional's Guide To Understanding and Practice.* Gaithersburg, MD: Aspen Publishers; 1997:312-331.
497. Brown PJ; Cultural perspectives on the etiology and treatment of obesity. In: Stunkard AJ, Wadden TA, eds. *Obesity Theory and Therapy.* New York: Raven Press; 1993:179-195.
498. Foster GD, Wadden TA, Vogt RA. Resting energy expenditure in obese African American and Caucasian women. *Obes Res.* 1997;5:1-8.
499. Yanovski SZ, Reynolds JC, Boyle AJ, Yanovski JA. Resting metabolic rate in African-American and Caucasian girls. *Obes Res.* 1997;5:321-325.
500. Albu JB, Murphy L, Frager DH, Johnson JA, Pi-Sunyer FX. Visceral fat and race-dependent health risks in obese nondiabetic premenopausal women. *Diabetes.* 1997;46:456-462.
501. Allison DB, Edlen-Nezin L, Clay-Williams G. Obesity among African American women: prevalence, consequences, causes, and developing research. *Womens Health.* 1997;3:243-274.
502. Andersson B, Seidell J, Terning K, Bjorntorp P. Influence of menopause on dietary treatment of obesity. *J Intern Med.* 1990;227:173-181.
503. Kumanyika SK, Obarzanek E, Stevens VJ, Hebert PR, Whelton PK. Weight-loss experience of black and white participants in NHLBI-sponsored clinical trials. *Am J Clin Nutr.* 1991;53:1631S-1638S.
504. Wing RR, Anglin K. Effectiveness of a behavioral weight control program for blacks and whites with NIDDM. *Diabetes Care.* 1996;19:409-413.
505. Wylie-Rosett J, Wassertheil-Smoller S, Blaufox MD, et al. Trial of antihypertensive intervention and management: greater efficacy with weight reduction than with a sodium-potassium intervention. *J Am Diet Assoc.* 1993;93:408-415.
506. Andersen T, Astrup A, Quaade F. Dexfenfluramine as adjuvant to a low-calorie formula diet in the treatment of obesity: a randomized clinical trial. *Int J Obes Relat Metab Disord.* 1992;16:35-40.
507. Guy-Grand B, Apfelbaum M, Crepaldi G, Gries A, Lefebvre P, Turner P. International trial of long-term dexfenfluramine in obesity. *Lancet.* 1989;2:1142-1145.
508. Finer N. Body weight evolution during dexfenfluramine treatment after initial weight control. *Int J Obes Relat Metab Disord.* 1992;16 (Suppl 3):S25-S29.
509. Willims R.A., Foulsham BM; Weight reduction in osteoarthritis using phentermine. *Practitioner.* 1981; 225:231-232.
510. Bray GA, Ryan DH, Gordon D, Heidingsfelder S, Cerise F, Wilson K. A double-blind randomized placebo-controlled trial of sibutramine. *Obes Res.* 1996;4:263-270.
511. Weintraub M, Ginsberg G, Stein EC, et al. Phenylpropanolamine OROS (Acutrim) vs. placebo in combination with caloric restriction and physician-managed behavior modification. *Clin Pharmacol Ther.* 1986;39:501-509.
512. Schteingart DE. Effectiveness of phenylpropanolamine in the management of

- moderate obesity. *Int J Obes Relat Metab Disord.* 1992;16:487-493.
513. Jones SP, Smith IG, Kelly F, Gray JA. Long term weight loss with sibutramine. *Int J Obes.* 1995;19 (Suppl 2):41 [abstract 071].
514. Weintraub M, Sundaresan PR, Schuster B, et al. Long-term weight control study. IV (weeks 156 to 190). The second double-blind phase. *Clin Pharmacol Ther.* 1992;51:608- 614.
515. Andersen T, Backer OG, Stokholm KH, Quaade F. Randomized trial of diet and gastroplasty compared with diet alone in morbid obesity. *N Engl J Med.* 1984;310:352- 356.
516. Andersen T, Backer OG, Astrup A, Quaade F. Horizontal or vertical banded gastroplasty after pretreatment with very-low-calorie formula diet: a randomized trial. *Int J Obes.* 1987;11:295-304.
517. Brolin RE, Kenler HA, Gorman JH, Cody RP. Long-limb gastric bypass in the super-obese. a prospective randomized study. *Ann Surg.* 1992;215:387-395.
518. Naslund I. The size of the gastric outlet and the outcome of surgery for obesity. *Acta Chir Scand.* 1986;152:205-210.
519. Hall JC, Watts JM, O'Brien PE, et al. Gastric surgery for morbid obesity. The Adelaide Study. *Ann Surg.* 1990;211:419-427.
520. Laws HL, Piantadosi S. Superior gastric reduction procedure for morbid obesity: a prospective, randomized trial. *Ann Surg.* 1981;193:334-340.
521. Lechner GW, Callender AK. Subtotal gastric exclusion and gastric partitioning: a randomized prospective comparison of one hundred patients. *Surgery.* 1981;90:637-644.
522. Pories WJ, Flickinger EG, Meelheim D, Van Rij AM, Thomas FT. The effectiveness of gastric bypass over gastric partition in morbid obesity: consequence of distal gastric and duodenal exclusion. *Ann Surg.* 1982;196:389-399.
523. Gastrointestinal surgery for severe obesity. National Institutes of Health Consensus Development Conference Statement. *Am J Clin Nutr.* 1992;55:615S-619S.
524. Busetto L, Perini P, Giantin V, et al. Relationship between energy expenditure and visceral fat accumulation in obese women submitted to adjustable silicone gastric banding (ASGB). *Int J Obes Relat Metab Disord.* 1995;19:227-233.
525. Zumoff B, Strain GW, Miller LK, et al. Plasma free and non-sex-hormone-binding-globulin-bound testosterone are decreased in obese men in proportion to their degree of obesity. *J Clin Endocrinol Metab.* 1990;71:929-931.
526. Gallagher D, Visser M, Sepulveda D, Pierson RN, Harris T, Heymsfield SB. How useful is body mass index for comparison of body fatness across age, sex, and ethnic groups? *Am J Epidemiol.* 1996;143:228-239.
527. Heymsfield SB, Allison DB, Heshka S, Pierson RN Jr. Assessment of human body composition. In: Allison DB, ed. *Handbook of Assessment Methods For Eating Behavior and Weight Related Problems: Measures, Theory, and Research.* Thousand Oaks, CA: Sage Publications; 1995:515-560.
528. Stensland SH, Margolis S. Simplifying the calculation of body mass index for quick reference. *J Am Diet Assoc.* 1990;90:856.
529. Abate N, Garg A, Peshock RM, Stray-Gundersen J, Adams-Huet B, Grundy SM.

- Relationship of generalized and regional adiposity to insulin sensitivity in men with NIDDM. *Diabetes*. 1996;45:1684-1693.
530. Lean ME, Han TS, Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ*. 1995;311:158-161.
531. Sjostrom L, Kvist H, Cederblad A, Tylen U. Determination of total adipose tissue and body fat in women by computed tomography, ⁴⁰K, and tritium. *Am J Physiol*. 1986;250:E736- E745.
532. Jensen MD, Kanaley JA, Reed JE, Sheedy PF. Measurement of abdominal and visceral fat with computed tomography and dual-energy x-ray absorptiometry. *Am J Clin Nutr*. 1995;61:274-278.
533. Despres JP, Moorjani S, Ferland M, et al. Adipose tissue distribution and plasma lipoprotein levels in obese women. Importance of intra-abdominal fat. *Arteriosclerosis*. 1989;9:203-210.
534. Pouliot MC, Despres JP, Nadeau A, et al. Visceral obesity in men. Associations with glucose tolerance, plasma insulin, and lipoprotein levels. *Diabetes*. 1992;41:826-834.
535. Abate N, Garg A, Peshock RM, Stray-Gundersen J, Grundy SM. Relationships of generalized and regional adiposity to insulin sensitivity in men. *J Clin Investigation*. 1995;96:88-98.
536. Goodpaster BH, Thaete FL, Thaete JA, Simoneau, Kelley DE. Subcutaneous abdominal fat and thigh muscle composition predict insulin sensitivity independently of visceral fat. *Diabetes*. 1997;46:1579-1585.
537. Lemieux S, Prud'homme D, Bouchard C, Tremblay A, Despres J. A single threshold value of waist girth identifies normal-weight and overweight subjects with excess visceral adipose tissue. *Am J Clin Nutr*. 1996;64:685-693.
538. National Center for Health Statistics. NHANES III Anthropometric Procedures Video. Washington, DC: U.S. Government Printing Office; 1996. Stock Number 017-022-01335-5.
539. Gordon T, Doyle JT. Weight and mortality in men: the Albany Study. *Int J Epidemiol*. 1988;17:77-81.
540. Hamm P, Shekelle RB, Stamler J. Large fluctuations in body weight during young adulthood and 25-year risk of coronary death in men. *Am J Epidemiol*. 1989;129:312-318.
541. Lindsted K, Tonstad S, Kuzma JW. Body mass index and patterns of mortality among Seventh-day Adventist men. *Int J Obes*. 1991;15:397-406.
542. Norgan NG, Jones PR. The effect of standardising the body mass index for relative sitting height. *Int J Obes Relat Metab Disord*. 1995;19:206-208.
543. Conway JM, Yanovski SZ, Avila NA, Hubbard VS. Visceral adipose tissue differences in black and white women. *Am J Clin Nutr*. 1995;61:765-771.
544. Potts J, Simmons D. Sex and ethnic group differences in fat distribution in young United Kingdom South Asians and Europids. *J Clin Epidemiol*. 1994;47:837-841.
545. Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure. The Sixth Report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure (JNCVI). *Arch Intern Med*.

- 1997;157:2413-2446.
546. American Diabetes Association. ADA Clinical Practice Recommendations 1998 . *Diabetes Care*. 1998;21(Suppl 1):S1-S93.
 547. Fiore MC, Bailey WC, Cohen SJ. Smoking Cessation. U.S. Department of Health and Human Services, PHS, Agency for Health Care Policy and Research. Clinical Practice Guidelines No. 18. 1996.
 548. Paffenbarger RS Jr, Hyde RT, Wing AL, Lee IM, Jung DL, Kampert JB. The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *N Engl J Med*. 1993;328:538-545.
 549. NIH Consensus conference. Triglyceride, high-density lipoprotein, and coronary heart disease. NIH Consensus Development Panel on Triglyceride, High-Density Lipoprotein, and Coronary Heart Disease. *JAMA*. 1993;269:505-510.
 550. Wadden TA, Wingate BJ. Behavioral assessment and treatment of markedly obese patients. In: Wadden TA, VanItallie TB, eds. *Treatment of the Seriously Obese Patient*. New York: Guilford Press; 1992:290-330.
 551. Stunkard AJ. Talking with patients. In: Stunkard AJ, Wadden TA, eds. *Obesity Theory and Therapy*. New York: Raven Press; 1993:355-363.
 552. Wadden TA, Wingate BJ. Compassionate treatment of the obese individual. In: Brownell KD, Fairburn CG, eds. *Eating Disorders and Obesity*. New York: Basic Books; 1995:564- 571.
 553. Foster GD, Wadden TA, Vogt RA, Brewer G. What is a reasonable weight loss? Patients' expectations and evaluations of obesity treatment outcome. *J Consult Clinl Psychol*. 1997;65:79-85.
 554. Williamson DF, Serdula MK, Anada RF, Levy A, Byers T. Weight loss attempts in adults: goals, duration, and rate of weight loss. *Am J Pub Health*. 1992;82:1251-1257.
 555. Wing RR, Blair E, Marcus M, Epstein LHH, Harvey J. Year-long weight loss treatment for obese patients with type II diabetes: does including an intermittent very-low-calorie diet improve outcome? *Am J Med*. 1994;97:354-362.
 556. VanItallie TB, Yang M-U. Current concepts in nutrition: diets and weight loss. *N Engl J Med*. 1977;297:1158-1161.
 557. Yang M, VanItallie TB. Effect of energy restriction on body composition and nitrogen balance in obese individuals. In: Wadden TA, VanItallie TB, eds. *Treatment of the Seriously Obese Patient*. New York: Guilford Press; 1992:83-106.
 558. Very low-calorie diets. National Task Force on the Prevention and Treatment of Obesity, National Institutes of Health. *JAMA*. 1993;270:967-974.
 559. Tremblay A, Buemann B, Theriault G, Bouchard C. Body fatness in active individuals reporting low lipid and alcohol intake. *Eur J Clin Nutr*. 1995;49:824-831.
 560. Gruchow HW, Sobocinski KA, Barboriak JJ, Scheller JG. Alcohol consumption, nutrient intake and relative body weight among US adults. *Am J Clin Nutr*. 1985;42:289-295.
 561. de Castro JM, Orozco S. Moderate alcohol intake and spontaneous eating patterns of humans: evidence of unregulated supplementation. *Am J Clin Nutr*. 1990;52:246-253.
 562. Veenstra J, Schenkel JA, van Erp-Baart

- AM, et al. Alcohol consumption in relation to food intake and smoking habits in the Dutch National Food Consumption Survey. *Eur J Clin Nutr.* 1993;47:482-489.
563. Tremblay A, Wouters E, Wenker M, St-Pierre S, Bouchard C, Despres JP. Alcohol and a high-fat diet: a combination favoring overfeeding. *Am J Clin Nutr.* 1995;62:639-644.
564. Tremblay A, St-Pierre S. The hyperphagic effect of a high-fat diet and alcohol intake persists after control for energy density. *Am J Clin Nutr.* 1996;63:479-482.
565. Poppitt SD, Eckhardt JW, McGonagle J, Murgatroyd PR, Prentice AM. Short-term effects of alcohol consumption on appetite and energy intake. *Physiol Behav.* 1996;60:1063-1070.
566. Foltin RW, Kelly TH, Fischman MW. Ethanol as an energy source in humans: comparison with dextrose-containing beverages. *Appetite.* 1993;20:95-110.
567. Butrum RR, Clifford CK, Lanza E. NCI dietary guidelines: rationale. *Am J Clin Nutr.* 1988;48(3 Suppl):888-895.
568. United States. Public Health Service. Office of the Surgeon General. Nutrition and Health: A Report of the Surgeon General. Washington, DC: Government Printing Office; 1988:192.
569. NIH Consensus conference. Optimal calcium intake. NIH Consensus Development Panel on Optimal Calcium Intake. *JAMA.* 1994;272:1942-1948.
570. Wadden TA, Letizia KA. Predictors of attrition and weight loss in patients treated by moderate and severe caloric restriction. In: Wadden TA, VanItallie TB, eds. *Treatment of the Seriously Obese Patient.* New York: Guilford Press; 1992:383-410.
571. Foreyt JP, Goodrick GK. Factors common to successful therapy for the obese patient. *Med Sci Sports Exerc.* 1991;23:292-297.
572. Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion. Surgeon General's Report on Physical Activity and Health. Atlanta, GA: CDC; 1996.
573. Blair SN, Kampert JB, Kohl HW III, et al. Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. *JAMA.* 1996;276:205-210.
574. Kohl HW, Gordon NF, Villegas JA, Blair SN. Cardiorespiratory fitness, glycemic status, and mortality risk in men. *Diabetes Care.* 1992;15:184-192.
575. Tremblay A, Nadeau A, Despres JP, St-Jean L, Theriault G, Bouchard C. Long-term exercise training with constant energy intake. 2: effect on glucose metabolism and resting energy expenditure. *Int J Obes.* 1990;14:75-84.
576. Wing RR, Marcus MD, Blair EH, Burton LR. Psychological responses of obese Type II diabetic subjects to very low-calorie diet. *Diabetes Care.* 1991;14:596-599.
577. Pate RR, Pratt M, Blair SN, et al. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA.* 1995;273:402-407.
578. NIH Consensus Conference. Physical Activity and Cardiovascular Health. *JAMA.* 1996;276:241-246.
579. Wadden TA, Foster GD. Behavioral assessment and treatment of markedly obese patients. In: Wadden TA, VanItallie TB,

- eds. *Treatment of the Seriously Obese Patient*. New York: Guilford Press; 1992:290-330.
580. Wadden TA. The treatment of obesity. In: Stunkard AJ, Wadden TA, eds. *Obesity: Theory and Therapy*. New York: Raven Press; 1993:197-217.
581. Wilson GT, Fairborn CG. Treatments of eating disorders. In: Nathan PE, Gorman JM, eds. *A Guide To Treatments That Work*. New York: Oxford University Press; 1997:501- 530.
582. Schlundt DG, Hill JO, Pope-Cordle J, Arnold D, Virts KL, Katahn M. Randomized evaluation of a low fat ad libitum carbohydrate diet for weight reduction. *Int J Obes Relat Metab Disord*. 1993;17:623-629.
583. Tremblay A, Almeras N, Boer J, Kranenbarg EK, Despres JP. Diet composition and postexercise energy balance. *Am J Clin Nutr*. 1994;59:975-979.
584. King NA, Blundell JE. High-fat foods overcome the energy expenditure induced by high-intensity cycling or running. *Eur J Clin Nutr*. 1995;49:114-123.
585. Methods for voluntary weight loss and control. NIH Technology Assessment Conference Panel. *Ann Intern Med*. 1992;116:942-949.
586. Atanassoff PG, Weiss BM, Schmid ER, Tornic M. Pulmonary hypertension and dexfenfluramine. *Lancet*. 1992;339:436.
587. Brenot F, Herve P, Petitpretz P, Parent F, Duroux P, Simonneau G. Primary pulmonary hypertension and fenfluramine use. *Br Heart J*. 1993;70:537-541.
588. Voelker R. Obesity drug renews toxicity debate . *JAMA*. 1994;272:1087-1088.
589. Food and Drug Administration. *Draft Guidance Clinical Evaluation of Weight Control Drug*. Rockville, MD: FDA; 1996.
590. Bray GA. Use and abuse of appetite-suppressant drugs in the treatment of obesity. *Ann Intern Med*. 1993;119:707-713.
591. Galloway SM, Farquhar DL, Munro JF. The current status of antiobesity drugs. *Postgrad Med J*. 1984;60 (Suppl 3):19-26.
592. Long-term pharmacotherapy in the management of obesity. National Task Force on the Prevention and Treatment of Obesity. *JAMA*. 1996;276:1907-1915.
593. Weintraub M, Bray GA. Drug treatment of obesity. *Med Clin North Am*. 1989;73:237- 249.
594. MacDonald KG Jr, Long SD, Swanson MS, et al. The gastric bypass operation reduces the progression and mortality of non-insulin-dependent diabetes mellitus. *J Gastrointest Surg*. 1997;1:213-220.
595. Pories WJ, Swanson MS, MacDonald KG Jr, et al. Who would have thought it? An operation proves to be the most effective therapy for adult-onset diabetes mellitus. *Ann Surg*. 1995;222:339-350; discussion 350-352.
596. Harris TB, Savage PJ, Tell GS, Haan M, Kumanyika S, Lynch JC. Carrying the burden of cardiovascular risk in old age: associations of weight and weight change with prevalent cardiovascular disease, risk factors, and health status in the Cardiovascular Health Study. *Am J Clin Nutr*. 1997;66:837-844.
597. Tayback M, Kumanyika S, Chee E. Body weight as a risk factor in the elderly. *Arch Intern Med*. 1990;150:1065-1072.
598. Galanos AN, Pieper CF, Cornoni-Huntley

- JC, Bales CW, Fillenbaum GG. Nutrition and function: is there a relationship between body mass index and the functional capabilities of community-dwelling elderly? *J Am Geriatr Soc.* 1994;42:368-373.
599. Keil JE, Gazes PC, Sutherland SE, Rust PF, Branch LG, Tyroler HA. Predictors of physical disability in elderly blacks and whites of the Charleston Heart Study. *J Clin Epidemiol.* 1989;42:521-529.
600. Launer LJ, Harris T, Rumpel C, Madans J. Body mass index, weight change, and risk of mobility disability in middle-aged and older women. The epidemiologic follow-up study of NHANES I. *JAMA.* 1994;271:1093-1098.
601. Willett WC. Weight loss in the elderly: cause or effect of poor health? *Am J Clin Nutr.* 1997;66:737-738.
602. Galanis DJ, Harris T, Sharp DS, Petrovitch H. Relative weight, weight change, and risk of coronary heart disease in the Honolulu Heart Program. *Am J Epidemiol.* 1998;147:379-386.
603. Chaturvedi N, Fuller JH. Mortality risk by body weight and weight change in people with NIDDM. The WHO Multinational Study of Vascular Disease in Diabetes. *Diabetes Care.* 1995;18:766-774.
604. Iribarren C, Sharp DS, Burchfiel CM, Petrovitch H. Association of weight loss and weight fluctuation with mortality among Japanese-American men. *N Engl J Med.* 1995;333:686-692.
605. Pamuk ER, Williamson DF, Madans J, Serdula MK, Kleinman JC, Byers T. Weight loss and mortality in a national cohort of adults, 1971-1987. *Am J Epidemiol.* 1992;136:686-697.
606. Rumpel C, Harris TB, Madans J. Modification of the relationship between the Quetelet index and mortality by weight-loss history among older women. *Ann Epidemiol.* 1993;3:343-350.
607. Wallace JI, Schwartz RS, LaCroix AZ, Uhlmann RF, Pearlman RA. Involuntary weight loss in older outpatients: incidence and clinical significance. *J Am Geriatr Soc.* 1995;43:329-337.
608. Ensrud KE, Cauley J, Lipschutz R, Cummings SR. Weight change and fractures in older women. Study of Osteoporotic Fractures Research Group. *Arch Intern Med.* 1997;157:857-863.
609. Langlois JA, Harris T, Looker AC, Madans J. Weight change between age 50 years and old age is associated with risk of hip fracture in white women aged 67 years and older. *Arch Intern Med.* 1996;156:989-994.
610. Shapses SA, Heymsfield SB, Ricci TA. Voluntary weight reduction increases bone turnover and loss. In: Burckhardt P, Heaney R, Dawson-Hughes B, eds. *Nutritional Aspects of Osteoporosis.* New York: Springer-Verlag; 1998.
611. Michel BA, Lane NE, Bloch DA, Jones HH, Fries JF. Effect of changes in weight-bearing exercise on lumbar bone mass after age fifty. *Ann Med.* 1991;23:397-401.
612. Marcus R, Drinkwater B, Dalsky G, et al. Osteoporosis and exercise in women. *Med Sci Sports Exerc.* 1992;24(Suppl 6):S301-S307.
613. Gerace TA, Hollis J, Ockene JK, Svendsen K. Smoking cessation and change in diastolic blood pressure, body weight, and plasma lipids. MRFIT Research Group. *Prev Med.* 1991;20:602-620.

614. Klesges RC, Meyers AW, Klesges LM, La Vasque ME. Smoking, body weight, and their effects on smoking behavior: a comprehensive review of the literature. *Psychol Bull.* 1989;106:204-230.
615. Williamson DF, Madans J, Anda RF, Kleinman JC, Giovino GA, Byers T. Smoking cessation and severity of weight gain in a national cohort. *N Engl J Med.* 1991;324:739-745.
616. Pirie PL, Murray DM, Luepker RV. Gender differences in cigarette smoking and quitting in a cohort of young adults. *Am J Public Health.* 1991;81:324-327.
617. Pomerleau CS, Kurth CL. Willingness of female smokers to tolerate postcessation weight gain. *J Subst Abuse.* 1996;8:371-378.
618. Hall SM, Tunstall CD, Vila KL, Duffy J. Weight gain prevention and smoking cessation: cautionary findings. *Am J Public Health.* 1992;82:799-803.
619. Spring B, Pingotore R, Kesler K. Cigarette smoking and body weight: strategies to minimize weight gain after smoking cessation. *Int J Obes.* 1992;16:S19-S23.
620. Perkins KA, Epstein LH, Marks BL, Stiller RL, Jacob RG. The effect of nicotine on energy expenditure during light physical activity. *N Engl J Med.* 1989;320:898-903.
621. Stamford BA, Matter S, Fell RD, Papanek P. Effects of smoking cessation on weight gain, metabolic rate, caloric consumption, and blood lipids. *Am J Clin Nutr.* 1986;43:486-494.
622. Gilbert RM, Pope MA. Early effects of quitting smoking. *Psychopharmacology (Berl).* 1982;78:121-127.
623. Spring B, Wurtman J, Gleason R, Wurtman R, Kessler K. Weight gain and withdrawal symptoms after smoking cessation: a preventive intervention using d-fenfluramine. *Health Psychol.* 1991;10:216-223.
624. Gross J, Stitzer ML, Maldonado J. Nicotine replacement: effects of postcessation weight gain. *J Consult Clin Psychol.* 1989;57:87-92.
625. Klesges RC, Klesges LM, Meyers AW, Klem ML, Isbell T. The effects of phenylpropranolamine on dietary intake, physical activity, and body weight after smoking cessation. *Clin Pharmacol Ther.* 1990;47:747-754.
626. Hurt RD, Sachs DPL, Clover ED, et al. A comparison of sustained-release bupropion and placebo for smoking cessation. *N Engl J Med.* 1997;337:1195-1202.
627. Perri MG, Lauer JB, Yancey DZ, et al. Effects of peer support and therapist contact on long-term weight loss. *J Consult Clin Psych.* 1995;55:615-617.
628. Broussard BA, Johnson A, Himes JH, et al. Prevalence of obesity in American Indians and Alaska Natives. *Am J Clin Nutr.* 1991;53:1535S-1542S.
629. Hodge AM, Dowse GK, Toelue P, Collins VR, Imo T, Zimmet PZ. Dramatic increase in the prevalence of obesity in western Samoa over the 13 year period 1978-1991. *Int J Obes Relat Metab Disord.* 1994;18:419-428.
630. Health Care Financing Administration. Medicare Current Beneficiary Survey: 1991. Percentage of medicare beneficiaries who are overweight, by race, gender, and education. Published Data. 1991.
631. Avila P, Hovell MF. Physical activity training for weight loss in Latinas: a controlled trial. *Int J Obes Relat Metab Disord.*

- 1994;18:476-482.
632. Cousins JH, Rubovits DS, Dunn JK, Reeves RS, Ramirez AG, Foreyt JP. Family versus individually oriented intervention for weight loss in Mexican-American women. *Public Health Rep.* 1992;107:549-555.
 633. Daniel EL. A multi-intervention weight management program for low-income rural women. *J Am Diet Assoc.* 1989;89:1310-1311.
 634. Domel SB, Alford BB, Cattlett HN, Gench BE. Weight control for black women. *J Am Diet Assoc.* 1992;92:346-348.
 635. Domel SB, Alford BB, Cattlett HN, Rodriguez ML, Gench BE. A pilot weight control program for Hispanic women. *J Am Diet Assoc.* 1992;92:1270-1271.
 636. Fox RA, Haniotes H, Rotatori A. A streamlined weight loss program for moderately retarded adults in a sheltered workshop setting. *Appl Res Ment Retard.* 1984;5:69-79.
 637. Heath GW, Wilson RH, Smith J, Leonard BE. Community-based exercise and weight control: diabetes risk reduction and glycemic control in Zuni Indians. *Am J Clin Nutr.* 1991;53:1642S-1646S.
 638. Kanders BS, Ullmann-Joy P, Foreyt JP, et al. The black American lifestyle intervention (BALI): the design of a weight loss program for working-class African-American women. *J Am Diet Assoc.* 1994;94:310-312.
 639. Kumanyika SK, Brancato J, Brewer A, et al. Interventions in the Trials of Nonpharmacologic Intervention in the Elderly. Effective approaches to weight and sodium reduction among older adults. *Circulation.* 1996;94:I-690.
 640. Kumanyika SK, Charleston JB. Lose weight and win: a church-based weight loss program for blood pressure control among black women. *Patient Educ Counseling.* 1992;19:19-32.
 641. Lasco RA, Curry RH, Dickson VJ, Powers J, Menes S, Merritt RK. Participation rates, weight loss, and blood pressure changes among obese women in a nutrition-exercise program. *Public Health Rep.* 1989;104:640-646.
 642. McNabb W, Quinn M, Kerver J, Cook S, Karrison T. The PATHWAYS church-based weight loss program for urban African-American women at risk for diabetes. *Diabetes Care.* 1997;20:1518-1523.
 643. McNabb WL, Quinn MT, Rosing L. Weight loss program for inner-city black women with non-insulin-dependent diabetes mellitus: Pathways. *J Am Diet Assoc.* 1993;93:75-77.
 644. Mount MA, Kendrick OW, Draughon M, Stitt KR, Head D, Mount R. Group participation as a method to achieve weight loss and blood glucose control. *J Nutr Educ.* 1991;23:25-29.
 645. Mulrow C, Bailey S, Sonksen PH, Slavin B. Evaluation of an audiovisual diabetes education program: Negative results of a randomized trial of patients with non-insulin-dependent diabetes mellitus. *J Gen Intern Med.* 1987;2:215-219.
 646. Pleas J. Long-term effects of a lifestyle-change obesity treatment program with minorities. *J Natl Med Assoc.* 1988;80:747-752.
 647. Shintani TT, Hughes CK, Beckham S, O'Connor HK. Obesity and cardiovascular risk intervention through and ad libitum feeding of traditional Hawaiian diet. *Am J Clin Nutr.* 1991;53:1647S-1651S.

648. Sullivan J, Carter JP. A nutrition-physical fitness intervention program for low-income black parents. *J Natl Med Assoc.* 1985;77:39-43.
649. Gleadhill IC, Schwartz AR, Schubert N, Wise RA, Permutt S, Smith PL. Upper airway collapsibility in snorers and in patients with obstructive hypopnea and apnea. *Am Rev Respir Dis.* 1991;143:1300-1303.
650. Gold AR, Schwartz AR, Wise RA, Smith PL. Pulmonary function and respiratory chemosensitivity in moderately obese patients with sleep apnea. *Chest.* 1993;103:1325-1329.
651. Block AJ, Boysen PG, Wynne JW, Hunt LA. Sleep apnea, hypopnea, and oxygen desaturation in normal subjects. A strong male predominance. *N Engl J Med.* 1979;300:513-517.
652. Guilleminault C, Dement WC. *Sleep Apnea Syndromes. Conference on Sleep Apnea Syndromes.* New York: Liss; 1978.
653. Schmidt-Nowara WW. Cardiovascular consequences of sleep apnea. *Prog Clin Biol Res.* 1990;345:377-385.
654. Norton PG, Dunn EV. Snoring as a risk factor for disease: an epidemiological survey. *BMJ (Clin Res Ed).* 1985;291:630-632.
655. Jennum P, Hein HO, Suadicani P, Gyntelberg F. Cardiovascular risk factors in snorers. A cross-sectional study of 3,323 men aged 54 to 74 years: the Copenhagen Male Study. *Chest.* 1992;102:1371-1376.
656. Stradling JR, Crosby JH. Predictors and prevalence of obstructive sleep apnoea and snoring in 1001 middle aged men. *Thorax.* 1991;46:85-90.
657. Guilleminault C, Quera-Salva MA, Partinen M, Jamieson A. Women and the obstructive sleep apnea syndrome. *Chest.* 1988;93:104-109.
658. Connolly HM, Crary JL, McGoon MD, et al. Valvular heart disease associated with Fenfluramine—Phentermine. *N Engl J Med.* 1997;337:581-588.
659. Hedner J. Vascular function in OSA. *Sleep.* 1996;19(10 Suppl):S213-S217.
660. Tofler GH, Brezinski D, Schafer AI, et al. Concurrent morning increase in platelet aggregability and the risk of myocardial infarction and sudden cardiac death. *N Engl J Med.* 1987;316:1514-1518.
661. He J, Kryger MH, Zorick FJ, Conway W, Roth T. Mortality and apnea index in obstructive sleep apnea. Experience in 385 male patients. *Chest.* 1988;94:9-14.
662. Partinen M, Jamieson A, Guilleminault C. Long-term outcome for obstructive sleep apnea syndrome patients. Mortality. *Chest.* 1988;94:1200-1204.
663. Harman EM, Wynne JW, Block AJ. The effect of weight loss on sleep-disordered breathing and oxygen desaturation in morbidly obese men. *Chest.* 1982;82:291-294.
664. Charuzi I, Ovnat A, Peiser J, Saltz H, Weitzman S, Lavie P. The effect of surgical weight reduction on sleep quality in obesity-related sleep apnea syndrome. *Surgery.* 1985;97:535-538.
665. Peiser J, Lavie P, Ovnat A, Charuzi I. Sleep apnea syndrome in the morbidly obese as an indication for weight reduction surgery. *Ann Surg.* 1984;199:112-115.
666. Sugerma HJ, Fairman RP, Baron PL, Kwentus JA. Gastric surgery for respiratory insufficiency of obesity. *Chest.* 1986;90:81-86.
667. Rubinstein I, Colapinto N, Rotstein LE,

- Brown IG, Hoffstein V. Improvement in upper airway function after weight loss in patients with obstructive sleep apnea. *Am Rev Respir Dis.* 1988;138:1192-1195.
668. Smith PL, Gold AR, Meyers DA, Haponik EF, Bleecker ER. Weight loss in mildly to moderately obese patients with obstructive sleep apnea. *Ann Intern Med.* 1985;103:850- 855.
669. Aubert-Tulkens G, Culee C, Rodenstein DO. Cure of sleep apnea syndrome after long-term nasal continuous positive airway pressure therapy and weight loss. *Sleep.* 1989;12:216- 222.
670. Pasquali R, Colella P, Cirignotta F, et al. Treatment of obese patients with obstructive sleep apnea syndrome (OSAS): effect of weight loss and interference of otorhinolaryngoiatric pathology. *Int J Obes.* 1990;14:207-217.
671. Suratt PM, McTier RF, Findley LJ, Pohl SL, Wilhoit SC. Effect of very low-calorie diets with weight loss on obstructive sleep apnea. *Am J Clin Nutr.* 1992;56:182S-184S.
672. Schwartz AR, Gold AR, Schubert N, et al. Effect of weight loss on upper airway collapsibility in obstructive sleep apnea. *Am Rev Respir Dis.* 1991;144:494-498.
673. American Thoracic Society. Indications and standards for use of nasal continuous positive airway pressure (CPAP) in sleep apnea syndromes. *Am J Resp Crit Care Med.* 1994;150:1738-1745.
674. Andersson B, Elam M, Wallin BG, Bjorntorp P, Andersson OK. Effect of energy-restricted diet on sympathetic muscle nerve activity in obese women. *Hypertension.* 1991;18:783- 789.
675. Andersson OK, Fagerberg B, Hedner T. Haemodynamic adjustment to weight reduction—separate effects of energy versus salt restriction. *J Hypertens Suppl.* 1983;1:35-37.
676. Collins RW, Anderson JW. Medication cost savings associated with weight loss for obese non-insulin-dependent diabetic men and women. *Prev Med.* 1995;24:369-374.
677. Darne B, Nivarong M, Tugaye A, et al. Hypocaloric diet and antihypertensive drug treatment. A randomized controlled clinical trial. *Blood Press.* 1993;2:130-135.
678. Gillett PA, Eisenman PA. The effect of intensity controlled aerobic dance exercise on aerobic capacity of middle-aged, overweight women. *Res Nurs Health.* 1987;10:383-390.
679. The treatment of mild hypertension study. A randomized, placebo-controlled trial of a nutritional-hygienic regimen along with various drug monotherapies. *Arch Intern Med.* 1991;151:1413-1423.
680. Jeffery RW, Gillum R, Gerber WM, Jacobs D, Elmer PJ, Prineas RJ. Weight and sodium reduction for the prevention of hypertension: a comparison of group treatment and individual counseling. *Am J Public Health.* 1983;73:691-693.
681. Kanaley JA, Andresen-Reid ML, Oenning L, Kottke BA, Jensen MD. Differential health benefits of weight loss in upper-body and lower-body obese women. *Am J Clin Nutr.* 1993;57:20-26.
682. Kumanyika SK. The impact of obesity on hypertension management in African American. *J Health Care Poor Underserved.* 1997;8:352-364.
683. Mellies MJ, Vitale C, Jandacek RJ, Lamkin GE, Glueck CJ. The substitution of sucrose polyester for dietary fat in obese,

- hypercholesterolemic outpatients. *Am J Clin Nutr.* 1985;41:1-12.
684. Page RC, Harnden KE, Cook JT, Turner RC. Can lifestyles of subjects with impaired glucose tolerance be changed? A feasibility study. *Diabet Med.* 1992;9:562-566.
685. Ready AE, Drinkwater DT, Ducas J, Fitzpatrick DW, Brereton DG, Oades SC. Walking program reduces elevated cholesterol in women postmenopause. *Can J Cardiol.* 1995;11:905-912.
686. Rigaud D, Ryttig KR, Angel LA, Apfelbaum M. Overweight treated with energy restriction and a dietary fibre supplement: a 6-month randomized, double-blind, placebo-controlled trial. *Int J Obes.* 1990;14:763-769.
687. Rissanen A, Pietinen P, Siljamaki-Ojansuu U, Piirainen H, Reissel P. Treatment of hypertension in obese patients: efficacy and feasibility of weight and salt reduction programs. *Acta Med Scand.* 1985;218:149-156.
688. Ryttig KR, Rossner S. Weight maintenance after a very low calorie diet (VLCD) weight reduction period and the effects of VLCD supplementation. A prospective, randomized, comparative, controlled long-term trial. *J Intern Med.* 1995;238:299-306.
689. Singh RB, Niaz MA, Bishnoi I, Singh U, Begum R, Rastogi SS. Effect of low energy diet and weight loss on major risk factors, central obesity and associated disturbances in patients with essential hypertension. *J Hum Hypertens.* 1995;9:355-362.
690. Singh RB, Niaz MA, Ghosh S. Effect on central obesity and associated disturbances of low-energy, fruit- and vegetable-enriched prudent diet in north Indians. *Postgrad Med J.* 1994;70:895-900.
691. Singh RB, Rastogi SS, Sircar AR, Mehta PJ, Sharma KK. Dietary strategies for risk-factor modification to prevent cardiovascular diseases. *Nutrition.* 1991;7:210-214.
692. Stokholm KH, Nielsen PE, Quaade F. Correlation between initial blood pressure and blood pressure decrease after weight loss: a study in patients with jejunoileal bypass versus medical treatment for morbid obesity. *Int J Obes.* 1982;6:307-312.
693. Walker KZ, O'Dea K, Nicholson GC, Muir JG. Dietary composition, body weight, and NIDDM. Comparison of high-fiber, high-carbohydrate, and modified-fat diets. *Diabetes Care.* 1995;18:401-403.
694. Wing RR, Jeffery RW, Burton LR, Thorson C, Kuller LH, Folsom AR. Change in waist-hip ratio with weight loss and its association with change in cardiovascular risk factors. *Am J Clin Nutr.* 1992;55:1086-1092.
695. Astrup A, Breum L, Toubro S, Hein P, Quaade F. The effect and safety of an ephedrine/caffeine compound compared to ephedrine, caffeine and placebo in obese subjects on an energy-restricted diet. A double blind trial. *Int J Obes Relat Metab Disord.* 1992;16:269-277.
696. Connacher AA, Jung RT, Mitchell PE. Weight loss in obese subjects on a restricted diet given BRL 26830A, a new atypical beta adrenoceptor agonist. *BMJ (Clin Res Ed).* 1988;296:1217-1220.
697. Daly PA, Krieger DR, Dulloo AG, Young JB, Landsberg L. Ephedrine, caffeine and aspirin: safety and efficacy for treatment of human obesity. *Int J Obes Relat Metab Disord.* 1993;17 (Suppl 1):S73-S78.

698. Marin P, Holmang S, Jonsson L, et al. The effects of testosterone treatment on body composition and metabolism in middle-aged obese men. *Int J Obes Relat Metab Disord.* 1992;16:991-997.
699. Rasmussen MH, Andersen T, Breum L, Gotzsche PC, Hilsted J. Cimetidine suspension as adjuvant to energy restricted diet in treating obesity. *BMJ.* 1993;306:1093-1096.
700. Coon PJ, Bleecker ER, Drinkwater DT, Meyers DA, Goldberg AP. Effects of body composition and exercise capacity on glucose tolerance, insulin, and lipoprotein lipids in healthy older men: a cross-sectional and longitudinal intervention study. *Metabolism.* 1989;38:1201-1209.
701. Franz MJ, Monk A, Barry B, et al. Effectiveness of medical nutrition therapy provided by dietitians in the management of non-insulin-dependent diabetes mellitus: a randomized, controlled clinical trial. *J Am Diet Assoc.* 1995;95:1009-1017.
702. Milne RM, Mann JI, Chisholm AW, Williams SM. Long-term comparison of three dietary prescriptions in the treatment of NIDDM. *Diabetes Care.* 1994;17:74-80.
703. Ready AE, Fitzpatrick DW, Boreskie SL, Hrycaiko DW. The response of obese females to low impact exercise and diet counselling. *J Sports Med Phys Fitness.* 1991;31:587-595.
704. Singman HS, Berman SN, Cowell C, Maslansky E, Archer M. The Anti-Coronary Club: 1957 to 1972. *Am J Clin Nutr.* 1980;33:1183-1191.
705. Suter E, Marti B, Gutzwiller F. Jogging or walking—comparison of health effects. *Ann Epidemiol.* 1994;4:375-381.
706. Viddal KO. Intestinal bypass. A randomized, prospective clinical study of end-to-side and end-to-end jejunoileal bypass. *Scand J Gastroenterol.* 1983;18:627-634.
707. Williams PT, Wood PD, Haskell WL, Vranizan K. The effects of running mileage and duration on plasma lipoprotein levels. *JAMA.* 1982;247:2674-2679.
708. Wing RR, Marcus MD, Salata R, Epstein LH, Miaskiewicz S, Blair EH. Effects of a very low-calorie diet on long-term glycemic control in obese Type 2 diabetic subjects. *Arch Intern Med.* 1991;151:1334-1340.
709. Wing RR, Blair E, Marcus M, Epstein LH, Harvey J. Year-long weight loss treatment for obese patients with type II diabetes: does including an intermittent very low-calorie diet improve outcome? *Am J Med.* 1994;97:354-362.
710. Drent ML, Larsson I, William-Olsson T, et al. Orlistat (Ro 18-0647), a lipase inhibitor, in the treatment of human obesity: a multiple dose study. *Int J Obes Relat Metab Disord.* 1995;19:221-226.
711. Pedrinola F, Szejnsznajd C, Lima N, Halpern A, Medeiros-Neto G. The addition of dexfenfluramine to fluoxetine in the treatment of obesity: a randomized clinical trial. *Obes Res.* 1996;4:549-554.
712. Connolly VM, Gallagher A, Kesson CM. A study of fluoxetine in obese elderly patients with type 2 diabetes. *Diabet Med.* 1995;12:416-418.
713. Alpert MA, Terry BE, Lambert CR, et al. Factors influencing left ventricular systolic function in nonhypertensive morbidly obese patients, and effect of weight loss induced by gastroplasty. *Am J Cardiol.* 1993;71:733-737.
714. Cairella G, Cairella M, Marchini G. Effect

- of dietary fibre on weight correction after modified fasting. *Eur J Clin Nutr.* 1995;49 (Suppl 3):S325-S327.
715. Foster GD, Wadden TA, Peterson FJ, Letizia KA, Bartlett SJ, Conill AM. A controlled comparison of three very low-calorie diets: effects on weight, body composition, and symptoms. *Am J Clin Nutr.* 1992;55:811-817.
716. Hakala P. Weight reduction programmes at a rehabilitation centre and a health centre based on group counselling and individual support: short- and long-term follow-up study. *Int J Obes Relat Metab Disord.* 1994;18:483-489.
717. Han TS, Richmond P, Avenell A, Lean ME. Waist circumference reduction and cardiovascular benefits during weight loss in women. *Int J Obes Relat Metab Disord.* 1997;21:127-134.
718. Lean ME, Han TS, Prvan T, Richmond PR, Avenell A. Weight loss with high and low carbohydrate 1200 kcal diets in free living women. *Eur J Clin Nutr.* 1997;51:243-248.
719. Miura J, Arai K, Tsukahara S, Ohno M, Ikeda Y. The long-term effectiveness of combined therapy by behavior modification and very low-calorie diet: 2 years followup. *Int J Obes.* 1989;13 (Suppl 2):73-77.
720. Walker KZ, O'Dea K, Johnson L, et al. Body fat distribution and non-insulin-dependent diabetes: comparison of a fiber-rich, high-carbohydrate, low-fat (23%) diet and a 35% fat diet high in monounsaturated fat. *Am J Clin Nutr.* 1996;63:254-260.
721. Foster GD, Wadden TA, Feurer ID, et al. Controlled trial of the metabolic effects of a very low-calorie diet: short- and long-term effects. *Am J Clin Nutr.* 1990;51:167-172.
722. Golay A, Allaz AF, Morel Y, de Tonnac N, Tankova S, Reaven G. Similar weight loss with low- or high-carbohydrate diets. *Am J Clin Nutr.* 1996;63:174-178.
723. Rossner S, Flaten H. VLCD versus LCD in long-term treatment of obesity. *Int J Obes Relat Metab Disord.* 1997;21:22-26.
724. Jarrett RJ, Keen H, Murrells T. Changes in blood pressure and body weight over ten years in men selected for glucose intolerance. *J Epidemiol Community Health.* 1987;41:145-151.
725. Gwinup G. Weight loss without dietary restriction: efficacy of different forms of aerobic exercise. *Am J Sports Med.* 1987;15:275-279.
726. Hespel P, Lijnen P, Van Hoof R, et al. Effects of physical endurance training on the plasma renin-angiotensin-aldosterone system in normal man. *J Endocrinol.* 1988;116:443-449.
727. King AC, Taylor CB, Haskell WL, DeBusk RF. Influence of regular aerobic exercise on psychological health: a randomized, controlled trial of healthy middle-aged adults. *Health Psychol.* 1989;8:305-324.
728. Karvetti RL, Knuts LR. Effects of comprehensive rehabilitation on weight reduction in myocardial infarction patients. *Scand J Rehabil Med.* 1983;15:11-16.
729. Liao Y, Emidy LA, Gosch FC, Stamler R, Stamler J. Cardiovascular responses to exercise of participants in a trial on the primary prevention of hypertension. *J Hypertens.* 1987;5:317-321.
730. Darga LL, Carroll-Michals L, Botsford SJ, Lucas CP. Fluoxetine's effect on weight loss in obese subjects. *Am J Clin Nutr.* 1991;54:321-325.

731. Elmer PJ, Grimm R Jr, Laing B, et al. Lifestyle intervention: results of the Treatment of Mild Hypertension Study (TOMHS). *Prev Med.* 1995;24:378-388.
732. Foreyt JP, Ramirez AG, Cousins JH. Cuidando El Corazon—a weight-reduction intervention for Mexican Americans. *Am J Clin Nutr.* 1991;53:1639S-1641S.
733. Hogan RB, Johnston JH, Long BW, et al. A double-blind, randomized, sham-controlled trial of the gastric bubble for obesity. *Gastrointest Endosc.* 1989;35:381-385.
734. Marcus MD, Wing RR, Ewing L, Kern E, McDermott M, Gooding W. A double-blind, placebo-controlled trial of fluoxetine plus behavior modification in the treatment of obese binge-eaters and non-binge-eaters. *Am J Psychiatry.* 1990;147:876-881.
735. Stamler J, Briefel RR, Milas C, Grandits GA, Caggiula AW. Relation of changes in dietary lipids and weight, trial years 1-6, to changes in blood lipids in the special intervention and usual care groups in the Multiple Risk Factor Intervention Trial. *Am J Clin Nutr.* 1997;65:272S-288S.
736. Wing RR, Shoemaker M, Marcus MD, McDermott M, Gooding W. Variables associated with weight loss and improvements in glycemic control in type II diabetic patients in behavioral weight control programs. *Int J Obes.* 1990;14:495-503.
737. DeLucia JL, Kalodner CR, Horan JJ. The effect of two nutritional software programs used as adjuncts to the behavioral treatment of obesity. *J Subst Abuse.* 1988-1989;1:203-208.
738. DeLucia JL, Kalodner CR. An individualized cognitive intervention: does it increase the efficacy of behavioral interventions for obesity? *Addict Behav.* 1990;15:473-479.
739. Heitzmann CA, Kaplan RM, Wilson DK, Sandler J. Sex differences in weight loss among adults with type II diabetes mellitus. *J Behav Med.* 1987;10:197-211.
740. Jeffery RW, Forster JL, French SA, et al. The Healthy Worker Project: a work-site intervention for weight control and smoking cessation. *Am J Public Health.* 1993;83:395-401.
741. Jeffery RW, Forster JL, Snell MK. Promoting weight control at the worksite: a pilot program of self-motivation using payroll-based incentives. *Prev Med.* 1985;14:187-194.
742. Jones SE, Owens HM, Bennett GA. Does behaviour therapy work for dietitians? An experimental evaluation of the effects of three procedures in a weight reduction clinic. *Hum Nutr Appl Nutr.* 1986;40:272-281.
743. Kalodner CR, DeLucia JL. The individual and combined effects of cognitive therapy and nutrition education as additions to a behavior modification program for weight loss. *Addict Behav.* 1991;16:255-263.
744. Schwartz SH, Inbar-Saban N. Value self-confrontation as a method to aid in weight loss. *J Pers Soc Psychol.* 1988;54:396-404.
745. Breum L, Astrup A, Andersen T, et al. The effect of long-term dexfenfluramine treatment on 24-hour energy expenditure in man. A double-blind placebo controlled study. *Int J Obes.* 1990;14:613-621.
746. Finer N, Finer S, Naoumova RP. Drug therapy after very-low-calorie diets. *Am J Clin Nutr.* 1992;56:195S-198S.
747. Pasquali R, Cesari MP, Melchionda N, Stefanini C, Raitano A, Labo G. Does ephedrine promote weight loss in low-energy-adapted obese women? *Int J Obes.*

- 1987;11:163-168.
748. Weintraub M, Hasday JD, Mushlin AI, Lockwood DH. A double-blind clinical trial in weight control. Use of fenfluramine and phentermine alone and in combination. *Arch Intern Med.* 1984;144:1143-1148.
 749. Weintraub M, Sundaresan PR, Schuster B, et al. Long-term weight control study. II (weeks 34 to 104). An open-label study of continuous fenfluramine plus phentermine versus targeted intermittent medication as adjuncts to behavior modification, caloric restriction, and exercise. *Clin Pharmacol Ther.* 1992;51:595-601.
 750. Bitsch M, Skrumsager BK. Femoxetine in the treatment of obese patients in general practice. A randomized group comparative study with placebo. *Int J Obes.* 1987;11:183-190.
 751. Fernandez-Soto ML, Gonzalez-Jimenez A, Barredo-Acedo F, Luna del Castillo JD, Escobar-Jimenez F. Comparison of fluoxetine and placebo in the treatment of obesity. *Ann Nutr Metab.* 1995;39:159-163.
 752. Goldstein DJ, Rampey AH Jr, Enas GG, Potvin JH, Fludzinski LA, Levine LR. Fluoxetine: a randomized clinical trial in the treatment of obesity. *Int J Obes Relat Metab Disord.* 1994;18:129-135.
 753. Cook RF, Howard AN, Mills IH. Low-dose mianserin as adjuvant therapy in obese patients treated by a very-low-calorie diet. *Int J Obes.* 1981;5:267-272.
 754. Steel JM, Munro JF, Duncan LJ. A comparative trial of different regimens of fenfluramine and phentermine in obesity. *Practitioner.* 1973;211:232-236.
 755. Sax L. Yohimbine does not affect fat distribution in men. *Int J Obes.* 1991;15:561-565.
 756. Benjamin SB, Maher KA, Cattau EL Jr, et al. Double-blind controlled trial of the Garren-Edwards gastric bubble: an adjunctive treatment for exogenous obesity. *Gastroenterology.* 1988;95:581-588.
 757. Mathus-Vliegen EM, Tytgat GN, Veldhuyzen-Offermans EA. Intra-gastric balloon in the treatment of super-morbid obesity. Double-blind, sham-controlled, crossover evaluation of 500-milliliter balloon. *Gastroenterology.* 1990;99:362-369.
 758. Bray GA, Gray DS. Obesity. Part I—Pathogenesis. *West J Med.* 1988;149:429-441.
 759. Centers for Disease Control and Prevention. Number and percentage of children and adolescents who were overweight by gender and race/ethnicity: United States NHANES III, 1988-1994. *Morb Mortal Wkly Rep.* 1997.
 760. National Center of Health Statistics. Center of Disease Prevention and Control, National Health and Nutrition Examination Survey, Phase III. Prevalence of overweight in U.S. adults, by race/ethnicity, gender, and education, 1988-1991.
 761. Centers for Disease Control. Cardiac valvulopathy associated with exposure to Fenfluramine or Dexfenfluramine: U.S. Department of Health and Human Services interim public health recommendation, November 1997. *MMWR Morb Mortal Wkly Rep.* 1997;46:1061-1065.
 762. American Psychiatric Association. Task Force on DSM-IV. *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV.* 4th ed. Washington, DC: The Association; 1994.
 763. Landin K, Stigendal L, Eriksson E, et al. Abdominal obesity is associated with an impaired fibrinolytic activity and elevated

plasminogen activator inhibitor-I.
Metabolism. 1990;39:1044-1048.

764. Reisin E., Frolich ED, Messerli FH.
Dreslinski GR, et al. Cardiovascular
changes after weight reduction in obesity
hypertension. Ann Int Med. 1983;98:315-
319.
765. Tuck MI, Sowers J, Dornfield L, et al. The
effect of weight reduction on blood pres-
sure plasma renin activity and plasma
aldosterone level in obese patients. N Eng J
Med 1981;304:930-933.
766. Rocchini AP, Key J, Bondie D, et al. The
effect of weight loss on the sensitivity of
blood pressure to sodium in obese adoles-
cents. N Eng. J med 1989;321:580-585.
767. Landsberg L, Krieger DR, Obesity, metab-
olism and the sympathetic nervous system.
Am J Hypertension 1989;2:125s-132s.
768. Jacobs DB, Sowers JR, et al. Effects of
weight reduction on cellular cation metab-
olism and vascular resistance.
Hypertension. 1993;21:308-314.