



Enabling Medication Management Through Health Information Technology



Agency for Healthcare Research and Quality
Advancing Excellence in Health Care • www.ahrq.gov

**Evidence-Based
Practice**

**Health Information
Technology**

Evidence Report/Technology Assessment

Number 201

Enabling Medication Management Through Health Information Technology

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
540 Gaither Road
Rockville, MD 20850
www.ahrq.gov

Contract No. 290-2007-10060-I

Prepared by:

McMaster Evidence-based Practice Center
Hamilton, ON Canada

Investigators:

K. Ann McKibbin, Ph.D.
Cynthia Lokker, Ph.D.
Steve M. Handler, M.D., Ph.D.
Lisa R. Dolovich, Pharm.D., M.Sc.
Anne M. Holbrook, Pharm.D., M.D.
Daria O'Reilly, Ph.D.
Robyn Tamblyn, Ph.D.
Brian J. Hemens, B.Sc.Pharm., M.Sc.
Runki Basu, B.Sc. (Hons.), M.A.
Sue Troyan, B.A., R.T.
Pavel S. Roshanov, B.Sc.
Norman P. Archer, Ph.D.
Parminder Raina, Ph.D.

AHRQ Publication No. 11-E008-EF
April 2011

This report is based on research conducted by the McMaster Evidence-based Practice Center under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. 290-2007-10060-I). The findings and conclusions in this document are those of the author(s) who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ. No statement in this article should be construed as an official position of the Agency for Healthcare Research and Quality or of the U.S. Department of Health and Human Services.

The information in this report is intended to help health care decision-makers; patients and clinicians, health system leaders, and policymakers, make well-informed decisions and thereby improve the quality of health care services. This report is not intended to be a substitute for the application of clinical judgment. Decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information, i.e., in the context of available resources and circumstances presented by individual patients.

This report may be used, in whole or in part, as the basis for development of clinical practice guidelines and other quality enhancement tools, or as a basis for reimbursement and coverage policies. AHRQ or U.S. Department of Health and Human Services endorsement of such derivative products may not be stated or implied.

This document is in the public domain and may be used and reprinted without permission except those copyrighted materials noted for which further reproduction is prohibited without the specific permission of copyright holders.

No investigators have any affiliations or financial involvement (e.g., employment, consultancies, honoraria, stock options, expert testimony, grants or patents received or pending, or royalties) that conflict with material presented in this report.

Suggested citation: McKibbin KA, Lokker C, Handler SM, Dolovich LR, Holbrook AM, O'Reilly D, Tamblyn R, Hemens BJ, Basu R, Troyan S, Roshanov PS, Archer NP, Raina P. Enabling Medication Management Through Health Information Technology. Evidence Report/Technology Assessment No. 201. (Prepared by the McMaster University Evidence-based Practice Center under Contract HHS 290-2007-10060-I). AHRQ Publication No. 11-E008-EF. Rockville MD: Agency for Healthcare Research and Quality. April 2011.

Acknowledgments

The research team would like to thank those who helped with screening, abstracting, and article retrievals: Danika Walden, Nathan McKibbin, Catherine Salmon, Jan Burke-Gaffney, Connie Freeborn, Jamie O'Donnell, Rizwan Bhaloo, Bryan Cheeseman, Hafsa Jan Qureshi, and Pouyan Ahangar. We would like to offer special thanks to Mary Gauld, Maureen Rice, and Roxanne Cheeseman for assistance and guidance with project management and editorial help. The project would not be complete without their wisdom, experience, good will, and sense of humor. Further thanks to Dr. Chris Gibbons and Dr. Paul Gorman who provided insights into their work on previous AHRQ evidence reports at the start of this project. We acknowledge the hard work of Nicholas Hobson, our computer programmer, for creating our evolving systematic review management software. The report is stronger and more focused because of the continued expert guidance of Rebecca Roper of the Agency for Healthcare Research and Quality as our Task Order Officer (TOO).

Our Technical Expert Panel provided valuable insights and challenges as well as ways to meet them. Our technical experts were David Bates, Doug Bell, Ken Boockvar, Chris Gibbons, Joy Grossman, Jerry Gurwitz, Joe Hanlon, Kevin Johnson, John Poikonen, Gordon Schiff, Bimla Schwarz, and Dennis Tribble. Their contact information is included in Appendix D. They represent a broad range of expertise and experience and the report is stronger because of them. Another group of experts who have been extremely helpful at improving the analyses of our data were our technical reviewers. Some of those who reviewed our first draft were members of the Technical Expert Panel: Joy Grossman, Joe Hanlon, Kevin Johnson, Bimla Schwartz, John Poikonen, Dennis Tribble, and our TOO Rebecca Roper. Other expert reviewers were Anne Bobb, Elizabeth Chrischilles, Alan Flynn, and Kevin Marvin. Their contact information is also in Appendix D.

Enabling Medication Management Through Health Information Technology

Structured Abstract

Objective. The objective of the report was to review the evidence on the impact of health information technology (IT) on all phases of the medication management process (prescribing and ordering, order communication, dispensing, administration and monitoring as well as education and reconciliation), to identify the gaps in the literature and to make recommendations for future research.

Data sources. We searched peer-reviewed electronic databases, grey literature, and performed hand-searches. Databases searched included MEDLINE,[®] EMBASE,[®] CINAHL[®] (Cumulated Index to Nursing and Allied Health Literature), Cochrane Database of Systematic Reviews, International Pharmaceutical Abstracts,[®] Compendex,[®] INSPEC[®] (which includes IEEE[®]), Library and Information Science Abstracts,[®] E-Prints in Library and Information Science,[®] PsycINFO,[®] Sociological Abstracts,[®] and Business Source[®] Complete. Grey literature searching involved Internet searching, reviewing relevant Web sites, and searching electronic databases of grey literatures. AHRQ also provided all references in their e-Prescribing, bar coding, and CPOE knowledge libraries.

Methods. Paired reviewers looked at citations to identify studies on a range of health IT used to assist in the medication management process (MMIT) during multiple levels of screening (titles and abstracts, full text and final review for assignment of questions and data abstraction). Randomized controlled trials and cohort, case-control, and case series studies were independently assessed for quality. All data were abstracted by one reviewer and examined by one of two different reviewers with content and methods expertise.

Results. 40,582 articles were retrieved. After duplicates were removed, 32,785 articles were screened at the title and abstract phase. 4,578 full text articles were assessed and 789 articles were included in the final report. Of these, 361 met only content criteria and were listed without further abstraction. The final report included data from 428 articles across the seven key questions. Study quality varied according to phase of medication management. Substantially more studies, and studies with stronger comparative methods, evaluated prescribing and monitoring. Clinical decision support systems (CDSS) and computerized provider order entry (CPOE) systems were studied more than any other application of MMIT. Physicians were more often the subject of evaluation than other participants. Other health care professionals, patients, and families are important but not studied as thoroughly as physicians. These nonphysicians groups often value different aspects of MMIT, have diverse needs, and use systems differently. Hospitals and ambulatory clinics were well-represented in the literature with less emphasis placed on long-term care facilities, communities, homes, and nonhospital pharmacies. Most studies evaluated changes in process and outcomes of use, usability, and knowledge, skills, and attitudes. Most showed moderate to substantial improvement with implementation of MMIT. Economics studies and those with clinical outcomes were less frequently studied. Those articles that did address economics and clinical outcomes often showed equivocal findings on the effectiveness and cost-effectiveness of MMIT systems. Qualitative studies provided evidence of

strong perceptions, both positive and negative, of the effects of MMIT and unintended consequences. We found little data on the effects of forms of medications, conformity, standards, and open source status. Much descriptive literature discusses implementation issues but little strong evidence exists. Interest is strong in MMIT and more groups and institutions will implement systems in the next decades, especially with the Federal Government's push toward more health IT to support better and more cost-effective health care.

Conclusions. MMIT is well-studied, although on closer examination of the literature the evidence is not uniform across phases of medication management, groups of people involved, or types of MMIT. MMIT holds the promise of improved processes; clinical and economics studies and the understanding of sustainability issues are lacking.

Contents

| | |
|---|------|
| Executive Summary | ES-1 |
| Introduction..... | 1 |
| Scope and Purpose of the Systematic Review | 1 |
| Key Questions (KQs)..... | 2 |
| Background..... | 4 |
| Methods..... | 6 |
| Recruitment of Technical Experts and Peer Reviewers..... | 6 |
| Key Questions..... | 6 |
| Analytic Framework | 7 |
| Literature Search Methods..... | 8 |
| Sources..... | 9 |
| Search Terms and Strategies..... | 9 |
| Organization and Tracking of the Literature Search..... | 9 |
| Title and Abstract Review | 9 |
| Defining Medication Management Health IT..... | 10 |
| Data Abstraction | 11 |
| Assessment of Study Quality..... | 11 |
| Data Synthesis..... | 13 |
| Data Entry and Quality Control | 13 |
| Grading the Evidence..... | 13 |
| Peer Review | 13 |
| Results..... | 14 |
| Results of the Literature Search..... | 14 |
| KQ1. Within and Across the Phases of Medication Management Continuum, What Evidence Exists That Health IT Applications Are Effective?..... | 16 |
| Effectiveness Studies Overall | 16 |
| Strengths and Limitations of the Evidence | 17 |
| Process Changes—Prescribing | 21 |
| Summary of the Findings for Process Changes | 21 |
| General Study Characteristics..... | 21 |
| Outcomes | 23 |
| Summary..... | 25 |
| Order Communication | 26 |
| Summary of the Findings for Process Changes | 26 |
| Strengths and Limitations of the Evidence | 26 |
| General Study Characteristics..... | 26 |
| Outcomes | 26 |
| Summary..... | 27 |
| Dispensing..... | 28 |
| Summary of the Findings for Process Changes | 28 |
| Strengths and Limitations of the Evidence | 28 |
| General Study Characteristics..... | 29 |
| Outcomes | 29 |
| Summary..... | 30 |
| Administering | 30 |

| | |
|--|----|
| Summary of the Findings for Process Changes | 30 |
| Strengths and Limitations of the Evidence | 30 |
| General Study Characteristics | 31 |
| Outcomes | 31 |
| Monitoring | 32 |
| Summary of the Findings for Process Changes | 32 |
| General Study Characteristics | 33 |
| Outcomes | 34 |
| Reconciliation, Discharge Summaries, and Education | 35 |
| Summary of the Findings for Process Changes | 35 |
| Combined Phases of Medication Management | 36 |
| Summary of the Findings for Process Changes | 36 |
| PDA's | 37 |
| Summary of the Findings for PDA's | 37 |
| Intermediate Outcomes | 37 |
| Summary of the Findings | 37 |
| Strengths and Limitations of the Evidence | 38 |
| General Study Characteristics | 38 |
| Outcomes | 40 |
| Economic Outcomes | 44 |
| Full Economic Evaluations | 44 |
| Partial Economic Evaluations | 45 |
| Economics Summary | 50 |
| Clinical Outcomes | 51 |
| Summary of the Findings | 51 |
| Outcomes | 54 |
| Prescribing—Strengths and Limitations of the Evidence | 54 |
| Prescribing—General Study Characteristics | 55 |
| Prescribing—Clinical Outcomes | 55 |
| Qualitative Studies | 57 |
| Summary of Findings | 57 |
| Strengths and Limitations of Evidence | 57 |
| Population Level Outcomes | 65 |
| Composite Outcomes | 65 |
| Variation in Impact Depending on Medication Type or Form | 66 |
| Summary of the Findings | 66 |
| Strengths and Limitations of the Evidence | 67 |
| General Study Characteristics | 67 |
| Outcomes | 68 |
| Unintended Consequences of MMIT Applications | 69 |
| Summary of the Findings | 69 |
| Strengths and Limitations of the Evidence | 69 |
| General Study Characteristics | 70 |
| Outcomes | 70 |
| KQ2. What Knowledge or Evidence Deficits Exist Regarding Needed Information To Support Estimates of Cost, Benefit, Impact, and Net Value With Regard To Enabling Health | |

| | |
|---|----|
| IT Applications in Terms of Prescribing, Order Transmission, Dispensing, Administering, and Monitoring, as Well as Reconciliation, Education, and Adherence? Discuss Gaps in Research, Including Specific Areas That Should Be Addressed and Suggest Possible Public and Private Organizational Types To Perform the Research and/or Analysis..... | 71 |
| Introduction..... | 71 |
| General Gaps..... | 73 |
| Summary..... | 79 |
| KQ3. What Critical Information Regarding the Impact of Health IT Applications Implemented To Support the Phases of Medication Management Is Needed To Give Clinicians (Physicians, Nurses, Psychologists, Dentists), Pharmacists, Health Care Administrators, Patients, and Their Families a Clear Understanding of the Value Proposition Particular to Them?..... | 79 |
| Summary of the Findings..... | 81 |
| Financial Benefits | 81 |
| Clinical Benefits..... | 81 |
| Organizational Benefits | 82 |
| Conclusions..... | 83 |
| KQ4. What Evidence Exists Regarding the Impact of the Characteristics of Medication Management Health IT Applications, Such as Open Source, Proprietary, Conformity With Federal and Other Interoperability Standards, and Being Certification Commission for Healthcare Information Technology (CCHIT) Certified, Impact, Likelihood for Purchase, Implementation, and Use of Such IT Applications..... | 83 |
| Summary of the Findings..... | 83 |
| Strengths and Limitations of the Evidence | 85 |
| General Study Characteristics..... | 85 |
| Outcomes | 85 |
| KQ5. What Factors Influence Sustainability (Use and Periodic Updates) of Health IT Applications That Support a Phase of Medication Management Continuum (Prescribing, Dispensing, Administering, and Patients’ Taking of Medications)? | 88 |
| Sustainability of Health IT and Medication Management Systems..... | 88 |
| Future Sustainability of Health IT and Medication Management Systems | 89 |
| Conclusions..... | 89 |
| 5a. To What Extent Does the Evidence Demonstrate That Health Care Settings (Inpatient, Ambulatory, Long-Term Care, etc.) Influence Implementation, Uuse, and Effectiveness of Such Health IT Applications?..... | 92 |
| Implementation | 92 |
| Effectiveness..... | 93 |
| Use | 93 |
| 5b. What is the Impact (Challenges, Merits, Costs, and Benefits) of Having Electronic Access to Patients’ Computerized Medication Records, Formulary Information, Billing Information, Laboratory Records in the Quality and Safety of Care Provided by Health IT Applications That Support at Least One Phase of the Continuum of Medication Management (Prescribing, Dispensing, Administering, and Patients Taking of Medications)?..... | 93 |
| KQ6. Two-Way Prescriber and Pharmacy Electronic Data Interchange (e-Prescribing) | |
| (a) What Evidences Exists Demonstrating the Barriers and Drivers of Implementation of Complete EDI That Can Support the Prescription, Transmittal and Receipt, and Perfection | |

| | |
|--|-----|
| Process of e-Prescriptions? | |
| (b) How Do Barriers, Facilitators, and Economic Incentives Vary Across Pharmacists, Physicians, and Other Relevant Stakeholders With Respect to Adoption and Use of Complete EDI (e-Prescribing/Ordering With e-Transmission)? | 94 |
| Summary of the Findings | 94 |
| Summary of Evidence | 96 |
| KQ7. What Evidence Exists Regarding the Extent of Integration of Electronic Clinical Decision Support (CDS) in a Health IT System for Prescribing and Dispensing of Medications? | |
| To What Extent Does the Use of CDSS in a Health IT System for Prescribing and Dispensing of Medications Impact the Various Outcomes of Interest Including Health Care Process, Intermediate and Clinical? | 96 |
| Summary of the Findings: All Phases of Medication Management | 96 |
| Discussion | 100 |
| Summary of Key Findings | 100 |
| KQ1. Effectiveness | 100 |
| KQ2. Gaps in Evidence and Knowledge | 100 |
| KQ3. Value Proposition for Implementers and Users | 101 |
| KQ4. System Characteristics | 101 |
| KQ5. Sustainability | 101 |
| KQ6. Two-Way EDI | 102 |
| KQ7. RCTs in CDSS | 102 |
| CDSS | 105 |
| Future Research | 109 |
| Need for High Quality Evidence | 111 |
| Need for Well-Designed Research | 111 |
| Conclusions | 116 |
| References | 118 |
| Acronyms and Abbreviations | 160 |

Tables

| | |
|--|------|
| Table A. Research Design for studies across the Phases of Medication Management and Education and Reconciliation | ES-6 |
| Table B. Settings for the Phases of Medication Management and Reconciliation and Education | ES-6 |
| Table C. Clinicians Evaluated in Outcomes Studies of Medication Management Phases, Education, and Reconciliation | ES-7 |
| Table D. Main Health IT Studied by Medication Management Phase and Education and Reconciliation | ES-7 |
| Table 1. Research Design for Studies Across the Phases of Medication Management and Education and Reconciliation | 16 |
| Table 2. Settings for the Phases of Medication Management and Reconciliation and Education | 18 |
| Table 3. Clinicians Evaluated in Outcomes Studies of Medication Management Phases, Education, and Reconciliation | 18 |
| Table 4. Patients and Caregivers Studied by Phase of Medication Management and Education and Reconciliation | 19 |

| | |
|--|-----|
| Table 5. Main Health IT Studied by Medication Management Phase and Education and Reconciliation | 20 |
| Table 6. Health IT Integrated With the Health IT Being Studied..... | 20 |
| Table 7. Research Methods of Studies That Evaluated Process Changes Associated With the Prescribing Phase of MMIT | 21 |
| Table 8. Summary of the Number of Studies Reporting Statistically Significant Process Changes in Studies of Prescribing by Process for Hospital and Ambulatory Based Studies | 24 |
| Table 9. Summary of the Number of Statistically Significant Process Changes in Studies of Order Communication by Process for Hospital and Ambulatory Based Studies | 28 |
| Table 10. Intermediate Outcomes Across the Phases for Medication Management | 38 |
| Table 11. Study Designs Used in Studies Measuring Intermediate Outcomes Across the Phases for Medication Management..... | 38 |
| Table 12. Clinician Study Participants in Studies Assessing Intermediate Outcomes Across the Phases of Medication Management | 39 |
| Table 13. Patient Study Participants in Studies Assessing Intermediate Outcomes Across the Phases of Medication Management | 39 |
| Table 14. Research Design for Studies Across the Phases of Medication Management and Education and Reconciliation That Address Clinical Outcomes as Their Main Outcomes | 52 |
| Table 15. Summary of the Number of Studies Reporting Statistically Significant Differences in Clinical Primary Endpoints Between Study Groups for Hospital and Ambulatory Based Studies..... | 52 |
| Table 16. Number of Studies Across the Medication Management Phases Using MMIT To Assist in the Management of Specific Drugs or Drug Classes..... | 68 |
| Table 17. Summary of Gaps and Needs Across Key Questions..... | 72 |
| Table 18. Frequency of Medication Management Phases Studies Plus Reconciliation and Education..... | 74 |
| Table 19. Frequency of Research Designs for Included Studies | 74 |
| Table 20. Number of Studies That Evaluated the Effects of MMIT on Clinicians (the Clinicians Were the Major Focus of the Outcomes of the Articles)..... | 75 |
| Table 21. Frequency With Which Patients or Caregivers Across Age Groups Were Studied as the Main Focus of an Article (How MMIT Affects Patients)..... | 75 |
| Table 22. Study Settings in Which the MMIT Application Was Studied (Studies Could Take Place in More Than One Setting)..... | 76 |
| Table 23. Technologies That Were the Main Focus of the Studies of MMIT | 76 |
| Table 24. Summary of the Evidence in Relation to the CITL Value Framework | 80 |
| Table 25. Number of Articles Addressing System Type in Relation to Likelihood To Purchase, Implement, or Use an MMIT System | 84 |
| Table 26. List of Articles Addressing Various Features That Were Instrumental in the Decision To Purchase, Implement, and Use | 86 |
| Table 27. Frequency of Core Informatics Journal Articles That Mention Sustainability to the End of 2009 | 89 |
| Table 28. Issues of Consideration and/or Further Exploration in Future Research..... | 109 |
| Table 29. Study Design of Included Studies Across the Medication Management Phases (Plus Education and Reconciliation)..... | 111 |

Figures

| | |
|--|----|
| Figure 1. A Functional Model of the Medication Management Continuum Created by Bell et al. | 2 |
| Figure 2. Conceptual Model Addressing the Seven Key Questions: Enabling Medication Management Through Health IT. | 8 |
| Figure 3. Literature Flow of Medication Management Studies..... | 15 |
| Figure 4. Trends in Publication of Articles Relating to the Phases of Medication Management Across Years Until Searching Was Completed in June 2010..... | 17 |
| Figure 5. Summary Overview of Meaningful Use Objectives | 91 |

Appendixes

| | |
|---|--|
| Appendix A. Exact Search Strings | |
| Appendix B. Sample Screening and Data Abstraction Forms | |
| Appendix C. Evidence Tables | |
| Appendix D. Technical Expert Panel and Peer Reviewers | |
| Appendix E. Excluded Studies | |
| Appendix F. Glossary of Terms | |

Executive Summary

Background

Medication management is a continuum that covers all aspects of prescription medications. Medication management includes prescribing and ordering, order communication (or order transmission) between prescribers and pharmacists, dispensing, administering, and monitoring, as well as reconciliation, adherence, and education.¹ Medication management is complex and costly and enhances the health and well-being of more than half of the population in the developing world. Health information technology (health IT) holds great promise to improve the quality of health care and reduce potential and real errors in medication management while at the same time providing cost-effective care. The Agency for Healthcare Research and Quality (AHRQ) is committed to summarizing and providing the evidence base for health IT. It has produced evidence summaries on health IT related to costs and benefits;² barriers and drivers of health IT for the elderly, chronically ill, and underserved;³ the impact of consumer informatics applications;⁴ and telemedicine.⁵ AHRQ also has contracted for evidence summaries on the use of health IT in decisionmaking,⁶ patient-centered care,⁷ and decision support for health care decisionmaking.⁸ The contracted reports will be available through www.healthit.AHRQ.gov in mid-2011. Although these reports often mention medication management, the body of published evidence on all aspects of the medication management process and how it is affected by multiple health IT systems has not been consolidated. A single document is needed to summarize the evidence evaluating the effects of health IT on the medication management process across providers, settings, patients, and research methods.

The objectives of this report are to:

1. Review the literature on the effects of health IT on medication management.
2. Synthesize available evidence regarding the effectiveness and effects of health IT in all phases of medication management as well as reconciliation and education.
3. Identify gaps in the literature.
4. Make recommendations for future research.

For the purposes of this review, medication management includes the processes that encompass the five phases of the medication process (i.e., prescribing and ordering, order communication, dispensing, administering, and monitoring) across groups of health professionals, patients, and their informal caregivers, and two aspects of quality with respect to medication management across the five phases of medication management (medication reconciliation and education, both postprofessional education of training and patient education related to medication management). Medication management can also include procurement, storage, and reporting from the first assessment of patients to determine their need for drugs through to optimal care and monitoring after the drugs are prescribed. The organization of the information in this report is based on the Bell framework of the five phases across the continuum of medication management and reconciliation and education.¹

To address the goals of this report, we further define medication management health IT (MMIT) applications as electronic systems that (1) collect, process, or exchange health information about patients; (2) are integrated with existing health IT systems such as electronic health records or electronic medical record (EMR) systems; and (3) provide advice or suggestions to either the health care provider or the patients and their families on issues or

decisions related to medication management. We recognize that functional elements of the MMIT will vary across particular implementation approaches within a given phase of medication management. Many of the MMIT applications we found were designed to encompass more than one phase of medication management. The sophistication of the systems, degree of integration of the health IT into workflow systems, and the broad range of settings in which a particular health IT is implemented and used are also complex and varied. Many health professionals, support staff, patients, and patients' families were involved in medication management in the studies assessed.

The evidence assessing MMIT is large, diffuse, and published across many disciplines. People who can benefit from the knowledge in this report include health professionals, researchers, administrators, and other decisionmakers and those who develop and implement health IT applications. This report is timely because of the Federal emphasis on the use of health IT to improve health care while at the same time making health and wellness care more cost effective and safer. Seven questions structure this evidence report. Within reporting related to the questions, sections are based on phases of medication management. Reporting is done to address the multiple settings where medication management is important, the range of health care providers who deliver and support care using medications, and classes of medications, specific drugs, or a broad spectrum of medications.

Key Questions (KQs)

KQ1. Effectiveness

Within all phases of the medication management continuum, what evidence exists that health IT applications are effective in improving:

- a. Health care processes,
- b. Other intermediate outcomes (e.g., satisfaction with system, usability, knowledge, skills, and attitude),
- c. Costs and economic outcomes,
- d. Clinical outcomes for patients,
- e. Population level outcomes, and
- f. Composite outcomes.
- g. To what extent does the impact of health IT on improving health care processes, other outcomes, costs and economics, and clinical outcomes vary depending on the type of medication (controlled or noncontrolled substance) or the form of the medication (e.g., oral, injection, intravenous)?

KQ2. Gaps in Knowledge or Evidence

What knowledge or evidence deficits exist to support estimates of cost, benefit, impact, and net value with regard to health IT applications in all phases of medication management?

KQ3. Value Proposition for Implementers and Users

What critical information regarding the impact of health IT applications implemented to support the phases of medication management is needed to give clinicians, health care facility administrators, patients, and their families a clear understanding of the value proposition particular to them?

KQ4. System Characteristics

What evidence supports or refutes the impact of any of: open source, homegrown, proprietary, local configuration ability, system configuration ability, conformity with standards being Certification Commission for Healthcare Information Technology (CCHIT) certified, system architecture, or feature set on the decision to purchase, implement, or use health IT in medication management systems?

KQ5. Sustainability

What factors influence sustainability of health IT applications that support a phase of the medication management continuum?

- a. What evidence exists to demonstrate that health care settings (ambulatory, long-term care, etc.) influence implementation, use, and effectiveness of such health IT applications?
- b. What is the impact (challenges, merits, costs, and benefits) of having electronic access to patient data on the quality and safety of care provided by health IT applications that support at least one phase of the continuum of medication management?

KQ6. Two-Way Prescription Electronic Data Interchange (EDI)

In a two-way electronic data interchange (EDI) between the prescribers and pharmacists:

- a. What evidence exists demonstrating the barriers and drivers of implementation of complete EDI that can support the prescription, transmittal and receipt, and perfection process of e-Prescriptions?
- b. How do barriers, facilitators, and economic incentives vary across pharmacists, physicians, and other relevant stakeholders with respect to adoption and use of complete EDI (e-Prescribing/ordering with e-Transmission)?

KQ7. Randomized Controlled Trials (RCTs) of Clinical Decision Support Systems (CDSS)

What evidence exists regarding the extent of integration of electronic clinical decision support in a health IT system for the prescribing, dispensing, and administering of medications, and to what extent does the use of clinical decision support systems impact the various outcomes (e.g., health care process, intermediate, cost and economics, and clinical) of interest?

Methods

We anticipated finding few RCTs across all phases of medication management and MMIT applications. Studies that employ other research methods can also provide valuable evidence for understanding MMIT applications. We therefore included studies employing a range of research methodologies. We restricted our analysis to hypothesis-driven studies with group comparisons and appropriate statistical analysis in addition to qualitative studies with explicit methods for KQ1: Effectiveness. The only methodological limit was for assessment of the effect of CDSSs on prescribing, for which sufficient RCTs were available to provide evidence for synthesis.

Through consultation with our internal team and AHRQ, we determined that the answers to KQ2: Gaps in Knowledge or Evidence and KQ3: Value Proposition for Implementers and Users would become evident from our review of the evidence in KQ1: Effectiveness. We supplemented these articles with other studies addressing values propositions by stakeholders. KQ4: System

Characteristics addresses the impact of MMIT application features on the likelihood that the systems will be purchased, implemented, and used. The evidence for this question comes from studies of all designs that measure implementation, use, and purchasing decisions. KQ5: Sustainability addresses the factors influencing the sustainability of MMIT applications, specifically the impact of the setting and access to other electronic data within integrated systems on health care quality and safety. To identify articles that addressed this question, the team, in consultation with AHRQ, used the definition of sustainability by Humphreys et al.,⁹ which restricted our choice of articles to only a few. Their definition of sustainability was the ability of a health service to provide ongoing access to appropriate quality care in a cost- and health-effective manner. KQ6: Two-Way Prescription EDI relates to the barriers and facilitators to complete EDI between prescribers and pharmacies during the time between prescription writing and dispensing and how these vary across stakeholders. The best evidence available for KQ6 is found in articles studying EDI between prescribers and pharmacies that include original data (qualitative or quantitative). Because insufficient evidence was found on two-way EDI, we included one-way EDI as well. KQ7: RCTs of CDSS addresses the extent to which CDSS systems are integrated into health IT systems for medication management and the impact on outcomes as described in KQ1: Effectiveness. As a team we felt that adequate evidence was available to address this issue so that we could limit our scope to RCTs.

Given the broad range of questions and outcomes addressed, we searched peer-reviewed electronic databases by first using textwords relating to the various types of health IT applied to medication management (Appendix A of the full report). These searches were then combined with a search using subject headings related to the five medication management phases plus reconciliation and education as well as specific health IT application terms (e.g., CDSS). We combined these medication management terms with computer and technology terms. When possible, we excluded letters, editorials, commentaries, and animal studies. Because our interest was in all study designs, we did not limit based on methodology. We also put no limits on language or time to capture the global literature and early studies.

Databases searched included MEDLINE, Embase, CINAHL (Cumulated Index to Nursing and Allied Health Literature), Cochrane Database of Systematic Reviews, International Pharmaceutical Abstracts, Compendex, Inspec (which includes IEEE Xplore), Library and Information Science Abstracts, E-Prints in Library and Information Science, PsycINFO, Sociological Abstracts, and Business Source Complete. We also looked for eligible studies by reviewing grey literature sites, performing hand searches of pertinent reviews, querying our experts, and by reviewing the AHRQ National Resource Center for Health IT Knowledge Library resources (available at: http://healthit.ahrq.gov/portal/server.pt/community/knowledge_library/653).

The search results were downloaded into Reference Manager version 10 (ISI ResearchSoft) and uploaded into a customized systematic review management system (Health Information Research Unit, McMaster University).

Studies were eligible for inclusion if they used health IT in any aspect of the medication management process. We included articles on MMIT only if the system was integrated with at least one existing health IT system and if they processed patient-specific information and provided advice or suggestions. A critical inclusion requirement was the integration of information.

Personal digital assistants (PDAs), which integrated patient-specific information provided by either the clinicians or the patients, were analyzed to assist in medication management decisions

(by request of AHRQ). This exception is made because PDAs and hand-held devices are considered an important, and perhaps unique, means of improving health care quality in relation to medications. The use of PDAs to manage medications is especially important for clinicians and patients who are in settings that do not have large, sophisticated, and integrated information systems. Other stand-alone devices with no integration of information with another health IT were excluded. Articles on all five phases of the management process plus medication reconciliation and postprofessional education related to MMIT were included. Once we tagged the articles for content, we assessed whether those that passed our inclusion criteria were pertinent to specific key questions. Many articles were analyzed in several phases of medication management and sections of the report.

Studies were classified as being observational, case-control, cohort, or RCTs. The quality of included studies was assessed using the same criteria employed by Jimison et al. in their AHRQ report.³ RCT scoring was based on the Delphi consensus work done by Verhagan and colleagues.¹⁰ This scale is referred to in this report as the Verhagen/AHRQ RCT quality scale. Observational studies with before–after, time series, surveys, or qualitative methods were not assessed for quality because few well-validated instruments exist. Bibliographies of systematic and narrative reviews were examined to identify studies, and select reviews were integrated into sections of the report.

Data were abstracted from relevant articles and tagged for applicability to the various key questions. Given the range of questions addressed, data abstraction was performed by a core group of staff and entered into online data abstraction forms. One reviewer did the abstraction, and a second, senior reviewer checked its accuracy. The authors of this report performed a final check on the abstracted data. The reviewers were not blinded to the identity of the article authors, institutions, or journal. Data abstraction was difficult in many instances because of the lack of accepted definitions and absence of important features of the study or MMIT application. For example, we identified problems with the differences between computerized provider order entry (CPOE) for ordering and e-Prescribing systems. Definitions for medication errors and related terms were often inconsistently used. To make data abstraction easier, we established working definitions, which can be found in Appendix F of the full report.

Meta-analysis was not performed on any data because of the heterogeneity of the studies in terms of interventions, populations, technologies used, and outcomes measured, as well as the presence of mostly descriptive and observational studies.

Throughout the project, the core team sought feedback from the internal advisors, our Task Officer from AHRQ, and the Technical Expert Panel.

Results

Our literature search retrieved 40,582 articles. After duplicates were removed, 32,785 articles were screened at title and abstract stage. From a full-text screen of 4,578 articles, we identified 789 articles that were eligible for inclusion in this report. Of these articles, 361 met only our inclusion criteria for content and did not have group comparisons, hypothesis testing, or appropriate analysis. These are listed in the bibliography of the report. Across the seven key questions, we synthesized the information from 428 articles.

KQ1. Effectiveness

All outcomes. KQ1: Effectiveness contains 379 studies assessing changes in process, intermediate outcomes, clinical outcomes, and economic and cost outcomes. The majority of

studies were observational, with a fair number of RCTs for prescribing and monitoring phases (Table A). Fifty-three qualitative studies are included in this total. Prescribing and monitoring were the most frequently studied phases of medication management (Table A), with hospital and ambulatory care settings well-represented to the near exclusion of long-term care, home, and community (Table B).

Though dealing with prescriptions and medications, pharmacists were poorly represented in studies, most focused on physicians (Table C). CDSS and CPOE systems were the most often studied MMIT technologies (Table D).

Table A. Research design for studies across the phases of medication management and education and reconciliation

| Design | P | OC | D | A | M | E | R |
|---------------|-----|----|----|----|----|---|---|
| RCT | 69 | 1 | 2 | 2 | 37 | 1 | 1 |
| Cohort | 13 | 2 | 2 | 1 | 6 | 0 | 1 |
| Observational | 144 | 18 | 10 | 26 | 29 | 2 | 4 |
| Qualitative | 37 | 5 | 3 | 10 | 5 | 0 | 0 |
| Total | 263 | 26 | 17 | 39 | 77 | 3 | 6 |

Note: some studies cross more than one phase.

Column headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Abbreviations: RCT = randomized controlled trial

Table B. Settings for the phases of medication management and reconciliation and education

| Setting | P | OC | D | A | M | E | R |
|---|-----|----|----|----|----|---|---|
| Ambulatory care (e.g., clinic, doctors office) | 94 | 6 | 2 | 1 | 40 | 2 | 0 |
| Community (e.g., school, community center) | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| Home | 2 | 0 | 0 | 2 | 5 | 0 | 0 |
| Hospital | 164 | 12 | 9 | 34 | 36 | 1 | 6 |
| Long-term care | 4 | 0 | 2 | 3 | 1 | 0 | 0 |
| Pharmacy | 11 | 13 | 10 | 2 | 4 | 0 | 1 |

Note: some studies cross more than one phase or setting.

Column headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Table C. Clinicians evaluated in outcomes studies of medication management phases, education, and reconciliation

| Clinicians | P | OC | D | A | M | E | R |
|---|----|----|---|----|---|---|---|
| Primary care physicians | 25 | 2 | 1 | 0 | 3 | 1 | 0 |
| Specialists | 11 | 0 | 0 | 0 | 1 | 0 | 0 |
| Hospitalists | 18 | 0 | 0 | 0 | 2 | 0 | 0 |
| Other physicians | 7 | 0 | 0 | 0 | 2 | 0 | 0 |
| Physicians undifferentiated | 26 | 1 | 0 | 1 | 3 | 0 | 1 |
| Nurses | 20 | 1 | 0 | 16 | 2 | 0 | 1 |
| Midlevel practitioners (e.g., PA, NP, MW) | 6 | 0 | 0 | 0 | 2 | 0 | 0 |
| Pharmacists | 13 | 6 | 5 | 2 | 1 | 0 | 1 |
| Other health professionals | 10 | 0 | 2 | 4 | 1 | 0 | 0 |
| Hospital administrators | 4 | 1 | 0 | 1 | 0 | 0 | 0 |

Note: some studies cross more than one phase and clinician type.

Column headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Abbreviations: MW = midwife, NP = nurse practitioner, PA = physician assistant

Table D. Main health IT studied by medication management phase and education and reconciliation

| Health IT | P | OC | D | A | M | E | R |
|--|-----|----|---|----|----|---|---|
| CDSS/reminders | 177 | 8 | 4 | 5 | 63 | 1 | 1 |
| CPOE/POE system | 90 | 12 | 5 | 9 | 11 | 0 | 0 |
| e-Prescribing | 31 | 10 | 3 | 4 | 2 | 1 | 1 |
| Order transmission of the prescription to and from doctor to pharmacy electronically | 2 | 3 | 0 | 0 | 0 | 0 | 0 |
| Pharmacy information system | 2 | 3 | 4 | 1 | 0 | 1 | 1 |
| Barcoding medication administering | 1 | 0 | 2 | 20 | 0 | 0 | 0 |
| Barcoding dispensing | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| eMAR, e-TAR | 2 | 2 | 2 | 14 | 0 | 0 | 0 |
| Other | 13 | 2 | 3 | 7 | 14 | 1 | 5 |
| Personal digital assistants or hand-helds | 7 | 1 | 0 | 4 | 5 | 0 | 1 |

Note: some studies cross more than one phase and technology.

Column headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Abbreviations: CDSS = Clinical decision support system, CPOE = Computerized provider order entry, POE = Provider order entry, eMAR = Electronic Medication Administration Record system, eTAR = Electronic Treatment Administration Record system

The results from this section suggest that care processes such as medication errors, time for tasks, workflow and knowledge, skills, and attitudes can be improved with the use of MMIT. The evidence is strongest specifically during the prescribing and monitoring phases. Few studies evaluated clinical outcomes associated with the use of MMIT. Those that did often did not show statistically significant improvements in clinical outcomes. Most of the studies with statistically

significant differences in clinical outcomes found small differences. The small number of articles with data on clinical outcomes is probably due, at least in part, to the difficulty in evaluating and establishing a direct association between the use of MMIT and clinical outcomes. This difficulty arises because of the distant nature of the outcome compared with the application of the health IT. Other contributing factors could also be considered.

Much of the relatively new research is addressing the type of research needed to come to a realistic and useful assessment of MMIT: pilot and demonstration projects and quantitative studies. Limited evidence suggests that MMIT can likely be cost effective, although most of the economic data come from cost analyses, which were often incomplete and seldom from head-to-head cost-effectiveness, cost utility or cost-benefit trials.

A substantial body of qualitative literature indicates support for the use of health IT in the various phases of medication management by a number of health care providers and patient groups. Survey studies of satisfaction and use reflect similar findings of acceptance and satisfaction, although most indicated room for improvement. Issues relating to changing care practices and workflow are frequently mentioned. The studies also provide useful summaries of unintended consequences of MMIT applications, which are discussed in detail in the full report.

Process changes. Most of the studies evaluating MMIT applications provided data on changes in process (225 of 378). Distribution in the number of studies across the five phases, plus reconciliation and education, was not equal. Prescribing was studied in 174 studies, order communication in 16 studies, dispensing in 9 studies, administering in 19 studies, and monitoring in 47 studies. Four studies evaluated reconciliation and one studied patient education. Studies often evaluated more than one phase.

Prescribing. The prescribing phase is well studied (174 studies), especially in hospital (61 percent of studies) and ambulatory care settings (39 percent). Long-term care centers (one study) and community and home settings (no studies) are not well studied. Physicians are by far the most studied group of health professionals. More studies are needed that evaluate nonphysician use of MMIT, specifically pharmacists, mental health professionals, nurses, and other nonphysician prescribers, as well as patients and their caregivers. Many of the studies of health care providers who were not physicians were purely descriptive of the people involved with them, and the systems themselves.

Based on the studies of process changes, CDSS and CPOE systems can play an important role in making prescribing and ordering more accurate, improving record keeping, and speeding up and improving communication. Both systems, either alone or, more often, integrated, are well studied (multiple studies with strong methods). Other MMIT applications lack evidence, especially those that involve nurses, pharmacists, and patients and their families.

MMIT in prescribing is associated with improvements in patient safety-related processes of the prescribing process, especially in hospital-based studies (87 percent, 52 of 60 studies), and somewhat less in ambulatory-based studies (68 percent, 28 of 41 studies). Errors related to prescribing and ordering were reduced in hospital-based studies (68 percent, 15 of 22 studies), but prescribing errors were not studied as often in ambulatory settings (two of two studies were positive). Reductions in time were related to the time taken to order or prescribe or the speed of the prescribing-to-administering processes. Most reductions in time were not seen as often in hospital-based studies (four of seven studies positive), but were positive more often in ambulatory settings (four of five studies). Adherence to treatment guidelines, reminders, and

recommended practice was improved in hospital studies (83 percent, 19 of 23) and to a lesser but still significant extent in ambulatory studies (64 percent, nine of 14 studies). Workflow was not evaluated in these studies of changes in process, although issues of workflow are addressed in qualitative studies in other sections of this report.

Order communication. Order communication, like dispensing, is one of the two medication management phases with the least number of studies—only 16 were identified. Two-way EDI holds promise to increasing the effectiveness of perfecting the prescription/order interactions between clinician prescribers and pharmacists. Currently, evidence on one-way communication predominates. The changes in process were also varied (two studies of errors, two of prescribing changes, five on time considerations, and three on workflow). Most studies were done using quantitative observational methods and all showed positive results.

Dispensing. Nine studies (three RCTs) assessed process improvements in dispensing. All process changes that were evaluated were found to be positive: four on modifications of the drugs that the pharmacists dispensed, three on errors, two on workflow, and one on adherence to good practice. With these few studies and multiple outcomes, evidence is limited on the role of MMIT in improving dispensing. This supports the findings of a Canadian health technology assessment report on MMIT that evaluated hospital dispensing and administering medications in hospitals.¹¹

Administering. Many articles dealing with administering medications were not included in this report because they were descriptive and did not include comparative data. Nineteen studies, 1 RCT, 1 cohort study, and 17 quantitative observational studies, were included. All studies were set in hospitals and included nurses. The MMIT systems were well integrated into multiple hospital IT systems. Error-reduction goals were common in the studies and almost always found to be improved (8 of 13 studies of errors). Errors were mixed, as some related to transcription and some to timing of administration, while some identified more serious errors. Four studies showed no improvement in errors while one study showed increases in errors, mostly related to timing of administration.¹² Four of five studies showed reductions in time from ordering to administering medication. Two studies evaluated the allocation of nursing time: one showed change and one did not in the proportion of time spent on various nursing tasks, including direct patient care, with the introduction of integrated MMIT for medication administering.

Monitoring. In our analysis, 70 percent (33 of 47 studies) of the included studies were associated with a 50 percent improvement in half or more process measures. Of these studies, most targeted physicians exclusively (34 studies), were conducted in academic institutions (33 studies), were developed for use in the ambulatory care setting (28 studies), focused on the adult population (36 studies), and provided CDSS with alerts or reminders to support chronic disease management (12 studies).

Studies that involved laboratory-based medication monitoring were most likely (76 percent of the time) to be associated with a greater than 50 percent improvement in a process outcome(s) than sign- or symptom-based medication monitoring. The most successful types of studies focused on changing prescriber behavior, improving response time to generated alerts, and improving the diagnosis and management of chronic diseases.

Reconciliation. Two systematic reviews and four studies provided evidence for improved reconciliation of medications with health IT. Reconciliation is the matching of medication lists over time, from different health care systems or from different prescribers. The evidence on reconciliation of medication lists is sparse, especially for systems that are fully integrated and capable of providing electronic comparisons of historical and current medications for individual patients at hospital discharge or on transfer to other facilities. All four studies showed improvements in agreement among lists of medications and two extended the evaluation to show improved prescribing¹³ and reduced errors.¹⁴

Unintended consequences. Eighteen studies provided data on adverse effects or unintended consequences. Two qualitative studies identified classes or categories of unintended consequences of health IT, many of which apply to MMIT applications. Some unintended consequences are minor, and some are major. In addition, some are seen to be positive and helpful. Some consequences are serious. For example, a small but statistically significant increase in mortality was seen in a children's hospital that installed a CPOE system that did not match workflow needs.¹⁵ A similar study showed another children's hospital that did not see the same increase in mortality in admitted children after their careful planning and implementation of health IT.¹⁶ Several authors contend that all health IT has unintended consequences. Formal evaluations of health IT installations should seek these unintended consequences and report them in their publications related to the evaluation. The importance of unintended consequences of MMIT also depends on the severity of the event, the degree of invasiveness of the MMIT, and the extent to which the use of the MMIT system disrupts existing workflow and processes. Consideration of formal reporting of serious unintended consequences might benefit all involved in development and implementation of MMIT systems. The qualitative studies in this report supplied a richer understanding of the adverse effects of MMIT, and they can form a strong base for more qualitative and quantitative studies of unintended consequences.

Education. Education related to MMIT centers on three aspects: formal informatics training during professional education or after graduation, training to use the MMIT systems, and improved outcomes based on knowledge and skills because of the use of the MMIT systems for health care providers, patients, and their families. This report does not include preprofessional or professional education related to the use and understanding of MMIT systems or certification in informatics or eHealth, all important aspects of MMIT application development and integration. Although we sought articles assessing postprofessional education related to changes in process associated with MMIT systems, we did not identify any articles that met our criteria. Training in the use of systems was often mentioned in articles but was not evaluated. Only one article was related to the educational component of MMIT systems for patient and family use, and it was associated with improved clinical outcomes. More information on health care professional and patient education is included in the sections of this report dealing with intermediate outcomes.

Intermediate outcomes. Intermediate outcomes deal with use, usability, education, knowledge, skills, and attitudes. Most studies with intermediate main endpoints focused on measuring use, correlates of use, perceptions, and satisfaction in the prescribing phase (26 of 42 studies). As for changes in process, clinicians and prescribing were well-studied. Use, perceptions, and satisfaction were reported to be improved. Factors such as ease of use, perceived usefulness, and improved quality of care predominated. Satisfaction and attitudes varied depending on the role of

the health care provider. Variation in needs and roles of health professionals with respect to use of health IT are real and should be considered when choosing or implementing any new IT system. Usability studies with comparison groups are sparse but can provide useful suggestions to improve systems. Usability studies are often difficult to generalize or transfer across settings, in part because MMIT effectiveness is linked strongly to the culture, institutional leadership, and other situation specific factors. Therefore, applicability of findings related to usability is problematic in MMIT applications.

Economic outcomes. Five of 31 articles dealing with costs conducted comprehensive economic evaluations (costs and consequences). Two evaluated a CPOE system and three evaluated CDSS. Most of the studies that included monetary data (22 of 31 studies) were partial economic evaluations in the form of cost analyses (assessing costs of alternatives without analysis of effectiveness or efficacy). Most of these partial economic analyses assessed costs of prescribed medications with the MMIT system compared with not having the MMIT system.

Several studies found that health IT interventions may offer cost advantages despite their increased acquisition costs. These studies showed that over time, a net benefit accrued based on cost reductions resulting from the MMIT (such as lower adverse drug events (ADEs), drug costs, and laboratory test usage). However, given the uncertainty that surrounds the cost and outcomes data, and limited study designs available in the literature, it is difficult to reach any definitive conclusion as to whether the additional costs and benefits represent value for money.

Clinical outcomes. A total of 76 studies sought to measure improvement in clinical outcomes or reduction in ADEs, of which 26 (34 percent) reported significant benefits of health IT. One reported harm—a small but clinically important increase in mortality when an inflexible CPOE was implemented in a children’s hospital.¹⁵ Because of the seriousness of the implications of this study, many people reviewed this article and its methods.¹⁷ A later and similar study showed that with careful planning another children’s hospital did not see the same increase in mortality in admitted children after the implementation of a health IT.¹⁶

An additional two studies implemented CDSSs to reduce costs and assessed whether reductions in drug use increased mortality¹⁵ and length of stay.¹⁸ Both studies lacked sufficient power to conduct a valid assessment.

Studies that used laboratory-, sign- and symptom-based monitoring approaches were mostly clinician based. If the MMIT monitoring was used to identify and intervene with patients with actual problems (e.g., excess blood pressure) or needed care (e.g., hemoglobin A1c monitoring), this appears to be more effective than CDSS approaches that identified theoretical problems (potential for ADEs), particularly if patients are also sent reminders and decision support recommendations.

Highly targeted interventions, which focused on specific problems that provide problem-related specific interventions, appear to be more effective than more diffusely focused systems such as CDSS and CPOE. Some of these highly targeted interventions involved CDSS tools for improving the effectiveness of anticoagulants (proportion of days with blood clotting parameters within the therapeutic range), improving the choice, route, and duration of antibiotics, and reducing ADEs related to antibiotic use, and most were successful.

Studies that have been successful in improving patient outcomes target high risk and vulnerable populations who have poor disease control, lack sufficient access to health care providers to manage their condition or subpopulations with sufficient economic resources to

respond to the CDSS intervention. The effect of similar CPOE systems on mortality can vary substantially as a function of the extent to which implementation strategies disrupt or delay critical activities in the clinical setting and demand additional time for order entry from clinical staff. Critically ill patients (i.e., those who are most vulnerable) are most likely to be affected by dysfunctional technology and implementation strategies.

Qualitative studies. Qualitative studies seek to understand phenomena and answer questions of why and how as well as to gain insights into real life situations. They often study the more human or “soft” side of health and health care. The preceding sections concentrated on studies with quantitative outcomes. Fifty-three qualitative studies are included in this section. Patient safety was the main health aspect evaluated in qualitative studies. Before MMIT implementation most studies found that clinicians expected that MMIT would improve patient safety and once implemented most clinicians felt that MMIT had improved safety.

The qualitative studies focused on system design including workflow changes, challenges with the system interface, and new communication processes—all of which can generate new kinds of medical errors, which in some cases were detrimental to patient safety.

Early implementers associated MMIT with a lot of self-reported “hard work” by those who were expected to use the new systems. These people, most often health professionals, struggled, often independently, with limited guidance with respect to planning and implementation tactics during preparation for and implementation of the MMIT applications. During planning and early implementation, the users often experienced unanticipated effects. Frequently, the initial stage was disruptive and, consequently, clinicians found provision of care to be more challenging with the MMIT system than without. However, after the initial stage was over, the attitudes of the care providers changed, and the potential benefits of the system become clearer to most. Of special note is that the implementation of MMIT systems generated emotional responses in a broad range of health professionals, both positive and negative. For example, strong feelings were associated with reminders and alerts¹⁹ and CPOE.²⁰

MMIT implementation did not just mean that a clinician needed to learn a new IT system, but the implementation also affected most of the other parts of the delivery of care processes, including how the interdisciplinary care team worked together.

KQ2. Knowledge and Evidence Gaps

We identified gaps in the report, some that we expected and some that we did not. We address the question of knowledge deficits across phases and outcomes, settings and participants, grouping similar gaps together.

Phases of medication management. Because of the preponderance of publications on the prescribing and monitoring phases, they are less in need of more study than the other phases of order communication, dispensing and administering, and medication reconciliation. In addition, the educational or training requirements for effective use of MMIT applications by health professionals need to be studied as well as education related to patients as new MMIT applications are developed for their use.

Research methods. MMIT applications are complex interventions and need to be studied in pragmatic (i.e., does it work in real settings?) evaluation projects and using complex interventions methods. The applications also should ideally be studied by teams of researchers

with, or teams that seek consultation from, those who have experience in clinical practice, research methods, statistical analysis, and informatics training and experience. Qualitative studies are also vital to understand the complex nature of how systems are used and valued, especially across groups of health professionals who often have different needs and expectations.

Health care providers. Physicians are well studied. Nurses, midlevel practitioners (nurse practitioners, physician assistants, midwives), pharmacists, other prescribers such as dentists and mental health practitioners, and hospital administrators need studies directed at their needs, practice patterns, and health IT tools.

Patients. Many studies included data related to patients, usually in the measurement and reporting of process changes and other outcomes. Few studies, however, concentrated on how the MMIT systems directly affected patients and clinical outcomes important to them. Traditionally, MMIT systems were developed as clinician and administrator tools. Patient and family use of MMIT systems is becoming more important, and this gap in our understanding needs to be addressed.

Settings. Hospitals and ambulatory care settings are well studied. Gaps exist in our knowledge of the effectiveness of MMIT in long-term care facilities, the community, and homes. Long-term care facilities most need strong qualitative and quantitative studies because they rely heavily on medication. Homes, schools, and other community settings will also become more important with shifting care to more self-reliance in relation to wellness care and chronic disease management.

Health IT. Much research has gone into evaluating CDSS and CPOE systems, either alone or integrated. For example, 77 of 88 RCTs evaluated some aspect of CDSSs. Other MMIT applications, especially those that are used by nonphysicians or outside the prescribing and monitoring phases, lack evidence. Examples with little evidence on effectiveness are bar coding for administering and dispensing, pharmacy information systems, electronic medication administration record systems, and fully integrated comprehensive information systems.

Process changes. Patient safety processes such as error reductions and improvement in prescribing have a strong evidence base. Issues related to workflow, communication changes, and unintended consequences are understudied. More study of laboratory-based monitoring of medications, especially in facilities that have highly integrated information systems, is important. More qualitative and controlled studies are needed as well as multicenter studies and those that use methods developed by groups focusing on health technology assessment (HTA). These HTA methods include integrated reports that bring together research syntheses, modeling of processes and full economic reports, and cost studies. Often these HTA reports do not, but can, involve additional collection of evidence.

Intermediate outcomes. More study is needed on the importance of usability testing in all stages of development and use. This must be done with all users and not just segments of those involved in using MMIT. Usability studies have not traditionally been generalizable or transferrable but more limited to a specific setting. AHRQ might consider a research program in how to make these usability studies more applicable to multiple institutions, training in usability

methods, collection of usability tools and completed studies, and research into the need for standards of usability testing for new or modified systems. Usability studies must also include all users of systems. For example, systems that have been optimized only for physician users are usually systems that nurses and other health professionals have difficulty using. Workarounds have often been unofficially implemented by users instead of system modifications and improvements.

Clinical outcomes. Findings associated with improvement in clinical outcomes are still equivocal. These studies are difficult to do well, expensive, and time consuming, but they must be done. Multicentered trials planned by strong teams of experienced people from multiple backgrounds are vital.

Cost and economic outcomes. Although many studies exist that list costs and outcomes, few comprehensive and definitive studies of the economic value of MMIT applications exist. Both the potential for improvement and the costs of implementing and maintaining these systems are huge. Again, well-planned studies with broad input from many stakeholders are necessary for understanding the true worth of MMIT applications. HTA or other studies that integrate costs and consequences of MMIT systems would be ideal.

Qualitative. Qualitative studies have provided much valuable information about MMIT. Gaps in qualitative knowledge center on the lack of qualitative studies that address the effects of MMIT on health outcomes. In addition, very few qualitative studies examined the effects of MMIT from the perspective of the patient.

KQ3. Value Proposition for Implementers and Users

Value proposition is determined from a balance of financial, clinical, and organizational benefits. A clear assessment of each of these from the viewpoint of each stakeholder is needed to make a clear value judgment. For each stakeholder—and many are involved with MMIT implementation—the relative importance of these three elements is different. Values will also vary depending on the setting and the type of technology employed. Multiple stakeholders, some of whom may be distant from the MMIT, need to be considered in any value proposition study. Based on the evidence in KQ1: Effectiveness, knowledge about the three elements needed to make value judgments is slowly accumulating. We cite only 31 papers in this section, although some of our assessments come from sections of this report that have included more studies. Gains in productivity and process of care outcomes have been shown, but good evidence of improvement in patient outcomes with MMIT is weak or lacking. The body of economic literature is still sparse and lacks vigorous study. We found little theoretical work or actual studies that were done to determine what each stakeholder takes into account to reach value proposition judgments related to MMIT.

KQ4. System Characteristics

Few studies (n = 21) demonstrated evidence of the impact of the characteristics of MMIT applications on the likelihood to purchase, implement, and use such IT applications. No studies assessed open-source health IT applications, with only one study each on conformity with standards and CCHIT-certified systems. Twenty of the articles related to the prescribing and ordering phase. Almost all of the articles suggest that feature sets of health IT applications have

been instrumental in reaching decisions to adopt MMIT applications. Certain features of systems improve the likelihood of purchase, implementation, and use of MMIT. The literature, however, is sparse and observational in nature. Most often authors described barriers and concerns toward implementation and acceptance rather than characteristics of MMIT that could facilitate implementation, purchase, and use of such systems. Authors seldom provided enough details about the technology to form conclusions about the value of feature sets and system characteristics. Head-to-head comparisons of systems differing in their features were not found.

KQ5. Sustainability

Our literature review revealed three important findings: sustainability is frequently mentioned in the core biomedical informatics literature, it is poorly defined, and none of the articles included in this evidence report explicitly studied sustainability. These findings are not entirely surprising. A previous AHRQ-sponsored evidence report that assessed the costs and benefits of health IT in pediatrics found only one article that explicitly discussed sustainability.²¹

Future research would be beneficial for many if a study or group would develop an operational definition of sustainability that could be used to study its determinants. Moreover, it is likely that the Health Information Technology for Economic and Clinical Health (HITECH) Act of 2009 will lead to improvements and sustainability of health IT applications that specifically support the medication management continuum through meaningful use.

We have summarized a body of literature that uses surveys to detect patterns in the characteristics of people and organizations that are more likely to implement various technologies. These surveys are often the basis for further study into barriers and facilitators to increasing uptake and adoption.

Integration of MMIT with other systems was an inclusion criterion for our report (except for PDAs that analyzed patient-specific data). Some technologies were integrated with a greater number of components than others. Frequently, the descriptions of the systems were inadequate to fully determine how the systems were connected. Access to various other information sources, most notably laboratory reports, enhanced the performance and acceptance of the MMIT applications.

KQ6. Complete Two-Way Electronic Data Interchange

No reports documenting the use of complete two-way EDI systems were found. Evidence from the limited set of one-way, e-Prescribing studies was extrapolated to identify possible key facilitators and barriers to completely electronic, two-way, e-Prescribing systems. Possible facilitators include monetary or other incentives to providers, a permissive regulatory environment, and the existence of an established standard for prescription EDI. Barriers included the low rate of EMR adoption in the United States, regulatory and legal uncertainties, and inadequate consideration of the effects of e-Prescriptions on pharmacists and pharmacies and their processes. While answering this question, we found that the Bell model does not represent the two-way communication between pharmacists and prescribers—it shows only a one-way linear movement of information.

KQ7. Effectiveness of CDSS

Seventy-seven RCTs were designated as primarily studying CDSS related to medication management and integrated with other health IT. These studies involved 4,709 providers and 828,441 patients in total. All studies assisted with at least the prescribing or monitoring phases of

medication management. Overall, we found a lack of RCTs addressing electronic decision support integrated with other types of health IT. Statistically significant process changes were often shown in these RCTs. Only a small minority of these focus on clinical outcomes, however. Studies with clinical outcomes are those that are most important to guide decisionmaking of patients' providers and policymakers about the usefulness and need for MMIT interventions. A very small number of studies reported improvement in clinical outcomes.

Discussion

The literature of MMIT presents challenges. It is diffused across multiple disciplines, and much of it is descriptive in nature. We also found that although studies with strong methods exist, they are not uniformly dispersed across phases of medication management, people, settings, or health IT applications.

The literature would be stronger if standardized definitions of issues like medication errors, adverse effects, MMIT applications, and sustainability were implemented. The evidence of effectiveness can be made stronger with directed evaluation funding. With direction the evaluations could be encouragement for studies to be done appropriately and not just on small budgets or by the system developers. Training in research skills as part of informatics training may also enhance the evidence on the effectiveness of MMIT. We noted problems in study methods and often found studies that lacked sufficient numbers for valid statistical analyses and assessment of implications.

Despite the challenges in the evidentiary base for MMIT, it is a vital, vibrant, and a proven component of health and health informatics—at least for improving the processes of care that include patient safety. Qualitative studies have provided data on expectations, hopes, changes in how care is delivered, and the need for deep understanding of the effects of MMIT applications in planning for and implementing them. We are much wiser for bringing this literature together into one resource. Moving forward and with the advent of new systems, greater emphasis on eHealth to improve health care and health care delivery, and the move to more patient-centered care, it is an exciting time for development and integration of MMIT applications.

References

1. Bell DS, Cretin S, Marken RS, et al. A conceptual framework for evaluating outpatient electronic prescribing systems based on their functional capabilities. *J Am Med Inform Assoc* 2004;11(1):60–70.
2. Shekelle PG, Morton SC, Keeler EB. Costs and Benefits of Health Information Technology. Evidence Report/Technology Assessment No. 132 (Prepared by the Southern California Evidence-based Practice Center under Contract No. 290-02-0003). Rockville, MD: Agency for Healthcare Research and Quality, April 2006. AHRQ Publication No.06-E006.
3. Jimison H, Gorman P, Woods S, et al. Barriers and Drivers of Health Information Technology Use for the Elderly, Chronically Ill, and Underserved. Evidence Report/Technology Assessment No. 175 (Prepared by the Oregon Evidence-based Practice Center under Contract No. 290-02-0024). Rockville, MD: Agency for Healthcare Research and Quality, November 2008. AHRQ Publication No. 09-E004. Available at: <http://www.ahrq.gov/clinic/tp/hitbartp.htm>

4. Gibbons MC, Wilson RF, Samal L, et al. Impact of Consumer Health Informatics Applications. Evidence Report/Technology Assessment No. 188 (Prepared by Johns Hopkins University Evidence-based Practice Center under contract No. HHS 290-2007-10061-I). Rockville, MD: Agency for Healthcare Research and Quality, October 2009. AHRQ Publication No. 09(10)-E019. Available at: <http://www.ahrq.gov/downloads/pub/evidence/pdf/chiapp/impactchia.pdf>
5. Hersh WR, Hickam DH, Severance SM, et al. Telemedicine for the Medicare Population: Update. Evidence Report/Technology Assessment No. 131 (Prepared by the Oregon Evidence-based Practice Center under Contract No. 290-02-0024). Rockville, MD: Agency for Healthcare Research and Quality, February 2006. AHRQ Publication No. 06-E007.
6. Agency for Healthcare Research and Quality. Enabling Health Care Decision Making through the Use of Health Information Technology (Health IT). Systematic Review Protocol. Rockville, MD: Agency for Healthcare Research and Quality, March 2010. Available at: <http://www.ahrq.gov/clinic/tp/knownmgttp.htm>.
7. Agency for Healthcare Research and Quality. Enabling Patient-Centered Care through Health Information Technology (Health IT). Systematic Review Protocol. Rockville, MD: Agency for Healthcare Research and Quality, March 2010. Available at: <http://www.ahrq.gov/clinic/tp/pcchhttp.htm>.
8. Lohbach D. Enabling Health Care Decision Making through the Use of Health Information Technology (Health IT). Expected release date December 2010. Rockville, MD: Agency for Healthcare Research and Quality, March 2010. Available at: <http://www.ahrq.gov/clinic/tp/knownmgttp.htm>
9. Humphreys JS, Wakerman J, Wells R. What do we mean by sustainable rural health services? Implications for rural health research. *Aust J Rural Health* 2006;14(1):33-5.
10. Verhagen A, de Vet H, de Bi R, et al. The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *J Clin Epidemiol* 1998;51(12):1235-41.
11. Perras C, Jacobs P, Boucher M, et al. Technologies to reduce errors in dispensing and administration of medication in hospitals: Clinical and economic analyses. Technology Report Number 121. Ottawa: Canadian Agency for Drugs and Technologies in Health; 2009.
12. Morriss F, Abramowitz P, Nelson S, et al. Effectiveness of a barcode medication administration system in reducing preventable adverse drug events in a neonatal intensive care unit: a prospective cohort study. *J Pediatr* 2009;154(3):363-8.
13. Grasso BC, Genest R, Yung K, et al. Reducing errors in discharge medication lists by using personal digital assistants. *Psychiatr Serv* 2002;53(10):1325-6.
14. Poole D, Chainakul J, Pearson M, et al. JHQ 177 medication reconciliation: a necessity in promoting a safe hospital discharge. *J Healthc Qual* 2006;28(3):12-19.
15. Han YY, Carcillo JA, Venkataraman ST, et al. Unexpected increased mortality after implementation of a commercially sold computerized physician order entry system. *Pediatrics* 2005;116(6):1506-12.
16. Keene A, Ashton L, Shure D, et al. Mortality before and after initiation of a computerized physician order entry system in a critically ill pediatric population. *Pediatr Crit Care Med* 2007;8(3):268-71.
17. Gesteland PH, Nebeker JR, Gardner RM. These are the technologies that try men's souls: common-sense health information technology. *Pediatrics* 2006;117(1):216-17.
18. Lesprit P, Duong T, Girou E, et al. Impact of a computer-generated alert system prompting review of antibiotic use in hospitals. *J Antimicrob Chemother* 2009;63(5):1058-63.
19. Van Ast JF, Talmon JL, Renier WO, et al. Development of diagnostic reference frames for seizures. Part 2: are seizure descriptions discriminative? *Int J Med Inf* 2003;70(2-3):293-300.

20. Weir C, Lincoln M, Roscoe D, et al. Dimensions associated with successful implementation of a hospital based integrated order entry system. Proc Annu Symp Comput Appl Med Care 1994:653–7.
21. Goldzweig CL, Towfigh A, Maglione M, et al. Costs and benefits of health information technology: new trends from the literature: since 2005, patient-focused applications have proliferated, but data on their costs and benefits remains sparse. Health Aff (Millwood) 2009 Mar–Apr;28(2): Supplement 1: w282–293:Supplement-93.

Introduction

Scope and Purpose of the Systematic Review


The Agency for Healthcare Research and Quality (AHRQ) has considerable interest in health information technology (health IT). They have contracted several reports that are published or will soon be published. These evidence summaries on health information technology (IT) are related to costs and benefits;² barriers and drivers of health IT for the elderly, chronically ill and underserved;³ the impact of consumer informatics applications;⁴ and telemedicine.⁵ Currently AHRQ has also contracted for the use of health IT in decisionmaking,⁶ in patient centered care⁷ and decision support for health care decisionmaking.⁸ These latter reports will be available early 2011. AHRQ asked the McMaster University Evidence-based Practice Center to generate an evidence report outlining the impact of health IT on the medication management process. Medication management is a major component of the health care system. Currently, approximately 10 percent of the health care budget in the United States is spent on prescription medications.²²

To structure this evidence report we use the framework of medication management as presented by Bell and colleagues.¹ They model the medication management continuum into the five phases of this evidence report; Figure 1 is a pictorial representation of the medication management phases.¹

The first phase of the continuum is **prescribing** medications by clinicians who have assessed the patients' conditions and needs. The second phase is to transmit the prescription to the pharmacists who work with the prescriber to clarify and verify the order (referred to as '**order communication**' in this report to capture the complexity of the communication that occurs between prescriber and pharmacy). The next step is **dispensing** the medication in its required form and dose, followed by **administering** the medications to the patient. **Monitoring** is the final phase where ongoing oversight occurs to address the changing medication needs and situation of the individual. **Reconciliation** of the medications taken by individuals and postprofessional **education** or training related to medication management IT (MMIT) are additional aspects (as opposed to phases) of the medication management cycle covered by this report. Reconciliation is a process whereby a patient has their medication lists verified for completeness and accuracy when the patient moves from hospital to home or to a nursing home, or is involved with multiple care providers. Reconciliation can improve care by using health IT to ensure accuracy of medication lists, identification of gaps and conflicts in prescription, and provide timely and efficient transfer of patients and their medication data. Education is also important in MMIT systems for both improving knowledge and skills of patients and care providers and to enable timely implementation and optimal use of MMIT systems. Therefore, AHRQ requested inclusion of both reconciliation and education in this report in addition to the five medication management phases.

This report includes clinicians, patients, informal caregivers, and administrators. All care settings are also covered: home, community, primary care and specialty clinics, all levels of hospitals, long-term care facilities, and pharmacies of all types. This report does not focus on the health insurance or pharmacy industries.

Figure 1. A functional model of the medication management continuum created by Bell et al.¹



Includes the major activities involved in medication management and forming the basis of our medication management phases. We refer to the transmission of the order/prescription and the bi-directional communication between prescriber and pharmacy staff as “order communication.” Used with permission BMJ Publishing Group.

Health IT holds great potential to improve the quality of health care and reduce potential and real errors while at the same time providing cost effective care. The coverage of this report is broad, reflecting the scope and breadth of health IT and the processes of medication management. This report centers on health IT applications that focus on medication management such as e-Prescribing applications, computerized provider order entry (CPOE), bar-coded medication administration (BCMA), pharmacy-based health IT, electronic medication administration record systems (eMAR), and other MMIT tools. Smaller health IT applications such as hand-held devices that provide calculations for dosing, as well as MMIT applications integrated with other health IT systems, such as electronic health or medical records systems (EHRs and EMRs), health information systems, hospital information systems, and personal health records (PHRs), and others as identified, are included. For inclusion, the MMIT had to be integrated into the health care system electronically and process patient-specific information that then provided direction for that patient’s care. This integration requirement meant that stand-alone devices such as smart infusion pumps and glucose monitors were not included unless they were integrated with other health IT. This requirement allowed the report to concentrate on MMIT systems and separate out these medical devices with some computing capabilities.

Because health IT is a new discipline that crosses many domains, definitions are not always standard. Therefore we have defined terms related to health IT and other issues in this report in a glossary, labeled as Appendix F.

Key Questions (KQs)

KQ1. Effectiveness. Within and across the phases of the medication management continuum (prescribing, order communication, dispensing, administering, and monitoring, plus reconciliation and education aspects) what evidence exists that health IT applications are effective in improving:

- a. Health care process changes (e.g., adherence to guidelines, changes in prescribing behavior, changes in patient monitoring activities, errors, efficiency),
- b. Other intermediate outcomes (e.g., use, measures correlated with use, satisfaction with system, usability, knowledge, skills, and attitudes),
- c. Costs and economic outcomes,

- d. Clinical outcomes for patients (e.g., physiological measures, adverse drug events, length of stay, mortality, quality of life, other patient events),
- e. Population level outcomes, and
- f. Composite outcomes.
- g. To what extent does the impact of health IT on improvement of the health care process, other outcomes, costs and economics, and clinical outcomes vary depending on the type of medication (e.g., controlled or noncontrolled substance) or the form of the medication (e.g., oral, injection, intravenous)?

KQ2. Gaps in Knowledge or Evidence. What knowledge or evidence deficits exist regarding needed information to support estimates of cost, benefit, impact, and net value with regard to enabling health IT applications in terms of prescribing, order transmission, dispensing, administering and monitoring, and adherence? Discuss gaps in research, including specific areas that should be addressed, and suggest possible public and private organizational types to perform the research, analysis, or both.

KQ3. Value Proposition. What critical information regarding the impact of health IT applications implemented to support the phases of medication management is needed to give clinicians (e.g., physicians, nurses, psychologists, dentists, and pharmacists), health care facility administrators, patients, and their families a clear understanding of the value proposition particular to them?

KQ4. System Characteristics. What evidence supports or refutes the impact of any of: open source, home grown, proprietary, local configuration ability, system configuration ability, conformity with U.S. Federal or other interoperability standards, conformity with other standards from other jurisdictions, being Certification Commission for Healthcare Information Technology (CCHIT) certified, system architecture, or feature set on the decision to purchase, implement, or use health IT in medication management systems?

KQ5. Sustainability. What factors influence sustainability (use and periodic updates) of health IT applications that support a phase of medication management continuum: prescribing, order communication, dispensing, administering and monitoring, plus reconciliation and education?

- a. To what extent does the evidence demonstrate that health care settings (e.g., inpatient, ambulatory, long-term care) influence implementation, use, and effectiveness of such health IT applications?
- b. What is the impact (e.g., challenges, merits, costs, and benefits) of having electronic access to patients' computerized medication records (current and past), EHRs and PHRs, formulary information (inpatient and outpatient issues), billing information, laboratory records, and other electronic patient data in the quality and safety of care provided by health IT applications that support at least one phase of the continuum of medication management (i.e., prescribing and ordering, transmission and verification, dispensing, administering and monitoring and adherence)?

KQ6. Two-way EDI for Order Communication. It has been recognized that implementation and use of a complete, two-way electronic data interchange (EDI) (e-prescribing with e-transmission) between the prescribers' electronic medical records (EMRs), including CPOE and

other health IT within EMR, and other similar systems or stand-alone e-prescribing systems, retail, and mail-order pharmacy prescribing systems have been limited. In many instances, health IT systems that facilitate prescribing are used at the point-of-care and are combined with nonelectronic modalities for transmission of prescriptions, such as paper, facsimile, voice, and telephone. On the pharmacy side, prescriptions being received may not automatically populate the pharmacy prescribing system, instead appearing in the fax printer or in a different computer program than the one the pharmacist regularly uses to fill prescriptions, requiring the pharmacist to manually retype the prescription information into the pharmacy's electronic system. This e-Prescribing with e-transmission also includes order clarification with electronic communication between the prescribers and pharmacists.

- a. What evidences exists demonstrating the barriers and drivers of implementation of complete EDI that can support the prescription, transmittal, receipt, and order clarification process of e-Prescriptions?
- b. How do barriers, facilitators, and economic incentives vary across pharmacists, physicians, and other relevant stakeholders with respect to adoption and use of complete EDI (e-Prescribing/ordering with e-Transmission)?

KQ7. RCTS of CDSS. What evidence exists regarding the extent of integration of electronic clinical decision support systems (CDSS) in a health IT system for the prescribing, dispensing, and administering of medications? To what extent does the use of the CDSS for prescribing/ordering, order communication, and dispensing of medications impact the various outcomes of interest, including health care process, intermediate, costs and economics, and clinical endpoints? CDSSs are broadly defined to include medical and pharmacy systems, reminders, and monitoring.

Background

Medication management is a complex and expensive process with high potential for both benefit and harm. Ninety percent of American seniors and 58 percent of nonelderly adults rely on medications daily. The average cost of prescription drugs per clinic visit in the United States in 1996 was \$79. By 2006, this had doubled to \$161. Nationally, all prescription drugs costs are projected to be \$246.3 billion for 2010. Substantial increases in medication costs are expected until at least 2019 based on the aging population²³ and increased demand for medications (72 percent increase from 1997-2007).²⁴ The introduction of newer, high cost, nongeneric, and specialty drugs also adds to the projected increases. The amount of new more complex medications also places a substantial cognitive burden on health professionals who prescribe and oversee these medications. Genomics research and its role in medication choices for individualized health care are also going to become more important in the next decades. Health IT can play a strong information support role to help deal with this increased cognitive load and provide efficiencies for provision of prescription medications, control, and recording of use.

In addition to increasing costs, medications can cause substantial health problems. Incorrect choice of medications and over or under use leads to less than optimal care. The U.S. Institute of Medicine (IOM)²⁵ report on medication errors estimates that errors occur in all levels and locations of care. Estimates for hospitalized patients show 1.5 to 10 errors per 100 opportunities for errors for prescribing and 2.4 to 11 errors per 100 opportunities per dose for dispensing. This translates to approximately one error per patient per hospital day. Error rates in long-term care

prescribing are calculated to be from 6 to 20 errors per 100 opportunities per dose. Ambulatory care studies show that up to 21 percent of prescriptions have errors.

Pediatric patients present special challenges in that doses must often be adjusted for body weight and age. As an example, one study showed that errors in acetaminophen use in the emergency department for children were 22 per 100 doses ordered.²⁶ Elderly patients also have special prescribing and drug monitoring needs based on issues related to aging, multiple conditions, the need for several medications, and often, decreased kidney function.

Pharmacist errors in order communication and dispensing also occur. Cheung and colleagues²⁷ reviewed the literature of dispensing errors and found that overall errors occurred in the range of 0.2 to 0.8 percent, although the number varied depending on how the errors were detected and reported. The task of medication administering by nurses, other health care providers, as well as patients and families, have also been shown to have associated errors. Many of the errors in medication management described above are preventable. The IOM report shows that preventable errors often constitute 20 to 50 percent of all errors. In addition to mortality, errors and inappropriate use are costly, often cause a huge drain on health care resources, and contribute to substantial morbidity and challenges to well-being.

Historically, the first MMIT application was published in 1979 as a decision support system to help in prescribing appropriate antibiotics.²⁸ The first RCT was done 5 years later.²⁹ Health IT has tremendous potential to improve care associated with medication management. For example, the Center for Information Technology Leadership (CITL), in their Value of Computerized Provider Order Entry in Ambulatory Care³⁰ report that potential savings from implementation of CPOE in ambulatory care prescribing and its ability to detect errors would provide savings in the U.S. of \$28 billion annually. Other MMIT applications are projected to have similar cost savings and improved care. However, the promises of health IT have not always been obtained after installation. For example, Mollen and colleagues³¹ reviewed CDSSs for prescribing and found 37 reports that successfully showed changed health care provider behavior. Only five of these studies noted improvements in patient outcomes. Similarly, Eslami and colleagues³² reviewed studies of CPOE applications in outpatient medication ordering. Of 67 studies, only 21 dealt with safety. Most of the evidence they identified used observational study methods. They showed that although CPOE and other information systems are often costly, some evidence supports medication safety benefits. However, they also note that some studies have data that support increased error rates and adverse drug events (ADEs) with CPOE implementation. Kaushal and colleagues³³ show that e-Prescribing with CDSS reduced errors from 52.5 to 6.6 per 100 prescriptions in ambulatory care. Paoletti and colleagues³⁴ reduced errors from 2.9 percent to 1.6 percent in a U.S. general hospital with the implementation of BCMA and eMAR.

Many groups have studied various components of the medication management process and the effects of multiple health IT systems and programs across settings and populations. However, the body of evidence that evaluates the actual, and not projected, effect of a broad range of MMIT applications and the medication management process is not available in one document or Web site. This evidence report is designed to be that summary.

Methods

The objective of this report is to review and synthesize the available evidence regarding the effectiveness and effects of health IT on all phases of medication management, as well as reconciliation and education. The report considers a broad range of health ITs and medication management processes and concentrates on those people involved in direct clinical care: physicians, pharmacists, dentists, nurses, and other health professionals; patients and their informal caregivers; and health care administrators across all health care settings and levels of care.

Recruitment of Technical Experts and Peer Reviewers

The Medication Management through Health Information Technology (MMIT) team was made up of experts from McMaster University, the University of Pittsburgh, and McGill University. Expertise of the group included medical informatics, primary care, geriatrics, internal medicine, pharmacy, conduct of clinical trials, and systematic literature reviews. Our Technical Expert Panel (TEP) was comprised of 12 external experts from diverse professional backgrounds including medication safety, health information technology in medication management, consumer informatics, and pharmacy. Their clinical expertise included specialization in pharmacy, geriatrics, reproductive health, pediatrics, and primary care. The TEP was involved in the development of the project by helping to refine the questions, focus the scope, solidify and streamline definitions, and approve modified plans and project direction. The members of the TEP and the external reviewers are listed in Appendix E. We also sought advice from other AHRQ Evidence Based Practice Centers who had completed health IT evidence summaries.

Key Questions

The core team worked with the external advisors, the TEP, and representatives of the AHRQ to refine the key questions (KQ) presented in the “Scope and Purpose of the Systematic Review” section of Chapter 1. Before searching for the relevant literature, the content of the questions was clarified, the concepts were defined, and the types of evidence that would be included in the review were ascertained.

KQ1. Effectiveness addresses the evidence that health IT applications improve a broad range of outcomes when health IT is applied to medication management (five phases plus the impact of postprofessional and patient education and reconciliation among those phases). Studies that reported changes in process, cost and economics, intermediate, qualitative, and clinical patient outcomes are included.

Much literature addresses the use of health IT in medication management. To address the MMIT question using the best available research findings, two limitations were placed on the included articles. First, only hypothesis-driven articles were included. For quantitative articles this meant that those with comparison groups and appropriate statistical analysis were analyzed in this report. Qualitative studies were included if they reported use of recognized qualitative methods. Many other articles met our inclusion criteria for content and measured an outcome of interest but they were not hypothesis-driven; the report lists these citations in the KQ1: Effectiveness section of Chapter 3: Results.

KQ2. Gaps in Knowledge or Evidence addresses knowledge and evidence deficits regarding needed information to support estimation of costs, benefits, impact, and net value regarding MMIT applications.

KQ3. Value Proposition requires the identification of information about the MMIT applications needed for each stakeholder to have a clear understanding of the value proposition particular to them. It was determined that the answers to KQ2: Gaps and KQ3: Value Proposition would become evident from the review of the evidence in KQ1: Effectiveness, although studies addressing values propositions by stakeholders are also included.

KQ4. System Characteristics addresses the impact of MMIT application features on the likelihood that the systems will be purchased, implemented, and used. This evidence comes from studies measuring implementation, use, and purchasing decisions. Studies of all designs are included.

KQ5. Sustainability addresses the factors influencing the sustainability of MMIT applications, specifically: (a) the impact of the type of setting, and (b) the impact of access to other electronic data on health care quality and safety. Sustainability is not well-defined. The definition of sustainability provided by Humphreys et al.,⁹ “the ability of a health service to provide ongoing access to appropriate quality care in a cost effective and health-effective manner” was incorporated. This definition restricted the number of articles that were included in this review. The topic of sustainability is one that needs further research in defining and further analyses of existing systems.


KQ6. Two-way EDI relates to the barriers and facilitators to complete two-way electronic data interchange (EDI) between prescribers and pharmacists and how these factors vary across stakeholder groups. Through discussions with experts and the MMIT writing group we determined that the evidence would be sparse in this category. Any article studying EDI communication (one- and two-way) that includes original data (qualitative or quantitative) is included in the report.

KQ7. RCTs of CDSS addresses the extent to which clinical decision support is integrated into health IT systems for medication management and the impact of CDSS on process and health outcomes. Because of the size of the literature and the improved level of evaluation rigor and generalizability or applicability of RCTs, only RCTs are included. This question included changes in process as well as the broad range of outcomes included in KQ1: Effectiveness (clinical outcomes, behavior change, and costs and economics) across the phases of medication management as well as reconciliation and education.

Analytic Framework

To provide a focus and structure for this review, an analytical model that incorporated the key component for seven key questions was developed. This provided direction for the literature search and guidance for the data abstraction and reporting (Figure 2).

Figure 2. Conceptual model addressing the seven key questions: enabling medication management through health IT



CDSS = computer decision support system, EMR = electronic medical records system, e-RX = e-prescribing, BCMA = bar code medication administration, CPOE = computer provider order entry, PIS = pharmacy information system, PDAs = personal digital assistant devices, eMAR = electronic medication administration records

Literature Search Methods

In the course of searching the literature, reference sources were identified; a search strategy for each source was formulated, executed, and documented (see Appendix A, Exact Search Strings). For the searching of electronic databases, database-appropriate subject headings and text-words were used. Given the broad range of questions and outcomes that the report addresses, searches were performed by first using text-words relating to the various types of health IT applied to medication management. These searches were combined with both medication management terms and computer and technology terms. No limits based on methodological terms were used as all study designs were considered. A number of grey literature resources and AHRQ resources were also searched (see Appendix A, Exact Search Strings).

The search strategies were peer reviewed by a librarian following the Peer Review of Electronic Search Strategies (PRESS) checklist process for systematic review searches.³⁵ The TEP and internal team provided references from their personal files. The reference lists of review articles were screened for eligibility.

Sources

The following databases were searched: MEDLINE,[®] EMBASE,[®] CINAHL[®] (Cumulated Index to Nursing and Allied Health Literature), Cochrane Database of Systematic Reviews, International Pharmaceutical Abstracts,[®] Compendex,[®] INSPEC[®] (which includes IEEE[®]), Library and Information Science Abstracts,[®] E-Prints in Library and Information Science,[®] PsycINFO,[®] Sociological Abstracts,[®] and Business Source[®] Complete. The search terms used are presented in Appendix A.

Supplemental searches targeting grey literature sources were conducted and included New York Academy of Medicine, SIGLE, U.S. HHS Health Information Technology, Health Technology Assessment reports from the U.K. Centre for Reviews and Dissemination, ProQuest Dissertations, National Library for Health United Kingdom (includes Bandolier), ProceedingsFirst, PapersFirst, National Technical Information Service, and Google. As part of the grey literature search, AHRQ made all references in their e-Prescribing, bar coding, and CPOE knowledge libraries available.

Search Terms and Strategies

Terms related to specific MMIT applications and in combination with both medication management terms and more general computer and technology terms, were prepared. The MEDLINE[®] search formed the basis for all other databases, but searches were edited as needed depending on the features of the database being used. When possible, letters, editorials or commentaries, and animal studies were excluded electronically. No limits were placed on language or time to capture the global literature and early studies.

Organization and Tracking of the Literature Search

Searching was done in the fall of 2009 and updated in early summer 2010. The results of the searches were downloaded into Reference Manager[®] version 10 (ISI ResearchSoft) and uploaded into our customized systematic review management system (Health Information Research Unit, McMaster University). The system is Web -based. It allows management of the systematic review process with improved auditing and control capabilities including automatic production of tables and tabulations. The system stores the full text of articles in portable document format (PDF) and tracks duplicates, results of title and abstract review, which articles were included or excluded with reasons, and data abstraction levels.

Title and Abstract Review

The study team reviewed titles and abstracts of all articles retrieved using prepared data abstraction forms (Appendix B, Sample Screening and Data Abstraction Forms). Two blinded, independent reviewers from a team of reviewers conducted title and abstract reviews in parallel. Both reviewers had to indicate that the article was to be excluded for it to be removed. Both reviewers also had to agree on inclusion for the article to be promoted to the next level. In the case of disagreements, a third reviewer determined if the article was to be promoted to the next level of screening.

This first review level was designed to detect all articles that reported on medication management with health IT assisting in the medication management process. Reviewers were instructed to consider applications as health IT if they were integrated with other information systems (rather than stand-alone applications or devices), with the systems being more than

passive vehicles for data transfer. We defined health IT as electronic systems that collect, process, or exchange health information about patients and formal caregivers. We included articles only if the MMIT was integrated with at least one health IT system, such as EHR or EMR systems, and that it processed patient-specific information and provided advice or suggestions to either the health care provider or the patients and their families on issues related to health or wellness care. We excluded stand-alone devices (no integration) with the exception of personal digital assistants (PDAs) or handheld devices into which clinicians or patients entered patient-specific information to assist in medication management. PDAs are an important focus for AHRQ. All articles about transmission or order communication between pharmacist and clinical prescriber were also included and tagged as Electronic Data Interchange (EDI).

Review articles were passed through to the second level of screening. Once identified, the bibliographies of the reviews were screened for articles with potential for inclusion and their citations were put through the screening process starting at the title and abstract level if they had not already been captured by the original search. The systematic reviews were also included in the answers to the seven key questions where appropriate.

Defining Medication Management Health IT

To be clear on what kinds of applications were included in MMIT, the following outline for MMIT applications was devised and used by screeners.

MMIT systems or programs were included if:

- The computer or technology processed patient-specific information,
- The information provided by the system was relevant to one of the five phases of medication management or two ancillary aspects (education and reconciliation):
 - Prescribing or ordering medications,
 - Order communication (transmission, clarification, verification),
 - Dispensing,
 - Administering (by health care provider, patient, or caregiver),
 - Monitoring (signs, symptoms, or laboratory data to ascertain patient adherence, adverse events, or the need for medication adjustment),
 - Education (of patients or care providers, but not preprofessional education),
 - Reconciliation of medication lists,
- Someone (e.g., patient, caregiver, family, health care professional) received information in return that was, or could be, linked to patient-specific information used in decisionmaking,
- The technology was part of, or linked to, another electronic information system,
- The article contained outcome data related to one of the areas of interest set out in the key questions.

Articles were to be excluded if they were health IT systems or programs and:

- The IT component was only Web or local browsing of general health information databases or information resources (e.g., online textbooks),
- The system acted as a conduit of information only (except order communication of prescriptions between health care providers and pharmacists),
- Systems where no feedback was provided for patient care (e.g., surveys),

- The system did not help with medication management decisionmaking or provide information about any of the medication management phases (prescribing, order communication, dispensing, administering, and monitoring), or education and reconciliation,
- Systems that made measurements but did not process the information,
- Stand-alone devices that do not integrate with information systems (except PDAs using patient specific information),
- The health IT application was used only to extract data (e.g., pill bottles that track opening and closing, smart infusion pumps not tied to other systems, studies using EMRs for data collection if the data were for quality improvement or other related tasks but not direct patient care).

Data Abstraction

Given the range of questions addressed, data abstraction was performed by a core group of staff for KQ1 and KQ7. Abstraction was done by one reviewer, and the accuracy was checked by a second reviewer. The authors of the report performed a final check on the abstracted data. The reviews were not blinded in terms of the article authors, institutions, or journal.

- For all articles, reviewers abstracted information on general study characteristics: study design, the intervention, study population, setting, disease, drugs of interest, and description of the MMIT application (see Appendix B).
- Outcomes data were abstracted from the articles that were applicable to KQ1: Effectiveness and KQ7: RCTs of CDSS regarding the MMIT application impact on a health, health care process, or other intermediate outcomes.
- We abstracted only the main endpoints (major endpoints) that authors indicated as such. If no main endpoint measures were indicated, we abstracted data on outcomes related to medication management and clinical outcomes and relied on the order that those outcomes were presented in the results section, methods description, or abstract.
- We saw great variation in the way outcomes and statistical methods were reported by article authors, even when using similar systems. As a result, for this report it was recorded whether the main endpoint was positively changed by the intervention (noted as + in Appendix C, Evidence Tables). The main endpoint could also be unchanged (noted as = in Appendix C, Evidence Tables). Some studies reported a negative effect where the predefined outcome was found to be in the opposite direction sought (noted as – in Appendix C, Evidence Tables). For example, measuring an increased time to prescribe when the MMIT system was developed to reduce prescriber time. In addition, those studies that identified unintended consequences (adverse effects) of the MMIT systems are summarized in their own section. If more than one main endpoint was reported, the positive and negative referred to the direction of the majority of outcomes.

Articles addressing KQ4: System Characteristics, KQ5: Sustainability and KQ6: two-way prescription EDI were abstracted separately to capture relevant outcome data.

Assessment of Study Quality

The included studies were assessed on the basis of the quality of their reporting of relevant data. Quantitative studies were assessed using the same criteria employed by Jimison et al.,³ in a

previous AHRQ report. RCT scoring was based on Delphi consensus work by Verhagen and colleagues,¹⁰ and is referred to in this report as the ‘Verhagen/AHRQ RCT quality scale.’ Quality assessments of applicable articles were performed by more experienced reviewers to maintain consistency and accuracy. Studies with before-after, time series, surveys, and qualitative methods were not assessed for quality because few well-validated instruments exist and the study design itself is considered lower on the hierarchy of evidence.

Method assessments used for articles of the relevant design:

Verhagen/AHRQ RCT quality scale (scored out of nine)

1. Was the assignment to the treatment groups really random?
2. Was the treatment allocation concealed?
3. Were the groups similar at baseline in terms of prognostic factors?
4. Were the eligibility criteria specified?
5. Were outcome assessors blinded to the treatment allocation?
6. Was the care provider blinded?
7. Was the patient blinded?
8. Were the point estimates and measure of variability presented for the main endpoint measure?
9. Did the analyses include an intention to treat analysis

Cohort studies (scored out of ten)

1. Was there sufficient description of the groups and the distribution of prognostic factors?
2. Are the groups assembled at a similar point in their disease progression?
3. Is the intervention/treatment reliably ascertained?
4. Were the groups comparable on all important confounding factors?
5. Was there adequate adjustment for the effects of these confounding variables?
6. Was a dose response relationship between intervention and outcome demonstrated?
7. Was outcome assessment blind to exposure status?
8. Was followup long enough for the outcomes to occur?
9. What proportion of the cohort was followed-up?
10. Were drop out rates and reasons for drop out similar across intervention and unexposed groups?

Case-control studies (scored out of nine)

1. Is the case definition explicit?
2. Has the disease state of the cases been reliably assessed and validated?
3. Were the controls randomly selected from the source of population of the cases?
4. How comparable are the cases and controls with respect to potential confounding factors?
5. Were interventions and other exposures assessed in the same way for cases and controls?
6. How was the response rate defined?
7. Were the nonresponse rates and reasons for nonresponse the same in both groups?
8. Is it possible that over-matching has occurred in that cases and controls were matched on factors related to exposure?
9. Was an appropriate statistical analysis used (matched or unmatched)?

Case series (scored out of six)

1. Is the study based on a representative sample selected from a relevant population?
2. Are the criteria for inclusion explicit?
3. Did all individuals enter the survey at a similar point in their disease progression?
4. Was followup long enough for important events to occur?
5. Were outcomes assessed using objective criteria or was blinding used?
6. If comparisons of subseries are being made, was there sufficient description of the series and the distribution of prognostic factors?

Data Synthesis

Evidence tables with article details were created and ordered by key question, subquestion, and medication management phase as applicable (Appendix C). This offered another opportunity to check abstracted elements with the original articles; any errors were brought to the attention of the abstractors of the specific section for correction. Meta-analyses were not performed on any data because of the heterogeneity of the studies, as well as the nature of the observational studies in most sections.

Data Entry and Quality Control

General study data for each article was abstracted by one staff member and entered into the online data abstraction forms (Appendix B). Second reviewers were generally more experienced members of the research team, and one of their main priorities was to check the quality and consistency of the first reviewers' answers and to perform the quality assessment where required.

Grading the Evidence

Because so much of the material was derived from observational studies, we did not provide grades for the evidence beyond quality scoring of the RCTs, cohort, case-control, and case series studies.

Peer Review

Throughout the project, the core team sought feedback from internal advisors and technical experts. These technical experts were members of the TEP and other content and methodology experts as needed. The report was reviewed in several stages, comments considered and incorporated into this final report. Members of the TEP and the peer reviewers are listed in Appendix E. Many of the TEP members also reviewed the initial version of the document. Both the members of the TEP and the review panel provided valuable comments and have made the final document stronger.

Results


Results of the Literature Search

The literature search retrieved 40,582 articles for screening for inclusion; this includes 93 from hand searches and 408 from the grey literature (Figure 3). We excluded 7,797 duplicates and screened 32,785 titles and abstracts. A total of 4,578 articles were screened at full-text. Reasons for exclusion at this stage included inaccessible copies of full-text, articles in foreign languages that were not translated, unretrievable theses, studies not using integrated technologies or technologies that did not impact on medication management (“not MMIT”), studies that were not a primary study with first hand observations (often review articles), or studies not measuring an outcome of interest to our key questions.

We found 361 articles which met our content criteria for Key Question (KQ) 1: Effectiveness but did not use formal qualitative methods or have comparison groups with hypothesis testing or appropriate statistical analyses (quantitative studies). These articles are not included in the synthesis but they are integrated into the report bibliography with the other articles that were synthesized. This left 428 articles that are synthesized in the evidence report.

A total of 377 articles are quantitatively synthesized in KQ1: Effectiveness. One article was included as evidence for KQ3: Value Proposition although 30 articles were cited directly and many more were used in some of the summaries described in this section; 21 articles were synthesized for KQ4: System Characteristics, 24 for KQ5: Sustainability, 33 for KQ6: Two-way Prescription EDI and 77 for KQ7: RCTs of CDSS. A number of articles were included in more than one key question response as they addressed more than one aspect of medication management.

Figure 3. Literature flow of medication management studies



To address the various outcomes measures or interest in the seven key questions, 428 articles were synthesized. An additional 361 articles met content criteria for integrated technology enabling medication management, but these did not meet methodological criteria of either formal qualitative methods or those with comparison groups and appropriate statistical analysis; these studies are included in the bibliography.

KQ1. Within and across the phases of medication management continuum, what evidence exists that health IT applications are effective?

Effectiveness Studies Overall

KQ1: Effectiveness includes the largest number of articles for any of the seven key questions. Our searching for KQ1 concentrated on content with no limits on methods. The articles were divided into two groups: (1) all qualitative studies were included in the analysis that forms the basis of this report; and (2) all quantitative studies were included in the analysis section if they included a comparison group and data on each group and if they contained statistical methods defined by statistical testing, a statement of hypotheses-based research defined *a priori*, or both.

Articles that met these criteria for qualitative or quantitative studies were analyzed in this report (n = 377). An additional 51 studies that did not meet the above methods criteria were included in KQs 3 to 6; the literature on these topics was sparse. Articles that met content criteria but that did not meet these methodological criteria are included in our bibliography (n = 361) but not in any of the tables nor are they analyzed in the report.³⁶⁻³⁹⁶

The final analysis for addressing KQ1 included 379 articles. Substantial variation exists in the concentration of evidence and content across issues related to MMIT. Table 1 shows the numbers of studies within each of the five phases plus reconciliation and education by study design. By far, more studies are done in the prescribing phase (n = 263) with a substantial number done in monitoring (n = 77). Dispensing is the phase that is least studied and little evidence exists on education and reconciliation. Figure 4 depicts the trends in publication frequency of articles included for analysis in the report. We saw a dramatic increase in publication of MMIT studies after 2000, most notably in studies dealing with prescribing.

Table 1. Research design for studies across the phases of medication management and education and reconciliation


| Design | P | OC | D | A | M | E | R |
|---------------|-----|----|----|----|----|---|---|
| RCT | 69 | 1 | 2 | 2 | 37 | 1 | 1 |
| Cohort | 13 | 2 | 2 | 1 | 6 | 0 | 1 |
| Observational | 144 | 18 | 10 | 26 | 29 | 2 | 4 |
| Qualitative | 37 | 5 | 3 | 10 | 5 | 0 | 0 |
| Total | 263 | 26 | 17 | 39 | 77 | 3 | 6 |

Note some studies cross more than one phase. See Appendix C, Evidence Table 16 for references to the included articles in each cell.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

RCT = Randomized controlled trial

Figure 4. Trends in publication of articles relating to the phases of medication management across years until searching was completed in June 2010



Strengths and Limitations of the Evidence

Table 1 illustrates that a variety of research methods were used in the studies, with the majority using observational methods. A substantial numbers of RCTs and qualitative studies were included. The large number of observational studies is reflective of the nature of the domain in that many of the articles retrieved were more often directed at the observational description or evaluation of existing systems rather than based on classical research methods of hypothesis-driven projects.

Settings. Settings where studies were performed also show variation. Table 2 includes the settings for studies across the medication management phases plus education and reconciliation. Studies often reported multiple settings. Most studies were set in hospitals and ambulatory care. Few studies were done in the community (n = 1), home (n = 5), or long-term care (n = 8). The lack of comparative, hypothesis-driven studies set in pharmacies is offset by a larger group of pharmacy studies that were descriptive in nature. Despite the lack of studies set in pharmacies, many studies relied on pharmacies and pharmacists.

Table 2. Settings for the phases of medication management and reconciliation and education

| Setting | P | OC | D | A | M | E | R |
|--|-----|----|----|----|----|---|---|
| Ambulatory care (e.g., clinic, doctors office) | 94 | 6 | 2 | 1 | 40 | 2 | 0 |
| Community (e.g., school, community centre) | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| Home | 2 | 0 | 0 | 2 | 5 | 0 | 0 |
| Hospital | 164 | 12 | 9 | 34 | 36 | 1 | 6 |
| Long-term care | 4 | 0 | 2 | 3 | 1 | 0 | 0 |
| Pharmacy | 11 | 13 | 10 | 2 | 4 | 0 | 1 |

Note some studies cross more than one phase or setting. See Appendix C, Evidence Table 17 for references to the included articles in each cell.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Clinicians. Physicians were the most represented clinicians studied (Table 3). Many of the health professionals functioned in primary care and other ambulatory settings. Often studies did not differentiate among specialties or included many specialties in a single study. Nurses were most often studied in the administering phase and pharmacists were involved in order communication. We did not identify any studies that evaluated dentists and found few studies of mental health professionals or midlevel practitioners (e.g., nurse practitioners, midwives, and physician assistants).

Table 3. Clinicians evaluated in outcomes studies of medication management phases, education, and reconciliation

| Clinicians | P | OC | D | A | M | E | R |
|--|----|----|---|----|---|---|---|
| Primary care physicians | 25 | 2 | 1 | 0 | 3 | 1 | 0 |
| Specialists | 11 | 0 | 0 | 0 | 1 | 0 | 0 |
| Hospitalists | 18 | 0 | 0 | 0 | 2 | 0 | 0 |
| Other Physicians | 7 | 0 | 0 | 0 | 2 | 0 | 0 |
| Physicians undifferentiated | 26 | 1 | 0 | 1 | 3 | 0 | 1 |
| Nurses | 20 | 1 | 0 | 16 | 2 | 0 | 1 |
| Mid level practitioners (e.g., PA, NP, MW) | 6 | 0 | 0 | 0 | 2 | 0 | 0 |
| Pharmacists | 13 | 6 | 5 | 2 | 1 | 0 | 1 |
| Other health professionals | 10 | 0 | 2 | 4 | 1 | 0 | 0 |
| Hospital administrators | 4 | 1 | 0 | 1 | 0 | 0 | 0 |

Note some studies cross more than one phase and clinician type. See Appendix C, Evidence Table 18 for references to the included articles in each cell.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

MW=midwife, NP=nurse practitioner, PA=physician assistant

Patient population studied. Patients studied represented those who were most likely to need medication: adults, middle aged people, and those over the age of 65 years. Infants, children, and

adolescents were also studied but to a lesser extent (Table 4). Monitoring and reconciliation concentrated on older persons.

Table 4. Patients and caregivers studied by phase of medication management and education and reconciliation

| Patients | P | OC | D | A | M | E | R |
|------------------------|----------|-----------|----------|----------|----------|----------|----------|
| Infants (0 to 2 years) | 6 | 1 | 1 | 1 | 1 | 0 | 0 |
| Children (2 to 12) | 11 | 1 | 1 | 1 | 3 | 0 | 0 |
| Adolescents (13 to 18) | 15 | 0 | 0 | 0 | 9 | 1 | 1 |
| Adults (19 to 44) | 40 | 0 | 0 | 0 | 23 | 2 | 1 |
| Middle age (45 to 64) | 56 | 1 | 0 | 1 | 38 | 3 | 1 |
| Geriatric (65 plus) | 60 | 2 | 2 | 1 | 32 | 2 | 3 |
| Undifferentiated | 9 | 0 | 0 | 0 | 8 | 0 | 0 |

Note some studies cross more than one phase and patient group. See Appendix C, Evidence Table 19 for references to the included articles in each cell.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Technology. Tables 5 and 6 list the MMIT applications studied. Table 5 includes those MMIT applications that were the main focus of the study while Table 6 includes the MMIT that were integrated with the MMIT studied. The CDSS and reminder systems were most common in prescribing and monitoring. CPOE and e-Prescribing were also commonly used. Systems associated with pharmacy use were less commonly studied. Considering integration, the health IT in medication management is well-integrated with comprehensive systems such as EMRs and hospital information systems as well as other components of the broader health IT domain, remembering that integration with a health IT system was a requirement for inclusion in our review. Although prescribing again is the major phase studied, the other phases are represented.

Table 5. Main health IT studied by medication management phase and education and reconciliation

| Health IT | P | OC | D | A | M | E | R |
|--|-----|----|---|----|----|---|---|
| CDSS/reminders | 177 | 8 | 4 | 5 | 63 | 1 | 1 |
| CPOE/POE system | 90 | 12 | 5 | 9 | 11 | 0 | 0 |
| e-Prescribing | 31 | 10 | 3 | 4 | 2 | 1 | 1 |
| Order transmission of the prescription to and from doctor to pharmacy electronically | 2 | 3 | 0 | 0 | 0 | 0 | 0 |
| Pharmacy information system | 2 | 3 | 4 | 1 | 0 | 1 | 1 |
| Barcoding medication administering | 1 | 0 | 2 | 20 | 0 | 0 | 0 |
| Barcoding dispensing | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| eMAR, e-TAR | 2 | 2 | 2 | 14 | 0 | 0 | 0 |
| Other | 13 | 2 | 3 | 7 | 14 | 1 | 5 |
| Personal digital assistants or hand-helds | 7 | 1 | 0 | 4 | 5 | 0 | 1 |

Note some studies cross more than one phase and technology. See Appendix C, Evidence Table 20 for references to the included articles in each cell.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

CDSS = Clinical decision support system, CPOE = Computerized provider order entry, POE = Provider order entry, eMAR = Electronic Medication Administration Record system, eTAR = Electronic Treatment Administration Record system

Table 6. Health IT integrated with the health IT being studied

| Integrated Health IT | P | OC | D | A | M | E | R |
|---------------------------------|-----|----|---|----|----|---|---|
| EHR/EMR system | 120 | 11 | 5 | 15 | 39 | 1 | 3 |
| Formulary | 11 | 1 | 1 | 3 | 3 | 0 | 0 |
| Pharmacy | 50 | 9 | 8 | 15 | 18 | 0 | 2 |
| CPOE/POE system | 58 | 0 | 1 | 7 | 12 | 0 | 2 |
| Hospital information system | 54 | 4 | 3 | 9 | 13 | 1 | 1 |
| Laboratory system | 49 | 3 | 1 | 3 | 27 | 1 | 0 |
| Imaging systems | 22 | 2 | 1 | 3 | 7 | 1 | 0 |
| CDSS/reminders | 24 | 1 | 0 | 1 | 6 | 0 | 0 |
| Billing/administration system | 8 | 0 | 0 | 1 | 4 | 0 | 0 |
| Insurance | 5 | 0 | 0 | 0 | 1 | 0 | 0 |
| Personal health records systems | 1 | 0 | 0 | 0 | 2 | 0 | 0 |
| Patient decision support system | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| Barcoding system | 2 | 0 | 1 | 0 | 0 | 0 | 0 |
| Not specified | 32 | 4 | 4 | 6 | 8 | 0 | 0 |
| Other | 15 | 3 | 0 | 5 | 6 | 1 | 1 |

Note some studies cross more than one phase and integrated with more than one technology. See Appendix C, Evidence Table 21 for references to the included articles in each cell.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

EHR = electronic health records system, EMR = electronic medical records system, CDSS = Clinical decision support system, CPOE = Computerized provider order entry, POE = Provider order entry

Summary. In summary, prescribing is the major medication management phase studied. Studies were often evaluative rather than research centered in nature, as reflected in the number of

observational studies. Substantial numbers of RCTs and qualitative studies exist. Most often studies were set in hospitals and in ambulatory care facilities. Few studies were set in pharmacies, although most of the articles showed interactions with pharmacists and pharmacies. Long-term care, community settings, and homes were not often studied. CDSS systems were the most common MMIT application studied. CPOEs were also studied substantially. The MMIT applications studied were very often embedded within a larger EMR, hospital, or pharmacy information system and integrated with other health IT applications. Many different MMIT systems were studied although again few were done outside the prescribing and monitoring phases and variation existed in the number of studies of each kind of health IT. Physicians were the health professionals most often studied. Few patients were evaluated.

Process Changes—Prescribing

Summary of the Findings for Process Changes

Of the 378 articles that have outcomes associated with MMIT, 174 (46 percent) are reports of the evaluation of processes in the prescribing phase of medication management (Appendix C, Evidence Table 1). Because prescribing and ordering are substantially different in the hospitals and ambulatory settings, the remainder of this section will provide analyses with the articles divided into hospital-based studies (n = 107) and ambulatory care-based studies (n = 67). The community- and home-based studies are included with ambulatory care. Only one study in this section was done in a long-term care facility.³⁹⁷

General Study Characteristics

Strengths and Limitation of the Evidence. The studies of process changes in MMIT based in hospital settings have a lower proportion of RCTs than ambulatory care studies. The 107 hospital-based studies are comprised of 19 RCTs (18 percent of hospital studies),³⁹⁸⁻⁴¹⁶ 84 observational studies,^{18,417-499} 3 cohort studies,⁵⁰⁰⁻⁵⁰² and 1 mixed methods study.⁵⁰³

The 67 articles set in primary care, communities, and home (ambulatory care) were studied using 40 RCTs (61 percent of nonhospital studies),⁵⁰⁴⁻⁵⁴³ 2 cohort studies,^{544,545} 1 case control study,⁵⁴⁶ 1 mixed methods study,⁵⁴⁷ and 22 observational studies.^{431,548-568} The long-term care study was an RCT.³⁹⁷

Table 7. Research methods of studies that evaluated process changes associated with the prescribing phase of MMIT

| Design | Hospital Based | Ambulatory Care Based | Long Term Care |
|---------------------------|----------------|-----------------------|----------------|
| RCT | 19 | 40 | 1 |
| Cohort | 3 | 2 | 0 |
| Case control | 0 | 1 | 0 |
| Qualitative/Mixed Methods | 1 | 1 | 0 |
| Observational | 84 | 23 | 0 |

RCT = Randomized controlled trial

Patient population. Not all studies included descriptive data on patients. Of those that did, the patient populations reflected the pattern of medication use with more studies including

participants who were, on average, over the age of 44 years. All age groups were studied in both hospital and ambulatory settings.

Thirty-seven studies set in hospitals provided descriptive data on participants. Four studies included infants (birth to 2 years),^{418,463,467,491} five studies included children (2 to 11 years),^{411,418,437,467,569} six studies included adolescents (12 to 18 years),^{411,418,437,446,467,490} eight studies evaluated adults (19 to 44 years),^{408,411,418,437,446,467,490,492} 22 studies included middle age participants (45 to 64 years),^{399,401-403,406,414,425,428,430,431,433,445,448,449,454,467,475,481,485,489,490,493} and 27 studies included geriatric participants (65 years and up).^{399,401-403,406,407,413,414,416,425,428,430,431,433,440,445,448,449,453,467,481,483,485,486,489,490,493}

Thirty-six of the 67 studies done in ambulatory settings included descriptive data on the patient population studied. Many studies included participants in a wide range of ages. Of the 36 studies that included patient information, one studied infants (up to 2 years of age),⁵⁵⁹ two evaluated medication management in children (2 to 11 years),^{539,559} three evaluated adolescents (12 to 18 years),^{504,537,539} 15 were of adult population (19 to 44 years),^{504,505,508,512,515,522,527,528,530,531,537,541,543,545,565} 24 studied patients in the middle age range (45-64 years),^{504,505,508,510,513,515,518-520,522,526-528,530,531,537,538,541-543,545,557,565,568} and 22 studies include geriatric patients (65 years and up).^{505,507,510,513,515,519,522,524,528,530,531,533,541-543,545,552,565,566,568} The single long-term care study did not describe its patients.³⁹⁷

Clinicians studied. Only 11 of the hospital studies included descriptive information on clinicians. Several of these included multiple groups of health professionals: hospitalists^{398,405,407,452,454,486} other physicians,^{407,415,487,488,502} other health professionals,⁴⁵² and nurses.⁴³⁹ Many of the other studies evaluated clinicians but did not provide sufficient demographic information for analysis or discussion.

Few of the studies set in ambulatory care provided substantial information on clinicians. Those clinicians who were specifically described were primary care clinicians,^{506,510,514,529,532,534,535,552,563,566} other physicians,^{514,525,535,544,547,553,554} nurses and midlevel practitioners (physicians assistants, nurse practitioners, advanced practice nurses, and midwives),^{535,561} and pharmacists.⁵¹⁸

Technology. Technology associated with studies set in hospital settings often evaluated several integrated MMIT systems, although some studies included only one MMIT. Individual MMIT applications included CDSS systems,^{18,397-399,401-405,407-418,420,422-428,430,431,433,435,448,449,451-454,458,460-462,464,466,468-478,480-483,485,486,489-502} CPOE systems,^{400,406,417,419,421,427,429,432,437,438,441-444,446-450,453,455,457,459,463,466-469,479-484,487,488,491,493,503} pharmacy information system,⁴³⁸ computerized unit dose drug dispensing system,⁴⁶⁵ e-prescribing,^{434,436,439} medication safety reporting system,⁴⁵⁶ an internet electronic diary for patients,⁴⁰⁸ and eMAR systems.^{439,456}

Studies set in ambulatory care also studied a range of MMIT applications with the majority of the MMIT applications based on CDSS systems. The MMIT applications were CDSS,^{504-531,533-543,545,546,548,550-560,563-566,568,570} CPOE,^{507,527,532,540,551,558,567} e-Prescribing,^{540,544,547,549,561,562} and a pharmacy information system.⁵⁰⁷

These MMIT systems that were the focus of study were also integrated with a range of other MMIT applications, most of which were some form of EHR or EMR systems. The hospital-based studies included integration with bar-coding systems,⁴⁹³ billing or administration,^{400,406,446,474,485} CDSS,^{417,429,432,442,448,461,462,479} CPOE,^{398,400,402,405,407,410,411,413,415,416,423,426,427,430,433,435,440,445,448,454,458,460-}

462,464,471,472,480,486,490,492,494-497,500,501 EHR and EMR systems,^{18,398,405,407,409,411,415,416,418,422,423,430,431,434,436-438,442,444,445,447,448,453,455,460,462,466,473,475,479,480,483,489,492,498,501} hospital information systems,^{400,403,406,412,414,415,424,425,428,438,442,443,450,451,453,456,461,465,467-470,474,478,481,482,487,488,491,499} imaging systems,^{405,437,442,443,456,463,466-468,470,475,483} laboratory systems,^{18,401,405,407,409,415,418,420,437,442,444,446,452,453,456,461,466,470,473,475,477,483,485} formulary systems,^{406,442,443} and an integrated pharmacy system.^{401,404,415,417,420,434,436,439,441-443,463,473,477,479,485,493,502}

Systems integrated within ambulatory studies were also mostly EMR and EHR systems. The ambulatory studies included billing and administrative systems,^{504,511,512,532,564,567} CPOE,^{511,512,518,522,529,532,548,550,564,570} EHR and EMR systems,^{505-507,510-513,515,517-524,526-528,531,533,535,536,538-540,542,543,545,546,548,550,551,554,556-561,564,566,568} formulary systems,^{544,549} hospital information systems,^{508,509,541,565} imaging systems,^{513,532,538} insurance systems,^{542,549} laboratory systems,^{504,509,513,516,538,541,548,550,555} and pharmacy systems.^{504,508,511,516,518,541,548,552,562}

Outcomes

Prescribing changes. Sixty-one studies evaluated changes in prescribing in hospital settings. Fifty-three (87 percent) showed statistically significant improvements in at least half of its main endpoints. Categorizing these 53 articles into groups based on study methods, 11 were RCTs,^{398,399,402-405,408-410,413,416} two were cohort studies,^{501,502} and 40 were observational.^{417,423,424,426,428,430,431,440,443,446,447,451,452,456,458-460,462,464,466-473,475,478,482,486,489,490,492,496-499,547,571} Eight (13 percent) of the 61 did not show statistically significant changes. Two were RCTs^{411,414} and six were observational studies.^{418,420,445,449,477,485}

Thirty-two studies set in ambulatory care found improvements in prescribing with the MMIT as defined by at least half of the main endpoints being positive. Twenty-three were RCTs.^{504,506-508,511-515,517,522,525,526,528-531,533,535-537,539,541} One was a case-control study.⁵⁴⁶ Seven were observational.^{549,553,554,562,565,566,568} Eleven studies set in ambulatory care settings sought to determine if prescribing was improved with the introduction of MMIT and they did not demonstrate differences. Of these, five were RCTs,^{510,520,523,524,538} two were cohort studies,^{544,545} and four were observational.^{557,560,564,572}

Errors. Twenty-two articles studied prescribing errors in hospital settings. Fifteen showed statistically significant improvement in at least half of the main endpoints. Two were RCTs^{400,405} and 13 were observational.^{421,434,438,439,450,453,454,457,465,474,479,491,493} Seven did not show statistically significant improvements: a mixed methods study⁵⁰³ and six observational reports.^{419,432,436,444,455,495} Two studies reported errors related to ambulatory care studies and both were positive, one using CPOE,⁵⁵⁸ and one using CDSS,⁵⁵⁹ both in observational studies.

Time considerations. Seven studies evaluated time considerations in hospital settings. Two observational studies showed statistically significant improvements in considerations of time.^{439,488} One study found a statistically significant increase in time to prescribe.⁴⁸⁷ One study evaluated mean time on antimicrobial management but did not do statistical testing.⁴⁰¹ Three observational studies did not show any differences in time.^{419,441,481} Five studies assessed time savings in the ambulatory care settings. Four were positive: an RCT of CPOE and CDSS that reduced time to respond to alert situations,⁵²⁷ a cohort study on time spent on asthma management,⁵⁴⁵ and two observational studies, one on e-Prescribing⁵⁶¹ on time spent on

computer and paper tasks and one on time spent on ordering laboratory testing for monitoring.⁵⁵⁰ One RCT showed that time spent on patient care did not decrease with the introduction of a CPOE system.⁵³²

Adherence to guidelines, reminders, and recommended practice. Twenty-four studies measured improvements in compliance with guidelines, reminders, and recommended practices in hospital based studies. Nineteen identified statistically significant improvements in compliance: four RCTs,^{407,412,521,573} one cohort study,⁵⁰⁰ and 14 observational studies.^{424,425,427,429,435,437,438,447,461,470,476,483,484,494} Four did not find any differences in compliance: one RCT⁴¹⁵ and three observational studies.^{442,448,480} One observational study⁴²² showed a decrease in adherence after the introduction of a CDSS system into a hospital EHR.

Thirteen studies that took place in nonhospital settings (primary care, community, and homes) considered compliance with guidelines, reminders, or recommended practice. Seven were RCTs of which five showed positive results for at least half of the main endpoints.^{505,509,534,542,543} Two RCTs^{518,519} did not identify a difference for measured compliance. One cohort study⁵⁴⁴ did not show a difference in compliance with the formulary using e-Prescribing and one mixed methods study⁵⁴⁷ reported no change in physician compliance with drug alert overrides. Four of four observational studies reported improvements in compliance with guidelines, reminders, or recommended practices.^{548,556,563,566}

Workflow. No studies set in hospitals studied workflow as one of their main endpoints that were changes in process. Two studies set in ambulatory care studied workflow. One study using CDSS reminders showed a significant reduction in missed followup appointments that had been scheduled by nurses.⁵⁵⁵ An RCT of CDSS and e-Prescribing did not affect the rate of callbacks generated between physicians and pharmacists.⁵⁴⁰

Table 8. Summary of the number of studies reporting statistically significant process changes in studies of prescribing by process for hospital and ambulatory based studies

| Process Category | RCTs | Cohort and Case-Control Studies | Observational Studies | Mixed Methods Studies |
|---|------|---------------------------------|-------------------------|-----------------------|
| Change in Prescribing (85 of 104 showed benefit) | | | | |
| Hospital Studies | 11+ | 2+ | 40+ | 0 |
| | 2= | 0 | 6= | 0 |
| Ambulatory | 24+ | 1+ | 7+ | 0 |
| | 5= | 2= | 4= | 0 |
| Errors in Prescribing (17 of 24 showed benefit) | | | | |
| Hospital Studies | 2+ | 0 | 13+ | 0 |
| | 0 | 0 | 6= | 1= |
| Ambulatory | 0 | 0 | 2+ | 0 |
| | 0 | 0 | 0 | 0 |
| Time Considerations (6 of 12 showed benefit) | | | | |
| Hospital Studies | 0 | 0 | 2+ | 0 |
| | 0 | 0 | 4= 1- ⁴⁸⁷ | 0 |
| Ambulatory | 1+ | 1+ | 2+ | 0 |
| | 1= | 0 | 0 | 0 |

Table 8. Summary of the number of studies reporting statistically significant process changes in studies of prescribing by process for hospital and ambulatory based studies (continued)

| Process Category | RCTs | Cohort and Case-Control Studies | Observational Studies | Mixed Methods Studies |
|--|----------|---------------------------------|--------------------------------|-----------------------|
| Compliance with Guidelines, Reminders, and Recommended Practice (28 of 37 showed benefit) | | | | |
| Hospital Studies | 4+ 1= | 1+ 0 | 14+ 3= 1- ⁴²² | 0 0 |
| Ambulatory | 5+ 2= | 1= | 4+ 0 | 0 1= |
| Workflow in Prescribing (1 of 2 showed benefit) | | | | |
| Hospital Studies | 0 0 | 0 0 | 0 0 | 0 0 |
| Ambulatory | 0 1= | 0 0 | 1+ 1= | 0 0 |

+ indicates that half or more of the main endpoints were shown to be statistically significant.

= indicates that at least half of the main endpoints were statistically not significant.

- indicates that the main endpoints were statistically significant in the opposite direction projected at study start.

The citations to the studies listed above are in the text preceding this table.

Ambulatory studies included those that were done outside hospitals including homes and communities.

Summary

Much research has been done to evaluate changes in process related to prescribing in hospital settings and ambulatory care situations. The research is varied in methods although many RCTs exist. A higher proportion of RCTs are done in the ambulatory care studies than in hospitals. Clinicians are the most studied. Pharmacists are often included in studies but are less frequently the major thrust of analyses. Many MMIT applications are studied in the prescribing phase. These prescribing MMIT applications are also frequently supported or integrated with EMR, EHR and hospital information systems. CDSS systems are often studied and frequently integrated with CPOE or e-Prescribing systems. Pharmacy-based MMIT applications generally lack evidence.

With respect to the process changes measured in the prescribing studies, changes in prescribing and compliance with reminders, guidelines, and standard practice are the most common outcomes for hospital- and ambulatory-based studies (Table 8). The RCTs were concentrated on evaluating changes in prescribing with some of them assessing compliance. The RCTs were positive 80 percent of the time while the observational studies were positive 77 percent of the time. Studies done in ambulatory care settings have not evaluated errors as outcome measures. Quantitative workflow studies are generally absent across all settings. MMIT in prescribing is associated strongly with improvements in prescribing and also associated, but to a lesser extent, with reducing errors and improving compliance with guidelines, reminders, and recommended practice. Time reductions or changes are not as often improved and workflow improvement assessments are lacking evidence.

Systems that provide information support, such as CDSS and CPOE systems, are combinations of technical capabilities and a knowledge base. The content of this knowledge base is probably more important than the technical aspects. Research findings and scientific evidence (i.e., evidence-based content) are difficult to compile and even more difficult to keep current. We did not find evaluations of the knowledge base of the systems or comments on updating, although some of the systems depended on clinical practice guidelines for their evidence base. Similarly, outcomes that were associated with correct knowledge such as adherence to best practice guidelines were also not often evaluated to show that they were accurate and current. Future research must address how this need for strong evidence to support the knowledge base of

CDSSs, provide evidence backing for order sets for CPOE systems, and clinical practice guidelines on which to base best practice can be best met.

Order Communication

Summary of the Findings for Process Changes

Order communication is less well-studied than prescribing with only 16 studies: two RCTs,^{540,574} and 14 observational studies.^{438,552,567,575-585} (Appendix C, Evidence Table 2) Order communication involves clinicians and pharmacists. The oldest study in this group was published in 1999,⁵⁷⁴ reflecting the recent advances in communication applications related to MMIT.

Strengths and Limitations of the Evidence

The evidence in this section is predominantly observational with two RCTs. The studies were mainly based on large sample sizes, from 39 clinicians⁵⁷⁸ to almost one million prescriptions.⁵⁷⁷ The outcomes, most often measures of efficiency and changing work patterns, were usually reported as being positive.

General Study Characteristics

Participants. All studies included physicians, other prescribers, and pharmacists. The main unit of analysis in 12 of the 16 studies was prescriptions, orders, and medications. The main unit of analysis for the other four studies were patients,^{552,578} pharmacists,⁵⁸⁰ and clinicians.⁵⁷⁴ The patients were of geriatric age (65 years or greater) or adults (45 to 64 years),⁵⁷⁸ or geriatric alone.⁵⁵² Except for these two articles, all others included undifferentiated patients.

Location. All studies included a pharmacy. Most studies were hospital-based but one study was of three mail order pharmacies,⁵⁷⁵ two were HMO pharmacies only,^{552,574} and three were in community pharmacies.^{579,585,586} Nine studies were set in hospitals,^{438,574,576,578,580-584} and four in primary care.^{540,577,585,587}

Drugs and diseases. Thirteen studies did not concentrate on one disease or disorder. One study each evaluated venous thromboembolism,⁵⁷⁸ cancer,⁵⁸⁴ and HIV/AIDS.⁵⁸⁵

Technology. The MMIT in the order communication phase is varied: six CDSS,^{438,540,552,577,580,582} eight CPOE,^{438,567,576,578,580-582,584} two eMAR systems,^{581,583} four e-Prescribing,^{540,575,579,585} one e-transmission,⁵⁷⁵ and two pharmacy information systems.^{438,574} Several studies included more than one MMIT as the major focus of the study. The ordering systems (CPOE and e-Prescribing), however, predominate.

The following MMIT applications were integrated in 15 of the 16 studies: one CDSS,⁵⁶⁷ nine EMRs,^{438,540,574,576-578,580,581,585} three hospital information systems,^{438,576,582} two imaging systems,^{574,581} three laboratory systems,^{574,578,581} six pharmacy systems,^{552,577,579,583-585} and one eMAR system.⁵⁸⁰ The nine EMRs and three hospital information systems reflect maturing of the MMIT systems with respect to order communication.

Outcomes

No process changes were presented for adherence with guidelines, monitoring, or preventive care. One article described decreases in prescribing of contraindicated drug-drug combinations in ambulatory settings.⁵⁷⁷ Another looked at the agreement between pharmacists and family physicians (need for clarification of prescriptions) with and without e-Transmission of prescriptions, again in the ambulatory setting.⁵⁷⁵ All other process changes that were the main focus of the order communication articles dealt with errors and efficiencies.

Errors. Two hospital studies addressed errors (Table 9). Mahoney et al.⁴³⁸ showed a decrease in drug-allergy violations, excessive dose, incomplete or unclear orders, and therapeutic duplication with the introduction of a CPOE and CDSS system into a pediatric standalone and another general hospital. Varkey et al.⁵⁶⁷ found an increase in the frequency of intercepted prescription errors after the introduction of another CPOE and CDSS system into the Mayo Clinic ambulatory practices.

Prescribing. Two studies showed improvements in prescribing with increased interaction between pharmacists and physicians (Table 9).^{552,577}

Efficiency and workflow. Five hospital-based studies sought to change response times (Table 9). Four showed decreased times for processing and validating orders.^{576,578,581,584} One found increased time with the introduction of a CPOE and a CDSS system.⁵⁸² Another found an increased time to checking the prescription with an e-Prescribing system compared with a paper based system (11 vs. 6 minutes, $p < 0.01$).⁵⁷⁵ Some changes were substantial. For example, a decrease from 115 minutes to 5 minutes for verification of a prescription in a study by Wielthrolter and colleagues.⁵⁸⁴ Three hospital studies found changes in work flows and processes with the introduction of a pharmacy information system⁵⁷⁴ and a CPOE and CDSS system with eMAR integration.⁵⁸⁰ Fewer callbacks occurred with the introduction of a CPOE and CDSS system integrated into a hospital EMR.⁵⁴⁰

Pearce et al.⁵⁸⁵ completed an ambulatory care based setting (three HIV clinics and two private pharmacies) and found decreased time to respond to a refill request and changes in communication patterns with MMIT involved in order communication. Mitchell and colleagues⁵⁸³ found that an eMAR system was associated with more accurate and complete recording of information. Ekeldahl and colleagues showed that the rate of picking up prescriptions did not change with the introduction of an e-Prescribing system.⁵⁷⁹

Summary

The evidence for the effects of MMIT on order communication comes from a limited number of studies, many of which were observational. The studies often include large numbers of participants or prescribing events. Most of the process evaluations show improvements, often in efficiency related to times and changing work patterns (Table 9).

Table 9. Summary of the number of statistically significant process changes in studies of order communication by process for hospital and ambulatory based studies

| Process Category | RCTs | Observational Studies |
|---|------|-----------------------|
| Errors (2 of 2 showed benefit) | | |
| Hospital Studies | 0 | 1+ |
| | 0 | 0 |
| Ambulatory | 0 | 1+ |
| | 0 | 0 |
| Prescribing (2 of 2 showed benefit) | | |
| Hospital Studies | 0 | 0 |
| | 0 | 0 |
| Ambulatory | 0 | 2+ |
| | 0 | 0 |
| Efficiencies and Time Considerations (6 of 8 showed benefit) | | |
| Hospital Studies | 0 | 4+ |
| | 0 | 2 ^{-575,582} |
| Ambulatory | 0 | 2+ |
| | 0 | 1= |
| Workflow Issues (4 of 4 showed benefit) | | |
| Hospital Studies | 2+ | 2+ |
| | 0 | 0 |
| Ambulatory | 0 | 0 |
| | 0 | 0 |

+ half or more of the major endpoints showed statistically significant improvements.

= less than half of the major endpoints showed statistically significant improvements.

- the outcomes were found to be in the opposite direction (a true negative study).

RCT = Randomized Controlled Trial

Dispensing

Summary of the Findings for Process Changes

Some overlap and duplication of studies occurs between the dispensing phase of medication management and the phases of order communication and administering. Nine studies were identified as evaluating dispensing (Appendix C, Evidence Table 3).^{438,507,552,574,585,586,588-590}

Much diversity was seen across these studies. In addition to these articles, a health technology assessment (HTA) report from the Canadian Agency for Drugs and Technologies in Health (CADTH) produced a systematic review on errors in dispensing and administering in hospitals in 2009.¹¹ It sought to assess the effectiveness and economic impact of MMIT applications designed to improve medication dispensing and administering in hospitals.

Strengths and Limitations of the Evidence

Study methods included three RCTs,^{507,574,588} one cohort study,⁵⁸⁶ and five observational studies.^{438,552,585,589,590} The HTA report found 30 studies on dispensing, administering, or both, most of which were done using observational methods. In addition, many of these studies evaluated technologies that were older, no longer available, or only available in Europe. Overall the authors of the report stated that the evidence on the effectiveness of MMIT for improving medication dispensing is lacking, of poor quality, and has limited applicability.¹¹ The year of publication of the nine papers in this AHRQ document were more recent: 1997,⁵⁸⁹ 1999,⁵⁷⁴ 2001,⁵⁵² 2007,^{438,507,586} 2008,⁵⁸⁸ and 2010.^{585,590}

General Study Characteristics

Participants. Raebel and colleagues⁵⁰⁷ and Halkin and colleagues⁵⁵² reported data based on patients as the unit of study. Both studies included patients older than 65 years. Two reports studied pharmacists.^{574,588} All others reported data on medications or prescribing events as their unit of analysis.

Location. Four studies were set in community pharmacies^{552,585,586,588} and one in an HMO pharmacy.⁵⁰⁷ The other study locations were pharmacies and clinics in hospitals.^{438,574,589,590}

Drugs and diseases. Aspirin for patients with diabetes was studied,⁵⁸⁸ and two others targeted groups of medications with high potential for interactions.^{507,552} One study included e-Prescribing in three HIV clinics and two private pharmacies.⁵⁸⁵ One article concentrated on drugs that are used heavily by seniors.⁵⁰⁷ All others studied the range of prescriptions available in the pharmacies.

Technology. The technology described in these dispensing studies varied considerably. Four studies evaluated pharmacy information systems,^{438,507,574,588} three looked at eMAR systems,^{585,589,590} and CDSS,^{438,507,552} one at an e-Prescribing system,⁵⁸⁶ and one evaluated a CPOE application.⁴³⁸ These systems were integrated with an EHR or EMR system,^{438,507,574,585} hospital information system,^{438,589} a pharmacy system,^{552,585,590} a laboratory system,⁵⁷⁴ a formulary,⁵⁸⁹ or a CPOE system.⁵⁹⁰ The HTA report includes a good description of their evaluation of MMIT that provides additional background information on administering and dispensing MMIT applications.⁵⁹¹

Outcomes

Each of the main endpoints for the trials was found to be positive. Efficiency, monitoring, and preventive care outcomes were not reported in the nine studies.

Errors. Three of the four hospital based studies assessed errors and all showed improvements with eMAR,⁵⁸⁹ CPOE, CDSS and a pharmacy information system,⁴³⁸ and a pharmacy information system with CPOE.⁵⁹⁰ None of the ambulatory care studies assessed errors. The HTA report provided some evidence that BCMA is associated with reduced errors for dispensing (pharmacists) and administering (nursing), with the BCMA either as a stand-alone system or integrated with other health IT applications. Evidence on other outcomes or technologies in dispensing was found to be lacking or inconclusive.¹¹

Adherence to guidelines. For pharmacists who were prompted electronically to suggest aspirin to patients with diabetes when they were filling other prescriptions, the use of aspirin increased.⁵⁸⁸

Changes in medications to be administered. Four of the four ambulatory studies demonstrated statistically significant improvements in what drugs were dispensed. Alerts to pharmacists improved dispensing of medications with high potential for interactions in an HMO pharmacy,⁵⁵² while the use of contraindicated medications decreased with most of the decrease associated with amitriptyline in another study.⁵⁰⁷ Refill utilization was improved⁵⁸⁵ and aspirin use increased

while pharmacists were being prompted to include aspirin use when dispensing medications for patients with diabetes.⁵⁸⁸

Other process changes. Murray and colleagues⁵⁷⁴ showed changes in workflow for pharmacists (more time interacting and problem solving) and who they interacted with (more time interacting with peers and physicians). Workflow was also changed in another study using a pharmacy information system.⁵⁷⁴ A commercial EMR system reduced the time to process and fill a refill request for HIV medication.⁵⁸⁵ Nilsson and colleagues⁵⁸⁶ showed that acute prescriptions were picked up more often for an e-Prescribing system compared with a paper-based system (91 percent vs. 85 percent, $p < 0.01$).

Summary

Few reports studied dispensing. Three of these nine studies were RCTs. All studies showed statistically significant improvements in process. External evidence suggests that the existing studies dealing with dispensing are weak and dated, with reports of currently used MMIT applications not being readily available.

Administering

Summary of the Findings for Process Changes

Nineteen studies measured changes in process associated with the administering phase of medication management (Appendix C, Evidence Table 4). All deal with nurses and either pharmacists or physicians. The technology is complex, integrated and often part of a complete package of a hospital information system or an EMR or EHR system. All studies but two^{12,592} were done using observational methods.

As noted in the dispensing section, CADTH produced a systematic review in 2009 on errors in dispensing and administering in hospitals.¹¹ This HTA report assessed the effectiveness and economic impact of MMIT applications designed to improve medication dispensing and administering. They found that the evidence on medication administering with MMIT was based on observational studies and that many of the studies were done on systems that have been updated or are no longer available. Many studies and descriptive papers that reported on medication administering and health IT, including Bar Code Medication Administration (BCMA) were reviewed for this report and were rejected. Most of the rejection decisions were because the medication administration system was stand-alone and not integrated with other MMIT applications. This nonintegration was especially true for older studies—most of the more recent studies show medication administering systems that are integrated. One good example of this integration is by Helmons and colleagues.⁵⁹³ Nonintegrated systems are not included in this report, as integration with other MMIT applications was an inclusion criterion.

Strengths and Limitations of the Evidence

One document was an RCT⁵⁹² and one was a cohort study.¹² All of the others were observational studies.^{34,438,439,581,583,589,593-602} Two were published in the late 1990s and 12 of the 19 were published since 2004.

General Study Characteristics

Participants. All but one study included nurses. Three studies included pharmacists,^{439,589,598} and four discussed physicians.^{465,592,593,596} The main focus of the study was medications or prescriptions,^{34,438,439,465,581,589,592-595,598,599,601,602} nurses^{597,600} and patients: infants¹² and those whose ages were unspecified.⁵⁹⁶ Medications were not limited to a specific drug or class of drugs except for one study of the need for antibiotics⁵⁹⁶ and one study of aspirin use.⁵⁹²

Location. All of the studies but one were set in hospitals: acute care or tertiary,^{439,465,581,596,598,600-602} critical care units,^{12,593-595} pediatric standalone hospitals,^{438,465} general hospitals,^{34,438} other specialty hospitals,^{465,581} and the emergency department.⁵⁹⁷ Pediatric hospitals or wards were often studied: neonatal ward and adult ICU,⁵⁹⁵ general pediatrics,⁴³⁸ and pediatric nephrology.⁴⁶⁵ One of the studies was done in an ambulatory setting,⁵⁹² and none were done in long-term care, community, or home settings.

Technology. The MMIT applications that were the focus of the administering phase of medication management were varied: automated drug dispensing system,⁵⁹⁹ BCMA systems,^{12,34,593,594,598,600,601} eMAR systems,^{12,34,439,583,589,601,602} CDSS,^{438,592,596} computerized unit dose drug dispensing system,⁴⁶⁵ CPOE,^{438,581,595,597} e-Prescribing,^{439,602} and a pharmacy information system.⁴³⁸

The MMIT systems that are integrated with these systems above are most likely to be hospital wide or pharmacy systems: a CPOE system,⁵⁹³ EHR and EMR systems,^{438,581,592,593,596,601} hospital information system,^{34,438,465,589,598} imaging systems,^{581,597} laboratory systems,^{581,597} eMAR,⁵⁹⁴ a formulary,⁵⁸⁹ and pharmacy information systems.^{12,34,439,583,595,597,599,601}

Outcomes

Errors. Thirteen studies evaluated administering errors. The issue of errors in administering drugs using MMIT is complex as many errors identified in MMIT systems are related to transcription and timing. These easily measured errors may be masking other more substantial errors. Eight studies had major endpoints that were found to be positive in reporting decreased errors.^{438,439,465,581,589,594,601,602} Another measured variances (differences between the order and administered medications) and found significant reductions after the introduction of a CPOE system integrated with the pharmacy and eMAR systems in a hospital.⁵⁹⁵ The relative risk reduction in many of the studies was high and often approximately 40 to 50 percent. Four studies had endpoints that were not found to show statistically significant improvements.^{34,583,593,598} Another hospital-based study showed increased errors, mostly related to a BCMA system,¹² because the BCMA system recorded issues such as timing of medication administering more accurately. The HTA report from CADTH also provided evidence that BCMA reduced errors in administering medications in hospitals.¹¹

Efficiency. Efficiency is also important in medication administering. Four of five studies that measured efficiencies showed improvements. One study showed that time from ordering to administering in a hospital setting decreased from 90 minutes before implementation of comprehensive MMIT systems to 11 minutes.⁴³⁸ Another article that measured time efficiencies had similar reductions (79 percent vs. 89 percent of medications were administered within 1 hour of ordering).⁴³⁹ An eMAR system reduced time from ordering to administering from 325 to 88

minutes.⁵⁸¹ Shirley and colleagues⁵⁹⁹ did not find a change in time to administering after implementation of an eMAR. No changes in time allocation were seen for direct patient care and medication administration after a BCMA system integration for hospital nurses.⁶⁰⁰ In contrast, Banet and colleagues described a system that integrated CPOE and eMAR and showed that nurses spent less time on paper documentation and searching for charts and more time on working with computers and charting in patient rooms with no changes in documentation time overall or time spent on direct patient care.⁵⁹⁷

Adherence to guidelines. One study with an anesthesia medication system had improvements in adherence to administering antibiotics during surgery.⁵⁹⁶ Shirley and colleagues⁵⁹⁹ found improved adherence to scheduled dosing. Persell and colleagues⁵⁹² identified no difference in self-reported aspirin use.

Other changes in process. Helmons and colleagues⁵⁹³ found no changes in error rates (they had few errors at baseline) but measured improved charting and labeling.

Summary. Although few studies evaluated administering with the use of MMIT, most of the 19 showed improvements, mostly in the realm of errors and efficiencies. Results were mixed with respect to whether the MMIT systems for drug administering altered the time nurses spent on various tasks.

Monitoring

Summary of the Findings for Process Changes

Medication monitoring can be defined as the process of assessing a patient's response to a medication and documenting its outcomes.⁶⁰³ Suboptimal medication monitoring describes a common pathway of systems failures that underlie monitoring errors and can be categorized as over, under, or inappropriate monitoring. Medication monitoring errors generally refer to one of three situations: inadequate **laboratory** evaluation of drug therapies, or a delayed or failed response by the clinician to **symptoms** (patient reported aspects of their disease or disorder), or to clinician observed or measured **signs** of the condition or of drug toxicity, or laboratory evidence of toxicity.⁶⁰⁴ Therefore, for the purposes of this report, we divided the health IT interventions designed to improve medication monitoring into studies that enhance **laboratory-, sign-, or symptom-based** medication monitoring. In the clinician and patient encounter the patient reports **symptoms** they are experiencing (e.g., fatigue, sudden weight gain, or dizziness) and the clinician observes or measures **signs** of the disease or disorder (e.g., blood pressure, heart rate, fever). Clinicians integrate information gained from assessments of symptoms, signs, and results of laboratory tests to determine disease status, often putting varying weights on the three aspects.

Previous systematic reviews provided information on the impact of health IT on medication monitoring.⁶⁰⁵⁻⁶⁰⁷ However, these systematic reviews are limited to a specific type of medication monitoring system (e.g., clinical event monitors), a single practice setting (e.g., ambulatory or acute care), or are more than 5 years old. This evidence report yielded a total of 47 articles describing health IT intervention designed to improve one or more change in process related to the medication monitoring phase in the acute, ambulatory, or long-term care settings (Appendix

C, Evidence Table 5).^{397,401,402,407,412,437,442,446,461,472,473,477,481,505,511,515,516,518-520,526-528,534,537,541,543,553-555,608-624}

For consistency, author-reported changes in process were selected. By definition, a study which showed statistically significant changes in at least half of its main endpoints was considered a positive study. Overall, 70 percent (33 of 47 studies) of the articles were rated as positive studies.^{397,401,402,407,412,437,461,472,473,477,505,515,516,527,528,537,541,554,555,608,610,612-623}

Study methods included 30 RCTs^{397,401,402,407,412,505,511,515,516,518-520,526-528,534,537,541,543,609-613,616-620,624} and 17 observational studies.^{437,442,446,461,472,473,477,481,553-555,608,614,615,621-623} Monitoring, along with CDSS, are the two areas that include the highest proportion of RCTs.

General Study Characteristics

Intervention targets. Nearly three-quarters (72 percent; 34 of 47) of the health IT medication monitoring interventions targeted physicians exclusively.^{397,401,402,407,412,437,442,446,461,472,481,505,511,515,520,527,528,534,543,553,554,609-611,613,615-620,622-624}

Eight of these studies targeted physicians along with other health care professionals,^{518,519,526,537,541,555,612,621} four targeted pharmacists,^{473,477,516,614} and one targeted nurses.⁶⁰⁸

Location. The overwhelming majority of health IT medication monitoring interventions studies (70 percent; 33 of 47) were conducted in an academic setting.^{397,401,402,407,412,437,442,446,461,472,473,477,481,505,515,518-520,527,534,554,608-611,614-619,621,622} Of those that

were conducted in an academic setting, 19 of these studies^{402,407,437,446,477,505,515,518,519,527,554,609-611,614-616,618,619} came from the following benchmark institutions: (1) Brigham and Women's Hospital/Partners Health Care, (2) LDS Hospital/Intermountain Health Care, (3) the Department of Veterans Affairs, and (4) the Regenstrief Institute.

The preponderance of studies (59 percent; 28 of 47) took place in the ambulatory care setting.^{472,505,511,515,516,518-520,526-528,534,537,541,543,553-555,609-613,616,617,619,620,624} Eighteen of the studies took place in the acute care,^{401,402,407,412,437,442,446,461,473,477,481,608,614,615,618,621-623} and one in the nursing home setting.³⁹⁷

Patient populations studied. The vast majority (n = 36) of the health IT interventions targeted the adult population.^{397,401,402,407,412,461,472,481,505,511,515,516,518-520,526-528,534,537,541,543,554,609-612,614-617,619-622,624}

Only four of the 44 articles were conducted in the pediatric population,^{437,442,446,553} and two targeted both adult and pediatric patients.^{477,613} Five studies did not explicitly mention the study patient population studied.^{473,555,608,618,623}

Type of medication monitoring. The majority (n = 29) of the health IT interventions focused on *laboratory-based medication monitoring*.^{397,401,402,407,412,442,461,472,473,477,481,511,515,516,527,528,534,537,541,543,555,609,611,612,614,615,619,620,623}

Five studies^{505,518,526,613,624} targeted *sign-based* (clinician observed or measured aspects of the disease process) *medication monitoring*. While three interventions focused on *symptom-based medication monitoring* (patient reported symptoms),^{520,608,621} ten studies^{437,446,519,553,554,610,616-618,622} provided a combination of *laboratory-*, *sign-*, or *symptom-based medication monitoring*.

A significant degree of overlap (n = 36) of health IT interventions that involved *laboratory-*, *sign-*, or *symptom-based medication monitoring* along with the prescribing phase of the medication use process existed.^{397,402,412,437,446,461,472,473,477,481,505,515,518,519,526-528,534,537,541,543,553-555,610,613-620,622-624}

Prescribing was most commonly associated with *laboratory-based medication monitoring* (n = 30),^{397,402,412,446,461,472,473,477,481,515,519,527,528,534,537,541,543,553-555,610,614-620,622,623} followed by *sign-based medication monitoring* (n = 15),^{437,446,505,518,519,526,553,554,610,613,616-618,622,624} and *symptom-based medication monitoring*.^{437,553} This overlap was most often a result of the evaluation of clinical practice guidelines, order sets, or both that contain prescribing and monitoring elements.

Drugs and diseases. Twenty-four of the health IT medication monitoring interventions studies dealt with chronic disease management such as asthma,^{446,553} asthma and chronic obstructive pulmonary disease,^{518,613} congestive heart failure and coronary artery disease,⁵¹⁹ deep venous thromboembolism,⁴⁰² depression,⁵²⁰ diabetes,^{537,610,619} diabetes and coronary artery disease,^{554,616} HIV,⁵²⁷ hyperlipidemia,^{515,528,534,541,543} hypertension,^{505,526,624} and multiple common chronic conditions.^{617,620,623} Sixteen studies addressed potentially nephrotoxic,^{397,615} hepatotoxic,²⁴¹ or cardiotoxic⁴⁷³ medications with a narrow therapeutic index,^{442,461,555,618} and certain laboratory and medication combinations.^{407,412,481,511,516,609,611,612} Four provided guidance about potentially inappropriate antibiotic management,^{401,477,614,622} and three provided information about pain management.^{437,608,621}

Technology. Almost all of the included studies regarding MMIT interventions (91 percent; 43 of 47)^{397,401,402,407,412,442,461,472,473,477,481,505,511,515,516,518-520,526-528,534,537,541,543,553,555,608,609,611-624} used a CDSS with alerts or reminders. Three studies used a CPOE system without alerts^{437,442,446} and one study involved the use of a personal health record (PHR).⁶¹⁰ Twelve of the studies used interruptive alerts to display and prompt the clinician for an immediate response while providing patient care.^{397,407,412,472,481,505,543,608,609,611,613,624}

Outcomes

As noted above, more than two-thirds (33 of 47) of the interventions were associated with a positive process outcome. A number of themes emerged from the variety of interventions that were conducted in various health care settings, using varying degrees of technological sophistication, and providing information to a number of health care professionals, as well as directly to patients.

By type of medication monitoring. The majority of the health IT interventions focused on *laboratory-based medication monitoring*.^{397,401,402,407,412,442,461,472,473,477,481,511,515,516,527,528,534,537,541,543,555,609,611,612,614,615,619,620,623} Of these 29 studies, 22^{397,401,402,407,412,461,472,473,477,515,516,527,528,537,541,555,612,614,615,619,620,623} or 76 percent of these interventions showed statistically significant changes in at least half of these main endpoints. Two^{505,613} of the five^{505,518,526,613,624} studies (40 percent) that targeted *sign-based medication monitoring* showed that greater than 50 percent of the process endpoints improved. Of the three interventions that focused on *symptom-based monitoring*,^{520,608,621} two^{608,621} resulted in statistically significant changes in at least half of their main process endpoints. Ten studies^{437,446,519,553,554,610,616-618,622} provided a combination of *laboratory-*, *sign-*, or *symptom-based monitoring*, and seven^{437,554,610,616-618,622} or 70 percent showed statistically significant changes in at least half of their main process endpoints.

By type of intervention. One of the most frequently reported types of intervention (n = 12) provided decision support to improve chronic disease management (i.e., prescribing, monitoring,

and clinical endpoints).^{505,520,526,528,537,541,543,554,555,610,616,624} The type of chronic diseases varied based on patient population, but included the management of asthma, chronic obstructive pulmonary disease, depression, diabetes, hyperlipidemia, and hypertension. Overall, 67 percent of these interventions resulted in a statistically significant change in at least half of its major endpoints. Another common intervention (n = 10) assessed the *adherence to guideline recommendations* for a variety of acute and chronic medical conditions including asthma, atrial fibrillation, coronary artery disease, cardiovascular disease, congestive heart failure, chronic obstructive pulmonary disease, depression, diabetes, glucose regulation in the ICU, pain management, and peripheral vascular disease.^{402,412,446,518,519,553,617,620,621,623} Overall, 60 percent of these interventions resulted in statistically significant change in at least half of its main endpoints.

Other common interventions (n = 8) included providing alerts and reminders to obtain laboratory testing for newly prescribed or chronically used medications.^{407,442,472,511,516,609,611,612} Overall, 50 percent of these interventions showed a statistically significant change in at least half of their main endpoints.

Seven studies targeted *changing prescribing behavior* by providing *laboratory-*, *sign-*, or *symptom-based monitoring* information for potentially nephrotoxic medications, medications for asthma and COPD, and hyperlipidemia.^{473,505,515,534,613,614,622} Overall, 86 percent of these interventions resulted in improvements in at least half of the major process changes reported as endpoints. Another metric commonly assessed was the *response time* to a variety of alerts (n = 7) including the management of narrow therapeutic index and potentially nephrotoxic medications, initiation of primary and secondary prevention, and time to pain assessment and management.^{461,477,481,527,615,618,619} Overall, 71 percent of these interventions showed statistically significant improvements in at least half of its main endpoints.

Finally, two interventions assessed *pain management* including error reassessment rate and patient controlled analgesia order set use.^{437,608} Overall, both of these interventions showed statistically significant changes in at least half of its main endpoints.

In our analysis, 70 percent (33 of 47 studies) of the included studies showed statistically significant changes in at least half of their main endpoints. Of these studies, the majority targeted physicians exclusively (n = 34), were conducted in academic institutions (n = 33), were developed for use in the ambulatory care setting (n = 28), focused on the adult population (n = 36), and provided CDSS with alerts or reminders to support chronic disease management (n = 12). When compared with *sign-* or *symptom-based medication monitoring*, *laboratory-based medication monitoring* studies were most likely (76 percent of the time) to be associated with a statistically significant change in at least half of its main endpoints. Moreover, these *laboratory-based medication monitoring* studies were conducted in a variety of health care settings including ambulatory, acute, and long-term care. The most successful types of studies focused on changing prescriber behavior, improving response time to generated alerts, and improving the diagnosis and management of chronic diseases.

Reconciliation, Discharge Summaries, and Education

Summary of the Findings for Process Changes

Reconciliation. Reconciliation of medications using MMIT is a complex process. Some of this stems from the complexity of medication management itself. Another issue is the challenge of interoperability of health IT across health care systems. The problem of medication

reconciliation is especially acute for patients who receive care across settings: from hospitals, specialists, and primary care—most often the elderly and those with multiple health challenges. Two review articles provide documentation of the difficulties of effective medication reconciliation using health IT and the lack of published evidence to support its value.^{625,626}

Four studies on medication reconciliation are included (Appendix C, Evidence Table 6).^{13,14,627,628} One was a cohort study⁶²⁸ and the others are quantitative observational. All were set in hospitals with the reconciliation done at discharge or transfer to another facility. One hospital was a Statepsychiatric hospital¹³ and the others were general hospitals.

One study was PDA-based,¹³ one was based on an e-MAR system,¹⁴ and the others were based on integrated systems: CDSS and COPE within an EMR,⁶²⁷ and an e-Prescribing system integrated with a pharmacy information system.⁶²⁸

All studies showed substantial improvement in agreement among records of medications provided by various clinicians involved in the care of the patients (Appendix C, Evidence Table 6). For example, one Dutch study showed improvements in agreement on prescriptions between the pharmacists and general practitioners with e-Prescribing compared with paper systems at discharge (31 percent vs. 49 percent) and at 10 days after discharge (33 percent vs. 53 percent).⁶²⁸ Grasso and colleagues¹³ showed a decrease in errors in the psychiatric hospital with the use of PDAs for reconciliation compared with paper summaries (rate of errors before PDAs was 22 percent compared with 8 percent after). Poole and colleagues¹⁴ also showed improvements in prescribing (more therapeutic drug duplications were identified and resolved with an automated discharge medication worksheet for physicians).

In summary, although few studies exist on MMIT for medication reconciliation, the four included showed substantial improvements in the ability to electronically reconcile medication lists and make the necessary adjustments resulting in reduced errors and better prescribing.

Education. Only one article targeted the education associated with MMIT systems and measured change in processes as their main endpoint (see Appendix C, Evidence Table 6).⁵³⁷ This RCT showed that combining patient education with submission of blood glucose levels to ambulatory care clinicians showed improvements in prescribing as well as improved hemoglobin A_{1c} levels. Most of the articles targeting educational aspects of medication management that measured changes in knowledge are covered in the section with intermediate outcomes.

Combined Phases of Medication Management

Summary of the Findings for Process Changes

Although some studies in this report assessed systems that covered the whole medication management process (five phases plus reconciliation and education), only one provided cross-phase study with changes in process. This observational study by Mahoney and colleagues⁴³⁸ took place in a U.S. pediatric hospital and an affiliated acute care hospital. The study started in 2002 and was completed in 2006 with publication in 2007. The hospitals included a full EMR system that incorporated CPOE, CDSS, and the pharmacy information system into one clinical information technology (hospital information system). All aspects of the medication management system were addressed electronically. An analysis of 1.4 million orders after implementation as compared with a similar number before implementation showed reductions in drug allergy violations, excessive doses, incomplete or unclear orders, and therapeutic duplication.

PDA's

Summary of the Findings for PDA's

We included 13 studies using PDA's.^{13,514,534,553,563,593,629-635} The studies often covered multiple medication management phases, such as prescribing (n = 7), order communication (n = 1), administering (n = 3), and monitoring phases (n = 6), as well as reconciliation (n = 1). Outcome measures focused on process and other intermediate measures, only two measured patient outcomes (blood glucose levels in both cases).^{630,631} Eight of the studies included a CDSS component.^{514,534,553,563,593,630,631,634} Two applications were tied to handheld BCMA units,^{593,635} and two were used for e-Prescribing.^{639,632} Most interventions targeted specific diseases such as diabetes,^{630,631} asthma,^{553,634} cancer,⁶³³ high blood pressure,⁵³⁴ psychiatric patients,¹³ or the use of certain classes of medications such as nonsteroidal anti-inflammatory drugs⁵¹⁴ and antibiotics.⁵⁶³ Two studies were qualitative,^{629,632} two mixed methods,^{633,635} five observational,^{13,553,563,593,631} and four were RCTs.^{514,630,633,634} Of the quantitative studies, five reported significant improvements as a result of the intervention^{13,563,630,631,634} and four reported no significant effects.^{514,534,553,593} An RCT of adherence to nonsteroidal anti-inflammatory drug prescribing guidelines in an ambulatory clinic showed stable levels of safe prescribing in the intervention group and a deterioration in the control group given PDA's without the guidelines.⁵¹⁴ Similarly, a PDA which provided physicians with Framingham scores and recommendations for patients at risk of high blood pressure, found no difference in levels of screening of patients and no effect on lipid management.⁵³⁴ A PDA-based CDSS for international asthma guidelines improved quality-of-life scores for patients and cost reductions.⁶³⁴ A crossover RCT of diabetic patient use of an insulin regimen dosage optimizer showed improvement in blood glucose levels during the phase when the advice was switched on.⁶³⁰

Intermediate Outcomes

Summary of the Findings

Articles measuring intermediate outcomes as their main endpoint were selected. We focused on the intermediate outcomes of: use; measures which were correlated with use (such as ease of use of the system, perceptions of users of the system, computer experience, etc.); knowledge, skills, and attitudes of the users; satisfaction; and usability (Table 10). Few hypothesis-driven studies with comparison groups assessed such intermediate outcomes as their main measure; 42 studies published in 44 articles were retrieved (Appendix C, Evidence Table 7). Only six studies were RCTs with quality scores from two to seven out of nine. The study results tended to show positive levels of satisfaction and use and measured a number of correlates of both to determine driving factors barriers, or both. Some negative impacts of systems on work processes were found.

Table 10. Intermediate outcomes across the phases for medication management

| Outcomes | P | OC | D | A | M | E | R |
|---|---|----|---|---|---|---|---|
| Use | 5 | 0 | 1 | 1 | 2 | 0 | 0 |
| Measures correlated with levels of use | 6 | 0 | 1 | 0 | 0 | 0 | 0 |
| Knowledge, Skills, and Attitudes of users (usually measured as perceptions) | 5 | 3 | 0 | 3 | 1 | 1 | 1 |
| Satisfaction | 9 | 1 | 1 | 2 | 0 | 0 | 1 |
| Usability | 3 | 0 | 0 | 0 | 0 | 0 | 0 |

Note some studies cross more than one phase.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Strengths and Limitations of the Evidence

Of the 43 included studies, 25 were observational, nine mixed methods or qualitative, six RCTs, and three cohort (Table 11). The RCTs rated two,^{636,637} four,⁶³⁸ six,⁶³⁹ and seven^{540,640} out of eight on the methods quality scale. The cohort studies scored three,⁶⁴¹ five,⁶⁴² and six⁶⁴³ out of nine. Studies of complex interventions often covered more than one phase of medication management.^{540,644-647}

Table 11. Study designs used in studies measuring intermediate outcomes across the phases for medication management

| Design | P | OC | D | A | M | E | R |
|------------------------------|----|----|---|---|---|---|---|
| RCT | 5 | 1 | 0 | 0 | 1 | 0 | 0 |
| Cohort | 1 | 0 | 0 | 0 | 2 | 0 | 0 |
| Observational | 17 | 2 | 2 | 5 | 1 | 1 | 1 |
| Qualitative or mixed methods | 4 | 1 | 0 | 3 | 1 | 0 | 0 |

Note some studies cross more than one phase. See Appendix C, Evidence Table 22 for references.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

RCT = Randomized controlled trial

General Study Characteristics

Study participants tended to be practicing clinicians (Table 12). Most of the studies were conducted in hospitals (n = 27) or primary care (n = 17), one in long-term care, and four in pharmacies, and assessed intermediate outcomes for health care staff. Twenty-two of the studies were performed in academic settings. Prescribing was the most commonly studied phase of medication management, but each other phase was represented. Three systems used hand-held devices. CDSS, e-Prescribing, and CPOE systems were most commonly studied. Most studies did not report on the proprietary nature of their systems, 17 studied commercial systems and seven were home grown. Many studies looked to correlate use of medication management systems with other factors. Only nine studies assessed intermediate outcomes for patients (Table 13).^{633,637,639,641,642,648-651}

Table 12. Clinician study participants in studies assessing intermediate outcomes across the phases of medication management

| Clinicians | P | OC | D | A | M | E | R |
|-----------------------------|---|----|---|---|---|---|---|
| Primary care physicians | 6 | 1 | 0 | 0 | 0 | 0 | 0 |
| Specialists | 6 | 0 | 0 | 0 | 0 | 0 | 0 |
| Hospitalists | 4 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other Physicians | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| Physicians undifferentiated | 9 | 0 | 0 | 0 | 1 | 0 | 1 |
| Nurses | 7 | 0 | 0 | 6 | 0 | 0 | 1 |
| Mid-level practitioners | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Pharmacists | 2 | 3 | 1 | 1 | 0 | 0 | 1 |
| Other health professionals | 4 | 0 | 0 | 1 | 0 | 0 | 0 |
| Hospital administrators | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Note some studies cross more than one phase.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Table 13. Patient study participants in studies assessing intermediate outcomes across the phases of medication management

| Patients | P | OC | D | A | M | E | R |
|------------------------|---|----|---|---|---|---|---|
| Infants (0 to 2 years) | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Children (2 to 12) | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Adolescents (13 to 18) | 2 | 0 | 0 | 0 | 0 | 0 | 1 |
| Adults (19 to 44) | 2 | 0 | 0 | 0 | 2 | 1 | 1 |
| Middle age (45 to 64) | 1 | 0 | 0 | 0 | 4 | 1 | 1 |
| Geriatric (65 plus) | 2 | 0 | 0 | 0 | 2 | 1 | 1 |
| Undifferentiated | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Note some studies cross more than one phase.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

Prescribing and ordering. Twenty-six studies looked at intermediate outcomes for interventions aimed at the prescribing phase (see Appendix C, Evidence Table 22). CDSS (n = 12), CPOE (n = 11), and e-Prescribing systems (n = 6) formed the bulk of the primary systems studied. Three studies assessed usability issues related to CPOE or CDSS.^{638,647,652} One study focused on the use of standards for medical history, formulary, and benefits.⁶⁵³ Satisfaction and correlates of satisfaction were measured in ten studies;^{636,637,644,645,654-659} use and measures correlated with use were studied in 11 studies.^{534,643,649,650,653,660-665} Pirnejad and colleagues⁶⁶⁶ used mixed methods to determine the impact of CPOE on the collaboration of nurses and physicians in hospitals. Glassman and colleagues⁶⁶⁷ looked at the impact of drug-drug interaction alerts on physician knowledge over time. Two studies assessed perceptions of technology on work.^{656,658} Participants were generally health care providers, located in either hospitals (16 studies), primary care (ten studies), or both, and one pharmacy.⁵⁴⁰ The majority were performed in the United States.

Order communication. Four studies looked at the order communication phase;^{540,645,668,669} three focussed on e-transfer of prescriptions,^{540,645,668} and all studied the perceptions of pharmacy staff as well as other stakeholders. Rupp and Warholak⁶⁴⁵ administered a survey and followed up with interviews of American chain community pharmacy staff to assess their attitudes towards e-Prescribing and recruited a sample of 1094 pharmacists, technicians, and interns from 276 pharmacies. Porteous and colleagues⁶⁶⁸ surveyed 494 patients, 145 general practitioners, and 148 pharmacists, and held interviews and focus groups to assess peoples' views regarding the upcoming implementation of e-Transfer of prescription information in the United Kingdom. On

the other hand, Kirking and Thomas⁶⁶⁹ performed a survey looking at pharmacists' attitudes towards computer technology used to detect and prevent adverse drug interactions, and correlated their findings with pharmacist computer use. Their sample included 218 pharmacists in Michigan using one of two pharmacy computer systems and a group of nonusers. Johnson and colleagues⁵⁴⁰ assessed the perceived usefulness of alerts and override comments appended to e-Prescriptions.

Dispensing. Two studies looked at dispensing. Chan⁶⁴⁶ looked at factors associated with the use of drug dispensing and eMAR systems in nursing homes by analyzing surveys of long-term care facilities. Rupp and Warholak⁶⁴⁵ assessed pharmacist personnel staff views of e-Prescribing.

Administering. Eight studies assessed technologies used at the administering phase. O'Morrow,⁶⁷⁰ Hurley,⁶⁷¹ Holden,^{672,673} Topps,⁶⁷⁴ and their colleagues assessed American nurses' attitudes and satisfaction toward bedside point-of-care BCMA technologies to verify drug administering. The usage patterns of BCMA verification in five medical departments of a Dutch hospital were tracked.⁶⁷⁵ The perceptions of the effects of a newly implemented CPOE in two groups of 211 Dutch nurses previously using different paper prescription systems were assessed by Niazkhani et al.⁶⁴⁴ Chan⁶⁴⁶ assessed factors associated with medication administration records use in nursing homes.

Monitoring. Four of the five monitoring studies focused on patient self-monitoring. Weingart and colleagues⁶⁴¹ measured the use of PatientSite, a patient internet portal, by 416 patients in three primary care practices to facilitate communication between physicians and patients regarding medication adherence and adverse effect rates. In an RCT involving 117 patients, Ross and colleagues⁶³⁹ provided patient online records for heart failure patients and assessed self-efficacy. McCann and colleagues⁶³³ performed an RCT on 112 cancer patients with a mobile phone application to monitor their chemotherapy toxicity symptoms compared with standard care and measured perceived benefits. Schmidt and colleagues⁶⁴² tested patient adherence using a telemonitoring intervention which included a beeping medication box integrated with the patient's EHR data. The final study assessed the usability of a CDSS Antibiotic Wizard in an ICU using an ergonomic survey tool to detect deficiencies in the system as viewed by 40 physicians.⁶⁴⁷

Reconciliation. One study evaluated patient and physician satisfaction and perceptions of a discharge reconciliation application.⁶⁵¹

Education. The study by Liu and colleagues⁶⁴⁸ is the only study focusing on the education aspect of medication management. Their study provides hospitalized patients in a Taiwanese hospital with a system that integrates their pharmacy, EMR, and CPOE information into an education tool to increase their knowledge of their medication regimens. Knowledge was assessed in pre-post surveys of 154 patients and they reported perceived knowledge gains.

Outcomes

Prescribing. Many studies measured use descriptively (e.g., presenting the percentage of time e-Prescribing was used for writing prescriptions), but did not meet our criteria for having comparison groups and being hypothesis-driven. From our included studies, use and measures

that were correlated with use were frequently measured. Rogers and colleagues⁶⁶⁰ found that usage rates increased following iterative changes made to a CDSS system based on feedback from users. Shannon and colleagues⁶⁴³ reported a significantly higher rate of e-Prescribing after emergency department physicians were allocated hand-held PDAs with e-Prescribing software. Access to medication history was more frequently used for patients with low socioeconomic status and a greater number of medications.⁶⁵⁰

Some studies found that ease of use and perceived usefulness relating to improved care and care processes of the MMIT applications were positively correlated with level of use.^{653,661-664} Kralewski and colleagues⁶⁶⁵ found that use of e-Prescribing systems was correlated with such cultural factors in primary care practices as trust, adaptability, and business orientation. Wang and colleagues⁶⁵³ found that a positive performance measure based on ease, efficiency, and care was correlated with nonuse of an e-Prescribing system incorporating standards for medication history, benefits, and formulary. Use of a CDSS in an Australian hospital was positively correlated with computer sophistication and access to laboratory data and negatively with years of experience.⁶⁶¹ Musser and Tchong⁶⁴⁰ measured preference and use of a graphic interface compared with a text-based interface for an anesthetic CPOE; clinicians used the graphic interface more often, but both interfaces had their proponents. A study assessing the frequency of use of three common pediatric order sets found differential use rates, with asthma order sets used significantly more often than both appendectomy and community-acquired pneumonia order sets.⁶⁴⁹

Usability. Three studies looked at usability and also included data on comparison groups.^{638,647,652} Rosenbloom and colleagues⁶³⁸ found that highly visible hyperlinks significantly increased the use of educational material and patient information. Rohrig⁶⁴⁷ and Li and colleagues⁶⁵² used usability testing to identify issues in a CDSS and CPOE system respectively. Their results were used to inform new iterations of their existing systems.

Satisfaction. Ten studies measured satisfaction as a main outcome. Satisfaction with various systems used by various health care providers tended to be positive.^{644,645,656-658} Satisfaction was lower in an intervention group of residents provided with CDSS within an e-Prescribing system in a small RCT, but they only used the system for 2.8 percent of their prescriptions.⁶³⁶ No difference in satisfaction levels were detected for patients or physicians using a discharge CPOE application compared with usual care.^{637,659}

Differences in satisfaction and perceptions of the systems were found between nurses and physicians;^{656,657} medical and surgical staff;^{654,657} and residents compared with physicians.⁶⁵⁴ Perceptions of the system impact on work were also found to be different among health care providers.^{656,658} Other factors correlated with satisfaction included computer sophistication, experience, training, system characteristics, and perceived improvements in care.⁶⁵⁴⁻⁶⁵⁷

Knowledge. Glassman and colleagues⁶⁶⁷ found no change in physician knowledge of selected drug-drug interactions over 2 years of using a CDSS with alerts in 97 American primary care physicians, although most preferred having the system.

Attitudes. Pirnejad and colleagues^{666,676} studied nurses' and physicians' attitudes to the impact of CPOE on the nurse-physician collaboration in the medication process. They found that the original paper-based system and the new CPOE system supported their work and collaboration

differently. The new system led to problems in the synchronization and feedback aspects of the joint medication care, leading to the recognition that new systems do not always directly replace the work entailed in old systems and that care processes can be negatively impacted.⁶⁶⁶

Attitudes toward MMIT often varied by groups of users. Junior students were more positive about CPOE than interns and residents.⁶⁷⁷ Similarly, using a diffusion of innovations model, Rahimi and colleagues⁶⁷⁸ found that a CPOE was perceived to work better for nurses than physicians (57 percent vs. 13 percent); further, more physicians felt that the system was not adapted to their practice and more would have liked a return to the old system, compared with the nurses.

Johnson and colleagues⁵⁴⁰ measured perceptions of pharmacists to appended alerts and override comments on e-prescriptions; they found some information (e.g., allergy alerts) more useful than others (e.g., insurance status).

Order communication. Of the four studies assessing the communication phase, outcomes assessed included only satisfaction⁶⁴⁵ and attitudes.^{540,668,669}

Rupp and Warholak⁶⁴⁵ found that chain community pharmacy personnel who dispensed e-Prescriptions were generally satisfied with e-Prescribing and rated e-Prescriptions more favorably than paper prescriptions on seven criteria related to safety, efficiency, effectiveness, communication, and relationships with patient and prescriber. They further produced 11 best practice recommendations to improve e-Prescribing in a community pharmacy setting. In the United Kingdom, before e-Transfer of prescription information being implemented, Porteous and colleagues⁶⁶⁸ found that various stakeholders viewed e-Transfer as a good idea (68 percent of patients [95 percent confidence interval (CI), 64 percent to 72 percent], 83 percent of general practitioners [95 percent CI, 77 percent to 89 percent], and 87 percent of community pharmacists [95 percent CI, 82 percent to 92 percent]). Concerns were expressed about security and sharing of confidential information. Benefits revolved around improved repeat prescription processes, convenience, and a greater role for pharmacists in medication management.

The potential for pharmacy systems to assist pharmacists in detecting adverse drug interactions by having greater access to patient information in the form of patient medication profiles was assessed by Kirking⁶⁶⁹ in a survey study asking pharmacists using two systems and a third group using no system how often they detected potential drug interactions and how often they contacted prescribers. Computer users reported an average of twice as many detected interactions per week (16.1 vs. 8.7, ns) and had significantly more contacts with prescribers per week (21.5 vs. 16, $p < 0.05$). The majority of the differences were the result of users of one of the unnamed computer systems, while the other groups had use rates similar to the noncomputer group.

Dispensing. One study suggests that drug dispensing and eMAR technologies were used more in nursing homes with higher occupancy rates; fewer metropolitan than rural homes using systems.⁶⁴⁶ Rupp and Warholak⁶⁴⁵ presented best practice recommendations for community pharmacies using e-Prescribing based on surveys showing satisfaction with e-Prescribing in community chain pharmacies.

Administering. Administering phase articles focused on nurses using BCMA systems⁶⁷⁰⁻⁶⁷⁵ or eMAR systems.⁶⁴⁶ O'Morrow⁶⁷⁰ found no differences in the attitudes of 17 nurses regarding patient care, charting, computer benefits, computer capability, computer characteristics, legal

issues, or management tools before and after implementation of a BCMA system. Hurley and colleagues,⁶⁷¹ on the other hand, found significant improvements on a satisfaction scale of 1,087 nurses after implementation of a similar system for efficacy, safety, care, and access factors. Holden et al.^{672,673} assessed nurses' perceptions and acceptance of BCMA; perceived ease of use and perceived usefulness; predicted satisfaction with the process before and after BCMA.⁶⁷³ In their second study, nurses' perceptions of the medication administering process changed with the implementation of BCMA compared with a control group; while perceived safety, accuracy and consistency in checking patient identification improved, ease of use, usefulness, and efficiency were perceived to decrease.⁶⁷² Topps and colleagues⁶⁷⁴ looked at nurse, pharmacist, and respiratory technicians' perceptions of BCMA before and after implementation; surveys after implementation showed that the staff felt that fewer medication errors occurred with a smoother administering of medication; they did, however, perceive that more time was spent administering medications, which took time away for other patient care. Overall, satisfaction and perceived benefits were improved in the study, by Niazkhani and colleagues,⁶⁴⁴ of nurses who went from two paper-based prescribing systems to a CPOE system. Perception of effects did depend on which previous paper system they were used to, and workflow support was perceived as worse by both groups.

Van Onzenoort and colleagues⁶⁷⁵ measured usage of bar code point-of-care systems by nurses and found that only 55 percent of 23,492 drug administrations were verified using the system; use depended on department, drug route, nurses available, nurse age, and timing of administering.

Monitoring. Most monitoring phase interventions were geared toward patients and showed positive effects on the intermediate outcomes of use, knowledge (self-efficacy), and satisfaction. PatientSite patient internet portal had a 48 percent response rate to index messages and a higher rate of ADE reporting via site (13 percent vs. 3 percent nonresponders, $p = 0.01$).⁶⁴¹ Ross and colleagues⁶³⁹ found that online records for heart failure patients improved self-efficacy (91 percent vs. 85 percent $p = 0.08$) and satisfaction. Chemotherapy patients using a mobile phone symptom system reported a number of benefits: better communication, better symptom management, and reassurance of physician access.⁶³³ Finally, patients who telemonitored their congestive heart failure issues consistently used a beeping medication box integrated with their EHR to increase adherence to their regimen.⁶⁴²

One study assessed usability; the Rohrig⁶⁴⁷ usability study of Antibiotic Wizard showed good usability. Physicians did report some weaknesses in the design of health IT which were to be used to inform future versions.

Reconciliation. A study of satisfaction with a reconciliation system found that patients reported satisfaction for self-reported perceptions of clear instructions on what medications to take, how much and how often the medications were to be taken, other instructions on taking the medication, potential side effects, and general understanding of the medications. Health care provider perceptions of satisfaction with reconciliation and instructions did not differ for five factors except for three factors reported by physician assistants and nurse practitioners. Physician assistants and nurse practitioners reported that patients had clearer instructions on discharge ($p = 0.01$); how much, how often, and when to take their medications at home ($p = 0.05$); and the medication discharge process was viewed as being sufficient for them as caregivers ($p = 0.0003$).⁶⁵¹

Education. Use of an integrated pharmaceutical system to provide information to patients to understand the pharmacological properties of their medications resulted in significantly improved patient knowledge after use of the system.⁶⁴⁸

Economic Outcomes

The introduction of health IT in the medication management process holds the promise of increasing efficiencies, improving quality of care, and reducing costs. However, even if these technologies are effective, they are expensive to implement and maintain and thus a review of the economic literature to determine cost-effectiveness and value for money for such interventions is warranted.

All studies passing the inclusion criteria that were considered to be cost or economics studies were reviewed and categorized into two groups based on the type of economic evaluation used in the analysis: (1) full economic evaluations; and (2) partial economic evaluations. A full economic evaluation is the comparative analysis of alternative courses of action in terms of **both** costs and consequences. Therefore, the economic evaluations which identify, measure, value, and compare the costs and consequences of the alternative being considered were further classified into one of the three categories: (1) cost-effectiveness analysis; (2) cost-utility analysis; and 3) cost-benefit analysis.⁶⁷⁹ The label, partial economic evaluation, indicates that the studies do not entirely fulfill both of the necessary conditions for a full economic evaluation (i.e., costs and consequences). However, cost analyses can provide useful information on ‘upfront’ costs compared with ‘downstream’ cost avoidance.⁶⁷⁹ For this reason, both full economic evaluations and cost analyses were included in this review. In each of these classifications, articles were further categorized by setting (i.e., hospital or community).

Descriptive information on the populations, interventions evaluated, the study year, perspective, and country of study were abstracted for each study. Data specific to the costs and effectiveness of each comparison were also abstracted and summarized in Appendix C, Evidence Tables 8a and 8b.

Full Economic Evaluations

Only five of the 31 (16 percent) economic articles reviewed conducted economic evaluations that provided information on the incremental costs and the incremental effects of an MMIT application. The following section reports the findings of five economic evaluations dealing with the use of CPOE (n = 2) and CDSS systems (n = 3) for improving prescribing practices for various conditions (Appendix C, Evidence Tables 8a).

Hospital. The potential economic consequences of implementing an eMAR system were estimated in a study using data from various literature sources.⁶⁸⁰ In a tertiary care hospital setting, the projected incremental effectiveness of the eMAR was 261 ADEs averted over the 10-year time horizon compared with the standard paper ordering approach. Given that the incremental cost of the new electronic medication ordering system was USD\$3.3 million during that same period, the incremental cost-effectiveness ratio was USD\$12,700 per ADE averted.

A 1-year RCT in a hospital family medicine center evaluated the effect of three reminder systems on compliance with tetanus vaccination.⁵³⁰ A computer-generated physician reminder system was found to cost \$0.43 per additional vaccination recorded compared with usual care.

The telephone reminders to the patients cost \$5.43 per additional vaccination, while the mailed letter reminder to the patients to recommend tetanus vaccination was \$6.05 versus standard care.

Community. A group of Norwegian researchers⁵¹⁰ conducted a cost-effectiveness analysis alongside an RCT involving 146 general practices from two separate geographical areas. The objective of the evaluation was to compare the costs and effects of a multifaceted intervention, including computerized reminders to physicians, aimed at improving prescribing of antihypertensive and cholesterol-lowering drugs compared with the passive dissemination of guidelines. The cost per additional patient started on a thiazide rather than another antihypertensive agent in the intervention group was compared with usual care. Over the 1-year study period, the authors calculated that the incremental cost-effectiveness ratio of the intervention was USD\$454 per additional patient started on thiazides. It was found that reduced drug expenditures based on increased use of thiazides did not outweigh the costs of the intervention. The authors commented that if the effect was sustained for a second year, the intervention would have been expected to lead to savings.

A Spanish study published in 2005⁶³⁴ evaluated the cost-effectiveness of a CDSS designed to promote guidelines for the treatment of asthma. Over the 1-year study period, the authors found that from a societal perspective, the intervention dominated standard care (i.e., less costly and more effective). From the health care payer perspective, the incremental cost-effectiveness ratio was €1 per percentage point reduction in the St. Georges Respiratory Questionnaire.

Setting not stated. Using information obtained from a systematic review of the literature, Karnon et al.⁶⁸¹ developed a decision tree model to estimate the net benefits of three interventions aimed at reducing medication errors (i.e., CPOE, ward pharmacists, and bar coding), either through prevention or detection. Based on estimated quality of life utility decrements associated with experiencing a preventable ADE, it was concluded that the CPOE had a mean net benefit of GBP £31.5 million, ward pharmacists of GBP £27.25 million, and bar-coding of GBP £13.1 million over a 5-year time horizon with the intervention and maintenance costs included in their model. It was noted that the monetary value of lost health needed to be included for the interventions to have a high probability of producing positive net benefits.

Partial Economic Evaluations

Most of the economic literature reported the results of partial evaluations (26 of 31 studies, 84 percent). All of these evaluations took the form of cost analyses. In other words, the costs of the alternatives were examined separately and the effectiveness, efficacy, or both measures were not used in the analyses, which results in an inability to answer efficiency questions about an intervention.

Hospital. A computerized ADE surveillance system was used to help identify and prevent specific types of ADEs in patients in hospitals.⁶⁸² The authors compared the length of stay in hospital of patients incurring an ADE with a historical control group of inpatients who did not have ADEs, and showed that the average length of stay for patients with severe ADEs was 20 days, 13 days for patients with moderate ADEs, and five days for those with no ADEs. This translated into a cost of USD\$38,007 for patients with severe ADEs compared with USD\$22,474 for patients with moderate and USD\$6,320 for patients with no ADEs. Given this significant difference in the length and cost of hospitalization between patients with severe and moderate

ADEs, the authors concluded that this *suggests* that the prevention and reduction of ADEs could reduce the length and cost of hospitalization for certain patients. However, it is important to acknowledge that the cases were not matched for disease severity and that no direct cost analysis was made of the ADEs prevented by the system compared with before the implementation of the system.

The same author measured the effect of a CDSS aimed at improving the use of and reducing the cost of antibiotics in four separate studies. The first was conducted in an academic, tertiary, private hospital and the average cost for 24 hours of antibiotic therapy recommended by the CDSS was USD\$10.85 less per patient than what was actually prescribed by physicians.⁴⁰⁹ The same CDSS was evaluated in two studies that took place in a 12-bed shock/trauma/respiratory ICU. The 7-month pilot study revealed a mean reduction in the cost of antibiotics of USD\$87.03 per patient compared with the preintervention period.⁶⁸³ The other ICU study was 12 months in duration and the mean cost of antibiotics for the computer regimen followed, regimen overridden, and no CDSS, respectively was USD\$102 compared with USD\$340 and \$427, while the cost of hospitalization was USD\$26,315 compared with USD\$35,283 and USD\$44,865.⁴⁷⁵ Finally, an antibiotic-dose monitor was incorporated into the CDSS to check the renal function of patients to identify those who were potentially receiving excessive dosages of antibiotics.⁶¹⁴ The patients in the intervention group received fewer mean doses of study antibiotics at a lower average cost (USD\$80.62) than patients during the preintervention period (USD\$92.96) of this 12-month study. If this reduction of USD\$12.34 per patient is summed for all 4,483 patients in the intervention period, this would result in a total decrease in cost of more than USD\$55,000 a year.

Another CDSS by Barrenfanger and colleagues,⁶⁸⁴ designed to improve antibiotic prescribing by electronically notifying the pharmacist of potential problems with a patient's antimicrobial therapy, was introduced in a 450-bed community teaching hospital and evaluated over a 5-month time period. The study compared patients whose microbiologic data were processed in the normal manual manner in the pharmacy to patients whose microbiological data were processed using the computer software. The study patients were matched by diagnosis related groups to patients in the control group. Additionally, the control group patients were adjusted for severity to make the groups more comparable. The study group had an average total standard cost of USD\$13,294 per patient; the severity adjusted control group had an average total standard cost of USD\$16,106 per patient, a decrease of USD\$2,812 per patient in the study group. By using these severity adjusted data, the estimated variable cost savings annually from the improvement of interventions is USD\$2,932,000 (2,000 inpatients for whom susceptibility testing is done multiplied by \$1,466). If the list price of the CDSS (USD\$44,500) was subtracted from the expected annual cost savings from the use of the program to improve interventions (USD\$2,932,000), the resulting savings (USD\$2,887,500) was still substantial in the first year.

A 3-month RCT was designed to evaluate the effect of a CDSS for the management of antimicrobial utilization in a 648-bed tertiary care academic hospital.⁴⁰¹ Antimicrobial utilization was managed by an existing antimicrobial management team using the system in the intervention arm and without the system in the control arm. The Web-based system was developed to alert the AMT of potentially inadequate antimicrobial therapy (a "back-end" or postprescription review). Expenditures for antimicrobial drugs were USD\$285,812 for the intervention group and USD\$370,006 in the control arm, for a savings of USD\$84,194 (23 percent) overall or \$37.64 per patient.

An antiinfective decision support tool, designed specifically for a pediatric population, was introduced in a 26-bed ICU in an academic hospital.⁴⁶⁹ During the 6-month period before CDSS installation, all patient care orders from the physicians were handwritten. The study found no difference in hospital costs in the period before CDSS installation (USD\$28,257.67) compared with the time after CDSS installation (USD\$25,032.11) or in antiinfective costs per patient (USD\$274.79 in the control group compared with USD\$289.60 in the intervention group).

An evaluation of a CDSS on appropriate antibiotic treatment used a cohort study followed by a multicentre, cluster RCT.³⁹⁹ The cohort study compared the advice of the CDSS with physician performance with respect to appropriate empirical antibiotic treatment and costs. The RCT compared hospital wards using the CDSS compared with antibiotic monitoring without the CDSS. In the cohort study, all cost components, except those related to expected adverse events, were significantly lower for the treatments suggested by the CDSS compared with those used by physicians. Total antibiotic costs were €289 lower per patient for CDSS, a relative decrease of 48 percent. In the RCT, the use of the CDSS resulted in significantly lower antibiotic costs in intervention versus control wards, the difference originating from lower ecological costs in intervention wards in Israel and Italy. Direct antibiotic costs, as well as costs incurred by observed adverse events, were similar.

A Canadian study in an orthopedic institution assessed the safety and potential cost savings of a computerized, laboratory-based program (i.e., CPOE and CDSS) to manage inpatient warfarin therapy after major joint arthroplasty.⁶⁸⁵ The authors estimated that the potential savings per patient of CAD\$5.50 per day was due to a reduction in nursing time, for a total annual figure of CAD\$55,836. It is important to note that the cost estimates and potential cost savings are speculative and are meant to be illustrative and not conclusive in nature.

A computerized order set within an CPOE was designed to manage pediatric inpatients with asthma.⁴⁴⁶ A before-after study of the system found no significant difference in the total inpatient costs among the groups before and after intervention. The hospital charges were USD\$3,567 and USD\$3,759, while the pharmacy charges were USD\$373 and USD\$429 in the groups before and after intervention, respectively.

The costs associated with the implementation of a CPOE and CDSS system over 10 years (1993 to 2002) were measured in a 720-adult bed, tertiary care academic hospital.⁶⁸⁶ Using data on the reductions in items such as ADEs, drug costs, and laboratory test usage, it was estimated that the system saved the hospital USD\$28.5 million over the 10-year period, even after including the capital and operational costs of USD\$11.8 million. The authors stated that it took over 5 years to realize a net benefit and over 7 years to realize an operating budget benefit.

Chertow et al.⁴⁶⁸ studied the effect of adding a CDSS to an existing CPOE for prescribing drugs to patients with renal insufficiency in a hospital setting. The authors measured the difference between the intervention and control groups in hospital and pharmacy costs and found no differences between the groups (USD\$4,881 compared with USD\$4,968 in total costs for the intervention and the control groups, respectively).

An evaluation of the introduction of a CPOE and eMAR system on the delivery of health care in an academic health system was done using a before-after design.⁵⁸¹ Based on total costs per admission, no significant difference was seen in any of the U.S. hospitals in the system.

A cost analysis of the implementation of an expensive CPOE (i.e., total capital cost of implementation was USD\$2.9 million and operating costs were USD\$2.3 million) in the management of surgical patients in an academic, multispecialty hospital was done by Stone et al.⁴¹⁹ Based on the data from 6 months before and 6 months after the intervention, a

redistribution of workload was found. The personnel changes resulted in a savings of USD\$445,500. The authors also noted that because of considerable gains in efficiencies (e.g., time necessary to have orders accessible to nursing, radiology, and laboratory), this implementation would likely result in long-term cost savings and improved quality of care.

An RCT done in 1993 assessed the effects of a network of microcomputer workstations for writing all inpatient orders (i.e., CPOE) on health care resource utilization.⁶⁸⁷ The overall aim of the CPOE was to encourage cost-effective ordering and to reduce costs. Using the costs associated with inpatient charges (i.e., bed, tests, and drugs), it was determined that total charges per admission were significantly less (USD\$887) for the intervention teams than for the control teams, with similar differences in all types of charges. The authors claim that if these effects were extrapolated to all medicine service admissions at that hospital, the projected savings in charges per year would be \$3 million in 1993 U.S. dollars. It was noted that the workstation network hardware costs were approximately USD\$20,000 per ward, with additional costs for installation and maintenance.

In two separate RCTs, Tierney et al. evaluated the effect of a CDSS that provided guidelines for the treatment of patients with ischemic heart disease or chronic heart failure⁵¹⁹ and patients with asthma or chronic obstructive pulmonary disease.⁵¹⁸ In both studies, care recommendations were displayed electronically to either physicians, pharmacists, or both physicians and pharmacists, compared with no care recommendations. In the heart disease study, the patients in the group receiving only the physician intervention had significantly elevated total health care charges (physician only: USD\$6,302, compared with pharmacist only: USD\$7,387, compared with physician and pharmacist: USD\$7,639, compared with control: USD\$7,025). In the asthma and chronic obstructive lung disease study, the authors found no difference in total costs (i.e., total inpatient and outpatient charges) across groups (physician only: USD\$8,006, compared with pharmacist only: USD\$5,333, compared with physician and pharmacist: USD\$5,652, compared with control: USD\$5,800).

A recent publication by Pointek and colleagues⁶⁸⁸ measured the impact of an ADE alert system on cost and quality outcomes in seven community hospitals within a health network. The ADE alerts were triggered in real time, which enabled immediate pharmacy intervention. The results showed a statistically significant decrease in average pharmacy department costs per patient (USD\$867 versus USD\$826, $p < 0.001$) from before to after implementation. In contrast, the external control group had a significant increase in pharmacy department costs (USD\$734 versus USD\$797, $p = 0.029$). Drug costs decreased significantly from baseline (USD\$360 versus USD\$337, $p < 0.001$) in the study group. Conversely, there were significant increases in drug costs in the external control group (USD\$401 versus USD\$429, $p = 0.029$). The authors applied the observed percentage of cost decrease from baseline exhibited by the study group to both the internal and external control groups' results and found that this yielded a combined pharmacy department cost savings estimate in excess of USD\$11 million. It was noted that these savings coincided with only modest quality improvements in projected mortality rates and length of stay. An important limitation in this study is that it did not compare ADE rates before and after implementation of the system.

Community. McMullin and colleagues^{689,690} published two papers that evaluated the impact of a CDSS on prescription costs on a range of medications used in primary care. The first study was a retrospective cohort study using pharmacy claims data, which found that the average cost per new and refilled prescriptions was USD\$4.99 lower in the intervention group, with the 6-month

savings being USD\$3,450 per clinician. A 6-month extension of this study showed a 12-month savings on new prescriptions of USD\$109,897.

A cluster, unblinded, pragmatic (i.e., real world) RCT was conducted in a routine clinical setting, to assess the cost and effectiveness of a CDSS based on recommendations of the European Society of Cardiology and other societies for hypercholesterolemia management in comparison with usual care for patients with hypercholesterolaemia.⁵²⁸ The total direct costs of hypercholesterolaemia management (i.e., physician visits, laboratory analyses, and the lipid-lowering drugs prescribed) for the intervention and control groups were calculated. The impact on total costs was markedly different in the two groups: €264,658 in the usual care group and €170,061 in the intervention group.

Ornstein et al.⁶⁹¹ set out to measure the impact of displaying prescription cost information in a computer-based patient record system at the time of prescribing on reducing drug costs by family physicians. When compared with a 6-month period where cost information was not displayed, it was concluded that no impact was found on overall drug costs to patients that could be related to the intervention. The mean cost per prescription in the control period was USD\$21.83, and in the intervention period was USD\$22.03.

Weingart et al.⁶⁹² designed an empirical study to understand the potential benefits of medication safety alerts generated by an e-Prescribing system in ambulatory care. Using a modified Delphi technique and data on 1.8 million prescriptions, the authors estimated that e-Prescribing alerts possibly averted 133 to 846 ADEs. These alerts could have avoided health care resource utilization in a number of areas (e.g., hospitalizations, emergency room visits), for a total savings to the system of USD\$141,012 to USD\$1,012,386. An expert panel reviewed a sample of common drug interaction alerts, estimating the likelihood and severity of ADEs associated with each alert, the likely injury to the patient and the health care resource utilization required to address each ADE. The analysis estimated that the cost savings due to the e-Prescribing by using third-party-payer and publically available information was USD\$402,619 (inter quartile range [IQR] \$141,012-\$1,012,386) with an average cost savings per clinician of USD\$173 (IQR \$61-\$436).

Community and hospital. One group of researchers developed a CDSS that used the clinical information contained in administration claims data from physicians, hospitals, pharmacies, and laboratories to identify common errors in care and departures from widely accepted clinical guidelines.⁶²⁰ This differs from the other CDSSs discussed in this section, in that the CDSS was not deployed within a hospital setting or within an integrated delivery system in which EHR systems provided the backbone of clinical information. The authors conducted a 12-month RCT to test the hypothesis that the claims-driven CDSS could increase compliance with evidence-based practices and effect improvements in patient outcomes as measured by decreased hospitalization and attendant cost. The sentinel system was designed as a rule-based artificial intelligence engine combined with an automatic message generator that conveys clinical recommendations and supporting literature to treating physicians. Nine hundred and eight clinical recommendations were issued to the intervention group. Among those in both groups who triggered recommendations, there were 19 percent fewer hospital admissions in the intervention group compared with the control group ($p < 0.001$). Charges among those whose recommendations were communicated were USD\$77.91 per member per month lower and paid claims were USD\$68.08 per member per month lower than among controls compared with the baseline values ($p = 0.003$ for both). According to the paper, the intervention cost USD\$1.00 per

member per month to deploy and was associated with lower paid claims of USD\$8.07 per member per month in the intervention group compared with controls, suggesting an eightfold return on investment from the payer perspective. However, it is important to note that this study was not intended as a formal cost-effectiveness analysis or cost savings analysis in that they did not directly measure costs at the patient or caregiver level, nor did they consider noneconomic costs or benefits.

An extension of this analysis was published 3 years later.⁵⁰⁴ This study used data from two additional years to analyze the effect of the intervention on resource utilization. This evaluation showed that the intervention reduced the average total charges (i.e., billing, pharmacy, and laboratory data) in the study group by USD\$24.77 per member per month compared with the group without the CDSS.

Economics Summary

Most of the studies (84 percent) reviewed that evaluated the economics of MMIT would not be considered full economic evaluations. Full economic evaluation studies measure the cost per successful patient outcome over time, whereas cost analyses measure only the costs of the alternatives examined. Cost analyses can provide useful information on ‘upfront’ costs compared with ‘downstream’ cost avoidance but an ideal economic evaluation would explicitly measure all direct health care costs (e.g., capital costs, health professional’s time) and direct nonhealth care costs (e.g., home care services, transportation), as well as indirect costs (e.g., productivity gains or losses related to illness or death by the patients and caregivers) that could be affected by the intervention of interest. It is important to be aware that the greatest costs of these health ITs are associated with the purchase of new software (capital outlay) to add to preexisting EMR systems, as well as implementation costs (e.g., management, clinical team involvement, training costs, maintenance costs), which were not included in the cost side of the economic evaluation in most studies. Additionally, the full enumeration of the total costs needs to be synthesized with the consequences or outcomes of the intervention (i.e., cost-effectiveness analysis, cost-utility analysis, and cost-benefit analysis). The effectiveness of any given system is dependent on the system’s design, implementation, the users of the system, and the setting into which the system is being introduced. Adoption of newer technologies needs to be based on formal evaluation of whether the additional health benefit (effectiveness) is worth the additional cost. Given the tension between the clinical benefits of integrated CPOE and CDSS systems and the high upfront costs, decisionmakers deciding whether to implement them need to better understand how and when financial benefits of such systems accrue (e.g., short-term compared with long-term benefits). These types of analyses are important for well-informed decisionmaking.

In summary, a few of the studies reviewed found that health IT interventions may offer cost advantages despite their increased acquisition costs compared with care provided without the health IT. However, given the uncertainty that surrounds the cost and outcomes data, and limited study designs available in the literature, it is difficult to reach any definitive conclusion as to whether the additional costs and benefits represent value for money. It is necessary that sophisticated concurrent prospective economic evaluations be conducted in the real world to address whether health IT interventions in the medication management process are actually cost-effective.

Clinical Outcomes

Summary of the Findings

Among the clinical outcomes assessed in 76 articles (Table 14), 54 percent reported significant benefits (Table 15). Studies that used monitoring approaches to identify and intervene with patients with actual problems (e.g., excessive blood pressure, increase in creatinine after being placed on a nephrotoxic drug) or needed care (e.g., hemoglobin A_{1c} monitoring) appear to be more effective than CDSS approaches that identify theoretical problems (potential for adverse drug events). The effectiveness of monitoring interventions in ambulatory care is enhanced (or only effective) if patients are also sent reminders and decision support recommendations.

Highly targeted interventions, focused on specific problems that provide problem-related specific interventions appear to be more effective than a more diffusely focused CDSS integrated with a CPOE system (e.g., nonpatient-specific guidelines for cardiovascular risk reduction).

Many studies have evaluated CDSS tools for improving the effectiveness of anticoagulants (proportion of days in therapeutic anticoagulant range) and improving the choice, route, duration of antibiotics, and reducing ADEs related to antibiotic use and most are successful.

Studies that have been successful in improving patient outcomes target high risk and vulnerable populations who have poor disease control,^{515,610,624,693} lack sufficient access to health care providers to manage their condition,⁴⁰⁸ or subpopulations with sufficient economic resources to respond to the CDSS intervention.⁶⁹⁴

While high risk groups have the potential to show the greatest benefits of IT, one study, which implemented a CPOE (prescribing, dispensing, and order communication system) in a children's hospital, reported substantial harm—a 270 percent relative increase in mortality after CPOE was implemented (2.8 percent vs. 6.6 percent unadjusted, adjusted OR 3.81, 95 percent CI 1.94 to 5.55).¹⁵ This before-after study and its methods have been debated¹⁷ and its conclusions contested. However, the increase in mortality they found provides important lessons about CPOE implementation, particularly in settings which include high-risk patients. Critically ill patients are most likely to benefit from IT but also most likely to be affected by dysfunctional technology and implementation strategies because delays in definitive treatment can increase the risk of mortality. As other groups have shown that CPOE systems either have no effect or a nonsignificant reduction in mortality in children's hospitals,¹⁶ the disparity in findings likely relates to the extent to which both the technologies and implementation strategies have disrupted or delayed critical activities in the clinical setting, and demanded additional time for order-entry from clinical staff.

Two studies that implemented computerized decision support CDSS drug use increased mortality,¹⁵ and length of stay.¹⁸ Both studies lacked sufficient power to conduct a valid assessment.

Table 14. Research design for studies across the phases of medication management and education and reconciliation that address clinical outcomes as their main outcomes

| Design | P | OC | D | A | M | E | R |
|-------------------------|----|----|---|---|----|---|---|
| RCT | 21 | 0 | 0 | 1 | 21 | 1 | 1 |
| Cohort and case control | 7 | 0 | 0 | 0 | 6 | 0 | 0 |
| Observational | 27 | 2 | 1 | 2 | 13 | 0 | 0 |
| Qualitative | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 55 | 2 | 1 | 3 | 40 | 1 | 1 |

Note some studies cross more than one phase. See Appendix C, Evidence Table 16 for references to the included articles in each cell.

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

RCT = Randomized Controlled Trial

Table 15. Summary of the number of studies reporting statistically significant differences in clinical primary endpoints between study groups for hospital and ambulatory based studies

| Clinical Endpoints | RCTs | Cohort and Case-Control Studies | Observational Studies |
|--|------------|---------------------------------|-------------------------------|
| Mortality (0 of 7 showed benefit) | | | |
| Hospital Studies | 1 = | 1 = | 4 = 1 ⁻¹⁵ |
| Ambulatory | 0 | 0 | 0 |
| Quality of Life (1 of 5 showed benefit) | | | |
| Hospital Studies | 0 | 0 | 0 |
| Ambulatory | 1+ 3 = | 1 = | |
| Length of stay (7 of 14 showed benefit) | | | |
| Hospital Studies | 1 = | 1+ 1 = | 6+ 4 = 1 ⁻¹⁸ |
| Ambulatory | 0 | 0 | 0 |
| Adverse Drug Events (8 of 10 showed benefit) | | | |
| Hospital Studies | 1+ 1 = | 0 | 6+ 1 = |
| Ambulatory | 0 | 1+ | 0 |
| Physiological (18 of 32 showed benefit) | | | |
| Hospital Studies | 0 | 4+ | 5+ 2 = |
| Ambulatory | 6+ 11 = | 1+ 1 = | 2+ |
| Other adverse events (e.g., readmissions, hospitalizations, etc.) (six of 16 showed benefit) | | | |
| Hospital Studies | 1+ 3 = | 1+ 3 = | 2+ 2 = |
| Ambulatory | 2+ 1 = | 0 | 1 = |

+ indicates that half or more of the main endpoints were shown to be positively statistically significant.

= indicates that at least half of the main endpoints were statistically not significant.

- indicates that half or more of the main endpoints were shown to be negatively statistically significant.

RCT Randomized controlled trial

Strengths and Limitations of the Evidence

Overall, 28 of 76 (37 percent) studies assessing clinical endpoints were RCTs, and the mean quality rating was 4.4 out of 9 (range two to seven).^{401-403,407,408,515,518-520,524,526-}

^{528,537,541,543,610,620,624,630,634,637,695-700} Low ratings are because most RCTs of health IT cannot be blinded, and the majority are cluster RCTs, where equivalence in the distribution of measured and unmeasured confounders (clinician and patient characteristics) cannot be assured. Statistical

adjustment for differences in the intervention and control groups has not been conventionally advocated even though it is likely required for unbiased comparisons.

The remaining studies were cohort, case control or observational; the majority were before-after studies or variants of this approach. Typically, in the before-after variant design, three time periods were assessed. Preintervention outcomes were compared with outcomes evaluated at two time periods of after implementation intervention. These comparisons sought to assess changes in care and the care processes associated with the interventions that were subsequently introduced. Only one study was a true time-series.⁴⁸² In most of the before-after studies, no adjustment was done for differences in patient mix or cointerventions in the time periods with and without the intervention. Unless a systematic trend for changes in the patient population mix was shown, this problem may have minimal effect on the reported results. The only exception is with length of stay, where well-documented trends in reductions in length of stay due to many factors unrelated to IT interventions are shown. For these outcomes, the positive benefits in reductions of length of stay shown in nine of 15 studies that measured this outcome are likely overestimated.

While the absence of a contemporaneous comparable control group is a problem with all before-after studies, the creation of control groups by comparing intervention patients to those that do not participate, or do not have a problem, to those that do is fundamentally far more likely to introduce major bias in the comparison (e.g., comparing patients with alerts to no alerts,¹⁸ pharmacists volunteering to provide the intervention compared with those that do not volunteer,⁶⁹⁴ and other similar problems^{642,701}). The direction of the bias will depend on the study. Volunteers in any study tend to have better outcomes than nonvolunteers, and selecting patients with problems compared with those that do not will ensure that at least both will regress to the mean—people with problems get better and those with no problems get worse, resulting in an overestimation of the effect of most interventions.

Many of the observational studies suffered from selecting an outcome that was distantly or only marginally related to the intervention. Almost all of the studies that measured quality-of-life, length of stay, and all cause ADEs were examples of this problem. Gurwitz and colleagues⁶⁹⁷ were able to show that only one-third of ADEs could have been prevented by the CDSS alerts that were provided. Moreover, in a substantial proportion of negative studies, minimal adoption was evident. The clinicians failed to adjust therapy or treatment to match the recommendations, and thus it was not surprising to find that the interventions had no effect on outcomes. Finally, the rate of some outcomes such as readmission, mortality, and nosocomial infections were too low to detect clinically meaningful differences if they had existed.

General Study Characteristics

A total of 76 studies assessed improvements in clinical endpoints or reduction in adverse events (Appendix C, Evidence Table 9).^{15,16,18,401-}

403,407,408,423,425,430,437,446,452,453,455,459,467,468,475,482,489,501,515,518-520,524,526-

528,537,541,543,545,550,555,581,610,614,615,620,624,630,631,634,637,641,642,682-685,688,693-714

Prescribing and monitoring are the phases that were well-studied with respect to clinical outcomes (95 percent of all studies). Forty included the monitoring phase, only two evaluated clinical outcomes associated with order communication,^{15,581} three studied drug administering^{581,630,693} and one each looked at dispensing,¹⁵ reconciliation,⁶⁹⁵ and a cell phone-based diabetes management program for educational purposes.⁵³⁷ A total of 85 different endpoints were assessed for different aspects of MMIT (Table 14).

Outcomes

Prescribing. Clinical outcomes have always been the most important and the most difficult to measure and study in health IT applications. The studies must be done in clinical settings which are complex and nonstandard. The health IT must be held constant to fit the traditional model of clinical trials and this does not reflect the reality of clinical practice and technology use. It is also difficult to ascertain if a technology can affect clinical outcomes—drugs, surgeries, and other similar interventions are easier to tie to outcomes. The health IT is a “long distance” from actual clinical care with many steps and factors involved and the latency between exposure to the health IT and outcome.

Many studies have been done on the effect of health IT on prescribing. Consequently, many systematic reviews have addressed the effects of these applications on clinical outcomes. Two Cochrane reviews have been done. One addresses onscreen point-of-care computer reminders on outcomes of clinical importance. The review by Shojania and colleagues⁷¹⁵ found some clinical improvements across studies with blood pressure (being reduced by a mean of 1.0 mmHg). Durieux and colleagues⁷¹⁶ showed small improvements in time to therapeutic stabilization, risk of toxic drug levels and length of hospital stay (mean decrease of 0.4 days). Another eight reviews show similar findings for clinical outcomes: more changes in process and some limited and rather small improvements in clinical outcomes: alerts and prompts to improve prescribing behaviors (five studies, three showed statistically significant improvements),⁷¹⁷ CDSSs to improve prescribing in older adults (two studies, mixed outcomes),⁷¹⁸ CDSSs on medication safety (five CPOE and seven CDSS of which three did not show improvements),⁷¹⁹ e-Prescribing in hospitals (23 of 25 studies showed medication error reduction and four of seven with reduced ADEs of 35 percent to 98 percent),⁷²⁰ CPOE in pediatric and ICUs (12 studies of proven error reduction but no effect on clinical outcomes),⁷²¹ CPOE in neonatal ICUs (reductions in errors but little if any effect on clinical outcomes),⁷²² outpatient CPOE (five studies of medication safety of which one showed reductions),⁷²³ and CPOE with CDSS to reduce ADEs in hospitals and ambulatory settings (10 studies: five showed improvements, four showed trends and one was not significant).⁷²⁴

Prescribing—Strengths and Limitations of the Evidence

The evidence in this section is weak although many RCTs exist (Table 14). Numbers of participants in the trials are often small, studies are short term, and are often done by those who have developed and implemented systems. To support the potential for bias in assessment of a health IT by its developers, Garg and colleagues⁷²⁵ completed a large and well-done systematic review of CDSSs. Their review evaluated RCTs of CDSSs for improving practitioner performance and patient outcomes. Using the data on practitioner performance, they found that if the trialists evaluated their own CDSS the trials were successful in 51 of 69 studies (74 percent). If the trialists were independent of the system being evaluated (i.e., not the developers), only five of 18 trials were positive (28 percent). The odds ratio (OR) adjusted for trial quality for a successful trial designed to improve practitioner performances if the evaluator was the developer was 6.6 (95 percent CI 1.7 to 26.7). The only other predictor of success besides the evaluator being the developer, in improving provider performance was if the users of the CDSSs were prompted to use the system automatically (adjusted OR 2.8, CI 1.2 to 7.1). It is difficult, however, to separate out developer bias from system effectiveness as they are confounded.

Commercial systems often do not have the resources to show changes in clinical outcomes and therefore this proof of clinical effectiveness is not completed.

Twenty-one RCTs studied the prescribing phase.^{401-403,407,408,515,518-520,524,526-528,537,541,543,630,637,697-699} In addition, seven cohort or case-control studies,^{501,545,701,702,709,710,712} and 27 observational studies^{16,18,423,425,430,437,446,452,453,455,459,467,468,475,482,489,550,555,631,683,688,693,703,704,706,708,714} also looked at the prescribing phase and reported clinical outcomes. Only the RCTs will be discussed below because of their strength of evidence. Four studies were done in the late 1990s.^{407,527,630,699} All of the rest were done after 2000.

Prescribing—General Study Characteristics

Participants. Because these studies evaluated clinical outcomes, all assessed patients and their caregivers. All RCTs used cluster randomization (clinicians or care units) to avoid problems of contamination (where the same caregiver is asked to use decision-support for a random half of patients but not the remainder).

Location. One study was done in a long-term care center,⁶⁹⁷ one was set in homes,⁶³⁰ and five were set in hospitals.^{401-403,407,637} All of the others were done in ambulatory care settings.

Drugs and diseases. Most studies evaluated specific diseases or conditions: asthma,⁵¹⁹ high cholesterol levels,^{515,528,699} hospitalized patients at risk of deep venous thrombosis or pulmonary embolism,⁵¹⁹ depression,⁵²⁰ infections in hospitalized patients,⁴⁰³ high blood pressure,^{526,699} and HIV.⁵²⁷

Technology. All studies involved CDSS. Six also included CPOE.^{402,407,519,527,697,699} A PDA was also featured in the home-based article.⁶³⁰

Prescribing—Clinical Outcomes

Please also see the section on CDSS (KQ7: RCTs of CDSS) for additional description of clinical outcomes. As seen in the systematic reviews, fewer articles address clinical outcomes than address process or other outcomes such as satisfaction and attitudes. Many of the studies that did evaluate clinical outcomes also did not find the expected improvements.

Adverse drug events. Gurwitz and colleagues⁶⁹⁷ found that the rate of ADEs and preventable ADEs were not decreased with implementation of a CDSS and CPOE system in a long-term care setting.

Disease related outcomes. A number of studies looked at disease outcomes. Kucher and colleagues⁴⁰² found that fewer patients at risk for venous thromboembolism were diagnosed with either deep venous thrombosis and pulmonary embolism at 90 days with the introduction of CDSS and CPOE in an academic hospital. Zanetti and colleagues⁴⁰³ studied prophylactic antibiotics in prolonged cardiac surgery. This RCT found similar rates of infection in both study groups. Of note, both control and intervention groups reduced their rates of infection. Rollman and colleagues⁵²⁰ addressed identification of depression in adult ambulatory care and found that the CDSS did not affect the rates of depression in the control or intervention groups in an RCT. Both groups improved their depression scores over time to the same extent.

Hospital stay. Three RCTs on prescribing looked at hospital length of stays. One RCT did not find differences in quality-of-life scores, hospitalizations, emergency department visits, or heart failure exacerbations.⁵¹⁹ Safran and colleagues,⁵²⁷ in another RCT of CPOE and CDSS for clinic patients in an academic setting, did find a lower hospitalization rate for intervention (reminders) group (44 percent vs. 35 percent, RRR 26 percent, $p = 0.04$). Hospital length of stay was not different in the RCT by Overhage and colleagues⁴⁰⁷ of CDSS and CPOE (eight days for both groups), nor in the study by McGregor and colleagues.⁴⁰¹

Physiological measures. Eleven RCT studies of the prescribing phase addressed physiological outcomes: hypertension in two articles and both showed no difference,^{526,699} high cholesterol with some positive findings in one⁵⁴¹ but not the other four^{515,524,528,543} and two with reductions in blood glucose levels.^{537,630} A study on depression found no change in patient depression rating scores.⁵²⁰ One study with asthma patients found improved lung function and airway hyperresponsiveness.⁴⁰⁸

Order communication. Order communication issues seemed to be at the heart of this before-after study of a children's hospital by Han and colleagues¹⁵ which showed increases in mortality after introduction of CPOE and CDSS integrated within a hospital information system. This is an important study and has garnered much discussion in the literature of its methods and findings with respect to the increase in mortality (2.8 percent vs. 6.6 percent unadjusted, adjusted OR 3.81, 95 percent CI 1.94 to 5.55). Length of stay showed improvement in one hospital, but not another in a study of CPOE implementation by Mekjjan and colleagues.⁵⁸¹

Dispensing and administering. The study by Han and colleagues¹⁵ evaluated dispensing; while three studies addressed administering.^{581,630,693} One study was an integrated system in the Ohio State University Health System (James Cancer Center and three other tertiary care hospitals). The hospital information system included laboratory, imaging, dietary, eMAR, and CPOE as well as all EMR capabilities. They found a reduced length of stay for patients with heart disease (14 percent) and transplant patients (15 percent) but not for those with cardiothoracic surgery or those in the cancer center.⁵⁸¹ Holdsworth et al.⁶⁹³ found significant reductions in ADEs following the implementation of a CPOE system in a pediatric population. The third administration study was a cross-over RCT of diabetic patients using a hand-held insulin regimen optimizer, which showed improvements in blood glucose levels when patients received advice through the device.⁶³⁰

Monitoring. Most of the prescribing interventions were integrated with hospital clinical information systems or EMR systems. This provided the opportunity to use existing structured electronic information to assist clinicians in identifying patients who needed a change in their treatment plan. The system made recommendations that suited the particular patient profile. Starting with monitoring of treatment choices for antimicrobial therapy in relation to antibiotic choice, a wide range of clinically useful monitoring and prescription and treatment recommendation options have been studied including those aimed at improving chronic disease management (e.g., Asthma-Critic), providing early detection of adverse events (e.g., creatinine monitoring for nephrotoxic effects), and glycemic and coagulation monitoring to predict and recommend optimal dose changes. Of the 21 RCTs that included the monitoring phases, 15 were

set in the prescribing phase. The issues related to RCTs and observational studies have been addressed in the general overview of studies in this area.

Reconciliation. One RCT at two academic hospitals studied a computerized reconciliation program that was integrated into a CPOE system and that required a process redesign.⁶⁹⁵ They found a reduction in unintentional discrepancies between preadmission medication and admission or discharge medication that had potential for harm (1.44 vs. 1.05 potential ADEs per patient, absolute risk reduction 0.72, 95 percent CI 0.52 to 0.99).

Education. In another RCT, Grant and colleagues⁶¹⁰ studied a PHR system for patients with diabetes that was integrated into a fully functioning EMR (laboratory, imaging, CDSS, and pharmacy). Patient education was a major, but not the only, component of the PHR. No change was noted for hemoglobin A_{1c} levels, although it is important to note that the patients were fairly well-controlled at baseline (7.1 percent vs. 7.2 percent, $p = 0.45$).

Qualitative Studies

Summary of Findings

Fifty-three articles that were complete or partially qualitative studies were identified that dealt broadly with MMIT (Appendix C, Evidence Table 10).^{20,439,503,540,547,597,629,632,633,635,652,666,671,674,726-764} No qualitative studies were identified that directly addressed the effect of an MMIT system on intermediate health care outcomes for any phase of the medication management continuum (prescribing, order communication, dispensing, administering and monitoring, as well as reconciliation, education, and adherence).

Strengths and Limitations of Evidence

The primary limitation of synthesizing qualitative studies to gain a deeper insight into the effect of MMIT applications in improving other intermediate health care outcomes within and across the medication management continuum is that no qualitative studies are available that directly address this question. Most of the qualitative studies identified examine the expectations or experiences of implementing an MMIT system on the process (but not the outcomes) of medication prescribing. These studies identify a large number of benefits to the health care delivery processes as well as a large number of barriers to uptake and use of the various systems studied. The strengths of the amalgam of evidence are that similar themes were identified across studies, health care settings were assessed by more than one study, studies were carried out in settings across the care continuum, study participants included physicians, pharmacists, nurses, other health care providers as well as some administrative management personnel, and multiple different types of qualitative data collection approaches including interviews, focus groups, observations, and document reviews were used across the set of studies evaluated. A small number of qualitative studies were available that examined MMIT systems on the processes of care for other phases of the medication management continuum.

MMIT is tremendously complicated and at the same time undeniably valuable. Strong and varied evaluations are vital and we have many evaluations of MMIT already. These evaluations show important changes to process. Clinical outcomes are more often mixed or nonexistent. We also see, in our evaluation studies, unintended consequences of MMIT and surprising results

such as the increased mortality in a children's hospital after a poor implementation of a set of MMIT applications.¹⁵ Because of these challenges in our results of quantitative studies, we include a fuller discussion of some of the qualitative studies in MMIT. These qualitative studies hold the promise of understanding more richly how MMIT is and should be used. The following paragraphs provide descriptions of some of the more important qualitative studies and their findings.

Prescribing and ordering. No qualitative studies were identified that directly addressed the effect of an MMIT system on intermediate health care outcomes. However, many qualitative studies provided evidence and examined positive and negative expectations and experiences of how an MMIT system designed to improve prescribing of medications could affect medication errors and medication safety,^{20,439,503,547,629,632,652,666,676,726,727,729-731,733-736,738-740,742,745,746,748,749,751-753,755,759,761-767} which could be considered as precursors to intermediate health care outcomes.

Before system implementation. Positive and negative expectations of an MMIT system designed to improve prescribing and ordering of medications were identified by physicians and other health care providers or staff in hospital or ambulatory clinic staff prior to system implementation. Some positive expectations were that an MMIT would reduce medication errors,^{727,733} increase pharmacological knowledge available,⁷⁴⁰ provide educational benefits,^{726,727} improve patient confidentiality,⁷²⁷ be flexible (e.g., prescribing from any location),^{666,729} allow for customization or tailoring to the individual prescriber or the patient (e.g., patient reminders), allow switching the system on and off,^{729,733} be concise,^{729,733} provide access to other areas of a medical chart,⁷⁴⁰ save time,⁷⁴⁰ and incorporate valuable allergy, dosing, and interaction alerts.^{745,746} Pharmacists felt that MMIT would facilitate new collaborations among physicians, pharmacists, and nurses.⁷⁴⁶

One group of physician study participants had a positive attitude towards implementation of a CDSS, provided that they had some control over the system.⁷²⁹ Many groups studied could be described as hopeful but cautious^{727,729,733,740,748} while others, mainly physicians (although they were who was studied most often), were skeptical.^{726,735,761,765} Hospital pharmacists felt that MMIT would allow them to spend more time with patients and improve collaborative working relationships with physicians and nurses.⁷⁴⁶

Only one study was identified that used qualitative methods to solicit patient views before implementation of an MMIT system focused on improving prescribing.⁷⁴⁸ Patients on a general surgery ward were interviewed before implementation of an e-Prescribing and an eMAR system. Their attitudes about the current paper-based system were generally positive and many had a mistrust of computer systems in general. However, they anticipated advantages of the e-Prescribing and eMAR system in terms of time, improving accuracy and efficiency, and decreasing mistakes. Patients identified that an electronic system may be an advantage for staff when the first language is not English.

Despite the willingness of many of the participants to use a new MMIT system designed to improve prescribing of medications including CPOE, some negative expectations were that the MMIT system would impair existing interactions and relationships among health care providers and between physicians and patients (e.g., diminishing patient contact because they need to leave the consulting room to enter the prescriptions),^{726,727,729,740,762} the ability to cope with the new system,⁷²⁶ implementation would be onerous,⁷²⁶ the costs of the system,⁷²⁷ especially to the health profession including the time efficiency and workload redistribution,⁷³⁵ the technical

challenges such as data entry time, software compatibility and updating,^{727,729} problems with availability or level of technical support,^{727,735} social and cultural barriers,⁷²⁷ deskilling of staff (people becoming dependent on the system for routine decisionmaking without understanding the background, reasons, or consequences of the decision made by the MMIT),⁷²⁶ need for more security,⁷²⁶ errors in prescribing such as decisionmaking errors,⁷²⁷ transcription errors,⁷²⁷ or overconfidence errors,⁷²⁷ that the system would not remove medication errors but could even create new errors,⁷³⁵ obscured responsibilities,⁷²⁹ loss of own reasoning and clinical autonomy,⁷²⁹ ensuring that the patient and not the computer would have the leading role in the encounter,⁷²⁹ difficulty with knowledge management (e.g., too much information or erroneous information—'garbage in–garbage out'),^{729,740} including prescribing alerts that were redundant or repetitive, of low priority, or difficult to interpret,⁷³⁹ resistance towards change,⁷²⁹ computer shortages,⁷⁴⁰ and altering workflow routines.⁷⁴⁰

Some underlying key needs for an MMIT system designed to improve prescribing of medications would be that the system would not diminish the patient provider relationship,^{726,727,729,740} be easy to use,⁷³³ flexible, concise, and customizable,^{729,733} clinically and technically trustworthy, reliable, and fast,^{729,733} integrated into other relevant systems,⁷⁴⁰ workflow needs to be maximized including development of new workflows^{726,727,739,735,740} and there be enough time and resources available to support implementation.^{726,727,735,740}

A Delphi survey was done in the United Kingdom to identify and reach consensus on the key clinical issues involving patient safety for which general practitioners in primary care might benefit from MMIT support, particularly in relation to medicines management. The key themes that emerged were importance of computerized alerts, need to minimize spurious alerts making it difficult to override critically important alerts, having audit trails of such overrides, support for safe repeat prescribing, effective computer–user interface, importance of call and recall, and need for safety reports. User interface, repeat prescribing, need to be able to run safety reports, and other safety issues were also agreed upon.⁷³⁸

After system implementation. The reporting of how MMIT systems designed to improve prescribing of medications improved intermediate health outcomes were also sparse among the qualitative studies. Drug alerts, including drug interaction alerts, were stated to be beneficial to improve patient safety.^{547,632,666,767} E-Prescribing triggered a variety of clinician behaviors (other than terminating or changing a prescription) that may improve patient safety.⁶³² One study identified 22 previously unexplored medication error sources users reported to be facilitated by CPOE which would likely have a detrimental effect on health outcomes.⁷⁵² These were grouped as (1) information errors generated by fragmentation of data and failure to integrate the hospital's several computer and information systems, and (2) human-machine interface flaws reflecting machine rules that do not correspond to work organization or usual behaviors such as selecting the wrong patient because a list is alphabetical versus by team or floor, or unclear log on and log off procedures or processes so that the next person does work using the previous person's permissions.

During or after system implementation, physicians, nurses, and other health care providers or staff in hospital or ambulatory clinic found that health IT improved safety alerts,⁶⁶⁶ provided useful drug alerts including drug interaction alerts, which appeared even if a different prescriber had ordered some of the drugs,⁵⁴⁷ allowed physicians to prescribe electronically from everywhere in the hospital, improved on features of a previous paper-based system,⁶⁶⁶ were user

friendly and allowed benefits when the system was integrated,⁷⁵¹ and was designed to take into account diverse cultures.⁷⁵¹

Physicians in one study felt that an electronic CPOE system improved the quality of care for patients because they got faster access to information and more up-to-date information, they received automatic reminders, and they could speed up care because knowing what had already been done would allow them to reduce the number of duplicate procedures carried out.⁷⁶¹ The multiple checks within the system also could lead to improved patient safety.⁷⁶¹ Better communication among physicians and structured reports for patients (e.g., discharge summaries) were also felt to improve quality of care.⁷⁶¹

In another study, physicians, physician assistants, and nurse practitioners felt that a computerized patient record system-based pain CDSS played a very positive role in assisting them with patient care. They reported that this was because data were more legible, could be accessed remotely, and reminders provided helpful decision support. However, they also reported that at the same time as being helpful, the reminder system was considered time consuming, redundant, and the speed of the system slow.⁷⁶³ Medical trainees also reported that an MMIT system provided valuable educational content such as geriatrics pharmacology review and nonpharmacologic treatment options.⁷³¹

MMIT systems designed to improve prescribing of medications also generated some challenges that could be categorized as challenges: (1) with the computer system and software, (2) of the interaction between the MMIT system and the health care provider working with the system, (3) of the effects of the system on the collaborative working relationships of the health care team, and (4) of the MMIT working within the local context and environment. Computer system and software challenges include difficulties with user rights, inflexibilities, and displacements with the use of CPOE,^{734,764} CPOE design failures, especially a faulty computer interface, lack of connection with other parallel systems, inadequacy of decision-support, and human errors occurring in interactions with the computer,^{503,727,759,761,767} difficulties with the text presentation (e.g., too much information presented, data density), too many decisions that needed to be made at one time, unappealing color scheme and lack of notation, caution or problems with a prescription,^{652,761,767} interface problems,^{750,755} content problems,⁷⁵⁵ and increased data entry time.⁷²⁷

Challenges generated from the interaction of the MMIT system and the health care provider working within the system included the need to take up new tasks and increased demands on the clinicians with the CPOE system),^{734,764} maintaining complete lists of patients and their medications,^{632,736} poor recording of data within the record such as allergy information,⁶³² propagation of errors if information was cut and pasted compared with creating new information and other mistakes,^{755,761} transcription errors,⁷²⁷ getting patient-specific formulary data,⁷³⁶ encouragement to ignore interactions alerts as many were viewed as too trivial or unnecessary, which indicated that sensitivity and specificity required improvements,^{547,632,753} initial difficulties with the technical components of the system,^{439,767} awkward prescription writing leading to workarounds,⁶³² unfamiliarity with the disease codes in the system,⁷⁵¹ difficulties with finding information in the chart because of multiple places where the information was stored,⁷⁶¹ reducing clinical situation awareness,⁷⁵⁹ overconfidence,⁷²⁷ and increased workload.⁷⁵⁰

Challenges generated by the MMIT system that affected collaborative working relationships of the health care team included damaging the workflow, synchronization and feedback mechanisms between nurses and physicians,^{666,676,750} altering the pace, sequencing, and dynamics

of clinical activities,^{750,759} and providing only partial support for the work activities of all types of clinical personnel.⁷⁵⁹

Challenges related to the MMIT working within the local context and environment included external implementation challenges (e.g., communication with pharmacists and vendor support),⁷³⁶ lack of computer resources,⁷⁵¹ the need to keep their EHR systems up to date,⁷⁵¹ poorly reflecting organizational policy and procedure,^{759,766} doctors' concerns that their views and opinions about the design and implementation of the new system had not been adequately addressed,⁷⁶⁴ high cost,⁷²⁷ social and cultural barriers,⁷²⁷ and problems with technical support.⁷²⁷

A number of studies related that participants felt it took longer to prepare a prescription using the MMIT system compared with conventional pen and paper.^{761,764,767} One study identified that physicians and nurses in an acute care setting found that CPOE did not meet naive and early expectations.⁷³⁴ Some adverse effects of the CPOE system were noted.

Attitudes towards MMIT systems in the early stages were mixed.⁷⁶⁸ Over time, and with experience of making the system work for them, attitudes changed to become more balanced and the potential benefits of the system become clearer to most.^{439,676,761} Some physician participants felt the MMIT was more efficient during consultation and led to better quality, while others were felt it took longer and took away from patient focus.⁷⁵¹ Physician users tended to provide comments related to the culture of professional quality (feeling that the computer facilitated quality). Alternately, those physicians that chose not to use the system tended to provide comments that focused on human relations. For example, they reported on their relationships with their patients that they felt were detrimentally affected by computer use.⁷⁵¹ Some physicians felt that MMIT helped physicians become more cost conscious by suggesting therapies that were less costly. This cost savings however, only directly benefited insurers and not the clinicians, patients, or health care facility.⁷⁵¹ Some physicians felt the MMIT systems improved their personal performance by allowing them to log on to the system from anywhere including home, while others felt this was an intrusion into their home life.^{761,767} Basic formatting and organization of information such as information that was legible, could improve order accuracy, or all in one place was seen as a benefit to MMIT.⁷⁶¹ MMIT applications were also felt to improve interdisciplinary work by improving communication with colleagues,⁷⁶¹ and having everyone reading from the same page.⁷⁶⁷

Alerts and reminders are important components of MMIT for prescribing. Important themes of these alerts or reminders in EMR systems included themes of efficiency, usefulness, information content, user interface, workflow, and training.^{742,745} Effectiveness focused on the positive effect of alerts on allergy awareness and patient education.⁷⁴⁵ Efficiency related to ensuring that the alerts and reminders were efficient, useful, and did not waste time.^{742,745} Usefulness concerned whether the alerts were helpful and appropriate.⁷⁴² Information content was concerned with accurate, comprehensive, timely, rich, and accessible information.^{742,745} The user interface was felt to be important for smooth and efficient work and provision of valuable information that was accurate and provided quickly.^{742,745}

The value of e-Prescribing alerts was diminished by the quantity of irrelevant and inappropriate alerts.⁶³² Workflow issues related to the information being available when and only when needed.⁷⁴² The need for training to improve the use of alert was noted.⁷⁴⁵ Attitudes to evidence-based guidelines were also seen as an important factor as to how alerts would be taken up, with physicians preferring that alerts be severity-rated, that only substantial ones should appear, and that user interface design be enhanced.⁷⁴⁵ The biggest surprise from a set of focus groups (reported in 2002) with a group of clinicians (physicians, physician assistants, and nurse

practitioners) was the considerable negative emotion associated with alerts and reminders (feelings of being criticized, embarrassment, guilt, frustration, annoyance, and anger).⁷⁴² One study of a set of three successful and three unsuccessful CPOE implementations across six hospitals identified 14 facilitating factors and 14 barriers comparing successful and unsuccessful implementation.²⁰ More people from the successful hospitals group reported supportive administering and heads of medical sections, direct involvement of physicians, mandatory implementation, adequate training, and sufficient hardware facilitated success. In terms of barriers, only inadequate hardware and lack of ability to easily complete patient transfer and advance admission orders (medical records package) differentiated the successful compared with unsuccessful groups. Changes involved in instituting a physician CPOE system are system wide and involve individual as well as organizational factors.²⁰ One study was identified that determined how clinicians use information management strategies during adaptation to an established CPOE system.⁷⁴⁹ User created strategies identified that information overload must be carefully managed and communication is vital and is often negatively affected by new systems.

Only one study using qualitative methods to solicit patient views after implementation of an MMIT system that focused on improving prescribing was identified.⁷⁴⁸ Patients on a general survey ward were interviewed after implementation of an e-Prescribing and administering system. Concerns were identified including loss of personal touch, not understanding the system, and perceived extra time needed if nursing staff had to check the drugs prescribed on the computer.⁷⁴⁸ Despite the concerns raised, on balance the feedback provided by patients was that generally they did not have a strong opinion (assessment) either positively or negatively as to whether MMIT would impact the quality of medication prescribing compared with paper-based process.⁷⁴⁸

MMIT also impacted the professionalization of pharmacy. The effects of a health IT system that generated an e-Prescription on the professionalization of community pharmacists were improving the analytical capacity of the pharmacists and physicians, greater dissemination of therapeutics and professional knowledge, better integration of process tasks, increased process automation, elimination of intermediaries, facilitation of the interpretation of prescriptions, increased tracking capability, and greater informational capability improves relevance and meaningfulness of interaction and improves quality of information transmitted.⁷³⁰ E-Prescribing has tremendous capacity to change and improve pharmacists' professional work and interactions.⁷³⁰ One study showed that overly ambitious expectations sometimes lead to failed implementation.⁶²⁹

Order communication. Seven qualitative studies specifically addressed the implementation of an MMIT system to affect the order communication and verification of prescriptions.^{540,597,671,732,736,746,752} None of these studies focused specifically on how MMIT affected intermediate health care outcomes. All of the studies addressed implementation issues. Nursing perspectives based on implementation of a BCMA system within the hospital setting found that an MMIT system was more time consuming but the nurses acknowledged that it produced a positive benefit because the extra time available was wisely spent to assure verification, generating an increased sense of safety for the patients,⁶⁷¹ or made improvements in the clarity of orders, organization of time their tasks, improved efficiency and standardization of documentation provided by templates, general improvement in emergency department processes, and decreased number of verbal orders and time searching for charts.⁵⁹⁷ One study identified 22 previously unexplored medication error sources that users reported to be facilitated by CPOE

including errors related to order communication and verification such as information errors generated by fragmentation of data and failure to integrate the hospital's several computer and information systems.⁷⁵²

These findings were consistent with another study carried out in a long-term care setting where numerous workarounds associated with the implementation of an eMAR and medication safety practices in nursing homes, were identified related to the technology itself creating unintentional blocks including slow wireless speed and the need to print each order on a separate page.⁷³² Organizational processes such as the limited resource of fax machines were also identified.⁷³² In the ambulatory setting limited electronic connectivity of e-Prescribing systems to pharmacies or pharmaceutical benefits managers (who administrate pharmacy prescriptions) meant that despite one-way electronic (non-fax) communication of prescription information from the practice there was still conventional communication (e.g., fax) back from pharmacies for clarifications and renewals.⁷³⁶ Pharmacist perspectives about a commercial e-Prescribing system revealed barriers to that systems' ability to maintain complete lists of patients and their medications, use of CDSS, and getting patient-specific formulary data.⁷³⁶ Factors associated with these issues related to product limitations, external implementation challenges (e.g., communication with pharmacists and vendor support), and physician preferences on specific product features.⁷³⁶ A system that appended alerts and comments to the bottom of e-Prescriptions and was designed to reduce pharmacy callbacks did not reduce the number of callbacks but did change the nature of the callbacks.⁵⁴⁰ Hospital pharmacy leaders with and without CPOE entry system experience all believed CPOE would improve patient safety through the allergy, dosing, and interaction alerts which they saw as valuable to medication management processes.⁷⁴⁶ Some expressed concern that poor design or implementation could lead to increased errors.⁷⁴⁶ Most believed the system would lead to improved efficiencies facilitating more time spent with patients.⁷⁴⁶ Most felt CPOE would improve working relationships with physicians and nurses by facilitating new collaborations.⁷⁴⁶

Medication dispensing and administering. Ten qualitative studies focused on evaluating health IT applications to improve medication dispensing and administering including studies of BCMA,^{635,671,674,728,743,754,756} PDA,⁷⁶⁹ eMAR,^{732,754} CPOE,⁵⁹⁷ and automated medication dispensing.⁷⁴⁴ All of these studies focused on evaluation of the process of care delivery before or after implementation of the systems.

Before implementation of a bar-code point-of-care eMAR system a group of pediatric nurses working in an American pediatric hospital provided qualitative responses to questions as part of a survey.⁶⁷⁴ Themes derived from the survey done before implementation indicated that the nurses felt that medications would be given in a timely manner with less error, but may result in an increase in time with this increase in safety, along with more reported errors, but fewer errors in administering actual meds (near misses). The surveys collected after implementation indicated that the staff felt there were fewer medication errors with a smoother administering of medication.⁶⁷⁴ Implementation of MMIT applications for medication dispensing and administering generated substantial number of nonIT workarounds.^{728,732} In one study done in a hospital setting, these workarounds were categorized into omission of process steps (seven workarounds), steps performed out of sequence (one workaround), and unauthorized process steps (seven workarounds).⁷²⁸ Probable causes for these workarounds included technology, task, organizational, patient, and environmental related causes.⁷²⁸ A further study examined how nurses integrated BCMA and an eMAR system into everyday clinical practice and found that the

implementation of new IT in the clinical setting can be disruptive to existing patterns of articulation work, or work that coordinates the activities of people across time and space.⁷⁵⁴

Another study of a system put in place in a long term care institution identified workarounds related to the technology itself and organizational processes.⁷³² The workarounds occurred at new medication order entry, communication with the pharmacy, and administering.⁷³² The technology introduced intentional blocks (safety features such as excessive dose blocking, dual documentation, and ADE monitoring) that led to workarounds related to the technology itself and organizational processes.⁷³² Organization process blocks leading to workarounds included the double checking of preparation and administration documents.⁷³² Integrating BCMA systems within real-world clinical workflows requires critical attention to ensure that technology safety features are used as intended and that nonIT systems are designed to support this use.^{728,732}

Nursing perspectives about a BCMA, eMAR system integrated with pharmacy, CPOE, and electronic charting in a hospital after implementation found that in terms of access, the nurses appreciated greater access to medications and information (e.g., policies, guidelines, drug resources, patient files), but identified some delays in getting medications from the pharmacy.⁶⁷¹ Another study carried out in a hospital and long-term care setting found that nurses were surprised that BCMA generated unanticipated side effects such as confusion created by automated removal of medications by BCMA, degraded coordination between nurses and physicians, and dropping activities such as not scanning wristbands or medications to reduce workload during busy periods.⁷⁴³ One study conducted interviews with nurses before and after the implementation of a BCMA. Before implementation most nurses expected the system to improve patient safety and after BCMA implementation most of the nurses reported that they felt BCMA improved safety although a number of concerns remained about the cumbersome and technical aspects of the system itself.⁷⁵⁶

After an automated medication dispensing system was installed interviews with all workers and managers who were affected (nurses, pharmacy managers, pharmacists, pharmacy technicians, hospital administrators, and patient care managers) resulted in themes of distrust, resistance, miscommunication, unrealistic expectations (skepticism that it reduced medication errors), speed and scale of implementation, concurrent changes, inadequate support, and social factors.⁷⁴⁴ Nursing perspectives were mostly positive on the use of a mobile PDA with a bar-code reader used to obtain medication profiles of patients and then uses as a decision support to identify drug therapy problems (e.g., drug interactions) for elderly home care patients, despite some system usability issues with the machine.⁶³⁵ Furthermore, some patients showed an interest when they saw the results from the electronic assessment.⁶³⁵

One ethnographic case study identified that the physician–nurse communications, mechanisms to ensure cooperation, and the procedures for preparing and administering the medications are the key process areas to address before implementing a system to augment the nursing administering of medications.⁷⁶²

Monitoring. Four qualitative studies assessed the clinician^{737,747} and patient^{633,760} perspective on the use of MMIT for medication monitoring. None addressed the effect of the systems on intermediate health care outcomes. The use of MMIT systems both facilitated and generated barriers to the process of patient monitoring by clinicians.^{737,747} One mobile phone-based system study showed that the MMIT system was well-accepted by patients as a mechanism to monitor symptoms for chemotherapy related toxicity.⁶³³

Adherence. No qualitative studies examined the effect of MMIT systems on medication adherence. MMIT systems facilitated patient monitoring⁷³⁷ by clinicians, however, barriers were reported to using health IT systems for patient monitoring.⁷⁴⁷ EMR with e-Prescribing facilitated monitoring and communication between physicians and patients with respect to the process of care that included checking active and inactive prescriptions and new and refill prescriptions, names of medication, and other medication themes (ordering and refilling prescriptions, mail-order issues, adherence, self regulation, alternate over the counter medication use issues).⁷³⁷ Clinicians caring for patients with HIV/AIDS using a CPOE and CDSS system integrated with the hospital, pharmacy, and laboratory systems identified six barriers to using reminders, including workload, time to document, reminders not applying, inapplicability to the situation, training shortcomings, quality of provider-patient interaction, and use of paper forms.⁷⁴⁷

Patients' perceptions and experiences were studied based on their use of a mobile phone-based advanced symptom management system (ASyMS[®]) for chemotherapy-related toxicity monitoring and management.⁶³³ Patients with lung, breast, or colorectal cancer who used the system generally felt that, with training, the handset was straightforward and easy to use, entering data twice a day for 14 days was acceptable, the system did not impact on patients' daily routines, and the set of six symptoms that were recorded on the handset were adequate (although some patients did indicate that they would have liked the opportunity to report other symptoms). They were very happy with the alerting facility of the system often reporting that they felt 'secure' in the knowledge that someone was being alerted about their symptoms, the real time, quick response rate of the data collection and alerting facility was viewed positively.⁶³³ However, one patient viewed the alerting system negatively, as she felt this part of the system was not sufficiently individually tailored.⁶³³

Another study focused on the patient perceptions of MMIT by studying a home telemonitoring device for ulcerative colitis that included their list of medications and questions designed to gather medication side effects.⁷⁶⁰ Patients felt that the system improved safety, feeling that the program 'would catch something I might not recognize' or help them 'respond quickly to a threat' to their health.⁷⁶⁰

Other studies with qualitative findings were also found.^{757,758,770}

Population Level Outcomes

Only one study met our inclusion criteria that assessed population level outcomes as a primary endpoint (Appendix C, Evidence Table 11). Yu and colleagues⁷¹² in 2009 conducted a case-control study using actual reportable ADEs from a relatively large number of pediatric hospitals, comparing the rates of ADEs between cases and controls in hospitals with various degrees of CPOE implementation. The study found that patients from hospitals without CPOE were 42 percent more likely to experience a reportable ADE after adjusting for comorbidities; thus a significant benefit is associated with CPOE implementation.

Composite Outcomes

Only one included study assessed a composite outcome as their primary endpoint (Appendix C, Evidence Table 11). Holbrook and colleagues⁷⁷¹ performed an RCT of 511 adult patients with type 2 diabetes receiving either usual care or an intervention involving shared access by patient and primary care provider to a Web -based diabetes tracker. The tracker interfaced with the providers' EMR and a phone reminder system, which sent monthly reminders for medications, laboratory reports, or physician visits. The main endpoint of process composite score for checks

of glycosylated hemoglobin, blood pressure, low density lipoprotein cholesterol, albuminuria, body mass index, foot surveillance, exercise, and smoking improved significantly more in the intervention group than in the control group (1.33 vs. 0.06 composite score scale; difference 1.27, 95 percent CI 0.79 to 1.75, $p < 0.001$).

Variation in Impact Depending on Medication Type or Form

Summary of the Findings

Although most studies looked at medication management in general, regardless of drug families, types or forms, 135 articles dealt with one or a few drugs or drug classes.^{18,399,401,403-405,409-411,414,416,418,420-428,430,431,433,437,440-442,444-449,451,452,454,458-464,466,469-473,475-478,481,482,486,489,491,492,494,496-499,501,502,505-512,514-517,520-525,530,534,535,538,542,545,546,548,553,555,557,559,562-566,568,570,577,578,582,588,592,596,608,611-615,618,621,622,624,630,631,633,647,661,683-685,698,701,702,705,713,731,772}

Prescribing and monitoring phases were again most often studied, with few studies looking at order communication, dispensing, or administering, and none on education (Table 16). No included studies addressed the issue of sound-alike or look-alike drugs, and four dealt with altering prescribing of generic drugs over name brand.^{414,458,510,535}

Specifically, 30 articles focused on antibiotics,^{18,399,401,403,405,409,418,423,426,428,451,452,458,460,464,469,475,477,482,506,523,525,562,563,596,614,647,661,683,684} seven on vaccinations,^{404,410,411,424,478,530,566} two on respiratory medications,^{446,613} three on psychotropics,^{476,502,520} two on nonnarcotic pain relievers,^{514,773} three on lipid-lowering agents,^{515,517,706} two on corticosteroids,^{462,553} 12 on cardiovascular drugs,^{414,448,449,505,509,510,521,522,534,588,592,624} and four on insulin.^{466,630,631,703} Narrow therapeutic index drugs were considered in 20 studies,^{421,425,427,447,461,463,470-472,481,507,512,555,577,612,618,633,685,701,702} and controlled substances in seven.^{437,445,486,501,535,564,731}

The form of medications was rarely mentioned, and was detected in only 18 studies.^{405,433,456,460,464,470,496,530,538,545,548,559,578,630,675,701,713,772} Prescribing changes from one drug form to another was the focus of two of these.^{460,464}

We focused here on narrow therapeutic index, controlled drugs, and the forms of drugs. The 20 studies reporting on narrow therapeutic index drugs overwhelmingly measured process ($n = 15$) and clinical outcomes ($n = 5$), only two measured costs,^{612,685} and one study was a qualitative assessment of patients on chemotherapy.⁶³³ The effect of the MMIT systems was generally positive on the main outcomes of process change measures of prescribing or laboratory monitoring changes. Clinical outcomes frequently were better with the use of the MMIT, but in some instances no change was observed.^{425,702} Systems used to assist monitoring or prescribing of narrow therapeutic index drugs were all either CDSS or CPOE systems.

Six of the seven studies on controlled substances measured changes in process, four of which showed a positive impact.^{437,486,501,535} Only two measured clinical outcomes with mixed results.^{437,501} The controlled substance interventions were some form of reminders, alerts, or CDSS in all cases but one, which dealt with order sets for opioids in CPOE.⁴³⁷

The evidence in this small selection of articles indicates that health IT interventions designed to influence the management of patients taking narrow therapeutic index or controlled drugs have positive impacts in terms of changes in process; results are less clear for clinical outcomes with a number of studies showing no change.

Only two studies targeted changing the form of a drug,^{460,464} both of which employed a CDSS and had positive results. Due to a lack of reporting of the form of medication being studied, we can make no conclusions about the variation in effectiveness of MMIT by drug form.

Strengths and Limitations of the Evidence

Narrow therapeutic index drugs. Of the 20 narrow therapeutic index drug studies, three are RCTs^{507,612,618} with quality scores eight, seven, and seven out of nine respectively. Three cohort studies are included^{685,701,702} with low quality scores of three, two, and three out of 10 respectively. The remaining studies were observational^{421,425,427,447,461,463,470-472,481,512,555,577} or mixed methods.⁶³³

Controlled substances. The evidence for managing controlled substances rests on seven studies. The quality scores for the one RCT,⁵³⁵ one nonrandomized controlled trial,⁴⁴⁵ and one cohort study⁵⁰¹ were generally low. The other four studies included a qualitative study,⁷³¹ and three observational studies.^{437,486,564}

General Study Characteristics

Narrow therapeutic index drugs. The narrow therapeutic index drug studies took place in hospitals (n = 14), ambulatory care (n = 6), and one at home. The drugs included digoxin,^{461,618} chemotherapy,^{421,447,633} anticoagulants,^{425,427,470,471,481,512,555,685,701,702} and others^{463,472,612} (Table 16). Three studies included CPOE interventions to assist with inpatient dosing,^{421,447,701} one on side effect monitoring by patients,⁶³³ and the remainder were CDSS alerts or reminder systems. Studies on anticoagulants measured adherence to prescribing and monitoring guidelines facilitated by some form of computer decision support.^{425,463,470,471,481,685} Two studies were of alerts sent to pharmacists for prescriptions written in primary care; one for prescriptions of drugs determined to be inappropriate for elderly patients⁵⁰⁷ and one for drug-drug interactions.⁵⁷⁷ One study implemented order sets within a CPOE for dosing of gentamicin and caffeine in the neonatal ICU, and assessed errors and drug turn-around times.⁴⁶³ Niiranen⁵⁵⁵ studied a computer-based warfarin followup system used by nurses to ease the burden on clinic physicians. Otherwise, prescribing physicians were most often the target of the alerts, reminders, or dosing support.

Table 16. Number of studies across the medication management phases using MMIT to assist in the management of specific drugs or drug classes

| Drug category | # of Studies | P | OC | D | A | M |
|---|--------------|----|----|---|---|----|
| Controlled | 7 | 0 | 0 | 0 | 0 | 0 |
| Benzodiazepines | 5 | 5 | 0 | 0 | 0 | 1 |
| Opioids | 4 | 4 | 0 | 0 | 1 | 2 |
| Narrow therapeutic index | 20 | 0 | 0 | 0 | 0 | 0 |
| Anticoagulants | 10 | 6 | 0 | 0 | 0 | 5 |
| Antineoplastic | 3 | 2 | 0 | 0 | 0 | 1 |
| Cardiovascular | 2 | 1 | 0 | 0 | 0 | 2 |
| Multiple narrow therapeutic index drugs | 1 | 0 | 1 | 0 | 0 | 0 |
| Psychotropic/hypnotics | 1 | 1 | 0 | 1 | 0 | 0 |
| Insulin | 4 | 4 | 0 | 0 | 1 | 2 |
| Antibiotics | 30 | 26 | 0 | 0 | 1 | 6 |
| Anti-infective (HIV) | 1 | 1 | 0 | 0 | 0 | 0 |
| Vaccines | 7 | 7 | 0 | 0 | 0 | 2 |
| Other | 37 | 32 | 0 | 0 | 1 | 16 |
| Antidote | 1 | 0 | 0 | 0 | 0 | 1 |
| Antiemetics | 1 | 1 | 0 | 0 | 0 | 0 |
| Anti-hyperglycemic drugs | 1 | 0 | 0 | 0 | 0 | 0 |
| Cardiovascular | 12 | 11 | 0 | 1 | 0 | 0 |
| Corticosteroids | 2 | 2 | 0 | 0 | 0 | 1 |
| Lipid lowering | 4 | 3 | 0 | 0 | 0 | 2 |
| Non-narcotic pain relievers | 2 | 2 | 0 | 0 | 0 | 0 |
| Psychotropic | 3 | 3 | 0 | 0 | 0 | 1 |
| Respiratory drugs | 2 | 2 | 0 | 0 | 0 | 2 |

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring
Some studies encompassed more than one phase. No studies looked at education for a specific drug or drug class.

Controlled substances. Two studies of controlled substances occurred in primary care settings. In an RCT, Fortuna and colleagues⁵³⁵ assessed the effect of computerized prescribing alerts on the prescription rates of heavily marketed hypnotics and benzodiazepines compared with their generic counterparts in 257 physicians. Smith and colleagues⁵⁶⁴ implemented a CDSS module to reduce prescribing on nonpreferred drugs in elderly patients in 15 primary care clinics. The other five studies were performed in hospital settings and used CDSS interventions^{445,486,501,731} and order sets in a CPOE⁴³⁷ geared towards prescribing physicians.

Outcomes

Narrow therapeutic index drugs. The interventions aimed at pharmacists both resulted in significant reductions in inappropriate prescribing. Raebel and colleagues⁵⁰⁷ reported a relative risk reduction of 16 percent inappropriate prescribing for elderly patients, and Humphries et al.⁵⁷⁷ reported a 31 percent relative risk reduction in drug-drug interaction prescribing. The studies of CDSS dealing with narrow therapeutic or narrow therapeutic index drugs frequently resulted in better laboratory monitoring of patients,^{461,472,612,618} prescribing adherence,^{427,470-472} dosing,^{447,555,618} or avoidance of errors.^{421,463,512} Cordero and colleagues⁴⁶³ found reduced errors and quicker medication turnaround times with the use of CPOE ordering and dosing in the neonatal ICU. Negative results were found by Riggio⁴⁸¹ with longer times to stop heparin treatment in patients experiencing heparin induced thrombocytopenia following implementation of an alert for 100 patients. Time from alert to laboratory test and start of direct thrombin inhibitor treatment did not vary before and after the implementation.

Clinical outcomes were measured in six studies. We considered positive studies to have at least 50 percent of the outcomes as being significantly impacted by the technology. Under this measure, four of the studies did not show significant impact of the technologies on patient outcomes^{425,555,685,702} though they tended towards being positive. Balcezak and colleagues⁷⁰¹ found better prescribing of heparin when a computerized nomogram was used by prescribers, but the nomogram was only used for 10 percent of prescriptions written. The highest quality evidence comes from Raebel,⁵⁰⁷ White,⁶¹⁸ and Feldstein⁶¹² and their colleagues who all showed positive, significant impacts of the technologies on narrow therapeutic index drug management.

Controlled drugs. The primary care RCT by Fortuna and colleagues⁵³⁵ found a significant decrease in the prescribing rates of heavily marketed drugs with the implementation of an alert plus education intervention, with a relative risk reduction of 74 percent. The observational study by Smith and colleagues⁵⁶⁴ in 15 primary care clinics to reduce prescribing on nonpreferred drugs in elderly patients showed a significant decrease in exposure of elderly patients to nonpreferred drugs, but no change in nonelderly patients, and a nonsignificant positive trend of preferred drugs in elderly patients.

The hospital-based quantitative studies showed generally positive process measures,^{437,486,501} with improved adherence to dosing in two,^{486,501} and better monitoring of patient pain levels.⁴³⁷ Morrison and colleagues⁴⁴⁵ found no change in prescription rates for laxatives to patients on opioids. Clinical outcomes were only measured by Peterson⁵⁰¹ and Wrona⁴³⁷ and their colleagues. Peterson and colleagues found no change in length of stay or rate of altered status, but a significant reduction in falls ($p = 0.001$). Wrona and colleagues found improved respiratory rate in patients on morphine and hydromorphone with order sets outlining monitoring and documentation requirements.

Unintended Consequences of MMIT Applications

Summary of the Findings

The unintended consequences of health IT are important and often not well-studied. (Note that this section is not about drug-related ADEs.) These unintended consequences associated with an MMIT are often identified after a system is implemented, despite careful planning and installation. Unintended consequences can be minor or major and they can be viewed as being helpful to the installation or detrimental. Eighteen studies were identified that reported unintended consequences of MMIT installations (Appendix C, Evidence Table 12).^{15,16,450,457,480,503,508,732,734,743,752,759,774-779} Because we report only those outcomes that the authors reported as the primary or main findings of the study, this listing of articles on unintended consequences is likely not comprehensive.

Strengths and Limitations of the Evidence

One study is a large observational study of medication errors reported to MEDMARX facilities that covers all the phases of medication management.⁷⁷⁴ As in previous sections of this report most of the studies evaluated prescribing. All of the remaining 17 studies (one RCT,⁵⁰⁸ eight quantitative observational studies,^{15,16,450,457,480,732,775,777} six qualitative studies,^{734,759,776,778-780} and two mixed methods studies^{503,752}) evaluated prescribing. Several of these studies also evaluated other phases. The order communication phase was evaluated in two studies, one

observational¹⁵ and one qualitative study.⁷³² Dispensing was studied in one observational study.¹⁵ Administering has one observational study¹⁵ and two qualitative studies.^{732,743} No studies of unintended consequences evaluated the monitoring phase or education and reconciliation.

General Study Characteristics

Participants. Most of the studies were done at an institution level rather than a patient or provider level. Raebel and colleagues⁵⁰⁸ studied medications with potential for harm to pregnant women and Han and colleagues¹⁵ studied admissions to a children's hospital after implementation of a CPOE system. Nurses were evaluated in two studies,^{732,743} and the rest of the studies included a range of clinicians.

Location. All studies were done in single hospitals or groups of hospitals. One study was done in a long-term care center.⁷³²

Drugs and diseases. Raebel and colleagues⁵⁰⁸ studied drugs with potential for harm to the fetus in pregnant women (category D and X medications). All other studies included all medications.

Technology. All of studies but two involved CDSS and CPOE systems integrated with EMR systems, dispensing systems or pharmacy information systems. The two studies that did not include CDSS and CPOE systems involved BCMA⁷⁴³ and eMAR systems.⁷³² They were both integrated within a hospital-based information system.

Outcomes

Ash and colleagues list a number of unintended consequences of MMIT and other health IT systems.⁷⁷⁸ These unintended consequences were categorized into direct compared with indirect, desirable compared with undesirable, and anticipated compared with unanticipated occurrences. Ash and colleagues contend that most unintended consequences center on errors, security concerns, and issues related to alerts, workflow, ergonomics, interpersonal relations, and reimplementation (e.g., updates). They also assert that all health IT systems will have unintended consequences.

Mortality. The University of Pittsburgh study of increased mortality with the introduction of an inflexible CPOE system is an example of a very serious unintended consequence.¹⁵ Because of the seriousness of the implications of this study, many people reviewed this article. Much attention has been given to this article and its methods.¹⁷ Another similar study shows that with careful planning, another children's hospital did not see the same increase in mortality in admitted children after careful implementation of health IT.¹⁶

Errors. New and different types of errors were identified as unintended consequences in three studies.^{450,457,503} Although most MMIT systems are associated with decreased errors, not all of the systems sought to determine **new or different** types of errors—they most often studied existing types and classes of medication errors. One study felt that problems with communication would probably lead to errors in medication management,⁷⁷⁵ and another study postulated the same increase in errors based on challenges to existing and changing roles.⁷³⁴ The study of use of inappropriate medications during pregnancy was stopped early because the system was not

accurate enough, causing the system to “miss” notification of drugs that should have been alerted and to give alerts that were not needed.⁵⁰⁸

Prescribing. Prescribing was not addressed specifically, although alert fatigue was a common theme in the studies of unintended consequences of MMIT.^{480,752}

Efficiency. Ash and colleagues⁷⁷⁶ list 47 types of unintended consequences and Koppell and colleagues⁷⁵² list 22. Ash and colleagues go on to verify that the types of unintended consequences they found were common in institutions outside those that she and her colleagues studied.⁷⁷⁷ Unintended consequences were related to roles,^{734,743,752,776,781} communication,^{775,779} workflow alterations or automation of poor existing workflows,^{752,759,779} inflexibility of the new system,^{743,752,759} poor content or poor display of content,^{752,759,776} alert fatigue,^{480,776,779} and overdependence on the system.⁷⁷⁹ Rather than fix the system, most often workarounds were instituted by clinical staff.^{732,743}

Summary. Seventeen of the 18 studies listed above report serious unintended consequences of MMIT in multiple categories. From these studies we see that unintended consequences exist for many health IT projects regardless of the quality of the implementation or the amount of planning that went into the project. Although consequences were viewed as being positive or negative, both provided useful information for those interested in MMIT implementation.

KQ2. What knowledge or evidence deficits exist regarding needed information to support estimates of cost, benefit, impact, and net value with regard to enabling health IT applications in terms of prescribing, order transmission, dispensing, administering, and monitoring, as well as reconciliation, education, and adherence? Discuss gaps in research, including specific areas that should be addressed and suggest possible public and private organizational types to perform the research and/or analysis.

Introduction

We identified gaps in the report—some that we expected and some not. We address gaps by the key questions (Table 17). In this section, some overlap exists with Chapter 5 (Future Research). Most of the gaps cross multiple phases of medication management. Where an issue is more strongly associated with a phase we mention the phase or other aspect (e.g., reconciliation).

Table 17. Summary of gaps and needs across key questions

| Gaps and Needs in Evidence and Knowledge | Strengths and Substantial Evidence |
|---|--|
| KQ1. Effectiveness: Medication Management Phases and other Processes | |
| Order Communication, especially two-way e-communication Dispensing Administering Reconciliation Education and training for professionals and patients System wide MMIT applications | Prescribing* Monitoring |
| KQ1. Effectiveness: Research Methods | |
| Controlled trials with comparative groups Trials with strong methods regardless of research method used Trials of whole systems and also components of MMIT Studies with outcomes important to patients Studies with population based outcomes Studies that address issues related to evaluation of complex interventions Pragmatic trials Multicenter trials Studies done by others besides developers Qualitative studies, especially of patients and families Evaluations of the evidence content of MMIT applications Knowledge translation (translational research) studies Understanding applicability of MMIT applications in relation to the complexities of MMIT systems Research teams or consultations that include clinicians, researchers, and informaticians as well as all major stakeholders | Observational studies Descriptive studies Studies that measure changes in process Studies done in one institution or location |
| KQ1. Effectiveness: Participants and Settings | |
| Nurses Pharmacists and other pharmacy personnel Other health professionals Patients and families especially in home situations Children Pharmacies, especially those outside of hospitals Long term care facilities Community Homes Specialty clinics Population based studies | Physicians* Hospital based settings Primary care/ambulatory care settings |
| KQ1. Effectiveness: Health IT Systems | |
| Fully integrated MMIT systems MMIT used by nonphysicians MMIT used by patients and families (patient-based systems) MMIT in relation to health information exchange systems | CDSS* CPOE |
| KQ1. Effectiveness: Reporting | |
| Reports of harms and other unintended consequences of MMIT systems and system integration Lack of consensus on terminology related to IT and MMIT Lack of standardization in reports of studies of MMIT despite having published standards | |

Table 17. Summary of gaps and needs across key questions (continued)

| Gaps and Needs in Evidence and Knowledge | Strengths and Substantial Evidence |
|---|--|
| KQ1: Effectiveness: Outcomes and other study endpoints | |
| Usability studies, especially those that can be generalized or transferred Workflow effects on both functional and dysfunctional groups before implementation Unintended consequences, with emphasis on major ones | Changes in process |
| KQ1: Effectiveness: Costs and Economics Studies | |
| Strong and full economics studies that include both costs and consequences Cost and economics studies of non-CDSS and CPOE systems | CDSS CPOE |
| KQ3: Value Proposition for Implementers and Users | |
| Full economic analyses Personal values of multiple stakeholders and what makes them decide to buy or use a system Patients and their families and their values Effectiveness research and pragmatic trials with an emphasis on outcomes important to patients The effect of MMIT on risk mitigation | |
| KQ4: System Characteristics | |
| Reporting and standardization of reporting of system characteristics, feature sets Head to head comparisons of systems taking into account their features and characteristics Health information exchanges and MMIT | |
| KQ5: Sustainability | |
| Operational definition of sustainability accepted by multiple stakeholders Studies that measure and report sustainability that are comparative across many groups and issues | |
| KQ6: Two-way EDI | |
| Studies of complete two-way EDI Studies of the effects of e-Prescribing on pharmacists, pharmacy personnel and patients and their families | The effects of e-Prescribing on hospital and primary care physicians |
| KQ7: RCTs of CDSS | |
| Strong trials on clinical outcomes | Trials of CDSS |

*substantial strength

General Gaps

Medication management phases. The literature places a great emphasis on studying the **prescribing** phase of medication management, with 263 of our included studies falling in that phase (Table 18). We feel that more study should be done on the phases of **order communication, dispensing, and administering**. In addition, the **educational requirements** for effective use of MMIT applications by health professionals needs to be studied. The evidence on the need to train patients and their families on how best to use MMIT systems as well as incorporating disease-specific information and management education into patient-based MMIT applications is needed.

Reconciliation of medications is vital, especially at the time of transfer to another health care setting, including transfer to and from home and community. Little evidence is available that MMIT systems are capable of and effective at doing this medication reconciliation and making adjustments to regimens. Challenges with system interoperability and standardized

representation of medication data make effective reconciliation using MMIT applications difficult.

Order communication is ripe for more research and development, especially in two-way communication to improve and speed up “perfection” of orders and prescriptions.

Table 18. Frequency of medication management phases studies plus reconciliation and education

| Phase | Frequency |
|----------------------|-----------|
| Prescribing | 263 |
| Order communication | 26 |
| Dispensing | 17 |
| Administering | 39 |
| Monitoring | 77 |
| Education | 3 |
| Reconciliation/Other | 6 |

Research methods. This same pattern of disparity for the number of studies in the medication management phases exists for the distribution of study methods. Most included studies are quantitative observational studies (Table 19). Although these studies provide good evidence for understanding and evaluating MMIT applications, more studies with control groups are needed to provide stronger methods where appropriate. MMIT applications are “complex” interventions and can be considered to be programmatic and pragmatic in their evaluation. Future research using methods appropriate for these complex interventions are needed. Studies of full MMIT systems and components of MMIT systems are needed.

Many studies were not powered to find the differences sought. We also identified other issues in study methods including inappropriate analyses, labeling of methods, and adjusting data sets in some of the observational studies. For example, studies seeking to identify such factors as barriers or facilitators of use of health IT systems did not report adjustment for multiple comparisons (e.g., Bonferroni corrections, bootstrapping, or Monte Carlo simulations). Some studies addressing feature preferences tested for 40 or more associations without adjustment. The authors of sections of this report also have commented on incorrect choice of statistical analysis techniques in some studies that could have led to positive findings that are not justified. Studies need strong statistical and methodological advice. We also agree with Bernstam and other informatics researchers and educators who suggest that research into MMIT systems needs to include those with informatics and research training and experience.⁷⁸²

Another gap in the research realm is the absence of formal study of MMIT in relation to knowledge translation (translational research). Much evidence exists on many aspects of MMIT.

Table 19. Frequency of research designs for included studies

| Design | Frequency |
|--------------------------------|-----------|
| RCT | 88 |
| Cohort or case-control studies | 21 |
| Observational studies | 207 |
| Qualitative studies | 39 |
| Total | 355 |

RCT Randomized controlled trial

Participants

Health care providers. Physicians are well-studied. Nurses, pharmacists, midlevel practitioners (e.g., nurse practitioners, physician assistants, advance practice nurses, and midwives), and hospital administrators are not (Table 20). Studies that include mental health professionals are also lacking. Studies that include nonphysician clinicians are not focused on the unique needs of the participants. The important issue of nursing workarounds that have developed to deal with systems that match physician but not nursing needs is also inadequately studied. Use and usability studies need to include all health professionals who use MMIT systems and studies need to be done that will allow knowledge gained in usability studies to be transferred to other settings.

Table 20. Number of studies that evaluated the effects of MMIT on clinicians (the clinicians were the major focus of the outcomes of the articles)

| Provider | Frequency |
|---|-----------|
| Physicians, undifferentiated | 28 |
| Primary care physicians | 29 |
| Specialists | 11 |
| Hospitalists | 18 |
| Other Physicians | 8 |
| Midlevel practitioners (physician assistants, nurse practitioners, advance practice nurses, midwives) | 7 |
| Nurses | 36 |
| Pharmacists | 23 |
| Hospital administrators | 5 |
| Other health professionals | 17 |

Patients. The age range of patients impacted by the MMIT were generally well-represented across age groups with notable concentration among those who require more prescription medications (e.g., middle age and geriatrics) (Table 21). However, the special needs of medication management for children such as age- and weight-based dosing were not adequately pursued. More study of pediatric patients would be beneficial. Many of these patient-specific studies used data from patients to evaluate MMIT systems and their functioning in hospitals and primary care settings. However the needs of the patients and their families to manage medications outside of hospitals and clinics were not studied. This lack of evaluation of MMIT systems that patients and families will use at home and the effects of these systems on patient care and outcomes is an important gap that will only grow because of the advent of new systems, improvements in existing ones, and the move of patient centered care, chronic disease management with the aid of health IT, and continued time and money pressures on health care providers. Qualitative studies that address pharmacists as well as patient needs and opportunities and important outcomes were also lacking.

Table 21. Frequency with which patients or caregivers across age groups were studied as the main focus of an article (how MMIT affects patients)

| Patient | Frequency |
|-------------------------------------|-----------|
| Infants (0 to 2 years) | 9 |
| Children (2 to 12 years) | 13 |
| Adolescents (13 to 18 years) | 20 |
| Adults (19 to 44 years) | 52 |
| Middle age (45 to 64 years) | 80 |
| Geriatric (65 years plus) | 85 |
| Patients with Undifferentiated Ages | 14 |

Settings. Hospitals and ambulatory care, but not necessarily specialty clinics, are also well-represented in the studies of this report (Table 22). The gaps are in other settings. Very few pharmacies or long-term care facilities were studied. Many existing articles on pharmacies and pharmacists were excluded because of lack of comparative data or integration of MMIT. Long-term care facilities, community locations, and homes also need formal evaluation to determine the effectiveness and use of MMIT applications for their constituents. One study evaluated outcomes at the population level.⁷¹² MMIT applications tend to target individuals and few of them measure population level effects. Research into the effect of MMIT on populations is challenging and research will have to be carefully planned.

Table 22. Study settings in which the MMIT application was studied (studies could take place in more than one setting)

| Setting | Frequency |
|---|-----------|
| Hospital | 224 |
| Ambulatory care (primary care offices and clinics and specialty hospital-based clinics) | 119 |
| Community | 1 |
| Home | 6 |
| Long term care facilities | 8 |
| Pharmacies | 28 |

Health IT systems. CDSS and CPOE systems are well-studied, most often in the prescribing and monitoring phases (Table 23). All other MMIT applications lack evidence of their effectiveness, especially in terms of workflow, communication, and clinical outcomes. Many studies did not report important details of the MMIT application itself, making the studies in this report more difficult to synthesize. From the descriptions in the articles we felt that descriptions of the system, including components and implementation issues such as training could have been added but they were not.

Another substantial gap that we noted is that the content of the MMIT systems was not studied. Systems like CDSS and CPOE and functions like drug-drug interactions and the knowledge base that reminders, alerting systems, and order sets are based on need a strong, evidence-based foundation of knowledge that is based on health research and reliably updated and disseminated. Assessment of the need for and value of this clinical evidence base was absent.

Table 23. Technologies that were the main focus of the studies of MMIT

| Technology | Frequency |
|--|-----------|
| CPOE/POE system | 102 |
| CDSS/CDS/CCDS/reminders | 213 |
| E-Prescribing | 41 |
| Order communication of the prescription to/from doctor to pharmacy | 3 |
| Pharmacy information system | 7 |
| Other | 35 |
| Barcoding-medication administering | 20 |
| Barcoding-dispensing | 1 |
| eMedication administration system (eMAR, eTAR) | 15 |

CPOE = Computerized provider order entry, POE = Provider order entry, CDSS = Computerized decision support systems, CDS = Computerized decision support, CCDS = Computerized clinician decision support

Health information exchange. Health information exchange is defined as the movement of health information across organizations using nationally accepted standards was not studied in any of the documents retrieved. Medication management is complex and challenged by interoperability of systems, and like reconciliation, it has not been evaluated in MMIT studies that we identified in this document.

Reporting. Through the process of data abstraction, we found problems with standardization and expanded inclusion of data elements in terms of reporting health IT studies. We feel that authors should be encouraged to strive for publication in the peer-reviewed literature rather than trade publications and news magazines. We also feel that authors should include more data in their publications of MMIT interventions. Lacks appear in descriptions of what was in place for medication management before implementation of MMIT systems (baseline data), full explanations of the MMIT system and its implementation process, settings, including culture, and participants (both health professionals and patients and their families). A guideline for writing evaluation reports in health IT, the STARE-HI, was published in 2009.⁷⁸³ We recommend that this document be used for planning and reporting research studies of MMIT. A list of the STARE-HI elements follows:

1. Title
2. Abstract
3. Keywords
4. Introduction
 - a. Scientific background
 - b. Rationale for the study
 - c. Objectives of study
5. Study context
 - a. Organizational setting
 - b. System details and system in use
6. Methods
 - a. Study design
 - b. Theoretical background
 - c. Participants
 - d. Study flow
 - e. Outcome measures or evaluation criteria
 - f. Methods for data acquisition and measurement
 - g. Methods for data analysis
7. Results
 - a. Demographic and other study coverage data
 - b. Unexpected events during the study
 - c. Study findings and outcome data
 - d. Unexpected observations
8. Discussion
 - a. Answers to study questions
 - b. Strengths and weaknesses of the study
 - c. Results in relation to other studies
 - d. Meaning and generalizability/applicability of the study

- e. Unanswered and new questions
- 9. Conclusion
- 10. Authors' contribution
- 11. Competing interests
- 12. Acknowledgements
- 13. References
- 14. Appendices

Another of the challenges in this report to do with retrieval of studies from the bibliographic databases and also for abstraction and combining data, were inconsistencies in the use of terminology. We observed differences in how authors categorized medication errors, ADEs, and therapeutic failures. Several authors are seeking consensus on terminology in health IT. These definitional aspects are also addressed in the STARE-HI reporting guidelines listed above.⁷⁸³ Most of the studies in this evidence report do not follow these guidelines.

Benefit and impact. Benefit and impact are similar but not identical. In the pharmaceutical world benefit can be thought of as being “can it work” often under ideal situations (i.e., efficiency research). Much of the evidence answering KQ1: Effectiveness is of this kind of research: evaluation of a project, often near its implementation and for a short period of time. Many of these studies attest to the fact that for process and other soft outcomes, many of the MMIT systems do work.

Impact, or pragmatic studies, refer to measuring the effect of an intervention in the real world. Very few studies in this report are in this category. Trials of this nature are complex, long-term, have large numbers of people/situations being studied, and are done on mature and well-functioning systems. These trials are costly to complete and require maturity in the systems. Their location is likely best at those centers in the United States that have established and mature health care systems that have solid support for technology, strong research teams, experience with qualitative and quantitative methods and expertise in collaborative projects that include clinicians, experienced informaticians, and patients and their families.

The gaps for completing benefit studies include the medication management phases of order communication, dispensing, and administering; people besides physicians (pharmacists, nurses, other health care professionals, patients and families, vulnerable populations); nonhospital settings (long-term care facilities, community, pharmacies, and home settings); generics, forms of medication, and controlled substances; MMIT applications beyond CDSSs; and dispensing, administering, adherence tools, and patient involved health IT.

Cost and economics. Cost and economics are complex issues and important to many people, groups, organizations, and governments. To complete a comprehensive economic evaluation (e.g., cost-effectiveness, cost-utility, or cost-benefit analysis) one needs to quantify all costs and benefits within a given perspective (e.g., societal). Strong economic evaluations can piggyback on an RCT, or an economic model may be developed with data from a number of sources. Well-designed studies with an economic evaluation component included, is the best way to move forward in this area.

Many studies have provided cost data, but useful economic data involves far more input. An example of a cost study with data that is limited in its use is by Chisolm and colleagues,⁴⁴⁶ who did a before-after study of children with asthma in a children's hospital. Their pharmacy charges

were \$373 before CPOE with standardized order sets were put in place, and \$429 after implementation.

Therefore, the gaps for estimates of costs in this report of MMIT are almost identical to those listed above. In addition, we identified gaps in research quality centering on research design and analysis. We need highly trained and experienced researchers and economists to complete useful and usable cost and economics studies in the complex and changing domain of MMIT.

Summary

This report identified broad based strengths and gaps in the MMIT literature. Many of the major endpoints sought were found to show positive and statistically significant improvements, especially those that dealt with process and issues related to use, usability, knowledge, skills, and attitudes. Clinical endpoints and full economic evaluations were lacking. We also identified gaps in the study of the phases of medication, people involved, locations of studies, and research methods. We also identified areas where these gaps are becoming more important such as patient and family needs and opportunities related to MMIT, complete MMIT systems, and interoperability. Much research has been done in MMIT, and moving forward needs directed and careful planning and vision to fill gaps in our evidence base, harness the best established and new research methods, and build on what we already know to embrace new and advancing abilities of MMIT.

KQ3. What critical information regarding the impact of health IT applications implemented to support the phases of medication management is needed to give clinicians (physicians, nurses, psychologists, dentists), pharmacists, health care administrators, patients, and their families a clear understanding of the value proposition particular to them?

The value propositions of health IT applications have been difficult to quantify with more of a focus in recent years on framing how best to consider and measure it.⁷⁸⁴ Menachemi and Brooks⁷⁸⁵ review the benefits and costs of EHRs and associated patient safety technologies. They have found that studies assessing the benefits of the technologies in process and clinical outcomes are far more frequent than those assessing the return on investment. This trend is supported by the considerable evidence presented in the current report; while we include numerous studies assessing process changes and clinical outcomes, the body of evidence on cost-effectiveness is sparse. A number of barriers to measuring return on investment in health technologies exist. Technologies do not result in a direct income stream and the benefits often accrue to organizations other than the ones making the investment as, for example, clinical benefit to patients and financial benefits to payers rather than the hospitals making the investments.⁷⁸⁵ Investments in health IT produce a fundamentally different kind of asset to health care providers, and the technologies and changes they bring are so complex that it is difficult to measure their benefits.⁷⁸⁴ Certainly the body of literature looking at return on investment for the various technologies covered in this report, across the various settings, is very limited.

We use the Center for Information Technology Leadership's (CITL) value framework (Table 24), which defines value as the sum of a technology's financial, clinical, and organizational benefits.⁷⁸⁶ This fits well with the definition used by AHRQ whereby "value" is defined as "clinical, organizational, financial, or other benefits derived from the adoption, utilization, and

diffusion of health IT less the costs of achieving these benefits” (<http://grants.nih.gov/grants/guide/rfa-files/RFA-HS-04-012.html>). The same considerations for stakeholder value propositions are elements outlined by Ash and colleagues⁷⁸⁷ as important themes to consider when implementing a system, specifically CPOE. We recognize that this framework does not include patients as an element, but we believe that the framework could be applied to the patient perspective and incorporate value propositions for patients where applicable.

The required information to make an assessment of benefits is different depending on the stakeholder. The costs incurred by primary care physicians in practice will be different and balanced against different organizational benefits than those incurred in hospitals, and influenced by factors such as practice size, the sophistication of the technology, and others.⁷⁸⁶ Similarly, what constitutes benefits to a patient will be different from that of other users. Subramanian and colleagues⁷⁸⁸ have looked at costs and benefits to a number of stakeholders using CPOE with CDSS in long-term care facilities. Their process sheds light on the facets that we need to understand and study before we can make sweeping generalizations about value of health IT application. They identified the various stakeholders, the potential costs and benefits of the health IT, and factors which could affect costs and benefits. Ideally, such an assessment would be available for each stakeholder using each technology in each setting. This is not often the case so realistically we will broadly look at factors taken into account in making a value assessment and determine what we know and where the gaps lie.

Table 24. Summary of the evidence in relation to the CITL value framework⁷⁸⁶

| CITL Value Framework ⁷⁸⁶ | | Current knowledge |
|-------------------------------------|---------------------------------------|---|
| Financial | Cost reductions | <ul style="list-style-type: none"> • The limited cost analyses evidence available indicates that MMIT may offer some cost advantages despite acquisition costs. • Cost of system purchase, implementation and maintenance are rarely reported in the primary literature • Investors in the technology do not always reap the rewards • It is difficult to reach any definitive conclusion as to whether the additional costs and benefits represent value for money due to a lack of high quality, full economic evaluations. |
| | Revenue enhancements | <ul style="list-style-type: none"> • MMIT rarely results in increased revenue • Not within the purview of this report |
| | Productivity gains | <ul style="list-style-type: none"> • Some improvements seen in length of stay • Some evidence of increased efficiency • Qualitative evidence of workflow and health care provider relationship disruptions • Seldom measured as main endpoints |
| Clinical | Care process advances | <ul style="list-style-type: none"> • A significant body of literature indicating positive, if modest, improvements • Still a lack of quality studies with strong methods • Clear indications that unintended consequences can impact the value of the systems to stakeholders |
| | Improved patient outcomes | <ul style="list-style-type: none"> • Shortage of quality studies with strong methods • Some technologies show a positive effect on patient outcomes • Often measured as secondary outcomes and lack power |
| Organizational | Stakeholder satisfaction improvements | <ul style="list-style-type: none"> • Qualitative and quantitative support for improved satisfaction and perceptions for a range of stakeholders, including patients, using a number of different technologies in various settings • Even when positive satisfaction is reported, improvements to the systems are often suggested |
| | Risk mitigation | <ul style="list-style-type: none"> • Not within the purview of this report |

Summary of the Findings

Given that the value framework is the sum of financial, clinical, and organizational benefits, the current body of literature summarized in our systematic review in KQ1: Effectiveness would indicate that too many unanswered questions exist to make a true value assessment for the different stakeholders and technologies in the applicable settings.

Financial Benefits

Cost reductions. The data on cost savings from the use of health IT in medication management are sparse. The few studies included in our review suggested that some cost savings may exist, which could be substantial over time. The economic information looks more favorable after the technology has been in place for an extended period of time so that the large upfront investment gets spread over time and then do we start to see a return on investment. However, a full economic evaluation requires the comparative analysis of alternative courses of action in terms of both costs and consequences, which provides the best information for making a decision to adopt an intervention or not, and very few of these have been rigorously completed in this field. We don't have good evidence of a positive return on investment. Also, the initial expenditure and ongoing costs were rarely reported and the included cost analyses were based on projections of savings given reported changes in care processes rather than improved clinical outcomes for patients.

Revenue enhancements. No studies that quantified revenue enhancements were captured for this review. Because of the nature of health IT assets, they do not per se bring about additional revenues to the investors.

Productivity gains. Evidence captured in KQ1: Effectiveness suggests that some productivity gains are achieved, often measured as improvements in efficiency in care processes.^{429,457,463} Gains achieved by reductions in outcomes such as lengths of stay or rehospitalizations have been less successful, though Durieux and colleagues⁷¹⁶ do report a significant decline in hospital length of stay in a review of drug dosing decision support technologies. A number of studies reported positive improvements in efficiency outcomes such as drug turnaround times,^{584,586,628} and time to administering drugs.^{439,600} One study reported that nurses spent about the same time on computer documentation as paper documentation.⁵⁶¹ In our review, efficiencies were rarely the main endpoints of any of the studies; they were frequently reported as secondary outcomes or additional measures analyzed, but without any assessment of the power of the analysis. Because of the quality of the studies, it is difficult to attribute true productivity gains except in the cases of some well-established systems as suggested by Chaudhry and colleagues.⁶⁰⁷ The qualitative evidence indicates that stakeholders believe that gains in productivity have occurred.^{439,547,632}

Clinical Benefits

Care processes. Certainly this aspect of values is the most studied across the phases of medication management, with 379 studies included in our review in KQ1: Effectiveness. These studies included a number of settings and stakeholders, and most reported improvements in processes of prescribing changes, adherence to guidelines or quality measures, error reductions, preventive care procedures done, and monitoring initiated. However, the studies were often observational and often had small sample sizes. In more than 80 percent of the cases in which an

improvement in process was sought, it was found to be positive. The findings of improvement were consistent across settings, levels of care, providers, and medication management phase. We report a positive effect in the use of MMIT in the prescribing and monitoring of controlled and toxic drugs as well. To balance this positive nature of the results, a growing body of evidence delineates unintended consequences of some technologies that will also contribute to the value proposition of stakeholders.^{632,734,752}

Patient clinical outcomes. We reported on 78 studies that assessed clinical outcomes as their primary endpoints, the majority of which focused on prescribing and monitoring phases. About half of these studies reported positive effects of the MMIT on patient outcomes. However, when clinical measures were the primary endpoint, often no differences between the intervention and control groups in the higher quality studies were seen (see Table 15). The strongest evidentiary weight for clinical outcomes is found in the use of CDSSs for the prescribing and monitoring phases, and the overall benefit is somewhat positive but most often mixed.⁷²⁵ The measurement of clinical outcomes is often so far removed from the MMIT intervention that it becomes difficult to make general conclusions about their efficacy, and adoption rates are still quite low. We found that efficacy was greater in interventions targeting specific populations or applications. Thus, a value assessment on patient outcomes would warrant a look at specific technologies, populations, and settings beyond the scope of this report.

Organizational Benefits

Stakeholder satisfaction. For implementation, adoption, and ongoing use of any technology to be successful, the people using the system need to find it useful, usable, and nondisruptive. Many studies have looked at workflow issues, satisfaction, and perceptions of users with regard to health IT. Our review includes only those providing qualitative data or comparison groups. The literature on satisfaction indicates that generally the stakeholders studied were satisfied with the technologies of interest, namely CPOE, CDSS, and e-Prescribing.^{651,654-656,656-658} Some studies, however, found no differences in satisfaction.^{651,659} Levels of satisfaction and positive perceptions were shown to be positively correlated with measures such as ease of use, productivity, quality of care, and reliability.^{654-657,661,673} Our review of the qualitative research in the area shows that the implementation of MMIT generates emotion, both positive and negative. MMIT implementation did not just mean that a clinician needed to learn a new IT system but it also affected most of the other parts of the delivery of care processes including how the interdisciplinary care team worked together. When determining the proposition values, the type of technology and how well it meets expectations and workflow are important considerations for users, greatly impacting their perceptions and openness to adoption/use.

Some literature has focused on comparing perceptions and attitudes of different health care providers, such as nurses compared with physicians and trainees;^{656,678} and residents compared with physicians using the same technologies.^{654,657,677} The findings from these studies indicate that perceptions of the benefits of MMIT can depend on the role of the user. The type of system and how it affects health care providers' work will impact how satisfied these stakeholders are with the technologies.

For any one technology or setting, insufficient data exist to determine levels of satisfaction among all stakeholders. From the literature we see that satisfaction and perceptions of the MMIT can vary according to provider role, setting, and technology, and no overall answer to the

question of stakeholder satisfaction exists. We have a deficiency of comprehensive studies of patients as stakeholders.

Risk mitigation. No literature was captured on risk mitigation in relation to the use of MMIT. A focus of the greater body of research, especially commentaries and narrative reviews, is on the use of technologies to reduce medication errors. Such benefits could have repercussions on risk mitigation, but also needs to be balanced with the fact that some technologies have been shown to result in new kinds of errors.

Conclusions

Only one study attempted to look at the value propositions across stakeholders in the use of MMIT and they concluded that to facilitate adoption of CDSSs in the long term care setting, financial incentives to both the institutions and physicians should be considered.⁷⁸⁸ Certainly, from the literature, we see no clear understanding of what information is needed from the standpoint of each stakeholder. We can surmise from studies that physicians consider cost, usability, patient improvements, and easy integration into workflow as important factors to consider before they purchase MMIT technologies. Hospital administrators place emphasis on other aspects such as costs, return on investment, and organizational change. The relative importance of these factors will vary among physicians practicing in different settings, with cost being more important to physicians in private practice than in hospitals, and other related issues. Capitation rates will also be a factor for physicians and will vary across U.S. states.⁷⁸⁶ Similarly, the importance of these factors will vary among pharmacists depending on their practice setting and the type of technology. For patients, convenience, usability, portability, and patient-centered functionality have been reported as important factors in their value assessment of consumer health IT.⁴ For MMIT, patients will likely be concerned with reduced medication costs, avoidance of ADEs, and improved disease management, although no studies evaluated their value-based concerns. Work needs to be done to identify the needed critical information before we can truly assess what is missing.

From the information garnered in this report, a growing body of evidence supports the use of some technologies (e.g., CDSSs) in prescribing and monitoring, which show positive changes in process, while large gaps in knowledge of the impact of the use of MMIT for other applications still exist (see KQ2: Gaps in Knowledge).

KQ4. What evidence exists regarding the impact of the characteristics of medication management health IT applications, such as open source, proprietary, conformity with Federal and other interoperability standards, and being Certification Commission for Healthcare Information Technology (CCHIT) certified, impact, likelihood for purchase, implementation, and use of such IT applications.

Summary of the Findings

Few studies ($n = 21$)^{45,48,632,653,661,663,667,789-802} demonstrated evidence of the impact of the characteristics of MMIT applications on likelihood to purchase, implement, and use such IT applications (Table 25). Little substantial evidence was found from studies that assessed open source health IT applications that met our inclusion criteria. Only two articles discussed

conformity with standards^{653,801} and one the Certification Commission for Healthcare Information Technology (CCHIT) certified system.⁸⁰⁰ Such system characteristics as the use of proprietary IT systems was suggested by seven articles^{45,632,653,663,798-800} and homegrown IT application by one article.⁶⁶⁷ Two reported on a stand-alone e-Prescribing system.^{653,798} Most of the articles suggest that the decision to adopt health IT applications has been influenced by the feature sets of health IT applications. Each of the 21 articles included in this section established evidence on likelihood to use, one on purchase,⁸⁰⁰ and five on implementation.^{653,789,791,793,798} A sizeable number (n = 20) of articles were on the prescribing and ordering phases, with only one on the administering phase of medication management.⁴⁵

The findings of the articles included in our study suggest that certain features of systems improve the likelihood of purchase, implementation, and use of MMIT. However, the literature is sparse and evidence from studies with stronger methods that can address this question is lacking. Most often authors spoke about barriers and concerns towards implementation and acceptance rather than characteristics of MMIT that could facilitate implementation, purchase, and use of such systems. Insufficient details were given about the technology they were studying. Head-to-head comparisons of systems differing in these features were not found.

A systematic review on CDSS revealed that widespread dissemination of appropriate CDSS might improve clinical practice, but providing information in electronic format alone does not ensure uptake.⁸⁰³ Fundamental issues related to system characteristics included the availability and accessibility of hardware, technical support and training, system integration into clinical workflow, timeliness of clinical messages, and acceptance of the system by various stakeholders.⁸⁰³ Another review involving descriptions of 112 information systems identified that for successful implementation, core components were order entry, guideline adherence, and decision support.⁸⁰⁴ Involving end users in the development process was also shown to be a key to success.⁸⁰⁴ However, these systematic reviews did not explore whether health IT system characteristics like proprietary or homegrown, system configuration, system characteristics, CCHIT certified, conformity with interoperability standard and standalone or integrated had any impact on purchase, implementation, or use.^{803,804}

Table 25. Number of articles addressing system type in relation to likelihood to purchase, implement, or use an MMIT system

| Systems | Likelihood To Purchase | Likelihood To Implement | Likelihood To Use |
|--|--------------------------------|------------------------------------|---|
| CPOE alone | 0 | 0 | Hospital ⁶⁶³ |
| CPOE with CDSS | 0 | Hospital ⁷⁹³ | Hospital ^{45,667,793} |
| CDSS | 0 | 0 | Hospital ⁶⁶¹ |
| e-Prescribing | 0 | Ambulatory Care ^{653,798} | Expert Panel ⁸⁰¹ Mixed set up ^{48,802} Ambulatory Care ^{632,653,798} |
| Pharmacy information system | 0 | 0 | Hospital ⁴⁵ |
| EMR/EHR/ clinical information systems | Ambulatory Care ⁸⁰⁰ | Hospital ⁷⁸⁹ | Primary Care ^{794,797} Ambulatory Care ^{799,800} Hospital ^{789,790} Mixed set up ⁷⁹² |
| Health IT (Type of system not specified) | 0 | Hospital ⁷⁹¹ | Hospital ⁷⁹¹ Primary Care ⁷⁹⁵ Mixed set up ⁷⁹⁶ |

Strengths and Limitations of the Evidence

Most of the studies were surveys (n = 18), although two used qualitative research methods^{632,801} and one collected data from scientific literature, organizations, government, and professional reports.⁷⁹⁷ Therefore, the strength of the evidence is relatively weak. Nineteen articles were published in the original literature and one was from the grey literature.⁸⁰⁰

General Study Characteristics

Participants. More than half of the studies (n = 13)^{48,653,661,663,667,790,792,794-797,799,802} evaluated physicians as the user of the technology. One article each included pharmacists,⁶⁶¹ nurses,⁶⁶⁷ directors and the leader of IT application users,⁴⁵ chief information officer,⁷⁹¹ pharmacy directors,^{45,793} and two administrative and other medical staff.^{661,667} Two reported combinations of different types of health care providers.^{632,798} One study convened a panel of technical experts representing organizations having direct experience in implementing e-Prescribing standards.⁸⁰¹ The size of the studies ranged from 14 to 18,600 participants.

Study setting. In most of the studies, the participants were primarily from hospitals,^{661,663,789-791,793} and some were set in pharmacies,⁴⁵ ambulatory care,^{632,653,667,798-800} and primary care.^{794,795,797} Four evaluated a combination of various settings.^{48,792,796,802}

Technology. Primarily five groups of health IT systems, namely, CPOE, CPOE with CDSS, CDSS, e-Prescribing, EHRs, and five other systems were studied.

Research methods. Research methods were weak: eighteen articles were surveys, two used qualitative research,^{801,805} while one used data from scientific literature, organizations, government, and professional reports.⁷⁹⁷

Outcomes

Identification of feature sets. Bell and colleagues conducted an expert panel consensus that resulted in 60 specific functional recommendations for e-Prescribing to improve patients' health outcomes and reduce costs.⁸⁰⁶ This list of features is useful for those considering an assessment in this area. We identified that one or more of these recommended features were the driving forces toward possible purchase, implementation, and use of health IT applications. Major features addressed in most of the articles were medication lists,^{208,632,653,663,789,792,794,798,801} dosing calculations,^{661,789,792,793,799} CDSSs (alerts and messages for allergies, drug-drug interaction, drug approval),^{45,632,653,661,667,789-794,796,798-801} e-Prescribing,^{48,208,632,653,667,790,792,794-798,802} and order communication of the prescription to pharmacies.^{45,632,792,794,798} Other factors were access to laboratory test results,^{45,789,790,792,794,797,799,800,802} implementation of guidelines,^{661,789,792,795,796,799} transcription services,⁷⁹¹ formularies,^{653,793,795,801} tallman letters and change of color to differentiate between look-alike drug name pairs,⁴⁵ integration with another system like BCMA, pharmacy information systems, etc.,^{45,791,793} and medication reconciliation (Table 26).^{792,793} (Tallman letters are the use of capitals to help guarantee recognition of differences between drugs with similar names, as for example, NovoLOG and NovoLIN, and HumaLOG and HumuLIN, helps differentiate these products.)

Table 26. List of articles addressing various features that were instrumental in the decision to purchase, implement, and use

| Features | Number of Studies Addressing the Features |
|---|--|
| Medication list | 9 ^{632,653,663,789,792,794,798,799,801} |
| eDosing calculations | 5 ^{661,789,792,793,799} |
| Clinical decision support (alerts and messages for allergies, drug-drug interaction, drug approval) | 16 ^{45,632,653,661,667,789-794,796,798-801} |
| e-Prescribing | 13 ^{48,632,653,667,790,792,794-799,802} |
| Order communication of prescription to pharmacies | 5 ^{45,632,792,794,798} |
| Access to laboratory test results | 9 ^{45,789,790,792,794,797,799,800,802} |
| Implementation of guidelines | 6 ^{661,789,792,795,796,799} |
| Transcription services | 1 ⁷⁹¹ |
| Formulary information | 4 ^{653,793,795,801} |
| Tallman letters and change of color to differentiate between look-alike drug name pairs, | 1 ⁴⁵ |
| Integration with another system (e.g., BCMA, pharmacy information systems etc.) | 3 ^{45,791,793} |
| Medication reconciliation | 2 ^{792,807} |

BCMA = Bar coded medication administration

Standards and conformity. Wang and colleagues⁶⁵³ suggest that mandating the use of standards is necessary but not sufficient for achieving the desired effects of e-Prescribing. Bell and colleagues evaluated two standards (Medication History Standard and Formularies and Benefits Standards) from the U.S. National Council for Prescription Drug Programs (NCPDP) that were considered as initial standards for e-Prescribing under Medicare.⁸⁰¹ Another study considered CCHIT certified IT applications to be the deciding factor for likelihood to purchase, implement, and use.⁸⁰⁰ The 2008 Healthcare Information and Management Systems Society Analytic Ambulatory health IT survey reported commercial, proprietary EMR systems were being used without any one vendor being the dominant leader.⁸⁰⁰ Apart from these two articles,^{653,800} four other articles^{45,632,663,799} reported the use of commercial proprietary systems with medication management health feature sets.

Use. All the studies addressing the decision to use were based primarily on one or more of the feature sets discussed above (Table 26). Three studies were on CPOE systems with CDSS capabilities^{45,667,793} and one on a CPOE system alone.⁶⁶³ The features that were more important were allergy checking, drug interactions, medication formulation, interface with the pharmacy information system and direct order communication, and integration with the BCMA and laboratory systems. Two studies were on CDSS,^{661,667} with one being integrated with CPOE.⁶⁶⁷ Their important features were e-Prescribing, drug-drug interactions, calculation of dosing, and access to additional information.

Seven studies were based on EHR, EMR, or clinical information systems^{789,790,792,794,797,799,800} with one study⁷⁸⁹ mentioning that a sizeable number of hospitals reported having implemented several key functionalities of CPOE and CDSS. The important features were CPOE with CDSS, electronically available laboratory test results, medication lists, e-Prescribing with electronic transmittal of prescriptions to pharmacies, access to reference materials, and dosing calculations.

Three studies were on general health IT systems.^{791,795,796} Grossman et al.,⁷⁹⁶ found the percentage of physicians reporting access to clinical activities such as obtaining guidelines,

generating reminders, and writing e-Prescriptions increased from 2000-2001 to 2004-2005 ($p < 0.05$). Six studies^{45,48,632,653,798,801,802} were on e-Prescribing with one being integrated with another system and hand-held access.⁶³² Some of the more important features addressed by these studies were e-Prescribing, medication lists, drug interaction and allergy alerts, receiving laboratory results electronically, changing doses, formularies, and order communication of prescription to pharmacies. According to the study by Bell and colleagues,⁸⁰¹ implementation of medication history standard and formulary and benefit standards in e-Prescribing would likely enhance usability of such systems if standard implementation was improved.

The qualitative study by Weingart and colleagues⁶³² found that the most valuable aspects of e-Prescribing in ambulatory care were the ease of changing doses, renewing prescriptions, ensuring legibility, and transmitting prescriptions to in- and out-of-state pharmacies. Participants were dissatisfied with the unreliability of transmitting prescriptions successfully to the pharmacy, creating medication lists, recording of allergy information, and quantity of irrelevant and inappropriate alerts. Despite their complaints about alerts, participants preferred to continue receiving alerts as a safeguard against missing a major interaction.

The studies of such health IT systems as pharmacy information systems found that important features were CDSS alerts, interface with the laboratory system, and tallman letters and change of color to differentiate between look-alike drug name pairs.⁴⁵

Purchase. One article reported on likelihood to purchase in a group of which half of the respondents of that survey were planning to purchase a CCHIT certified EMR system.⁸⁰⁰ The important features were electronic connectivity for laboratory test results and orders, nursing and physician orders for medications, and prescription refills.

Implementation. Five articles reported on the likelihood to implement an MMIT system.^{653,789,791,793,798} Larger hospitals, those located in urban areas, and teaching hospitals are more likely to implement EHR systems.⁷⁸⁹ Collectively, the important features were allergy checking, drug interactions, medication history, dosing calculation, medication formulation, and availability of laboratory test results.

Wang and colleagues⁶⁵³ conducted a descriptive field study of ten commercially available ambulatory e-Prescribing systems, to compare the functional capabilities offered by commercial ambulatory electronic system with 60 expert panel recommendations suggested by Bell and colleagues.^{798,806} Data were collected from vendors by telephone interview and at sites where the systems were functioning, through direct observation of the systems and through personal interviews with prescribers and technical staff. Five of the systems were full EHR systems and five were nonEHR systems. Among the 60 e-Prescribing recommendations by Bell and colleagues,⁸⁰⁶ 6 nine recommendations were not implemented by any of the ten systems.⁷⁹⁸ These included recommendations that would require e-Prescribing systems to handle prescription fulfillment data (their recommendations 10, 47, and 48), to use more complex drug benefit data (recommendation 22), and to use more advanced drug knowledge bases (recommendations 26 and 49).^{798,806} Prescribing systems that were part of EHR systems implemented more recommendations than did stand-alone nonEHR systems. Considering all 60 recommendations, the median EHR-based system fully implemented 60 percent, whereas the median nonEHR system fully implemented 35 percent ($p = 0.09$). Including partial and full support together, median implementation levels were 72 percent for EHR systems and 46 percent for nonEHR systems ($p = 0.06$). On average, the systems fully implemented 50 percent of the recommended

capabilities, with individual systems ranging from 26 percent to 64 percent implementation. Only 15 percent of the recommended capabilities were not implemented by any system.

Level of care. Six studies evaluated systems for ambulatory care.^{632,653,667,798-800} Features of the six ambulatory care studies centered around clinician experience using commercial proprietary systems, with CDSS capabilities being the most common feature used.

KQ5. What factors influence sustainability (use and periodic updates) of health IT applications that support a phase of medication management continuum (prescribing, dispensing, administering, and patients' taking of medications)?

Sustainability of Health IT and Medication Management Systems

AHRQ seeks to support activities that can demonstrate the effect of health IT on important outcomes relating to quality, safety, efficiency, and effectiveness. Moreover, AHRQ places priority on initiatives to identify and overcome barriers to health IT implementation and adoption and to foster long-term sustainability.⁸⁰⁸ Intuitively, a system's sustainability refers to its capacity to continue providing value. Sustaining the benefit of health IT applications may require ongoing resources for maintenance and updating, training and support for those who use the systems, as well as institutional support that encompasses planning, implementation and maturing of the systems, and replacement as needed. Thus the concept of sustainability raises questions about the long-term viability of many health IT interventions, as well as important concerns about the potential health impact of migrating existing processes to less sustainable or costly forms.

We conducted an additional comprehensive review of the literature to find a suitable operational definition and set of metrics of sustainable health IT. We found this necessary because we did not have, and could not readily find a prespecified definition that was widely accepted or supported in the literature of health IT.

In search of a definition of sustainability relevant to health IT, we did additional searching in the core informatics journals using the key term "sustainability" to identify articles that have discussed the concept (Table 27). Some articles defined sustainability quite narrowly (e.g., the decline of prescribing improvements once experimental alerts were removed from a system that had integrated CPOE and CDSS systems). We believe that the most relevant available definition comes from Humphreys and colleagues,⁹ who defined sustainability as the ability of a health service to provide ongoing access to appropriate quality care in a cost effective and health-effective manner.

Our literature reviews revealed three important findings: although sustainability is mentioned frequently in the core informatics literature, it is poorly and infrequently defined, and none of the articles identified in the primary literature searching done to produce this evidence report explicitly studied sustainability. These findings were not entirely surprising. A previous AHRQ-sponsored Evidence Report that assessed the costs and benefits of health IT in pediatrics found only one article that explicitly discussed sustainability.²¹

Table 27. Frequency of core informatics journal articles that mention sustainability to the end of 2009

| Informatics Journal | Frequency of mention of sustainability to the end of 2009 |
|---|--|
| Journal of the American Medical Informatics Association | 28 |
| Journal of Medical Internet Research | 90 |
| Medical Decision Making | 3 |
| Journal of Biomedical Informatics | 9 |
| International Journal of Medical Informatics | 22 |
| Methods of Information in Medicine | 30 |
| BMC Medical Informatics and Decision Making | 9 |

Future Sustainability of Health IT and Medication Management Systems

In 2009, the United States passed the Health Information Technology for Economic and Clinical Health (HITECH) act to authorize incentive payments through Medicaid and Medicare to clinicians and hospitals when they use electronic health records (EHRs) for patient care. The legislation ties payments specifically to the achievement of advances in health care processes and outcomes. Starting in 2011, the HITECH act will make available incentive payments totaling up to \$27 billion over 10 years. This legislation will require substantial collaboration between health IT workforce professionals including those from IT, health information management, and biomedical informatics to accomplish its goals.

According to Dr. David Blumenthal, the U.S. National Coordinator for Health Information Technology at the Department of Health and Human Services, “[this legislation] will lead us toward improvements and sustainability of our health care system that can only be attained with the help of a reliable and secure nationwide electronic health information system.”⁸⁰⁹ HITECH’s goal is not just based on the adoption, but also on the “meaningful use” of EHRs. Meaningful use is defined by a set of core health IT objectives that constitute an essential starting point, as well as an additional menu of activities which providers and hospitals will choose to implement during 2011 to 2012 (see Figure 5).⁸⁰⁹ Overall, these features should help clinicians make better medical decisions and potentially avoid preventable errors.

This legislation may lead to improvements and sustainability of health IT applications that specifically support the medication management continuum. For example, to receive incentive payments, eligible professionals (e.g., physicians, optometrists, podiatrists, and chiropractors), and hospitals (e.g., acute care hospitals and critical care access hospitals) must implement and use the following *core set* of objectives that relate to medication management: CPOE, e-Prescribing, implementation of at least one decision support rule, and maintenance of active medication and allergy lists. Eligible professionals and hospitals may implement and use the additional *menu set* of objectives that relate to medication management: incorporation of clinical laboratory test results in the EHRs, performance of medication reconciliation across care settings, and sending reminders to patients for followup care.

Conclusions

We conducted an additional literature review of the core informatics journals to identify articles that have discussed sustainability related to MMIT systems and found that while sustainability is not infrequently mentioned in this informatics literature, it is often poorly defined, and none of the articles included in this evidence report explicitly discussed sustainability. Future research should develop an operational definition of sustainability that can

be used to study its determinants. Moreover, it is likely that the HITECH Act of 2009 will lead to improvements and sustainability of health IT applications that specifically support the medication management continuum through meaningful use.

Figure 5. Summary overview of meaningful use objectives

| Summary Overview of Meaningful Use Objectives.* | |
|---|---|
| Objective | Measure |
| Core set† | |
| Record patient demographics (sex, race, ethnicity, date of birth, preferred language, and in the case of hospitals, date and preliminary cause of death in the event of mortality) | More than 50% of patients' demographic data recorded as structured data |
| Record vital signs and chart changes (height, weight, blood pressure, body-mass index, growth charts for children) | More than 50% of patients 2 years of age or older have height, weight, and blood pressure recorded as structured data |
| Maintain up-to-date problem list of current and active diagnoses | More than 80% of patients have at least one entry recorded as structured data |
| Maintain active medication list | More than 80% of patients have at least one entry recorded as structured data |
| Maintain active medication allergy list | More than 80% of patients have at least one entry recorded as structured data |
| Record smoking status for patients 13 years of age or older | More than 50% of patients 13 years of age or older have smoking status recorded as structured data |
| For individual professionals, provide patients with clinical summaries for each office visit; for hospitals, provide an electronic copy of hospital discharge instructions on request | Clinical summaries provided to patients for more than 50% of all office visits within 3 business days; more than 50% of all patients who are discharged from the inpatient department or emergency department of an eligible hospital or critical access hospital and who request an electronic copy of their discharge instructions are provided with it |
| On request, provide patients with an electronic copy of their health information (including diagnostic test results, problem list, medication lists, medication allergies, and for hospitals, discharge summary and procedures) | More than 50% of requesting patients receive electronic copy within 3 business days |
| Generate and transmit permissible prescriptions electronically (does not apply to hospitals) | More than 40% are transmitted electronically using certified EHR technology |
| Computer provider order entry (CPOE) for medication orders | More than 30% of patients with at least one medication in their medication list have at least one medication ordered through CPOE |
| Implement drug–drug and drug–allergy interaction checks | Functionality is enabled for these checks for the entire reporting period |
| Implement capability to electronically exchange key clinical information among providers and patient-authorized entities | Perform at least one test of EHR's capacity to electronically exchange information |
| Implement one clinical decision support rule and ability to track compliance with the rule | One clinical decision support rule implemented |
| Implement systems to protect privacy and security of patient data in the EHR | Conduct or review a security risk analysis, implement security updates as necessary, and correct identified security deficiencies |
| Report clinical quality measures to CMS or states | For 2011, provide aggregate numerator and denominator through attestation; for 2012, electronically submit measures |
| Menu set‡ | |
| Implement drug formulary checks | Drug formulary check system is implemented and has access to at least one internal or external drug formulary for the entire reporting period |
| Incorporate clinical laboratory test results into EHRs as structured data | More than 40% of clinical laboratory test results whose results are in positive/negative or numerical format are incorporated into EHRs as structured data |
| Generate lists of patients by specific conditions to use for quality improvement, reduction of disparities, research, or outreach | Generate at least one listing of patients with a specific condition |
| Use EHR technology to identify patient-specific education resources and provide those to the patient as appropriate | More than 10% of patients are provided patient-specific education resources |
| Perform medication reconciliation between care settings | Medication reconciliation is performed for more than 50% of transitions of care |
| Provide summary of care record for patients referred or transitioned to another provider or setting | Summary of care record is provided for more than 50% of patient transitions or referrals |
| Submit electronic immunization data to immunization registries or immunization information systems | Perform at least one test of data submission and follow-up submission (where registries can accept electronic submissions) |
| Submit electronic syndromic surveillance data to public health agencies | Perform at least one test of data submission and follow-up submission (where public health agencies can accept electronic data) |
| Additional choices for hospitals and critical access hospitals | |
| Record advance directives for patients 65 years of age or older | More than 50% of patients 65 years of age or older have an indication of an advance directive status recorded |
| Submit of electronic data on reportable laboratory results to public health agencies | Perform at least one test of data submission and follow-up submission (where public health agencies can accept electronic data) |
| Additional choices for eligible professionals | |
| Send reminders to patients (per patient preference) for preventive and follow-up care | More than 20% of patients 65 years of age or older or 5 years of age or younger are sent appropriate reminders |
| Provide patients with timely electronic access to their health information (including laboratory results, problem list, medication lists, medication allergies) | More than 10% of patients are provided electronic access to information within 4 days of its being updated in the EHR |

* This overview is meant to provide a reference tool indicating the key elements of meaningful use of health information technology. It does not provide sufficient information for providers to document and demonstrate meaningful use in order to obtain financial incentives from the Centers for Medicare and Medicaid Services. The regulations and filing requirements that must be fulfilled to qualify for the Health IT financial incentive program are detailed at www.cms.gov.

† These objectives are to be achieved by all eligible professionals, hospitals, and critical access hospitals in order to qualify for incentive payments.

‡ Eligible professionals, hospitals, and critical access hospitals may select any five choices from the menu set.

Source: New England Journal of Medicine, 2010.⁸⁰⁹ Used with permission.

5a. To what extent does the evidence demonstrate that health care settings (inpatient, ambulatory, long-term care, etc.) influence implementation, use, and effectiveness of such health IT applications?

Implementation

Reports of implementation tend to be opinion pieces or descriptive studies. A number of articles looked at some or all of implementation, adoption rates, and factors related to adoption. These focused mostly on CPOE in hospitals,^{789,810-814} e-Prescribing, or ambulatory CPOE in primary care.^{48,172,794,796,815-819} These articles did not meet our criteria for inclusion in KQ1: Effectiveness because of methods limitations. The general findings for hospitals show that implementation and adoption are generally greater in larger, academic, urban, public hospitals. Adoption in primary care practices tends to increase with younger, recent medical grads, larger practice size, and also with more specialized physicians. Yet, overall, actual usage of the systems is low with varying rates across MMIT systems, facilities, and groups of people.

Poon and colleagues^{820,821} discussed a number of barriers to CPOE implementation in U.S. hospitals, and provided recommendations to overcome the barriers based on experiences in successful hospitals. They categorized barriers into physician and organizational resistance, cost and lack of capital, and vendor or product immaturity. They further provided recommendations to overcome these barriers.

Ash et al.⁷⁸⁷ provided a consensus statement with a list of categories and considerations for a successful implementation of CPOE (see <http://cpoe.org/> for more details). Their predominant themes to consider are: motivation for implementation; CPOE vision, leadership, and personnel; costs; integration; value to users; project management and implementation staging; the technology; ongoing staff training, support, and evaluation. These themes reflect the considerations for the values propositions of the various stakeholders as addressed in KQ3: Value Proposition.

The implementation of a new health IT can have unintended consequences, often as a result of the interaction between the technology and the sociotechnical system within which it is implemented. This would include the workflows, culture, social interactions, and technologies in place. Harrison et al.⁸²² have modeled this interactive sociotechnical system to help users and implementers understand how and where unintended consequences could arise within a particular setting. Some examples of unintended consequences have been reported in this document, and our sections on intermediate and qualitative research in KQ1: Effectiveness describe some information on workflows, social interactions, communication, and interdisciplinary work challenges with MMIT implementation. Assessing the potential impact of a new MMIT system using the framework of Harrison et al.⁸²² could easily help to avoid some of the unintended consequences reported to date.

Beyond CPOE and e-Prescribing systems, the useable literature on implementation and adoption of other technologies is negligible. Some surveys did look at the adoption of various health IT applications for patient safety. Menachemi and colleagues⁸²³ measured rates of adoption of CPOE, BCMA applications, pharmacy IT systems, pharmacy dispensing, EHRs, PDAs and CDSSs in Floridian acute care hospitals. Pharmacy systems were widely adopted (85 percent for IT systems and 64 percent for dispensing), and the rest ranged from 12 to 40 percent. They also studied health IT adoption in U.S. pediatric hospitals and found a fairly high level of adoption (almost 50 percent for EHRs, 40 percent CPOE, and 36 percent CDSS). Furukawa and colleagues⁸²⁴ used national survey data to measure adoption of technologies across the United

States. They found a range of levels of adoption across technologies, from 62 percent for automated drug dispensing to five percent for BCMA. Their analysis supports the findings that hospital size, teaching status, hospital or clinical ownership, and system membership are associated with adoption. Robinson et al.⁷⁹⁰ found that adoption of 19 health IT capabilities, including MMIT, was higher when practices were evaluated for pay-for-performance and public reporting purposes, and in practices participating in quality improvement initiatives. From our review of the qualitative literature, we find that unintended negative consequences, the need to develop workarounds including changes to workflow, and the resultant negative emotions generated with MMIT implementation are important to recognize and deal with in order to improve the success of implementation.

The American Society of Health-System Pharmacists (ASHP) have published the results of an ongoing series of surveys assessing the adoption and use of pharmacy informatics applications in U.S. hospitals and trends in pharmacy practice.⁸²⁵ Again the studies are descriptive surveys and not included in our KQ1: Effectiveness. They do, however, show how hospitals are progressing rapidly in their adoption of health IT used in their pharmacies.^{826,827}

Effectiveness

Due to the observational nature of many of the studies assessing health IT across settings, this question is difficult to answer. Hospitals and primary care are well-studied, especially for the two phases of prescribing and ordering, and monitoring. Gaps are seen in the other phases of medication management, and education and reconciliation. Further, some MMIT applications are well-studied while others, such as those that pharmacists and nurses use, are less well-evaluated. A limited number of studies are carried out in long-term care settings, pharmacies, or with patients at home, or other community settings. Many of the hospital- and clinic-based studies tended to show improvements in process with some, but limited, evidence of clinical improvements. These research gaps in settings, MMIT applications, and health professionals, together with proof of effectiveness are similar to the deficiencies seen with cost-effectiveness or other types of economics studies, especially full economic analyses which are required to address and satisfy the definition of sustainability used here.

Use

From KQ1: Effectiveness we see that the study of use of systems is rarely done in a completely rigorous way. Articles that measure use tend to frame it in the context of adoption and implementation, looking merely to ascertain if systems are used, not how they are used and if they are being used appropriately. Few measure levels of use. Again, the definition of sustainability is not met without the inclusion of economics studies.

5b. What is the impact (challenges, merits, costs, and benefits) of having electronic access to patients' computerized medication records, formulary information, billing information, laboratory records in the quality and safety of care provided by health IT applications that support at least one phase of the continuum of medication management (prescribing, dispensing, administering, and patients taking of medications)?

Almost all of the MMIT applications we report were integrated with at least one other system. The systematic review in KQ1: Effectiveness addresses the effect of these integrated

technologies on a range of outcomes, many related to patient safety and health care quality. Evidence is available to address prescribing and also monitoring. The other phases are not well-evaluated. The study of patient access to their medication records and integration of these data into clinic and hospital information systems (EMRs and EHRs) is exciting. Some evidence exists that the use of MMIT integrated into clinician-based systems holds much promise and will be an exciting area of research in the next decade. This is especially important with the efforts by the U.S. government to improve health care delivery and to implement health IT systems to enhance this expanded delivery.

KQ6. Two-Way Prescriber and Pharmacy Electronic Data Interchange (e-Prescribing)

(a) What evidences exists demonstrating the barriers and drivers of implementation of complete EDI that can support the prescription, transmittal and receipt, and perfection process of e-Prescriptions?

(b) How do barriers, facilitators, and economic incentives vary across pharmacists, physicians and other relevant stakeholders with respect to adoption and use of complete EDI (e-Prescribing/ordering with e-transmission)?

All studies eligible for inclusion in this evidence report were reviewed to determine if they evaluated two-way, complete EDI between prescribers and pharmacies, commonly referred to as e-Prescribing. To be considered a true one-way e-Prescribing system the article had to describe a computer system used by a prescriber to generate a prescription (authorization to supply drug) that is transmitted electronically to a pharmacy information system. Further, for the system to be considered a two-way e-Prescribing system it had to be capable of transmitting a message from dispenser to prescriber by electronic means. This criterion is broadly consistent with the definition of e-Prescribing promulgated under the U.S. Medicare Modernization Act of 2003.⁸²⁸ We did not consider systems used for inpatients of a hospital to be an e-Prescribing system; these technologies are reviewed elsewhere in the report, often under the rubric of CPOE systems.

Summary of the Findings

Thirty-three reports^{434,549,561,575,579,584-586,645,668,724,730,736,797,800,801,806,829-844} were checked for eligibility and only one⁵⁸⁵ met the above criteria for inclusion for bidirectional e-Prescribing systems. Nearly all systems self-described by investigators as “e-Prescribing” allowed physicians or other prescribers to generate a prescription through a software application that were later reproduced in paper form prior to being dispensed by a pharmacist (incomplete one-way e-Prescribing). One report⁵⁸⁵ described an interrupted time-series study of a two-way e-Prescribing system intended to reduce the time required for prescribers to respond to pharmacist queries and refill requests. The authors did not describe any barriers or facilitators to uptake of the system used in the small pilot study.

We have extrapolated key themes from the data available on one-way or incomplete one-way e-Prescribing to describe potential barriers and drivers to implementation of complete two-way EDI. These data may be useful indicators of issues that would need to be addressed before widespread implementation of two-way EDI would be expected to yield benefits for stakeholders. The following facilitators and barriers are listed in order of high to low frequency of mention in the reviewed literature.

Facilitators

1. ***External monetary or other incentives to prescribers.*** Nearly all reports of e-Prescribing implementations in the United States described some financial incentive that was offered to prescribers to adopt an e-Prescribing system.⁸³⁹ In most of those cases where no financial incentive was offered, the system was adopted by a health system that required its prescribers to adopt the system.
2. ***Supportive regulatory environment.*** Formal endorsement by regulators such as the State Boards of Pharmacy or Medicine seemed necessary enablers for prescribers to adopt e-Prescribing systems.^{736,839}
3. ***Existence of some standard for prescription electronic data interchange.*** A set of messaging standards to enable the electronic flow of prescription information between diverse software platforms have been developed for use in the prescribing and order communication processes.^{834,836,845} While not all standards have been judged suitable for implementation,⁸³⁹ the core set of standards currently available should facilitate further development and testing of e-Prescribing solutions.

Barriers

1. ***Incomplete consideration of the effects of e-Prescribing on pharmacists and pharmacies.*** Most evaluations of one-way e-Prescribing systems conducted in the United States focused almost entirely on the e-Prescribing system from the perspective of the prescriber, the prescriber's staff, or both.^{736,833-836,838,839} Several of these reports described a lack of awareness of the e-Prescribing process on the part of pharmacies and pharmacists and a subsequent need to educate pharmacists on the specific e-Prescribing process adopted by the prescriber.^{736,835} Pharmacists and pharmacy staff generally reported that e-Prescribing systems negatively impacted their workflow.^{645,833,834} The authors of an AHRQ commissioned report⁸³⁹ on e-Prescribing pilot projects conclude that the prescribing workflow advantages observed for prescribers using e-Prescribing may actually reflect a burden shift to pharmacists. While reduced pharmacy to prescriber callback rates are touted as a potential advantage to e-Prescribing, the highest quality evidence available did not support a reduced callback rate.⁵⁷⁵ A sample of e-Prescribing prescriptions sent to selected pharmacies in Denmark was prospectively compared with a sample of handwritten prescriptions sent to the same set of pharmacies. The investigators' adjusted analysis indicated a significantly higher likelihood (relative risk, 1.7; 95 percent CI, 1.3 to 2.2) of pharmacy callbacks to prescribers for electronic compared with paper prescriptions.⁵⁷⁵ This finding is especially significant as nearly two-thirds of prescriptions are transmitted electronically in Denmark.^{575,797} All pharmacies in Denmark are government owned and therefore likely share the same IT infrastructure.
2. Pharmacists are an essential part of the medication use process and better integration of e-Prescribing and pharmacy information systems through, at a minimum, one-way complete electronic data interchange should be a focus of further research.
3. ***Regulatory and legal uncertainties.*** Some participants in U.S. studies were unsure whether complete one-way e-Prescribing was permitted under relevant State laws.^{736,839} e-Prescriptions for controlled substances were not evaluated as a component of the reviewed studies because of the perceived prohibition on the use of e-Prescribing for these drugs under relevant State and/or Federal laws. Prescribers were also concerned

that notification by pharmacies of prescription fill status (filled or not filled) could increase their exposure to malpractice claims.⁸³⁹

4. **Low preexisting adoption rate of EMRs and EHRs.** Nearly all of the systems evaluated in the United States described the use of prescription writing software limited to generating e-Prescriptions, but without any other clinical record keeping functionality.^{736,839} These systems generated prescriptions and retrieved pharmacy dispensing histories while requiring providers to concurrently maintain paper-based medical records. Prescribers report deferring adoption of e-Prescribing (prescription writing) systems in favor of complete EMR systems that include e-Prescribing functionality.⁸³⁹ Thus the low rate of EMR adoption in the U.S. likely decreases the rate of e-Prescribing adoption. Addressing barriers to EMR adoption⁸⁰⁰ may increase the rate of e-Prescribing amongst physicians and other providers.

Summary of Evidence

No reports documenting the use of complete two-way EDI (prescribing) systems were located in the literature search for this report. Evidence from the limited set of one-way e-Prescribing studies was extrapolated to identify possible key facilitators and barriers to completely electronic, two-way e-Prescribing systems. Possible facilitators include monetary or other incentives to providers, a permissive regulatory environment, and the existence of enabling technical standards necessary for e-Prescribing. Barriers identified included the low rate of EMR adoption in the United States, regulatory and legal uncertainties, and inadequate consideration of the effects of e-Prescriptions on pharmacists and pharmacies.

KQ7. What evidence exists regarding the extent of integration of electronic clinical decision support (CDS) in a health IT system for prescribing and dispensing of medications?

To what extent does the use of CDSS in a health IT system for prescribing and dispensing of medications impact the various outcomes of interest including health care process, intermediate and clinical?

Summary of the Findings: All Phases of Medication Management

Seventy-seven RCTs in total were designated as primarily studying CDSSs related to medication management and with integration with other health ITs.^{397-399,401-405,407,409-416,504-531,533-543,592,609,611-613,616-620,624,630,634,636-638,697-700,771}

Full details of the studies are contained in Appendix C, Evidence Tables 13-15. These studies involved 4,709 providers and approximately 828,441 patients in total (numbers were not specified in all articles). Patients included were primarily adults, with only two studies addressing issues specific to children. Seven studies addressed seniors exclusively. Currently, AHRQ has contracted with Duke University to prepare an evidence report focused on CDSS due for release in 2011.⁸

All studies assisted with at least the prescribing (71 percent) or monitoring (29 percent) phases of medication management. Notably, none concentrated solely on the order communication, dispensing, or administering phases of medication management. Reconciliation and education were also not addressed.

The studies were much more likely to focus on process changes than clinical (patient-important) outcomes. Furthermore, many studies did not report directly which outcome was their

main endpoint—a fundamental flaw. A total of 36 articles measured changes in process as their main endpoints, 24 of which were deemed to have positive results—meaning that at least 50 percent of the changes in process measured showed that the MMIT improved medication management. Only five of 34 studies measuring clinical outcomes, whether a main endpoint or not, had a statistically significant impact on a clinical outcome. These five RCTs were all published recently (since January 2005), addressed primarily the prescribing and monitoring phases of MMIT, and a variety of disease and drug target groups, usually in an outpatient setting.^{402,537,541,620,634} Where clinical outcomes were thought to be designated main endpoints, 12 of 16 studies showed no differences in clinical outcomes between intervention and control groups.^{403,518-520,526,528,624,630,637,697,699,700} No study was able to demonstrate a positive impact on mortality.

Regarding integration of the CDSS, authors used various descriptions of other components of the integration. EMRs, EHRs, and hospital information systems were specified in 41 of the studies, and CPOE was integrated with CDSS in 10 studies, seven of which specified CPOE in addition to the EMR.

Strengths and Limitations

As per our inclusion criteria, all trials used randomization for allocation. However, by applying the Verhagen/AHRQ RCT quality scale,¹⁰ the overall quality of methods of the studies was generally only fair at best with a mean quality score of 4.4 out of total possible nine points. One of the most important features to avoid bias, allocation concealment was only described to a minimally acceptable degree by 25 studies. Twenty articles scored six or more and none of the studies scored the maximum nine points. Mean followup of the studies was 9.9 months. Twenty-four studies (31 percent) used a cluster design. This design is prone to bias. Cluster numbers are often small, and therefore, if clusters initially randomized to control group drop out, or participants within the clusters (who are known to be in the intervention or control group) are selected in a biased manner, trial results may not be valid.

Overall, high quality is lacking from RCTs that address CDSS integrated with other types of health IT. Only a small minority of these focus on clinical outcomes—those outcomes that are most important to guide decisions of patients’ providers and policymakers about these interventions. Furthermore, a very small number report improvement in these clinical outcomes.

General Study Characteristics

Of the 77 trials, 46 (60 percent) were rated as impacting primarily the prescribing phase of medication management, 12 (16 percent) aimed primarily at medication monitoring, 15 (19 percent) tried to impact both phases and one addressed administering. Three trials (4 percent) attempted to influence a mix of prescribing, monitoring, order communication, and administering phases of medication management.

The setting for the studies was judged to be ambulatory care in 53 (69 percent), or hospital-based in 19 (25 percent), with a small minority based in long term care (two (3 percent)), or other settings (three (4 percent)) such as community or home. Approximately half (36 or 47 percent) of these studies were identified as associated with academic institutions.

Health care providers were a target of the CDSS in 64 studies and included physicians in most cases where targets were specified. However, many studies did not address the specific type of provider targeted by the intervention. Three studies identified pharmacists as one of the

intervention targets and one study targeted nurses specifically. Patients were named as targets of the intervention in 22 studies, 13 of which exclusively targeted patients.

A wide variety of diseases and drugs were studied as the topic of the CDSS. Of the 42 studies where disease targets were mentioned, 19 dealt with vascular disease including risk factors, eight with diabetes, six with asthma, and four with infections, including HIV. Drug topics were evaluated in 42 studies—19 were vascular medications, 13 antibiotics or vaccines, and five addressed multiple medications. The CDSS system was known to be ‘home grown’ in 26 studies, a commercially available product in 14, a hybrid of both in four, and unknown in 33.

Thirty-five CDSS were thought to be integrated with an EMR or EHR system. Fourteen were integrated with CPOE or prescription writing systems, another 17 with a laboratory or imaging system and ten other multiple systems.

Characteristics of each of the CDSS were beyond the scope of this review, so it is unclear whether any signals from these RCTs indicated how a system should be designed, installed, maintained and training supplied, to optimize the chance of success. Similarly, we were not able to critique the suitability of control groups in this systematic review, which were typically described as usual care.

Outcomes

Of the 77 studies, 54 indicated in some way that they had a primary or main outcome and only 16 appeared to have designated a clinical outcome as a main endpoint. Clinical outcomes were defined liberally as any clinical morbidity, mortality, quality of life, adverse event, or clinical surrogate such as improved LDL cholesterol levels. Only eight studies addressed mortality in any way; none had a significant effect.

Overall, only five studies noted a positive change in clinical outcomes.^{402,537,541,620,634} All were published since early 2005. Four of the five took place in an outpatient setting.^{537,541,620,634} The studies addressed venous thromboembolism prophylaxis,⁴⁰² asthma control,⁶³⁴ cholesterol management,⁵⁴¹ diabetes care,⁵³⁷ and recommended drugs.⁶²⁰ The mean quality score of these five studies was only 4.8 out of nine. Two studies with the highest methodologic quality (six out of nine) are further described. One evaluated a CDSS which calculated venous thromboembolism risk and recommended venous thromboembolism prophylaxis when the risk was high thus improving their main endpoint of venous thromboembolism rates in a group of inpatients primarily with cancer. The other used a university affiliated managed care plan data to identify gaps in recommended drug therapy and monitoring to recommend drugs to stop or add, or for monitoring to take place. However, this analysis was based on a post-hoc outcome applied to a subgroup of the original participants and the changes in hospitalization are very high given the small change in recommendation use. In summary, we found no consistent impact of CDSS on clinical outcomes, and the quality of the studies is generally inadequate.

In 38 studies, a process endpoint was determined to be a main endpoint. In 26 cases, the process was judged to be positively affected; with improvement in at least 50 percent of the process measures reported. The changes in process measured in these studies generally dealt with reminders about recommended medications or vaccines,^{403,404,407,410,509,525,530,535,536} dose adjustments,^{398,412} recommended laboratory monitoring for medications prescribed or chronic disease management,^{412,504,513,516,612,619,771} ‘inappropriate’ medications avoided,^{397,413,416,507,508,512,533} and other similar outcomes. Some of the alerts or reminders were based on established guidelines, while others were assessing more locally derived quality measures and standards of care.

Only one of the studies we reviewed scored at least eight out of nine for the AHRQ methods quality assessment. Terrell et al.⁴¹⁶ randomized 63 emergency physicians to receive or not receive alerts to disrupt intended prescriptions of nonrecommended medications for seniors to be discharged from the emergency department. The CDSS resulted in a small decrease in the number of visits with a nonrecommended prescription from 3.9 percent to 2.6 percent (95 percent CI 0.34 to 0.89, $p = 0.02$). No clinical outcomes were measured in this study.

One article measured a composite score in which a shared CDSS to support the primary care of diabetes improved the process of care and some clinical markers of the quality of diabetes care.⁷⁷¹ One other study evaluated whether actively or passively displaying context-sensitive links to infrequently accessed educational materials and patient information using an inpatient CPOE would affect access rates to the materials, and found that the active alerts were more effective.⁶³⁸

Notably, the negative effects of the CDSS intervention were virtually never reported. Specifically, only two studies referred to any harm incurred by the study intervention.^{508,630} This implicates a major publication bias, a result of not requiring studies to measure and report on harm.

In terms of costs, 11 studies reported that they had intended to measure costs or cost-effectiveness. However, no full cost-effectiveness analysis was found as part of the RCT. Separate publications on resource utilization are covered in the KQ1: Effectiveness section on economics outcomes.

Summary

In summary, despite it being 34 years since the first RCT⁶¹⁹ in 1976, in this important area of health IT research, little high quality evidence shows a consistently positive effect of CDSS on clinical outcomes. Implementers, developers, and funders of MMIT applications need to continue to produce and rely on the best possible research evaluating outcomes important to people and institutions. The informatics world can strengthen their abilities to determine value for money in MMIT projects by obtaining input during planning for research projects from health technology appraisal methods and those who have expertise in clinical care, research methods, informatics, statistics, and stakeholders who will be affected by the MMIT system.⁸⁴⁶

Discussion

Summary of Key Findings

We have presented the results of a systematic review of the literature regarding the use of health IT to enable all phases of medication management as well as reconciliation and education. We have focused on MMIT systems that were integrated with other health IT systems. Our review identified a total of 789 studies dealing with health IT and medication management. Three hundred and sixty-one of these articles were only listed in the bibliography of this report and were not synthesized because they did not include comparative data, statistical methods, or qualitative methods. The remaining 428 articles were synthesized after being identified from an initial retrieval of 40,582 articles. We used these 428 articles to address the seven key questions (KQs). Overall, we found that the literature on MMIT applications was heterogeneous. The majority were based on observational methods, often with identifiable opportunity for bias (e.g., descriptive before and after studies without statistical adjustment for time trends or group differences). Research methods were not uniform across MMIT applications, with 77 of 88 RCTs studying CDSS.

KQ1. Effectiveness

Process and other outcomes related to use and satisfaction with MMIT were often improved, especially for prescribing and ordering and the monitoring phases. Improvements in the appropriateness of prescribing and decreased errors (e.g., correct doses and timing, better choices of antibiotics, fewer drug-drug interaction potentials, and corrected doses related to body weight or liver function) seem to be consistently shown. Changes in workflow, improvements in communication, and improved efficiencies such as time reductions are also positive, although fewer studies addressed these types of outcomes. Clinical endpoints were sometimes found to be improved with the use of MMIT, more often in the observational studies than in controlled clinical trials. CDSS applications and, to a lesser extent, CPOE systems have been shown to be useful, especially when studying prescribing and monitoring in hospitals and clinics. Notable was the identification of strong emotions expressed by users of CPOE (clinicians), both positive and negative, which were reported in the qualitative studies. A number of unintended consequences of the technologies were found, some of which were unfortunate and some of which were beneficial. Few cost studies and full economic evaluations were identified. Those articles that were included found that health IT interventions may offer cost advantages despite their increased acquisition costs. Proof of clinical improvements and economic effectiveness through the use of MMIT is lacking. However, given the uncertainty that surrounds the cost and outcomes data, and limited study designs available in the literature, it is difficult to reach any definitive conclusion as to whether the additional costs and benefits represent value for money.

KQ2. Gaps in Evidence and Knowledge

The major gap in the research is true full economic evaluations, weighing all costs and benefits of the various MMIT technologies across all settings and participants. For the effectiveness research, we found gaps were related to setting (few studies were carried out in pharmacies, long-term care facilities, homes, or communities), people (few studies assessing outcomes for pharmacists, nurses, nurse practitioners, physician assistants, other health professionals including dentists and psychologists, or patients and families), and MMIT

technologies (rigorous studies of all but CDSS and CPOE, and especially those related to dispensing and administering, were sparse). Prescribing and monitoring were relatively well-studied while order communication, dispensing, administering, reconciliation, and education were understudied.

Gaps were also found in the sophistication and complexity of the quantitative research methods. Many of the studies initially identified were descriptive in nature. These are listed in the bibliography of this report. Qualitative studies and the quantitative studies that were hypothesis-based and comparative were analyzed. A good number of the studies, including those that were more strongly controlled (e.g., RCTs and cohort studies), often had methodology or reporting flaws or both including inconsistent use of standard methods for identifying and describing their methods, poorly justified or incorrect choices, or poor application of statistical tests and failure to adjust for group differences or cluster randomization. We also often found underpowered studies and situation-specific studies that were difficult to generalize or transfer to other settings or situations.

In addition, we found substantial deficiencies in reporting data important to the understanding of published studies. Although we identified data deficiencies in many aspects of studies, most serious were in descriptions of baseline data related to what was in place with respect to medication management in the health care setting before implementation of the MMIT system and descriptions of the MMIT implementation itself. Context is important for understanding studies and assessing their potential for application; detailed information on the setting and participants was also not often provided in studies.

KQ3. Value Proposition for Implementers and Users

Value propositions are determined by the balance of financial, clinical and organizational benefits. Limited data were available to address these issues comprehensively. Of note, we found that the various stakeholders had very different needs, perceptions, and access to MMIT systems and this must be addressed in valuing systems.

KQ4. System Characteristics

Different features of MMIT are important to various groups and settings. Very few studies (n = 21) reported on the specific feature sets of the systems being used and their links to purchase, implementation, and use. Few head-to-head comparisons using comparative effectiveness analysis methods, for example, were found. The evidence identified uses both qualitative and quantitative methods to gain an understanding of which features are important to users and stakeholders. Of note, we found that desired feature sets differed between the planning phase (perceived to be of value) and after implementation (based on actual use).

KQ5. Sustainability

Sustainability is vital to health IT. Before it can be fully understood and studied, it must be defined. For this document we chose to use a definition of sustainability that suggests sustainable systems are cost effective and clinically-effective. Because the evidence on economics data are lacking, we can add only a small amount of information on the sustainability of MMIT applications. Some data exist on effectiveness and use. We have included some data on patterns and characteristics that are important to use, including data on barriers and facilitators of successful implementations and ongoing system use. Use is higher in physicians, larger and

better funded organizations, hospital settings, some larger primary care groups, and in academic medical centers.

KQ6. Two-Way EDI

Very little evidence exists on bidirectional communication between pharmacists and physicians to enhance the order communication process. Extrapolation of data on one-way communication, factors that work to increase electronic communication on medications between prescribers and pharmacists are external incentives, a supportive regulatory environment, and existence of standards for prescribing EDI. Three factors work against effective EDI: incomplete consideration of the workflow and financial effects of e-Prescribing on pharmacists and pharmacies, regulatory and legal uncertainties, and low adoption rates of EDI capable EMR systems. Further development and evaluation of two-way EDI technologies for outpatient order communication regarding drugs is required to facilitate adoption. Pharmacies and pharmacists should take a more active role in the EDI development and evaluation process.

KQ7. RCTs in CDSS

CDSS applications are well-studied although problems with methods and reporting exist. CDSS is probably the best studied type of MMIT in terms of studies with strong methods and a sufficient number of studies to provide reliable answers to research questions of any of the MMIT applications. The first RCT was published in 1976, over 35 years ago. Of the 88 RCTs in this document, 77 are on CDSSs. The quality and sophistication of study methods, analysis, and reporting of the RCTs has improved over time, and there tends to be more measurement of clinical outcomes. However, evaluations of health care delivery, such as comparisons of effectiveness of treatment or prevention methods (e.g., drugs, services, and medical devices) are held to a higher standard than the types of reported research projects included in this evidence report. The studies in this report, while they met their own research objectives, collectively do not contain the research designs and associated clarity of findings to be able to definitively inform patients, clinicians, and policymakers regarding the effectiveness and overall impact of CDSS applications. Furthermore, the more rigorous and transferable research conducted tends to show no or limited effect on patient-important clinical outcomes.

This report was not designed to evaluate specific MMIT applications. In addition, MMIT interventions were not catalogued and characterized in great detail. Therefore we found no obvious themes that would suggest that a certain type of MMIT intervention with a certain type of implementation for a certain type of user in a particular setting would be successful. However, the following areas of commonality emerged in our analysis.

General

1. Research to date has concentrated on measurement of process changes and descriptive and pilot studies. In addition, some studies based on stronger methods have failed on issues such as adequate concealment of allocation and blinding, poor understanding of some methods, lack of adjustment of groups, and statistical challenges. Processes in health care are poor surrogates for clinical- or patient-important clinical outcomes, therefore it is important that new studies address clinical outcomes and use the most appropriate methods, and use them correctly, to adequately study MMIT applications. Researchers should also be encouraged to consider the generalizability or transferability of their results for all of their projects. Researchers in health IT could strengthen their

studies by using interdisciplinary teams with representation of multiple stakeholders, learning from other domains such as health technology assessment and economics, and with better reporting of their studies and results.

2. Standard and accepted definitions are lacking for MMIT applications, as well as standards for presenting the results of studies of health IT applications. Definitions are inconsistent for MMIT applications (e.g., e-Prescribing, CPOE, EMR, or EHR hospital information systems), study designs (e.g., observational or before-after), and outcomes (e.g., adverse drug events, adverse drug effects, prescribing errors, or errors per patient, 100 orders, day, hospital day, or physician). This has made identification of studies, data abstraction, synthesis of evidence, and presentation of findings challenging. Many study reports did not include important information that would have made this report stronger. Noticeable deficiencies centered on the MMIT application, its setting within the institution, training and implementation details, and maintenance and updating information. Professional associations interested in MMIT are pushing for standardization of definitions. AHRQ can join this movement for more standardization of terminology and definitions.

Effectiveness

1. Interventions most frequently targeted prescribing and monitoring stages of the medication use process.
2. Physicians who provided care in the hospital and ambulatory care settings were most likely to be the target of the intervention.
3. CDSS and CPOE applications were the most frequently studied type of health IT application studied. Seventy-seven of the 88 RCTs in this report study CDSSs.
4. Improvements in prescribing accuracy and decreased errors such as appropriate scheduling and choice of medications, prescribing taking into account weight-based dosing and dosing based on liver function, avoidance of drug-drug interactions and potential allergies and in being in accordance with guideline recommendations were consistently identified as improvements with the use of MMIT. Workflow, communication, interaction with peers and time considerations were found to be improved less often.
5. Studies that used health IT to identify and intervene on patients with actual problems (e.g., elevated blood pressure) or needed care (e.g., hemoglobin A_{1c} monitoring) appear to be more effective than CDSS approaches that identify potential problems (e.g., potential adverse drug events). This was particularly true when patient-centered principles were employed, such as providing patients with reminders and decision support recommendations about their current health status. However, this may be alternatively explained by the greater difficulty in measuring outcomes, such as potential for ADEs.
6. Studies that have been successful in improving a patient's clinical outcomes target high risk and vulnerable populations who have poor disease control,^{515,610,624,693} lack sufficient access to health care providers to manage their condition,⁴⁰⁸ or subpopulations with sufficient economic resources to respond to the CDSS intervention.⁶⁹⁴
7. The effect of similar CPOE systems on mortality can vary substantially as a function of the extent to which implementation strategies disrupt or delay critical activities in the clinical setting, or demand additional time for order-entry from clinical staff.

8. Highly targeted interventions, focused on specific medical problems appear to be demonstrated as more effective than more diffusely focused CDSS and CPOE. Again this may be due to the greater difficulty in measuring the outcomes of diffusely focused CDSS and CPOE in the generally smaller sample size (and inadequate power) studies that were identified.

Qualitative

1. No qualitative studies were identified that directly addressed the effect of an MMIT system on intermediate health care outcomes for any phase of the medication management continuum (prescribing, order communication, dispensing, administering and monitoring). Patient safety was the main health outcome mentioned in qualitative studies. Before MMIT implementation, most studies found that clinicians expected that the MMIT system would improve patient safety. Once implemented, most clinicians felt that MMIT did improve patient safety.
2. Differences in study outcomes for similar qualitative studies across settings were not apparent, suggesting that findings from qualitative studies could be transferrable across settings.
3. Despite the willingness of many of the participants to use a new MMIT system designed to improve prescribing and ordering of medications including CPOE, reservations were expressed by some implementers that the MMIT system and the resulting change in workflow would impair existing interactions and relationships among health care providers and between physicians and patients.
4. MMIT systems often substantially facilitated clinicians' monitoring of patients' adherence with their prescribed medication regimen.⁷³⁷ However, barriers were reported to using health IT systems for medication monitoring in some situations.⁷⁴⁷ For example, clinicians caring for patients with HIV/AIDS using a CPOE and CDSS system integrated with the hospital, pharmacy, and laboratory systems identified six barriers to using reminders, including workload, time to document, reminders that did not apply, inapplicability of reminders to the situation, lack of training to teach the users how best to use the new or modified system, quality of provider-patient interaction, and use of paper forms.⁷⁴⁷
5. From qualitative studies, system design including workflow changes, challenges with the system interface and new communication processes demonstrated that without adequate attention to system changes, the new kinds of medical errors with potential detrimental impact to patient safety could occur. Unintended negative consequences including the need to develop workarounds (one-off or nonstandardized changes) to workflow and the frustration generated in some studies with MMIT implementation are important to recognize and deal with to improve the success of implementation.
6. MMIT implementation did not just mean that a clinician needed to learn a new IT system but it also affected most of the other parts of the delivery of care processes, including how the interdisciplinary care team worked together.

Economics and Costs

1. Cost analyses can provide useful information on 'upfront' costs compared with 'downstream' cost avoidance if they explicitly measure all direct health care costs (e.g., capital costs, health professionals' time), direct nonhealth care costs (e.g., home care

services, transportation) as well as indirect costs (e.g., productivity gains or losses) that could be affected by the intervention of interest.

2. It is important to be aware that the greatest reported costs associated with these health IT are associated with the purchase of new software to add to preexisting EMR systems, as well as implementation costs (e.g., management, clinical team involvement, training costs) and maintenance costs. This assumes a large investment has already been made to purchase, implement, and maintain an MMIT system.
3. The full enumeration of the total costs needs to be synthesized with the consequences or outcomes of the intervention (i.e., cost-effectiveness analysis, cost-utility analysis, and cost-benefit analysis). Full economic evaluations linked to clinical outcomes need to be done.
4. Adoption of newer technologies needs to be based on formal evaluation of whether the additional health benefit (effectiveness) is worth the additional cost. Given the tension between the clinical benefits of CPOE and CDSS and the high up-front costs, decisionmakers deciding whether to implement CPOE and CDSS need to better understand how and when financial benefits of such systems accrue (e.g., short-term compared with long-term benefits). These types of analyses are important for well-informed decisionmaking.

Unintended Consequences

1. Unintended consequences, both positive and negative, were found across many of the studies as main endpoints, or were alluded to in others. Some were minor and some much more serious. A tracking system of major and clinically important unintended consequences would be useful for many audiences and should be considered by system developers and funding agencies.

CDSS

1. Many studies have evaluated CDSS tools for improving the effectiveness of anticoagulants (proportion of days in therapeutic range for anticoagulants) and improving the choice, route, duration of antibiotics, and reducing ADEs related to antibiotic use, and most are successful.
2. Sophisticated CDSS and other advanced clinical support should be built to insure added clinical value without being burdensome to those who use them. These sophisticated systems are difficult to develop, implement, and maintain.

Values

1. Values related to MMIT systems and implementations need to be determined from all stakeholders. Clinicians, administrators, and likely patients and their families have different values and place varying importance on each.

System Characteristics

1. Most often authors spoke about barriers and concerns towards implementation and acceptance, rather than characteristics of MMIT that could facilitate implementation, purchase, and use of such systems.

Two-Way EDI

1. Two-way EDI between prescribers and pharmacists is not common. Both facilitators and barriers exist that impact movement to implementation of e-Prescribing and two-way communication designed to enhance and streamline prescription optimization.

Limitations

Our review has a number of limitations. With the exception of PDA applications using patient-specific input, we focused on applications that enable medication management and that are integrated with other health IT systems. A number of technologies, such as smart intravenous pumps, bar-code scanners, and reporting systems for diabetes or asthma monitoring were not integrated with other health IT systems and were thus excluded. Indexing of individual articles in electronic databases is poor. Although we tried to be thorough in our search methods, we feel that we did not capture all potential articles—a very difficult task in new and multidisciplinary areas of study.

Further, we concentrated only on the main or major endpoints reported in studies with comparison groups and hypothesis testing. Given the heterogeneity in the literature, it was often difficult to discern main endpoints; where possible we determined main endpoints as those declared as such, or those that were the basis of power calculations (infrequently), or were stated to be main outcome measures in the abstract or objectives. We identified instances where the main endpoint was not clear. In these cases we gave priority to outcomes related to medication management and clinically important patient outcomes. We did not test the replicability of our abstraction of these outcomes.

Because of the lack of clear definitions on some of the technologies and issues associated with health IT, we were unable to address some key questions as thoroughly as we would have liked. This is especially noted in KQ5 relating to sustainability and KQ3 on value propositions. We feel that these are important issues for all health IT, that need to be addressed to effectively answer questions about ongoing use and effectiveness of these technologies.

It has proven difficult to synthesize the evidence on such a range of technologies, implemented in a number of settings and used by various stakeholders. Each intervention is so complex that it is often difficult to tell which studies are assessing the same processes. Also, outcome measures used by authors were variable. For example, similar outcomes such as prescribing changes were measured as changes in daily doses; prescribing rates per hospital, per physician, per 1,000 patient days, etc. The number of orders and compliance rates were difficult to extract and synthesize.

Our ability to draw conclusions is also reliant on the quality of the evidence we have found. In most cases, the research relies on observational studies, with RCTs and other methodologies with stronger controls only available on a select group of health ITs and phases of medication management. Even in the case of CDSSs, a lack of RCTs addressing electronic decision support integrated with other types of health IT still exists. Only a small minority of these studies focus on clinical outcomes—the endpoints that are most important to guide decisions by patients, providers and policymakers, about adopting these interventions. Furthermore, a very small number report improvement in these clinical outcomes.

We found great variation in the level of description of the health IT employed, with studies frequently lacking details on standards, hardware, integration, implementation dates and processes, and other similar factors. A large number of studies neglected to report the study dates

(see Evidence Tables in Appendix C). We repeat Chaudhry's call for a set of standards for reporting on health IT research.⁶⁰⁷

Although the absence of a contemporaneous comparable control group is a problem with all observational studies, the creation of control groups by comparing intervention patients to those that do not participate, or do not have a problem to those that do is fundamentally far more likely to introduce major bias in the comparison (e.g., comparing patients with alerts to no alerts,¹⁸ pharmacists volunteering to provide the intervention compared with those that do not volunteer,⁶⁹⁴ and other similar problems⁷⁰¹). The direction of the bias will depend on the study.

Many observational studies suffered from selecting an outcome that was distantly or only marginally related to the intervention. Length of stay, all-cause ADEs were examples of this problem. Gurwitz and colleagues⁶⁹⁷ were able to show that only one-third of ADEs could have been prevented by the CDSS alerts provided. Moreover, in a substantial proportion of negative studies, minimal adoption was seen. The clinicians failed to adjust therapy or treatment based on recommendations, and thus it is not very surprising to find that the interventions had no effect on outcomes. Finally, the rate of some outcomes such as readmission, mortality, and nosocomial infections was too low to detect clinically meaningful differences if they had existed with the numbers involved in the study.

Implications

The strength of this document lies in the breadth of health IT applications used across the phases of medication management, and in the organization of those findings, both through synthesizing the body of evidence by key questions and a tabular presentation of those findings. A review of this scope for MMIT has not been completed previously. We searched for literature across many domains and reviewed a substantial number of studies. The implications of the report fall within the purview of future research, policy, and evaluation. We have detailed gaps in evidence in KQ2 and future research needs in Chapter 5.

Important implications of this evidence report exist for health care decisionmakers, especially AHRQ and the U.S. National Coordinator of Information Technology. A large amount of health care spending in the United States is currently being funneled into development and implementation of various health ITs. Certainly the burden of evidence is towards positive effects on process changes and measures of satisfaction and perceived benefits among users. These early indications are logical precursors to changes in demonstrated effects in benefits such as quality of care and clinical outcomes, economic benefits, or both as the technologies advance and mature. A lack of proven effectiveness in improving patient outcomes and a lack of studies on value and cost-effectiveness still exist. Currently, most systems are in their infancy and need to be continued to be scrutinized for effectiveness and safety.

Because MMIT systems specifically, and health IT in general, are expensive to develop, support, and update, it is essential that these burgeoning health ITs be rigorously assessed for cost-effectiveness and clinical-effectiveness. This effectiveness information is essential for policymakers who are allocating scarce health care resources which have multiple competing priorities. Computerization of health care will continue with the adoption of more and newly developed MMIT and other health IT applications. Clinicians, researchers, policy advisors, and health administrators should be prepared for a major investment of time and resources for implementation and use. They need to consider direct and indirect effects on health care processes such as altered work flows, adverse patient outcomes, and indirect costs. Because of the paucity of successful clinical outcome studies, and the heterogeneity of the systems, the

specific interventions, and measures of effectiveness, this systematic review has been unable to clarify which factors of topic, design, or implementation may assist in the success of the MMIT.

Administrators will be able to plan for implementation better using the quantitative and qualitative findings and results. They will also be able to use this report to balance their expectations of MMIT installations and interact better with vendors and consultants.

Researchers and research funders will have a roadmap of the evidence that supports the effectiveness of MMIT applications, an outline of gaps, and lists of remaining challenges. Researchers should be aware of quality and reporting issues related to research methods as described in this review, as well as the need for research teams to include expertise or consultation from all clinician groups affected by the technology, informaticians, and those with research skills in a wide range of methodologies (research synthesis, complex interventions, pragmatic trials, usability studies, statistical planning and analysis, health technology assessment methods, and knowledge translation skills). Researchers and evaluators also need to adhere to established publication guidelines such as the STARE-HI guidelines⁷⁸³ for presenting results of their studies, to ensure that readers will have the information they need to plan for implementation of MMIT systems.

The meaningful use objectives should also be deployed in all projects and implementations. Research funders can direct their programs and reinforce use of standard definitions, reporting standards, and meaningful use objectives. They can also encourage multicenter trials and those that have potential for broad applicability. Adherence of the MMIT systems to local, regional, and national standards is also important to encourage and foster. At the same time, incremental studies which show the transferability and reproducibility of findings from one study to other health care settings, systems (vendors), and health care issues (type of disease or patient and setting) should also be encouraged.

Although the strength and breadth of the body of evidence supporting the usefulness of MMIT for improving health care is not uniform across people, places, and technology, it still is substantial. We can learn much from reviewing the original studies and systematic reviews on MMIT. We also feel that the content of this report can help us leverage our existing knowledge of MMIT to a broader audience and that this can improve the health and health IT effectiveness for many people in various health care settings.

Future Research

We reviewed a large body of literature from many domains. From a content point of view, medication management information technology (MMIT) is well-covered, although coverage in the literature is not uniform for all aspects of MMIT. Effective medication management is important for many people and costly for individuals and society. Medications themselves are changing and becoming more complex with the emergence of new drugs and the integration of health information and genomics research to set the stage for individualized health care. As the population ages, we start to rely more on medications, and polypharmacy becomes standard. At the same time that the management of drugs and medications is becoming more complex and costly, the move to health IT is occurring at an increasing rate and with increasing sophistication. Newer health IT applications hold tremendous potential for patients through their health care providers and also with the move to self-management of chronic diseases, patient-centered care, personal health record systems tethered to electronic medical record systems (EMRs), and automatic monitoring devices built into smart homes to increase and prolong independence.

We provide some future directions for consideration (Table 28). We saw much that was exciting and challenging in the evidential base of MMIT in this report. Future research should be conducted in those areas we have identified that can build on the existing evidence, address the gaps that have become evident, and to support trends that can improve the quality, efficiency, and cost of health care. The section on KQ2: Knowledge and Evidence Gaps has additional and supporting information.

Table 28. Issues of consideration and/or further exploration in future research

| |
|--|
| <p>Research Methods:</p> <ul style="list-style-type: none">• Research studies with control groups, statistically appropriate comparisons, and sufficient power and funding to produce unequivocal answers. These studies should recognize that MMIT applications need to be treated as complex interventions and evaluated as pragmatic studies (i.e., can they work in real life situations and settings).• We need large overarching trials of complete systems, and we also need smaller scale research and evaluation of the components of MMIT systems. Studies of components, such as two-way communication between pharmacists and prescribers or email between caregivers and patients are important to aid in our understanding of the contribution that each makes towards building a complete MMIT system (complex interventions).• Multicenter studies. Most studies seem to focus within a single organization using the same system and often done by those who built or developed the application. Multicenter studies can be supported, including involvement of centers that use different systems. A single study can yield valuable information about the system deployed as well as the organizational culture around the acceptance and use of the system, but understanding and enabling of generalizability or applicability and interoperability are more likely to occur with multicenter studies.• Studies and guidance of how best to conduct usability studies and how to make their results applicable and available to others with the same or similar applications, target populations, and clinical settings. Tool kits, training sessions, and encouragement to publish usability studies are important steps towards improved usability testing and transfer of knowledge related to the findings of these usability studies.• Adherence to standardized reporting and communication guidelines such as STARE-HI for published articles and technical reports. Consistency in reporting details of systems include substantial details and descriptions of the features and characteristics of the MMIT system, and how it fits into existing systems, priorities, and cultures of the institution; settings and user groups; exact details of the interaction of the system with clinicians and patients; and concise reporting of the outcomes assessed.• Research into studying how best to collect and analyze existing health data from patient care records (e.g., EMRs and EHRs) to produce new knowledge related to treatment outcomes, prognostic information and other related health issues. Newer methods to collect (harvest and analyze) research data from clinical health IT systems deserve further study taking into account ethics, privacy, and security issues." |
|--|

Table 28. Issues of consideration and/or further exploration in future research (continued)

Research Needs:

- Studies for order communication, dispensing and administering phases, and related aspects of medication management such as post-professional and professional education, electronic medication reconciliation, and health information exchange methods and standards.
- Studies in pharmacy settings to better understand how MMIT can be used to improve interprofessional communication, communication between pharmacists and patients, and prescribing outcomes.
- Studies that focus on patient-centered MMIT applications, such as medication adherence and automated and self-reported measures of monitoring medications tied to integrated systems.
- Studies of issues related to standards and interoperability and how these affect generalizability or transferability, and the spread of MMIT across institutions and geographic regions.
- Studies targeting nonphysicians including pharmacists, advanced practitioners (e.g., nurse practitioners and physician assistants), nurses, mental health professionals, and patients, as well as formal and informal caregivers who might use MMIT applications as part of providing care.
- RCTs and other studies with appropriate methods that concentrate, if possible, on clinical outcomes related to the use of medications and detailed costs. Special consideration needs to be given to adherence to accepted research methods and newer research methods such as cluster randomization.
- Studies of MMIT that leverage existing sources of electronic data such as clinical chemistry, hematology, and therapeutic drug monitoring across various health care settings to improve laboratory-based medication monitoring.
- Recognition that genomic data will likely have a major effect on choices of medications once the research has evolved to the extent where drug treatment decisions can be made for individuals based on their genetic profiles. This genomic information will become an essential part of the data in the next generation of CDSSs and will likely need to be evaluated as such.
- Qualitative and mixed methods studies on the effects of MMIT from the perspective of the patients and their existing needs and values and the implications of developing MMIT applications.
- Qualitative and mixed methods studies to provide a greater understanding of the role, function, and effects of MMIT on clinician workflow, inter- and intra-personal communication and satisfaction.
- Studies in older adults who reside in long-term care settings (e.g., nursing homes, assisted living, home-based primary care) and studies that centre on the geriatric population and those with complex care needs related to medications.
- Studies with pediatric populations in inpatient and outpatient settings.
- Improved methods of dissemination of MMIT research methodologies, strategies, and results. Those interested in MMIT can learn much from those who have expertise in clinical and translational research and knowledge translation (i.e., application of research findings) using improvement science principles.
- Comparative effectiveness research to compare the effect of more than one type of MMIT on process or outcomes.
- Data on unintended positive and negative consequences of MMIT applications should be collected and disseminated with priority given to those consequences that have substantial potential for harm or benefit or occur frequently.
- Sophisticated concurrent prospective economic evaluations conducted in the real world to address whether MMIT interventions are cost effective are vital for policymakers and decisionmakers.
- Studies of the ability to apply standard health technology appraisal methods to improve the ability to determine value for money of MMIT interventions to show if these methods should be adopted.
- Study the value of feature sets of the various technologies and how they impact purchasing and use for multiple MMIT intervention stakeholders and insure that results are applicable to multiple stakeholders. Studies must include multiple stakeholders: clinicians, other health care providers, patients, caregivers, administrators, vendors, computer programmers, etc.
- Study of how best to keep systems that rely on a strong knowledge base (e.g., CDSS, CPOE, order sets, drug-drug interaction programs) current with new scientific knowledge (i.e., guaranteeing the fidelity of evidence-based knowledge resources)
- Develop a more relevant operational definition of sustainability related to MMIT applications, and require future studies to state explicitly how they intend on studying and reporting on these results.
- Consensus meeting of experts on the types of preferred research methods to ascertain effectiveness and ensure production and reporting of quality evidence.
- Series of related studies, building sequentially, testing interventions across facilities, vendors, and settings to improve applicability and transferability of research findings.

Need for High Quality Evidence

High quality evidence is lacking in many MMIT phases, care settings, and populations (Table 29). Despite the fact that many RCTs exist in the MMIT literature, they are concentrated in certain areas: 77 of 88 RCTs evaluated CDSSs. The prescribing and monitoring phases have a strong base of studies and systematic reviews. Almost completely lacking were studies in the other phases. For this report we provide the numbers of studies and research methods used (Table 29). In addition, we used the bibliographies and summaries from more than 100 systematic and narrative review articles for this report.

Table 29. Study design of included studies across the medication management phases (plus education and reconciliation)

| Design | P | OC | D | A | M | E | R |
|---------------|-----|----|----|----|----|---|---|
| RCT | 70 | 2 | 3 | 2 | 37 | 1 | 1 |
| Cohort | 13 | 1 | 1 | 1 | 6 | 0 | 1 |
| Observational | 146 | 19 | 10 | 26 | 29 | 2 | 4 |
| Qualitative | 37 | 5 | 3 | 10 | 5 | 0 | 0 |
| Total | 266 | 27 | 17 | 39 | 77 | 3 | 6 |

Column Headings: P = Prescribing, OC = Order Communication, D = Dispensing, A = Administering, M = Monitoring, E = Education, R = Reconciliation

RCT Randomized controlled trial

Almost half of the articles that met the content criteria for MMIT did not include a comparison group, hypotheses testing or qualitative methods. These articles were not formally evaluated but are listed in the bibliography. We need well-designed research studies with control groups and appropriate analysis. Research needs center on issues related to:

- Appropriate methods for MMIT and the needs of the stakeholders. Important issues are multicentered studies, research teams with a broad experience or consultation in research methods, statistical analyses, clinical care, and informatics, and that represent the interests of all groups involved in the use of the MMIT system.
- Recognition that MMIT applications are complex interventions and their evaluation must reflect their use in real life settings and situations (i.e., complex interventions methods and pragmatic trials).
- Qualitative and mixed methods studies to understand issues related to workflow, communications, interdisciplinary collaboration and care processes, and patient use and values.

We also found that certain technologies (CDSSs, either stand-alone or integrated with CPOE systems) are well-studied. Although much data exist, as these CDSS and CPOE systems evolve and have greater penetration among health care settings, we should continue to evaluate their effectiveness. The tools that are outside the prescribing phase of medication management and the health IT tools that pharmacists, nurses, other health care professionals, and patients use are less well-studied—fewer studies and weaker methods.

Need for Well-Designed Research

Despite having 88 RCTs in the MMIT literature base, many of the studies have weaknesses. This is shown by the low quality scores, most of which were in the range of four to five out of nine points. In addition, we saw errors and poor methods in published studies. For example, most of the RCTs with clinical outcomes (n = 28) used cluster randomization methods to allocate

clinical units or clinicians to study groups, but analyzed and reported results based on patients or medication events. Many authors did not test or adjust for clustering so that complex analyses could be accomplished appropriately. We also identified problems with poor application of methods in most other research studies.

Training informaticians in research methodology and statistical methods is crucial. Many programs sponsored by the U.S. National Library of Medicine and other institutions are graduating health informaticians. Training programs are content-rich because of the breadth of the field. Formal training and experience in the research methods and statistical analyses components of the training initiatives might be useful to determine what is being taught and if it is sufficient to produce researchers who are competent in evaluating MMIT and other health IT systems.

By settings and levels of care. Adult hospitals were relatively well-studied and we have sufficient evidence to show that MMIT systems for ordering medications improve processes and reduce medication errors. Adult ambulatory care clinics were also well-represented in the literature, although studies of errors and error prevention have not been done. Additional studies are especially needed in the nursing home setting, where some 1.6 million people receive care annually, and concern continues about the quality of pharmaceutical care, the frequency of polypharmacy, and an insufficient health care workforce with a poorly developed safety culture. Other long-term care settings such as assisted living and home-based primary care also need more research. The number of older adults continues to increase rapidly and they frequently have multiple comorbidities resulting in complex medication regimens, polypharmacy, and ADEs. Studies conducted in pediatric hospitals are warranted because these patients are particularly vulnerable to medication errors and those medication errors that do occur have three times the potential to cause harm.⁸⁴⁷ Community pharmacies and the newer mail-order and online pharmacy services were not studied. Evaluating these settings may be problematic because of their commercial nature. Homes and other residential or community settings will become more important to study with the spread of patient-centered medicine and associated technologies such as PHRs and remote MMITs.

Monitoring. Our data suggest that interventions that focused on *laboratory-based* medication monitoring (22 of 29 studies) were associated with the most number of interventions, and showed statistically significant changes in at least half of its main endpoints. We recommend additional research in this area especially because laboratory data are readily available in most clinical settings, and studies in the acute, ambulatory, and nursing home settings suggest that failure to act on available laboratory information accounts for a substantial number of ADEs.⁸⁴⁸⁻⁸⁵⁰ With the integration of more health IT systems and the move to more patient directed care, systematic monitoring will become even more important.

Practitioners and patients. Nurses and pharmacists are not studied as thoroughly as physicians. Mental health professionals and other health care workers who prescribe, including dentists, are studied even less than nurses and pharmacists. Each group of health professionals reports different needs for their MMIT and health IT tools (e.g., specialist physicians compared with primary care, nurses compared with physicians in hospital wards need compatible but different MMIT tools for ordering, dispensing, and administering). These differences need to be studied and applied in building, evaluating, and implementing MMIT applications. Nurse practitioners,

advance practice nurses and physician assistants, and allied and other health professionals should also be the target of MMIT interventions, especially because they play an increasing role in providing primary and subspecialty care, especially in the United States. The move to patient-centered care and chronic disease management also make the study of patients and their informal caregivers an important area for research and development.

Unique needs of evaluation of health IT. MMIT applications are neither simple nor isolatable components that can be easily studied as such. Research methods to evaluate MMIT applications should be based on principles of complex interventions. For example, the U.K. Medical Research Council provides a framework for individuals to consider when planning complex intervention projects (<http://www.mrc.ac.uk/Utilities/Documentrecord/index.htm?d=MRC003372>).

In addition, MMIT applications are often being changed and modified to address problems or to implement fixes or upgrades. This ever changing aspect of health IT poses challenges for health researchers. Classical evaluation and research methods dictate that what is being evaluated needs to be stable over the time period of the study. “Fast” analysis methods may need to be developed.

MMIT systems, as for any IT system, are easiest to study in laboratory settings. Because health care is so complex, the study of MMIT systems must be done in real settings. Pragmatic trials methods (trials done in real life situations) may also need to be applied in many research projects related to MMIT systems.

Commercial interests also complicate the evaluation of MMIT and health IT. This makes research harder to do and provides barriers to the most common government-based funding sources. AHRQ has provided funding for many unique and valuable health IT applications that have included much evaluation of MMIT. They should be encouraged to continue to provide leadership in this domain.

Another challenge to research methods is that often the existing evaluations have been done by system developers or implementers. This ownership can cloud an evaluator’s vision (i.e., bias) or at least stand in the way of publication of negative findings. Some evidence exists that evaluation of one’s own system contributes to biases towards the system being found to be positive.⁷²⁵

Knowledge translation/translational research. Because many of the studies in MMIT are small descriptive studies and often done by developers, generalizability or transferability are often not ideal. Researchers in this domain of “getting research into practice” (i.e., knowledge translation or translational research) can provide tools and insights in two areas. The first is how to harness existing knowledge tool development such as building the knowledge base behind health IT systems such as CDSS, CPOE systems, and other knowledge summaries such as clinical practice guidelines and order sets. Second, those involved in knowledge translation can assist in the dissemination of studies of MMIT applications, or provide leadership in how we can disseminate studies of health IT.⁷²⁵

MMIT applications do not stand alone. MMIT implementation has had substantial impact on communication, interpersonal, and inter-professional relationships, and development of unintended consequences. In some cases, issues such as rage against the machine, guilt, embarrassment associated with reminders and alerts, and frustration have been reported. The qualitative and mixed methods studies summarized in the KQ1: Effectiveness section are

examples of studies that have shown how MMIT and health IT can and do affect individuals on a personal level. More of these studies of the effects of these technologies on people, clinicians, and individuals need to be done in various settings and with all technologies. Workflow and communication are ideally studied using qualitative and mixed methods.

Sustainability. Sustainability is tremendously important to MMIT and other health IT applications. See also KQ5: Sustainability. We could not find an agreed upon definition and used one from Australia: “the ability of a health service to provide ongoing access to appropriate quality care in a cost effective and health-effective manner.” The informatics domain needs to have an agreed upon definition of sustainability. Once this is established, research needs to be done to identify our current “sustained” systems and determine the factors that are associated with them. Qualitative and quantitative studies are essential and they need to be done by people with strong content and methods background and sufficient financial backing. Partnerships among Federal groups (e.g., AHRQ, Office of the National Coordinator for Health Information Technology, Health Resources and Services Administration, and Centers for Medicare and Medicaid Services), vendors, professional organizations (e.g., Healthcare Information and Management Systems Society, American Medical Informatics Association, and the major pharmacy associations such as the American Pharmacists Association, American Society of Health System Pharmacists, Academy of Managed Care Pharmacy, American College of Clinical Pharmacy, American Society of Consultant Pharmacists, National Community Pharmacists Association, and National Association of Chain Drug Stores), researchers, and others could work together to address the sustainability challenge. We also need studies of successful MMIT applications as well as systematic study of failures. Perhaps the HITECH Act of 2009 will lead to improvements and sustainability of health IT applications that specifically support the medication management continuum through meaningful use.

Standards and certification. We were asked to provide the evidence on the influence of standards and certification and how they affect MMIT systems. This evidence is sorely lacking. Standards are necessary for interoperability and smooth functioning of existing systems and large scale integration of data at State and national levels. Leadership, probably more than research efforts, continues to be needed in this domain.

Measurement and definitional issues. Other gaps in the evidence that need addressing are definitional or measurement issues. Because health IT is an interdisciplinary field, standard definitions are crucial. Producers and users of research and evaluations function best when everyone is using the same terms with the same parameters. One simple example we found is a formal working definition of the difference between an e-Prescribing system and a medication-based CPOE system. Some European literature described a system as e-Prescribing, while the same system in the United States would be classified as COPE. Another idea that seemed to cause confusion among authors and readers is the use of EMR or EHR systems, and hospital information systems. Consistency in reporting and communicating MMIT information is also important.

Clinical practice guidelines, CPOE, and CDSS. One final issue that seems unresolved centers on the evidentiary nature and strength of the knowledge base that forms the foundation of CDSS applications, order sets, and most other MMIT systems. Some systems linked to established

clinical practice guidelines, but we did not find studies that addressed the strength of the evidence base of MMIT systems. We feel that a strong, reliable, consistent, fully disseminated, and continually updated evidence base for MMIT and other health IT systems is vital. More emphasis has been placed on the mechanics of these systems than content. Establishment of standards and content for the knowledge base is something that is potentially more important than the mechanics of these decision-support systems. The U.S. National Library of Medicine could provide leadership here. They have already built strong knowledge management tools such as their Unified Medical Language System that knits together multiple vocabularies in a machine processable form. They have also developed other information handling and processing tools, and techniques such as natural language processing capabilities of medical text, RxNorm (a standardized electronic nomenclature for clinical drugs and drug delivery devices), and codified drug allergy information provision and transfer. Their work in genomics and proteomics is also important once an individual's genetic information is ready for useful integration into our health IT and MMIT systems to provide individualized medicine.

Conclusions

Our evidence review on the use of health information technology (health IT) in enabling medication management derives from a summary of 428 studies assessing the use of integrated health IT that assisted with at least one phase of the medication management process, and associated aspects of education and reconciliation. We define the medication management process as having five phases; prescribing, order communication, dispensing, administering, and monitoring. For this report we also included medication reconciliation, education, and adherence.

Key Question (KQ) 1: Effectiveness assessed the effectiveness of health IT on changes in process and intermediate, cost, and clinical outcomes. We limited our studies to those that used qualitative methods or included comparison groups, hypothesis testing, and appropriate statistical analysis. We did this in an effort to limit inclusion to studies that used research methods and had data that we could use to draw conclusions. The majority (378 of 428 studies) of the evaluated studies were included in KQ1: Effectiveness. Even so, our findings indicate that RCTs and other methodologies with controlled populations are lacking in adequate details and robust methods, which result in an only incremental addition to the evidence base for the use of MMIT. Most of the studies in this evidence report are quantitative observational assessments, often using historical controls.

The evidence from these studies indicate positive effects on improving process, often measured as improvements in medication orders during the prescribing and monitoring phases. The bulk of this evidence of improvement is shown in studies set in hospitals. We also found improvements associated with MMIT systems related to use, usability, knowledge, skills, and attitudes. These cumulated changes can, but may not always, lead to efficiency and cost gains. On the other hand, little work is being done on the other phases of medication management with integrated health IT. Some IT applications used in dispensing and administering are stand-alone technologies and, by definition, not included in this report.

We found little evidence of significant effects on clinical outcomes; possibly because of the small number of events, the outcomes being far removed from the application of the technology; or that they were often not the main endpoints of the studies included in the review. We do not know if MMIT applications are clinical- and cost effective because of a lack of sound economic data.

The qualitative literature highlighted positive and negative perceptions and satisfaction with the integrated health IT applications, supporting much of the literature on the importance of the effects of the technologies on workflow and the working relationships of the users.

We have identified a number of gaps in the evidence of the effects of MMIT applications: most notably the order communication, dispensing, and administering, as well as reconciliation and education. Inpatient care is well-studied, followed by ambulatory care. A low number of studies assessed long-term care and effects on pharmacies, especially those outside the hospital setting. Gaps in research also exist for studies that evaluate MMIT that are not computerized decision support systems (CDSSs) or computerized provider order entry (CPOE) systems. The domain of patient and informal caregiver access to MMIT applications, especially those applications that are integrated with such existing clinical applications as electronic medical record systems (EMRs), will be an exciting and promising new domain of study. One major gap is the assessment of MMIT tools that are used by nonphysicians.

The value of the MMIT systems needs to be summed up across financial, clinical, and organizational components. The values proposition for each stakeholder will be different based

on their own value set, and what is important to each has not been well-studied. Though some evidence suggests positive financial and organizational gains, these gains are not universal and will depend on the technology, the setting, and the impact on the stakeholders using them. Clinical benefit is proving difficult to assess. Rigorous studies are needed to truly assess economic and other values.

This broad scoped review sought to include a large number of health ITs across an array of settings and users. As such, the literature was heterogeneous and difficult to synthesize. Based on our findings, we feel that it is important to note that the burden of proof of the value of MMIT is somewhat limited, but promising. We feel also that decisionmakers must be aware of the potential for negative impacts of the technologies and carefully consider these possibilities during implementation and provide for continued monitoring across all stakeholder groups. . The evidence base of MMIT applications is strong and varied, and it can be further strengthened by using multicentered studies, building an integrated body of evidence which will demonstrate the transferability and applicability of the systems, and multidisciplinary teams of researchers, or at least consultation input from clinicians; methodologists, including biostatisticians; informaticians; particular health IT users and representatives of various stakeholders. We can also learn much from those who work with complex interventions, pragmatic trials, research syntheses, knowledge translation or translation research challenges, and health technology assessment studies to enhance the construction, conduct, and communication of health IT implementation and use research findings.

References

1. Bell DS, Cretin S, Marken RS, et al. A conceptual framework for evaluating outpatient electronic prescribing systems based on their functional capabilities. *J Am Med Inform Assoc* 2004;11(1):60-70.
2. Shekelle PG, Morton SC, Keeler EB. Costs and Benefits of Health Information Technology. Evidence Report/Technology Assessment No. 132. AHRQ Publication No.06-E006. Rockville, MD: Agency for Healthcare Research and Quality. April 2006.
3. Jimison H, Gorman P, Woods S, et al. Barriers and Drivers of Health Information Technology Use for the Elderly, Chronically Ill, and Underserved. Evidence Report/Technology Assessment No. 175 (Prepared by the Oregon Evidence-based Practice Center under Contract No. 290-02-0024). AHRQ Publication No. 09-E004. 175. Rockville,MD: Agency for Healthcare Research and Quality. November 2008. Available at: <http://www.ahrq.gov/clinic/tp/hitbartp.htm>
4. Gibbons MC, Wilson RF, Samal L et al. Impact of Consumer Health Informatics Applications. Evidence Report/Technology Assessment No. 188. AHRQ Publication No. 09(10)-E019. Agency for Healthcare Research and Quality. October 2009. Available at: <http://www.ahrq.gov/downloads/pub/evidence/pdf/chiapp/impactchia.pdf>
5. Hersh WR, Hickam DH, Severance SM et al. Telemedicine for the Medicare Population: Update. Evidence Report/Technology Assessment No. 131. AHRQ Publication No. 06-E007. Rockville, MD: Agency for Healthcare Research and Quality: 2006.
6. Agency for Healthcare Research and Quality. Enabling Health Care Decision Making through the Use of Health Information Technology (Health IT). <http://www.ahrq.gov/clinic/tp/knownmgtp.htm>.
7. Agency for Healthcare Research and Quality. Enabling Patient-Centered Care through Health Information Technology (Health IT). <http://www.ahrq.gov/clinic/tp/pcchittp.htm>.
8. Lohbach D. Enabling Health Care Decision Making through the Use of Health Information Technology (Health IT). due December 2010. AHRQ; 2010. <http://www.ahrq.gov/clinic/tp/knownmgtp.htm>
9. Humphreys JS, Wakerman J, Wells R. What do we mean by sustainable rural health services? Implications for rural health research. *Aust J Rural Health* 2006;14(1):33-5.
10. Verhagen A, de Vet H, de Bi R, et al. The Delphi List: A Criteria List for Quality Assessment of Randomized Clinical Trials for Conducting Systematic Reviews Developed by Delphi Consensus. *J Clin Epidemiol* 1998;51(12):1235-41.
11. Perras C, Jacobs P, Boucher M et al. Technologies to reduce errors in dispensing and administration of medication in hospitals: Clinical and economic analyses. Technology Report Number 121. Ottawa: Canadian Agency for Drugs and Technologies in Health; 2009.
12. Morriss F, Abramowitz P, Nelson S, et al. Effectiveness of a Barcode Medication Administration System in Reducing Preventable Adverse Drug Events in a Neonatal Intensive Care Unit: A Prospective Cohort Study. *J Pediatr* 2009;154(3):363-8.
13. Grasso BC, Genest R, Yung K, et al. Reducing errors in discharge medication lists by using personal digital assistants. *Psychiatr Serv* 2002;53(10):1325-6.
14. Poole D, Chainakul J, Pearson M, et al. JHQ 177 medication reconciliation: A necessity in promoting a safe hospital discharge. *J Healthc Qual* 2006;28(3):12-9.
15. Han YY, Carcillo JA, Venkataraman ST, et al. Unexpected increased mortality after implementation of a commercially sold computerized physician order entry system. *Pediatrics* 2005;116(6):1506-12.

16. Keene A, Ashton L, Shure D, et al. Mortality before and after initiation of a computerized physician order entry system in a critically ill pediatric population. *Pediatr Crit Care Med* 2007;8(3):268-71.
17. Gesteland PH, Nebeker JR, Gardner RM. These are the technologies that try men's souls: common-sense health information technology. *Pediatrics* 2006;117(1):216-7.
18. Lesprit P, Duong T, Girou E, et al. Impact of a computer-generated alert system prompting review of antibiotic use in hospitals. *J Antimicrob Chemother* 2009;63(5):1058-63.
19. van Ast JF, Talmon JL, Renier WO, et al. Development of diagnostic reference frames for seizures. Part 2: are seizure descriptions discriminative? *Int J Med Inf* 2003;70(2-3):293-300.
20. Weir C, Lincoln M, Roscoe D, et al. Dimensions associated with successful implementation of a hospital based integrated order entry system. *Proc Annu Symp Comput Appl Med Care*. 1994:653-7.
21. Goldzweig CL, Towfigh A, Maglione M, et al. Costs and benefits of health information technology: new trends from the literature: since 2005, patient-focused applications have proliferated, but data on their costs and benefits remains sparse. *Health Aff (Millwood)* 2009 Mar-Apr;28(2): Supplement 1: w282-93:Supplement-93.
22. Truffer C, Keenan S, Smith S, et al. Health spending projections through 2019: The recession's impact continues. *Health Aff (Millwood)* 10 A.D.;29(3):522-9.
23. Agency for Healthcare Research and Quality. Prescription Medicines-Mean and Median Expenses per Person with Expense and Distribution of Expenses by Source of Payment: United States, 2007. 2007. Agency for Healthcare Research and Quality, Prescription Medicines-Mean and Median Expenses per Person with Expense and Distribution of Expenses by Source of Payment: United States, 2007. Medical Expenditure Panel Survey Component Data
24. Henry J.Kaiser Family Foundation. Prescription drug trends. September 2008. http://www.kff.org/rxdrugs/upload/3057_07.pdf.
25. Aspden P, Wolcott JA, Bootman L, et al. Preventing Medication Errors. 1. Washington: National Academies Press; 2007.
26. Kozer E, Scolnik D, Macpherson A, et al. Variables associated with medication errors in pediatric emergency medicine. *Pediatrics* 2002;110(4):737-42.
27. Cheung KC, Bouvy ML, De Smet PA. Medication errors: the importance of safe dispensing. *Br J Clin Pharmacol* 2009;67(6):676-80.
28. Yu VL, Fagan LM, Wraith SM, et al. Antimicrobial selection by a cexpertsomputer: A blinded evaluation by infectious disease. *JAMA* 1979;242(12):1279-82.
29. Gehlbach SH, Wilkinson WE, Hammond WE, et al. Improving drug prescribing in a primary care practice. *Med Care* 1984;22(3):193-201.
30. Executive preview: The value of CPOE in ambulatory settings. 2002. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=29895>
31. Mollon B, Chong J, Jr., Holbrook AM, et al. Features predicting the success of computerized decision support for prescribing: a systematic review of randomized controlled trials. *BMC Med Inform Decis Mak* 2009;9:11
32. Eslami S, de Keizer NF, Abu-Hanna A. The impact of computerized physician medication order entry in hospitalized patients--a systematic review. *Int J Med Inf* 2008;77(6):365-76.
33. Kaushal R, Kern L, Barrón Y, et al. Electronic prescribing improves medication safety in community-based office practices. *J Gen Intern Med* 2010;25(6):530-6.
34. Paoletti RD, Suess TM, Lesko MG, et al. Using bar-code technology and medication observation methodology for safer medication administration. *Am J Health Syst Pharm* 2007;64(5):536-43.
35. Sampson M, McGowan J, Lefebvre C et al. PRESS: Peer Review of Electronic Search Strategies. Ottawa: Canadian Agency for Drugs and Technologies in Health; 2008.

36. Hanuscak TL, Szeinbach SL, Seoane-Vazquez E, et al. Evaluation of causes and frequency of medication errors during information technology downtime. *Am J Health Syst Pharm* 2009;66(12):1119-24.
37. Fowler SB, Sohler P, Zarillo DF. Bar-code technology for medication administration: medication errors and nurse satisfaction. *Medsurg Nurs* 2009;18(2):103-9.
38. Lesselroth BJ, Felder RS, Adams SM, et al. Design and implementation of a medication reconciliation kiosk: the Automated Patient History Intake Device (APHID). *J Am Med Inform Assoc* 2009;16(3):300-4.
39. Pattacini C, Rivolta GF, Di Perna C, et al. A web-based clinical record 'xl'Emofilia' for outpatients with haemophilia and allied disorders in the Region of Emilia-Romagna: features and pilot use. *Haemophilia* 2009;15(1):150-8.
40. Di Pentima MC, Chan S, Eppes SC, et al. Antimicrobial prescription errors in hospitalized children: role of antimicrobial stewardship program in detection and intervention. *Clin Pediatr (Phila)* 2009;48(5):505-12.
41. Gardner B, Graner K. Pharmacists' medication reconciliation-related clinical interventions in a children's hospital. *Jt Comm J Qual Patient Saf* 2009;35(5):278-82.
42. Stock R, Scott J, Gurtel S. Using an electronic prescribing system to ensure accurate medication lists in a large multidisciplinary medical group. *Jt Comm J Qual Patient Saf* 2009;35(5):271-7.
43. Lesselroth B, Adams S, Felder R, et al. Using consumer-based kiosk technology to improve and standardize medication reconciliation in a specialty care setting. *Jt Comm J Qual Patient Saf* 2009;35(5):264-70.
44. Murphy DM, Vercruyse RA, Bertucci TM, et al. Reducing hyperglycemia hospitalwide: the basal-bolus concept. *Jt Comm J Qual Patient Saf* 2009;35(4):216-23.
45. Barnett J, Jennings H. Pharmacy information systems in Canada. *Stud Health Technol Inform* 2009;143:131-5.
46. Bedouch P, Allenet B, Grass A, et al. Drug-related problems in medical wards with a computerized physician order entry system. *J Clin Pharm Therapeut* 2009;34(2):187-95.
47. Bepko RJ, Jr., Moore JR, Coleman JR. Implementation of a pharmacy automation system (robotics) to ensure medication safety at Norwalk hospital. *Qual Manag Health Care* 2009;18(2):103-14.
48. Pagan JA, Pratt WR, Sun J. Which physicians have access to electronic prescribing and which ones end up using it? *Health Policy* 2009;89(3):288-94.
49. Weir CR, McCarthy CA. Using implementation safety indicators for CPOE implementation. *Jt Comm J Qual Patient Saf* 2009;35(1):21-8.
50. Agrawal A, Wu WY. Reducing medication errors and improving systems reliability using an electronic medication reconciliation system. *Jt Comm J Qual Patient Saf* 2009;35(2):106-14.
51. Cummings A, Parker CD, Kwapniowski LA, et al. Using mobility technology to improve pharmacist workflow in the PICU rounding process. *J Healthc Inf Manag* 2009;22(4):39-43.
52. Denai MA, Mahfouf M, Ross JJ. A hybrid hierarchical decision support system for cardiac surgical intensive care patients. Part I: Physiological modelling and decision support system design. *Artif Intell Med* 2009;45(1):35-52.
53. Warholak TL, Rupp MT. Analysis of community chain pharmacists' interventions on electronic prescriptions. *J Am Pharm Assoc* 2009;49(1):59-64.
54. Karuppan M, Karuppan CM, Craig C. Point-of-care technologies: a case for resource integration. *Health Care Manager* 2009;28(1):38-43.
55. Mille F, Schwartz C, Brion F, et al. Analysis of overridden alerts in a drug-drug interaction detection system. *Int J Qual Health Care* 2008;20(6):400-5.
56. Neafsey PJ, Anderson E, Peabody S, et al. Beta testing of a network-based health literacy program tailored for older adults with hypertension. *Comput Inform Nurs* 2008;26(6):311-9.

57. Huntman L, Ward L, Read D, et al. Analysis of allergy alerts within a computerized prescriber-order-entry system. *Am J Health Syst Pharm* 2009;66(4):373-7.
58. Mannheimer B, Ulfvarson J, Eklof S, et al. A clinical evaluation of the Janus Web Application, a software screening tool for drug-drug interactions. *Eur J Clin Pharmacol* 2008;64(12):1209-14.
59. Eguale T, Tamblyn R, Winslade N, et al. Detection of adverse drug events and other treatment outcomes using an electronic prescribing system. *Drug Saf* 2008;31(11):1005-16.
60. Park CS, Kim TB, Kim SL, et al. The use of an electronic medical record system for mandatory reporting of drug hypersensitivity reactions has been shown to improve the management of patients in the university hospital in Korea. *Pharmacoepidemiol Drug Saf* 2008;17(9):919-25.
61. Markert A, Thierry V, Kleber M, et al. Chemotherapy safety and severe adverse events in cancer patients: strategies to efficiently avoid chemotherapy errors in in- and outpatient treatment. *Int J Cancer* 2009;124(3):722-8.
62. Sard BE, Walsh KE, Doros G, et al. Retrospective evaluation of a computerized physician order entry adaptation to prevent prescribing errors in a pediatric emergency department. *Pediatrics* 2008;122(4):782-7.
63. Munger MA, Stoddard GJ, Wenner AR, et al. Safety of prescribing PDE-5 inhibitors via e-medicine vs traditional medicine. *Mayo Clin Proc* 2008;83(8):890-6.
64. Meidl TM, Woller TW, Iglar AM, et al. Implementation of pharmacy services in a telemedicine intensive care unit. *Am J Health Syst Pharm* 2008;65(15):1464-9.
65. van der SH, Aarts J, van Gelder T, et al. Turning off frequently overridden drug alerts: limited opportunities for doing it safely. *J Am Med Inform Assoc* 2008;15(4):439-48.
66. Tamblyn R, Huang A, Taylor L, et al. A randomized trial of the effectiveness of on-demand versus computer-triggered drug decision support in primary care. *J Am Med Inform Assoc* 2008;15(4):430-8.
67. Bonnabry P, Despont-Gros C, Grauser D, et al. A risk analysis method to evaluate the impact of a computerized provider order entry system on patient safety. *J Am Med Inform Assoc* 2008;15(4):453-60.
68. Toth-Pal E, Wardh I, Strender LE, et al. A guideline-based computerised decision support system (CDSS) to influence general practitioners management of chronic heart failure. *Inform Prim Care* 2008;16(1):29-39.
69. Jaspers MW, Peute LW, Lauteslager A, et al. Pre-post evaluation of physicians' satisfaction with a redesigned electronic medical record system. *Stud Health Technol Inform* 2008;136:303-8.
70. Fursse J, Clarke M, Jones R, et al. An automated personalised intervention algorithm for remote patient monitoring. *Stud Health Technol Inform* 2008;136:181-6.
71. Nolen AL, Rodes WD. Bar-code medication administration system for anesthetics: effects on documentation and billing. *Am J Health Syst Pharm* 2008;65(7):655-9.
72. Chuo J, Hicks RW. Computer-related medication errors in neonatal intensive care units. *Clin Perinatol* 2008;35(1):119-39.
73. Lapinsky SE, Holt D, Hallett D, et al. Survey of information technology in Intensive Care Units in Ontario, Canada. *BMC Med Inform Decis Mak* 2008;8:5
74. Rose AJ, Shimada SL, Rothendler JA, et al. The accuracy of clinician perceptions of "usual" blood pressure control. *J Gen Intern Med* 2008;23(2):180-3.
75. Glintborg B, Poulsen HE, Dalhoff KP. The use of nationwide on-line prescription records improves the drug history in hospitalized patients. *Br J Clin Pharmacol* 2008;65(2):265-9.
76. Martens JD, van der WT, Winkens RA, et al. Feasibility and acceptability of a computerised system with automated reminders for prescribing behaviour in primary care. *Int J Med Inf* 2008;77(3):199-207.
77. Gilhuly TJ, Hutchings SR, Dumont GA, et al. Development and pilot testing of the Neuromuscular Blockade Advisory System. *Comput Methods Programs Biomed* 2008;89(2):179-88.

78. Krogh PR, Rough S, Thomley S. Comparison of two personal-computer-based mobile devices to support pharmacists' clinical documentation. *Am J Health Syst Pharm* 2008;65(2):154-7.
79. Pham DQ, Pham AQ, Ullah E, et al. Evaluating the appropriateness of thromboprophylaxis in an acute care setting using a computerised reminder, through order-entry system. *Int J Clin Pract* 2008;62(1):134-7.
80. Siteman E, Businger A, Gandhi T, et al. Physicians value patient review of their electronic health record data as a means to improve accuracy of medication list documentation. *AMIA* 2007;1116
81. Huang Y, Reichley RM, Noirod LA, et al. Automated dose checking and intervention for bariatric patients. *AMIA* 2007;983
82. Heard KM, Hollands JM, Noirod LA, et al. Alerts to improve chart documentation for National Quality Measures. *AMIA* 2007;971
83. Collins S, Bakken S, Cimino JJ, et al. Information needs related to antibiotic prescribing while using CPOE. *AMIA* 2007;916
84. Aronsky D, Johnston PE, Jenkins G, et al. The effect of implementing computerized provider order entry on medication prescribing errors in an emergency department. *AMIA* 2007;863
85. Stutman HR, Fineman R, Meyer K, et al. Optimizing the acceptance of medication-based alerts by physicians during CPOE implementation in a community hospital environment. *AMIA* 2007;701-5.
86. Linder J, Schnipper JL, Volk LA, et al. Clinical decision support to improve antibiotic prescribing for acute respiratory infections: results of a pilot study. *AMIA* 2007;468-72.
87. Lerma G, V, Poveda Andres JL, Font N, I, et al. Alerts system associated with computerized physician order entry: analysis and identification of improvement points. *Fam Hosp* 2007;31(5):276-82.
88. Nugent C, Finlay D, Davies R, et al. The next generation of mobile medication management solutions. *Int J Electron Healthc* 2007;3(1):7-31.
89. Trivedi MH, Rush AJ, Gaynes BN, et al. Maximizing the adequacy of medication treatment in controlled trials and clinical practice: STAR(*)D measurement-based care. *Neuropsychopharmacology* 2007;32(12):2479-89.
90. Seidling HM, Al Barmawi A, Kaltschmidt J, et al. Detection and prevention of prescriptions with excessive doses in electronic prescribing systems. *Eur J Clin Pharmacol* 2007;63(12):1185-92.
91. Lapane KL, Dube CE, Schneider KL, et al. Misperceptions of patients vs providers regarding medication-related communication issues. *Am J Manag Care* 2007;13(11):613-8.
92. FitzHenry F, Peterson JF, Arrieta M, et al. Medication administration discrepancies persist despite electronic ordering. *J Am Med Inform Assoc* 2007;14(6):756-64.
93. Grizzle AJ, Mahmood MH, Ko Y, et al. Reasons provided by prescribers when overriding drug-drug interaction alerts. *Am J Manag Care* 2007;13(10):573-8.
94. La C, V, Hellstern-Hauerslev C. Evaluating the implementation and use of a computerized physician order entry system: a case study. *Stud Health Technol Inform* 2007;130:75-9.
95. Kuno E, Hadley TR, Rothbard AB. Costs of implementing a computerized prescription system in a public mental health agency. *Psychiatr Serv* 2007;58(10):1351-4.
96. Rocha BH, Langford LH, Towner S. Computerized management of chronic anticoagulation: three years of experience. *Stud Health Technol Inform* 2007;129(Pt:2):2-6.
97. Collins S, Currie L, Patel V, et al. Multitasking by clinicians in the context of CPOE and CIS use. *Stud Health Technol Inform* 2007;129(Pt:2):2-62.
98. Estellat C, Colombet I, Vautier S, et al. Impact of pharmacy validation in a computerized physician order entry context. *Int J Qual Health Care* 2007;19(5):317-25.
99. Jayawardena S, Eisdorfer J, Indulkar S, et al. Prescription errors and the impact of computerized prescription order entry system in a community-based hospital. *Am J Ther* 2007;14(4):336-40.

100. Lapane KL, Dube C, Schneider KL, et al. Patient perceptions regarding electronic prescriptions: is the geriatric patient ready? *J Am Geriatr Soc* 2007;55(8):1254-9.
101. Oswald S, Caldwell R. Dispensing error rate after implementation of an automated pharmacy carousel system. *Am J Health Syst Pharm* 2007;64(13):1427-31.
102. Cosgrove SE, Patel A, Song X, et al. Impact of different methods of feedback to clinicians after postprescription antimicrobial review based on the Centers For Disease Control and Prevention's 12 Steps to Prevent Antimicrobial Resistance Among Hospitalized Adults. *InfectControl Hosp Epidemiol* 2007;28(6):641-6.
103. Vardi A, Efrati O, Levin I, et al. Prevention of potential errors in resuscitation medications orders by means of a computerised physician order entry in paediatric critical care. *Resuscitation* 2007;73(3):400-6.
104. Cochran GL, Jones KJ, Brockman J, et al. Errors prevented by and associated with barcode medication administration systems. *Jt Comm J Qual Patient Saf* 2007;33(5):293-301.
105. Malone DC, Abarca J, Skrepnek GH, et al. Pharmacist workload and pharmacy characteristics associated with the dispensing of potentially clinically important drug-drug interactions. *Med Care* 2007;45(5):456-62.
106. Bermejo VT, Perez Menendez CC, Grupo de trabajo nuevas tecnologias de la SEFH (TECNO), et al. The application of new technologies to hospital pharmacy in Spain. *Farm Hosp* 2007;31(1):17-22.
107. Indermitte J, Beutler M, Bruppacher R, et al. Management of drug-interaction alerts in community pharmacies. *J Clin Pharm Therapeut* 2007;32(2):133-42.
108. Galvin L, McBeth S, Hasdorff C, et al. Medication bar coding: to scan or not to scan? *Comput Inform Nurs* 2007;25(2):86-92.
109. Staggers N, Kobus D, Brown C. Nurses' evaluations of a novel design for an electronic medication administration record. *Comput Inform Nurs* 2007;25(2):67-75.
110. O'Neill L, Klepack W. Electronic medical records for a rural family practice: a case study in systems development. *J Med Syst* 2007;31(1):25-33.
111. Hartzema AG, Winterstein AG, Johns TE, et al. Planning for pharmacy health information technology in critical access hospitals. *Am J Health Syst Pharm* 2007;64(3):315-21.
112. Killelea BK, Kaushal R, Cooper M, et al. To what extent do pediatricians accept computer-based dosing suggestions? *Pediatrics* 2007;119(1):e69-e75
113. Ko Y, Abarca J, Malone DC, et al. Practitioners' views on computerized drug-drug interaction alerts in the VA system. *J Am Med Inform Assoc* 2007;14(1):56-64.
114. Skibinski KA, White BA, Lin LI, et al. Effects of technological interventions on the safety of a medication-use system. *Am J Health Syst Pharm* 2007;64(1):90-6.
115. Parker BM, Henderson JM, Vitagliano S, et al. Six sigma methodology can be used to improve adherence for antibiotic prophylaxis in patients undergoing noncardiac surgery. *Anesth Analg* 2007;104(1):140-6.
116. Ulanimo VM, O'Leary-Kelley C, Connolly PM. Nurses' perceptions of causes of medication errors and barriers to reporting. *J Nurs Care Qual* 2007;22(1):28-33.
117. Barron WM, Reed RL, Forsythe S, et al. Implementing computerized provider order entry with an existing clinical information system. *Jt Comm J Qual Patient Saf* 2006;32(9):506-16.
118. Reichley R, McKinnon PS, Seaton T, et al. Adherence to dyslipidemia treatment guidelines in diabetic patients. *AMIA* 2006;1071
119. Poon EG, Keohane C, Featherstone E, et al. Impact of barcode medication administration technology on how nurses spend their time on clinical care. *AMIA* 2006;1065
120. Collins S, Currie L, Bakken S, et al. Interruptions during the use of a CPOE system for MICU rounds. *AMIA* 2006;895
121. Sintchenko V, Coiera E. Decision complexity affects the extent and type of decision support use. *AMIA* 2006;724-8.

122. Martins SB, Lai S, Tu S, et al. Offline testing of the ATHENA Hypertension decision support system knowledge base to improve the accuracy of recommendations. *AMIA 2006*;539-43.
123. Lobach DF, Willis JM, Macri JM, et al. Perceptions of Medicaid beneficiaries regarding the usefulness of accessing personal health information and services through a patient Internet portal. *AMIA 2006*;509-13.
124. Jensen J. The effects of Computerized Provider Order Entry on medication turn-around time: a time-to-first dose study at the Providence Portland Medical Center. *AMIA 2006*;384-8.
125. Del Fiol G, Rocha RA, Clayton PD. Infobuttons at Intermountain Healthcare: utilization and infrastructure. *AMIA 2006*;180-4.
126. Bouaud J, Seroussi B, Falcoff H, et al. Design factors for success or failure of guideline-based decision support systems: an hypothesis involving case complexity. *AMIA 2006*;71-5.
127. Zamora N, Carter M, Saull-McCaig S, et al. The benefits of the MOE/MAR implementation: a quantitative approach. *HEALTHC Q 2006*;10:77-83.
128. Bradley VM, Steltenkamp CL, Hite KB. Evaluation of reported medication errors before and after implementation of computerized practitioner order entry. *J Healthc Inf Manag 2006*;20(4):46-53.
129. Reifsteck M, Swanson T, Dallas M. Driving out errors through tight integration between software and automation. *J Healthc Inf Manag 2006*;20(4):35-9.
130. Greenberg A, Kramer S, Welch V, et al. Cancer Care Ontario's computerized physician order entry system: a province-wide patient safety innovation. *HEALTHC Q 2006*;9:108-13.
131. Walsh KE, Adams WG, Bauchner H, et al. Medication errors related to computerized order entry for children. *Pediatrics 2006*;118(5):1872-9.
132. Webb AL, Flagg RL, Fink AS. Reducing surgical site infections through a multidisciplinary computerized process for preoperative prophylactic antibiotic administration. *Am J Surg 2006*;192(5):663-8.
133. Nace GS, Graumlich JF, Aldag JC. Software design to facilitate information transfer at hospital discharge. *Inform Prim Care 2006*;14(2):109-19.
134. Judge J, Field TS, DeFlorio M, et al. Prescribers' responses to alerts during medication ordering in the long term care setting. *J Am Med Inform Assoc 2006*;13(4):385-90.
135. Kilbridge PM, Campbell UC, Cozart HB, et al. Automated surveillance for adverse drug events at a community hospital and an academic medical center. *J Am Med Inform Assoc 2006*;13(4):372-7.
136. Menachemi N, Lee SC, Shepherd JE, et al. Proliferation of electronic health records among obstetrician-gynecologists. *Qual Manag Health Care 2006*;15(3):150-6.
137. McInnes DK, Saltman DC, Kidd MR. General practitioners' use of computers for prescribing and electronic health records: results from a national survey. *Med J Aust 2006*;185(2):88-91.
138. Thursky KA, Buising KL, Bak N, et al. Reduction of broad-spectrum antibiotic use with computerized decision support in an intensive care unit. *Int J Qual Health Care 2006*;18(3):224-31.
139. Shaw GM, Chase JG, Wong J, et al. Rethinking glycaemic control in critical illness--from concept to clinical practice change. *Journal of the Australasian Academy of Critical 2006*;8(2):90-9.
140. Hornick TR, Higgins PA, Stollings C, et al. Initial evaluation of a computer-based medication management tool in a geriatric clinic. *American Journal Geriatric Pharmacotherapy 2006*;4(1):62-9.
141. Patterson ES, Rogers ML, Chapman RJ, et al. Compliance with intended use of Bar Code Medication Administration in acute and long-term care: an observational study. *Hum Factors 2006*;48(1):15-22.

142. Skrepnek GH, Armstrong EP, Malone DC, et al. Workload and availability of technology in metropolitan community pharmacies. *J Am Pharm Assoc* 2006;46(2):154-60.
143. Eliasson M, Bastholm P, Forsberg P, et al. Janus computerised prescribing system provides pharmacological knowledge at point of care - design, development and proof of concept. *Eur J Clin Pharmacol* 2006;62(4):251-8.
144. Asaro PV, Sheldahl AL, Char DM. Embedded guideline information without patient specificity in a commercial emergency department computerized order-entry system. *Acad Emerg Med* 2006;13(4):452-8.
145. Tamblyn R, Huang A, Kawasumi Y, et al. The development and evaluation of an integrated electronic prescribing and drug management system for primary care. *J Am Med Inform Assoc* 2006;13(2):148-59.
146. Ward MM, Evans TC, Spies AJ, et al. National Quality Forum 30 safe practices: priority and progress in Iowa hospitals. *Am J Med Qual* 2006;21(2):101-8.
147. Fitzmaurice DA. Oral anticoagulation control: the European perspective. *J Thromb Thrombolysis* 2006;21(1):95-100.
148. Bonnabry P, Cingria L, Ackermann M, et al. Use of a prospective risk analysis method to improve the safety of the cancer chemotherapy process. *Int J Qual Health Care* 2006;18(1):9-16.
149. Poon EG, Jha AK, Christino M, et al. Assessing the level of healthcare information technology adoption in the United States: a snapshot. *BMC Med Inform Decis Mak* 2006;6:1
150. Shah NR, Seger AC, Seger DL, et al. Improving acceptance of computerized prescribing alerts in ambulatory care. *J Am Med Inform Assoc* 2006;13(1):5-11.
151. Shah NR, Seger AC, Seger DL, et al. Improving override rates for computerized prescribing alerts in ambulatory care. *AMIA* 2005;1110
152. Reynolds K, Peres A, Tatham JM. The impact on patient safety of free-text entry of nursing orders into an electronic medical record in an integrated delivery system. *AMIA* 2005;1095
153. Menke JA, Rich D, McClead RE. Physician response to CPOE allergy information alerts. *AMIA* 2005;1051
154. Quinn MM, Mannion J. Improving patient safety using interactive, evidence-based decision support tools. *Jt Comm J Qual Patient Saf* 2005;31(12):678-83.
155. Taylor LK, Kawasumi Y, Bartlett G, et al. Inappropriate prescribing practices: the challenge and opportunity for patient safety. *HEALTHC Q* 2005;8:81-5.
156. Gandhi TK, Bartel SB, Shulman LN, et al. Medication safety in the ambulatory chemotherapy setting. *Cancer* 2005;104(11):2477-83.
157. Stevenson KB, Barbera J, Moore JW, et al. Understanding keys to successful implementation of electronic decision support in rural hospitals: analysis of a pilot study for antimicrobial prescribing. *Am J Med Qual* 2005;20(6):313-8.
158. Lawton G, Shields A. Bar-code verification of medication administration in a small hospital. *Am J Health Syst Pharm* 2005;62(22):2413-5.
159. Hillman JM, Given RS. Hospital implementation of computerized provider order entry systems: results from the 2003 leapfrog group quality and safety survey. *J Healthc Inf Manag* 2005;19(4):55-65.
160. Mirco A, Campos L, Falcao F, et al. Medication errors in an internal medicine department. Evaluation of a computerized prescription system. *Pharm World Sci* 2005;27(4):351-2.
161. Rochon PA, Field TS, Bates DW, et al. Computerized physician order entry with clinical decision support in the long-term care setting: insights from the Baycrest Centre for Geriatric Care. *J Am Geriatr Soc* 2005;53(10):1780-9.
162. Gordon JO, Hadsall RS, Schommer JC. Automated medication-dispensing system in two hospital emergency departments. *Am J Health Syst Pharm* 2005;62(18):1917-23.

163. Beuscart-Zephir MC, Pelayo S, Anceaux F, et al. Impact of CPOE on doctor-nurse cooperation for the medication ordering and administration process. *Int J Med Inf* 2005;74(7-8):629-41.
164. Kushniruk AW, Triola MM, Borycki EM, et al. Technology induced error and usability: the relationship between usability problems and prescription errors when using a handheld application. *Int J Med Inf* 2005;74(7-8):519-26.
165. Reichley RM, Seaton TL, Resetar E, et al. Implementing a commercial rule base as a medication order safety net. *J Am Med Inform Assoc* 2005;12(4):383-9.
166. Morris CJ, Savelyich BS, Avery AJ, et al. Patient safety features of clinical computer systems: questionnaire survey of GP views. *Qual Safe Health Care* 2005;14(3):164-8.
167. Suarez-Varela UJ, Beltran CC, Molina LT, et al. Computer-aided prescribing: from utopia to reality. *Aten Primaria* 2005;35(9):451-6.
168. Nebeker JR, Hoffman JM, Weir CR, et al. High rates of adverse drug events in a highly computerized hospital. *Arch Intern Med* 2005;165(10):1111-6.
169. Ragan R, Bond J, Major K, et al. Improved control of medication use with an integrated bar-code-packaging and distribution system. *Am J Health Syst Pharm* 2005;62(10):1075-9.
170. Awaya T, Ohtaki K, Yamada T, et al. Automation in drug inventory management saves personnel time and budget. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2005;125(5):427-32.
171. Lederman RM, Parkes C. Systems failure in hospitals-using Reason's model to predict problems in a prescribing information system. *J Med Syst* 2005;29(1):33-43.
172. Pizzi LT, Suh DC, Barone J, et al. Factors related to physicians' adoption of electronic prescribing: results from a national survey. *Am J Med Qual* 2005;20(1):22-32.
173. Spina JR, Glassman PA, Belperio P, et al. Clinical relevance of automated drug alerts from the perspective of medical providers. *Am J Med Qual* 2005;20(1):7-14.
174. Burt CW, Hing E. Use of computerized clinical support systems in medical settings: United States, 2001-03. *Adv Data* 2005;(353):1-8.
175. Levick D, Lukens HF, Stillman PL. You've led the horse to water, now how do you get him to drink: managing change and increasing utilization of computerized provider order entry. *J Healthc Inf Manag* 2005;19(1):70-5.
176. Smith WD, Hatton RC, Fann AL, et al. Evaluation of drug interaction software to identify alerts for transplant medications. *Ann Pharmacother* 2005;39(1):45-50.
177. Burke JP, Pestotnik SL. Antibiotic use and microbial resistance in intensive care units: impact of computer-assisted decision support. *J Chemother* 1999;11(6):530-5.
178. Perlini S, Piepoli M, Marti G, et al. Treatment of chronic heart failure: an expert system advisor for general practitioners. *Acta Cardiol* 1990;45(5):365-78.
179. Stewart K, Loftus S, DeLisle S. Prescription of amiodarone through a computerized template that includes both decision support and executive functions improves the monitoring for toxicities. *AMIA* 2003;1020
180. Rich D, Menke J, Fisher D. Dose range checking in a computer order entry system. *AMIA* 2003;985
181. Bartlett G, Tamblyn R, Huang A, et al. Evaluation of standardized tasks for primary care physicians using the MOXXI electronic prescribing and integrated drug management system. *AMIA* 2003;786
182. Bailey TC, Noirot LA, Christensen EM, et al. Validation of automated event triggers using laboratory values related to two problem-prone drugs. *AMIA* 2003;781
183. Horsky J, Kaufman DR, Patel VL. The cognitive complexity of a provider order entry interface. *AMIA* 2003;294-8.
184. Fung KW, Vogel LH. Will decision support in medications order entry save money? A return on investment analysis of the case of the Hong Kong hospital authority. *AMIA* 2003;244-54.
185. Cheng CH, Goldstein MK, Geller E, et al. The Effects of CPOE on ICU workflow: an observational study. *AMIA* 2003;150-4.

186. Perel A, Berkenstadt H, Ziv A, et al. Anaesthesiologists' views on the need for point-of-care information system in the operating room: a survey of the European Society of Anaesthesiologists. *Eur J Anaesthesiol* 2004;21(11):898-901.
187. Twiggs JE, Fifield J, Jackson E, et al. Treating asthma by the guidelines: developing a medication management information system for use in primary care. *Disease Management* 2004;7(3):244-60.
188. Anderson S, Wittwer W. Using bar-code point-of-care technology for patient safety. *J Healthc Qual* 2004;26(6):5-11.
189. Fischer MA, Lilly CM, Churchill WW, et al. An algorithmic computerised order entry approach to assist in the prescribing of new therapeutic agents: case study of activated protein C at an academic medical centre. *Drug Saf* 2004;27(15):1253-61.
190. Yousef ZR, Tandy SC, Tudor V, et al. Warfarin for non-rheumatic atrial fibrillation: five year experience in a district general hospital. *Heart* 2004;90(11):1259-62.
191. Silverman JB, Stapinski CD, Huber C, et al. Computer-based system for preventing adverse drug events. *Am J Health Syst Pharm* 2004;61(15):1599-603.
192. Taylor LK, Tamblyn R. Reasons for physician non-adherence to electronic drug alerts. *Stud Health Technol Inform* 2004;107(Pt:2):2-5.
193. Kushniruk A, Triola M, Stein B, et al. The relationship of usability to medical error: an evaluation of errors associated with usability problems in the use of a handheld application for prescribing medications. *Stud Health Technol Inform* 2004;107(Pt:2):2-6.
194. Horsky J, Kaufman DR, Patel VL. Computer-based drug ordering: evaluation of interaction with a decision-support system. *Stud Health Technol Inform* 2004;107(Pt:2):2-7.
195. Beuscart-Zephir MC, Pelayo S, Degoulet P, et al. A usability study of CPOE's medication administration functions: impact on physician-nurse cooperation. *Stud Health Technol Inform* 2004;107(Pt:2):2-22.
196. Chan AS, Coleman RW, Martins SB, et al. Evaluating provider adherence in a trial of a guideline-based decision support system for hypertension. *Stud Health Technol Inform* 2004;107(Pt:1):1-9.
197. Trivedi MH, Kern JK, Grannemann BD, et al. A computerized clinical decision support system as a means of implementing depression guidelines. *Psychiatr Serv* 2004;55(8):879-85.
198. Apkon M, Leonard J, Probst L, et al. Design of a safer approach to intravenous drug infusions: failure mode effects analysis. *Qual Safe Health Care* 2004;13(4):265-71.
199. Collins CD, Pedersen CA, Schneider PJ, et al. Effect on amphotericin B lipid complex use of a clinical decision support system for computerized prescriber order entry. *Am J Health Syst Pharm* 2004;61(13):1395-9.
200. Roumie CL, Grogan EL, Falbe W, et al. A three-part intervention to change the use of hormone replacement therapy in response to new evidence. *Ann Intern Med* 2004;141(2):118-25.
201. Fair MA, Pane F. Pharmacist interventions in electronic drug orders entered by prescribers. *Am J Health Syst Pharm* 2004;61(12):1286-8.
202. Morera T, Gervasini G, Carrillo JA, et al. Using a computerized drug prescription screening system to trace drug interactions in an outpatient setting. *Ann Pharmacother* 2004;38(7-8):1301-6.
203. Anton C, Nightingale PG, Adu D, et al. Improving prescribing using a rule based prescribing system. *Qual Safe Health Care* 2004;13(3):186-90.
204. Rosenbloom ST, Talbert D, Aronsky D. Clinicians' perceptions of clinical decision support integrated into computerized provider order entry. *Int J Med Inf* 2004;73(5):433-41.
205. de Lusignan S, Singleton A, Wells S. Lessons from the implementation of a near patient anticoagulant monitoring service in primary care. *Inform Prim Care* 2004;12(1):27-33.

206. Grayson ML, Melvani S, Kirsa SW, et al. Impact of an electronic antibiotic advice and approval system on antibiotic prescribing in an Australian teaching hospital. *Med J Aust* 2004;180(9):455-8.
207. Krampera M, Venturini F, Benedetti F, et al. Computer-based drug management in a bone marrow transplant unit: a suitable tool for multiple prescriptions even in critical conditions. *Br J Haematol* 2004;125(1):50-7.
208. Potts AL, Barr FE, Gregory DF, et al. Computerized physician order entry and medication errors in a pediatric critical care unit. *Pediatrics* 2004;113(1:Pt:1):59-63.
209. Manotti C, Pattacini C, Quintavalla R, et al. Computer assisted anticoagulant therapy. *Pathophysiol Haemost Thromb* 2003;33(5-6):366-72.
210. Short D, Frischer M, Bashford J. The development and evaluation of a computerised decision support system for primary care based upon 'patient profile decision analysis'. *Inform Prim Care* 2003;11(4):195-202.
211. Rogers JE, Wroe CJ, Roberts A, et al. Automated quality checks on repeat prescribing. *Br J Gen Pract* 2003;53(496):838-44.
212. Weingart SN, Toth M, Sands DZ, et al. Physicians' decisions to override computerized drug alerts in primary care. *Arch Intern Med* 2003;163(21):2625-31.
213. Horsky J, Kaufman DR, Oppenheim MI, et al. A framework for analyzing the cognitive complexity of computer-assisted clinical ordering. *J Biomed Inform* 2003;36(1-2):4-22.
214. King WJ, Paice N, Rangrej J, et al. The effect of computerized physician order entry on medication errors and adverse drug events in pediatric inpatients. *Pediatrics* 2003;112(3 Pt 1):t-9
215. Wells BJ, Lobel KD, Dickerson LM. Using the electronic medical record to enhance the use of combination drugs. *Am J Med Qual* 2003;18(4):147-9.
216. Larrabee S, Brown MM. Recognizing the institutional benefits of bar-code point-of-care technology. *Jt Comm J Qual Patient Saf* 2003;29(7):345-53.
217. France DJ, Miles P, Cartwright J, et al. A chemotherapy incident reporting and improvement system. *Jt Comm J Qual Patient Saf* 2003;29(4):171-80.
218. Klibanov OM, Eckel SF. Effects of automated dispensing on inventory control, billing, workload, and potential for medication errors. *Am J Health Syst Pharm* 2003;60(6):569-72.
219. Dobscha SK, Anderson TA, Hoffman WF, et al. Strategies to decrease costs of prescribing selective serotonin reuptake inhibitors at a VA Medical Center. *Psychiatr Serv* 2003;54(2):195-200.
220. Payne TH, Nichol WP, Hoey P, et al. Characteristics and override rates of order checks in a practitioner order entry system. *Proceedings / AMIA* 2002;602-6.
221. Oppenheim MI, Vidal C, Velasco FT, et al. Impact of a computerized alert during physician order entry on medication dosing in patients with renal impairment. *Proceedings / AMIA* 2002;577-81.
222. Noirot LA, Blickensderfer A, Christensen E, et al. Implementation of an automated guideline monitor for secondary prevention of acute myocardial infarction. *Proceedings / AMIA* 2002;562-6.
223. Kinn JW, Marek JC, O'Toole MF, et al. Effectiveness of the electronic medical record in improving the management of hypertension. *J Clin Hypertens* 2002;4(6):415-9.
224. Bizovi KE, Beckley BE, McDade MC, et al. The effect of computer-assisted prescription writing on emergency department prescription errors. *Acad Emerg Med* 2002;9(11):1168-75.
225. Magnus D, Rodgers S, Avery AJ. GPs' views on computerized drug interaction alerts: questionnaire survey. *J Clin Pharm Therapeut* 2002;27(5):377-82.
226. Kuilboer MM, van Wijk MA, Mosseveld M, et al. Feasibility of AsthmaCritic, a decision-support system for asthma and COPD which generates patient-specific feedback on routinely recorded data in general practice. *Fam Pract* 2002;19(5):442-7.

227. Schrezenmeir J, Dirting K, Papazov P. Controlled multicenter study on the effect of computer assistance in intensive insulin therapy of type 1 diabetics. *Comput Methods Programs Biomed* 2002;69(2):97-114.
228. Lehman ML, Brill JH, Skarulis PC, et al. Physician Order Entry impact on drug turn-around times. *Proceedings / AMIA* 2001;359-63.
229. Ovhed I, Berglund J, Oistamo S, et al. A pop-up menu linked to a computerized drug prescribing system. Prescribing pattern's feedback via a simple and quick method. *Lakartidningen* 2001;98(50):5772-6.
230. Murff HJ, Kannry J. Physician satisfaction with two order entry systems. *J Am Med Inform Assoc* 2001;8(5):499-509.
231. Lee YL, Hsu CY, Hsieh D, et al. Development and deployment of a web-based physician order entry system. *Int J Med Inf* 2001;62(2-3):135-42.
232. Payne TH, Savarino J, Marshall R, et al. Use of a clinical event monitor to prevent and detect medication errors. *Proceedings / AMIA* 2000;640-4.
233. Brown S, Black K, Mrochek S, et al. RADARx: Recognizing, Assessing, and Documenting Adverse Rx events. *Proceedings / AMIA* 2000;101-5.
234. Abookire SA, Teich JM, Sandige H, et al. Improving allergy alerting in a computerized physician order entry system. *Proceedings / AMIA* 2000;2-6.
235. Franke L, Avery AJ, Groom L, et al. Is there a role for computerized decision support for drug dosing in general practice? A questionnaire survey. *J Clin Pharm Therapeut* 2000;25(5):373-7.
236. Glemaud I. Use of a physician order entry system to identify opportunities for intravenous to oral levofloxacin conversion. *Am J Health Syst Pharm* 2000;57:14-6.
237. Kailajarvi M, Takala T, Gronroos P, et al. Reminders of drug effects on laboratory test results. *Clin Chem* 2000;46(9):1395-400.
238. Luchins DJ, Klass D, Hanrahan P, et al. Computerized monitoring of valproate and physician responsiveness to laboratory studies as a quality indicator. *Psychiatr Serv* 2000;51(9):1179-81.
239. Gonzalez-Heydrich J, DeMaso DR, Irwin C, et al. Implementation of an electronic medical record system in a pediatric psychopharmacology program. *Int J Med Inf* 2000;57(2-3):109-16.
240. Liu Z, Sakurai T, Orii T, et al. Evaluations of the prescription order entry system for outpatient clinics by physicians in the 80 university hospitals in Japan. *Med Inform Internet Med* 2000;25(2):123-32.
241. Nightingale PG, Adu D, Richards NT, et al. Implementation of rules based computerised bedside prescribing and administration: intervention study.[see comment]. *BMJ* 2000;320(7237):750-3.
242. Levy M, Azaz-Livshits T, Sadan B, et al. Computerized surveillance of adverse drug reactions in hospital: implementation. *Eur J Clin Pharmacol* 1999;54(11):887-92.
243. Bolton PG, Usher HE, Mira M, et al. Information technology and general practice. A survey of general practitioner attitudes towards computerisation. *Aust Fam Physician* 1999;28 Suppl 1:S19-S21
244. Warner H, Jr., Reimer L, Suvinier D, et al. Modeling empiric antibiotic therapy evaluation of QID. *Proceedings / AMIA* 1999;Annual Symposium.:440-4.
245. Miller JE, Reichley RM, McNamee LA, et al. Notification of real-time clinical alerts generated by pharmacy expert systems. *Proceedings / AMIA* 1999;Annual Symposium.:325-9.
246. Grundmeier R, Johnson K. Housestaff attitudes toward computer-based clinical decision support. *Proceedings / AMIA* 1999;Annual Symposium.:266-70.
247. Chin HL, Wallace P. Embedding guidelines into direct physician order entry: simple methods, powerful results. *Proceedings / AMIA* 1999;Annual Symposium.:221-5.
248. Ash JS, Gorman PN, Hersh WR. Physician order entry in U.S. hospitals. *Proceedings / AMIA* 1998;Annual Symposium.:235-9.

249. Strasberg HR, Tudiver F, Holbrook AM, et al. Moving towards an electronic patient record: a survey to assess the needs of community family physicians. *Proceedings / AMIA 1998;Annual Symposium*.:230-4.
250. Raschke RA, Gollihare B, Wunderlich TA, et al. A computer alert system to prevent injury from adverse drug events: development and evaluation in a community teaching hospital.[see comment][erratum appears in JAMA 1999 Feb 3;281(5):420]. *JAMA* 1998;280(15):1317-20.
251. Monane M, Matthias DM, Nagle BA, et al. Improving prescribing patterns for the elderly through an online drug utilization review intervention: a system linking the physician, pharmacist, and computer.[see comment]. *JAMA* 1998;280(14):1249-52.
252. Urschitz M, Lorenz S, Unterasinger L, et al. Three years experience with a patient data management system at a neonatal intensive care unit. *J Clin Monit Comput* 1998;14(2):119-25.
253. Fitzmaurice DA, Hobbs FD, Murray ET. Primary care anticoagulant clinic management using computerized decision support and near patient International Normalized Ratio (INR) testing: routine data from a practice nurse-led clinic. *Fam Pract* 1998;15(2):144-6.
254. Eisenstein EL, Peterson ED, Jollis JG, et al. Assessing the value of newer pharmacologic agents in non-ST elevation patients: a decision support system application. *Proceedings/AMIA Annual Fall Symposium* 1997;273-7.
255. Anderson JG, Jay SJ, Anderson M, et al. Evaluating the potential effectiveness of using computerized information systems to prevent adverse drug events. *Proceedings/AMIA Annual Fall Symposium* 1997;228-32.
256. Gronroos PE, Irjala KM, Huupponen RK, et al. A medication database--a tool for detecting drug interactions in hospital. *Eur J Clin Pharmacol* 1997;53(1):13-7.
257. Rivkin S. Opportunities and challenges of electronic physician prescribing technology. *Med Interface* 1983;10(8):77-8.
258. Vedsted P, Nielsen JN, Olesen F. Does a computerized price comparison module reduce prescribing costs in general practice? *Fam Pract* 1997;14(3):199-203.
259. Chin HL, Krall M. Implementation of a comprehensive computer-based patient record system in Kaiser Permanente's Northwest Region. *MD Comput* 1997;14(1):41-5.
260. Weir C, Johnsen V, Roscoe D, et al. The impact of physician order entry on nursing roles. *Proceedings/AMIA Annual Fall Symposium* 1996;714-8.
261. Zarowitz BJ, Petitta A, Rudis MI, et al. Bar code documentation of pharmacotherapy services in intensive care units. *Pharmacotherapy* 1996;16(2):261-6.
262. Pestotnik SL, Classen DC, Evans RS, et al. Implementing antibiotic practice guidelines through computer-assisted decision support: clinical and financial outcomes. *Ann Intern Med* 1996;124(10):884-90.
263. Arbogast JG, Carr MC, Dodrill WH. Overcoming the limitations of proprietary computerized billing systems to enhance patient care. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1995;722-6.
264. Krall MA. Acceptance and performance by clinicians using an ambulatory electronic medical record in an HMO. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1995;708-11.
265. Jozefiak ET, Lewicki JE, Kozinn WP. Computer-assisted antimicrobial surveillance in a community teaching hospital. *Am J Health Syst Pharm* 1995;52(14):1536-40.
266. Borel JM, Rascati KL. Effect of an automated, nursing unit-based drug-dispensing device on medication errors.[see comment]. *Am J Health Syst Pharm* 1995;52(17):1875-9.
267. Puckett F. Medication-management component of a point-of-care information system. *Am J Health Syst Pharm* 1995;52(12):1305-9.
268. Schwarz HO, Brodowy BA. Implementation and evaluation of an automated dispensing system. *Am J Health Syst Pharm* 1995;52(8):823-8.

269. Schumock GT, Marwaha TR, McBride JM, et al. Automated order-entry mechanisms to influence prescribing. *Top Hosp Pharm Manage* 1994;14(3):21-9.
270. Mason RN, Pugh CB, Boyer SB, et al. Computerized documentation of pharmacists' interventions. *Am J Hosp Pharm* 1994;51(17):2131-8.
271. Sands DZ, Safran C. Closing the loop of patient care--a clinical trial of a computerized discharge medication program. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1994;841-5.
272. Morrell R, Wasiluskas B, Winslow R. Personal computer-based expert system for quality assurance of antimicrobial therapy. *Am J Hosp Pharm* 1993;50(10):2067-73.
273. Hunt B. Development of a MUMPS-based anticoagulant management system. *Br J Biomed Sci* 1993;50(2):117-24.
274. Love DJ, Schalk DC, Morgan MC. Computerized relational database for monitoring clozapine therapy. *Am J Hosp Pharm* 1993;50(8):1657-62.
275. Zarowitz BJ, Petitta A, Mlynarek M, et al. Bar-code technology applied to drug-use evaluation. *Am J Hosp Pharm* 1993;50(5):935-9.
276. Pestotnik SL, Classen DC, Evans RS, et al. Prospective surveillance of imipenem/cilastatin use and associated seizures using a hospital information system. *Ann Pharmacother* 1993;27(4):497-501.
277. Evans RS, Pestotnik SL, Classen DC, et al. Development of an automated antibiotic consultant. *MD Comput* 1993;10(1):17-22.
278. Gaucher M, Greer M. A nursing evaluation of unit dose and computerized medication administration records. *Can J Hosp Pharm* 1992;45(4):145-50.
279. Yates P, Stear M. A mainframe interfacing computer management system for the control of oral anticoagulant therapy. *Clinical & Laboratory Haematology* 1992;14(3):245-50.
280. Goldberg DE, Baardsgaard G, Johnson MT, et al. Computer-based program for identifying medication orders requiring dosage modification based on renal function. *Am J Hosp Pharm* 1991;48(9):1965-9.
281. Ishizuka H, Horiguchi M, Waki Y, et al. Computerized dispensing system: reducing the time of dispensing medicines. *Int J Biomed Comput* 1991;28(1-2):137-46.
282. Craghead RM, Wartski DM. An evaluative study of unclaimed prescriptions. *Hosp Pharm* 1991;26(7):616-7.
283. DeNeef P, Ellsworth A, Schneeweiss R. A system for drug utilization review in ambulatory care. *J Fam Pract* 1991;32(6):607-12.
284. Huntress JD, Possidente CJ, Harry DJ. Documenting pharmacists' interventions on a hospital's mainframe computer system. *Am J Hosp Pharm* 1990;47(12):2711-5.
285. Tsubaki T, Orii T, Sugiura M. Utilization of a computerized system at the pharmacy department of the University of Tokyo Hospital--impact of prescription order entry and computerized dispensing system. *Jpn Hosp* 1990;9:61-7.
286. Mahoney CD. Restructuring pharmacy services to reduce expenses without eliminating services. *Am J Hosp Pharm* 1990;47(3):579-84.
287. Pestotnik SL, Evans RS, Burke JP, et al. Therapeutic antibiotic monitoring: surveillance using a computerized expert system. *Am J Med* 1990;88(1):43-8.
288. Ryan PJ, Gilbert M, Rose PE. Computer control of anticoagulant dose for therapeutic management.[see comment]. *BMJ* 1989;299(6709):1207-9.
289. Kawahara NE, Jordan FM. Influencing prescribing behavior by adapting computerized order-entry pathways.[see comment]. *Am J Hosp Pharm* 1989;46(9):1798-801.
290. Barry GA, Bass GE, Jr., Eddlemon JK, et al. Bar-code technology for documenting administration of large-volume intravenous solutions. *Am J Hosp Pharm* 1989;46(2):282-7.

291. Ogura H, Sagara E, Iwata M, et al. Online support functions of prescription order system and prescription audit in an integrated hospital information system. *Med Inform (Lond)* 1988;13(3):161-9.
292. Hokanson JA, Guernsey BG, Bryant SG, et al. The feasibility of barcode-based dispensing quality assurance programs. *Drug Intell Clin Pharm* 1984;18(1):76-8.
293. Lomonte PJ, Besser RA, Thomas EC. Effect of decentralized computer order entry on medication turnaround time. *Am J Hosp Pharm* 1983;40(6):979-81.
294. Physician order entry reduces inappropriate vancomycin prescribing, VRE rates. *Formulary* 2001;36(7):
295. Bahl V. Developing dashboards to measure and manage inpatient pharmacy costs. *Am J Health Syst Pharm* 2007;64(17):1859-66.
296. Benkhaial A. Prescribing errors in patients with documented drug allergies: Comparison of ICD-10 coding and written patient notes. *Pharm World Sci* 2009;31(4):464-72.
297. Bomba D. Moving beyond implementation to sustained use of computers in general practice in Australia. *International Journal of Healthcare Technology and Management* 2004;6(1):83-90.
298. Brown MD. Implementation of an Emergency Department-based Transient Ischemic Attack Clinical Pathway: A Pilot Study in Knowledge Translation. *Acad Emerg Med* 2007;14(11):1114-9.
299. Chahal JK. The impact of drug-history clerking using computerised notes in the ICU on subsequent prescribing. *Pharmacy in Practice* 2007;17(1):14-8.
300. Chen C-I. Medical errors in a hospital in Taiwan: Incidence, aetiology and proposed solutions. *Journal on Information Technology in Healthcare* 2004;2(1):11-8.
301. Chu S. Electronic prescription: Standards and decision support issues. *Journal on Information Technology in Healthcare* 2004;2(6):
302. DuBeshter B, Griggs J, Angel C, et al. Chemotherapy dose limits set by users of a computer order entry system. *Hosp Pharm* 2006;41(2):136-42.
303. Ekedahl A, Wessling A, Melander A. Primary non-compliance with automated prescription transmittals from health care centres in Sweden. *J Soc Adm Pharm* 2002;19(4):137-40.
304. Garr DR, Jenkins R, Ornstein S, et al. The effect of routine use of computer-generated preventive reminders in a clinical practice. *Am J Prev Med* 1993;9(1):55-61.
305. Gimenes FRE, Grou CR, Miasso A, et al. Electronic prescription as contributing factor for hospitalized patients' safety. *Pharmacy Practice* 2006;4(1):13-7.
306. Grangeasse L, Fagnoni-Legat C, Chaigneau L, et al. Computerized prescribing of standardized chemotherapy schedules: Residual medication errors and pharmaceutical interventions. *Journal de Pharmacie Clinique* 2006;25(1):33-8.
307. Han Y, Carcillo J, Venkataraman S, et al. Erratum: Unexpected increased mortality after implementation of a commercially sold computerized physician order entry system. *Pediatrics* 2007;117(2):1506-12.
308. Hazlehurst B, Naleway A, Mullooly J. Detecting possible vaccine adverse events in clinical notes of the electronic medical record. *Vaccine* 2009;27(14):2077-83.
309. Macdonald D, Neville D, Priddle M. Community pharmacists' expectations of a pharmacy network: A baseline evaluation. *Canadian Pharmacists Journal* 2005;138(5):50-8.
310. Manzo J, Sinnett M, Sosnowski F, et al. Case study: Challenges, successes and lessons learned from implementing Computerized Physician Order Entry (CPOE) at two distinct health systems: Implications of CPOE on the pharmacy and the medication-use process. *Hosp Pharm* 2005;40(5):420-9.
311. Mehta R, Onatade R. Experience of electronic prescribing in UK hospitals: A perspective from pharmacy staff. *Pharmaceutical Journal* 2008;281(7511):79-82.
312. Menke JA, Broner C, Campbell D, et al. Computerized clinical documentation system in the pediatric intensive care unit. *BMC Med Inform Decis Mak* 2001;1:1-7.

313. Mungall DR, Anbe D, Forrester P, et al. A prospective randomized comparison of the accuracy of computer-assisted versus GUSTO nomogram-directed heparin therapy. *J Clin Pharm Therapeut* 1993;55(5):591-6.
314. Nicol N. Case study: An interdisciplinary approach to medication error reduction. *Am J Health Syst Pharm* 2007;64(14):S17-S20
315. Ortega A, Aguinagalde A, Lacasa C, et al. Efficacy of an adverse drug reaction electronic reporting system integrated into a hospital information system. *Ann Pharmacother* 2008;42(10):1491-6.
316. Poller L, Keown M, Ibrahim S, et al. A multicentre randomised assessment of the DAWN AC computer-assisted oral anticoagulant dosage program. *Thromb Haemost* 2009;101(3):487-94.
317. Pummer TL, Shalaby K, Erush S. Ordering off the menu: Assessing compliance with a nonformulary medication policy. *Ann Pharmacother* 2009;43(7-8):1251-7.
318. Reidel K, Tamblyn R, Patel V, et al. Pilot study of an interactive voice response system to improve medication refill compliance. *BMC Med Inform Decis Mak* 2008;8:46
319. Ruskin PE, van der Wende J, Clark C, et al. Feasibility of Using the Med-eMonitor System in the Treatment of Schizophrenia: A Pilot Study. *Drug Inf J* 2003;37(3):283-91.
320. Urquhart GA, Williams W, Tobias J, et al. Immunization information systems use during a public health emergency in the United States. *J Public Health Manag Pract* 2007;13(5):481-5.
321. Vaillancourt R, Goulet G, Pean Y, et al. Electronic discharge prescription: A retrospective survey. *Can J Hosp Pharm* 2007;60(2):91-9.
322. van der SH, Kowlesar R, Kloostwijk A, et al. Clinically relevant QTc prolongation due to overridden drug-drug interaction alerts: A retrospective cohort study. *Br J Clin Pharmacol* 2009;67(3):347-54.
323. Beard R, Candlish C. Is electronic prescribing the best system for preventing pharmacy dispensing errors? *BJHC & IM* 2007;24(1):15-8.
324. Brunt BA, Gifford IL. Patient safety, quality care, and service utilization with PLATO (Physician Leadership for Accurate and Timely Orders): A pilot study. *J Nurses Staff Dev* 2009;25(4):E11-E18
325. Burns P, Perkins DA, Larsen K, et al. The introduction of electronic medication charts and prescribing in aged care facilities: An evaluation. *Australas J Ageing* 2007;26(3):131-4.
326. Cassiani SHB, Freire CC, Gimenes FRE. Computerized physician order entry in a university hospital: Writing failure and users' opinions. *Rev Esc Enferm USP* 2003;37(4):51-60.
327. Dombkowski KJ, Leung SW, Clark SJ. Provider attitudes regarding use of an immunization information system to identify children with asthma for influenza vaccination. *J Public Health Manag Pract* 2007;13(6):567-71.
328. France DJ, Cartwright J, Jones V, et al. Improving pediatric chemotherapy safety through voluntary incident reporting: Lessons from the field. *J Pediatr Oncol Nurs* 2004;21(4):200-6.
329. Gandhi TK, Weingart SN, Seger AC, et al. Impact of basic computerized prescribing on outpatient medication errors and adverse drug events... reprinted from the proceedings of the 2001 AMIA Annual Symposium, with permission. *J Am Med Inform Assoc* 2002;9(6):s48-s49
330. Jha AK, Laguette J, Seger A, et al. Can surveillance systems identify and avert adverse drug events? A prospective evaluation of a commercial application. *J Am Med Inform Assoc* 2008;15(5):647-53.
331. O'Brien MS. Fellow project. Implementation of the EPIC electronic medical record/physician order-entry system. *J Healthc Manag* 2006;51(5):338-43.
332. Protti D. The benefits and impacts of the MOE/MAR implementation: A quantitative approach. *HEALTHC Q* 2009;10(83):77-83.
333. Wilson JW, Oyen LJ, Ou NN, et al. Hospital rules-based system: The next generation of medical informatics for patient safety. *Am J Health Syst Pharm* 2004;62(5):499-505.

334. Craig T, Mehta R. Clinician-computer interaction: Automated review of psychotropic drugs. *Am J Psychiatry* 1983;141(2):658-9.
335. Marini S, Hasman A, Huijjer H. Information technology for medication administration: Assessing bedside readiness among nurses in Lebanon. *Int J Evid Based Healthc* 2009;7(1):49-58.
336. Valdovinos M, Schroeder S. The effects of a computerized monitoring system of psychotropic medication use by people with developmental disabilities on staff reported side effects. *J Dev Phys Disabil* 2003;15(4):299-334.
337. Elson B. Electronic prescribing in ambulatory care: Market primer and implications for managed care pharmacy. *J Manag Care Pharm* 2001;7(Mar-Apr):115-20.
338. Fitzpatrick R, Cooke P, Southall C, et al. Evaluation of an automated dispensing system in a hospital, pharmacy dispensary. *Pharmaceutical Journal* 2005;274:763-5.
339. Laucka PV, Hoffman NB. Decreasing medication use in a nursing home patient-care unit. *Am J Hosp Pharm* 1992;49(Jan):96-9.
340. Morrell J, Podlone M, Cohen S. Receptivity of physicians in a teaching hospital to a computerized drug interaction monitoring and reporting system. *Med Care* 1977;15(Jan):68-78.
341. Papshev D, Peterson AM. Extent of electronic prescribing implementation as perceived by MCO pharmacy managers. *J Manag Care Pharm* 2002;8(1):41-7.
342. Perkins L, Hussein G, Leung B. Anemia and erythropoietin (rHuEPO) management program for optimal therapeutics and patient care. *Clinical Research and Regulatory Affairs* 2003;20(3):331-9.
343. Simborgn DW, Derewicz HJ. Highly automated hospital medication system: Five years' experience and evaluation. *Ann Intern Med* 1975;83(Sep):341-6.
344. Young AS, Mintz J, Cohen AN, et al. A network-based system to improve care for schizophrenia: The medical informatics network tool (MINT). *J Am Med Inform Assoc* 2004;11(5):358-67.
345. Aydin C, Ischar R. Effects of computerized order entry on communication between pharmacy and nursing. In Washington, DC, USA: Publ by IEEE; 1989. p.796-801.
346. Lutes K, Chang K, Baggili I. Diabetic e-management system (DEMS). In Las Vegas, NV, United states: Institute of Electrical and Electronics Engineers Computer Society; 2006. p.619-24.
347. Blish C, Proctor R, Fletcher S W and others. Physician acceptance of a computerized outpatient medication system in a teaching hospital group practice. In Silver Spring, MD, USA: IEEE Comput. Soc. Press; 1984. p.203-6.
348. Costa AL, de Oliveira MMB, Oliveria Machado R. An information system for drug prescription and distribution in a public hospital. *Int J Med Inf* 2004;73(4):371-81.
349. Gardner R M, Hulse R K, Larsen K G. Assessing the effectiveness of a computerized pharmacy system. In Los Alamitos, CA, USA: IEEE Comput. Soc. Press; 1990. p.668-72.
350. Goethe JW, Bronzino JD. An expert system for monitoring psychiatric treatment. *IEEE Eng Med Biol Mag* 1995;14(6):776-80.
351. Hoffer EP, Marble KD, Yurchak PM, et al. A computer-based information system for managing patients on long-term oral anticoagulants. *Comput Biomed Res* 1975;8(6):573-9.
352. Ogura H, Yamamoto K, Furutani H, et al. Online prescription order and prescription support in an integrated hospital information system. *Med Inform (Lond)* 1985;10(4):287-99.
353. Prokosch HU, Wong TW, Pryor TA. Medication ordering based on a predictive knowledge base. *Artif Intell Med* 1989;1(1):41-8.
354. Agrawal A, Wub W, Khachewatsky I. Evaluation of an electronic medication reconciliation system in inpatient setting in an acute care hospital. *Stud Health Technol Inform* 2007;129(2):1027-31.
355. Rizos A, Snyder R, Rothschild J and others. Multi-method approach for medication safety event detection in community hospitals. In Washington, DC): 2005. p. 1096.

356. Wang S, Kuperman G, Turetsky M, et al. User-definable medication favorites for an outpatient electronic medical record system. *J Biomed Inform* 2001;35:1055
357. Adams M, bates D, Coffman G et al. Saving lives, saving money: The imperative for CPOE in Massachusetts hospitals. Massachusetts Technology Collaborative and New England Healthcare Institute; 2008. <http://web3.streamhoster.com/mtc/cpoe20808.pdf>
358. Manzo J, Cusick D, Taylor R. Session 47: Measuring medication-related ROI and process improvement after implementing POE. Healthcare Information and Management Systems Society; 2001. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=877>
359. Miller M, Clark J, Lehmann C. Computer based medication error reporting: Insights and implications. *Qual Safe Health Care* 2006;15(3):208-13.
360. Computerized physician order entry: Costs, benefits and challenges. First Consulting Group; 2003. http://www.leapfroggroup.org/media/file/Leapfrog-CPOE_Costs_Benefits_Challenges.pdf
361. Apkon M, Mattera J, Lin Z, et al. A randomized outpatient trial of a decision-support information technology tool. *Arch Intern Med* 2005;165:2388-94.
362. Boling B, McKibben M, Hingl J, et al. Effectiveness of computerized provider order entry with dose range checking on prescribing errors. *J Patient Saf* 2005;1:190-4.
363. Lobach D, Hammond W. Computerized decision support based on a clinical practice guideline improves compliance with care standards. *Am J Med* 1997;102:89-98.
364. Yamauchi K, Ikeda M, Suzuki Y, et al. Evaluation of the order entry system by end users: A step to the new hospital information system. *Nagoya J Med Sci* 1994;57:19-24.
365. Evans K, Benham S, Garrard C. A comparison of handwritten and computer-assisted prescriptions in an intensive care unit. *Critical Care* 1998;2(2):73-8.
366. Dib J, Abdulmohsin S, Farooki M, et al. Effects of an automated drug dispensing system on medication adverse event occurrences and cost containment at SAMSO. *Hosp Pharm* 2006;41(12):1180-4.
367. Marini SD, Hasman A, Huijjer HA, et al. Nurses' attitudes toward the use of the bar-coding medication administration system. *Comput Inform Nurs* 2010;28(2):112-23.
368. Feifer RA, James JM. Geographic variation in drug safety: Potentially unsafe prescribing of medications and prescriber responsiveness to safety alerts. *J Manag Care Pharm* 2010;16(3):196-205.
369. Ali J, Barrow L, Vuylsteke A. The impact of computerised physician order entry on prescribing practices in a cardiothoracic intensive care unit. *Anaesthesia* 2010;65(2):119-23.
370. Champion TR, Jr., Waitman LR, May AK, et al. Social, organizational, and contextual characteristics of clinical decision support systems for intensive insulin therapy: A literature review and case study. *Int J Med Inf* 2010;79(1):31-43.
371. Zwart-van Rijkom JE, Uijtendaal EV, ten Berg MJ, et al. Frequency and nature of drug-drug interactions in a Dutch university hospital. *Br J Clin Pharmacol* 2009;68(2):187-93.
372. Lam AY, Rose D. Telepharmacy services in an urban community health clinic system. *J Am Pharm Assoc* 2009;49(5):652-9.
373. McCoy AB, Peterson JF, Gadd CS, et al. A system to improve medication safety in the setting of acute kidney injury: initial provider response. *AMIA* 2008;Annual:1051
374. Kerrison F, Fraser HS. The impact of uncertain diagnostic results on responses to a decision support system for TB drug prescribing. *AMIA* 2008;Annual:949
375. Wadhwa R, Fridsma DB, Saul MI, et al. Analysis of a failed clinical decision support system for management of congestive heart failure. *AMIA* 2008;Annual:773-7.
376. Kilbridge PM, Noirot LA, Reichley RM, et al. Computerized surveillance for adverse drug events in a pediatric hospital. *AMIA* 2008;16:607-12.

377. Handler SM, Hanlon JT, Perera S, et al. Assessing the performance characteristics of signals used by a clinical event monitor to detect adverse drug reactions in the nursing home. *AMIA 2008;Annual*:278-82.
378. Ramirez E, Carcas AJ, Borobia AM, et al. A pharmacovigilance program from laboratory signals for the detection and reporting of serious adverse drug reactions in hospitalized patients. *J Clin Pharm Therapeut* 2010;87(1):74-86.
379. Riccioli C, Leroy N, Pelayo S. The PSIP approach to account for human factors in Adverse Drug Events: preliminary field studies. *Stud Health Technol Inform* 2009;148:197-205.
380. Hartmann Hamilton AR, Anhoj J, Hellebek A, et al. Computerised Physician Order Entry (CPOE). *Stud Health Technol Inform* 2009;148:159-62.
381. Kucher N, Puck M, Blaser J, et al. Physician compliance with advanced electronic alerts for preventing venous thromboembolism among hospitalized medical patients. *Journal of Thrombosis & Haemostasis* 2009;7(8):1291-6.
382. van der SH, Mulder A, van Gelder T, et al. Drug safety alert generation and overriding in a large Dutch university medical centre. *Pharmacoepidemiol Drug Saf* 2009;18(10):941-7.
383. Khajouei R, de Jongh D, Jaspers MW. Usability evaluation of a computerized physician order entry for medication ordering. *Stud Health Technol Inform* 2009;150:532-6.
384. Hellstrom L, Waern K, Montelius E, et al. Physicians' attitudes towards ePrescribing--evaluation of a Swedish full-scale implementation. *BMC Med Inform Decis Mak* 2009;9:37
385. Weingart SN, Simchowitz B, Shiman L, et al. Clinicians' assessments of electronic medication safety alerts in ambulatory care. *Arch Intern Med* 2009;169(17):1627-32.
386. Milner KK, Healy D, Barry KL, et al. State mental health policy: implementation of computerized medication prescribing algorithms in a community mental health system. *Psychiatr Serv* 2009;60(8):1010-2.
387. Johansson P, Petersson G, Nilsson G. A mobile medicine decision support system for district nurses. *Stud Health Technol Inform* 2009;146:516-20.
388. McGee NM, Reeder G, Regan TS, et al. Private health plans perspectives: Electronic personal health records and electronic prescribing. *American Health and Drug Benefits* 2009;2(6):252-9.
389. Balust J, Egger Halbeis CB, Macario A. Prevalence of anaesthesia information management systems in university-affiliated hospitals in Europe. *Eur J Anaesthesiol* 2010;27(2):202-8.
390. Riaz I, Williams SD. Impact of a new electronic discharge system on the prevalence of prescribing errors. *Int J Pharm Pract* 2010;18(SUPPL. 1):22
391. Hogan SO, Kissam SM. Measuring meaningful use. *Health Aff (Millwood)* 2010;29(4):601-6.
392. Lesselroth B, Adams S, Felder R, et al. National patient safety goals. Using consumer-based kiosk technology to improve and standardize medication reconciliation in a specialty care setting. *Jt Comm J Qual Patient Saf* 2009;35(5):264-70.
393. Finkelstein J, Dennison C R. A Pilot Study of Home Automated Telemanagement (HAT) System in African Americans with Congestive Heart Failure. In Piscataway, NJ, USA: IEEE; 2010. p.90-4.
394. Lei T, Xingshe Z, Zhiwen Y and others. Adaptive Prompting based on Petri Net in a Smart Medication System. In Piscataway, NJ, USA: IEEE; 2010. p.328-33.
395. Rahimi B, Vimarlund V, Mokhtar R and others. Integrated electronic prescribing systems: pharmacists' perceptions of impact on work performance and patient safety. In Athens, Greece: WSEAS Press; 2009. p.301-6.
396. Barakah D M, Alwakeel S S. Impact of CPOE on physicians and dentists' work performance at king saud medical complex hospital: a case study. In Hong Kong, China: International Association of Engineers; 2009. p.296-8.

397. Field TS, Rochon P, Lee M, et al. Computerized clinical decision support during medication ordering for long-term care residents with renal insufficiency. *J Am Med Inform Assoc* 2009;16(4):480-5.
398. Peterson JF, Rosenbaum BP, Waitman LR, et al. Physicians' response to guided geriatric dosing: initial results from a randomized trial. *Stud Health Technol Inform* 2007;129(Pt:2):2-40.
399. Paul M, Andreassen S, Tacconelli E, et al. Improving empirical antibiotic treatment using TREAT, a computerized decision support system: cluster randomized trial. *J Antimicrob Chemother* 2006;58(6):1238-45.
400. Colpaert K, Claus B, Somers A, et al. Impact of computerized physician order entry on medication prescription errors in the intensive care unit: a controlled cross-sectional trial. *Critical Care (London, England)* 2006;10(1):R21
401. McGregor JC, Weekes E, Forrest GN, et al. Impact of a computerized clinical decision support system on reducing inappropriate antimicrobial use: a randomized controlled trial. *J Am Med Inform Assoc* 2006;13(4):378-84.
402. Kucher N, Koo S, Quiroz R, et al. Electronic alerts to prevent venous thromboembolism among hospitalized patients. *N Engl J Med* 2005;352(10):969-77.
403. Zanetti G, Flanagan HL, Jr., Cohn LH, et al. Improvement of intraoperative antibiotic prophylaxis in prolonged cardiac surgery by automated alerts in the operating room. *InfectControl Hosp Epidemiol* 2003;24(1):13-6.
404. Dexter PR, Perkins S, Overhage JM, et al. A computerized reminder system to increase the use of preventive care for hospitalized patients. *N Engl J Med* 2001;345(13):965-70.
405. Shojania KG, Yokoe D, Platt R, et al. Reducing vancomycin use utilizing a computer guideline: results of a randomized controlled trial.[see comment]. *J Am Med Inform Assoc* 1998;5(6):554-62.
406. Bates DW, Leape LL, Cullen DJ, et al. Effect of computerized physician order entry and a team intervention on prevention of serious medication errors.[see comment]. *JAMA* 1998;280(15):1311-6.
407. Overhage JM, Tierney WM, Zhou XH, et al. A randomized trial of "corollary orders" to prevent errors of omission. *J Am Med Inform Assoc* 1997;4(5):364-75.
408. Rasmussen LM, Phanareth K, Nolte H, et al. Internet-based monitoring of asthma: a long-term, randomized clinical study of 300 asthmatic subjects. *J Allergy Clin Immunol* 2005;115(6):1137-42.
409. Evans RS, Classen DC, Pestotnik SL, et al. Improving empiric antibiotic selection using computer decision support. *Arch Intern Med* 1994;154(8):878-84.
410. Dexter PR, Maharry K, Jones K, et al. Inpatient computer-based standing orders vs physician reminders to increase influenza and pneumococcal vaccination rates: A randomized trial. *JAMA* 2004;292(19):2366-71.
411. Fiks AG, Hunter K, Localio R, et al. Impact of electronic health record-based alerts on influenza vaccination for children with asthma. *Pediatrics* 2009;124(1):159-69.
412. Rood E, Bosman R, van der Spoel J, et al. Use of a computerized guideline for glucose regulation in the intensive care unit improved both guideline adherence and glucose regulation. *J Am Med Inform Assoc* 2005;12(2):172-80.
413. Terrell KM, Perkins AJ, Dexter PR, et al. Computerized decision support to reduce potentially inappropriate prescribing to older emergency department patients: A randomized, controlled trial. *J Am Geriatr Soc* 2009;57(8):1388-94.
414. Weir CJ, Lees KR, MacWalter RS, et al. Cluster-randomized, controlled trial of computer-based decision support for selecting long-term anti-thrombotic therapy after acute ischaemic stroke. *QJM: An International Journal of Medicine* 2003;96(2):143-53.

415. Overhage J, Tierney W, McDonald C. Computer reminders to implement preventive care guidelines for hospitalized patients. *Arch Intern Med* 1996;156(14):1551-6.
416. Terrell KM, Perkins AJ, Dexter PR, et al. Computerized decision support to reduce potentially inappropriate prescribing to older emergency department patients: a randomized, controlled trial. *J Am Geriatr Soc* 2009;57(8):1388-94.
417. Quinzler R, Schmitt SP, Pritsch M, et al. Substantial reduction of inappropriate tablet splitting with computerised decision support: a prospective intervention study assessing potential benefit and harm. *BMC Med Inform Decis Mak* 2009;9:30-5.
418. Wilkes JJ, Zaoutis TE, Keren R, et al. Treatment with oseltamivir in children hospitalized with community-acquired, laboratory-confirmed influenza: review of five seasons and evaluation of an electronic reminder. *J Hosp Med* 2009;4(3):171-8.
419. Stone WM, Smith BE, Shaft JD, et al. Impact of a computerized physician order-entry system. *J Am Coll Surg* 2009;208(5):960-7.
420. Sellier E, Colombet I, Sabatier B, et al. Effect of alerts for drug dosage adjustment in inpatients with renal insufficiency. *J Am Med Inform Assoc* 2009;16(2):203-10.
421. Small MD, Barrett A, Price GM. The impact of computerized prescribing on error rate in a department of Oncology/Hematology. *J Oncol Pharm Pract* 2008;14(4):181-7.
422. Scotton DW, Wierman H, Coughlan A, et al. Assessing the appropriate use of metformin in an inpatient setting and the effectiveness of two pharmacy-based measures to improve guideline adherence. *Qual Manag Health Care* 2009;18(1):71-6.
423. Liu SA, Chiu YT, Lin WD, et al. Using information technology to reduce the inappropriate use of surgical prophylactic antibiotic. *Eur Arch Otorhinolaryngol* 2008;265(9):1109-12.
424. Gerard MN, Trick WE, Das K, et al. Use of clinical decision support to increase influenza vaccination: multi-year evolution of the system. *J Am Med Inform Assoc* 2008;15(6):776-9.
425. Lecumberri R, Marques M, Diaz-Navarlaz MT, et al. Maintained effectiveness of an electronic alert system to prevent venous thromboembolism among hospitalized patients. *Thrombosis & Haemostasis* 2008;100(4):699-704.
426. Kim JY, Sohn JW, Park DW, et al. Control of extended-spectrum {beta}-lactamase-producing *Klebsiella pneumoniae* using a computer-assisted management program to restrict third-generation cephalosporin use. *J Antimicrob Chemother* 2008;62(2):416-21.
427. Sobieraj DM. Development and implementation of a program to assess medical patients' need for venous thromboembolism prophylaxis. *Am J Health Syst Pharm* 2008;65(18):1755-60.
428. Busing KL, Thursky KA, Black JF, et al. Improving antibiotic prescribing for adults with community acquired pneumonia: Does a computerised decision support system achieve more than academic detailing alone?--A time series analysis. *BMC Med Inform Decis Mak* 2008;8:35
429. Cunningham TR, Geller ES, Clarke SW. Impact of electronic prescribing in a hospital setting: a process-focused evaluation. *Int J Med Inf* 2008;77(8):546-54.
430. Rohrig R, Niczko EJ, Beutefuhr H, et al. Examination of computer assisted prescribing of an initial calculated antibiotic treatment. *Stud Health Technol Inform* 2008;136:63-8.
431. Cote GA, Rice JP, Bulsiewicz W, et al. Use of physician education and computer alert to improve targeted use of gastroprotection among NSAID users. *Am J Gastroenterol* 2008;103(5):1097-103.
432. Walsh KE, Landrigan CP, Adams WG, et al. Effect of computer order entry on prevention of serious medication errors in hospitalized children. *Pediatrics* 2008;121(3):e421-e427
433. Kooij FO, Klok T, Hollmann MW, et al. Decision support increases guideline adherence for prescribing postoperative nausea and vomiting prophylaxis. *Anesth Analg* 2008;106(3):893-8.
434. Jani YH, Ghaleb MA, Marks SD, et al. Electronic prescribing reduced prescribing errors in a pediatric renal outpatient clinic. *J Pediatr* 2008;152(2):214-8.

435. Karson AS, Campbell EJ, Panagou CM, et al. Using computerized provider order entry application to improve compliance with co-signature of verbal orders. *AMIA Proceedings* 2007;1004
436. Delgado SE, Soler VM, Perez Menendez-Conde C, et al. Prescription errors after the implementation of an electronic prescribing system. *Farm Hosp* 2007;31(4):223-30.
437. Wrona S, Chisolm DJ, Powers M, et al. Improving processes of care in patient-controlled analgesia: the impact of computerized order sets and acute pain service patient management. *Paediatr Anaesth* 2007;17(11):1083-9.
438. Mahoney CD, Berard-Collins CM, Coleman R, et al. Effects of an integrated clinical information system on medication safety in a multi-hospital setting. *Am J Health Syst Pharm* 2007;64(18):1969-77.
439. Franklin BD, O'Grady K, Donyai P, et al. The impact of a closed-loop electronic prescribing and administration system on prescribing errors, administration errors and staff time: a before-and-after study. *Qual Safe Health Care* 2007;16(4):279-84.
440. Agostini J, Shang Y, Inouye S. Use of a computer-based reminder to improve sedative-hypnotic prescribing in older hospitalized patients. *J Am Geriatr Soc* 2007;55:43-8.
441. Voeffray M, Pannatier A, Stupp R, et al. Effect of computerisation on the quality and safety of chemotherapy prescription. *Qual Safe Health Care* 2006;15(6):418-21.
442. Abboud PA, Ancheta R, McKibben M, et al. Impact of workflow-integrated corollary orders on aminoglycoside monitoring in children. *Health Informatics Journal* 2006;12(3):187-98.
443. Kaplan JM, Ancheta R, Jacobs BR, et al. Inpatient verbal orders and the impact of computerized provider order entry. *J Pediatr* 2006;149(4):461-7.
444. Eslami S, Abu-Hanna A, de Keizer NF, et al. Errors associated with applying decision support by suggesting default doses for aminoglycosides. *Drug Saf* 2006;29(9):803-9.
445. Morrison RS, Meier DE, Fischberg D, et al. Improving the management of pain in hospitalized adults. *Arch Intern Med* 2006;166(9):1033-9.
446. Chisolm DJ, McAlearney AS, Veneris S, et al. The role of computerized order sets in pediatric inpatient asthma treatment. *Pediatr Allergy Immunol* 2006;17(3):199-206.
447. Kim GR, Chen AR, Arceci RJ, et al. Error reduction in pediatric chemotherapy: computerized order entry and failure modes and effects analysis. *Arch Pediatr Adolesc Med* 2006;160(5):495-8.
448. Ozdas A, Speroff T, Waitman LR, et al. Integrating "best of care" protocols into clinicians' workflow via care provider order entry: impact on quality-of-care indicators for acute myocardial infarction. *J Am Med Inform Assoc* 2006;13(2):188-96.
449. Butler J, Speroff T, Arbogast PG, et al. Improved compliance with quality measures at hospital discharge with a computerized physician order entry system. *Am Heart J* 2006;151(3):643-53.
450. Shulman R, Singer M, Goldstone J, et al. Medication errors: a prospective cohort study of hand-written and computerised physician order entry in the intensive care unit. *Critical Care (London, England)* 2005;9(5):R516-R521
451. St Jacques P, Sanders N, Patel N, et al. Improving timely surgical antibiotic prophylaxis redosing administration using computerized record prompts. *Surgical Infections* 2005;6(2):215-21.
452. Sintchenko V, Iredell JR, Gilbert GL, et al. Handheld computer-based decision support reduces patient length of stay and antibiotic prescribing in critical care. *J Am Med Inform Assoc* 2005;12(4):398-402.
453. Oliven A, Michalake I, Zalman D, et al. Prevention of prescription errors by computerized, on-line surveillance of drug order entry. *Int J Med Inf* 2005;74(5):377-86.
454. Galanter WL, Didomenico RJ, Polikaitis A. A trial of automated decision support alerts for contraindicated medications using computerized physician order entry. *J Am Med Inform Assoc* 2005;12(3):269-74.

455. Upperman JS, Staley P, Friend K, et al. The impact of hospitalwide computerized physician order entry on medical errors in a pediatric hospital. *J Pediatr Surg* 2005;40(1):57-9.
456. Nash IS, Rojas M, Hebert P, et al. Reducing excessive medication administration in hospitalized adults with renal dysfunction. *Am J Med Qual* 2005;20(2):64-9.
457. Spencer DC, Leininger A, Daniels R, et al. Effect of a computerized prescriber-order-entry system on reported medication errors. *Am J Health Syst Pharm* 2005;62(4):416-9.
458. Bernstein SL, Whitaker D, Winograd J, et al. An electronic chart prompt to decrease proprietary antibiotic prescription to self-pay patients. *Acad Emerg Med* 2005;12(3):225-31.
459. Ali NA, Mekhjian HS, Kuehn PL, et al. Specificity of computerized physician order entry has a significant effect on the efficiency of workflow for critically ill patients. *Crit Care Med* 2005;33(1):110-4.
460. Hulgan T, Rosenbloom ST, Hargrove F, et al. Oral quinolones in hospitalized patients: an evaluation of a computerized decision support intervention. *J Intern Med* 2004;256(4):349-57.
461. Galanter WL, Polikaitis A, Didomenico RJ. A trial of automated safety alerts for inpatient digoxin use with computerized physician order entry. *J Am Med Inform Assoc* 2004;11(4):270-7.
462. Bogucki B, Jacobs BR, Hingle J, et al. Computerized reminders reduce the use of medications during shortages. *J Am Med Inform Assoc* 2004;11(4):278-80.
463. Cordero L, Kuehn L, Kumar RR, et al. Impact of computerized physician order entry on clinical practice in a newborn intensive care unit. *J Perinatol* 2004;24(2):88-93.
464. Fischer MA, Solomon DH, Teich JM, et al. Conversion from intravenous to oral medications: assessment of a computerized intervention for hospitalized patients. *Arch Intern Med* 2003;163(21):2585-9.
465. Fontan JE, Maneglier V, Nguyen VX, et al. Medication errors in hospitals: computerized unit dose drug dispensing system versus ward stock distribution system. *Pharm World Sci* 2003;25(3):112-7.
466. Achtmeyer CE, Payne TH, Anawalt BD. Computer order entry system decreased use of sliding scale insulin regimens. *Methods Inf Med* 2002;41(4):277-81.
467. Hwang JI, Park HA, Bakken S. Impact of a physician's order entry (POE) system on physicians' ordering patterns and patient length of stay. *Int J Med Inf* 2002;65(3):213-23.
468. Chertow GM, Lee J, Kuperman GJ, et al. Guided medication dosing for inpatients with renal insufficiency. *JAMA* 2001;286(22):2839-44.
469. Mullett CJ, Evans RS, Christenson JC, et al. Development and impact of a computerized pediatric antiinfective decision support program. *Pediatrics* 2001;108(4):e75
470. Teich JM, Merchia PR, Schmitz JL, et al. Effects of computerized physician order entry on prescribing practices. *Arch Intern Med* 2000;160(18):2741-7.
471. Durieux P, Nizard R, Ravaud P, et al. A clinical decision support system for prevention of venous thromboembolism: effect on physician behavior. *JAMA* 2000;283(21):2816-21.
472. Koide D, Ohe K, Ross-Degnan D, et al. Computerized reminders to monitor liver function to improve the use of etretinate. *Int J Med Inf* 2000;57(1):11-9.
473. McMullin ST, Reichley RM, Watson LA, et al. Impact of a Web-based clinical information system on cisapride drug interactions and patient safety.[see comment]. *Arch Intern Med* 1999;159(17):2077-82.
474. Bates DW, Teich JM, Lee J, et al. The impact of computerized physician order entry on medication error prevention. *J Am Med Inform Assoc* 1999;6(4):313-21.
475. Evans RS, Pestotnik SL, Classen DC, et al. A computer-assisted management program for antibiotics and other antiinfective agents.[see comment]. *N Engl J Med* 1998;338(4):232-8.

476. Goethe JW, Schwartz HI, Szarek BL. Physician compliance with practice guidelines. *Conn Med* 1997;61(9):553-8.
477. Evans RS, Pestotnik SL, Burke JP, et al. Reducing the duration of prophylactic antibiotic use through computer monitoring of surgical patients. *DICP* 1990;24(4):351-4.
478. Clancy CM, Gelfman D, Poses RM. A strategy to improve the utilization of pneumococcal vaccine. *J Gen Intern Med* 1992;7(1):14-8.
479. Igboechi CA, Ng C, Yang C, et al. Impact of computerized prescriber order entry on medication errors at an acute tertiary care hospital. *Hosp Pharm* 2003;38(3):227-31.
480. Lin C, Payne T, Nichol W, et al. Evaluating Clinical Decision Support Systems: Monitoring CPOE Order Check Override Rates in the Department of Veterans Affairs' Computerized Patient Record System. *J Am Med Inform Assoc* 2008;15(5):620-6.
481. Riggio JM, Cooper M, Leiby B, et al. Effectiveness of a clinical decision support system to identify heparin induced thrombocytopenia. *J Thromb Thrombolysis* 2009;28(2):124-31.
482. Madaras-Kelly KJ, Remington RE, Lewis PG, et al. Evaluation of an intervention designed to decrease the rate of nosocomial methicillin-resistant *Staphylococcus aureus* infection by encouraging decreased fluoroquinolone use. *INFECT CONTROL HOSP EPIDEMIOL* 2006;27(2):155-69.
483. Were MC, Abernathy G, Hui SL, et al. Using computerized provider order entry and clinical decision support to improve referring physicians' implementation of consultants' medical recommendations. *J Am Med Inform Assoc* 2009;16(2):196-202.
484. Yu FB, Menachemi N, Berner ES, et al. Full implementation of computerized physician order entry and medication-related quality outcomes: A study of 3364 hospitals. *Am J Med Qual* 2009;24(4):278-86.
485. Niemi K, Geary S, Quinn B, et al. Implementation and evaluation of electronic clinical decision support for compliance with pneumonia and heart failure quality indicators. *Am J Health Syst Pharm* 2009;66(4):389-97.
486. Griffey RT. Guided medication dosing for elderly emergency department patients using a real-time, computerized decision support tool. *Ann Emerg Med* 2009;54(3):265
487. Bates D, Boyle D, Teich J. Impact of computerized physician order entry on physician time. *Proceedings of the AMIA Symposium* 1994;996
488. Bates D, Shu K, Narasimhan D, et al. Comparing time spent writing orders on paper and physician computer order entry. *Proceedings of the AMIA Symposium* 2000;965
489. Novis SJ, Havelka GE, Ostrowski D, et al. Prevention of thromboembolic events in surgical patients through the creation and implementation of a computerized risk assessment program. *J Vasc Surg* 2010;51(3):648-54.
490. Maynard GA, Morris TA, Jenkins IH, et al. Optimizing prevention of hospital-acquired venous thromboembolism (VTE): prospective validation of a VTE risk assessment model. *J Hosp Med* 2010;5(1):10-8.
491. Kazemi A, Fors UG, Tofighi S, et al. Physician order entry or nurse order entry? Comparison of two implementation strategies for a computerized order entry system aimed at reducing dosing medication errors. *J Med Internet Res* 2010;12(1):e5
492. McCluggage L, Lee K, Potter T, et al. Implementation and evaluation of vancomycin nomogram guidelines in a computerized prescriber-order-entry system. *Am J Health Syst Pharm* 2010;67(1):70-5.
493. van Doormaal JE, van den Bemt PM, Zaal RJ, et al. The influence that electronic prescribing has on medication errors and preventable adverse drug events: an interrupted time-series study. *J Am Med Inform Assoc* 2009;16(6):816-25.
494. Riggio JM, Sorokin R, Moxey ED, et al. Effectiveness of a clinical-decision-support system in improving compliance with cardiac-care quality measures and supporting resident training. *Acad Med* 2009;84(12):1719-26.

495. Kadmon G, Bron-Harlev E, Nahum E, et al. Computerized order entry with limited decision support to prevent prescription errors in a PICU. *Pediatrics* 2009;124(3):935-40.
496. Xamplas RC, Itokazu GS, Glowacki RC, et al. Implementation of an extended-infusion piperacillin-tazobactam program at an urban teaching hospital. *Am J Health Syst Pharm* 2010;67(8):622-8.
497. Mattison ML, Afonso KA, Ngo LH, et al. Preventing Potentially Inappropriate Medication Use in Hospitalized Older Patients With a Computerized Provider Order Entry Warning System. *Arch Intern Med* 2010;170(15):1331-6.
498. Kooij FO, Klok T, Hollmann MW, et al. Automated reminders increase adherence to guidelines for administration of prophylaxis for postoperative nausea and vomiting. *Eur J Anaesthesiol* 2010;27(2):187-91.
499. Patterson R. A computerized reminder for prophylaxis of deep vein thrombosis in surgical patients. *Proceedings / AMIA* 1998;Annual Symposium.:573-6.
500. Paterno MD, Maviglia SM, Gorman PN, et al. Tiering drug-drug interaction alerts by severity increases compliance rates. *J Am Med Inform Assoc* 2009;16(1):40-6.
501. Peterson JF, Kuperman GJ, Shek C, et al. Guided prescription of psychotropic medications for geriatric inpatients. *Arch Intern Med* 2005;165(7):802-7.
502. Uttaro T, Finnerty M, White T, et al. Reduction of concurrent antipsychotic prescribing practices through the use of PSYCKES. *Adm Policy Ment Health* 2007;34(1):57-61.
503. Zhan C, Hicks RW, Blanchette CM, et al. Potential benefits and problems with computerized prescriber order entry: analysis of a voluntary medication error-reporting database. *Am J Health Syst Pharm* 2006;63(4):353-8.
504. Javitt JC, Rebitzer JB, Reisman L. Information technology and medical missteps: evidence from a randomized trial. *J Health Econ* 2008;27(3):585-602.
505. Hicks LS, Sequist TD, Ayanian JZ, et al. Impact of computerized decision support on blood pressure management and control: a randomized controlled trial. *J Gen Intern Med* 2008;23(4):429-41.
506. Martens JD, van der WT, Severens JL, et al. The effect of computer reminders on GPs' prescribing behaviour: a cluster-randomised trial. *Int J Med Inf* 2007;76(Suppl 3):S403-S416
507. Raebel MA, Charles J, Dugan J, et al. Randomized trial to improve prescribing safety in ambulatory elderly patients. *J Am Geriatr Soc* 2007;55(7):977-85.
508. Raebel MA, Carroll NM, Kelleher JA, et al. Randomized trial to improve prescribing safety during pregnancy. *J Am Med Inform Assoc* 2007;14(4):440-50.
509. Bailey TC, Noirot LA, Blickensderfer A, et al. An intervention to improve secondary prevention of coronary heart disease. *Arch Intern Med* 2007;167(6):586-90.
510. Fretheim A, Aaserud M, Oxman AD. Rational prescribing in primary care (RaPP): economic evaluation of an intervention to improve professional practice. *PLoS Medicine* 2006;3(6):e216
511. Palen TE, Raebel M, Lyons E, et al. Evaluation of laboratory monitoring alerts within a computerized physician order entry system for medication orders. *Am J Manag Care* 2006;12(7):389-95.
512. Feldstein AC, Smith DH, Perrin N, et al. Reducing warfarin medication interactions: an interrupted time series evaluation. *Arch Intern Med* 2006;166(9):1009-15.
513. Feldstein A, Elmer PJ, Smith DH, et al. Electronic medical record reminder improves osteoporosis management after a fracture: a randomized, controlled trial. *J Am Geriatr Soc* 2006;54(3):450-7.
514. Berner ES, Houston TK, Ray MN, et al. Improving ambulatory prescribing safety with a handheld decision support system: a randomized controlled trial. *J Am Med Inform Assoc* 2006;13(2):171-9.

515. Lester WT, Grant RW, Barnett GO, et al. Randomized controlled trial of an informatics-based intervention to increase statin prescription for secondary prevention of coronary disease. *J Gen Intern Med* 2006;21(1):22-9.
516. Raebel MA, Lyons EE, Chester EA, et al. Improving laboratory monitoring at initiation of drug therapy in ambulatory care: a randomized trial. *Arch Intern Med* 2005;165(20):2395-401.
517. Bloomfield HE, Nelson DB, van Ryn M, et al. A trial of education, prompts, and opinion leaders to improve prescription of lipid modifying therapy by primary care physicians for patients with ischemic heart disease. *Qual Safe Health Care* 2005;14(4):258-63.
518. Tierney WM, Overhage JM, Murray MD, et al. Can computer-generated evidence-based care suggestions enhance evidence-based management of asthma and chronic obstructive pulmonary disease? A randomized, controlled trial. *Health Serv Res* 2005;40(2):477-97.
519. Tierney WM, Overhage JM, Murray MD, et al. Effects of computerized guidelines for managing heart disease in primary care. *J Gen Intern Med* 2003;18(12):967-76.
520. Rollman BL, Hanusa BH, Lowe HJ, et al. A randomized trial using computerized decision support to improve treatment of major depression in primary care. *J Gen Intern Med* 2002;17(7):493-503.
521. Krall MA, Traunweiser K, Towery W. Effectiveness of an electronic medical record clinical quality alert prepared by off-line data analysis. *Stud Health Technol Inform* 2004;107(Pt:1):1-9.
522. Filippi A, Sabatini A, Badioli L, et al. Effects of an automated electronic reminder in changing the antiplatelet drug-prescribing behavior among Italian general practitioners in diabetic patients: an intervention trial. *Diabetes Care* 2003;26(5):1497-500.
523. Flottorp S, Oxman AD, Havelsrud K, et al. Cluster randomised controlled trial of tailored interventions to improve the management of urinary tract infections in women and sore throat. *BMJ* 2002;325(7360):367
524. Frances CD, Alperin P, Adler JS, et al. Does a fixed physician reminder system improve the care of patients with coronary artery disease? A randomized controlled trial. *West J Med* 2001;175(3):165-6.
525. Christakis DA, Zimmerman FJ, Wright JA, et al. A randomized controlled trial of point-of-care evidence to improve the antibiotic prescribing practices for otitis media in children. *Pediatrics* 2001;107(2):e15
526. Montgomery AA, Fahey T, Peters TJ, et al. Evaluation of computer based clinical decision support system and risk chart for management of hypertension in primary care: randomised controlled trial.[see comment]. *BMJ* 2000;320(7236):686-90.
527. Safran C, Rind D, Davis R, et al. Guidelines for management of HIV infection with computer-based patient's record. *The Lancet* 1995;346:341-6.
528. Cobos A, Cobos A, Vilaseca J, et al. Cost effectiveness of a clinical decision support system based on the recommendations of the European Society of Cardiology and other societies for the management of hypercholesterolemia: Report of a cluster-randomized trial. *Disease Management and Health Outcomes* 2005;13(6):421-32.
529. Davis RL, Wright J, Chalmers F, et al. A cluster randomized clinical trial to improve prescribing patterns in ambulatory pediatrics. *PLoS Clinical Trials* 2007;2(5):e25
530. Rosser WW, Hutchison B, McDowell I, et al. Use of reminders to increase compliance with tetanus booster vaccination. *Can Med Assoc J* 1992;146(6):911-7.
531. Van Wyk JT, van Wijk M, Sturkenboom M, et al. Electronic alerts versus on-demand decision support to improve dyslipidemia treatment: A cluster randomized controlled trial. *Circulation* 2008;117(3):371-8.
532. Overhage JM, Perkins S, Tierney WM, et al. Controlled trial of direct physician order entry: Effects on physicians' time utilization in ambulatory primary care internal medicine practices. *J Am Med Inform Assoc* 2001;8(4):361-9.

533. Tamblyn R, Huang A, Perreault R, et al. The medical office of the 21st century (MOXXI): Effectiveness of computerized decision-making support in reducing inappropriate prescribing in primary care. *Can Med Assoc J* 2003;169(6):549-56.
534. Bertoni A, Bonds D, Chen H, et al. Impact of a multifaceted intervention on cholesterol management in primary care practices. *Arch Intern Med* 2009;169(7):678-86.
535. Fortuna R, Zhang F, Ross-Degnan D, et al. Reducing the prescribing of heavily marketed medications: A randomized controlled trial. *J Gen Intern Med* 2009;24(8):897-903.
536. Frank O, Litt J, Beilby J. Opportunistic electronic reminders: Improving performance of preventive care in general practice. *Aust Fam Physician* 2004;33(1/2):87-90.
537. Quinn C, Clough S, Minor J, et al. WellDoc mobile diabetes management randomized controlled trial: Change in clinical and behavioral outcomes and patient and physician satisfaction. *Diabetes Technol Ther* 2008;10(3):160-8.
538. Linder JA, Schnipper JL, Tsurikova R, et al. Documentation-based clinical decision support to improve antibiotic prescribing for acute respiratory infections in primary care: A cluster randomised controlled trial. *Inform Prim Care* 2009;17(4):231-40.
539. Bell LM, Grundmeier R, Localio R, et al. Electronic health record-based decision support to improve asthma care: A cluster-randomized trial. *Pediatrics* 2010;125(4):e770-e777
540. Johnson KB, Ho Y-X, Cala CM, et al. Showing Your Work: Impact of annotating electronic prescriptions with decision support results. *J Biomed Inform* 2010;43(2):321-5.
541. Gilutz H, Novack L, Shvartzman P, et al. Computerized community cholesterol control (4C): meeting the challenge of secondary prevention. *The Israel Medical Association journal : IMAJ* 2009;11(1):23-9.
542. Tamblyn R, Reidel K, Huang A, et al. Increasing the detection and response to adherence problems with cardiovascular medication in primary care through computerized drug management systems: a randomized controlled trial. *Med Decis Making* 2010;30(2):176-88.
543. Gill JM, Chen YX, Glutting JJ, et al. Impact of decision support in electronic medical records on lipid management in primary care. *Popul Health Manag* 2009;12(5):221-6.
544. Ross SM, Papshev D, Murphy EL, et al. Effects of electronic prescribing on formulary compliance and generic drug utilization in the ambulatory care setting: a retrospective analysis of administrative claims data. *J Manag Care Pharm* 2005;11(5):410-5.
545. Miskulin DC, Weiner DE, Tighiouart H, et al. Computerized decision support for EPO dosing in hemodialysis patients. *Am J Kidney Dis* 2009;54(6):1081-8.
546. Kralj B, Iverson D, Hotz K, et al. The impact of computerized clinical reminders on physician prescribing behavior: evidence from community oncology practice. *Am J Med Qual* 2003;18(5):197-203.
547. Lapane KL, Waring ME, Schneider KL, et al. A mixed method study of the merits of e-prescribing drug alerts in primary care. *J Gen Intern Med* 2008;23(4):442-6.
548. Tang MB, Tan ES, Tian EA, et al. Electronic e-isotretinoin prescription chart: improving physicians' adherence to isotretinoin prescription guidelines. *Australas J Dermatol* 2009;50(2):107-12.
549. Fischer MA, Vogeli C, Stedman M, et al. Effect of electronic prescribing with formulary decision support on medication use and cost. *Arch Intern Med* 2008;168(22):2433-9.
550. Steele AW, Eisert S, Witter J, et al. The effect of automated alerts on provider ordering behavior in an outpatient setting. *PLoS Medicine* 2005;2(9):e255
551. Choi SS, Jazayeri DG, Mitnick CD, et al. Implementation and initial evaluation of a Web-based nurse order entry system for multidrug-resistant tuberculosis patients in Peru. *Stud Health Technol Inform* 2004;107(Pt:1):1-6.

552. Halkin H, Katzir I, Kurman I, et al. Preventing drug interactions by online prescription screening in community pharmacies and medical practices. *J Clin Pharm Therapeut* 2001;69(4):260-5.
553. Shiffman RN, Brandt C, Liaw Y, et al. A guideline implementation system using handheld computers for office management of asthma: Effects on adherence and patient outcomes. *Pediatrics* 2000;105(4):767-73.
554. Schnipper JL, McColgan KE, Linder JA, et al. Improving management of chronic diseases with documentation-based clinical decision support: results of a pilot study. *AMIA 2008;Annual:Symposium*
555. Niiranen ST, Yli-Hietanen JM. Analysis of computer-supported oral anticoagulant treatment follow-up workflow. *Conference Proceedings: 2008;4330-2.*
556. Bouaud J, Seroussi B, Antoine EC, et al. A before-after study using OncoDoc, a guideline-based decision support-system on breast cancer management: impact upon physician prescribing behaviour. *Stud Health Technol Inform* 2001;84(Pt:1):1-4.
557. Segarra-Newnham M. Tracking vaccination rates among HIV-positive patients with a computerized reminder system. *Hosp Pharm* 2003;38(8):758-62.
558. Devine EB, Wilson-Norton JL, Lawless NM, et al. The impact of an ambulatory CPOE system on medication errors. *AMIA 2008;928*
559. Ginzburg R, Barr WB, Harris M, et al. Effect of a weight-based prescribing method within an electronic health record on prescribing errors. *Am J Health Syst Pharm* 2009;66(22):2037-41.
560. de Jong JD, Groenewegen PP, Spreeuwenberg P, et al. Do decision support systems influence variation in prescription? *BMC Health Serv Res* 2009;9:9-20.
561. Hollingworth W, Devine EB, Hansen RN, et al. The impact of e-prescribing on prescriber and staff time in ambulatory care clinics: a time motion study. *J Am Med Inform Assoc* 2007;14(6):722-30.
562. Newby DA, Fryer JL, Henry DA. Effect of computerised prescribing on use of antibiotics. *Med J Aust* 2003;178(5):210-3.
563. Rubin MA, Bateman K, Donnelly S, et al. Use of a personal digital assistant for managing antibiotic prescribing for outpatient respiratory tract infections in rural communities. *J Am Med Inform Assoc* 2006;13(6):627-34.
564. Smith DH, Perrin N, Feldstein A, et al. The impact of prescribing safety alerts for elderly persons in an electronic medical record: an interrupted time series evaluation. *Arch Intern Med* 2006;166(10):1098-104.
565. Kitahata MM, Dillingham PW, Chaiyakunapruk N, et al. Electronic human immunodeficiency virus (HIV) clinical reminder system improves adherence to practice guidelines among the University of Washington HIV Study Cohort. *Clin Infect Dis* 2003;36(6):803-11.
566. Tang PC, LaRosa M, Newcomb C, et al. Measuring the effects of reminders for outpatient influenza immunizations at the point of clinical opportunity. *J Am Med Inform Assoc* 1999;6(2):115-21.
567. Varkey P, Aponte P, Swanton C, et al. The effect of computerized physician-order entry on outpatient prescription errors. *Manag Care Interface* 2007;20(3):53-7.
568. Ledwich LJ, Harrington TM, Ayoub WT, et al. Improved influenza and pneumococcal vaccination in rheumatology patients taking immunosuppressants using an electronic health record best practice alert. *ARTHRITIS RHEUM* 2009;61(11):1505-10.
569. Ngo VN, Davis RE, Lamy L, et al. A loss-of-function RNA interference screen for molecular targets in cancer. *Nature* 2006;441(7089):106-10.
570. Kirk RC, Li-Meng GD, Packia J, et al. Computer calculated dose in paediatric prescribing. *Drug Saf* 2005;28(9):817-24.
571. Dolor RJ, Yancy WS, Jr., Owen WF, et al. Hypertension Improvement Project (HIP): study protocol and implementation challenges. *Trials* 2009;10:13
572. Kerr T, Hogg RS, Yip B, et al. Validity of self-reported adherence among injection drug users. *J Int Assoc Physicians AIDS Care* 2008;7(4):157-9.

573. Walders N, Kopel SJ, Koinis-Mitchell D, et al. Patterns of quick-relief and long-term controller medication use in pediatric asthma. *J Pediatr* 2005;146(2):177-82.
574. Murray MD, Loos B, Tu W, et al. Work patterns of ambulatory care pharmacists with access to electronic guideline-based treatment suggestions. *Am J Health Syst Pharm* 1999;56(3):225-32.
575. Astrand B, Montelius E, Petersson G, et al. Assessment of ePrescription quality: an observational study at three mail-order pharmacies. *BMC Med Inform Decis Mak* 2009;9:8
576. Wess ML, Embi PJ, Besier JL, et al. Effect of a Computerized Provider Order Entry (CPOE) System on medication orders at a community hospital and university hospital. *AMIA Proceedings* 2007;Oct 11:796-800.
577. Humphries TL, Carroll N, Chester EA, et al. Evaluation of an electronic critical drug interaction program coupled with active pharmacist intervention. *Ann Pharmacother* 2007;41(12):1979-85.
578. Nam HS, Han SW, Ahn SH, et al. Improved time intervals by implementation of computerized physician order entry-based stroke team approach. *Cerebrovasc Dis* 2007;23(4):289-93.
579. Ekedahl A, Mansson N. Unclaimed prescriptions after automated prescription transmittals to pharmacies. *Pharm World Sci* 2004;26(1):26-31.
580. Senholzi C, Gottlieb J. Pharmacist interventions after implementation of computerized prescriber order entry. *Am J Health Syst Pharm* 2003;60(18):1880-2.
581. Mekhjian HS, Kumar RR, Kuehn L, et al. Immediate benefits realized following implementation of physician order entry at an academic medical center. *J Am Med Inform Assoc* 2002;9(5):529-39.
582. Beer J. Physician order entry: A mixed blessing to pharmacy? *J Oncol Pharm Pract* 2002;8(4):
583. Mitchell D, Usher J, Gray S, et al. Evaluation and audit of a pilot of electronic prescribing and drug administration. *Journal on Information Technology in Healthcare* 2004;2(1):19-29.
584. Wietholter J, Sitterson S, Allison S. Effects of computerized prescriber order entry on pharmacy order-processing time. *Am J Health Syst Pharm* 2009;66(15):1394-8.
585. Pearce DD, Opperman JM. Electronic medical record reduces HIV medication refill response time and emergency refills in a Latino community clinic. *Int J STD AIDS* 2010;21(3):184-6.
586. Nilsson JLG, Backstrom S, Sundstrom J. Electronically transferred prescriptions: Picked up faster than paper prescriptions. *Int J Pharm Pract* 2007;15(2):157-8.
587. Sommet A, Desplas M, Lapeyre-Mestre M, et al. Drug-induced yawning: a review of the French pharmacovigilance database. *Drug Saf* 2007;30(4):327-31.
588. Reeve JF, Tenni PC, Peterson GM. An electronic prompt in dispensing software to promote clinical interventions by community pharmacists: a randomized controlled trial. *Br J Clin Pharmacol* 2008;65(3):377-85.
589. Wilson AL, Hill JJ, Wilson RG, et al. Computerized medication administration records decrease medication occurrences. *Pharm Pract Manag Q* 1997;17(1):17-29.
590. Alvarez Diaz AM, Delgado SE, Perez Menendez-Conde C, et al. New technologies applied to the medication-dispensing process, error analysis and contributing factors. [Spanish]. *Farm Hosp* 2010;34(2):59-67.
591. Cocosila M, Coursaris C, Yufei Y. M-healthcare for patient self-management: a case for diabetics. *Int J Electron Healthc* 2004;1(2):221-41.
592. Persell SD, Denecke-Dattalo TA, Dunham DP, et al. Evidence-based medicine. Patient-directed intervention versus clinician reminders alone to improve aspirin use in diabetes: A cluster randomized trial. *Jt Comm J Qual Patient Saf* 2008;34(2):98-105.
593. Helmons PJ, Wargel LN, Daniels CE. Effect of bar-code-assisted medication administration on medication administration errors and accuracy in multiple patient care areas. *Am J Health Syst Pharm* 2009;66(13):1202-10.

594. DeYoung JL, Vanderkooi ME, Barletta JF. Effect of bar-code-assisted medication administration on medication error rates in an adult medical intensive care unit. *Am J Health Syst Pharm* 2009;66(12):1110-5.
595. Taylor JA, Loan LA, Kamara J, et al. Medication administration variances before and after implementation of computerized physician order entry in a neonatal intensive care unit. *Pediatrics* 2008;121(1):123-8.
596. Wax DB, Beilin Y, Levin M, et al. The effect of an interactive visual reminder in an anesthesia information management system on timeliness of prophylactic antibiotic administration. *Anesth Analg* 2007;104(6):1462-6.
597. Banet GA, Jeffe DB, Williams JA, et al. Effects of implementing computerized practitioner order entry and nursing documentation on nursing workflow in an emergency department. *J Healthc Inf Manag* 2006;20(2):45-54.
598. Low DK, Belcher JV. Reporting medication errors through computerized medication administration. *Comput Inform Nurs* 2002;20(5):178-83.
599. Shirley KL. Effect of an automated dispensing system on medication administration time. *Am J Health Syst Pharm* 1999;56(15):1542-5.
600. Poon EG, Keohane CA, Bane A, et al. Impact of barcode medication administration technology on how nurses spend their time providing patient care. *J Nurs Adm* 2006;38(12):541-9.
601. Poon EG, Keohane CA, Yoon CS, et al. Effect of bar-code technology on the safety of medication administration. *N Engl J Med* 2010;362(18):1698-707.
602. Climent C, Font-Noguera I, Poveda Andres JL, et al. Medication errors in a tertiary hospital with three different drug delivery systems. *Farm Hosp* 2008;32(1):18-24.
603. Handler SM, Nace DA, Studenski SA, et al. Medication error reporting in long term care. *Am J Geriatr Pharmacother* 2004;2(3):190-6.
604. Fillit H, Rockwood K, Woodhouse L. *Brocklehurst's Textbook of Geriatric Medicine and Gerontology*. 7th. Churchill Livingstone; 2010. Geriatric Pharmacotherapy and Polypharmacy.
605. Hayward GL, Parnes AJ, Simon SR. Using health information technology to improve drug monitoring: a systematic review. *Pharmacoepidemiol Drug Saf* 2009;18(12):1232-7.
606. Handler SM, Altman RL, Perera S, et al. A systematic review of the performance characteristics of clinical event monitor signals used to detect adverse drug events in the hospital setting. *J Am Med Inform Assoc* 2007;14(4):451-8.
607. Chaudhry B, Wang J, Wu S, et al. Systematic review: impact of health information technology on quality, efficiency, and costs of medical care. *Ann Intern Med* 2006;144(10):742-52.
608. Okon TR, Lutz PS, Liang H. Improved pain resolution in hospitalized patients through targeting of pain mismanagement as medical error. *J Pain Symptom Manage* 2009;37(6):1039-49.
609. Lo HG, Matheny ME, Seger DL, et al. Impact of non-interruptive medication laboratory monitoring alerts in ambulatory care. *J Am Med Inform Assoc* 2009;16(1):66-71.
610. Grant RW, Wald JS, Schnipper JL, et al. Practice-linked online personal health records for type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med* 2008;168(16):1776-82.
611. Matheny ME, Sequist TD, Seger AC, et al. A randomized trial of electronic clinical reminders to improve medication laboratory monitoring. *J Am Med Inform Assoc* 2008;15(4):424-9.
612. Feldstein AC, Smith DH, Perrin N, et al. Improved therapeutic monitoring with several interventions: a randomized trial. *Arch Intern Med* 2006;166(17):1848-54.
613. Kuilboer MM, van Wijk MA, Mosseveld M, et al. Computed critiquing integrated into daily clinical practice affects physicians' behavior--a randomized clinical trial with AsthmaCritic. *Methods Inf Med* 2006;45(4):447-54.

614. Evans RS, Pestotnik SL, Classen DC, et al. Evaluation of a computer-assisted antibiotic-dose monitor. *Ann Pharmacother* 1999;33(10):1026-31.
615. Rind DM, Safran C, Phillips RS, et al. Effect of computer-based alerts on the treatment and outcomes of hospitalized patients. *Arch Intern Med* 1994;154(13):1511-7.
616. Sequist TD, Gandhi T, Karson A, et al. A randomized trial of electronic clinical reminders to improve quality of care for diabetes and coronary artery disease. *J Am Med Inform Assoc* 2005;12(4):431-7.
617. DEMakis J, Beauchamp C, Cull W, et al. Improving residents' compliance with standards of ambulatory care: Results from the VA cooperative study on computerized reminders. *JAMA* 2000;284(11):1411-6.
618. White K, Lindsay A, Pryor T, et al. Application of a computerized medical decision-making process to the problem of digoxin intoxication. *J Am Coll Cardiol* 1984;4(3):571-6.
619. McDonald C. Use of a computer to detect and respond to clinical events: Its effect on clinical behavior. *Ann Intern Med* 1976;84(2):162-7.
620. Javitt J, Steinberg G, Locke T, et al. Using a claims data-based sentinel system to improve compliance with clinical guidelines: Results of a randomized prospective study. *Am J Manag Care* 2005;11:93-102.
621. Bertsche T, Askoxylakis V, Habl G, et al. Multidisciplinary pain management based on a computerized clinical decision support system in cancer pain patients. *Pain* 2009;147(1-3):20-8.
622. Chambers RM, Wilson JW, Estes LL. Computer-based monitoring as a tool for antimicrobial de-escalation. *Hosp Pharm* 2008;43(3):199-205.
623. Patel PV, Gilski D, Morrison J. Improving outcomes in high-risk populations using REACH: An inpatient cardiac risk reduction program. *Crit Pathw Cardiol* 2009;8(3):112-8.
624. Roumie CL, Elasy TA, Greevy R, et al. Improving blood pressure control through provider education, provider alerts, and patient education: a cluster randomized trial. *Ann Intern Med* 2006;145(3):165-75.
625. Bassi J, Lau F, Bardal S. Use of information technology in medication reconciliation: A scoping review. *Ann Pharmacother* 2010;44(5):885-97.
626. Bayoumi I HMHASI. Interventions to improve medication reconciliation in primary care. *Ann Pharmacother* 2009;43(10):1667-75.
627. Agrawal A, Wu WY. National patient safety goals. Reducing medication errors and improving systems reliability using an electronic medication reconciliation system. *Jt Comm J Qual Patient Saf* 2009;35(2):106-14.
628. van der Kam WJ, Meyboom dJ, Tromp TF, et al. Effects of electronic communication between the GP and the pharmacist. The quality of medication data on admission and after discharge. *Fam Pract* 2001;18(6):605-9.
629. Crosson JC, Isaacson N, Lancaster D, et al. Variation in electronic prescribing implementation among twelve ambulatory practices. *J Gen Intern Med* 2008;23(4):364-71.
630. Holman RR, Smale AD, Pemberton E, et al. Randomized controlled pilot trial of a handheld patient-oriented, insulin regimen optimizer. *Med Inform (Lond)* 1996;21(4):317-26.
631. Cook CB, Mann LJ, King EC, et al. Management of insulin therapy in urban diabetes patients is facilitated by use of an intelligent dosing system. *Diabetes Technology and Therapeutics* 2004;6(3):326-35.
632. Weingart SN, Massagli M, Cyrulik A, et al. Assessing the value of electronic prescribing in ambulatory care: A focus group study. *Int J Med Inf* 2009;78(9):571-8.

633. McCann L, Maguire R, Miller M, et al. Patients' perceptions and experiences of using a mobile phone-based advanced symptom management system (ASyMS) to monitor and manage chemotherapy related toxicity. *Eur J Cancer Care* 2009;18(2):156-64.
634. Plaza V, Cobas A, Ignacio-Garcia J, et al. Cost-effectiveness of an intervention based on the recommendations of the Global Initiative for Asthma (GINA) by a computerized clinical decision support: A trial with randomization of physicians. *Med Clin (Barc)* 2005;124(6):201-6.
635. Johansson PE, Petersson GI, Nilsson GC. Personal digital assistant with a barcode reader-A medical decision support system for nurses in home care. *Int J Med Inf* 2010;79(4):232-42.
636. Rothman BL, Sullivan AN, McDonald TW, et al. A randomized controlled trial of a computer-based physician workstation in an outpatient setting. Implementation barriers to outcome evaluation. *J Am Med Inform Assoc* 1996;3(5):340-8.
637. Graumlich JF, Novotny NL, Stephen NG, et al. Patient readmissions, emergency visits, and adverse events after software-assisted discharge from hospital: cluster randomized trial. *J Hosp Med* 2009;4(7):E11-E19
638. Rosenbloom ST, Geissbuhler AJ, Dupont WD, et al. Effect of CPOE user interface design on user-initiated access to educational and patient information during clinical care. *J Am Med Inform Assoc* 2005;12(4):458-73.
639. Ross SE, Moore LA, Earnest MA, et al. Providing a web-based online medical record with electronic communication capabilities to patients with congestive heart failure: randomized trial. *J Med Internet Res* 2004;6(2):e12
640. Musser R, Tchong J. Quantitative and qualitative comparison of text-based and graphical user interfaces for computerized provider order entry. *Proceedings of the AMIA Symposium* 2006;1041
641. Weingart SN, Hamrick HE, Tutkus S, et al. Medication safety messages for patients via the web portal: the MedCheck intervention. *Int J Med Inf* 2008;77(3):161-8.
642. Schmidt S, Sheikzadeh S, Beil B, et al. Acceptance of telemonitoring to enhance medication compliance in patients with chronic heart failure. *Telemed J E Health* 2008;14(5):426-33.
643. Shannon T, Feied C, Smith M, et al. Wireless handheld computers and voluntary utilization of computerized prescribing systems in the emergency department. *J Emerg Med* 2006;31(3):309-15.
644. Niazkhani Z, van der SH, Pirnejad H, et al. Same system, different outcomes: comparing the transitions from two paper-based systems to the same computerized physician order entry system. *Int J Med Inf* 2009;78(3):170-81.
645. Rupp MT, Warholak TL. Evaluation of e-prescribing in chain community pharmacy: best-practice recommendations. *J Am Pharm Assoc* 2008;48(3):364-70.
646. Chan S. Factors Associated With the Use of Electronic Information Systems for Drug Dispensing and Medication Administration Records in Nursing Homes. *J Am Med Dir Assoc* 2008;9(6):414-21.
647. Rohrig R, Beutefuhr H, Hartmann B, et al. Summative software evaluation of a therapeutic guideline assistance system for empiric antimicrobial therapy in ICU. *J Clin Monit Comput* 2007;21(4):203-10.
648. Liu CT, Yeh YT, Chiang IJ, et al. Development and evaluation of an integrated pharmaceutical education system. *Int J Med Inf* 2004;73(4):383-9.
649. McAlearney A, Chisolm D, Veneris S, et al. Utilization of evidence-based computerized order sets in pediatrics. *Int J Med Inf* 2006;75:501-12.
650. Kawasumi Y, Tamblyn R, Platt R, et al. Evaluation of the use of an integrated drug information system by primary care physicians for vulnerable population. *Int J Med Inf* 2008;77(2):98-106.
651. Kramer JS, Hopkins PJ, Rosendale JC, et al. Implementation of an electronic system for medication reconciliation. *Am J Health Syst Pharm* 2007;64(4):404-22.

652. Li Q, Douglas S, Hundt A et al. A heuristic usability evaluation of a computerized provider order entry (CPOE) technology. Proceedings of the IEA2006 Congress, 2006.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_204532_0_0_18/Lietal-2006-IEA-CPOEUsability.pdf
653. Wang CJ, Patel MH, Schueth AJ, et al. Perceptions of standards-based electronic prescribing systems as implemented in outpatient primary care: a physician survey. *J Am Med Inform Assoc* 2009;16(4):493-502.
654. Ghahramani N, Lendel I, Haque R, et al. User satisfaction with computerized order entry system and its effect on workplace level of stress. *J Med Syst* 2009;33(3):199-205.
655. Wilson JP, Bulatao PT, Rascati KL. Satisfaction with a computerized practitioner order-entry system at two military health care facilities. *Am J Health Syst Pharm* 2000;57(23):2188-95.
656. Weiner M, Gress T, Thiemann DR, et al. Contrasting views of physicians and nurses about an inpatient computer-based provider order-entry system. *J Am Med Inform Assoc* 1999;6(3):234-44.
657. Lee F, Teich JM, Spurr CD, et al. Implementation of physician order entry: user satisfaction and self-reported usage patterns. *J Am Med Inform Assoc* 1996;3(1):42-55.
658. Tan WS, Phang JS, Tan LK. Evaluating user satisfaction with an electronic prescription system in a primary care group. *Ann Acad Med Singapore* 2009;38(6):494-597.
659. Graumlich JF, Novotny NL, Nace GS, et al. Patient and physician perceptions after software-assisted hospital discharge: Cluster randomized trial. *J Hosp Med* 2009;4(6):356-63.
660. Rogers J, Jain NL, Hayes GM. Evaluation of an implementation of PRODIGY phase two. *AMIA Proceedings* 1999;604-8.
661. Zaidi ST, Marriott JL, Nation RL. The role of perceptions of clinicians in their adoption of a web-based antibiotic approval system: do perceptions translate into actions? *Int J Med Inf* 2008;77(1):33-40.
662. Sittig DF, Krall MA, Dykstra RH, et al. A survey of factors affecting clinician acceptance of clinical decision support. *BMC Med Inform Decis Mak* 2006;6:6
663. Lindenauer PK, Ling D, Pekow PS, et al. Physician characteristics, attitudes, and use of computerized order entry. *J Hosp Med* 2006;1(4):221-30.
664. Schectman JM, Schorling JB, Nadkarni MM, et al. Determinants of physician use of an ambulatory prescription expert system. *Int J Med Inf* 2005;74(9):711-7.
665. Kralewski JE, Dowd BE, Cole-Adeniyi T, et al. Factors influencing physician use of clinical electronic information technologies after adoption by their medical group practices. *Health Care Manage Rev* 2008;33(4):361-7.
666. Pirnejad H, Niazkhani Z, van der SH, et al. Impact of a computerized physician order entry system on nurse-physician collaboration in the medication process. *Int J Med Inf* 2008;77(11):735-44.
667. Glassman PA, Belperio P, Simon B, et al. Exposure to automated drug alerts over time: effects on clinicians' knowledge and perceptions. *Med Care* 2006;44(3):250-6.
668. Porteous T, Bond C, Robertson R, et al. Electronic transfer of prescription-related information: comparing views of patients, general practitioners, and pharmacists. *Br J Gen Pract* 2003;53(488):204-9.
669. Kirking D, Thomas J, Ascione F, et al. Detecting and preventing adverse drug interactions: The potential contribution of computers in pharmacies. *Soc Sci Med* 1986;22(1):1-8.
670. O'Morrow SD. Nurses' attitudes toward use of an automated medication management system. *Medical University of South Carolina - College of Health Professions* 2003; D H A 2000;14(2):1-13.
671. Hurley AC, Bane A, Fotakis S, et al. Nurses' satisfaction with medication administration point-of-care technology. *J Nurs Adm* 2007;37(7-8):343-9.

672. Holden RJ, Brown RL, Alper SJ, et al. That's nice, but what does IT do? Evaluating the impact of bar coded medication administration by measuring changes in the process of care. *Int J Ind Ergon* 2010;(accepted):
673. Holden R J, Scanlon M C, Brown R L and others. What is IT? New conceptualizations and measures of pediatric nurses' acceptance of bar-coded medication administration information technology. In Santa Monica, CA, USA: Human Factors and Ergonomics Society; 2008. p.768-72.
674. Topps C, Lopez L, Messmer PR, et al. Perceptions of pediatric nurses toward bar-code point-of-care medication administration. *Nurs Adm Q* 2005;29(1):102-7.
675. van Onzenoort HA, van de PA, Kessels AG, et al. Factors influencing bar-code verification by nurses during medication administration in a Dutch hospital. *Am J Health Syst Pharm* 2008;65(7):644-8.
676. Pirnejad H, Niazkhani Z, van der SH, et al. Evaluation of the impact of a CPOE system on nurse-physician communication--a mixed method study. *Methods Inf Med* 2009;48(4):350-60.
677. Tierney W, Overhage J, McDonald C, et al. Medical students' and housestaff's opinions of computerized order-writing. *Acad Med* 1994;69(5):386-9.
678. Rahimi B, Timpka T, Vimarlund V, et al. Organization-wide adoption of computerized provider order entry systems: a study based on diffusion of innovations theory. *BMC Med Inform Decis Mak* 2009;9:52
679. Drummond MF, Sculpher MJ, Torrance GW, et al. *Methods for the economic evaluation of health care programmes*. 3rd. Oxford: Oxford University Press; 2005. 2, Basic types of economic evaluation.
680. Wu RC, Laporte A, Ungar WJ. Cost-effectiveness of an electronic medication ordering and administration system in reducing adverse drug events. *J Eval Clin Pract* 2007;13(3):440-8.
681. Karnon J, McIntosh A, Dean J, et al. Modelling the expected net benefits of interventions to reduce the burden of medication errors. *J Health Serv Res Policy* 2008;13(2):85-91.
682. Evans RS, Pestotnik SL, Classen DC, et al. Prevention of adverse drug events through computerized surveillance. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1992;437-41.
683. Evans RS, Classen DC, Pestotnik SL, et al. A decision support tool for antibiotic therapy. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1995;651-5.
684. Barenfanger J, Short MA, Groesch AA. Improved antimicrobial interventions have benefits. *J Clin Microbiol* 2001;39(8):2823-8.
685. Macdonald D, Bhalla P, Cass W, et al. Computerized management of oral anticoagulant therapy: experience in major joint arthroplasty. *Can J Surg* 2002;45(1):47-52.
686. Kaushal R, Jha AK, Franz C, et al. Return on investment for a computerized physician order entry system. *J Am Med Inform Assoc* 2006;13(3):261-6.
687. Tierney W, Miller M, Overhage J, et al. Physician inpatient order writing on microcomputer workstations: effects on resource utilization. *JAMA* 1993;269(3):379-83.
688. Piontek F, Kohli R, Conlon P, et al. Effects of an adverse-drug-event alert system on cost and quality outcomes in community hospitals. *Am J Health Syst Pharm* 2010;67(8):613-20.
689. McMullin ST, Lonergan TP, Rynearson CS. Twelve-month drug cost savings related to use of an electronic prescribing system with integrated decision support in primary care. *J Manag Care Pharm* 2005;11(4):322-32.
690. McMullin ST, Lonergan TP, Rynearson CS, et al. Impact of an evidence-based computerized decision support system on primary care prescription costs. *Annals of Family Medicine* 2004;2(5):494-8.

691. Ornstein SM, MacFarlane LL, Jenkins RG, et al. Medication cost information in a computer-based patient record system. Impact on prescribing in a family medicine clinical practice. *Arch Fam Med* 1999;8(2):118-21.
692. Weingart SN, Simchowitz B, Padolsky H, et al. An empirical model to estimate the potential impact of medication safety alerts on patient safety, health care utilization, and cost in ambulatory care. *Arch Intern Med* 2009;169(16):1465-73.
693. Holdsworth MT, Fichtl RE, Raisch DW, et al. Impact of computerized prescriber order entry on the incidence of adverse drug events in pediatric inpatients. *Pediatrics* 2007;120(5):1058-66.
694. Chabot I, Moisan J, Gregoire JP, et al. Pharmacist intervention program for control of hypertension. *Ann Pharmacother* 2003;37(9):1186-93.
695. Schnipper JL, Hamann C, Ndumele CD, et al. Effect of an electronic medication reconciliation application and process redesign on potential adverse drug events: a cluster-randomized trial. *Arch Intern Med* 2009;169(8):771-80.
696. Ralston JD, Hirsch IB, Hoath J, et al. Web-based collaborative care for type 2 diabetes: a pilot randomized trial. *Diabetes Care* 2009;32(2):234-9.
697. Gurwitz JH, Field TS, Rochon P, et al. Effect of computerized provider order entry with clinical decision support on adverse drug events in the long-term care setting. *J Am Geriatr Soc* 2008;56(12):2225-33.
698. Murray MD, Harris LE, Overhage JM, et al. Failure of computerized treatment suggestions to improve health outcomes of outpatients with uncomplicated hypertension: results of a randomized controlled trial. *Pharmacotherapy* 2004;24(3):324-37.
699. Hetlevik I, Holmen J, Kruger O. Implementing clinical guidelines in the treatment of hypertension in general practice: Evaluation of patient outcome related to implementation of a computer-based clinical decision support system. *Scand J Prim Health Care* 1999;17(1):35-40.
700. Meigs J, Cagliero E, Dubey A, et al. A controlled trial of web-based diabetes disease management. *Diabetes Care* 2003;26(3):750-7.
701. Balcezak TJ, Krumholz HM, Getnick GS, et al. Utilization and effectiveness of a weight-based heparin nomogram at a large academic medical center. *Am J Manag Care* 2000;6(3):329-38.
702. Baroletti S, Munz K, Sonis J, et al. Electronic alerts for hospitalized high-VTE risk patients not receiving prophylaxis: a cohort study. *J Thromb Thrombolysis* 2008;25(2):146-50.
703. Boord JB, Sharifi M, Greevy RA, et al. Computer-based insulin infusion protocol improves glycemia control over manual protocol. *J Am Med Inform Assoc* 2007;14(3):278-87.
704. Del Beccaro MA, Jeffries HE, Eisenberg MA, et al. Computerized provider order entry implementation: no association with increased mortality rates in an intensive care unit. *Pediatrics* 2006;118(1):290-5.
705. Garthwaite EA, Will EJ, Bartlett C, et al. Patient-specific prompts in the cholesterol management of renal transplant outpatients: results and analysis of underperformance. *Transplantation* 2004;78(7):1042-7.
706. Takada M, Demizu M, Shibakawa M. Physicians' prescribing attitudes to combined therapy with statins and fibrates. *J Clin Pharm Therapeut* 2003;28(6):445-50.
707. Evans RS, Pestotnik SL, Classen DC, et al. Preventing adverse drug events in hospitalized patients. *Ann Pharmacother* 1994;28(4):523-7.
708. Vartak S, Crandall DK, Brokel JM, et al. Professional practice and innovation: transformation of emergency department processes of care with EHR, CPOE, and ER event tracking systems. *Health Information Management Journal* 2010;38(2):27-32.
709. Chen C, Chen K, Hsu CY, et al. A guideline-based decision support for pharmacological treatment can improve the quality of hyperlipidemia management. *Comput Methods Programs Biomed* 2010;97(3):280-5.

710. Fiumara K, Piovella C, Hurwitz S, et al. Multi-screen electronic alerts to augment venous thromboembolism prophylaxis. *Thrombosis & Haemostasis* 2010;103(2):312-7.
711. Pielmeier U, Andreassen S, Juliusen B, et al. The Glucosafe system for tight glycemic control in critical care: A pilot evaluation study. *J Crit Care* 2010;25(1):97-104.
712. Yu F, Salas M, Kim YI, et al. The relationship between computerized physician order entry and pediatric adverse drug events: a nested matched case-control study. *Pharmacoepidemiol Drug Saf* 2009;18(8):751-5.
713. Schnipper JL, Ndumele CD, Liang CL, et al. Effects of a subcutaneous insulin protocol, clinical education, and computerized order set on the quality of inpatient management of hyperglycemia: results of a clinical trial. *J Hosp Med* 2009;4(1):16-27.
714. Janssen B, Ludwig S, Eustermann H, et al. Improving outpatient treatment in schizophrenia: effects of computerized guideline implementation--results of a multicenter-study within the German research network on schizophrenia. *Eur Arch Psychiatry Clin Neurosci* 2010;260(1):51-7.
715. Shojania KG, Jennings A, Mayhew A, et al. The effects of on-screen, point of care computer reminders on processes and outcomes of care. (Review). *The Cochrane Library* 2009;Jul 8(3):
716. Durieux P, Trinquart L, Colombet I, et al. Computerized advice on drug dosage to improve prescribing practice. *COCHRANE DATABASE SYST REV* 2008;(3):CD002894
717. Schedlbauer A, Prasad V, Mulvaney C, et al. What evidence supports the use of computerized alerts and prompts to improve clinicians' prescribing behavior? *J Am Med Inform Assoc* 2009;16(4):531-8.
718. Yourman L, Concato J, Agostini JV. Use of computer decision support interventions to improve medication prescribing in older adults: a systematic review. *American Journal Geriatric Pharmacotherapy* 2008;6(2):119-29.
719. Ernstmann N, Ommen O, Neumann M, et al. Primary care physician's attitude towards the German e-health card project--determinants and implications. *J Med Syst* 2009;33(3):181-8.
720. Ammenwerth E. The Effect of Electronic Prescribing on Medication Errors and Adverse Drug Events: A Systematic Review. *J Am Med Inform Assoc* 2008;15(5):585-600.
721. van Rosse F, Maat B, Rademaker CM, et al. The effect of computerized physician order entry on medication prescription errors and clinical outcome in pediatric and intensive care: a systematic review. *Pediatrics* 2009;123(4):1184-90.
722. Chedoe I, Molendijk HA, Dittrich ST, et al. Incidence and nature of medication errors in neonatal intensive care with strategies to improve safety: a review of the current literature. *Drug Saf* 2007;30(6):503-13.
723. Eslami S, Abu-Hanna A, de Keizer NF. Evaluation of outpatient computerized physician medication order entry systems: a systematic review. *J Am Med Inform Assoc* 2007;14(4):400-6.
724. Wolfstadt JI, Gurwitz JH, Field TS, et al. The effect of computerized physician order entry with clinical decision support on the rates of adverse drug events: a systematic review. *J Gen Intern Med* 2008;23(4):451-8.
725. Garg AX, Adhikari NK, McDonald H, et al. Effects of computerized clinical decision support systems on practitioner performance and patient outcomes: a systematic review. *JAMA* 2005;293(10):1223-38.
726. Georgiou A, Ampt A, Creswick N, et al. Computerized Provider Order Entry--what are health professionals concerned about? A qualitative study in an Australian hospital. *Int J Med Inf* 2009;78(1):60-70.
727. Kazemi A, Ellenius J, Tofighi S, et al. CPOE in Iran--a viable prospect? Physicians' opinions on using CPOE in an Iranian teaching hospital. *Int J Med Inf* 2009;78(3):199-207.
728. Koppel R, Wetterneck T, Telles JL, et al. Workarounds to barcode medication administration systems: their occurrences, causes, and threats to patient safety. *J Am Med Inform Assoc* 2008;15(4):408-23.

729. Varonen H, Kortteisto T, Kaila M, et al. What may help or hinder the implementation of computerized decision support systems (CDSSs): a focus group study with physicians. *Fam Pract* 2008;25(3):162-7.
730. Motulsky A, Winslade N, Tamblyn R, et al. The impact of electronic prescribing on the professionalization of community pharmacists: a qualitative study of pharmacists' perception. *J Pharm Pharm Sci* 2008;11(1):131-46.
731. Agostini JV, Concato J, Inouye SK. Improving sedative-hypnotic prescribing in older hospitalized patients: provider-perceived benefits and barriers of a computer-based reminder. *J Gen Intern Med* 2008;23(Suppl 1):32-6.
732. Vogelsmeier AA, Halbesleben JR, Scott-Cawiezell JR. Technology implementation and workarounds in the nursing home. *J Am Med Inform Assoc* 2008;15(1):114-9.
733. Fields W, Snyder R. Community hospital CPOE system implementation: The lived experience of multi-disciplinary healthcare team members. *AMIA Proceedings* 2007;954
734. Wentzer HS, Bottger U, Boye N. Unintended transformations of clinical relations with a computerized physician order entry system. *Int J Med Inf* 2007;76(Suppl 3):S453-S461
735. McAlearney AS, Chisolm DJ, Schweikhart S, et al. The story behind the story: physician skepticism about relying on clinical information technologies to reduce medical errors. *Int J Med Inf* 2007;76(11-12):836-42.
736. Grossman JM, Gerland A, Reed MC, et al. Physicians' experiences using commercial e-prescribing systems. *Health Aff (Millwood)* 2007;26(3):w393-w404
737. Arar NH, Wen L, McGrath J, et al. Communicating about medications during primary care outpatient visits: the role of electronic medical records. *Inform Prim Care* 2005;13(1):13-22.
738. Avery AJ, Savelyich BS, Sheikh A, et al. Identifying and establishing consensus on the most important safety features of GP computer systems: e-Delphi study. *Inform Prim Care* 2005;13(1):3-12.
739. Feldstein A, Simon SR, Schneider J, et al. How to design computerized alerts to safe prescribing practices. *Jt Comm J Qual Patient Saf* 2004;30(11):602-13.
740. Bastholm RP, Andersen-Karlsson E, Arnhjort T, et al. Physicians' perceptions of possibilities and obstacles prior to implementing a computerised drug prescribing support system. *Int J Health Care Qual Assur Inc Leadersh Health Serv* 2004;17(4-5):173-9.
741. Ash JS, Sittig DF, Seshadri V, et al. Adding insight: a qualitative cross-site study of physician order entry. *Stud Health Technol Inform* 2004;107(Pt:2):2-7.
742. Krall MA, Sittig DF. Clinician's assessments of outpatient electronic medical record alert and reminder usability and usefulness requirements. *AMIA Proceedings* 2002;400-4.
743. Patterson ES, Cook RI, Render ML. Improving patient safety by identifying side effects from introducing bar coding in medication administration. *J Am Med Inform Assoc* 2002;9(5):540-53.
744. Novek J, Bettess S, Burke K, et al. Nurses' perceptions of the reliability of an automated medication dispensing system. *J Nurs Care Qual* 2000;14(2):1-13.
745. Ahearn MD, Kerr SJ. General practitioners' perceptions of the pharmaceutical decision-support tools in their prescribing software. *Med J Aust* 2003;179(1):34-7.
746. Lai JS, Yokoyama G, Louie C, et al. Impact of computerized prescriber order entry (CPOE) on clinical pharmacy practice: A hypothesis-generating study. *Hosp Pharm* 2007;42(10):931-8.
747. Patterson ES, Nguyen A, Halloran J, et al. Human Factors Barriers to the Effective Use of Ten HIV Clinical Reminders. *J Am Med Inform Assoc* 2004;11(1):50-9.
748. O'Grady K, Donyai P, Franklin BD. Patients' views about an electronic prescribing and drug administration system in secondary care. *BJHC & IM* 2006;23(7):15-8.

749. Weir CR, Nebeker JJR, Hicken BL, et al. A cognitive task analysis of information management strategies in a computerized provider order entry environment. *J Am Med Inform Assoc* 2007;14(1):65-75.
750. Saleem JJ, Patterson ES, Militello L, et al. Exploring barriers and facilitators to the use of computerized clinical reminders. *J Am Med Inform Assoc* 2005;12(4):438-47.
751. Boonstra A, Boddy D, Fischbacher M. The limited acceptance of an electronic prescription system by general practitioners: Reasons and practical implications. *New Tech Work Employ* 2004;19(2):128-44.
752. Koppel R, Metlay J, Cohen A, et al. Role of computerized physician order entry systems in facilitating medication errors. *JAMA* 2005;293(10):1197-203.
753. Vaziri A, Connor E, Shepherd I, et al. Are we setting about improving the safety of computerised prescribing in the right way? A workshop report. *Inform Prim Care* 2009;17(3):175-82.
754. Novak LL, Lorenzi NM. Barcode medication administration: supporting transitions in articulation work. *AMIA* 2008;Annual:515-9.
755. Graham TA, Kushniruk AW, Bullard MJ, et al. How usability of a web-based clinical decision support system has the potential to contribute to adverse medical events. *AMIA* 2008;Annual:257-61.
756. Buhner ED, Novak LL, Lorenzi NM. Implementing barcode medication administration: nurses' attitudes. *AMIA* 2008;Annual:1072
757. Nanji KC, Cina J, Patel N, et al. Overcoming barriers to the implementation of a pharmacy bar code scanning system for medication dispensing: a case study. *J Am Med Inform Assoc* 2009;16(5):645-50.
758. Schoville RR. Work-arounds and artifacts during transition to a computer physician order entry: what they are and what they mean. *J Nurs Care Qual* 2009;24(4):316-24.
759. Campbell EM, Guappone KP, Sittig DF, et al. Computerized provider order entry adoption: implications for clinical workflow. *J Gen Intern Med* 2009;24(1):21-6.
760. Cross RK, Cheevers N, Finkelstein J. Home telemanagement for patients with ulcerative colitis (UC HAT). *Dig Dis Sci* 2009;54(11):2463-72.
761. Holden RJ. Physicians' beliefs about using EMR and CPOE: In pursuit of a contextualized understanding of health it use behavior. *Int J Med Inf* 2010;79(2):71-80.
762. Beuscart-Zephir M-C, Pelayo S, Bernonville S. Example of a Human Factors Engineering approach to a medication administration work system: Potential impact on patient safety. *Int J Med Inf* 2010;79(4):e43-e57
763. Ruiz JG, Qadri SS, Nader S, et al. Primary care management of chronic nonmalignant pain in veterans: a qualitative study. *EDUC GERONTOL* 2010;36(5):372-93.
764. Fernando S, Georgiou A, Holdgate A, et al. Challenges associated with electronic ordering in the emergency department: a study of doctors' experiences. *Emergency Medicine Australasia* 2009;21(5):373-8.
765. Ash JS, Gorman PN, Lavelle M, et al. A cross-site qualitative study of physician order entry. *J Am Med Inform Assoc* 2003;10(2):188-200.
766. Pegram AA, Kennedy LD, Powell BL. Use of erythropoietin alpha after introduction of a treatment algorithm in an oncology clinic. *Ashp Midyear Clinical Meeting* 2000;35:
767. Ash J, Gorman P, Lavelle M, et al. Multiple perspectives on physician order entry. *Proceedings of the AMIA Symposium* 2000;27-31.
768. Khovanov AV, Nechaev VI, Krylov VV, et al. An objective relation approach in the modeling of dispensary registration of patients with tuberculosis. *Problemy Tuberkuleza I Boleznej Legkih* 2007;tuberk.(4):10-3.
769. Ronco G, Giubilato P, Naldoni C, et al. Extension of organised cervical cancer screening programmes in Italy and their process indicators: 2007 activity. *Epidemiol Prev* 2009;33 Suppl 2:41-56.
770. Barber N, Cornford T, Klecun E. Qualitative evaluation of an electronic prescribing and administration system. *Qual Safe Health Care* 2007;16(4):271-8.

771. Holbrook A, Thabane L, Keshavjee K, et al. Individualized electronic decision support and reminders to improve diabetes care in the community: COMPETE II randomized trial. *CMAJ Canadian Medical Association Journal* 2009;181(1-2):37-44.
772. Field TS, Rochon P, Lee M, et al. Costs associated with developing and implementing a computerized clinical decision support system for medication dosing for patients with renal insufficiency in the long-term care setting. *J Am Med Inform Assoc* 2008;15(4):466-72.
773. Brock TP, Smith SR. Using digital videos displayed on personal digital assistants (PDAs) to enhance patient education in clinical settings. *Int J Med Inf* 2007;76(11-12):829-35.
774. Santell JP, Kowiatek JG, Weber RJ, et al. Medication errors resulting from computer entry by nonprescribers. *Am J Health Syst Pharm* 2009;66(9):843-53.
775. Singh H, Mani S, Espadas D, et al. Prescription errors and outcomes related to inconsistent information transmitted through computerized order entry: a prospective study. *Arch Intern Med* 2009;169(10):982-9.
776. Ash JS, Sittig DF, Campbell EM, et al. Some unintended consequences of clinical decision support systems. *AMIA Proceedings* 2007;26-30.
777. Ash JS, Sittig DF, Poon EG, et al. The extent and importance of unintended consequences related to computerized provider order entry. *J Am Med Inform Assoc* 2007;14(4):415-23.
778. Ash JS, Sittig DF, Dykstra RH, et al. Categorizing the unintended sociotechnical consequences of computerized provider order entry. *Int J Med Inf* 2007;76(Suppl 1):S1-S7
779. Campbell EM, Sittig DF, Ash JS, et al. Types of unintended consequences related to computerized provider order entry. *J Am Med Inform Assoc* 2006;13(5):547-56.
780. Khong TK, Singer DR. Adverse drug reactions: current issues and strategies for prevention and management. *Expert Opin Pharmacother* 2002;3(9):1289-300.
781. Biffi M, Bertini M, Saporito D, et al. Automatic management of left ventricular stimulation: hints for technologic improvement. *Pacing Clin Electrophysiol* 2009;32(3):346-53.
782. Bernstam EV, Hersh WR, Sim I, et al. Unintended consequences of health information technology: a need for biomedical informatics. *J Biomed Inform* 2010;43(5):828-30.
783. Talmon J, Ammenwerth E, Brender J, et al. STARE-HI - Statement on reporting of evaluation studies in Health Informatics. *Int J Med Inform* 2009;78(1):1-9.
784. Vogel L. Finding Value from IT Investments: Exploring the Elusive ROI in Healthcare. *J Healthc Inf Manag* 2003;17(4):20-8.
785. Menachemi N, Brooks RG. Reviewing the benefits and costs of electronic health records and associated patient safety technologies. *J Med Syst* 2006;30(3):159-68.
786. Johnston D, Pan E, Walker J et al. Patient safety in the physician's office: Assessing the value of ambulatory CPOE. California HealthCare Foundation; 2004. <http://www.chcf.org/documents/ihealth/PatientSafetyInPhysiciansOfficeACPOE.pdf>
787. Ash JS, Stavri PZ, Kuperman GJ. A consensus statement on considerations for a successful CPOE implementation. *J Am Med Inform Assoc* 2003;10(3):229-34.
788. Subramanian S, Hoover S, Gilman B, et al. Computerized physician order entry with clinical decision support in long-term care facilities: costs and benefits to stakeholders. *J Am Geriatr Soc* 2007;55(9):1451-7.
789. Jha AK, DesRoches CM, Campbell EG, et al. Use of electronic health records in U.S. hospitals. *N Engl J Med* 2009;360(16):1628-38.
790. Robinson JC, Casalino LP, Gillies RR, et al. Financial incentives, quality improvement programs, and the adoption of clinical information technology. *Med Care* 2009;47(4):411-7.
791. Menachemi N, Brooks RG, Schwalenstocker E, et al. Use of health information technology by children's hospitals in the United States. *Pediatrics* 2009;123:80-4.

792. Menachemi N, Ford EW, Beitsch LM, et al. Incomplete EHR adoption: late uptake of patient safety and cost control functions. *Am J Med Qual* 2007;22(5):319-26.
793. Inquilla CC, Szeinbach S, Seoane-Vazquez E, et al. Pharmacists' perceptions of computerized prescriber-order-entry systems. *Am J Health Syst Pharm* 2007;64(15):1626-32.
794. Simon SR, Kaushal R, Cleary PD, et al. Physicians and electronic health records: a statewide survey. *Arch Intern Med* 2007;167(5):507-12.
795. Furukawa MF, Ketcham JD, Rimsza ME. Physician practice revenues and use of information technology in patient care. *Med Care* 2007;45(2):168-76.
796. Grossman JM, Reed MC. Clinical information technology gaps persist among physicians. *Issue Brief/Center for Studying Health System Change* 2006;(106):1-4.
797. Protti D, Wright G, Treweek S, et al. Primary care computing in England and Scotland: a comparison with Denmark. *Inform Prim Care* 2006;14(2):93-9.
798. Wang CJ, Marken RS, Meili RC, et al. Functional characteristics of commercial ambulatory electronic prescribing systems: a field study. *J Am Med Inform Assoc* 2005;12(3):346-56.
799. Menachemi N, Perkins R, van Durme D, et al. Examining the adoption of electronic health records and personal digital assistants by family physicians in Florida. *Inform Prim Care* 2006;14(1):1-9.
800. 2008 HIMSS/HIMSS analytics ambulatory healthcare IT survey. *Healthcare Information and Management Systems Society*; 2008. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68533>
801. Bell DS, Schueth AJ, Guinan JP, et al. Evaluating the technical adequacy of electronic prescribing standards: Results of an expert panel process. *AMIA* 2008;46-50.
802. Osborn M, Day R, Westbrook J. Are specialist physicians missing out on the e-Health boat? *INTERN MED J* 2009;39(10):655-61.
803. Moxey A, Robertson J, Newby D, et al. Computerized clinical decision support for prescribing: provision does not guarantee uptake. [Review] [78 refs]. *J Am Med Inform Assoc* 2010;17(1):25-33.
804. Dorr D, Bonner LM, Cohen AN, et al. Informatics systems to promote improved care for chronic illness: A literature review. *J Am Med Inform Assoc* 2007;14(2):156-63.
805. Wang J. New diagnosis of hypertension among celecoxib and nonselective NSAID users: A population-based cohort study. *Ann Pharmacother* 1 A.D.;41(6):937-43.
806. Bell DS, Marken RS, Meili RC, et al. Recommendations for comparing electronic prescribing systems: results of an expert consensus process. *Health Aff (Millwood)* 2004;Suppl:Web-305
807. Bashyam V, Divita G, Bennett DB, et al. A normalized lexical lookup approach to identifying UMLS concepts in free text. *Stud Health Technol Inform* 2007;129(Pt:1):1-9.
808. Agency for Healthcare Research and Quality. Expert Panel Meeting: Health Information Technology <http://www.ahrq.gov/data/hitmeet.htm>
809. Blumenthal D TM. The "Meaningful Use" Regulation for Electronic Health Records. *N Engl J Med* 2010;363(6):501-4.
810. Ahmad A. Key attributes of a successful physician order entry system implementation in a multi-hospital environment. *J Am Med Inform Assoc* 2002;9(1):16-24.
811. Aarts J. Implementation of computerized physician order entry in seven countries. *Health Aff (Millwood)* 2009;28(2):404-14.
812. Teufel RJ, Kazley AS, Basco WT, Jr. Early adopters of computerized physician order entry in hospitals that care for children: a picture of US health care shortly after the Institute of Medicine reports on quality. *Clin Pediatr (Phila)* 2009;48(4):389-96.
813. Cutler DM, Feldman NE, Horwitz JR. U.S. adoption of computerized physician order entry systems. *Health Aff (Millwood)* 2005;24(6):1654-63.

814. Longo DR, Hewett JE, Ge B, et al. Rural hospital patient safety systems implementation in two States. *J Rural Health* 2007;23(3):189-97.
815. Simon JS, Rundall TG, Shortell SM. Adoption of order entry with decision support for chronic care by physician organizations. *J Am Med Inform Assoc* 2007;14(4):432-9.
816. Menachemi N, Ford EW, Chukmaitov A, et al. Managed care penetration and other factors affecting computerized physician order entry in the ambulatory setting. *Am J Manag Care* 2006;12(12):738-44.
817. Spil TAM, Schuring R, Michel-Verkerke M. Electronic prescription system: Do the professional use it? *International Journal of Healthcare Technology and Management* 2004;6(1):32-55.
818. KLAS. Clinical automation in the physician office. FOCUS: e-Prescribing/Ambulatory EMR 2005. *HEALTHC Q* 2005;8(3):115-8.
819. Fischer MA, Vogeli C, Stedman MR, et al. Uptake of electronic prescribing in community-based practices. *J Gen Intern Med* 2008;23(4):358-63.
820. Poon EG, Blumenthal D, Jaggi T, et al. Overcoming barriers to adopting and implementing computerized physician order entry systems in U.S. hospitals. *Health Aff (Millwood)* 2004;23(4):184-90.
821. Poon EG, Blumenthal D, Jaggi T, et al. Overcoming the barriers to the implementing computerized physician order entry systems in US hospitals: perspectives from senior management. *AMIA* 2003;975
822. Harrison M, Koppel R, Bar-Lev S. Unintended consequences of information technologies in health care: An interactive sociotechnical analysis. *J Am Med Inform Assoc* 2007;14:542-9.
823. Menachemi N, Burke D, Brooks RG. Adoption factors associated with patient safety-related information technology. *J Healthc Qual* 2004;26(6):39-44.
824. Furukawa MF, Raghu T, Spaulding T, et al. Adoption of health information technology for medication safety in U.S. hospitals, 2006. *Health Aff (Millwood)* 2008;27(3):865-75.
825. Pedersen CA, Schneider PJ, Scheckelhoff DJ. ASHP national survey of pharmacy practice in hospital settings: dispensing and administration--2005. *Am J Health Syst Pharm* 2006;63(4):327-45.
826. Pedersen CA, Gumpfer KF. ASHP national survey on informatics: assessment of the adoption and use of pharmacy informatics in U.S. hospitals--2007. *Am J Health Syst Pharm* 2008;65(23):2244-64.
827. Pedersen CA, Schneider PJ, Scheckelhoff DJ. ASHP national survey of pharmacy practice in hospital settings: monitoring and patient education--2006. *Am J Health Syst Pharm* 2007;64(5):507-20.
828. Centre for Medicare and Medicaid Services. Part 423: Voluntary MEDICARE Prescription Drug Benefit Subpart D-Cost Control and Quality Improvement Requirements . *Electronic Code of Federal Regulations*.
<http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?c=ecfr&sid=956b1d55986299e9a20b885672924c9d&rgn=div8&view=text&node=42:3.0.1.1.10.4.6.5&idno=42>.
829. Ghinea G, Asgari S, Moradi A, et al. A Jini-based solution for electronic prescriptions. *IEEE Transactions on Information Technology in Biomedicine* 2006;10(4):794-802.
830. Craghead RM, Wartski DM. Effect of automated prescription transmittal on number of unclaimed prescriptions. *Am J Hosp Pharm* 1989;46(2):310-2.
831. Sugden B, Wilson R. Integrated care and electronic transmission of prescriptions: Experience of the evaluation of ETP pilots. *Health Informatics Journal* 2004;10(4):277-90.
832. Phul S, Cooper SL, Cantrill JA. Pharmacy services and patient choice: Insights into differences between patient groups. *Int J Pharm Pract* 2003;11(4):233-42.
833. Long term care: e-Prescribing standards pilot study. Final report. 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_227304_0_0_18/Long%20Term%20Care%20e-Prescribing%20Standards%20Pilot%20Study%20-%20Final%20Report.pdf

834. Electronic prescribing using a community utility: The eprescribing gateway. AHRQ; 2007.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_227306_0_0_18/Electronic%20Prescribing%20Using%20A%20Community%20Utility%20-%20The%20ePrescribing%20Gateway.pdf
835. Centers for Medicare & Medicaid services. Special Study: Pilot testing of electronic prescribing standards. 2007.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_227308_0_0_18/Ohio%20KePRO%20-%20UHMP%20-%20Final%20Report.pdf
836. Maximizing the effectiveness of e-prescribing between physicians and community pharmacies. Surescripts, LLC; 2006.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_227310_0_0_18/SureScripts%20-%20Final%20Report.pdf
837. Leavitt M. Pilot testing of initial electronic prescribing standards: Cooperative agreements. U.S. Department of Health and Human Services; 2003.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_227312_0_0_18/eRxReport_041607.pdf
838. Bell D, Schueth A, Crosson J et al. Pilot testing of electronic prescribing standards. AHRQ; 2006.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_227350_0_0_18/NJ-EPAC%20Pilot%20Testing%20of%20Electronic%20Prescribing%20Standards.pdf
839. Findings from the evaluation of e-prescribing pilot sites. AHRQ; 2007.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_227460_0_0_18/Findings%20From%20The%20Evaluation%20of%20OE-Prescribing%20Pilot%20Sites.pdf
840. Al Tajir GK, Kelly WN. Epidemiology, comparative methods of detection, and preventability of adverse drug events. *Ann Pharmacother* 2005;39(7-8):1169-74.
841. Smith AD. Barriers to accepting e-prescribing in the U.S.A. *Int J Health Care Qual Assur Inc Leadersh Health Serv* 2006;19(2-3):158-80.
842. Halamka J, Aranow M, Ascenzo C, et al. E-Prescribing collaboration in Massachusetts: early experiences from regional prescribing projects. *J Am Med Inform Assoc* 2006;13(3):239-44.
843. Davidson EJ, Chismar WG. Planning and managing computerized order entry: a case study of IT-enabled organizational transformation. *Top Health Inf Manage* 1999;19(4):47-61.
844. Ax F, Ekedahl A. Electronically transmitted prescriptions not picked up at pharmacies in Sweden. *Res Social Adm Pharm* 2010;6(1):70-7.
845. Kaphingst KA, Rudd RE, DeJong W, et al. Literacy demands of product information intended to supplement television direct-to-consumer prescription drug advertisements. *Patient Educ Couns* 2004;55(2):293-300.
846. Shcherbatykh, I, Holbrook, A., Thabane, L., et al. Methodologic Issues in Health Informatics Trials: The Complexities of Complex Interventions. *Journal of the American Medical Informatics Association*.
847. Fortescue EB, Kaushal R, Landrigan CP, et al. Prioritizing strategies for preventing medication errors and adverse drug events in pediatric inpatients. *Pediatrics* 2003;111(4:Pt:1):t-9
848. Gurwitz JH, Field TS, Harrold LR, et al. Incidence and preventability of adverse drug events among older persons in the ambulatory setting. *JAMA* 2003;289(9):1107-16.
849. Gurwitz JH, Field TS, Judge J. The incidence of adverse drug events in two large academic long-term care facilities. *Am J Med* 2005;118(3):251-8.
850. Schiff GD, Klass D, Peterson J, et al. Linking laboratory and pharmacy: opportunities for reducing errors and improving care. *Arch Intern Med* 2003;163(8):893-900.

Acronyms and Abbreviations

| Abbreviation | Words/Phrases |
|---------------------|---|
| ADE | Adverse Drug Event |
| AHRQ | Agency for Healthcare Research and Quality |
| AMIA | American Medical Informatics Association |
| ASHP | American Society of Health-System Pharmacists |
| BCM | Bar Code Medication |
| BCMA | Bar Code Medication Administration |
| CA | Cost Analysis |
| CBA | Cost Benefit Analysis |
| CCA | Cost Consequence Analysis |
| CCHIT | Commission for Healthcare Information Technology |
| CDSS | Clinical Decision Support System |
| CEA | Cost Effectiveness Analysis |
| CI | Confidence Interval |
| CINAHL | Cumulated Index to Nursing and Allied Health Literature |
| CITL | Center for Information Technology Leadership |
| CPOE | Computerized Provider Order Entry |
| CUA | Cost Utility Analysis |
| EDI | Electronic Data Interchange |
| EHR | Electronic Health Record |
| E-LIS | Eprints in Library and Information Science |
| e-MAR | Electronic Medication Administration Record |
| EMBASE | Excerpta Medica Database |
| EMR | Electronic Medical Records |
| e-TAR | Electronic Treatment Authorization Request |
| Health IT | Health Information Technology |
| HIV | Human Immunodeficiency Virus |
| HMO | Health Maintenance Organization |
| HTA | Health Technology Assessment |
| ICU | Intensive Care Unit |
| IEEE | Institute of Electrical and Electronics Engineers |
| IOM | Institute of Medicine |
| IPA | International Pharmaceutical Abstracts |
| IT | Information Technology |
| JAMIA | Journal of the American Medical Informatics Association |
| MEDLINE | Medical Literature Analysis and Retrieval System |
| MMIT | Medication Management Information Technology |
| MMS | Medication Management System |
| NICU | Neonatal Intensive Care Unit |
| PDA | Personal Digital Assistants |
| PHR | Personal Health Record |
| RCT | Randomized Controlled Trial |

Appendix A. Exact Search Strings

| |
|---|
| MEDLINE® Ovid MEDLINE(R) <1950 to September Week 2 2009> |
| Date searched: Sept 21-09 |
| Number of retrievals: 10767 |
| 1 electronic prescribing/ (61) 2 drug therapy, computer assisted/ (1151) 3 (electronic adj3 prescri*).mp. (351) 4 electronic medication*.mp. (116) 5 automated prescri*.mp. (25) 6 (automated adj3 medication*).mp. (72) 7 (online adj3 prescri*).mp. (39) 8 (online adj3 medication*).mp. (17) 9 e-prescri*.mp. (163) 10 eprescri*.mp. (12) 11 e-medication*.mp. (4) 12 emar*.mp. (169) 13 (bcma and (medication* or prescri* or drug)).mp. (24) 14 e-rx.mp. (11) 15 ((bar cod* or barcod*) and (prescri* or medication* or drug*)).mp. (280) 16 (computer* adj2 prescri*).mp. (310) 17 prescri* monitor*.mp. (89) 18 clinical pharmacy information systems/ (986) 19 prescri* order entry.mp. (56) 20 pharma* order entry.mp. (5) 21 computer* order entry.mp. (115) 22 automated dispens*.mp. (82) 23 or/1-22 (3403) 24 exp pharmaceutical services/ (38883) 25 exp medical errors/ (64473) 26 exp drug therapy/ (864020) 27 exp drug interactions/ (122756) 28 exp drug monitoring/ (9813) 29 exp medication systems/ (3386) 30 exp drug administration schedule/ (71159) 31 exp drug costs/ (9397) 32 exp dose-response relationship, drug/ (288539) 33 drug therapy, computer assisted/ (1151) 34 (prescri* or medication*).mp. (218668) 35 pharmacotherap*.mp. (14545) 36 pharmaceutical*.mp. (115354) 37 dispens*.mp. (18455) 38 exp therapeutic uses/ (3544181) 39 (safety or safe).mp. (310321) 40 error*.mp. (173803) 41 (adverse adj3 event*).mp. (46457) 42 (adverse adj3 effect*).mp. (79271) 43 mistake*.mp. (11308) 44 complication*.mp. (744423) 45 (risk adj5 manag*).mp. (21572) 46 (risk adj5 assess*).mp. (138608) 47 harm*.mp. (57502) 48 exp medical errors/ (64473) 49 safety management/ (11037) 50 patient safety/ (0) 51 medical error/ (8433) 52 medication error/ (7690) 53 risk management/ (11711) 54 risk assessment/ (109320) |

55 adverse drug reaction reporting systems/ (4027)
 56 or/24-55 (5187543)
 57 cdss.tw. (355)
 58 ccdss.tw. (2)
 59 (comput* adj3 decision support*).mp. (603)
 60 reminder system*.tw. (380)
 61 decision support systems, clinical/ (3072)
 62 reminder systems/ (1486)
 63 therapy, computer assisted/ (3599)
 64 decision making, computer assisted/ (2051)
 65 (comput* adj3 order entry).tw. (714)
 66 provider order entry.tw. (196)
 67 cpoe.tw. (492)
 68 clinician order entry.tw. (4)
 69 physician order entry.tw. (443)
 70 nurs* order entry.tw. (2)
 71 pharma* order entry.tw. (5)
 72 medical order entry systems/ (799)
 73 patient portal*.mp. (67)
 74 personal medical record*.mp. (42)
 75 personal health record*.mp. (215)
 76 (patient adj2 access* adj2 record*).mp. (728)
 77 (patient adj2 carried adj2 record*).mp. (3)
 78 (patient adj2 held adj2 record*).mp. (52)
 79 (patient adj2 shared adj2 record*).mp. (14)
 80 patient internet portal*.mp. (9)
 81 phr.mp. (484)
 82 ephr.mp. (11)
 83 exp medical records/ and patient access to record*.mp. (478)
 84 kiosk*.tw. (105)
 85 point-of-care systems/ (4135)
 86 computers, handheld/ (1396)
 87 Medical Records Systems, Computerized/ (15799)
 88 or/57-87 (31551)
 89 56 and 88 (7973)
 90 guideline adherence/ (12072)
 91 exp patient compliance/ (38269)
 92 (patient compliance or patient adherence).tw. (5892)
 93 (comput* or online or internet or electron*).mp. (1263766)
 94 or/90-92 (53419)
 95 94 and 93 (3358)
 96 56 and 95 (1775)
 97 23 or 89 or 96 (11560)
 98 97 not letter.pt. (11306)
 99 98 not editorial.pt. (11144)
 100 99 not news.pt. (11001)
 101 animal/ not (human/ and animal/) (3351990)
 102 100 not 101 (10767)

EMBASE Ovid EMBASE <1980 to 2009 Week 38>

Date searched: Sept 23-09

Number of retrievals: 8693

- 1 electronic prescribing/ (8)
- 2 computer assisted drug therapy/ (168)
- 3 electronic prescri*.mp. (256)
- 4 electronic medication*.mp. (101)
- 5 automated prescri*.mp. (22)
- 6 automated medication*.mp. (25)
- 7 online prescri*.mp. (12)
- 8 online medication*.mp. (6)

9 e-prescri*.mp. (105)
 10 eprescri*.mp. (5)
 11 e-medication*.mp. (2)
 12 emar.mp. (11)
 13 (bcma and (medication* or prescri* or drug*)).mp. (90)
 14 e-rx.mp. (6)
 15 ((bar cod* or barcod*) and (prescri* or medication*)).mp. (141)
 16 computer* prescri*.mp. (182)
 17 prescription monitor*.mp. (64)
 18 electronic medication administration record.sh. (2)
 19 computer assisted drug therapy/ (168)
 20 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 (1096)
 21 exp medical errors/ (31362)
 22 exp drug therapy/ (905532)
 23 exp drug interactions/ (141655)
 24 exp drug monitoring/ (28295)
 25 exp medication systems/ (4626)
 26 exp drug administration schedule/ (791038)
 27 exp drug costs/ (36734)
 28 exp dose-response relationship, drug/ (230877)
 29 drug therapy, computer assisted/ (168)
 30 (prescri* or medication*).mp. (209602)
 31 pharmacotherap*.mp. (15893)
 32 pharmaceutical*.mp. (86405)
 33 dispens*.mp. (13409)
 34 pharmacy/ (23073)
 35 (safety or safe).mp. (358398)
 36 error*.mp. (213771)
 37 (adverse adj3 event*).mp. (46207)
 38 (adverse adj3 effect*).mp. (74072)
 39 mistake*.mp. (8656)
 40 complication*.mp. (439160)
 41 (risk adj5 manag*).mp. (19630)
 42 (risk adj5 assess*).mp. (204644)
 43 harm*.mp. (48109)
 44 exp medication error/ (3076)
 45 drug safety/ (140603)
 46 patient safety/ (14509)
 47 risk assessment/ (186991)
 48 drug surveillance program/ (7404)
 49 drug monitoring/ (28295)
 50 or/21-49 (2850800)
 51 cdss.tw. (285)
 52 ccdss.tw. (0)
 53 (comput: adj3 decision support*).mp. (972)
 54 reminder system*.tw. (239)
 55 decision support system/ (1700)
 56 reminder systems/ (166)
 57 computer assisted drug therapy/ (168)
 58 cpoe.tw. (228)
 59 (comput* adj3 order entry).tw. (443)
 60 provider order entry.tw. (88)
 61 clinician order entry.tw. (1)
 62 physician order entry.tw. (265)
 63 nurs* order entry.tw. (1)
 64 pharma* order entry.tw. (6)
 65 hospital information system/ (1335)
 66 medical information system/ (7562)
 67 patient portal*.mp. (30)
 68 personal medical record*.mp. (27)
 69 personal health record*.mp. (109)
 70 (patient adj2 access* adj2 record*).mp. (51)

71 (patient adj2 carried adj2 record*).mp. (2)
 72 (patient adj2 held adj2 record*).mp. (48)
 73 (patient adj2 shared adj2 record*).mp. (14)
 74 patient internet portal*.mp. (4)
 75 electronic medical record/ and patient access.tw. (12)
 76 kiosk*.tw. (69)
 77 microcomputer/ (5437)
 78 electronic medical record/ (4542)
 79 or/51-78 (20325)
 80 50 and 79 (6242)
 81 (guideline adherence or adherence to guideline*).tw. (639)
 82 patient compliance/ (49247)
 83 (patient compliance or patient adherence).tw. (5413)
 84 or/81-83 (51850)
 85 (comput* or online or internet or electron*).mp. (939381)
 86 84 and 85 (2902)
 87 20 or 80 or 86 (9504)
 88 87 not letter.pt. (9125)
 89 88 not editorial.pt. (8694)
 90 89 not news.pt. (8694)
 91 animal/ not (human/ and animal/) (14494)
 92 90 not 91 (8693)

CINAHL via EBSCOhost

Date searched: Sept 25-09

Number of retrievals: 4692

S97 (S96 not PT(editorial))
 S96 (S24 or S88 or S94) not PT(letter)
 S95 (S24 or S88 or S94)
 S94 (S92 and S93)
 S93 (S89 or S90 or S91)
 S92 (comput* or online or internet or electron*)
 S91 (patient compliance or patient adherence)
 S90 MH(patient compliance+)
 S89 MH(guideline adherence)
 S88 (S59 and S87)
 S87 (S60 or S61 or S62 or S63 or S64 or S65 or S66 or S67 or S68 or S69 or S70 or S71 or S72 or S73 or S74 or S75 or S76 or S77 or S78 or S79 or S80 or S81 or S82 or S83 or S84 or S85 or S86)
 S86 MH (risk assessment)
 S85 MH (risk management)
 S84 MH (patient safety+)
 S83 (safety manage*)
 S82 (harm*)
 S81 (risk N5 assess*)
 S80 (risk N5 manage*)
 S79 (complication*)
 S78 (mistake*)
 S77 (adverse N3 effect*)
 S76 (adverse N3 event*)
 S75 (error*)
 S74 (safety or safe)
 S73 (MH "Therapeutics")
 S72 (dispense*)
 S71 (pharmaceutical*)
 S70 (pharmacotherap*)
 S69 (prescri* or medication*)
 S68 (drug cost*)
 S67 MH "drug therapy, computer assisted"
 S66 MH "dose-response relationship, drug"
 S65 MH "drug administration schedule"

S64 MH "medication systems"
 S63 MH "drug monitoring"
 S62 MH "drug interactions+"
 S61 MH "drug therapy+"
 S60 MH "treatment errors"
 S59 (S25 or S26 or S27 or S28 or S29 or S30 or S31 or S32 or S33 or S34 or S35 or S36 or S37 or S38 or S39 or S40 or S41 or S42 or S43 or S44 or S45 or S46 or S47 or S48 or S49 or S50 or S51 or S52 or S53 or S54 or S55 or S56 or S57 or S58)
 S58 (MH "patient record systems")
 S57 (MH "Computers, Hand-Held")
 S56 (point-of-care) or (point of care)
 S55 (kiosk)
 S54 MH(medical records+) and (patient access*)
 S53 MH(computerized patient record)
 S52 MH(medical records, personal)
 S51 "(ephr)"
 S50 "(phr)"
 S49 "(Patient internet portal)"
 S48 (MH "Patient Access to Records")
 S47 (patient N2 shared N2 record*)
 S46 (patient N2 held N2 record*)
 S45 (patient N2 carried N2 record*)
 S44 (patient N2 access* N2 record*)
 S43 (personal health record*)
 S42 (personal medical record*)
 S41 (patient portal*)
 S40 MH(electronic order entry)
 S39 (pharma* order entry)
 S38 (nurs* order entry)
 S37 (physician order entry)
 S36 (clinician order entry)
 S35 (cpoe)
 S34 (provider order entry)
 S33 (comput* N3 order entry)
 S32 MH(decision making, computer assisted)
 S31 MH(therapy, computer assisted)
 S30 MH(reminder systems)
 S29 MH(decision support systems, clinical)
 S28 (reminder system*)
 S27 (comput* N3 decision support*)
 S26 (ccdss)
 S25 (cdss)
 S24 (S1 or S2 or S3 or S4 or S5 or S6 or S7 or S8 or S9 or S10 or S11 or S12 or S13 or S14 or S15 or S16 or S17 or S18 or S19 or S20 or S21 or S22 or S23)
 S23 (automated dispens*)
 S22 (MH "Electronic Order Entry")
 S21 (computer* order entry)
 S20 (pharma* order entry)
 S19 (prescri* order entry)
 S18 (prescri* monitor*)
 S17 (computer* N2 prescri*)
 S16 ((bar cod* or barcod*) and (prescri* or medication* or drug*))
 S15 (e-rx)
 S14 (bcma)
 S13 (emar*)
 S12 (emedication*)
 S11 (emedication*)
 S10 (eprescri*)
 S9 (e-prescri*)
 S8 (online N3 medication*)
 S7 (online N3 prescri*)
 S6 (automated N3 medication*)

S5 automated prescri*
 S4 (electronic medication*)
 S3 (electronic N3 prescri*)
 S2 MH ("drug therapy, computer assisted")
 S1 MH ("clinical pharmacy information systems")

Cochrane Library

Date searched: Sept 29-09

Number of retrievals: 148

#1 MeSH descriptor Drug Therapy, Computer-Assisted, this term only

#2 MeSH descriptor Electronic Prescribing, this term only

#3 electronic prescri* OR "electronic medication*" OR "automated prescri*" OR "automated medication*" OR "online prescri*" OR "online medication*" OR "e-prescri*" OR "eprescri*" OR "e-medication*" OR "emar" OR (bcma and (medication* or prescri* or drug*)) OR "e-rx" OR ((bar cod* or barcod*) and (prescri* or medication)) OR "computer* prescri*" OR "prescription monitor*" :ti,ab,kw

#4 (#1 OR #2 OR #3)

IPA Abstracts

Date searched: Sept 21-09

Number of retrievals: 4387

(((((patient and compliance) or (patient and adherence) or (guideline and adherence) or (guideline and compliance)) and (comput* or online or internet or electron*)) or (computer assisted drug therapy or (electronic adj3 prescri*) or (electronic adj3 medication*) or (automated adj3 prescri*) or (automated adj3 medication*) or (online adj3 prescri*) or (online adj3 medication*) or e-prescri* or eprescri* or e-medication* or emar or bcma or e-rx or (bar cod* or barcod*) or (computer* adj2 prescrib*) or prescri* monitor* or information systems* or automated dispens* or cdss or (comput* adj3 decision support*) or (reminder system* or (clinic* adj3 decision support*) or (therapy and computer assist*) or (decision making and computer) or (comput* adj3 order entry) or (provider adj3 order entry) or cpoe or (clinician adj3 order entry) or (physician adj3 order entry) or (nurs* adj3 order entry) or (pharma* adj3 order entry) or (prescri* adj3 order entry) or order entry systems or patient portal* or (personal adj2 record*) or (patient adj2 access* adj2 record*) or (patient adj2 carried adj2 record*) or (patient adj2 held adj2 record*) or (patient adj2 shared adj2 record*) or patient internet portal*) or (phr or ephr or kiosk* or point of care or point-of-care or handheld)).af. (4387)

Compendex AND Inspec via Engineering Village

Date searched: Sept 28-09

Number of retrievals: 1503

(((((medication*)WN ALL)) NOT (({461.2} OR {804.1} OR {a8770e} OR {921} OR {a8730c} OR {716} OR {804} OR {922.2} OR {b7510d} OR {801} OR {802.3} OR {741.1} OR {461.5} OR {921.6} OR {801.2} OR {a8745h} OR {804.2} OR {701.1} OR {803} OR {931.2} OR {717}) WN CL)) NOT (({462} OR {723.4} OR {731.1} OR {c3385} OR {462.1} OR {b7510} OR {a8770} OR {c1140z} OR {a8730} OR {a8745d} OR {a8725} OR {622.3}) WN CL))

LISTA (1974-2009) via EBSCOhost

Date searched: Sept 25-09

Number of retrievals: 276

S3 (S2 and S3)

S2 (comput* or online or internet or electron*)

S1 medication* or prescri*

PsycINFO <1967 to September Week 3 2009>

Date searched: Sept 23-09

Number of retrievals: 3074

1 computer assisted therapy/ (177)

2 (electronic adj3 prescri*).mp. (16)

3 electronic medication*.mp. (16)
 4 automated prescri*.mp. (1)
 5 (automated adj3 medication*).mp. (2)
 6 (online adj3 prescri*).mp. (10)
 7 (online adj3 medication*).mp. (3)
 8 e-prescri*.mp. (3)
 9 eprescri*.mp. (0)
 10 e-medication*.mp. (4)
 11 emar*.mp. (16)
 12 (bcma and (medication* or prescri* or drug)).mp. (1)
 13 e-rx.mp. (1)
 14 ((bar cod* or barcod*) and (prescri* or medication* or drug*)).mp. (6)
 15 (computer* adj2 prescri*).mp. (12)
 16 prescri* monitor*.mp. (24)
 17 prescri* order entry.mp. (1)
 18 pharma* order entry.mp. (0)
 19 computer* order entry.mp. (3)
 20 automated dispens*.mp. (3)
 21 or/1-20 (292)
 22 pharmaceutical services.mp. (15)
 23 medical error*.mp. (231)
 24 exp drug therapy/ (82011)
 25 exp drug interactions/ (5922)
 26 drug monitor*.mp. (250)
 27 self monitoring/ (1987)
 28 medication system*.mp. (3)
 29 exp drug administration methods/ (6226)
 30 drug administration schedule.mp. (8)
 31 drug cost*.mp. (177)
 32 exp "Costs and Cost Analysis"/ (12017)
 33 (dose-response or dose response).mp. (3282)
 34 computer assisted therapy/ (177)
 35 exp "Prescribing (Drugs)"/ (2025)
 36 prescription drugs/ (1311)
 37 (prescri* or medication*).mp. (57547)
 38 pharmacotherap*.mp. (7804)
 39 pharmaceutical*.mp. (3033)
 40 dispens*.mp. (1442)
 41 therapeutic uses.mp. (259)
 42 (safety or safe).mp. (33813)
 43 error*.mp. (50722)
 44 (adverse adj3 event*).mp. (4502)
 45 (adverse adj3 effect*).mp. (8104)
 46 mistake*.mp. (5096)
 47 complication*.mp. (10447)
 48 (risk adj5 manag*).mp. (4103)
 49 (risk adj5 assess*).mp. (11152)
 50 harm*.mp. (22923)
 51 exp "quality of care"/ (5241)
 52 drug interactions/ (5922)
 53 "side effects (drug)"/ (16075)
 54 medication error*.mp. (153)
 55 risk management/ (2007)
 56 risk assessment/ (4872)
 57 (adverse adj3 drug adj3 reaction).mp. (110)
 58 client treatment matching/ (763)
 59 treatment planning/ (2623)
 60 or/22-59 (272739)
 61 cdss.tw. (75)
 62 ccdss.tw. (3)
 63 (comput* adj3 decision support*).mp. (126)
 64 reminder system*.tw. (49)

- 65 (comput* adj3 order entry).tw. (23)
- 66 provider order entry.tw. (7)
- 67 cpoe.tw. (8)
- 68 clinician order entry.tw. (0)
- 69 physician order entry.tw. (15)
- 70 nurs* order entry.tw. (0)
- 71 pharma* order entry.tw. (0)
- 72 patient portal*.mp. (6)
- 73 personal medical record*.mp. (3)
- 74 personal health record*.mp. (20)
- 75 (patient adj2 access* adj2 record*).mp. (18)
- 76 (patient adj2 carried adj2 record*).mp. (0)
- 77 (patient adj2 held adj2 record*).mp. (16)
- 78 (patient adj2 shared adj2 record*).mp. (2)
- 79 patient internet portal*.mp. (1)
- 80 phr.mp. (36)
- 81 ephr.mp. (0)
- 82 exp medical records/ and patient access to record*.mp. (4)
- 83 kiosk*.tw. (55)
- 84 exp expert systems/ (3289)
- 85 prompting/ (229)
- 86 computer assisted therapy/ (177)
- 87 intelligent agents/ (185)
- 88 information technology/ (1944)
- 89 computer applications/ (9604)
- 90 automated information processing/ (952)
- 91 exp information systems/ (14268)
- 92 microcomputers/ (1111)
- 93 computer peripheral devices/ (329)
- 94 or/61-93 (29083)
- 95 60 and 94 (2751)
- 96 (patient compliance or patient adherence).tw. (889)
- 97 treatment compliance/ (7750)
- 98 treatment guidelines/ and (complian* or adheren*).mp. [mp=title, abstract, heading word, table of contents, key concepts] (219)
- 99 (comput* or online or internet or electron*).mp. (109841)
- 100 or/96-98 (8282)
- 101 99 and 100 (323)
- 102 21 or 95 or 101 (3123)
- 103 102 not letter.dt. (3112)
- 104 103 not editorial.dt. (3074)

| |
|---|
| Sociological Abstracts via Scholar's portal |
| Date searched: Sept 28-09 |
| Number of retrievals: 489 |
| (KW=(comput* or electron* or online*) or KW=(internet or (information within 3 system*) or automat*) or KW=technolog*) And ((((KW=medic*) or(KW=prescri*) or(KW=pharma*) or(KW=drug*)) and((KW=monitor*) or(KW=administr*) or(KW=adhere*) or(KW=comply*) or(KW=complian*) or(KW=dispens*))) or((KW=(drug within 1 therap*)) or(KW=(drug within 1 safe*)) or(KW=(medical within 2 error*)) or(KW=(medication within 2 error*)) or(KW=(patient within 2 safe*)) or(KW=(order entry)) or(KW=(decision within 2 support)) or(KW=(adverse within 2 event*)) or(KW=(adverse within 2 effect*))) |

| |
|---|
| Business Source Complete via EBSCOhost |
| Date searched: Sept 29-09 |
| Number of retrievals: 1055 |
| S9 (S8 or S7) |

S8 (S1 and S6) Limiters - Publication Type: Academic Journal, Book, Primary Source Document
 Narrow by Subject0: - MEDICAL technology
 S7 (S1 and S6) Limiters - Publication Type: Academic Journal, Book, Primary Source Document; Narrow by
 Subject0: - MEDICAL care
 S6 (S4 or S5)
 S5 (drug N1 therap*) or (drug N1 safe*) or (medical N2 error*) or (medication N2 error*) or (patient N2 safe*) or
 (order entry) or (decision N2 support) or (adverse N2 event*) or (adverse N2 effect*)
 S4 (S2 and S3)
 S3 (medicin* or medic* or drug* or pharma*)
 S2 (monitor* or administ* or adhere* or comply* or complian* or dispens* or prescri*)
 S1 (comput* or electron* or online* or internet or (information N2 system*) or automat* or technolog*

| Grey Literature Source Date searched | Search Terms | Retrieved | Reviewed |
|---|---|---------------|---|
| New York Academy of Medicine 2-Nov-09 | "medication management " returned 1716 results.-first 40 reviewed | 1716 | 40 |
| | "medication management information " returned 156 results.-first 40 reviewed | 156 | 40 |
| | "kw, wrdl: technology and kw, wrdl: drug medication " returned 36 results.-all reviewed | 36 | 36 |
| | and kw, wrdl: medication drug " returned 3 results. | 3 | 3 |
| | su, wrdl: medical informatics returned 156 results -first 40 reviewed | 156 | 40 |
| | "computer and drug " returned 29 results. | 29 | 29 |
| | "computer and medication " returned 451 results- reviewed first 40 | 451 | 40 |
| SIGLE 25-Nov-09 | ((medication)) or (("medication management")) AND ((computer)) or ((informatics)) | 317 | all |
| US HHS Health Information Technology 27-Nov-09 | none- searched 'reports' page and selected reports possibly on topic | | all |
| Health Technology Assessment reports CRD 24-Nov-09 | ((medicat* or "medication management")) AND ((comput* or informatic*)) | 393 | all |
| ProQuest Dissertations 25-Nov-09 | ((medicat* or "medication management")) AND ((comput* or informatic*)) | 264 | all |
| National Library for Health UK, includes Bandolier 25-Nov-09 | (medicat* or "medication management") AND (comput* or informatic*). | 74 | all |
| ProceedingsFirst Nov-25-09 | (kw: medicat* and kw: medication w management) and (kw: comput* or kw: informatic*). | 16 | all |
| PapersFirst 25-Nov-09 | (kw: medicat* or kw: medication w management) and (kw: comput* or kw: informatic*). | 143 | all |
| National Technical Information Service 4-Nov-09 | ALL: medicine computer drug | 46 | 46 |
| Google 27-Nov-09 | medication management health information technology | about 651,000 | First 50 websites searched for relevant documents |
| AHRQ 30-Nov-09 | eRx, Bar Coding and CPOE knowledge libraries | 254 | 254 |

Appendix B. Sample Screening and Abstraction Forms

Defining Medication Management IT

To be clear on what kinds of applications we're including in MMIT, we've devised the following outline for MMIT applications:

Include:

Medication Management Health information technology systems/ programs where:

The computer/technology processes patient specific information in some way

AND

The information provided by the system is relevant to one of the following processes in medication management:

- Prescribing/ordering medications
- order communication
- dispensing
- administering (by health care provider or care giver)
- monitoring (patient adherence/compliance, adverse event surveillance)
- education (of patients, not pre-professional education)
- reconciliation

AND

Someone (patient, care giver, family, health care professionals) receives information in return that is/can be linked to patient-specific information and which is used in decision making

AND

The technology is part of or links to an information system

OR

The article is about transmission/order communication eg. Electronic Data Interchange (EDI)

AND

The article contains outcome data

Examples: CPOE, CDSS for prescribing, automated pharmacy systems coupled with CPOE

Exclude:

Health information technology systems/programs with:

The IT component is only web browsing of general health information databases

OR

The system acts as a conduit of information only (except transmission of prescriptions between Health Care Provider and Pharmacy)

OR

Systems where no feedback is provided (eg surveys)

OR

The system does not help with medication management decision-making or provide information about any of the 6 medication use/process steps (prescribe/transmit/dispense/administer/monitor/educate)

OR

Systems that make measurements but do not process the information

OR

Stand-alone devices that do not integrate with information systems

OR

HIT application is used only to extract data

Examples: pill bottles that track opening/closing, smart pumps not tied to other systems, studies using EHR for data collection only (eg. quality improvement tracking), ,

Inclusion/Exclusion Criteria

| Inclusion Criteria | Exclusion Criteria |
|---|---|
| Original or Review articles | Letters, editorials, news items |
| Relates to at least 1 step in the medication management process | Not related to the medication management process |
| Medication management assisted by Health IT | Health IT not involved in medication management |
| Computerized order entry, e-prescribing, computer decision support for medication management, barcode medication administration, medication reminders for patients and clinicians | HIT that is only used to extract data |
| Deals with a Medication Management System or an application that feeds into/out of a system | Devices that are stand-alone. They may administer drugs etc but are not tied to a medication management/information system. E.g. MEMS |
| Article Contains data | Article Does not contain data |
| Foreign language with data | Foreign language with no access to full-text articles |
| Articles on Electronic Data Interchange/electronic transmission between Health Care Providers and Pharmacies | Technologies that are passive, don't process information. eg. only take measurements, transmit data (except EDI) or administer drugs |
| Prescribed drugs or medications only | Over-the-counter drugs or prescribed devices |
| Medical, Dentistry, Nursing articles | Veterinary articles |
| PDA's or stand-alone devices that take personal patient data and provide decision support | |

Title and Abstract Screen Guide

This screening will establish if the articles are to continue to full-text screening (medication management and IT)

- 1) Is this article an original^a or review article^b?
 - a) If Yes, select and continue to Q2.
 - b) If No, select and submit
 - c) If Uncertain, select and continue to Q2.

- 2) Does the article relate to at least one step in the medication management process (prescribing/ordering, transmitting, order communication, administering, monitoring-patient or population, reconciliation or education)?
 - a) If Yes, select and continue to Q3.
 - b) If No, select and submit
 - c) If Uncertain, select and continue to Q3.
- 3) If you answered yes above, which step(s) in the medication management process are involved (select all that are relevant)?
 - a) Prescribing/ordering
 - b) Transmitting
 - c) Order communication (verification, transformation, and communication (perfecting))
 - d) Administering
 - e) Monitoring (patient or population)
 - f) Medication reconciliation
 - g) Patient education
- 4) Is a medication management process assisted by Health IT?
 - a) If Yes, select and continue to Q4.
 - b) If No, select and submit
 - c) If Uncertain, select and continue to Q4.
- 5) Is this article relevant for background information (can be included or excluded articles)?
 - a) Yes
 - b) No
 - c) Uncertain

Submit to move to the next article. If you have left a question unanswered and have not selected no or uncertain, the system will display an error and you will need to check for any unanswered questions.

^aAn original study is any full text article in which investigators report first-hand observations-quantitative or qualitative, except for case reports

^bA review article is any full text article that indicates the intent is to review, summarize, highlight (or similar terms) the literature on a particular topic. This intent may be explicitly stated in the text of the article or it may be bannered as a review, overview or meta-analysis in the title or in a section heading. This may not be clear from the abstract.

(case reports, general and miscellaneous articles (no stated purpose, no methods, not bannered review anywhere), secondary publications and abstracts only are not considered original or review)

Full-Text Screening Guide

- 1) Does this article contain data (quantitative or qualitative; simulated or real patient)? Look for tables and graphs OR THE WORD QUALITATIVE. (qualitative research ‘results’ are text-based so there will be no numbers, BUT they will describe their methods as qualitative...they actually talk to people.
 - a) Yes, continue
 - b) No, exclude and go to next article
 - c) Uncertain

- 2) Is the MM information technology integrated into an IT system? (If it is a stand-alone device or software that is not hooked into a MM system, then exclude). Integrated systems include EHRs, CPOE, CDSS, etc.
 - a) Yes, continue
 - b) No, exclude and go to next article
 - c) Uncertain, continue

- 3) Which medication management phases are reported on? (Mandatory to answer this question, and can choose more than one).

If it's about patients taking drugs or someone deciding what/how much or =when to take a drug, then include it. If it is about using a system to make decisions about drugs then include it. Remember that MM includes the decision support in making prescribing decisions all the way through to medication reconciliation, post-marketing surveillance and patient monitoring/education.

 - a) Medication management in general
 - b) Prescribing/ordering
 - c) Transmission, order communication
 - d) Dispensing
 - e) Administering
 - f) Monitoring
 - g) Education
 - h) Reconciliation
 - i) Uncertain
 - j) None – exclude

- 4) Please classify the article to the relevant categories below (mandatory to answer this question, and can choose more than one). These correspond with the 5 Key Questions being addressed in the review
 - a) Related to patient outcomes
 - b) Deals with costs, benefits etc of the system
 - c) Deals with values proposition to any of the users (*value issues that users consider when deciding to use the system “what benefit is there for me?”*)
 - d) Reports on system characteristics with related outcomes (*e.g. usability, validity, use, satisfaction etc*)
 - e) Deals with issues relating to sustainability of the system (*maintenance,*)
 - f) Reports on computerized decision support in medication management

Instructions for Key Questions algorithm to determine which KQ the article pertains to and if the article has comparison groups and is hypothesis driven to be applicable to KQ1.

1. First fill in the article UI and your initials at the top of the page.
2. Make sure the article fits our criteria thus far:
 - a. It is about an integrated technology (EXCEPT for PDAs or insulin pumps-they need to take individual patient data and provide MM support) that enables the management of the medication process-across all phases
 - b. It contains data from a study, which can be numerical (quantitative) or text data from a qualitative study where focus groups, delphi method, interviews etc were conducted and transcribed.
 - c. **If the article is to be excluded, please state the specific failing (review article, no data, not integrated, not MM) and draw a line through the blue sheet.**
3. Start on the left-hand algorithm.
4. Does the study assess the values people consider when determining whether or not a particular application is useful to them? They must have data on this (qualitative or quantitative).
 - a. These will likely be survey or interview studies with people's opinions, and can be patients, clinicians, pharmacists, hospital administrators etc.
 - b. If yes, make sure you circle the KQ3 box.
 - c. For **all** articles continue to the next question.
5. Does the study assess decisions to purchase, implement or use a system? They must have data on this (qualitative or quantitative).
 - a. If yes, does it further describe the systems characteristics (such as proprietary, home-grown), or information about certification or conformity, or flexibility in the system (ability to customize) etc?
 - b. If yes, make sure you circle the KQ4 box.
 - c. For **all** articles continue to the next question.
6. Does the study discuss sustainability of the system? OR does the study report on a system that is proven sustainable, ie. it has been in use for 3+ years, in a real practice setting. Discussion of sustainability could include financial sustainability, maintenance and updating issues, adaptability of the system. It can relate to financial, technological, socio-political or organizational factors. They must have data on this (qualitative or quantitative).
 - a. If yes, make sure you circle the KQ5 box.
 - b. For **all** articles continue to the next question.
7. Does the study address the electronic communication between the clinician and the pharmacy? They must have data on this (qualitative or quantitative).
 - a. If yes, make sure you circle the KQ6 box.
 - b. For **all** articles continue to the **next algorithm**
8. Does the study measure one of the following? Please refer to the outcomes rubric for guidance. If you have an unusual measurement and you don't know where it falls, make a note of it at the bottom of the assessment page: (see table on p 3 for guidance)
 - i. Process (about providing care)
 - ii. Other outcomes (satisfaction, skills etc)

- iii. Cost
 - iv. Clinical (patient related) outcomes
 - v. Population level (eg screening rates)
 - vi. Composite outcomes (are formed by combining individuals' scores on a collection of singular measures-usually in trials with a range of treatment effects)
- b. If no, STOP and circle on the far right the KQ that are addressed by the study that you found in the left-column of the algorithm. If there are NO KQ addressed, make sure you give the article to Cynthia.
 - c. Indicate the methodology used in the box at the bottom of the blue page
 - d. If yes, continue
9. Does the study have a comparison group?
- a. Assess if it has a comparison group. This can be a different time-point, a before-after, a control group, another intervention group. But they must be comparing one set of data to another.
 - b. If no, is the study qualitative? – If yes, circle KQ1 and continue to the bottom of the page and make a note in the outcomes section and methodology box.
 - c. If NO, STOP and circle on the far right the KQ that are addressed by the study that you found in the left-column of the algorithm. If there are NO KQ addressed, make sure you give the article to Cynthia.
 - d. If yes, continue
10. Is the study hypothesis driven? This means that they will state in their introduction the effect they think they will see based on their intervention, or they will statistically analyze/compare the groups. The presence of p-values or confidence intervals (CI) is another indicator that the study was hypothesis driven.
- a. If no, STOP and circle on the far right the KQ that are addressed by the study that you found in the left-column of the algorithm. **Please circle 'list for KQ1'.**
 - b. If yes, continue
11. Does the article report on a CDSS (clinical decision support system) defined as: “Clinical Decision Support systems link health observations with health knowledge to influence health choices by clinicians for improved health care”—we need them to be computerized and providing health information related to medication management.
- a. If no, Circle KQ1. Also circle on the far right all of the KQ that are addressed by the study that you found in the left-column of the algorithm. Continue to the bottom of the page and make a note in the outcomes section and methodology box.
 - b. If yes, continue
12. Is it a Randomized controlled trial?
- a. If no, Circle KQ1. Also circle on the far right all of the KQ that are addressed by the study. Continue to the bottom of the page and make a note in the outcomes section and methodology box.
 - b. If yes, circle KQ7, and KQ1 and ensure that all KQs addressed by the study are indicated in the far right. Continue to the bottom of the page and make a note in the outcomes section and methodology box.

Outcomes: follow the outcomes table for guidance. Indicate with 1° the primary outcomes if the authors make the distinction.

Methods: Follow the methods algorithm to determine the methodology used in the study.

Broadly, we will categorize outcomes into the following categories. The table gives some specific measures/examples to help guide you. There will be measures that are difficult to classify. Please make a note at the bottom of the blue page.

| Process | Other | Clinical |
|--|---|--|
| Errors <ul style="list-style-type: none"> ◦ Error rates ◦ Types of errors ◦ Potential ADEs ◦ Number of errors Efficiency <ul style="list-style-type: none"> ◦ Time related outcomes ◦ Utilization of care ◦ Provider time ◦ Time to dispensing Compliance/adherence <ul style="list-style-type: none"> ◦ To guidelines ◦ To order sets ◦ Care related Changes in prescribing decisions <ul style="list-style-type: none"> ◦ Altering dosages ◦ Changing preparation ◦ Changing Rx pattern Monitoring/surveillance <ul style="list-style-type: none"> ◦ Change in test ordering ◦ Inappropriate test ordering ◦ Screening rates Preventative care <ul style="list-style-type: none"> ◦ Vaccinations ◦ Population screening | Satisfaction/Usability K/S/A <ul style="list-style-type: none"> ◦ Knowledge ◦ Skills ◦ Attitudes Usage Other Cost Change in utilization Costs Length of stay assoc costs Qualitative Themes Text excerpts | Other clinical: <ul style="list-style-type: none"> ◦ Infection rate ◦ Length of stay ◦ Compliance Error effects: <ul style="list-style-type: none"> ◦ Adverse Drug Events (ADE) QoL Physiological measures: <ul style="list-style-type: none"> ◦ Hb1Ac ◦ BP ◦ INR Population Screening Composite One index that encompasses a number of measures eg. risk assessment values etc. |

Data Abstraction Form with Instructions

Data abstraction instructions

General Study Information:

| Question | General Study Information | Options | Instructions |
|----------|---|-------------------|--|
| 1.1 | First Author | (text) | Type in the last name of the first author |
| 1.2 | Year of publication | (text) | enter the year of publication |
| 1.3 | What phase of medication management is being reported on? | | can select more than 1 |
| | Prescribing | CheckBox | The process of a clinician deciding and ordering a medication for a patient |
| | Transmission, order communication | CheckBox | The bi-directional communication of the prescription and it's fine-tuning between clinician and pharmacist. Includes electronic data exchange |
| | Dispensing | CheckBox | The preparation of the prescription in the pharmacy and getting it to the patient |
| | Administering | CheckBox | The patient taking the drug. Can be administered by nurse, other clinician, patient or caregiver. |
| | Monitoring including patient adherence and compliance | CheckBox | Monitoring of patient taking drug for adverse events, reactions, compliance, adherence, and efficacy. Include lab monitoring or ordering of tests to monitor drug levels etc. |
| | Education of patients and clinicians but not pre-professional education | CheckBox | Pre-professional education includes nursing, medical, dental etc students learning their profession--they are excluded. Interns and residents are included as well as patients. Need to include the issue of medication as well as education around taking and reviewing medications. |
| | Other e.g. discharge summaries. MM in general, etc | CheckBox | Discharge summaries are provided when the patient transitions from one level of care to another including home. For example, from the surgical ward to home or a nursing home. Reconciliation can go here as well where clinicians and patients check that lists of drugs for a particular patient is complete and up to date. |
| 1.3.1 | Specify Other | text box | |
| 1.4 | What is the country address of the first author? | Manual Text Entry | Please use: US, Can, UK, NL, Eur, Asia, Aus, NZ, other |
| 1.5 | Select funding information | | This can usually be found just before the reference section or on the first page of the paper in small print, sometimes in the methods section. |
| | Internal funding | Radio--start off | This would be a statement that the division or group provided funding or if the study says things like "no external funding was used". |
| | External funding by grants, projects, contracts | Radio--start off | Funding section will indicate funding agency name |
| | External funding by industry, companies | Radio--start off | Funding section will indicate sponsoring company name |
| | Both internal and external | Radio--start off | |

| Question | General Study Information | Options | Instructions |
|-------------|--|------------------|--|
| | Not specified | Radio--start off | Use this when no funding information is provided in the paper. |
| | Other | Radio--start off | |
| 1.5.1 | Indicate Funding agency | textbox | Please note name of funding agency OR company |
| 1.6 | Is the article a general or systematic review summarizing a number of studies or the evidence on their question? | yes/no | If it is an original study, answer no. An original study reports first hand observations of a STUDY-having some kind of data. If this article is not about an original study, and it summarizes the evidence or is a systematic review with many studies included, then answer yes. If yes, STOP If no, continue to next question. |
| 1.6.1 | Is the article an original study, in which the authors report first hand observations, either qualitative or quantitative? | Yes/no. | |
| 1.6.1.1 | If the article is an original study, is it ONLY qualitative? | Yes/no | |
| 1.6.1.1.1 | If the article is an original study, is it ONLY quantitative? Does it report numerical findings? | Yes/no. | |
| 1.6.1.1.1.1 | Does the article reports mixed methods--both qualitative and quantitative? | Yes/no. | |
| 1.6.1.2 | Provide a clear description of the study, including all PICOM components | Open text | Here we need a very clear description of the study intervention, include the type of intervention(s), the groups involved. Try to be concise as possible. Often the aim of the study can help. Please include PICOM if possible (people, intervention, comparison, outcomes, method/design). Put this in paragraph form using 3-4 sentences. |
| 1.6.1.3 | Does the study have a comparison group | y/n | A comparison group can be the same population at a different time (e.g. before and after or time series), or it can be different groups of doctors (specialists vs. GPs etc) or it can be different groups of patient, different hospitals, clinic types, systems etc. |
| 1.6.1.3.1 | What is the intervention group being compared to? | | |
| | Usual care | Checkbox | Pre-implementation or baseline would be considered usual care, where care has not changed from the usual |
| | A control group | Checkbox | This does not include the 'before' for a pre-post implementation study or baseline in a time series study. |
| | Another technology/system | Checkbox | |
| | Other | Checkbox | |
| 1.6.1.3.1.1 | Specify Other | text box | describe the other comparison groups here |

| Question | General Study Information | Options | Instructions |
|----------|---------------------------|----------|---|
| 1.7 | What is the study design? | | |
| | RCT | Checkbox | “experimental design that studies the effect of an intervention or treatment using at least two groups: one that received the intervention and one that did not; participants ARE randomly assigned to a group (therapy, prevention)” |
| | Non-randomized trial | Checkbox | “experimental design that studies the effect of an intervention or treatment using at least two groups: one that received the intervention and one that did not; participants ARE NOT randomly assigned to a group (therapy, prevention)” |
| | Cohort study | Checkbox | Cohort study: involves establishing groups, often people, one of which is “exposed” (e.g. HIT) and one is not exposed. Both groups followed forward in time to determine if the outcomes of interest develop. |
| | Case control | Checkbox | A study where groups of people are formed, one of which has the outcome of interest (e.g. better prescribing) and one of which does not (not better prescribing). Often members in the groups are “matched” in relation to things like age or experience. People in both groups are evaluated to assess if the exposure of interest (e.g. EHRs) were present in the past. |
| | Time series | Checkbox | A study in which periodic measurements are obtained prior to, during, and following the introduction of an intervention or treatment in order to reach conclusions about the effect of the intervention. This usually has more than 2 time points. If only 2 points, 1 pre- and 1 post-implementation, then it is a before-after study. |
| | Before-after | Checkbox | A type nonrandomized study where data are collected before and after patients receive an intervention. Before-after studies can have a single arm where the before group is usual care or it can include a control group. |
| | Cross-sectional | Checkbox | involve observation of some subset of a population of items all at the same time, in which, groups can be compared |
| | Qualitative | Checkbox | Research that derives data from observation, interviews, or verbal interactions and focuses on the meanings and interpretations of the participants. Must say they are doing qualitative work-words like qualitative, themes, narrative, ethnography, phenomology. |
| | Mixed methods | Checkbox | an approach to professional research that combines the collection and analysis of quantitative and qualitative data |
| | Case series | CheckBox | “A medical research study that tracks patients or other participants with a known exposure given similar treatment or examines their medical records for exposure and outcome.” |

| Question | General Study Information | Options | Instructions |
|----------|--|----------|---|
| | Survey | CheckBox | A research method involving the use of questionnaires and/or statistical surveys to gather data about people and their thoughts and behaviors. |
| | Observational study | CheckBox | |
| 1.8 | Do you have any comments about the study/article not captured in the data? | text box | If there is something you cannot find/is wonky/you want to clarify etc. Use this comment box |
| 1.9 | Does this article have a companion(s)? | y/n | A companion is an article of the SAME study, with the SAME participants, that reports on different data. E.g. your article may have the economic data but they refer to another article reporting the clinical data. It does not include a study of the same system in the same setting but done with other patients/doctors/interventions. |
| 1.9.1 | if yes, provide the citation details: | text box | from the reference section, cut and past the reference for any companion article(s) |

Study Population:

| Question | Study Population | Options | Instructions |
|----------|--|----------|---|
| 2.1 | What is the primary unit of study and analysis? (select one of the following) | | info: this is the unit they report their data/numbers and analysis on |
| | Health care providers | CheckBox | |
| | Patients | CheckBox | if they're measuring a patient's blood, ADEs, admissions --unhitching that happens to the patient, select this |
| | Institutions | CheckBox | |
| | IT Systems | CheckBox | |
| | Medications, prescriptions, orders | CheckBox | |
| | Geographic regions | CheckBox | |
| | Other | CheckBox | |
| 2.1.1 | Specify other (text) | text box | |
| | Not answered | CheckBox | |
| 2.2 | Are people being measured as the unit of study? | y/n | |
| 2.2.1 | Are clinicians being studied? | y/n | Who is being studied? Select as many groups as long as they represent at least 10% of the sample or data presented. |
| 2.2.1.1 | Please Select the types of clinicians being studied. | | can select more than 1 |
| | Physicians undifferentiated or cannot sort out | CheckBox | |
| | Primary care physicians/GPs, Family physicians | CheckBox | |
| | Specialists | CheckBox | |
| | Hospitalists | CheckBox | |
| | Other Physicians | CheckBox | |

| Question | Study Population | Options | Instructions |
|----------|---|----------|---|
| | Mid level practitioners (PAs, NPs, APN, midwives) | CheckBox | |
| | Nurses | CheckBox | |
| | Psychologists, counselors | CheckBox | |
| | Pharmacists | CheckBox | |
| | Dentists | CheckBox | |
| | Hospital administrators | CheckBox | |
| | Other health professionals | CheckBox | |
| 2.2.2 | Are caregivers being studied? | y/n | |
| 2.2.3 | Are patients being studied? | y/n | |
| 2.2.3.1 | Please select the types of patients being studied. | | |
| | Infants (0 to 2 years) | CheckBox | |
| | Children (2 to 12) | CheckBox | |
| | Adolescents (13 to 18) | CheckBox | |
| | Adults (19 to 44) | CheckBox | |
| | Middle age (45 to 64) | CheckBox | |
| | Geriatric (65 plus) | CheckBox | |
| | Undifferentiated | CheckBox | |
| | Not Specified | CheckBox | |
| 2.3 | How many groups were studied?--not subgroups but original study groups e.g., number of groups randomized in an RCT or number of groups in a cohort study. | text box | How many groups were studied?--not subgroups but original study groups e.g., number of groups randomized in an RCT or number of groups in a cohort study. |
| 2.4 | Were the inclusion criteria given? | yes/no | Did they define the people/participants/population that was included in their sample? |
| 2.4.1 | What were the Inclusion criteria for the study (usually reported in the methods, copy and paste) | text box | copy and paste from methods when applicable |
| 2.5 | Were the exclusion criteria given? | yes/no | Did they define the people/participants/population that were purposely EXCLUDED from their sample? |
| 2.5.1 | What were the exclusion criteria for the study (usually reported in the methods, copy and paste) | text box | copy and paste from methods when applicable |

Setting:

| Question | Setting | Options | Instructions |
|----------|---|----------|------------------------------------|
| 4.1 | Did the study take place in a hospital? | y/n | (can select more than one setting) |
| 4.1.1 | Indicate the type(s) of hospital(s): | | |
| | Acute care/tertiary | CheckBox | |
| | Critical care units (CCU, ICU, NICU) | CheckBox | |
| | Emergency department | CheckBox | |
| | General hospital | CheckBox | |
| | Palliative/hospice | CheckBox | |
| | Pediatric stand alone hospital | CheckBox | |

| Question | Setting | Options | Instructions |
|----------|---|---------------|---|
| | Other specialty hospital (rehab, oncology) | CheckBox | |
| | Not specified | CheckBox | |
| | Do the authors report the number of beds? | y/n | |
| 4.1.2.1 | How many beds were reported? | text box | This should be the total of the beds for a multiple hospital study |
| 4.2 | Select any other settings in which the study took place: | | |
| | Ambulatory care (clinic, doctor's office, etc.) | CheckBox | |
| | Long term care (nursing homes) | CheckBox | |
| | Home | CheckBox | |
| | Community (school, community centre, etc.) | CheckBox | |
| | Pharmacy | CheckBox | |
| | None | CheckBox | |
| | Other | CheckBox | |
| 4.2.1 | Specify Other | text box | |
| | Not specified | CheckBox | |
| 4.3 | Is there a pharmacy involved? | y/n | A pharmacy or pharmacist must be directly involved in the study. (can select more than one). |
| 4.3.1 | Specify the type(s) of pharmacy: | | |
| | Inpatient hospital based | CheckBox | An inpatient pharmacy is located in a hospital and is the pharmacy system that serves patients while they are hospitalized--usually patients who are staying overnight in the hospital. |
| | Outpatient hospital based | CheckBox | This pharmacy is located in a hospital but provides drugs for those patients who are not hospitalized overnight. |
| | Other institution based | CheckBox | For example, in a nursing home setting. |
| | HMO pharmacy | CheckBox | |
| | Veterans Affairs Pharmacy | CheckBox | |
| | Pharmacy chain | CheckBox | |
| | Stand alone non chain store (e.g. family run) | CheckBox | |
| | Health insurance company based and Pharmacy Benefit Management (PBM) pharmacies | CheckBox | |
| | Other mail/email in pharmacies | CheckBox | |
| | Nuclear pharmacies (radioactive drugs) | CheckBox | |
| | Compounding pharmacies (<i>those that produce their own products using various chemicals and binders</i>) | CheckBox | |
| | Not specified | CheckBox | |
| | Other | CheckBox | |
| 4.3.1.1 | Specify other | text box | |
| 4.4 | Was the study conducted in an academic setting? Answer yes if at least 1 setting was academic e.g. University Hospital. | y/n/uncertain | look for names of Universities, or 'academic hospital', 'teaching hospital', "university hospital" etc |

Diseases/Drugs studied:

| Question | Drug/disease | Options | Instructions |
|-----------|---|------------------|---|
| 3.1 | Were any disease(s) or condition(s) studied? | y/n | they might not 'measure' the disease, but if they are looking at a specific population of patients with a certain disease or the intervention is to improve treatment for a certain disease, then select this. E.g. flu vaccines =flu; diabetes PHR=diabetes; CDSS for DVT prevention=DVT |
| 3.1.1 | Please indicate information about the disease(s)/condition(s) studied: | | |
| | All diseases and conditions | Radio--start off | articles that don't specify a condition but is about MM in any patient |
| | One disease or condition | Radio--start off | Articles specifically patients with 1 condition e.g. cancer, hypertension etc |
| | Not answered | Radio--start off | Articles that are not focused on patient MM e.g. articles about systems, settings etc |
| 3.1.1.1 | Which disease was specified | | |
| | Asthma | Radio--start off | |
| | Diabetes | Radio--start off | |
| | Heart diseases | Radio--start off | |
| | Deep venous thrombosis, pulmonary embolism, other clotting issues | Radio--start off | |
| | Depression/schizophrenia/all mental health | Radio--start off | |
| | Other | Radio--start off | |
| 3.1.1.1.1 | Specify Other | text box | |
| 3.2 | Were certain drugs or classes of drugs the focus of the study? | y/n | the study must focus on certain drugs or classes of drugs rather than MM in general |
| 3.2.1 | Specify the drugs, classes or families of drugs | textbox | |
| 3.2.2 | Were the drugs | | |
| | Toxic drugs/drugs with narrow therapeutic index (e.g. warfarin, digoxin) | Radio--start off | |
| | Controlled substance | Radio--start off | |
| | Both | Radio--start off | |
| 3.2.2.1 | Indicate the controlled substance: | text box | |
| | None of the above | Radio--start off | |
| | Not answered | Radio--start off | |
| 3.2.3 | Did the article focus on generic vs. brand names of drugs | y/n | |
| 3.2.3.1 | Indicate which drugs | text box | |
| 3.2.4 | Did the article focus on sound-alike or look-alike drugs? | y/n | |
| | Indicate which drugs | text box | |
| 3.3 | Did the article report on the preparation of a drug (the form in which it is administered)? | y/n | (can select more than one) |
| 3.3.1 | Indicate the preparations included: | | |
| | Oral | CheckBox | |

| Question | Drug/disease | Options | Instructions |
|----------|---------------|----------|--------------|
| | Intravenous | CheckBox | |
| | Injection | CheckBox | |
| | Inhaled | CheckBox | |
| | Parenteral | CheckBox | |
| | Topical | CheckBox | |
| | Rectal | CheckBox | |
| | Not Specified | CheckBox | |
| | Other | CheckBox | |
| 3.3.1.1 | Specify Other | TextBox | |

The technology:

| Question | Technology | Options | Instructions |
|----------|--|----------|--|
| 5.1 | Please indicate the nature of the producer of the Health IT system used: | | Health IT systems can be built by various people or groups. Early ones were built by an individual clinician who could program personal computers. Modern ones are often built/developed by commercial companies such as GE or Seimans. (can select more than one) |
| | Commercial | CheckBox | Is it a company--often for profit companies |
| | Home grown | CheckBox | Was it developed by those who are using it--e.g. by clinicians on the wards or by others working under clinicians, such as onsite programmers. |
| | Both | CheckBox | Can be a mix of commercial and home grown. Often a system might be started by an individual or group of individuals in a hospital or ward and when it shows promise and needs further development a commercial company may take over development and production of the system. |
| | Not specified | CheckBox | Check this if the article does not say who developed the system. |
| | Other | CheckBox | This could be a non-profit organization such as the World Health Organization. |
| 5.1.1 | specify other | text box | |
| 5.2 | Please indicate the nature of the MMS used: | | (can select more than one). This question is designed to look at the "ownership" issue in relation to the MMS. For example, is it Open Source so anyone can implement the system. (can select more than one) |

| Question | Technology | Options | Instructions |
|----------|---|----------|--|
| | Open source | CheckBox | From: Wikipedia. Software whose source code is published and made available to the public, enabling anyone to copy, modify and redistribute the source code without paying royalties or fees. Open source code evolves through community cooperation. These communities are composed of individual programmers as well as very large companies. Some examples of open source initiatives are Linux, Eclipse, Apache, Tomcat web server, Mozilla, and various projects hosted on SourceForge and elsewhere. For eHealth, one of the most prevalent is OSCAR for EHRs |
| | Proprietary-commercial | CheckBox | A commercial company "owns" the software and users need to buy or lease the product and its code |
| | Proprietary-academic | CheckBox | An academic institution "owns" the software/ product and users need to buy or lease the product and its code. |
| | Not Specified | CheckBox | Check yes if not specified |
| | Other | CheckBox | Should be very few of these... |
| 5.2.1 | Specify Other | text box | |
| 5.3 | What kind of conformity standards did the MMS meet (can select more than 1) | | MMSs must "interact" or integrate with many other information systems. Standards need to be implemented to insure that when the systems interact/interoperate they both function accurately and efficiently. MMSs that will be used by multiple users in multiple settings need to be developed based on standards. These standards can come from the computing, health, business, etc communities. HL7 is one often encountered standard. A good hint on the presence of standards will be the use of CAPITALS to describe things like standards and conformity. (can select more than one) |
| | Not specified | CheckBox | You will see this most often in articles. |
| | Local | CheckBox | This would be hospital wide system conformity with local standards and local IT and MMS components, systems. |
| | State/provincial | CheckBox | Some state/provincial standards exist. Most will be National although some like the BC, MB, AB, and ON HISC (Health Information Standards Council) standards exist. |
| | National-US | CheckBox | National standards include ANSI (American National Standards Institute) standards, HIPPA, PHIPPA |

| Question | Technology | Options | Instructions |
|----------|---|-------------------|--|
| | National-other | CheckBox | You may need to list stuff here. the UK, Australia, the Netherlands, Canada have National standards in place. |
| | International | CheckBox | Groups like ISO and HL7 go here |
| | Certification Commission for Healthcare Information Technology (CCHIT) | CheckBox | This is a US based standard and is one that needs to be addressed by the final report. |
| | Other | CheckBox | |
| 5.3.1 | Specify other | text box | |
| 5.4 | Did the system have a specific name? | y/n | |
| 5.4.1 | If so, What was the name of the MMS used? | text box | What was the system called? |
| 5.5 | Did the MMS system replace an existing system? | Yes/No/don't know | |
| 5.6 | What kind of MMS system is being studied in the article? Can select more than 1 | | Select the kind of system(s) used in the article. (can select more than one) |
| | Bacoding-medication administering | CheckBox | |
| | Bacoding-dispensing | CheckBox | |
| | eMedication administration system (eMAR, eTAR) | CheckBox | |
| | CPOE/POE system | CheckBox | |
| | CDSS/CDS/CCDS/reminders | CheckBox | |
| | Eprescribing | CheckBox | |
| | eTransmission-of the prescription to/from doctor to pharmacy | CheckBox | |
| | AMDD anesthesia medication dispensing system | CheckBox | |
| | Pharmacy information system | CheckBox | |
| | None | CheckBox | |
| | Other | CheckBox | |
| 5.6.1 | Specify Other | text box | |
| 5.7 | Is the system described as: | | can select more than 1 |
| | Stand alone | CheckBox | |
| | Integrated with another system or set of systems | CheckBox | |
| | a PDA/handheld access | CheckBox | |
| | Not Specified | CheckBox | |
| | Other | CheckBox | |
| 5.7.1 | specify other | text box | |
| 5.8 | What other system(s) is the MMS integrated with? (can select more than 1) | | can select more than 1. A strong (i.e. effective) MMS will integrate (i.e. talk to) multiple other systems. We need to know which systems that the MMS is integrated with. |
| | EHR/EMR system | CheckBox | |
| | Personal health records systems | CheckBox | |
| | CPOE/POE system | CheckBox | |
| | CDSS/CDS/CCDS/reminders | CheckBox | |
| | Billing/administration system | CheckBox | |
| | Laboratory system | CheckBox | |
| | Imaging systems | CheckBox | |
| | Patient decision support system | CheckBox | |
| | Pharmacy | CheckBox | |

| Question | Technology | Options | Instructions |
|----------|--|--------------------------------|--|
| | Formulary | CheckBox | |
| | Insurance | CheckBox | |
| | Barcoding system | CheckBox | |
| | Hospital information system | CheckBox | |
| | Other | CheckBox | |
| 5.8.1 | Specify Other | text box | |
| | not specified | CheckBox | |
| 5.9 | Is the MMS web-based? | | |
| | Fully we based | Radio--start off | |
| | Partly web based | Radio--start off | |
| | Not web based (internal system) | Radio--start off | |
| | not reported | Radio--start off | |
| 5.10. | What kind of computer equipment does the system use? | | can select more than 1 |
| | PCs for clinicians-including COWs (computers on wheels) | CheckBox | |
| | PCs for patients | CheckBox | |
| | PDAs for clinicians | CheckBox | |
| | PDAs for patients and caregivers | CheckBox | |
| | Touchscreens for clinicians | CheckBox | |
| | Touchscreens for patients and caregivers | CheckBox | |
| | Mobile carts | CheckBox | |
| | Kiosks | CheckBox | |
| | Robots | CheckBox | |
| 5.10.1 | Other | text box | |
| | Not Specified | CheckBox | |
| 5.11 | What is the source of patient data for processing by the technology? | | Where does the patient data come from that is used by the IT system? can select more than one. (can select more than one) |
| | EHRs/EMRs | CheckBox | |
| | Other internal-e.g. lab data, pharmacy records, PHRs | CheckBox | |
| | Third party vendor-e.g.. insurance database | CheckBox | |
| | Personal health records systems | CheckBox | |
| | Medical devices such as glucometers | CheckBox | |
| | Manual entry | CheckBox | |
| | Not specified | | |
| 5.12 | When was the system implemented | text box for date/not reported | enter the date of original implementation of the original system. Note the system could have been built upon and developed since that date. Month/year 00/0000 |
| 5.13 | What was the start date of the study? | text box for date/not reported | enter the Month/year the study began. 00/0000 |
| 5.14 | What was the end data of the study? | text box for date/not reported | enter the Month/year the study ended. 00/0000. for a survey, beginning and end date often the same |
| 5.15 | Was the system in use for more than 3 years? | y/n/can't determine | Is end of study date at least 3 years since the initial implementation? |

Outcomes:

| Question | Outcomes | Options | Instructions |
|----------|---|------------------------------|---|
| 7.1 | Did the authors clearly declare their primary outcome? | y/n | |
| 7.2 | What was the sample size of the study? | text box for number | |
| 7.3 | What was the unit for the sample size? | text box | |
| 7.4 | Were costs associated with the use of the MMIT system assessed? | y/n | |
| 7.4.1 | State the costs in as much detail as possible. | text box | |
| 7.5 | Were adverse effects of the HIT assessed? | yes/no/unstated/ | this would be a separate measurement of any negative impact of the technology on the MM process. Unintended consequences could be 1 example. See article 37. A negative effect on your primary outcome does NOT go here. |
| 7.5.1 | State the findings, including any assessment of clinical impact/relevance. State the findings in terms of % change among comparison groups, for the primary outcome only. | text box | if not statistically analyzed do not include this here |
| 7.5.2 | What was the p value of the analysis above? | options: ns/ p<.05/P<.001 | |
| 7.6 | Were the impacts of the MMIT on PROCESS outcomes measured? | y/n | report only on the primary outcome, or if not clear, then abstract the medication management outcomes. Process outcomes are associated with the care given to patients, relating to errors, efficiencies, adherence to guidelines, prescribing changes, monitoring of patients e.g. labs, ordering preventative care etc. |
| 7.6.1 | Indicate the process outcomes measured: | | |
| | Errors | checkbox | |
| | Efficiency | checkbox | |
| | Adherence/compliance with guidelines | checkbox | |
| | Changes in prescribing patterns | checkbox | |
| | Changes in monitoring/surveillance activities (e.g. lab test ordering) | checkbox | |
| | Preventative care | checkbox | |
| | Other | checkbox | |
| 7.6.1.1 | Specify other | text box | |
| 7.6.2 | State the general findings for process outcomes | text box | provide a general statement about the process outcomes results. Include details such as ...a reduction in inappropriate prescribing, less time to administration, etc. |
| 7.6.3 | Define outcome 1-usually the primary outcome | text box | the primary outcome is the one of most importance. Primary outcome can be determined by the aim/objective/ purpose or if there is a power calculation for it or if they say it is the 1° outcome |

| Question | Outcomes | Options | Instructions |
|---------------|---|----------|---|
| 7.6.4 | provide RRR statement-outcome 1 | text box | % vs. %, RRR, p |
| 7.6.5 | Was there a second outcome? | y/n | what was the second outcome/sequential important outcome |
| 7.6.5.1 | Define outcome 2 | text box | % vs. %, RRR, p |
| 7.6.5.2 | Provide RRR statement-outcome 2 | text box | % vs. %, RRR, p |
| 7.6.5.3 | Was there a third outcome? | y/n | what was the third outcome/sequential important outcome |
| 7.6.5.3.1 | Define outcome 3 | text box | % vs. %, RRR, p |
| 7.6.5.3.2 | Provide RRR statement-outcome 3 | text box | % vs. %, RRR, p |
| 7.6.5.3.3 | Was there a fourth outcome? | y/n | what was the fourth outcome/sequential important outcome |
| 7.6.5.3.3.1 | Define outcome 4 | text box | % vs. %, RRR, p |
| 7.6.5.3.3.2 | Provide RRR statement-outcome 4 | text box | % vs. %, RRR, p |
| 7.6.5.3.3.3 | Was there fifth outcome? | y/n | what was the fifth outcome/sequential important outcome |
| 7.6.5.3.3.3.1 | Define outcome 5 | text box | % vs. %, RRR, p |
| 7.6.5.3.3.3.2 | Provide RRR statement-outcome 5 | text box | % vs. %, RRR, p |
| 7.6.6 | Was the PROCESS outcome improved on the primary outcome measure (or more that 50% of the measures of process were improved if no primary outcome measure indicated) | y/n | |
| 7.7 | Were the impacts of the MMIT on 'OTHER' outcomes measured? | y/n | Again, the "other" outcome must be the primary outcomes, or if not indicated, related to medication management. |
| 7.7.1 | Indicate the 'other' outcomes measured: | | |
| | Use/usage | checkbox | |
| | Knowledge/skills/attitude | checkbox | |
| | Satisfaction | checkbox | |
| | Usability | checkbox | |
| | Other | checkbox | |
| 7.7.1 | Specify Other | text box | |
| 7.7.2 | State the general findings for other outcomes | text box | |
| 7.7.3 | Define outcome 1 - usually the primary outcome | text box | What was the primary outcome/most important outcome |
| 7.7.4 | Provide RRR statement-outcome 1 | text box | % vs. %, RRR, p |
| 7.7.5 | Was there a second outcome? | y/n | |
| 7.7.5.1 | Define outcome 2 | text box | What was the second outcome/sequential important outcome |
| 7.7.5.2 | Provide RRR statement-outcome 2 | text box | % vs. %, RRR, p |
| 7.7.5.3 | Was there a third outcome? | y/n | |
| 7.7.5.3.1 | Define outcome 3 | text box | What was the third outcome/sequential important outcome |

| Question | Outcomes | Options | Instructions |
|---------------|--|-------------------|---|
| 7.7.5.3.2 | Provide RRR statement-outcome 3 | text box | % vs. %, RRR, p |
| 7.7.5.3.3 | Was there a fourth outcome? | y/n | |
| 7.7.5.3.3.1 | Define outcome 4 | text box | What was the fourth outcome/sequential important outcome |
| 7.7.5.3.3.2 | Provide RRR statement-outcome 4 | text box | % vs. %, RRR, p |
| 7.7.5.3.3.3 | Was there a fifth outcome? | y/n | |
| 7.7.5.3.3.3.1 | Define outcome 5 | text box | What was the fifth outcome/sequential important outcome |
| 7.7.5.3.3.3.2 | Provide RRR statement-outcome 5 | text box | % vs. %, RRR, p |
| 7.7.6 | Was the other outcome improved on the primary outcome measure (or more that 50% of the measures of process were improved if no primary outcome measure indicated)? | yes/no/not tested | |
| 7.8 | Where the impacts of MMIT on patient CLINICAL outcomes measured? | y/n | This would apply for studies where patients are the unit of study. Measurements could be physiological e.g. blood pressure, Hb1ac etc, or adverse events, length of stay, mortality, quality of life etc. Again, the "other" outcome must be the primary outcomes, or if not indicated, related to medication management. |
| 7.8.1 | Indicate the patient clinical outcomes measured: | | |
| | Physiological measure | checkbox | |
| | Adverse drug events | checkbox | |
| | Length of stay | checkbox | |
| | Mortality | checkbox | |
| | Quality of Life | checkbox | |
| | Other patient events | checkbox | List anything else here that would be important (i.e. felt or appreciated) by patients including things like improved conception rates. |
| | Other | checkbox | |
| 7.8.1.1 | Specify Other | text box | |
| 7.8.2 | State the general findings for the patient clinical outcomes | text box | |
| 7.8.3 | Define outcome 1-the primary outcome | text box | The primary outcome is the one of most importance. Primary outcome can be determined by the aim/objective/ purpose or if there is a power calculation for it or if they say it is the 1° outcome |
| 7.8.4 | Provide RRR statement-outcome 1 | text box | % vs. %, RRR, p |
| 7.8.5 | Was there a second outcome? | y/n | |
| 7.8.5.1 | Define outcome 2 | text box | What was the second outcome/sequential important outcome |
| 7.8.5.2 | Provide RRR statement-outcome 2 | text box | % vs. %, RRR, p |
| 7.8.5.3 | Was there a third outcome? | y/n | |

| Question | Outcomes | Options | Instructions |
|---------------|--|----------|--|
| 7.8.5.3.1 | Define outcome 3 | text box | What was the third outcome/sequential important outcome |
| 7.8.5.3.2 | Provide RRR statement-outcome 3 | text box | % vs. %, RRR, p |
| 7.8.5.3.3 | Was there a fourth outcome? | y/n | |
| 7.8.5.3.3.1 | Define outcome 4 | text box | What was the fourth outcome/sequential important outcome |
| 7.8.5.3.3.2 | Provide RRR statement-outcome 4 | text box | % vs. %, RRR, p |
| 7.8.5.3.3.3 | Was there a fifth outcome? | y/n | |
| 7.8.5.3.3.3.1 | Define outcome 5 | text box | What was the fifth outcome/sequential important outcome |
| 7.8.5.3.3.3.2 | Provide RRR statement-outcome 5 | text box | % vs. %, RRR, p |
| 7.8.6 | Was the CLINICAL outcome improved on the primary outcome measure (or more that 50% of the measures of process were improved if no primary outcome measure indicated) | y/n | |
| 7.9 | Where the impacts of MMIT on population level outcomes measured? | y/n | Again, the population outcome must be the primary outcomes, or if not indicated, related to medication management. |
| 7.9.1 | How was this measured? | text box | |
| 7.9.2 | What did the study conclude? | text box | |
| 7.10. | Where the impacts of MMIT on composite outcomes measured? | y/n | Again, the composite outcome must be the primary outcomes, or if not indicated, related to medication management. |
| 7.10.1 | How was this measured? | text box | |
| 7.10.2 | What did the study conclude? | text box | |
| 7.11 | Did the study address decisions to buy/implement or use for any of the stakeholders? | y/n | |
| 7.11.1 | How was this measured? | text box | Include the measurement and method |
| 7.11.2 | What did the study conclude? | text box | |
| 7.12 | Did the study address values propositions for any of the stakeholders? | y/n | |
| 7.12.1 | How was this measured? | text box | |
| 7.12.2 | What did the study conclude? | text box | |
| 7.13 | Did the qualitative study produce codes or themes? | y/n | |
| 7.13.1 | Describe the resulting codes/themes | text box | |
| 7.13.2 | What did the study conclude? | text box | |
| 7.14 | Was the study about electronic communication between physicians and pharmacists? | y/n | |
| 7.14.1 | How was this measured? | text box | |
| 7.14.2 | What did the study conclude? | text box | |

Methods Quality Assessment:

| Question | Methods Assessments | Options | Instructions |
|-----------|---|--|---|
| 6.1 | the study contains quantitative data? | yes/no | if yes, continue to next question, if no, no need to complete this form |
| 6.1.1 | Is the quantitative study a Randomized controlled trial? | Yes/no | Is the quantitative study a Randomized controlled trial? [info button “experimental design that studies the effect of an intervention or treatment using at least two groups: one that received the intervention and one that did not; participants ARE randomly assigned to a group (therapy, prevention)”]. |
| 6.1.1.1 | 1. Was the assignment to the treatment groups really random? | Yes/no. | look for methods of randomization (random number generator, coin flip etc) |
| 6.1.1.2 | 2. Was the treatment allocation concealed? | Yes/no. | This means that the people involved in getting the people into the study did not have any information or knowledge of what group the person might be going into. Look for things like: |
| 6.1.1.3 | 3. Were the groups similar at baseline in terms of prognostic factors? | Yes/no. | did the groups have similar characteristics at baseline? |
| 6.1.1.4 | 4. Were the eligibility criteria specified? | Yes/no. | Were there clear criteria for inclusion/exclusion for the study population? |
| 6.1.1.5 | 5. Were outcome assessors blinded to the treatment allocation? | Yes/no. | Were the people measuring or analyzing data blinded to what groups the data came from? |
| 6.1.1.6 | 6. Was the care provider blinded? | Yes/no. | Did the clinician know what group the study sample belonged to? |
| 6.1.1.7 | 7. Was the patient blinded? | Yes/no. | (were the people blinded to the group they were in?) |
| 6.1.1.8 | 8. Were the point estimates and measure of variability presented for the primary outcome measure? | Yes/no. | This is a 2 part question. The first--point estimates means that we have some sort of summary number like average minutes per prescription or rate of errors with the new/old system. The second issue is the measure of variability. Look for SDs (standard deviation measures usually in the form of mean 20 minutes per patient +-23 cm) or confidence intervals--for example mean 24 minutes (95% CI 21 to 25 minuets). |
| Not asked | 9. Did the analyses include an intention to treat analysis | Yes/no. | Look for the phrase “Intention to treat” or ITT. It refers to analyzing people as they were randomized. For example patients allocated to surgery would be analyzed within the surgery group even if they had been too sick, say, to get the surgery and got the drug instead). |
| | Sum quality score from the above 8 questions | (automatic summation of 9 items “yes”) | [don't worry about this] |
| 6.1.2 | If not an RCT, does the article report a cohort study? | Yes/No | Cohort study: <i>involves establishing groups, often people, one of which is “exposed” (e.g. HIT) and one is not exposed. Both groups followed forward in time to determine if the outcomes of interest develop.</i> |
| 6.1.2.1 | 1. Is there sufficient description of the groups and the distribution of prognostic factors? | Yes/no. | This question should be answered yes if you see a table of data on the study participants, usually Table 1 at the front end of the article. |

| Question | Methods Assessments | Options | Instructions |
|----------|--|---|--|
| 6.1.2.2 | 2. Are the groups assembled at a similar point in their disease progression? | Yes/no. | This question only applies to studies of people who have a condition. The question needs to state how long the people in each of the groups had the disease or condition in question--e.g. the mean number of years since diagnosis. |
| 6.1.2.3 | 3. Is the intervention/treatment reliably ascertained? | Yes/no. | Is there a description of the medication management or health information technology--answer yes if more than a paragraph on each. |
| 6.1.2.4 | 4. Were the groups comparable on all important confounding factors? | Yes/no. | Again look for data in Tables that gives data on how comparable the groups were. |
| 6.1.2.5 | 5. Was there adequate adjustment for the effects of these confounding variables? | Yes/no. | You will not likely see this information as it refers to adjustment in the analyses. Look for terms like adjustment, adjusted, regression analyses. |
| 6.1.2.6 | 6. Was a dose-response relationship between intervention and outcome demonstrated? | Yes/no. | You will not likely find this information so say no. |
| 6.1.2.7 | 7. Was outcome assessment blind to exposure status? | Yes/no. | This means that the people analyzing/assessing the data don't know which group the data came from --that way they are not biased thinking that people in this group should do better than the other group |
| 6.1.2.8 | 8. Was follow-up long enough for the outcomes to occur? | Yes/no. | Use common sense here--for example, were the errors assessed say within the first month, or 6 months of implementing a new system. Say no if there is not time for the intervention (or new system) to have an effect on the situation. |
| 6.1.2.9 | 9. Was the proportion of follow-up >80%? | Yes/no. | This question is looking for an assessment of the number of people who were initially enrolled into the study AND the number who were available for assessment at the end of the study. Also called follow up rate or proportion or just follow up. |
| 6.1.2.10 | 10. Were drop-out rates and reasons for drop-out similar across intervention and unexposed groups? | Yes/no. | This question is asking "why" did people drop out or why they were "lost" to the study. |
| | Sum quality score for above 10 questions | (automatic summation of 10 items "yes") | [this will not show in the interface, but be done by the computer system independently) |
| 6.1.3 | If not an RCT or cohort study, does the article report on a case control study? | Radio--start off | Case control study: <i>A study where groups of people are formed, one of which has the outcome of interest (e.g. better prescribing) and one of which does not (not better prescribing). Often members in the groups are "matched" in relation to things like age or experience. People in both groups are evaluated to assess if the exposure of interest (e.g. EHRs) were present in the past.</i> |
| 6.1.3.1 | 1. Is the case (people with the outcome) definition explicit? | Yes/no. | See the question and methods section to ascertain if the people who are the cases (e.g. those with errors in prescriptions) are described. |

| Question | Methods Assessments | Options | Instructions |
|----------|--|--|---|
| 6.1.3.2 | 2. Has the disease state of the cases been reliably assessed and validated? | Yes/no. | If a disease or disorder mentioned, did they tell how it was ascertained--often using rules or standard definitions. |
| 6.1.3.3 | 3. Were the controls randomly selected from the source of population of the cases? | Yes/no. | This is often hard to tell. Look for the word "random" or some mention of how the controls were selected. Often this will be no. |
| 6.1.3.4 | 4. How comparable are the cases and controls with respect to potential confounding factors? | Yes/no. | Look for Table 1 or in the first paragraph of the results section. If some information on the comparability of the groups is listed answer yes. |
| 6.1.3.5 | 5. Were interventions and other exposures assessed in the same way for cases and controls? | Yes/no. | Were measurements taken the same for the controls and the case groups? |
| 6.1.3.6 | 6. How was the response rate defined? | Yes/no/n/a | Not applicable |
| 6.1.3.7 | 7. Were the non-response rates and reasons for non-response the same in both groups? | Yes/no/n/a | Not applicable |
| 6.1.3.8 | 8. Is it possible that over-matching has occurred in that cases and controls were matched on factors related to exposure? | Yes/no/n/a | Not applicable |
| 6.1.3.9 | 9. Was an appropriate statistical analysis used (matched or unmatched)? | Yes/no/n/a | Not applicable |
| | Sum score | (automatic summation of 9 items "yes") | [this will not show in the interface, but be done by the computer system independently) |
| 6.1.4 | If not an RCT, cohort study or case control, does the article report on a case series? | Radio--start off | If not an RCT, cohort study or case control, does the article report on a case series?[info button "a medical research study that tracks patients with a known exposure given similar treatment or examines their medical records for exposure and outcome."] |
| 6.1.4.1 | 1. Is the study based on a representative sample selected from a relevant population? | Yes/no. | Answer yes if the article explains why and how these cases were chosen. |
| 6.1.4.2 | 2. Are the criteria for inclusion explicit? | Yes/no. | Answer yes if they list what the criteria for choosing the sites listed? |
| 6.1.4.3 | 3. Did all individuals enter the survey at a similar point in their disease progression? | Yes/no. | Answer yes if a disease is present and they provide any information on how long the disease had been diagnosed. |
| 6.1.4.4 | 4. Was follow-up long enough for important events to occur? | Yes/no. | Likely not applicable . |
| 6.1.4.5 | 5. Were outcomes assessed using objective criteria or was blinding used? | Yes/no. | The outcomes such as error rates need to be assessed in a blinded manner to suit methodologists. Look for the terms blind:, mask:, placebo:, etc. |
| 6.1.4.6 | 6. If comparisons of sub-series are being made, was there sufficient description of the series and the distribution of prognostic factors? | Yes/no . | Answer yes if any information is given. If the analysis of the cases was down broken down into subcategories, such as men and women, children and adolescents, young or old.... |

| Question | Methods Assessments | Options | Instructions |
|----------|--|--|--|
| | Sum score | (automatic summation of 6 items "yes") | [[this will not show in the interface, but be done by the computer system independently) |
| 6.1.5 | If not any of the above, is this a before-after study? | Yes/No | A before-after study will have measures taken before and after implementation of a change. |
| 6.1.6 | If not any of the above, is the study a time-series? | Yes/No | |
| 6.1.7 | If not any of the above, is this an observational study? (can include a case study with data, survey etc). | Yes/No | An observational study is one where the researchers have no control over exposures and instead observe what happens to groups of people. |

Appendix C. Evidence Tables

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management

| Article Information | HIT Studied | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|--|---------|
| Abboud (2006) (Abboud et al. 187-198) Design: Before-after N = 336 orders Implementation: 04/2002 Study Start: 10/2003 Study End: 03/2004 | Integrated system CDSS/CDS/CCDS/reminders CPOE/POE system Integrated CDSS/CDS/CCDS/reminders EHR/EMR system, Formulary, Hospital information system, Imaging systems, Laboratory system, Pharmacy | Pediatric stand alone hospital, 423 Beds | antibiotics courses with no lab order* | no significant differences between the baseline and the corollary order periods on courses of antibiotics associated with no laboratory monitoring 31 (19.5%) vs. 31(17.5%), p = NS. | - |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: A1c = hemoglobin A1c; ACE = Angiotensin Converting Enzyme; ACEI = Angiotensin-Converting Enzyme Inhibitor; ADEs = Adverse Drug Events; ALT = Alanine Aminotransferase; AMI = Acute Myocardial Infarction; AR = Absolute Reduction; ARB = Angiotensin-II-Receptor Blocker; ARI = Acute respiratory infection; AST = Aspartate Aminotransferase; CC = Care Considerations; CCDS = Computerized Clinical Decision Support; CDS = Clinical / Computerized Decision Support; CDSS = Clinical Decision Support System; CHF = Congestive Heart Failure; CI = Confidence interval; CIT = Clinical information technology; COPD = Chronic Obstructive Pulmonary Disease; CPG = Clinical Practice Guidelines; CPOE = Computerized Provider Order Entry; DDI = Drug-drug Interaction; DS = Decision Support; DSS = Decision Support System; ED = Emergency Department; EHR = Electronic Health Record; e-MAR = Electronic Medication Administration Record; EMR = Electronic Medical Records; EP = Electronic Prescribing; e-RX = Electronic Prescribing; e-TAR = Electronic Treatment Authorization Request; GP = General Practitioner; h = Hour; HIT = Health Information Technology; HIV = Human Immunodeficiency Virus; hr = Hour; hrs = Hours; ICU = Intensive Care Unit; K = Potassium; LVSD = Left Ventricular Systolic Dysfunction; ME = Medication Error; Mg = Magnesium; min = Minute; MMR = Measles; Mumps and Rubella; N = Sample Size; n/a = Not Applicable; Np = Nurse Practitioner; NR = not reported; NS = NS; NSAID = Nonsteroidal anti-inflammatory drug; NSAIDS = Nonsteroidal anti-inflammatory drugs; OR = Odds ratio; OSUH = Ohio State University Health System; p = Probability; PCA = Patient-Controlled Analgesia; PDA = Personal Digital Assistants; PICU = Pediatric Intensive Care Unit; POE = Provider Order Entry; PONV = Postoperative Nausea and Vomiting; PRN = *pro re nata*; RCT = Randomized Controlled Trial; RR = Relative Risk; RRR = Relative Risk Reduction; RV = rule violation; UDDS = Unit Dose Drug Dispensing System; UTI = Urinary tract Infection; vs. = Versus; VTE = Venous Thromboembolism

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|---|---|----------|
| <p>Achtmeyer (2002) (Achtmeyer, Payne, and Anawalt 277-281) Design: Before-after N = 1,405 orders for supplemental insulin Implementation: 12/1998 Study Start: 12/1998 Study End: 07/1999</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated EHR/EMR system, Imaging systems, Laboratory system</p> | <p>Acute care/tertiary, 290 Beds Academic</p> | <p>rate of traditional sliding scale orders for supplemental insulin*</p> | <p>rate of traditional sliding scale orders for supplemental insulin in hospitalized patients was reduced when a quick-order CPOE/CDSS system was put in place (97.1% vs. 63.8%, RRR 34%, p <0.001).</p> | <p>+</p> |
| <p>Agostini (2007) (Agostini, Shang, and Inouye 43-48) Design: Before-after N = 24,509 patients Implementation: 04/2002 Study Start: 04/2002 Study End: 03/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system, Formulary</p> | <p>Acute care/tertiary, 944 Beds Academic</p> | <p>rate of prescribing of sedative-hypnotics*</p> | <p>Prescribing of sedative-hypnotics decreased from 2,208 per 12,356 (18%) patients preintervention to 1,832 per 12,153 (15%) postintervention (OR for the intervention = 0.82, 95% CI = 0.76–0.87), an 18% risk reduction (p <0.001 for pre/post difference).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|----------------------------------|--|---|--|----------|
| <p>Ali (2005) (Ali et al. 110-114) Design: Before-after N = 91 patients Implementation: 02/2000 Study Start: 05/2000 Study End: 05/2002</p> | <p>CPOE/POE system</p> | <p>Critical care units (CCU, ICU, NICU) 25 Beds Academic</p> | <p>mean number of orders for vasoactive drips per patient, mean number of orders for sedative infusions per patient</p> | <p>Compared to the initial CPOE, the redesign of the CPOE system to incorporate more complex order sets resulted in significantly fewer orders placed per patient (means) for vasoactive drips (4.8 vs. 2.2, p <0.01) and sedative infusions (6.4 vs. 2.9, p <0.01), as a measure of improved workflow efficiency.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|---|---|----------|
| <p>Bailey (2007) (Bailey et al. 586-590). Design: RCT N = 853 patients Implementation: 00/0000 Study Start: 02/2000 Study End: 05/2001</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system, Laboratory system</p> | <p>Acute care/tertiary, 1,385 Beds Inpatient hospital based, Academic</p> | <p>compliance rates: - patients discharged on a full complement regimen of secondary prevention medications* -ACE inhibitor*, -statins*, -aspirin -beta-blockers.</p> | <p>When individual drug class exclusions were considered, compliance rates increased for patients discharged on a full-complement regimen of secondary prevention medications (70.3% vs. 83.6%, RRR - 19%, p <0.001). Compliance rates for ACE inhibitor (83.6 vs. 89.9, RRR - 8%, p = 0.01) and statin use (89.3 vs. 94.2%, RRR - 5%, p = 0.02) were significantly higher, while rates for aspirin (96.5% vs. 96.4%, RRR 0%, p = 0.95) and beta-blockers (91.8% vs. 95.9%, RRR - 5%, p = 0.08) remained the same.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|---|----------|
| <p>Bates (1999) (Bates et al. 313-321) Design: Time series N = 1,817 admissions Implementation: 05/1993 Study Start: 10/1992 Study End: 04/1997</p> | <p>CDSS/CDS/CCDS/reminders Integrated Billing/administration system Hospital information system</p> | <p>Acute care/tertiary 700 Beds Academic</p> | <p>Rate of non-missed dose errors per 1,000 patient-days over 4 time periods*, Rate of non-missed dose errors per admission*</p> | <p>The rate of errors (other than missed dose) per 1000 patient-days fell from baseline across all time points for medication errors: non-missed-dose medication errors (142, 51.2, 74, 2666; p = 0.0001). The results were similar for non-missed-dose error rate per admission (0.64, 0.27, 0.28, 0.11, p = 0.0001). Non-intercepted serious medication errors declined significantly over time (7.6, 7.3, 1.7, 1.1, p = 0.0003).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|-----------------------------|---|--|----------|
| <p>Bates (1994) (Bates, Boyle, and Teich 996) Design: Before-after N = 62 Physicians (Medical Interns and 1st and 2nd year surgical residents) Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CPOE/POE system Integrated Hospital information system</p> | <p>Unspecified Hospital</p> | <p>Time spent ordering by medical interns*, Time spent ordering by surgical residents*, Time spent on daily and one-time orders*, Time spent on sets of orders*</p> | <p>When time spent ordering was compared between pre-order entry and post-order entry periods, the percent for medical interns increased from 5.3% to 10.5% (p <0.001) representing 44 additional minutes per day, while for surgical house officers the corresponding figures were an increase from 6.4% to 15.5% (p <0.001), 73 minutes per day. Daily and one-time orders accounted for the majority of this change, increasing almost threefold in percent total time (2.2% before, vs. 7.2% after order entry). However, sets of orders took less total time after order entry (1.7% vs. 3.1%).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|----------|
| <p>Bates (1998) (Bates et al. 1311-1316) Design: RCT N = 4,220 patients Implementation: 00/0000 Study Start: 02/1993 Study End: 07/1995</p> | <p>CPOE/POE system Integrated Billing/administration system, Formulary, Hospital information system</p> | <p>Acute care/tertiary, 726 Beds Academic</p> | <p>the rate of nonintercepted serious medication errors/1,000 patient days -phase 1 to 2*, the rate of nonintercepted serious medication errors/1,000 patient days -CPOE vs. CPOE+team, Transcription errors</p> | <p>In paired analyses comparing phase 1 and phase 2 (Table 2), the rate of nonintercepted serious medication errors fell 55%, from 10.7 events per 1,000 patient-days to 4.86 events (p = 0.01). For the RCT in the post-CPOE phase, comparing CPOE alone with CPOE plus team showed no significant difference in error rates (4.81 vs. 6.01, p = 0.49). Transcription errors (CPOE to paper in pharmacy) fell 84%, p <0.001.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--------------------------------------|---|---|----------|
| <p>Bell (2010) (Bell et al. e770-e777) Design: RCT N = 19,450 patients Implementation: 00/0000 Study Start: 04/2007 Study End: 04/2008</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care, Academic</p> | <p>proportion of children with asthma having at least 1 prescription for controller medication*, proportion of children with asthma having an up-to-date asthma care plan*, proportion of children with asthma having spirometry performed*</p> | <p>Increases in the number of prescriptions for controller medications, over time, was 6% greater (p = 0.006) and 3% greater for spirometry (p = 0.04) in the intervention urban practices. Filing an up-to-date asthma care plan improved 14% (p = 0.03) and spirometry improved 6% (p = 0.003) in the suburban practices with the intervention.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|--|----------|
| <p>Berner (2006) (Berner et al. 171-179) Design: RCT N = 59 internal medicine residents Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Handheld</p> | <p>Ambulatory care, Academic</p> | <p>proportion of unsafe NSAID prescribing per physician at followup</p> | <p>The proportion of cases per physician with unsafe NSAID prescriptions were similar at baseline for control (0.29) and intervention residents (0.27). At followup, the rates were statistically different, with lower proportions for intervention residents after adjustment for baseline rates (0.45 control vs. 0.23 intervention, p <0.05). Note that the control group prescribing degraded over time while the intervention group was stable.</p> | <p>-</p> |
| <p>Bernstein (2005) (Bernstein et al. 225-231) Design: Before-after N = 1,158 prescriptions Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system</p> | <p>Emergency department, Academic</p> | <p>percentage of proprietary antibiotics prescribed*</p> | <p>The percentage of proprietary antibiotics prescribed before and after insertion of the electronic prompt was 26.6% vs. 20.7%, RRR 22%, p = 0.03.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|------------------------|--|---|----------|
| <p>Bertoni (2009) (Bertoni et al. 678-686) Design: RCT N = 8,878 patients Implementation: 00/0000 Study Start: 06/2001 Study End: 04/2006</p> | <p>CDSS/CDS/CCDS/reminders Handheld</p> | <p>Ambulatory care</p> | <p>adherence to guideline-screening*, adherence to guideline-appropriate lipid management*</p> | <p>There was no difference in screening rates between the CDSS-PDA group and the control. The control group had a 10.8% drop in appropriate management from baseline, while the PDA group had a 1.1% drop, p <0.01. Stable adherence was observed in the PDA intervention group, whereas a decline in guideline adherence was observed in the control group.</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|----------|
| <p>Bloomfield (2005) (Bloomfield et al. 258-263) Design: RCT N = 9,105 patients Implementation: 04/2002 Study Start: 10/2001 Study End: 10/2003</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>rate of prescription lipid therapy (before-after)</p> | <p>rate of lipid therapy prescriptions increased significantly after implementation of the prompts in the intervention clinics (8.3% vs. 39.1%, RRR -371, p <0.0001) no statistically significant difference in prescription rates (40.7% for progress notes, 36.9% for patient letters, and 39.4% for reminders, p = 0.60) alternative logistic regression analysis, significant interaction between group and site, indicating that the efficacy of the prompts differed by site.</p> | <p>+</p> |
| <p>Bogucki (2004) (Bogucki et al. 278-280) Design: Before-after N = 2,124 orders for parenteral corticosteroids Implementation: 04/2002 Study Start: 04/2003 Study End: 07/2003</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CDSS/CDS/CCDS/ reminders CPOE/POE system EHR/EMR system, e-MAR</p> | <p>Pediatric stand alone hospital, 324 Beds</p> | <p>rate of methylprednisone ordering*</p> | <p>There was a significant reduction in methylprednisone prescribing following the implementation of the alert in relation to the total number of parenteral corticosteroids ordered (21.5% vs. 9.7%, RRR 55%, p <0.0001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|---|---------|
| Bouaud (2001) (Bouaud et al. 1-4) Design: Before-after N = 127 decisions/orders Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Ambulatory care Other | rate of compliance with CPG | Before using OncoDoc, physicians compliance with CPG was 61.42%. Using the system significantly increased actual compliance to 85.03% (p <0.0001). | + |
| Buising (2008) (Buising et al. 35) Design: Time series N = 740 patients Implementation: 01/2005 Study Start: 04/2003 Study End: 09/2006 | CDSS/CDS/CCDS/reminders Integrated Hospital information system | Acute care/tertiary, 350 Beds Academic | proportion of patients receiving appropriate antibiotic therapy* | proportion of patients receiving appropriate antibiotic therapy increased significantly between each time period (61.9% baseline vs. 68.7 academic detailing vs. 89.7 CDSS, pairwise comparisons p <0.01) associated ORs for having received the recommended empiric antibiotic therapy were 2.58 between baseline and CDSS periods and 2.03 between academic detailing and CDSS. | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|----------|
| <p>Butler (2006) (Butler et al. 643-653) Design: Before-after N = 1,827 patients (1,251 with CHF and 576 with AMI) Implementation: 07/2002 Study Start: 07/2001 Study End: 0.9/2003</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system</p> | <p>Acute care/tertiary, Academic</p> | <p>compliance rate: ACEI for LVSD*, compliance rate: ACEI for AMI*, compliance rate: aspirin for AMI*, compliance rate: beta-blocker for AMI*</p> | <p>Aspirin (95% vs. 95%, RRR 0%, NS), betablocker (88% vs. 95%, RRR -8%, NS), and ACEI (77% vs. 81%, RRR -5%, NS) use for AMI patients at the time of discharge in the pre-CPOE era was high and remained so in the CPOE period. Similarly for ACEI for CHF patients (74% vs. 87%, RRR -18%, NS). When examining indicators in the post-CPOE phase, rates were higher in patients for which the tool was used, vs. not used for all 4 medication related indicators (p <0.001).</p> | <p>-</p> |
| <p>Chertow (2001) (Chertow et al. 2839-2844) Design: Time series N = 19,982 admissions Implementation: 00/0000 Study Start: 09/1997 Study End: 04/1998</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Hospital information system Imaging systems</p> | <p>Acute care/tertiary, 720 Beds Academic</p> | <p>rate of appropriate prescribing*, rate of appropriate prescribing involving dosage alterations*, rate of appropriate prescribing involving frequency alterations*</p> | <p>The rate of appropriate prescribing was increased with CPOE/CDSS for all orders (51% intervention vs. 30% control, RRR 70%, p <0.001) by dose (67% vs. 54%, RRR 43%, p <0.001), or by frequency (59% vs. 35%, RRR 69%, p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|---|--|----------|
| <p>Chisholm (2003) (Chisholm et al. 199-206) Design: Before-after N = 790 children admitted to hospital with asthma exacerbations Implementation: 10/2002 Study Start: 11/2001 Study End: 12/2003</p> | <p>CPOE/POE system Integrated Billing/administration system, EHR/EMR system, Laboratory system</p> | <p>Pediatric stand alone hospital, 323 Beds</p> | <p>systemic corticosteroids use*, metered-dose inhaler use*</p> | <p>More use was made of systemic corticosteroids (OR 5.61, 95% CI 3.46 to 9.11) and of metered-dose inhalers (OR 1.42, CI 1.04 to 1.94) after implementation of standard order sets in the CPOE for asthma patients.</p> | <p>+</p> |
| <p>Choi (2004) (Choi et al. 1-6) Design: Before-after N = 307 patients Implementation: 02/2003 Study Start: 12/2002 Study End: 04/2003</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Error rates per patient*</p> | <p>Error rates per patient significantly declined in the intervention site following implementation of the nurse CPOE with CDSS (17.4% vs. 3.1%, RRR 82%, p = 0.0075). In the control group, error rates remained unchanged (8.6% vs. 6.9%, NS). At baseline, the control group rate was statistically lower than the intervention group (8.6% vs. 17.4%, p = 0.04).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|----------------------------------|--|--|---------|
| Christakis (2001) (Christakis et al. e15) Design: RCT N = 38 providers Implementation: 00/0000 Study Start: 03/0000 Study End: 05/0000 | CDSS/CDS/CCDS/ reminders Integrated online prescription writer | Ambulatory care, Academic | change in the frequency of antibiotic prescription* | For the primary outcome, providers in the intervention arm had a 44% change in the frequency with which they prescribed antibiotics for <10 days, whereas providers in the control arm had a 10% change, this change in behavior was significantly related to the intervention, although both groups improved (p <0.01). | + |
| Clancy (1992) (Clancy, Gelfman, and Poses 14-18) Design: Before-after N = 1,013 patients Implementation: 02/1985 Study Start: 11/1984 Study End: 05/1985 | CDSS/CDS/CCDS/ reminders Integrated Hospital information system | Acute care/tertiary, Academic | pneumococcal vaccination rate per admission* | Preimplementation of the reminder pneumococcal vaccination rate was 3.4% compared 45% post (p <0.0001). | + |
| Cobos (2005) (Cobos et al. 421-432) Design: RCT N = 2,221 patients Implementation: 04/2000 Study Start: 04/2000 Study End: 05/2002 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | proportion of patients prescribed lipid lowering drugs (secondary) | The proportion of patients prescribed lipid lowering drugs was significantly lower in the CDSS guideline intervention group (59.1% vs. 40.8%, RRR 31%, p <0.0001). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|--|--|----------|
| <p>Colpaert (2006) (Colpaert et al. R21) Design: RCT N = 2,510 prescriptions Implementation: 00/0000 Study Start: 03/2004 Study End: 04/2004</p> | <p>CPOE/POE system Integrated Billing/administration system, CPOE/POE system, Hospital information system, Laboratory system</p> | <p>Acute care/tertiary, Critical care units (CCU, ICU, NICU) 22 bed unit Beds Academic</p> | <p>rate of medication prescribing errors*, minor MPEs*, Intercepted MPEs *, Serious MPEs *</p> | <p>The incidence of MPEs was significantly lower in the computerized unit (C-U) compared with the paper based unit (PBU) [44/1,286 (3.4%) vs. 331/1,224 (27.0%); p <0.001]. There were significantly fewer minor MPEs in the C-U than in the PB-U [9 (0.7%) vs. 225 (18.4%); p <0.001]. Intercepted MPEs were also lower in the C-U [12 (0.9%) vs. 46 (3.8%); p <0.001]. Serious MPEs were also lower in C-U than PBU [23 (1.8%) vs. 60 (4.9%), p <0.001].</p> | <p>+</p> |
| <p>Cordero (2004) (Cordero et al. 88-93) Design: Before-after N = 211 infants Implementation: 02/2000 Study Start: 10/2001 Study End: 09/2002</p> | <p>CPOE/POE system Integrated Imaging systems, Pharmacy</p> | <p>Acute care/tertiary, Critical care units (CCU, ICU, NICU) Academic</p> | <p>medication turn-around times-caffeine*, medication error rate-gentamicin</p> | <p>The turn-around times for the pre- and post-CPOE loading dose of caffeine were 10.5 ± 9.8 and 2.8 ± 3.3 hours p <0.01, respectively. In the pre-CPOE period, there were 14 (13%) gentamicin prescription dosage errors, in the post-CPOE period there were 0</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|--|----------|
| <p>Cote (2008) (Cote et al. 1097-1103) Design: Before-after N = 601 adult patients Implementation: 00/0000 Study Start: 12/2005 Study End: 06/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Unspecified Hospital</p> | <p>rate of gastroprotection at discharge*, control vs. physician education vs. alert vs. alert plus education</p> | <p>The study sought the change in rate of gastroprotection at discharge for all patients; changes only occurred for the group that had both education and alerts compared to control (43% vs. 61%, RRR - 42%, p <0.001). Education alone (42%) or alerts alone (39%) did not change rates of gastroprotection.</p> | <p>-</p> |
| <p>Cunningham (2008) (Cunningham, Geller, and Clarke 546-554) Design: Before-after N = 1,040 order sets Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CPOE/POE system Integrated CDSS/CDS/CCDS/reminders</p> | <p>Acute care/tertiary, General Hospital 667 Beds</p> | <p>compliance to medication order sets*, minutes to first dose of antibiotics</p> | <p>Medication orders placed using CPOE were significantly more compliant with hospital protocols (80%) than paper based medication orders at both the CPOE hospital (63%) and the control hospital (64%), and first doses of antibiotics were delivered significantly faster when ordered with CPOE (180 min) than when placed using the standard paper-based system (326 min, p <.01).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|--------------------------------------|--|--|----------|
| <p>Davis (2007) (Davis et al. e25) Design: RCT N = 44 health care providers Implementation: 11/1999 Study Start: 11/1999 Study End: 12/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system</p> | <p>Ambulatory care, Academic</p> | <p>changed physician behavior in accordance with the intervention message screens*</p> | <p>Prescribing behavior in accordance with the evidence improved only marginally, by 1% in control group and 4% in the intervention group (absolute difference 3%, 95% CI 1%, 15%).</p> | <p>+</p> |
| <p>de Jong (2009) (de Jong et al. 9-20) Design: Cross-sectional N = 749,811 contacts Implementation: 00/1998 Study Start: 01/2001 Study End: 12/2001</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>proportion of prescriptions in accordance with DSS*, Herfindahl-Hirschman Index</p> | <p>GPs who use the DSS daily prescribe more according to the advice given in the DSS (89%) than GPs who do not use the DSS (75%, RRR 19%, p = 0.04). There was no significant difference between the Herfindahl-Hirschman Index for both groups (40.3 for daily users and 41.4 for non users, p = 0.3) the variation in prescriptions for a given diagnoses was comparable between groups.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|------------------------------|--|---|---------|
| Devine (2010) (Devine et al. 928) Design: Before-after N = 10,169 prescriptions Implementation: 03/2003 Study Start: 03/2002 Study End: 04/2006 | CDSS/CDS/CCDS/reminders CPOE/POE system Integrated EHR/EMR system | Ambulatory care | errors | Frequency of errors declined from 18.2% (Pre-CPOE) to 8.2% (post-CPOE), a reduction in adjusted odds of 70% (OR: 0.30; 95% CI 0.23 to 0.40), p <0.001. | + |
| Dexter (2001) (Dexter et al. 965-970) Design: RCT N = 3,416 patients Implementation: 00/0000 Study Start: 05/1997 Study End: 10/1998 | CDSS/CDS/CCDS/reminders Integrated Pharmacy | General Hospital Academic | proportion compliance: -pneumococcal vaccination*, -influenza vaccination* -subcutaneous heparin -aspirin at discharge | The use of the reminders led to a higher ordering rate all 4 preventive therapies for eligible patients; pneumococcal vaccination (0.8% vs. 35.8%, RRR - 4375%, p <0.001), influenza vaccination (1.0% vs. 51.4%, RRR - 5040%, p <0.001), subcutaneous heparin (18.9% vs. 32.2%, RRR -70%, p <0.001) and aspirin at discharge (27.6% vs. 36.4%, RRR - 32%, p <0.001). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|---|----------|
| <p>Dexter (2004) (Dexter et al. 2366-2371) Design: RCT N = 1,677 patients Implementation: 11/1998 Study Start: 11/1998 Study End: 12/1999</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system</p> | <p>General Hospital, Academic</p> | <p>rate of receipt of vaccination -influenza*, rate of receipt of vaccination - pneumococcal*</p> | <p>Patients in the standing order group received both vaccinations more often than patients in the pop-up reminder group; for the influenza vaccine 30% reminder vs. 42% standing order, p <0.001; for the pneumococcal vaccine 51% vs. 31%, p <0.001.</p> | <p>+</p> |
| <p>Durieux (2000) (Durieux et al. 2816-2821) Design: Time series N = 1,971 patients Implementation: 00/0000 Study Start: 12/1997 Study End: 07/1999</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, Hospital information system</p> | <p>Acute care/tertiary, 1,000 Beds Academic</p> | <p>rate of appropriate anticoagulant prescribing*</p> | <p>Physicians complied with guidelines in 82.8% of cases during control periods and in 94.9% of cases during intervention periods (RRR - 15%, p <0.001). During each intervention period, the proportion of appropriate prescriptions ordered increased significantly. Each time the CDSS was removed, physician compliance with guidelines reverted to that observed before initiation of the intervention.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|---|---|----------|
| <p>Eslami (2006) (Eslami et al. 803-809) Design: Cross-sectional N = 392 orders Implementation: 00/2002 Study Start: 05/2002 Study End: 12/2004</p> | <p>CPOE/POE system Integrated EHR/EMR system, Laboratory system</p> | <p>Acute care/tertiary, Critical care units (CCU, ICU, NICU) 28 in 3 units Beds</p> | <p>Dosing error*</p> | <p>The dose was wrong (i.e. there was >10% deviation from the guideline) in 73% (165/227) of the orders that used the default value (essentially as suggested by the CPOE) and in 77% (127/165) of the orders in which the default value was not administered (p = 0.4).</p> | <p>-</p> |
| <p>Evans (1998) (Evans et al. 232-238) Design: Before-after N = 1,681 patients Implementation: 00/0000 Study Start: 07/1992 Study End: 06/1995</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Imaging systems, Laboratory system</p> | <p>Acute care/tertiary, 520 Beds Academic</p> | <p>mean number of days with excessive antibiotic dosing*, usage rate of antiinfectives*</p> | <p>During the intervention period, there were significantly fewer days when doses of antiinfective agents were excessive than during the preintervention period (2.7 days vs. 5.9 days per patient, respectively; p <0.002). There was an increase in the use of antiinfectives following the intervention reminder (67% vs. 73%, RRR 9%, p <0.03).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|-------------------------------------|---|---|----------|
| <p>Evans (1990) (Evans et al. 351-354) Design: Before-after N = 7,656 patients Implementation: 00/0000 Study Start: 06/1985 Study End: 09/1986</p> | <p>CDSS/CDS/CCDS/reminders Hospital information system Integrated Laboratory system, Pharmacy</p> | <p>Unspecified Hospital</p> | <p>mean number of antibiotic doses per patient, proportion of patients receiving preoperative antibiotics, proportion of patients receiving antibiotics for too long,</p> | <p>Surgical patients received an average of 19 antibiotic doses before implementation of the implementation of the 'stop orders' and 13 after (p <0.001). There were non significant changes in the proportion of patients receiving preoperative antibiotics (64% vs. 66%, NS) or those receiving antibiotics for too long (40% vs. 35%, NS).</p> | <p>+</p> |
| <p>Evans (1994) (Evans et al. 878-884) Design: RCT N = 482 cultures Implementation: 00/000 Study Start: 07/1990 Study End: 01/1991</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Laboratory system</p> | <p>Acute care/tertiary 520 Beds</p> | <p>rate of prescribing antibiotics to which all of the isolated pathogens were susceptible</p> | <p>The computer group had a higher rate of prescribing antibiotics to which all of the isolated pathogens were susceptible (77% vs. 94%, RRR 22%, p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|------------------------|--|--|----------|
| <p>Feldstein (2006) (Feldstein et al. 1009-1015) Design: RCT N = 9,910 patients with 239 care providers in 15 primary care clinics Implementation: 12/2002 Study Start: 01/2000 Study End: 08/2004</p> | <p>CDSS/CDS/CCDS/reminders Integrated CDSS/CDS/CCDS/reminders CPOE/POE system, EHR/EMR system</p> | <p>Ambulatory care</p> | <p>interacting prescription rate (/10,000 warfarin users/month), slope for interacting prescription rate</p> | <p>When baseline trends were controlled for, the overall interacting prescription rate decreased immediately after the alerts were implemented, with an estimated reduction of 329.7 interacting prescriptions per 10,000 warfarin users in the first month (p = 0.002). The alerts also significantly changed the trend in the interacting prescription rate, with a preintervention increasing rate of 1.1 and a postintervention decreasing rate of 21.3 (slope change -22.4; p = 0.01). Academic detailing did not have an effect on interacting prescription rates.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|------------------------|--|--|----------|
| <p>Feldstein (2006) (Feldstein et al. 450-457) Design: RCT N = 311 women Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Laboratory system</p> | <p>Ambulatory care</p> | <p>rate of completion of BMD or medication for osteoporosis, The same pattern was evidence for medication only</p> | <p>control group had fewer women who had BMD completer or medication for osteoporosis compare with the reminder and reminder plus education groups (5.9% control, 51.5% reminders, and 33% reminders and education, p <0.01 for both comparisons with control RRR for reminders alone 690% and RRR for reminders and education 460%). The same pattern was evidence for medication only (5.0% control, 27.7% reminders and 20.2% reminders plus education; p <0.01 for comparisons with control.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---------------------------------------|---|--|----------|
| <p>Field (2009) (Field et al. 480-485) Design: RCT N = 833 patients (10 physicians and 213,967 patient days) Implementation: 00/000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated EHR/EMR system</p> | <p>Long term care (nursing homes)</p> | <p>proportion of appropriate orders*, proportion of inappropriate drugs avoided</p> | <p>The proportion of appropriate antidepressant order rates for patients with renal insufficiency was higher in the CDSS group (52% vs. 63%, OR 1.2, 95% CI 1.0 to 1.4). More inappropriate drugs were avoided (15% vs. 46%, OR 2.6, CI 1.4 to 5.0). Improvements were seen in frequency and missing information but not for doses in the CDSS group.</p> | <p>+</p> |
| <p>Fiks (2009) (Fiks et al. 159-169) Design: RCT N = 22,586 patients Implementation: 00/0000 Study Start: 10/2006 Study End: 05/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system, EHR/EMR system</p> | <p>Ambulatory care, Academic</p> | <p>rates of up-to-date influenza vaccination*, rates of captured opportunities for vaccination*</p> | <p>Rates of up-to-date influenza vaccination increased from 44.2% to 48.2% (control) and from 45.0% to 53.0% (intervention), a 4.0% (95% CI: -1.3% to 9.1%) greater but NS. Overall rates of captured opportunities for vaccination increased 3.8% (12.3% to 16.1%) control practices and 4.8% (14.4% to 19.2%) intervention sites, difference 1% (95% CI: -2.4% to 4.9%).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|---|---|----------------|
| Filippi (2003) (Filippi et al. 1497-1500) Design: RCT N = 15,343 patients Implementation: 00/0000 Study Start: 05/2001 Study End: 11/2001 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system, EHR/EMR system | Ambulatory care | Antipletlet drug treatment | number of treated patients significantly increased in the intervention group (OR 1.99, 95% CI 1.79 to 2.22). | + |
| Fischer (2003) (Fischer et al. 2585-2589) Design: Before-after N = 1,045 orders Implementation: 00/00 Study Start: 00/00 Study End: 00/00 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system | Acute care/tertiary, Inpatient hospital based Academic | defined daily dose -IV, defined daily dose –oral DDD | After implementation the use of IV medication (DDD) decreased by 11.1%, p = 0.002 and the oral drug use (DDD) increased by 3.7%, p = 0.002. | + |
| Fischer (2008) (Fischer et al. 2433-2439) Design: Before-after N = 12,625,276 prescriptions Implementation: 10/2003 Study Start: 10/2003 Study End: 5/2005 | e-Rx Integrated Formulary, Insurance | Not specified | rates of prescribing, tier 1*, rates of prescribing, tier 2*, rates of prescribing, tier 3* | 20% of prescriptions written by intervention physicians completed using e-Rx intervention group prescribed 1.4% more (95% CI, 0.6% to 2.0%) tier 1 medications, 0.3% fewer (95% CI, -0.8% to 0.2%) tier 2 medications, and 1.0% fewer (95% CI, -1.4% to -0.7%) tier 3 medications than the control group. | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|------------------------|---|---|----------|
| <p>Flottorp (2002) (Flottorp et al. 367) Design: RCT N = 26,826 Consultation Implementation: 00/0000 Study Start: 01/2000 Study End: 01/2001</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Use of antibiotics for sore throat, use of antibiotics for UTI</p> | <p>sore throat group 3% less likely to receive antibiotics after the intervention (49.5% vs. 43.8%, p = 0.032) UTI (43.4% vs. 46.3%, p = 0.639) Women with symptoms of UTI in the intervention group were 5.1% less likely to have a laboratory test ordered (55% vs. 49.8%, p = 0.046) For sore throat, the numbers were 39.7% vs. 42.0%, p = 0.638 proportion of telephone consultations sore throat: 1.2% greater in the control group than in the intervention group (14.1% vs. 12.9%, p = 0.128) proportion decreased for UTI (18.9% vs. 19.8%, p = 0.874)</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|--|----------|
| <p>Fontan (2003) (Fontan et al. 112-117) Design: Cross-sectional N = 4,532 prescriptions Implementation: 00/1988 Study Start: 02/1999 Study End: 03/1999</p> | <p>Computerized unit dose drug dispensing system (UDDS) Integrated Hospital information system</p> | <p>Other specialty hospital (rehab, oncology) Pediatric stand alone hospital 510 Beds</p> | <p>prescribing error rate administering error rate</p> | <p>Errors were decreased with the use of the eRX and computerized dispensing system compared with the hand-written prescriptions and ward distribution system. Prescribing errors were reduced from 87.9% to 10.6%, RRR 88%, p <0.00001. Administrative errors with time errors were reduced from 29.3% to 22.5%, RRR 23%, p <0.001.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|--------------------------------------|---|---|----------|
| <p>Fortuna (2009) (Fortuna et al. 897-903) Design: RCT N = 257 clinicians Implementation: 00/1997 Study Start: 03/2006 Study End: 03/2008</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, e-Rx</p> | <p>Ambulatory care, Academic</p> | <p>relative risk of prescribing heavily marketed medications*</p> | <p>The relative risk of prescribing heavily marketed medications in the alert-group during the intervention period was less than in the usual-care group (RRR 0.74; 95% CI 0.57 to 0.96; p = 0.02). The RR of prescribing heavily marketed hypnotics in the alert-plus-education group was less than in the usual-care group (RRR 0.74; 95% CI 0.58 to 0.97, p = 0.03). The prescribing of heavily marketed medications was similar in the alert-only group and the alert-plus-education group (RRR 1.02; 95% CI 0.80 to 1.29; p = 0.90).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|-----------------------------|---|---|----------|
| <p>Frances (2001) (Frances et al. 165-166) Design: RCT N = 63 physicians and 730 patients Implementation: 00/0000 Study Start: 03/1997 Study End: 06/1997</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Pharmacy</p> | <p>Ambulatory care</p> | <p>Receiving aspirin*, History of MI and receiving beta-blocker*, Receiving cholesterol-lowering agent*</p> | <p>proportion of patients had an active prescription for aspirin 37.9% vs. 35.1%, RRR 7%, p = 0.440, NS; proportion of patients with MI who had an active beta-blocker prescription 22.2% vs. 33.3%, RRR - 50%, p = 0.465, NS; proportion of patients receiving a cholesterol-lowering agent (73.2 % vs. 71.0%, RRR - 15%, p = 0.512)</p> | <p>-</p> |
| <p>Frank (2004) (Frank, Litt, and Beilby 87-90) Design: RCT N = 10,507 patients Implementation: 00/0000 Study Start: 03/1998 Study End: 03/1999</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>proportion of opportunities taken for preventive activity*</p> | <p>Reminders did not improve adherence to MMR and flu vaccinations, but there was a significant increase in tetanus immunization (1.5% vs. 2.8%, relative change 1.89, 95% CI 1.59, 2.25). and pneumococcal immunization rates (1.6% vs. 2.8%, relative change 1.70, 95% CI 1.10, 2.62). Two of 8 non-medication related preventive care recommendations were significantly improved as well.</p> | <p>+</p> |
| <p>Franklin (2007) (Franklin et al.</p> | <p>Automated Dispensing</p> | <p>Acute care/tertiary,</p> | <p>error rate for new prescriptions*, error rate</p> | <p>The prescription</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|---|---------|
| <p>279-284) Donyai (2008) (Conyai et al. 230-237) Barber (2007) (Barber, Cornford, and Klecun 271-278) Franklin (2008) (Franklin, Jakclin, and Barber 375-379) Franklin (2007) (Franklin et al. 133-139) Design: Before-after N = 4,803 medication orders Implementation: 06/2003 Study Start: 00/0000 Study End: 00/0000</p> | <p>Machine, e-Medication administration system (e-MAR, e-TAR) e-Rx Integrated Pharmacy</p> | <p>28 bed surgery ward of a teaching hospital Inpatient hospital based Academic</p> | <p>for drug administrations*, %administered <1hr (Franklin, Jacklin, and Barber 375-379), rate of pharmacist interventions (Donyai et al. 230-237), total pharmacy time taken on study ward</p> | <p>error rate for new orders dropped significantly after implementation of the system (3.8% vs. 2.0%, RRR 47%, p = 0.0004). Medication administration error rate also significantly declined (8.6% vs. 4.4%, RRR 49%, p = 0.0003). (Franklin, Jacklin, and Barber 375-379) Post-intervention medication timeliness was improved (% administered <1hr, 79% vs. 89%, p <0.001). (Donyai et al. 230-237) The rate of pharmacist interventions declined significantly after implementation (3.0% vs. 1.9%, AR 1.1 (95% CI 0.2,2.0). (Franklin et al. 133-139) Total pharmacy time taken on study ward increased after implementation (1h 8 min vs. 1h 38 min, p = 0.001). Pharmacists were required to endorse fewer orders (50% vs. 21%,</p> | |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|------------------------|--|---|----------|
| | | | | RRR 58%, p <0.0001) and endorsed fewer orders (55% vs. 30%, RRR 45%, p <0.0001). | |
| <p>Fretheim (2006) (Fretheim, Aaserud, and Oxman e216) Fretheim (2006) (Fretheim et al. e134) Design: RCT N = 139 practices and 501 physicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>thiazides prescription rates*, rates of cardiovascular risk assessment, proportion of patients achieving treatment goal at 3 months</p> | <p>Prescribing of thiazides increased in the reminders + group (11% vs. 15%, RRR 54%, p <0.001, RR 1.94 95% CI 1.49 to 2.49). The groups did not differ for cardiovascular risk assessment (RR 1.04, CI 0.60 to 1.71) or proportion that achieved treatment goal at 3 months (RR 0.98, CI 0.93 to 1.02).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--------------------------------------|---|--|----------|
| <p>Galanter (2004) (Galanter, Polikaitis, and Didomenico 270-277) Design: Before-after N = 620 patients Implementation: 00/0000 Study Start: 02/2001 Study End: 03/2002</p> | <p>CDSS/CDS/CCDS/reminders Integrated CDSS/CDS/CCDS/reminders CPOE/POE system Laboratory system</p> | <p>Acute care/tertiary, Academic</p> | <p>compliance with digoxin monitoring guidelines - synchronous alerts*, compliance with hypokalemia and hypomagnesemia treatment guidelines - synchronous alerts*, compliance with hypokalemia and hypomagnesemia treatment guidelines - asynchronous alerts*</p> | <p>Post implementation, synchronous alerts significantly increased test ordering for digoxin levels, K levels and Mg levels at 1 hr and 24 hrs (p <0.01 for all). Supplementatio n of Mg at 1 hour was significantly improved, but not at 24 hrs. Supplementatio n of K was not improved at 1 or 24 hrs. Synchronous alerts resulted in improved compliance at 1 hr and 24 hrs for both K and Mg supplementatio n (p <0.01).</p> | <p>+</p> |
| <p>Galanter (2005) (Galanter, Didomenico, and Polikaitis 269-274) Design: Before-after N = 410 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system</p> | <p>Acute care/tertiary, Academic</p> | <p>likelihood of a patient receiving contraindicated medication, compliance rates- housestaff compared to other clinicians</p> | <p>The likelihood of a patient receiving at least one dose of the contraindicated medication decreased from 89% to 47% after alert implementation (p <0.0001), RRR 47%. For the 226 alerts received by housestaff, the alert compliance rate was 42%; for the remaining clinicians the compliance rate was 38% (p = 0.54).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--------------------------------------|---|---|----------|
| <p>Gerard (2008) (Gerard et al. 776-779) Design: Time series N = 907 orders for flu vaccination Implementation: 00/2001 Study Start: 00/2003 Study End: 00/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system</p> | <p>General Hospital 464 Beds</p> | <p>acceptance rate of pre-selected orders, year 1 vs. year 2, acceptance rate of pre-selected orders, year 2 vs. year 3, vaccination rate, year 1 vs. year 2, vaccination rate, year 2 vs. year 3</p> | <p>During the intervention, physicians were significantly more likely to accept pre-selected vaccination orders, Year 1 (47%), Year 2 (77%), Year 3 (83%); however vaccine administration by nurses was suboptimal. EMR functionality improved, patient receipt of vaccine increased significantly, Year 1 [0/36; 0%], Year 2 [8/66; 12%], Year 3 [286/805; 36%].</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|------------------------|---|--|----------|
| <p>Gill (2009) (Gill et al. 221-226) Design: RCT N = 64,150 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 10/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Up-to-date lipid test*, Lipid medication if not at goal (high risk patients only)*</p> | <p>Outcomes improved for most measures from before to 1 year after the intervention (univariate analysis). However, after controlling for confounding variables and for clustering in multilevel modeling, only up-to-date lipid testing for high-risk patients was statistically better in the intervention group as compared to the control group (adjusted OR 15.0, p <0.05). Intervention status was NS</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|-------------------------------------|---|--|----------|
| <p>Gilutz (2009) (Gilutz et al. 23-29) Design: RCT N = 7,448 patients from 56 control and 56 intervention clinics Implementation: 00/0000 Study Start: 01/2000 Study End: 12/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system, Laboratory system, Pharmacy</p> | <p>Ambulatory care Academic</p> | <p>rate of adequate monitoring Positive treatment trend, overall up-titration rate in patients with LDL = 110 mg/dl</p> | <p>higher rate of adequate monitoring documented in intervention arm (54.8% vs. 48.7%, p <0.001). Medication initiation or up-titration recommended for patients with LDL levels above 110 mg/dl results showed overall positive trends were minimally more prominent in the intervention arm (59.1% vs. 53.7%, p <0.003). This difference constitutes a higher rate of drug initiation (2.5%), up-titration (1.8%) and avoiding drug cessation (1.1%). However, overall up-titration in patients with LDL = 110 mg/dl was poor, both in the intervention arm and in the control arm (8.6% vs. 7.4%,NS).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|--|----------|
| <p>Ginzburg (2009) (Ginzburg et al. 2037-2041) Design: Before-after N = 540 Patients Implementation: 00/0000 Study Start: 01/2005 Study End: 12/2005</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Medication error*</p> | <p>Significantly more medication errors were found in the preintervention group than in the postintervention group [(32.6% (n = 103) vs. 20.5% (n = 46), p = 0.002]. Significantly fewer strength overdosing errors occurred in the postintervention group (8.9% vs. 4.0%, p = 0.028).</p> | <p>+</p> |
| <p>Goethe (1997) (Goethe, Schwartz, and Szarek 553-558) Design: Time series N = 1,604 alerts Implementation: 00/0000 Study Start: 01/1994 Study End: 12/1996</p> | <p>CDSS/CDS/CCDS/reminders</p> | <p>Other specialty hospital (rehab, oncology) 130 Beds</p> | <p>alert rate, physician response rate to alerts*, compliance with alerts*</p> | <p>The rate of alerts went down in the second year (29% vs. 15%, RRR 48%, p <0.001), as did the rate of physician responses to the alerts (67% vs. 55%, RRR 18%, p <0.001) change in practice to comply with guidelines occurred 28% (year 1) compared to 21% (year 2)</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|-----------------------------------|--|--|----------------|
| Griffey (2009) (Griffey 265) Design: Time series N = 2,419 orders Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders Stand-Alone, CPOE/POE system | Emergency department, Academic | During use of the CDSS system, agreement with recommended doses was increased. | During use of the CDSS system, agreement with recommended doses was increased (23.0% for off and 31.4% for on, RRR 37%, p = 0.03) reduction similar for benzodiazepines (p = 0.03), opiates (p = 0.04), and NSAIDs (p = 0.0009). | + |
| Halkin (2001) (Halkin et al. 260-265) Design: Time series N = 775,186 prescriptions Implementation: 11/1997 to 00/1998 Study Start: 01/1998 Study End: 06/1999 | CDSS/CDS/CCDS/ reminders Integrated Pharmacy | Pharmacy, HMO pharmacy | rate of drug interaction prescriptions 90% pharmacies and 50% physicians compared with baseline, rate of drug interaction prescriptions 95% pharmacies and 90% physicians compared with baseline | Dispensing of drug interaction prescriptions was reduced by 21.1% and by 67.5% in periods II and III compared with period I (OR, 0.79; 95% CI, 0.75 to 0.83 and OR, 0.28; 95% CI, 0.26 to 0.30, respectively). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|------------------------|---|--|----------|
| <p>Hicks (2007) (Hicks et al. 429-441) Design: RCT N = 1,422 patients Implementation: 00/0000 Study Start: 07/2003 Study End: 02/2005</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Other, Academic</p> | <p>blood pressure controlled, receiving a recommended drug class medication within 1 week of the clinic visit adjusted</p> | <p>This study had 4 groups: usual care, CDS, NPs, and NPs+CDS. No difference was seen across all 4 groups for blood pressure readings: Usual care vs. CDS: 45% vs. 48% controlled, OR 0.96 (CI 0.78 to 1.19). Patients in the CDS group were more likely to have received a recommended drug class medication within 1 week of the clinic visit than patients in the usual care group: adjusted OR 1.32 (CI 1.09 to 1.61).</p> | <p>+</p> |
| <p>Hollingworth (2007) (Hollingworth et al. 722-730) Devine (2010) (Devine et al. 152-171) Design: Cross-sectional N = 146 health care providers (69 in phase 1 and 77 in phase 2) Implementation: 00/2003 Study Start: 02/2005 Study End: 01/2006</p> | <p>e-Rx Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>time spent on writing tasks (min/hr), paper vs. desktop vs. laptop, time spent on computer tasks (min/hr) paper vs. desktop vs. laptop, time spent on computer and writing tasks (min/hr), paper vs. desktop vs. laptop, More time in phase 2 compared with handwritten prescriptions for all prescriptions and new prescriptions but not for renewed prescriptions. (Devine et al. 152-171)</p> | <p>Prescribers at e-RX sites, both desktop and wireless laptops, spent significantly less time (minutes/hour) on writing tasks than their paper-based colleagues (8.7 paper vs. 5.5 desktop vs. 5.9 laptops, p <0.05), but more time on computer based tasks (3.8 vs. 7.4. vs. 8.1, p <0.05). Overall time on writing tasks and computer tasks together were not different among</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--------------------------------------|--|--|----------|
| | | | | the three formats (12.4 vs. 12.9. vs. 14.0, NS) and should not greatly disrupt workflow.[In the second phases (Deveine et al. 152-171) (point of care prescribing) the clinicians spend more time than handwritten prescribing for all prescriptions: 25 seconds more (99.5% CI 12 to 38), and more time for new prescriptions: 29 seconds more (CI 14 to 44) but not more time for renewed prescriptions: 13 seconds more (CI -13 to 39). | |
| <p>Hulgan (2004) (Hulgan et al. 349-357) Design: Time series N = 15,194 quinolone orders Implementation: 02/2002 Study Start: 02/2001 Study End: 01/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary, Academic</p> | <p>change in weekly proportion of oral quinolone orders*</p> | <p>55.5% orders were for oral quinolones before the intervention orders compared with 62.4% after (RRR -12%, p = NR). In the time-series analysis, the intervention increased the proportion of oral quinolone orders per week by 5.6% (95% CI 2.8 to 8.4%; p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|---|----------|
| <p>Hwang (2002) (Hwang, Park, and Bakken 213-223) Design: Time series N = 171 patients Implementation: 10/1999 Study Start: 06/1999 Study End: 05/2000</p> | <p>CPOE/POE system Integrated Hospital information system Imaging systems</p> | <p>Acute care/tertiary, 1,000 plus Beds Academic</p> | <p>number of daily orders per patient, number of daily medication orders, number of changed orders, number of cancelled orders, number of daily PRN orders</p> | <p>daily orders per patient significantly increased following POE system introduction compared to both 3- and 6-months post (10.9 vs. 17.4 vs. 19.9, p </0.0001) similar pattern observed for number of daily medication orders (4.2 vs. 6.6 vs. 6.1, p <0.0001) and PRN orders (2.9 vs. 7.9 vs. 8.3, p <0.0001) difference between 3 and 6 months after POE was NS for either measure. The number of changed orders (2.2 vs. 0.2 vs. 0.03, NS) and cancelled orders (3.3 vs. 2.3 vs. 2.2, NS)</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|---|----------|
| <p>Igboechi (2003) (Igboechi et al. 227-231) Design: Before-after N = 10,134 medication errors Implementation: 06/2001 Study Start: 06/1999 Study End: 05/2002</p> | <p>CPOE/POE system Integrated CDSS/CDS/CCDS/reminders EHR/EMR system, Pharmacy</p> | <p>Acute care/tertiary, 350 Beds Inpatient hospital based</p> | <p>total potential errors*, Illegible orders, incomplete orders, incorrect orders, drug therapy problems</p> | <p>The number of documented medication errors decreased postimplementation for total potential errors (p <0.001), illegible orders (p <0.001), incomplete orders, (p <0.001) and incorrect orders (p <0.001) but not for drug therapy problems (p = 0.289). Annual numbers were compared for each of the 2 years before implementation of CPOE and the year after CPOE.</p> | <p>+</p> |
| <p>Jacques (2005) (St Jacques et al. 215-221) Design: Before-after N = 287 procedures Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system</p> | <p>Acute care/tertiary, Academic</p> | <p>antibiotic redosing rate*</p> | <p>On-time antibiotic redosing increased significantly after the implementation of the computer reminder system (20% vs. 57%, RRR - 185%, p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|---|---|----------|
| <p>Jani (2008) (Jani et al. 214-218) Design: Before-after N = 2,222 prescribed drugs Implementation: 03/2006 Study Start: 07/2005 Study End: 07/2006</p> | <p>e-Rx Integrated EHR/EMR system, e-MAR, Pharmacy</p> | <p>Pediatric stand alone hospital, Ambulatory care</p> | <p>error rate*, error free visit rate</p> | <p>The overall prescribing error rate was 77.4% (95% CI = 75.3% to 79.4%) for handwritten items and 4.8% (95% CI = 3.4% to 6.7%) with e-Rx (RRR, 94%, p <0.001). Pre-e-Rx, 1153 items (73.3%; 95% CI = 71.1% to 75.4%) were missing essential information, and 194 items (12.3%; 95% CI = 10.8% to 14%) were judged to be illegible. Post-EP, only 9 items (1.4%; 95% CI = 0.7% to 2.6%) were missing essential information, and illegibility errors were eliminated. The number of patient visits that were error-free increased from 21% to 90% (69% difference; 95% CI = 64% to 73.4%; RRR - 324%, p <0.001) after the implementation of e-Rx.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|---|---------|
| Javitt (2008) (Javitt, Rebitzer, and Reisman 585-602) Design: RCT N = 39,508 patients Implementation: 01/2001 Study Start: 01/2001 Study End: 12/2001 | CDSS/CDS/CCDS/ reminders Integrated Billing/administration system Laboratory system, Pharmacy | Ambulatory care | resolution rate-add a drug alert*, resolution rate-stop a drug*, resolution rate -do a test* | Resolution rate for "add a drug" CCs was 8.6 % higher in the study group than the control group (p <0.05). There was, however, no significant difference in the resolution rates for "stop a drug" CCs (change -6%, NS). Resolution rates for "do a test" CCs were 5.8% higher in the study group, p <0.05. | + |
| Johnson (2010) (Johnson et al. 321-325) Design: RCT N = 3,285 patients Implementation: 00/0000 Study Start: 04/2007 Study End: 08/2007 | CDSS/CDS/CCDS/ reminders e-Rx Integrated EHR/EMR system | Ambulatory care, Pharmacy, Not specified, Academic | rate of callbacks generated* | There was no significant difference in the callback rates between the "SYW off" and the "SYW on" periods (0.4% vs. 0.45%; p = 0.47). | - |
| Kadmon (2009) (Kadmon, et al. 935-940) Design: Time series N = 5,000 Medication orders Implementation: 11/2004 Study Start: 09/2004 Study End: 09/2007 | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, Hospital information system | Acute care/tertiary, Critical care units (CCU, ICU, NICU) 12 bed PICU unit | total prescription error rate(combination of the 3 error types)*, pADE*, rule violations*, medication prescription errors* | Among the 5,000 prescriptions reviewed, 273 (5.5%) contained prescription errors. Implementation of CPOE associated with a slight, nonsignificant decrease in prescription error rate (between periods 1 and 2; 8.2% vs. 7.8%, p = 0.66). Decreases in rate of | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---------------------|----------------------------------|----------|-------------------|--|---------|
| | | | | <p>prescription errors after CDSS implementation were statistically significant between periods 2 and 3 (7.8% vs. 4.4%, $p = 0.0004$) and after prescription authorization between period 3 and 4 (4.4% vs. 1.4%, $p < 0.0001$). The rate of potential ADEs decreased slightly between periods 1 and 2 (from 2.5% to 2.4%, $p = 0.9$) and significantly in periods 3 and 4 (to 0.8% and 0.7%, respectively; $p < 0.005$). Rate of MPEs decreased slightly between periods 1 and 2 (from 5.5% to 5.3%, $p = 0.79$), but new types of MPEs appeared. A significant decrease in period 3 (to 3.8%; $p < 0.05$) and a dramatically significant decrease in period 4 (to 0.7%; $p < 0.0005$) was noted. 3 RVs were found</p> | |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|---|---------|
| | | | | between period 1 and 2 (0.002% vs. 0.001%, p = 0.3). No RVs were found in period 3 and 4. | |
| Kaplan (2006) (Kaplan et al. 461-467) Design: Time series N = n/a orders Implementation: 04/2002 Study Start: 12/2001 Study End: 01/2004 | CPOE/POE system Integrated Formulary, Hospital information system, Imaging systems, Pharmacy | Pediatric stand alone hospital 423 Beds | rate of verbal orders*, rate of unsigned verbal orders* | Overall, there was a significant decrease in the rates of verbal orders (from 22% to 10%) and unsigned verbal orders (from 43% to 9%) between the period before CPOE implementation and 21 months after CPOE implementation (p = 0.0001 for both). | + |
| Karson (2007) (Karson et al. 1004) Design: Time series N = 74,494 verbal orders Implementation: 0/0000 Study Start: 01/2005 Study End: 00/0000 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system | Acute care/tertiary, 900 Beds Academic | compliance with co-signing within 24 hours, compliance with co-signing by month end | At baseline, 49% of verbal orders were co-signed within 24 hours. This increased to 63% after the first intervention and 93% after the second intervention (p <0.001). At month end, the compliance rate was 61% at baseline, 94% after the first intervention and 98% after the second (p <0.001). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|---|----------|
| <p>Kazemi (2010) (Kazemi et al. e5) Design: Observational study N = 158 patients (neonates) Implementation: 00/2007 Study Start: 12/2007 Study End: 09/2008</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Hospital information system</p> | <p>Acute care/tertiary, 400 Beds Academic</p> | <p>rate of non-intercepted errors for orders (POE Errors vs. NOE Errors)*, rate of nonintercepted errors for ordered medication*, rate of nonintercepted errors for patient days*, rate of nonintercepted errors medication-days *</p> | <p>The rate of nonintercepted errors for orders decreased from 22.7% to 14.5% (RR 0.64; 95% CI 0.53 to 0.77, p <0.001). For ordered medication it dropped from 12.8% to 7.6% respectively (RR 0.60; 95% CI 0.50 to 0.71, p <0.001). However, the highest rate difference (9.5%) was seen when calculated according to patient days (24.5% vs. 15%, RR 0.61; 95% CI 0.49 to 0.77; p <0.001). The rate difference for medication-days were 5.8% (14.4% vs. 8.6%, RR 0.60; 95% CI 0.49 to 0.74; p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|--|----------|
| <p>Kim (2006) (Kim et al. 495-498) Design: Before-after N = 2,375 chemo orders Implementation: 02/2003 Study Start: 07/2001 Study End: 02/2004</p> | <p>CPOE/POE system Integrated EHR/EMR system</p> | <p>Other specialty hospital (rehab, oncology) Academic</p> | <p>rate of improper dosing on treatment plans, rate of improper dosing on orders, rate of treatment plans and orders not matching, rate of missing cumulative dose calculations, rate of incorrect dosing calculations</p> | <p>After CPOE deployment, daily chemotherapy orders were less likely to have improper dosing on orders (2.3% vs. 0.1%, RRR 97%, p <0.05), incorrect dosing calculations (5.8% vs. 0.5%, RRR 91%, p <0.05), missing cumulative dose calculations (18% vs. 5.7%, RRR 68%, p <0.05), and incomplete nursing checklists (4.8% vs. 2.5%, RRR 48%, p <0.05). There was no difference in the likelihood of improper dosing on treatment plans (4.0% vs. 2.6%, RRR 35%, NS) and a higher likelihood of not matching medication orders to treatment plans (1.1% vs. 6%, RRR -445%, p <0.05).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|---|---|---------|
| Kim (2008) (Kim et al. 416-421) Design: Time series N = not given not given Implementation: 00/2002 Study Start: 02/2004 Study End: 04/2006 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system | Acute care/tertiary, 750 Beds Academic | 3rd generation cephalosporin use (daily doses/1,000 patient days) | The use of third generation antibiotic cephalosporin use decreased significantly from 103.2 doses/1,000 patient days to 84.9 postimplementation. It increased once the feedback element was stopped (84.9 vs. 115.1, p <0.05). | + |
| Kirk (2005) (Kirk et al. 817-824) Design: Observational study N = 4,274 prescriptions Implementation: 00/2000 Study Start: 03/2003 Study End: 08/2003 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system | Acute care/tertiary, Ambulatory care Academic | error rate | The computer calculated dose error rate was 12.6% compared with the traditional prescription error rate of 28.2% (RRR 55%, p <0.001). | + |
| Kitahata (2003) (Kitahata et al. 803-811) Design: Time series N = 1,204 patients with HIV Implementation: 04/1998 Study Start: 03/1996 Study End: 09/1999 | CDSS/CDS/CCDS/reminders Integrated Hospital information system | Ambulatory care, Academic | rate of prophylaxis for mycobacterium avium complex infection, rate of prophylaxis for pneumocystis cairnii pneumonia | After implementation of the CDSS patients were more likely to be given prophylaxis for mycobacterium avium complex infection (Hazard Ratio 3.84, CI 1.58 to 9.32, = 0.003) but not for pneumocystis cairnii pneumonia (Hazard Ratio 1.14, CI 0.84 to 1.59, NS). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|---|---|----------|
| <p>Koide (1999) (Koide et al. 11-19) Design: Before-after N = 1,024 prescriptions for 111 patients and 68 physicians Implementation: 09/1994 Study Start: 09/1994 Study End: 09/1996</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system, Hospital information system, Laboratory system</p> | <p>Acute care/tertiary, 1,040 Beds Academic</p> | <p>rate of 'appropriate' prescribing (normal value of ALT or AST within 3 mon)ths</p> | <p>127/491 (25.9%) preintervention prescriptions were classified as 'appropriate'. 353/533 (66.2%) postintervention prescriptions were classified as 'appropriate'. This sudden increase in level of 40.3% occurring immediately after the start of the intervention was highly significant (p <0.0001).</p> | <p>+</p> |
| <p>Kooij (2008) (Kooij et al. 893-898) Design: Time series N = 1,565 patients Implementation: 00/0000 Study Start: 11/2005 Study End: 06/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>General Hospital, Academic</p> | <p>rate of prophylaxis, control vs. CDSS, rate of prophylaxis, CDSS vs. stopping CDSS</p> | <p>Patients who needed PONV prophylaxis were more likely to be prescribed medication if clinicians were provided with electronic DS (73%) than before DS (38%) or after the electronic DS was stopped (37%), p <0.001.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--------------------------------------|--|--|----------|
| <p>Kooij (2009) (Kooij et al. 187-191) Design: Before-after N = 5,652 patients Implementation: 00/0000 Study Start: 11/2005 Study End: 06/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Acute care/tertiary, Academic</p> | <p>percentage of patients who received dexamethasone*, percentage of patients who received granisetron*, percentage of patients who received both dexamethasone and granisetron*</p> | <p>Dexamethasone was given to 46% of the control period. In the decision support period, rate increased significantly to 95% and after deactivating the automated reminders, it decreased to 47% in the post decision support period (p <0.001) For granisetron, these percentages were 53%, 81%, and 51%, respectively (p <0.001). Percentage of patients receiving both medications was 39% in the control period, increased to 79% in the decision support group and decreased to 41% in the post decision support group (p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---------------------------------|--|--|---------|
| Kralj (2003) (Kralj et al. 197-203) Design: Case control N = 11,644 patient-physician encounter Implementation: April 2000 Study Start: 12/1999 Study End: 11/2001 | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Ambulatory care | changes in prescribing rates of erythropoietin between clinics at baseline compared with during the intervention group | The mean difference in prescribing rates between experimental and control clinic at baseline was 0.36 (p = 0.044). Whereas in the intervention period the difference in the rates between them almost tripled to .093 (p = 0.000). The rate of erythropoietin prescribing increased by 14.2% (p = 0.05) at the experimental clinic. It declined by 15.9% (p = 0.12, NS) in the control clinic. | + |
| Krall (2004) (Krall, Traunweiser, and Towery 1-9) Design: RCT N = 1,076 patients Implementation: 00/1994 Study Start: 01/2000 Study End: 02/2000 | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Acute care/tertiary | proportion of patients no longer eligible for alerts at the end of the month* | Following implementation of the alert, more patients were 'no longer eligible for alerts at the end of the month' (25.8% pre vs. 54.3% post, RRR - 103%, p <0.001). | + |
| Kucher (2005) (Kucher et al. 969-977) Design: RCT N = 2,506 patients Implementation: 00/0000 Study Start: 09/2000 Study End: 01/2004 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system, Hospital information system | Acute care/tertiary Academic | received pharmacological interventions | More patients in the CDSS group received pharmacological interventions. (13% vs. 24%, RRR 69%, p <0.001). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|----------------------------------|------------------------|--|--|----------|
| <p>Lapane (2008) (Lapane et al. 442-446) Design: Mixed methods N = 276 primary care prescribers and their staff Implementation: 00/2003 Study Start: 04/2006 Study End: 08/2006</p> | <p>e-Rx</p> | <p>Ambulatory care</p> | <p>self reported drug alert overrides*</p> | <p>22/145 prescribers (15%) reported overriding drug-allergy alerts most of the time or 'always' with variation in frequency of overriding drug alerts by e-Rx software system ranging from 9% to 50% (p = 0.656 for overall comparison by e-Rx software system). Nearly 1 in 4 respondents reported overriding drug-dose alerts 'most of the time' or 'always' (range 13% to 33%; p = 0.006). More than 40% indicated they override drug-drug interactions 'most of the time' or 'always' (range, 25% to 50%; p = 0.374).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|---|----------|
| <p>Lecumberri (2008) (Lecumberri et al. 699-704) Design: Time series N = 19,338 patients Implementation: 09/2005 Study Start: 01/2005 Study End: 06/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated hospital guidelines</p> | <p>Unspecified Hospital Academic</p> | <p>number of alerts, proportion of alerted patients receiving thromboprophylaxis</p> | <p>an electronic alert was sent to 32.8% and 32.2% of all hospitalized patients, respectively. Appropriate prophylaxis among alerted patients was ordered in 89.7% (2006) and 88.5% (in 2007) of surgical patients, and in 49.2% (in 2006) and 64.4% (in 2007) of medical patients.</p> | <p>-</p> |
| <p>Ledwich (2009) (Ledwich et al. 1505-1510) Design: Before-after N = 2,477 vaccine possibilities (patients) Implementation: 00/0000 Study Start: 10/2006 Study End: 12/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care, Academic</p> | <p>influenza vaccination rates, pneumococcal vaccination rates*</p> | <p>PostBPA influenza vaccination rates significantly increased (47% to 65%; p <0.001), at both sites. PostBPA pneumococcal vaccination rates likewise significantly increased (19% to 41%; p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|---|---|----------|
| <p>Lesprit (2009) (Lesprit et al. 1058-1063) Design: Observational study N = 932 prescriptions Implementation: 11/2006 Study Start: 11/2006 Study End: 10/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Laboratory system</p> | <p>Acute care/tertiary, 960 Beds Academic</p> | <p>actual duration of treatment in days compared to prescribed*</p> | <p>Of the 482 prescriptions requiring intervention, the physicians complied with 80.3% of the recommendations. There was a significant reduction in the actual duration of antibiotic treatment compared to the originally prescribed duration (8 to 7 days (p <0.0001).</p> | <p>+</p> |
| <p>Lester (2005) (Lester et al. 22-29) Design: RCT N = 235 patients and 14 clinicians Implementation: 07/2003 Study Start: 07/2003 Study End: 07/2004</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care Academic</p> | <p>proportion of patients receiving statins*, proportion of patients receiving statins at 1 yr*</p> | <p>At 1 month, more patients in the email group had received statins than control patients (3% vs. 15%, RRR 400, p <0.001). At 1 year the difference in receipt of statins had disappeared (17% vs. 25%, NS).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|--|--|----------|
| <p>Lin (2008) (Lin et al. 620-626) Design: Time series N = 1,123 high severity order checks Implementation: 00/1997 Study Start: 01/2001 Study End: 01/2006</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary, General Hospital, 444 Beds Ambulatory care, Long term care (nursing homes)</p> | <p>override rates-severe drug-drug alerts*, override rates-severe drug-allergy alerts*</p> | <p>There were 215 high severity order checks in 2001 (0.5% of orders) and 908 in 2006 (2.5% of orders). Rate of overrides for drug-drug checks remained the same between 2001 and 2006 (88% vs. 87%, NS). Rate of overrides for drug-allergy order checks increased significantly from 2001 to 2006 (69% vs. 81%, RRR - 17%, p <0.005).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|------------------------|--|---|----------|
| <p>Linder (2009) (Linder et al. 231-240) Design: RCT N = 111,820 patients Implementation: 00/0000 Study Start: 11/2005 Study End: 05/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Imaging systems, Laboratory system</p> | <p>Ambulatory care</p> | <p>Rate of antibiotic prescribing to patients with ARI *</p> | <p>In the intent-to-intervene analysis, clinicians prescribed antibiotics to 43% of patients with ARI diagnoses in control clinic compared to 39% in the intervention clinic (OR. 0.8; 95% CI 0.6 to 1.2; p = 0.30). The ARI Smart Form did not significantly reduce overall antibiotic prescribing, was used by 33% of intervention clinicians (86/262) at least once. Appropriate antibiotic prescribing rate was 88% (n = 990 visits) in the as-used analysis.</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|-------------------------------------|--|--|----------|
| <p>Liu (2008) (Liu et al. 1109-1112) Design: Time series N = 858 patients Implementation: 00/1989 Study Start: 01/2005 Study End: 12/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary</p> | <p>percentage of no prophylactic antibiotic after clean surgery, mean number of days of antibiotic treatment</p> | <p>In clean procedures, the percentage of no prophylactic antibiotic after surgery increased in the long run (overall 76% vs. 84% vs. 93%, no analysis); the increase was significant for 2 of the 4 surgery types (p <0.005). In clean-contaminated procedures, the duration of prophylactic antibiotic after surgery (mean number of days) was significantly reduced in 2 of the 3 surgery types (p <0.001).</p> | <p>+</p> |
| <p>Madaras-Kelly (2006) (Madaras-Kelly et al. 155-169) Design: Time series N = not reported Implementation: 00/0000 Study Start: 07/2001 Study End: 06/2004</p> | <p>CDSS/CDS/CCDS/CPOE/POE system Integrated Hospital information system</p> | <p>Acute care/tertiary, 87 Beds</p> | <p>use of antibiotics*</p> | <p>Use of aminopenicillin beta-lactam inhibitors, all fluoroquinolones and levofloxacin decreased while use of first-generation cephalosporins and trimethoprim sulfamethoxazole increased (p <0.05 for each).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|---|----------|
| <p>Mahoney (2007) (Mahoney et al. 1969-1977) Design: Before-after N = 2,843,135 inpatient medication orders Implementation: 02/2002 Study Start: 02/2002 Study End: 06/2006</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Pharmacy information system Integrated EHR/EMR system, Hospital information system</p> | <p>General Hospital, Pediatric stand alone hospital, 966 in 2 hospitals Beds Pharmacy, Inpatient hospital based, Academic</p> | <p>rate of: -drug allergy violations*, -excessive doses*, -incomplete or unclear orders*, -therapeutic duplication*</p> | <p>Medication errors decreased after implementation of the CIT with respect to drug allergy violations (OR 0.14, 95% CI 0.11 to 0.17, p <0.001), excessive doses (OR 0.68, 95% CI 0.62 to 0.74, p <0.001) and incomplete or unclear orders (0.35, 95% CI 0.32 to 0.38, p <0.001), but no decrease in therapeutic duplications. Turnaround time between drug ordering and administration decreased from 90 minutes to 11 minutes. The override rate also decreased (7.1 to 2.9%, RRR 59%, p = 0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|------------------------|--|--|----------|
| <p>Martens (2007) (Martens et al. S403-S416) Design: RCT N = 77 physicians (GPs) Implementation: 04/2004 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated, EHR/EMR system</p> | <p>Ambulatory care</p> | <p>quinolone prescriptions, inhaled corticosteroids for newly diagnosed COPD in patients >40 years, first choice drugs for sore throats GPs got either reminders on antibiotics, asthma, and COPD or cholesterol. Reminders were either to stop prescribing drugs or to prescribe a specific first-line drug.</p> | <p>No differences were seen for either group to prescribe a drug or in the cholesterol reminder group. GPs in the antibiotics, asthma and COPD group showed changes in 3 of 8 drug categories. Outcome measures were sum scores for drug volume: lower scores were improvements in prescribing. Reminder physicians prescribed fewer quinolones (4.6 (95% CI 2.8 to 8.1) vs. 1.5 (95% CI 0.8 to 2.2); fewer inhaled corticosteroids for newly diagnosed COPD in patients >40 yr (0.5 (95% CI 0.3 to 0.9) vs. 0.0 (95% CI 0 to 0.1), p = 0.00); and better first choice drugs for sore throats (0.8 (95% CI 0.3 to 2.4) vs. 0.2 (95% CI 0.0 to 9.4), p = 0.03.</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|--|----------|
| <p>Mattison (2010) (Mattison et al. 1331-1336) Design: Before-after N = not reported medication orders Implementation: 10/2004 Study Start: 06/2004 Study End: 08/2008</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system</p> | <p>Acute care/tertiary, 621 Beds Academic</p> | <p>rate of prescribing not-recommended medications*, rate of prescribing medications with recommended dosage reductions*</p> | <p>In before-and-after comparisons, the mean (SE) rate of prescribing not-recommended medications dropped from 11.56 (0.36) to 9.94 (0.12) orders per day (difference, 1.62 [0.33]; p <0.001). There were no appreciable changes in the rate of ordering medications for which only dose reduction was recommended or that were not targeted after CPOE implementation.</p> | <p>+</p> |
| <p>Maynard (2010) (Maynard et al. 10-18) Design: Time series N = 3,285 patients Implementation: 04/2006 Study Start: 00/2005 Study End: 00/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system</p> | <p>Acute care/tertiary, 350 Beds Academic</p> | <p>percent of patients on adequate prophylaxis*</p> | <p>The percent of patients on adequate prophylaxis improved in each of the 3 years from a baseline of 58% in 2005 to 78% in 2006 (unadjusted relative benefit = 1.35; 95% CI 1.28 to 1.43), and 93% in 2007 (unadjusted relative benefit =1.61; 95% CI 1.52 to 1.69).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|---|----------------|
| McCluggage (2010) (McCluggage et al. 70-75) Design: Before-after N = 522 patients Implementation: 02/2007 Study Start: 08/2006 Study End: 04/2007 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system | Acute care/tertiary, Academic | optimal regimen prescribed* | The percentage of patients whose initial vancomycin regimen matched the nomogram recommendation was higher in the postimplementation group compared with the preimplementation group (35.8% vs. 23.7%, p = 0.0028). | + |
| McGregor (2006) (McGregor et al. 378-384) Design: RCT N = 4,507 patients Implementation: 00/000 Study Start: 05/2004 Study End: 08/2004 | CDSS/CDS/CCDS/reminders Integrated Laboratory system, Pharmacy | Acute care/tertiary 648 Beds Inpatient hospital based, Academic | mean time spent on antimicrobial management | Team members spent 3.2 hours per day on management of antimicrobials with the decision support system compared with 4 hours per day without. No statistical testing was done. | + |
| McMullin (1999) (McMullin et al. 2077-2082) Design: Before-after N = 265 patients Implementation: 01/1996 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Laboratory system, Pharmacy | Acute care/tertiary, Pharmacy, Inpatient hospital based | rate of concomitant orders for contraindicated medications with cisapride* | The rate of ordering contraindicated drugs with cisapride was reduced with COPE (9% vs. 3.1%, RRR 65%, p <0.001). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|------------------------|---|--|----------|
| <p>Miskulin (2009) (Miskulin et al. 1081-1088) Design: Cohort study N = 8,941 patients Implementation: 00/2005 Study Start: 11/2005 Study End: 04/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>EPO use*, time spent on anemia management (hours per month)</p> | <p>After adjustment for center and baseline differences, the log weekly EPO dose in patients treated using CDS was 4% less than those dosed manually (RR 0.96; 95% CI, 0.77 to 1.18, NS). CDS was associated with a nearly 50% decrease (p <0.001) in the time spent by dialysis unit staff on anemia management. Units using the computerized protocol spent a median of 3 hours per month on anemia management units using manual dosing spent a median of 6.5 hours per month.</p> | <p>+</p> |
| <p>Montgomery (2000) (Montgomery et al. 686-690) Design: RCT N = 552 patients Implementation: 00/0000 Study Start: 09/1996 Study End: 09/1998</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>probability of patients taking 2 drugs, probability of patients taking 3 drugs</p> | <p>risk chart group alone compared to computer support group had a lower probability of patients taking 2 drugs (OR 0.5, 95% CI 0.2 to 0.9) p <0.05) or 3 drugs (OR 0.3, 95% CI 0.1 to 0.6, p <0.05).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|---|---|----------|
| <p>Morrison (2006) (Morrison et al. 1033-1039) Design: Time series N = 3,864 patients Implementation: 00/0000 Study Start: 02/2001 Study End: 02/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary, 1171 Beds Academic</p> | <p>meperidine prescription rate, rate of patients receiving a concomitant laxative with an opioid</p> | <p>rate of patients receiving a concomitant laxative with an opioid did not change with the introduction of the CDSS system (data for 5 groups, 24.7% of patients who needed a laxative, 27.8%, 32.1%, 26.8%, and 34.0%, all comparisons NS). Fewer patients received meperidine with the introduction of the CDSS system. For group 4 (CDSS and enhanced assessment compared with Group 1 control 44.2% vs. 25.4%, RRR 20%, p <0.05). For Group 5 vs. Group 1 the rate of meperidine use was even lower (44.4% vs. 11.9%, RRR 73%, p = 0.01).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|---|----------|
| <p>Mullett (2001) (Mullett et al. e75) Design: Before-after N = 1,758 patients Implementation: 02/1999 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Hospital information system</p> | <p>Critical care units (CCU, ICU, NICU) Pediatric stand alone hospital, 232 Beds Inpatient hospital based, Academic</p> | <p>per patient anti-infective doses, per patient number of anti-infectives, per patient anti-infective orders per course, mean subtherapeutic anti-infective days/100 patient days, mean excessive dosage anti-infective days/100 patient days</p> | <p>The rate of per person use of anti-infective agents did not differ for PICU doses (12.8 vs. 13.4, NS), PICU number of doses (1.85 vs. 1.97, NS) but did differ for PICU anti-infective orders per patient-anti-infective course (1.56 vs. 1.38, p <0.01). The mean number of subtherapeutic risk days decreased (7.35 vs. 4.7, p <0.001) as did the mean excessive dosage risk days (8.45 vs. 6.1, p <0.001).</p> | <p>+</p> |
| <p>Nash (2005) (Nash et al. 64-69) Design: Time series N = 39,440 doses Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Medication safety reporting system Integrated Hospital information system Laboratory system</p> | <p>Acute care/tertiary, 1171 Beds Academic</p> | <p>reduction in excessive dosing for the nursing intervention, reduction in excessive dosing for the pharmacist intervention</p> | <p>There was a reduction in the rate of excessive dosing in the participating units compared to the control unit in the nurse intervention (23% for baseline for the control group with 17% for the nurse intervention and 17% for the pharmacist interventions (p <0.05).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--------------------------------|---|----------|
| <p>Newby (2003) (Newby, Fryer, and Henry 210- 213) Design: Cross- sectional N = 1,667 prescriptions Implementation: 00/0000 Study Start: 10/0000 Study End: 11/0000</p> | <p>e-Rx Integrated Pharmacy</p> | <p>Pharmacy, Stand alone (e.g. family run)</p> | <p>rate of repeat ordering</p> | <p>The rate of repeat ordering was higher for all antibiotics if the original was written using an e-Rx system (40% for paper vs. 69% for initial e-Rx, adjusted OR 3.82, 95% CI 2.55 to 5.72, p <0.05). This same significant affect was seen for all 4 study antibiotics. The rate of filling prescriptions NS as reported by patients (69% if the prescription was on paper vs. 61% by e-Rx, NS).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|----------------------------|---|--|----------|
| <p>Niemi (2009) (Niemi et al. 389-397) Design: Before-after N = 5,076 patients Implementation: 00/0000 Study Start: 10/2006 Study End: 03/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated Billing/administration system, Imaging systems, Laboratory system, Pharmacy</p> | <p>Acute care/tertiary</p> | <p>antibiotic administration within four hours*, pneumonia vaccination status documentation*, appropriate pneumonia antibiotic selection*, ACE or ARB initiation*, provision of discharge instructions to patients*</p> | <p>Compliance with the medication related indicators for pneumonia measures were NS; antibiotic administration within four hours (83% vs. 87%, RRR - 5%), pneumonia vaccination status documentation (82% vs. 92%, RRR-12%), appropriate pneumonia antibiotic selection (93% vs. 92%, RRR 1%). After implementation, heart failure medication related quality indicators measures were not significantly for ACE or ARB initiation (95% vs. 98%, RRR - 4%) but there was a significant increase in compliance with the provision of discharge instructions to patients (84% vs. 95%, RRR - 13%, p <0.01).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|------------------------------|--|--|----------|
| <p>Niiranen (2008) (Niiranen and Yli-Hietanen 4330-4332) Design: Time series N = 18,818 patient followups Implementation: 03/2005 Study Start: 04/2005 Study End: 12/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated Laboratory system</p> | <p>Ambulatory care, Home</p> | <p>proportion of patient followups assigned by nurses, year 1 to 2, proportion of patient followups assigned by nurses, year 2 to 3</p> | <p>In general, the share of patient followups assigned by nurses was similar in year 1 and 2 (56.7% vs. 55.1%, RRR 3%, NS), and increased significantly between year 2 and 3 (55.1% vs. 58.7%, RRR -7%, p <0.001).</p> | <p>+</p> |
| <p>Novis (2010) (Novis et al. 648-654) Design: Before-after N = 800 patients Implementation: 08/2007 Study Start: 03/2007 Study End: 03/2008</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Acute care/tertiary</p> | <p>percentage of patients receiving pharmacological prophylaxis, percentage of patients receiving sequential compression devices and pharmaprophylaxis</p> | <p>The proportion of patients receiving the recommended pharmacological prophylaxis preoperatively more than doubled (14% to 36%, p <0.001) Overall, the percentage of at-risk patients receiving the recommended combined DVT prophylaxis of SCD and pharmacological prophylaxis increased nearly seven-fold (5% to 32%, p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|---|---|----------|
| <p>Oliven (2005) (Oliven et al. 377-386) Design: Cross-sectional N = 1,350 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Drug order database EHR/EMR system, Hospital information system, Laboratory system</p> | <p>Acute care/tertiary, 88 Beds Academic</p> | <p>Type 1 PEs per 100 hospitalization days, Type 2 PEs per 100 hospitalization days</p> | <p>The incidence of Type 1 PEs was 5.21 and 1.36 per 100 hospitalization days in the HW dept and CDOE dept, respectively (p <0.0001). Type 2 PEs were more common, 7.20 and 3.02 per 100 hospitalization days in the HWdept and CDOEdept, respectively (p <0.0001), and about 75% of them were due to few drug laboratory interactions.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--------------------------------------|--|---|----------|
| <p>Overhage (1996) (Overhage, Tierney, and McDonald 1551-1556) Design: RCT N = 24 practice teams Implementation: 10/1991 Study Start: 10/1992 Study End: 03/1993</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system, Hospital information system, Laboratory system, Pharmacy</p> | <p>General Hospital Academic</p> | <p>rates of compliance with preventive care recommendations*</p> | <p>control teams complied with 24% of the reminders compared with 23% for intervention teams (p = 0.78) When preventive care measures were analyzed individually, 2 significant differences were seen in compliance (24-hour urine protein and angiotensin-converting enzyme [ACE] inhibitor) between control and intervention teams. Assumed to be due to chance with multiple testing and because they were in the opposite directions.</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|---------------------------------------|---|---|----------|
| <p>Overhage (1997) (Overhage et al. 364-375) Design: RCT N = 86 physicians on 6 services (services randomized) Implementation: 00/0000 Study Start: 10/1992 Study End: 04/1994</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system, Laboratory system</p> | <p>General Hospital, Academic</p> | <p>immediate compliance with corollary ordering*, 24 hour compliance*, hospital stay compliance*</p> | <p>Intervention physicians ordered the corollary orders required by the guidelines twice as often as control physicians did when measured by immediate compliance (46.3% vs. 21.9%, RRR - 111%, p <0.0001). Significant differences between study and control physicians also appear in 24 hour compliance (50.4% vs. 29.0%, RRR - 74%, p <0.0001) and hospital-stay compliance (55.9% vs. 37.1%, RRR 51%, p <0.0001).</p> | <p>+</p> |
| <p>Overhage (2001) (Overhage et al. 361-369) Design: RCT N = 34 physicians Implementation: 00/1984 Study Start: 09/1996 Study End: 02/1998</p> | <p>CPOE/POE system Integrated Billing/administratio n system CPOE/POE system, Imaging systems</p> | <p>Ambulatory care, Academic</p> | <p>mean time spent in direct care per patient, minutes*, mean time spent in writing tasks per patient, minutes*</p> | <p>Time spent in direct care with a patient in minutes remained the same in the control (paper-based) and CPOE groups (15.8 vs. 16.1, NS). Time spent on writing tasks in minutes remained the same between groups (6.2 vs. 6.9, NS).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|---|----------|
| <p>Ozdas (2006) (Ozdas et al. 188-196) Design: Before-after N = 540 patients Implementation: 04/2003 Study Start: 08/2002 Study End: 09/2003</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated CDSS/CDS/CCDS/reminders CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary, 630 Beds Academic</p> | <p>rate of order set use for sensitive to AMI patients*, rate of order set use for confirmed AMI patients*</p> | <p>There was a significant increase in ACS order set use after the implementation of the Admission Advisor for 'sensitive to AMI' admissions (60% vs. 70%, RRR -17%, p = 0.009), and a non-significant increase for confirmed AMI patients (46% vs. 64%, RRR -39%, p = 0.07). For all suspected AMI admissions, ACS order set use yielded a significant increase in early aspirin ordering (77% vs. 91.2%, RRR -17%, p = 0.001) and an increase in trend toward significance in beta-blocker ordering (70% vs. 76%, RRR -9%, p = 0.07). A similar non-significant trend in aspirin (89% vs. 97%, RRR -9%, p = 0.07) and beta-blocker (81% vs. 88%, RRR -9%, p = 0.18) ordering behavior associated with a confirmed diagnosis of AMI.</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|------------------------|------------------------|---|----------|
| <p>Palen (2006) (Palen et al. 389-395) Design: RCT N = 26,586 index dispensings Implementation: 00/0000 Study Start: 11/2002 Study End: 10/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated CDSS/CDS/CCDS/reminders CPOE/POE system EHR/EMR system, Pharmacy</p> | <p>Ambulatory care</p> | <p>compliance rate</p> | <p>Difference between the control and intervention group physicians in the overall rate of compliance with ordering the recommended laboratory monitoring for prescribed study medications (NS). Laboratory monitoring was performed as recommended 56.6% of the time in the intervention group compared with 57.1% of the time in the control group (p = 0.31). Improved compliance favored the intervention group (71.2% vs. 62.3% [p = 0.003] for gemfibrozil; 75.7% vs. 73.9% [p = 0.05] for statins, 52.8% vs. 46% for colchicine [p = 0.05]; 42.9% vs. 0% for methotrexate [p = 0.03]).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|---|----------|
| <p>Paterno (2009) (Paterno et al. 40-46) Design: Cohort study N = 71,350 alerts Implementation: 00/1996 Study Start: 02/2004 Study End: 02/2005</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system</p> | <p>Acute care/tertiary, 1633 beds in 2 hospitals Academic</p> | <p>compliance rate with DDI alerts: -overall, -severe alerts, -moderately severe alerts</p> | <p>71,350 alerts were reviewed, of which 39,474 occurred at the non-tiered site and 31,876 at the tiered site. Compliance with DDI alerts was significantly higher at the site with tiered DDI alerts compared to the non-tiered site (29% vs. 10%, p <0.001). At the tiered site, 100% of the most severe alerts were accepted, vs. only 34% at the non-tiered site (p <0.001); moderately severe alerts were also more likely to be accepted at the tiered site (29% vs. 10%, p <0.001).</p> | <p>+</p> |
| <p>Patterson (1998) (Patterson 573-576) Design: Before-after N = 2,013 Patients (cases) Implementation: 00/0000 Study Start: 00/0000 Study End: 01/1998</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system</p> | <p>Acute care/tertiary, 520 Beds Academic</p> | <p>rate of DVT prophylaxis *</p> | <p>The preintervention rate of deep vein thrombosis (DVT) prophylaxis was 85.2%. With the introduction of the computerized reminder, compliance with DVT prophylaxis increased to 99.3% (85.2% vs. 99.3%, p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|---|----------|
| <p>Paul (2006) (Paul et al. 1238-1245) Design: RCT N = 3,529 patients in the RCT and 1,203 in the cohort study Implementation: 00/0000 Study Start: 05/2004 Study End: 11/2004</p> | <p>CDSS/CDS/CCDS/reminders</p> | <p>Acute care/tertiary, 424 Beds Academic</p> | <p>appropriate antibiotic prescribing increased</p> | <p>Appropriate antibiotic prescribing increased for both intention to treat analyzes (64.5% vs. 72.7%, RRR 13%, p <0.05) and for per protocol analyzes (64.5% vs. 85.1%, RRR 32%, p <0.05). The cohort study showed similar increases in improved prescribing (57% vs. 70%, p <0.001).</p> | <p>+</p> |
| <p>Peterson (2005) (Peterson et al. 802-807) Design: Cohort study N = 7,456 Medication orders Implementation: 00/0000 Study Start: 10/2001 Study End: 05/2002</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary, Critical care units (CCU, ICU, NICU) 720 Beds Academic</p> | <p>agreement with system recommended daily dose of psychotropic drugs for control vs. CPOE, incidence of dosing that was 10-fold greater than recommended for control vs. CDSS</p> | <p>The CDSS increased the prescription of the recommended daily dose (29% vs. 19%; RRR 58% p <0.001) reduced the incidence of dosing that was 10-fold greater than recommended (2.8% vs. 5.0%, RRR 48%; p <0.001).</p> | <p>+</p> |
| <p>Peterson (2007) (Peterson et al. 2-40) Design: RCT N = 9,111 medication orders Implementation: 00/0000 Study Start: 12/2005 Study End: 08/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary Critical care units (CCU, ICU, NICU) Emergency department, Not specified Academic</p> | <p>ratio between prescribed and recommended doses</p> | <p>Ratio between the prescribed dose and recommended dose showed that compared to controls the intervention group (reminders) received lower doses (3.0 vs. 2.5, p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|----------------|
| <p>Prescription in Ischaemic Stroke Management (PRISM) Study Group (2003) (Weir) et al. 143-153) Design: RCT N = 1,640 Patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/ reminders Integrated Hospital information system</p> | <p>Unspecified Hospital</p> | <p>relative risk reduction (RRR) in ischemic and hemorrhagic vascular events</p> | <p>Actual therapy prescribed vs. the option of 'no antiplatelet or anticoagulant therapy. Estimated RRR(%) for the control and intervention in the first phase was 16.7 (13.2 to 23.7) vs. 16.3 (15.2 to 21.2) (not significantly different) For the second phase it was 16.3 (13.1 to 23.8) vs. 16.7 (13.5 to 22.9) (NS).</p> | - |
| <p>Quinn (2008) (Quinn et al. 160-168) Design: RCT N = 30 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/ reminders Daibetes Management Tool Integrated Web-based data analytics and therapy optimization tools</p> | <p>Ambulatory care</p> | <p>changes in medication (medication intensified)</p> | <p>Patients using WDS were more likely to have physicians intensify diabetes medications (84.6% vs. 23.08%, p = 0.002).</p> | + |
| <p>Quinzler (2009) (Quinzler et al. 30-35) Design: Before-after N = 20,031 prescribed drugs Implementation: 00/2003 Study Start: 08/2006 Study End: 03/2007</p> | <p>CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated CDSS/CDS/CCDS/ reminders Pharmacy</p> | <p>Acute care/tertiary 1680 Beds Academic</p> | <p>proportion of prescriptions with inappropriate tablet splitting</p> | <p>The CDSS alert resulted in a significant reduction in prescriptions for inappropriate tablet splitting (2.7% vs. 1.4%, RRR 48%, p <0.001).</p> | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|--------------------------------------|--|--|----------|
| <p>Raebel (2005) (Raebel et al. 2395-2401) Design: RCT N = 10,169 dispensings Implementation: 00/0000 Study Start: 09/2002 Study End: 12/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated Laboratory system, Pharmacy</p> | <p>Ambulatory care, HMO pharmacy</p> | <p>percentage of dispensings with baseline monitoring</p> | <p>Recommended laboratory monitoring was completed in 74.7% (n=7,598) of dispensings at initiation of therapy. Compared to the usual care group, monitoring was higher in the intervention group (70% vs. 79%, RRR - 13%, p <0.001).</p> | <p>+</p> |
| <p>Raebel (2007) (Raebel et al. 977-985) Design: RCT N = 59,680 patients Implementation: 00/0000 Study Start: 05/2005 Study End: 05/2006</p> | <p>CDSS/CDS/CCDS/reminders Pharmacy information system Integrated EHR/EMR system</p> | <p>Ambulatory care, HMO pharmacy</p> | <p>new dispensings of targeted medications*, dispensings of targeted medications considered inappropriate*</p> | <p>In the analysis of all dispensings of targeted medications, there was a significant reduction of new dispensings of at least one targeted medication (2.2% vs. 1.8%, RRR 16%, p <0.002). For dispensings of targeted medications considered inappropriate, there was also a significant reduction with the use of the alerting system (1.5% vs. 1.1%, RRR 27%, p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|-------------------------------------|--|---|----------|
| <p>Raebel (2007) (Raebel et al. 440-450) Design: RCT N = 11,100 women Implementation: 00/0000 Study Start: 01/2003 Study End: 04/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system Pharmacy</p> | <p>Ambulatory care HMO pharmacy</p> | <p>the proportion of pregnant women dispensed a category D or X medication*, the total number of first dispensings of targeted medications</p> | <p>The alerts resulted in a 47% reduction in the proportion of pregnant patients receiving category D or X drugs (p <0.001) Intervention patients received 238 dispensings of unique targeted medications and usual care patients received 361 dispensings (p = 0.03). The study was stopped primarily due to 2 false-positive alert types: Misidentification of medications as contraindicated in pregnancy by the pharmacy information system and misidentification of pregnancy related to delayed transfer of diagnosis information.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|--|----------|
| <p>Rasmussen (2005) (Rasmussen et al. 1137-1142) Design: RCT N = 253 patients Implementation: 00/0000 Study Start: 00/2001 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/ reminders Integrated Internet based electronic diary</p> | <p>Ambulatory Care Academic</p> | <p>good compliance (use of medication always or almost always) Internet vs. specialist group</p> | <p>A significant improvement in compliance was observed for all groups, but good compliance was significantly higher (p <0.001) for both the Internet vs. the GP group and the specialist vs. the GP group. 4 of 4 measures of improve prescribing was noted in the internet group and the specialist group. The GP group also improved but to a lesser extent.</p> | <p>+</p> |
| <p>Riggio (2009) (Riggio et al. 124-131) Design: Before-after N = 100 patients with heparin induced thrombocytopenia Implementation: 06/2005 Study Start: 03/2004 Study End: 09/2006</p> | <p>CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated Hospital information system</p> | <p>Acute care/tertiary, 728 Beds Academic</p> | <p>time from platelet count criterion until heparin/enoxaparin stop* Time from platelet count criterion until 1st HIT laboratory test drawn* Time from platelet count criterion until direct thrombin inhibitor started*</p> | <p>Counter to expectations, the time (in days) taken from alert to heparin stop order was significantly higher after implementation (1.3 vs. 2.9, p = 0.04). There were no significant differences in time (in days) from alert to lab test (2.3 vs. 3.0, NS), nor time to start of treatment with direct thrombin inhibitor (19.3 vs. 15.0, NS).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|--|---------|
| Riggio (2009) (Riggio et al. 1719-1726) Design: Before-after N = 2,151 discharge measures Implementation: 00/2001 Study Start: 07/2005 Study End: 03/2008 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system, Hospital information system | Acute care/tertiary, 690 Beds Academic | overall compliance rate* | CDSS yielded a 26% increase in overall compliance with the cardiac discharge measures, from 76.8% in the preintervention period to 96.8% (p <0.001) in the postintervention period. | + |
| Rohrig (2008) (Rohrig et al. 63-68) Design: Before-after N = 156 patients Implementation: 00/1999 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system | Critical care units (CCU, ICU, NICU) 14 bed unit Beds Academic | rate of adequate treatment, rate of inadequate treatment | The frequency of adequate treatment increased from an average 47.8% in the pre-period to 66.5% in the post-period (p <0.01). Rate of inadequate treatment decreased from 34.2% to 18.5%. | + |
| Rollman (2002) (Rollman et al. 493-503) Design: RCT N = 200 Patients with documented major depression Implementation: 00/0000 Study Start: 04/1997 Study End: 12/1998 | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Ambulatory care | antidepressant prescribing rate (secondary) | Prescribing antidepressants (continuous use of change in prescriptions) did not differ across the 3 groups at 3 or 6 months. | - |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|---|----------|
| <p>Rood (2005) (Rood et al. 172-180) Design: RCT N = 484 patients Implementation: 04/2001 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system</p> | <p>Critical care units (CCU, ICU, NICU) 18 Beds Academic</p> | <p>adherence to glucose measurement timing recommendations*, adherence to insulin dose advice*</p> | <p>Rate of compliance with glucose measurement timing recommendations control-intervention-control (29% vs. 38% vs. 41% with period 2 and 3 greater than period 1, p = 0.05). During the intervention period, the rate for computerized group was higher than the control (36% vs. 40%, p = 0.05). Rate of compliance with insulin dose advice was higher in period 2 than 1, and then decreased significantly in period 3 (56% vs. 70% vs. 42%, p = 0.05). During the intervention period the rate for computerized group was higher than the control (64% vs. 77%, p = 0.05).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|-------------------------------------|--------------|---|--|---------|
| Ross (2005) (Ross et al. 410-415) Design: Cohort study N = 190 physicians Implementation: 00/0000 Study Start: 08/2001 Study End: 07/2002 | e-Rx Integrated Formulary | HMO pharmacy | Formulary compliance ratio*, Absolute generic utilization ratio,* Adjusted generic utilization ratio * | No differences between predominantly traditional prescribers and e-prescribers for formulary compliance (82.8% vs. 83.2%, p = 0.32) or absolute generic drug utilization (36.9% vs. 37.3%, p = 0.18) or adjusted generic drug utilization (74.3% vs. 74.7%, p = 0.27). | - |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---------------------------------------|--------------------------------------|--|--|----------|
| <p>Rosser (1992) (Rosser et al. 911-917) Design: RCT N = 8,069 patients Implementation: 00/0000 Study Start: 04/1985 Study End: 03/1986</p> | <p>CDSS/CDS/CCDS/reminders</p> | <p>Ambulatory care, Academic</p> | <p>rate of tetanus toxoid vaccination*</p> | <p>The rates of tetanus toxin given were 3.2% in control, 22.8% in physician reminder, 24% in telephone reminder, and 30.6% in the letter reminder. The differences in the recorded vaccination rate between the randomized control group and the three reminder groups are as follows: 19.6% in the physician reminder group (95% CI 17.1% to 22.2%, p <0.00001), 20.8% in the telephone reminder group (CI 18.3% to 23.5%, p <0.00001) and 27.4% in the letter group (CI 24.8% to 30.2%, p <0.00001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--------------------------------------|---|--|----------|
| <p>Rubin (2006) (Rubin et al. 627-634) Design: Observational study N = 99 primary care physicians Implementation: 01/2002 Study Start: 01/2002 Study End: 03/2004</p> | <p>CDSS/CDS/CCDS/reminders Integrated Handheld, Stand-Alone</p> | <p>Ambulatory care</p> | <p>change in rate of antibiotic prescribing according to recommendations*, change in rate of adherence to NOT prescribe antibiotic recommendations*</p> | <p>Adherence with CDSS recommendations increased from 79.3% in the first one-third of provider's cases to 82.0% in the second two-thirds (an increase of 2.7%; p <0.016). Total adherence was higher with diagnoses for which an antibiotic was not indicated (84.8% vs. 75.7% for diagnoses warranting antibiotics), and providers showed a significant improvement in adherence over time for cases not requiring antibiotics (an increase of 2.7%; p <0.039).</p> | <p>+</p> |
| <p>Safran (1995) (Safran et al. 341-346) Safran (1993) (Safran et al. 224-228) Design: RCT N = 349 patients with HIV Implementation: 00/0000 Study Start: 05/1992 Study End: 09/1993</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated EHR/EMR system</p> | <p>Ambulatory care, Academic</p> | <p>mean response time to alerts* mean response times to reminders*</p> | <p>Physicians who got alerts responded more quickly to them (mean 52 vs. 11 days, p <0.0001). Physicians who got reminders responded more quickly to them (mean 500 vs. 114 days, p = 0.0001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|--------------------------------------|--|---|----------|
| <p>Schnipper (2008) (Schnipper et al. Symposium) Design: Before-after N = 30 clinicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Antiplatelet prescribed or contraindication documented*, Beta-blocker prescribed *, Change in diabetic therapy if A1c >7.0 *</p> | <p>Antiplatelet prescribed or contraindication documented improved from 3.2% in the preintervention to 31.0% in the postintervention period, p <0.001. Beta-blocker prescribed or contraindication documented was 4.2 % in the preintervention compared to 66.7% in the post period, p = 0.03. Change in diabetic therapy if A1c >7.0 was 10.7% in the pre-period and 16.9% in the post period, p = 0.11.</p> | <p>+</p> |
| <p>Scotton (2009) (Scotton et al. 71-76) Design: Before-after N = 283 patients Implementation: 12/2003 Study Start: 03/2003 Study End: 01/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Acute care/tertiary, 606 Beds</p> | <p>proportion of cases with guideline violations</p> | <p>Contrary to expectations, the prescribing guidelines were violated significantly less frequently during baseline (27.4%) than after implementation of the reminder (34.3%), p <0.01.</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|---|----------|
| <p>Segarra-Newnham (2003) (Segarra-Newnham 758-762) Design: Before-after N = 211 Patients Implementation: 00/1997 Study Start: 00/1995 Study End: 07/2001</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Vaccination rate for pneumococcal vaccine*, Vaccination rate for tetanus vaccine*</p> | <p>Vaccination rates for enrolled before 1997 and after 1997 were 100% vs. 97% for pneumococcal vaccine (NS). However the vaccination rate for the same time period for tetanus vaccine was 100% vs. 61% due to national shortage of vaccine after 1997 (p <0.001).</p> | <p>-</p> |
| <p>Sellier (2009) (Sellier et al. 203-210) Design: Time series N = 942 prescriptions Implementation: 00/0000 Study Start: 08/2006 Study End: 08/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated Laboratory system, Pharmacy</p> | <p>Acute care/tertiary, 827 Beds Academic</p> | <p>rate of inappropriate first prescriptions*, overall rate of inappropriate prescriptions, rate of cancellation of prescriptions if no eGFR lab result was available</p> | <p>The rate of inappropriate first prescriptions did not differ significantly between intervention and control periods (19.9% vs. 21.3%, RRR 7%, p = 0.63); nor did the overall rate of inappropriate prescriptions (20.4% vs. 18.5%, RRR 9%, p = 0.37). The rate of cancellation of prescriptions if no eGFR lab result was available also did not differ between control and intervention periods (31.3% vs. 35%, RRR-12%, p = 0.62).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|---|--|----------|
| <p>Shiffman (2000) (Shiffman et al. 767-773) Design: Before-after N = 9 physicians Implementation: 00/0000 Study Start: 09/1996 Study End: 10/1998</p> | <p>CDSS/CDS/CCDS/ reminders Handheld</p> | <p>Ambulatory care</p> | <p>Adherence rate with metered-dose inhaler/nebulization*, rate of systemic corticosteroid prescriptions*</p> | <p>Adherence with metered-dose inhaler/nebulization rates did not differ between control and intervention (73% vs. 91%, NS), nor did rate of prescribing systemic corticosteroids (43% vs. 57%, NS).</p> | <p>-</p> |
| <p>Shojania (1998) (Shojania et al. 554-562) Design: RCT N = 396 physicians Implementation: 00/0000 Study Start: 06/1996 Study End: 03/1997</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system, Imaging systems, Laboratory system, Pharmacy</p> | <p>Acute care/tertiary, 720 Beds Academic</p> | <p>number of vancomycin orders/ prescriber*, mean duration of treatment prescribed per physician*, mean number of days of vancomycin per course of treatment*</p> | <p>The total number of orders for vancomycin for physicians in the control group was higher than in the intervention group (16.7 vs. 11.3 orders per physician, p = 0.04). Physicians in the intervention group prescribed vancomycin for 36% fewer days than physicians in the control group (26.5 vs. 41.2, p = 0.05). The number of days of vancomycin per course of treatment was also lower for the physicians in the intervention group, mean of 1.8 vs. 2.0 for the control group (p = 0.05).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|----------------------|---|---------|
| Shu (2001) (Bates et al. 965) Design: Before-after N = 44 Physicians (Interns) Implementation: 11/01998 Study Start: 09/1998 Study End: 06/1999 | CPOE/POE system Integrated Hospital information system | Acute care/tertiary, 820 Beds | time spent ordering* | The percentage of total time spent writing orders by medical interns between pre-CPOE and post-CPOE period increased from 2.1% to 9.0% (p <0.001). | - |
| Shulman (2005) (Shulman et al. R516-R521) Design: Time series N = 3,465 prescriptions over 4 times points Implementation: 04/2002 Study Start: 09/2001 Study End: 12/2002 | CPOE/POE system Integrated Hospital information system | Critical care units (CCU, ICU, NICU) 22 (in the ICU) Beds Academic | rate of ME* | The proportion of MEs before CPOE was 6.7% and 4.8% after CPOE introduction (RRR 28%, p <0.04) The proportion of MEs with CPOE varied over time after its introduction (p <0.001). Evidence also indicated the strong linear trend of a declining proportion of MEs over time (p <0.001). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|---|----------|
| <p>Silveira (2007) (Delgado et al. 223-230) Design: Before-after N = 4,814 orders (treatment lines) Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>e-Rx Integrated EHR/EMR system, Pharmacy</p> | <p>General Hospital, 53 beds in 2 wards of the hospital Beds</p> | <p>rate of errors: -medication data, -dose, -administration frequency/time, -route of administration, -nursing transcription</p> | <p>The EP system was associated with a lower rate of errors compared with the manual system for medical data (38% vs. 8%, RRR 79%, p <0.05), dosage (29% vs. 2%, RRR 92%, p <0.05), administration frequency/time (6% vs. 1%, RRR 83%, p <0.05) and route of administration (17% vs. 0%, RRR 99%, p <0.05). Nursing transcription errors were increased (18% vs. 21%, RRR 17% p <0.05) while drug interaction (2% vs. 3%) and treatment duration errors (1% vs. 1%) remained the same.</p> | <p>+</p> |
| <p>Sintchenko (2005) (Sintchenko et al. 398-402) Design: Before-after N = not reported n/a Implementation: 10/2002 Study Start: 04/2002 Study End: 03/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated Laboratory system</p> | <p>Acute care/tertiary, Critical care units (CCU, ICU, NICU) 800 (18 bed ICU) Beds Academic</p> | <p>antibiotic consumption (defined daily doses/1,000 patient days)*</p> | <p>Consumption of antibiotics in defined daily doses/1,000 patient days decreased significantly after implementation of the hand-held decision support tool (1,767 vs. 1,458, p = 0.04).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|----------------------------------|--------------------------------------|--|--|----------|
| <p>Small (2008) (Small, Barrett, and Price 181-187) Design: Cross-sectional N = 1,941 prescriptions Implementation: 01/2003 Study Start: 01/2005 Study End: 09/2005</p> | <p>CPOE/POE system</p> | <p>Acute care/tertiary, Academic</p> | <p>error rate*, types of errors, severity of errors, error rates among prescribers</p> | <p>For error rates using computerized vs. spreadsheets indicated a relative risk reduction of 42% (20% vs. 12%, RRR 42%, p <0.0001) The distribution of type of error differed significantly according to prescription method (p <0.001) and the distribution of severity of errors also differed significantly according to prescribing method (p <0.001).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|---|----------|
| <p>Smith (2006) (Smith et al. 1098-1104) Design: Time series N = no sample size given number of dispensings Implementation: 09/2000 Study Start: 10/1999 Study End: 12/2002</p> | <p>CDSS/CDS/CCDS/reminders Integrated CDSS/CDS/CCDS/reminders CPOE/POE system EHR/EMR system</p> | <p>Ambulatory care</p> | <p>number of dispensing of non-preferred drugs/10,000 population in elderly patients, number of dispensing of preferred drugs/10,000 population in elderly patients, number of dispensing of non-preferred drugs/10,000 population in non-elderly patients</p> | <p>Following the implementation of the drug-specific alerts, a large and persistent reduction (5.1 prescriptions per 10,000, $p = 0.004$) a 22% relative decrease from the month before alert implementation, in the exposure of elderly patients to nonpreferred medications was observed. We found no evidence of a decrease in use of nonpreferred agents for nonelderly patients. There was an upward, though non-significant trend in the use of preferred agents in elderly patients following the intervention ($p = 0.66$).</p> | <p>-</p> |
| <p>Sobieraj (2008) (Sobieraj 1755-1760) Design: Before-after N = 101 patients Implementation: 03/2007 Study Start: 07/2006 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated CPOE/POE system</p> | <p>Acute care/tertiary, 819 Beds Academic</p> | <p>compliance with ordering VTE prophylaxis</p> | <p>The addition of alerts for patients at risk of VTE and an education program resulted in a significant improvement in compliance with ordering VTE prophylaxis (49% vs. 93%, RRR -90%, $p < 0.001$).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|----------------------------------|---|---------|
| Spencer (2005) (Spencer et al. 416-419) Design: Before- after N = 5,063 medication errors Implementation: 10/2002 Study Start: 01/2002 Study End: 05/2003 | CPOE/POE system | Acute care/tertiary, 688 Beds Academic | reported errors per discharge | Implementation of CPOE on the two units was associated with a significant increase in reported errors, from 0.068 per discharge before CPOE implementation to 0.088 per discharge afterward (p = 0.01). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|------------------------|---|---|----------|
| <p>Steele (2005) (Steele et al. e255) Design: Before-after N = 54,206 patient visits Implementation: 12/2002 Study Start: 08/2002 Study End: 04/2003</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system, Laboratory system</p> | <p>Ambulatory care</p> | <p>percentage of: -time provider ordered the rule-associated laboratory test (for which alert was triggered and message displayed), - times medication order triggered but not completed (for an abnormal laboratory value), -times the provider ordered the rule-associated laboratory test (for alert that was triggered for a missing laboratory test)</p> | <p>Medication orders for which an alert was presented shows an increase in the percentage of time the provider ordered the rule-associated laboratory test (38.5% vs. 51.1%, p, 0.001) .When alert was for an abnormal laboratory value, percentage of times medication order triggered but was not completed increased from 5.6% at baseline to 10.9% during the intervention (p = 0.03). The largest effect was noticed when the alert was triggered for a missing laboratory test, the percentage of times the provider ordered the rule-associated laboratory test increased from 43.0% at baseline to 62.0% (p <0.001). All other outcomes did not have statistically significant change.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|-----------------------------|---|---|----------|
| <p>Stone (2009) (Stone et al. 960-967) Design: Before-after N = 18,884 procedures Implementation: 05/2007 Study Start: 12/2006 Study End: 05/2008</p> | <p>CPOE/POE system</p> | <p>Unspecified Hospital</p> | <p>medication error rates, Mean total time from placement of order to nurse receipt</p> | <p>Medication error rates did not decrease significantly from preimplementation to 6 or 12 months postimplementation (0.22% vs. 0.16 % vs. 0.21%, p = NS). Mean total time from placement of order to nurse receipt before implementation was significantly reduced (41.2 minutes vs. 27 seconds, p <0.01).</p> | <p>-</p> |
| <p>Tamblyn (2003) (Tamblyn et al. 549-556) Design: RCT N = 12,560 Patients Implementation: 00/0000 Study Start: 01/1997 Study End: 02/1998</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>rate of initiation of inappropriate drugs per 1,000 visits, Rate of discontinuation of inappropriate drugs per 1,000</p> | <p>During the study the number of new potentially inappropriate prescriptions per 1,000 visits was lower (52.2 v 43.8) in the CDS group than in the control group (RR 0.82, 95% CI 0.69 to 0.98). The rate of discontinuation of inappropriate drugs per 1,000 was not different: 67.4 vs. 71.4, (RR 1.06, 95% CI 0.089 to 1.26).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|------------------------|--|---|----------|
| <p>Tamblyn (2010) (Tamblyn et al. 176-188) Design: RCT N = 2,293 patients Implementation: 00/0000 Study Start: 04/2006 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Insurance, provincial beneficiary and prescription databases</p> | <p>Ambulatory care</p> | <p>rate of drug profile review, Changes in therapy</p> | <p>Significant increase in drug profile review in the intervention compared to the control group (44.5% vs. 35.5%;p <0:001). There was no statistically significant difference between the intervention and control group in the proportion of patients who had increases in therapy (28.5% vs. 29.1%; OR 0.98; p = 0.86).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--------------------------------------|--|--|----------|
| <p>Tang (1999) (Tang et al. 115-121) Design: Time series N = 2,484 patient visits Implementation: 07/1996 Study Start: 10/1995 Study End: 01/1998</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care, Academic</p> | <p>compliance rates with vaccination guidelines*-computer users compliance rates with vaccination guidelines*-paper users</p> | <p>Compliance rates did not increase in the first year for either group. For the computer users, compliance rates steadily increased year 2 to year 3 to year 4 (38.7% vs. 60.9%, RRR -57%, p = 0.001; 60.9% vs. 68.2%, RRR -12%, p = 0.02). For the paper group, year 2 to 3 saw a significant increase (28.5 vs. 37.0, p = 0.02), but year 3 to 4 saw a significant decrease (37.0% vs. 30.6%, p = 0.03). No comparisons between paper and computer were performed by the authors.</p> | <p>+</p> |
| <p>Tang (2009)¹⁶² Design: Before-after N = 1,762 patients Implementation: 01/2005 Study Start: 09/2004 Study End: 06/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system, Laboratory system, Pharmacy</p> | <p>Ambulatory care</p> | <p>overall compliance rate, pregnancy test ordering, Charting of cumulative dose, liver function and lipid profile test ordering</p> | <p>Introduction of e-isotretinoin chart resulted in marked improvement in physician compliance to all steps of the isotretinoin prescription process, with the overall compliance rate increasing from 57.5% to 97.8% (p <0.05) in the first year post-implementation. Of the female patients, 100%</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|---|----------|
| | | | | <p>were tested for pregnancy prior to starting isotretinoin therapy, an increase of 66.0% compared with the pre-implementation period (p <0.05). Charting of cumulative dose improved (an increase of 13.5% to 99.5%, p <0.05) so did liver function tests and lipid investigations (an increase of 3.8% to 100%). The results demonstrated close to 100% compliance with charting of cumulative dose of isotretinoin, pregnancy testing, liver function and lipid profile tests. The results sustained for more than 2 years from January 2005 to June 2007 [no analysis given past 1 year].</p> | |
| <p>Teich (2000)¹⁶³ Design: Time series N = not reported orders Implementation: 10/1993 Study Start: 00/0000 Study End:</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system, Imaging systems, Laboratory system</p> | <p>Acute care/tertiary, 720 Beds Academic</p> | <p>H2-blocker orders, variability in dosages, frequency of administration and exceeding maximum dosages, proportion of orders for 3x IV ondansetron, compliance with heparin ordered consequent to</p> | <p>Study 1: Nizatidine was used for <20% of all oral H2-blocker orders before implementation of the alert, vs. >95% after wards (p <</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---------------------|----------------------------------|----------|-------------------|---|---------|
| 00/0000 | | | bed rest | <p>0.001); this was sustained for year 1 and 2. The use of IV ranitidine increased from 0% before the intervention to 71% of intravenous H2-blocker orders (32/45) in the first week and to 97% or more from the fourth week onward.</p> <p>Study 2: Variability in standard deviation dosages across medications reduced by 11% following implementation of the dosage guidance application (p <0.001). Maintained over 3 years followup. Standard deviation of frequency of administration reduced by 30% post-implementation (p <0.001) and proportion of orders exceeding maximum dose decreased significantly from 2.1% to 0.56% post-implementation (p <0.001).</p> <p>Study 3: Orders for 3x IV ondansetron increased significantly after the</p> | |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|---|--|---------|
| | | | | preferred order was highlighted in the dosing list (5.9% vs. 93.5%, RRR - 1485%, p <0.001). Study 4: heparin ordering with bed rest increased from 23.9% to 46.9%. | |
| Terrell (2009) ¹⁶⁴ Design: RCT N = 5,162 Patients Implementation: 00/0000 Study Start: 01/2005 Study End: 07/2007 | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system | Acute care/tertiary, 450 Beds Academic | proportion of ED visits by seniors with an inappropriate medication, proportion of medications that were potentially inappropriate was also reduced | The decision support reduced the proportion of ED discharges that resulted in potentially inappropriate prescriptions (3.9% vs. 2.6%; p = 0.02; OR 0.55, 95% CI 0.34 to 0.89). The proportion of medications that were potentially inappropriate was also reduced, from 5.4% to 3.4% (p = .006; OR 0.59, CI 0.41 to 0.85). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|--|----------|
| <p>Terrell (2009)¹⁶⁵ Design: RCT N = 5,162 patient visits to 63 physicians Implementation: 00/0000 Study Start: 01/2005 Study End: 07/2007</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system</p> | <p>Acute care/tertiary, 450 Beds Academic</p> | <p>visits with an inappropriate medication prescription*, prescriptions that were inappropriate, n (%)</p> | <p>Primary Outcome: Decision support significantly reduced the proportion of ED discharges that resulted in a potentially inappropriate prescription (3.9% vs. 2.6%; p = 0.02; OR 0.55, 95% CI 0.34 to 0.89. This difference represents an absolute RR of 1.3% (95% CI 0.4 to 2.3). Secondary Outcome: When analyzed as a percentage of all medications prescribed by physician subjects, the proportion of medications that were potentially inappropriate was significantly reduced, from 5.4% to 3.4% (p = 0.006; OR 0.59, 95% CI 0.41 to 0.85), with an absolute reduction of 2.0% (95% CI 0.7 to 3.3).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|--|---------|
| Tierney (2003) ¹⁶⁶ Design: RCT N = 706 patients, 20 pharmacists, 94 physicians and 1 nurse practitioner Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Pharmacy | Ambulatory care, Outpatient hospital based Academic | compliance with cardiac care suggestions* | Neither the physician nor the pharmacist intervention had any significant effect on whether patients' cardiac care was compliant with the suggestions (p >0.8 across the 4 intervention groups by analysis of variance, with p > 0.7 and p >0.4 when testing the physician and pharmacist interventions separately). | - |
| Tierney (2005) ¹⁶⁷ Design: RCT N = 706 patients Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996 | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system EHR/EMR system, Pharmacy | Ambulatory care, Pharmacy Outpatient hospital based Academic | adherence to the care suggestions* | There were no differences between the four study groups in either adherence to the care suggestions, combined or individually (32% control, 32% physician intervention, 32% pharmacist intervention, 37% both interventions, NS). | - |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|---------|
| Upperman (2005) ¹⁶⁸ Design: Before-after N = Not reported ADE/1,000 doses Implementation: 00/2002 Study Start: 01/2002 Study End: 0/0000 | CPOE/POE system Integrated EHR/EMR system | Acute care/tertiary, Pediatric stand alone hospital, Academic | ADE rates per 1,000 before and after CPOE implementation | All ADEs before CPOE were 0.3 per 1,000 doses, whereas after CPOE ADEs were 0.37 per 1,000 doses (p = 0.3). | + |
| Uttaro (2007) ¹⁶⁹ Design: Cohort study N = 23 psychiatrists Implementation: 01/2004 Study Start: 01/2004 Study End: 03/2005 | CDSS/CDS/CCDS/reminders Stand-Alone New York State Office of Mental Health intranet, pharmacology resources | Other specialty hospital (rehab, oncology) | percentage of caseloads on 2 or more antipsychotics*, overall percentage of patients on 2 or more antipsychotics | Overall, there were moderately large reductions for most psychiatrists in the percentage of caseloads on 2 or more antipsychotics (56% vs. 36%, RRR 36%, p <0.01). There were significantly greater reductions in March 2005 for psychiatrists who had higher percentages of their caseloads on two or more concurrent antipsychotics in January 2004. The overall percentage of patients on 2 or more antipsychotics dropped significantly (54% vs. 36%, RRR 33%, p <0.01). | + |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|--|----------|
| <p>van Doormaal (2009)¹⁷⁰ Design: Time series N = 1,195 patients Implementation: 00/0005 Study Start: 07/2005 Study End: 05/2008</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Barcoding system, Pharmacy</p> | <p>Acute care/tertiary, 1,300 (Groningen); 600 (Tilburg and Waalwijk) Beds Academic</p> | <p>medication errors (ME)* preventable adverse drug events (pADEs)*</p> | <p>During the baseline period, 55% of all medication orders (MOs) contained at least one or more MEs, whereas during the postintervention period this was 17%; a significant immediate absolute reduction of 40.3% (95% CI: -45.13% to 35.48%). In the baseline period, 15.5% of admitted patients experienced one or more pADE, as opposed to 7.3% in the postintervention period. Decrease could not be attributed to CPOE/CDSS. The immediate change was NS (-0.42%, 95% CI: -15.52% to 14.68%) because of the observed underlying negative trend during the pre-CPOE period of -4.04% [95% CI: -7.70% to 0.38%] per month.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|--|----------|
| <p>Van Wyk (2007)¹⁷¹ Design: RCT N = 87,860 Patients Implementation: 00/0000 Study Start: 05/2004 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Percentage of patient treated</p> | <p>Of the patients requiring treatment, 66% were treated in the alerting arm, 40% in the on-demand arm, and 36% in the control arm. After adjustment for differences between arms, the likelihood of being treated was 40% higher in the alerting arm (adjusted RR = 1.40; 95% CI 1.15 to 1.70) and 19% higher (NS) in the on-demand arm in comparison to the control arm (adjusted RR = 1.19; 95% CI 0.94 to 1.50). A similar pattern was shown for the need for screening within the 3 groups.</p> | <p>+</p> |
| <p>Varkey (2007)¹⁷² Design: Cross-sectional N = 4,527 prescriptions Implementation: 00/0000 Study Start: 00/1996 Study End: 00/2002</p> | <p>CPOE/POE system Integrated CDSS/CDS/CCDS/reminders</p> | <p>Ambulatory care, Other institution based</p> | <p>frequency of intercepted prescription errors*</p> | <p>Statistically significant decrease in frequency of intercepted prescription errors among handwritten and computerized prescriptions was observed (7.4% vs. 4.9%, p = 0.0048).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|---|----------|
| <p>Voeffray (2006)¹⁷³ Design: Before-after N = 2,445 prescriptions Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Pharmacy</p> | <p>Acute care/tertiary, 850 Beds Pharmacy, Inpatient hospital based, Academic</p> | <p>Rate of error*</p> | <p>The average monthly error rate was 15% (95% CI 13% to 18%). After introduction of the CPOE system, the average monthly error rate (which included both computer orders and handwritten, amounted to 13% (95% CI 10% to 16%). This decrease in rate was not statistically different from the rate observed in the first period (p = 0.36). Postimplementation errors in the computerized group only was 0.6% (95% CI 0.3% to 1.4%).</p> | <p>-</p> |
| <p>Walsh (2008)¹⁷⁴ Design: Time series N = 627 admissions Implementation: 04/2002 Study Start: 09/2001 Study End: 05/2003</p> | <p>CPOE/POE system Integrated CDSS/CDS/CCDS/reminders</p> | <p>Critical care units (CCU, ICU, NICU) General Hospital, 59 Pediatric beds</p> | <p>rates non-intercepted serious medical errors*</p> | <p>The rates of errors did not differ for all errors (44.7 before vs. 50.9 errors per 1,000 patient days after COPE, NS), non-intercepted serious medical errors (23.1 before vs. 20.6 per 1,000 patient days after CPOE, NS), or serious medical errors (31.7 before vs. 33.0 per 1,000 patient days after CPOE, NS).</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|---|--|----------|
| <p>Were (2009)¹⁷⁵ Design: Before-after N = 40 Patients Implementation: 00/0000 Study Start: 12/2007 Study End: 04/2008</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated EHR/EMR system, Imaging systems, Laboratory system</p> | <p>Acute care/tertiary, 264 Beds Academic</p> | <p>acceptance of all recommendations, rate of acceptance of pharmacological recommendations</p> | <p>More recommendations were implemented in the reminders group (59% vs. 78%, RRR 32%, p = 0.01) The rate of acceptance of pharmacological recommendations was similar (51% vs. 77%).</p> | <p>+</p> |
| <p>Wilkes (2009)¹⁷⁶ Design: Before-after N = 84 patients Implementation: 06/2005 Study Start: 06/2005 Study End: 05/2006</p> | <p>CDSS/CDS/CCDS/reminders Integrated EHR/EMR system, Laboratory system</p> | <p>Acute care/tertiary, Pediatric stand alone hospital, 418 Beds</p> | <p>prescription rate among eligible patients, prescription rate -off-label</p> | <p>The rate of oseltamivir prescription did not change significantly for patients eligible for the drug (40% vs. 25%, RRR 38%, p = NS), or for off-label prescribing for patients not eligible for the drug (4% vs. 5%, RRR -24%, p = NS) following the implementation of a computerized reminder.</p> | <p>-</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|---------------------------------------|---|---|----------|
| <p>Wrona (2007)¹⁷⁷ Design: Observational study N = 536 PCA patients Implementation: 00/2003 Study Start: 01/2003 Study End: 03/2004</p> | <p>CPOE/POE system, Integrated EHR/EMR system, Imaging systems, Laboratory system</p> | <p>Pediatric stand alone hospital</p> | <p>rates of respiratory monitoring rates of oxygen saturation monitoring</p> | <p>Compared to the control group of 'no order set', patients in the Acute Pain Team Service had a higher rate of respiratory monitoring (43% vs. 66.3%, RRR - 54%, p <0.05) and oxygen saturation monitoring (86.1% vs. 98.6%, RRR - 15%, p <0.05). Compared to the control group of 'no order set', patients in the prescriber initiated PCA had higher respiratory rate monitoring (43% vs. 57.8%, RRR - 34%, p <0.05). No other comparisons were significant.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|---|--|----------|
| <p>Xamplas (2010)¹⁷⁸ Design: Before-after N = 96 patients Implementation: 02/2008 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/reminders Integrated CPOE/POE system, e-MAR, Pharmacy</p> | <p>Acute care/tertiary, 465 Beds Inpatient hospital based, Academic</p> | <p>Piperacillin–tazobactam days per 1,000 patient-days*, Piperacillin–tazobactam doses per 1000 patient-days*</p> | <p>While the number of piperacillin–tazobactam days per 1,000 patient days did not significantly change (124 ± 6.3 vs. 121 ± 12.6, $p = 0.389$) during the preintervention and postintervention periods, there was a significant reduction in the number of piperacillin–tazobactam doses per 1,000 patient-days during the postintervention period (457 ± 33.3 vs. 341 ± 35.7, $p < 0.001$).</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|----------------------------------|--|--|---|----------|
| <p>Yu (2009)¹⁷⁹ Design: Cross-sectional N = 3,364 hospitals Implementation: 00/0000 Study Start: 07/2003 Study End: 06/2004</p> | <p>CPOE/POE system</p> | <p>Unspecified Hospital, Not specified</p> | <p>11 medication quality indicators*</p> | <p>Among the 11 medication-related measures for acute myocardial infarction, heart failure and pneumonia, the mean performance on 6 cardiovascular-related measures was higher among CPOE hospitals ($p < 0.001$) vs. the comparison (nonCPOE) hospitals. Also, for one pneumonia measure, administering “Antibiotics within 4 hours of arrival for patients with pneumonia,” performance was lower for hospitals with full CPOE implementation ($p < 0.001$). Four quality indicators were not significantly different among the groups; 3 for pneumonia and administration of thrombolytic agent within 30 minutes for AMI. The differences are maintained when hospital teaching status and ownership and number of beds are taken into account.</p> | <p>+</p> |

Evidence Table 1. KQ1: Articles assessing primary process outcomes for all technologies assisting the prescribing phase of medication management (continued)

| Article Information | HIT Studied Integrated system | Settings | Outcomes Measured | Results | Outcome |
|---|---|-------------------------------------|--|---|----------------|
| Zanetti (2003) ¹⁸⁰ Design: RCT N = 273 patients Implementation: 00/0000 Study Start: 03/2000 Study End: 06/ 2000 | CDSS/CDS/CCDS/ reminders Integrated Hospital information system | Acute care/tertiary, Academic | appropriate redosing of antibiotics* | More patients in the alarm plus reminder group received appropriate redosing of antibiotics after > 240 minutes in surgery (adjusted OR 3.31, 95% CI 1.97 to 5.56, p <0.0001). | + |
| Zhan (2006) ¹⁸¹ Design: Mixed methods N = 138,922 number of errors/100,000 doses Implementation: 00/0000 Study Start: 01/2003 Study End: 12/2003 | CPOE/POE system | Unspecified Hospital | number of errors reported per 100,000 doses-inpatients*, number of errors reported per 100,000 doses-outpatients* | The number of errors reported per 100,000 doses was not different among non-CPOE (n=339) and CPOE facilities (n=120) for inpatients (mean 56 vs. 55, p = 0.9) or outpatients (mean 60 vs. 57, p = 0.8). | - |

Evidence Table 2. KQ1: primary process outcomes for all technologies assisting order communication

| Article Information | HIT Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|---|---------|
| Astrand (2009) ¹⁸² Design: Cross-sectional N = 14,365 prescriptions Implementation: 00/0000 Study Start: 02/2006 Study End: 03/2006 | e-Rx, e-Transmission-of the prescription to/from doctor to pharmacy | Pharmacy, Other mail/email in pharmacies | proportion of new prescriptions needing clarification* | Clarification contacts were made for 2.0% (147/7532) of new e-Prescriptions and 1.2% (79/6833) of new non-electronic prescriptions. RR 1.7 (95% CI 1.3 to 2.2) Increased RR was mainly due to 'Dosage and directions for use', RR 7.6 (95% CI 2.8 to 20.4) when compared to other clarification contacts. In all, 89.5% of the suggested pharmacist interventions were accepted by the prescriber, 77.7% (192/247) as suggested and an additional 11.7% (29/247) after a modification during contact with the prescriber. | - |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: CCDS = Computerized Clinical Decision Support; CDIX = Critical Drug Interaction Alert Program; CDS = Clinical / Computerized Decision Support; CDSS = Clinical Decision Support System; CI = Confidence interval; CPOE = Computerized Provider Order Entry; EHR = Electronic Health Record; e-MAR = Electronic Medication Administration Record; EMR = Electronic Medical Records; e-RX = Electronic Prescribing; e-TAR = Electronic Treatment Authorization Request; GP = General Practitioner; HIT = Health Information Technology; HMO = Health Maintenance Organization; N = Sample Size; NS = Not specified; OSUH = Ohio State University Health System; p = Probability; POE = Provider Order Entry; RCT = Randomized Controlled Trial; RR = Relative Risk; RRR = Relative Risk Reduction; vs. = Versus

Evidence Table 2. KQ1: primary process outcomes for all technologies assisting order communication (continued)

| Article Information | HIT Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|---|---------|
| Beer (2002) ¹⁸³ Design: Cross-sectional N = 836 Medication orders Implementation: 00/0000 Study Start: 06/2002 Study End: 06/2002 | CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Hospital information system | Pharmacy, Outpatient hospital based Academic | mean time required to complete prescription review for OpTx order * | The mean time required to complete the prescription review for an OpTx order was 11.11 min (95% CI 10.1 to 12.1; n = 140) compared to the mean time to review a paper order of 5.96 min (95% CI 5.6 to 6.4, p <0.001; n = 696). Therefore, the mean time required to review an order was increased by 5.15 min with the implementation of the direct electronic order entry system. | - |
| Ekedahl (2004) ¹⁸⁴ Design: Cross-sectional N = 24,6991 prescriptions Implementation: 00/0000 Study Start: 05/2000 Study End: 10/2001 | e-Rx Integrated Pharmacy | Pharmacy, Not specified | rate of non compliance (unclaimed "all other" prescriptions vs. unclaimed e- Prescriptions) | Rate of non compliance between unclaimed "all other" prescriptions 369/322754 (0.01%) vs. unclaimed e-Prescriptions 2,171/91,704 (2.37%). | - |
| Halkin (2001) ⁶⁵ Design: Time series N = 775,186 prescriptions Implementation: 11/1997 to 00/1998 Study Start: 01/1998 Study End: 06/1999 | CDSS/CDS/CCDS/reminders Integrated, Pharmacy | Pharmacy, HMO pharmacy | rate of drug interaction prescriptions when 90% of pharmacies and 50% of physicians compared with baseline, rate of drug interaction prescriptions when 95% of pharmacies and 90% of physicians compared with baseline | Dispensing of drug interaction prescriptions reduced by 21.1% and by 67.5% in periods II and III compared with period I (OR, 0.79; 95% CI 0.75 to 0.83 and OR, 0.28; 95% confidence limit, 0.26 to 0.30, respectively). | + |
| Humphries (2007) ¹⁸⁵ Design: Before-after N = not given prescriptions Implementation: 02/2002 Study Start: 07/2000 Study End: 05/2005 | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system Pharmacy | Ambulatory care, Outpatient hospital based | proportion of co-dispensings for interacting drugs per 10,000 prescriptions | The proportion of prescriptions of any of the 8 drug pairs decreased after implementation of CDIX for all 8 drugs (21.3 to 14.7 per 10,000 prescriptions, RRR 31%, (CI 12.7 to 49.5, p = 0.01). | + |

Evidence Table 2. KQ1: primary process outcomes for all technologies assisting order communication (continued)

| Article Information | HIT Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|--|---------|
| Johnson (2010) ⁷⁵ Design: RCT N = 3,285 patients Implementation: 00/0000 Study Start: 04/2007 Study End: 08/2007 | CDSS/CDS/CCDS/ reminders e-Rx Integrated EHR/EMR system | Ambulatory care, Pharmacy, Not specified, Academic | rate of callbacks generated* | There was no significant difference in the callback rates between the "SYW off" and the "SYW on" periods (0.4% vs. 0.45%; p = 0.47). | - |
| Mahoney (2007) ⁹⁹ Design: Before-after N = 2,843,135 inpatient medication orders Implementation: 02/2002 Study Start: 02/2002 Study End: 06/2006 | CDSS/CDS/CCDS/ reminders CPOE/POE system, Pharmacy information system Integrated EHR/EMR system Hospital information system | General Hospital, Pediatric stand alone hospital, 966 in 2 hospitals Beds Pharmacy Inpatient hospital based Academic | rate of -drug allergy violations*, -excessive doses*, -incomplete or unclear orders*, - therapeutic duplication* | Medication errors decreased after implementation of the CIT with respect to drug allergy violations (OR 0.14, CI 0.11 to 0.17, p <0.001), excessive doses (OR 0.68, CI 0.62 to 0.74, p <0.001) and incomplete or unclear orders (0.35, CI 0.32 to 0.38, p <0.001), but no decrease in therapeutic duplications. Turnaround time between drug ordering and administration decreased from 90 minuets to 11 minutes, NR. The override rate also decreased (7.1 to 2.9%, RRR 59%, p = 0.001). | + |
| Mekhjian (2002) ¹⁸⁶ Design: Before-after N = 28,898 patients Implementation: 05/2000 Study Start: 02/2000 Study End: 01/2001 | CPOE/POE system, e-Medication administration system (e-MAR, e-TAR) Integrated Dietary system EHR/EMR system Imaging systems Laboratory system | Acute care/tertiary, Other specialty hospital (rehab, oncology) Academic | medication turn-around time, proportion of verbal orders countersigned, rate of transcription errors | Combining the data showed that time from initiation of the prescription and administration was reduced after POE: mean 5:28 hours before vs. 1:51 hours after, RRR 64%, p <0.001. The proportion of signed verbal orders increased for both hospitals: OSUH 56.4% vs. 76%, RRR 76%, p <0.001 and James Cancer 72.8% vs. 99.0, RRR 36%, p <0.001. The volume of transcription errors was reduced after POE from 11.3% to 0%, RRR 100%, p <0.001. | + |

Evidence Table 2. KQ1: primary process outcomes for all technologies assisting order communication (continued)

| Article Information | HIT Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|---|---|---------|
| Mitchell (2004) ¹⁸⁷ Design: Cross-sectional N = 4,297 prescriptions Implementation: 0/1999 Study Start: 09/2002 Study End: 12/2002 | e-Medication administration system (e-MAR, e-TAR) e-Rx Integrated Formulary Pharmacy | Acute care/tertiary, Academic | 15 aspects of data completeness for e-MAR were sought with implementation of the e-MAR. | e-MAR was more accurate (more inclusion of important information) for nurses 9 of the 15 were statistically significantly improved including presence of dosing recommendations (30% v3 99%, RRR 230%, p <0.01) Errors detected by the pharmacist did not differ before and after implementation of the e-Rx system. Only minor errors were reduced with the system. | + |
| Murray (1999) ¹⁸⁸ Design: Cohort N = 11,102 observations of 28 pharmacists Implementation: 03/1995 Study Start: 11/1995 Study End: 01/1996 | Pharmacy information system Integrated EHR/EMR system Imaging systems Laboratory system | Acute care/tertiary, Pharmacy, Inpatient hospital based | distribution of pharmacist time on activities, functions and contacts*. | The electronic guidelines and reminders were associated with the overall distribution of activities (more time discussing information and less time checking and preparing prescriptions) p <0.001; overall functions (more time advising or discussing information or problem solving and less time filling prescriptions) p <0.001 and distribution of contacts (more time with other pharmacy personnel, patients, and clinicians and less time working alone) p <0.001. | + |
| Nam (2007) ¹⁸⁹ Design: Before-after N = 39 Patients Implementation: 00/0000 Study Start: 06/2003 Study End: 05/2005 | CPOE/POE system Integrated EHR/EMR system Laboratory system | Emergency department | Time to arrival to tPA treatment in minutes* | Time from arrival to tPA treatment was reduced by 23 minutes (from 79 to 56 minutes; p <0.01). | + |
| Nilsson (2007) ¹⁹⁰ Design: Cohort study N = 2,563 prescriptions (acute) Implementation: 00/0000 Study Start: 02/2005 Study End: 03/2005 | e-Rx Integrated Pharmacy | Pharmacy, Other | rate of prescription pick up by patients within 5 days* | e-RX accounted for 84% of the prescriptions. Among the patients with e-prescriptions 91% picked up their prescriptions in 5 days compared to 85% in the paper group, (RRR -7%, p <0.01). | + |

Evidence Table 2. KQ1: primary process outcomes for all technologies assisting order communication (continued)

| Article Information | HIT Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|--|---|---------|
| Pearce (2010) ¹⁹¹ Design: Before-after N = 332 medication refill orders Implementation: 05/2006 Study Start: 02/2006 Study End: 03/2007 | e-Rx Integrated EHR/EMR system Pharmacy | Ambulatory care, Pharmacy, Pharmacy chain | time to a response for refill request* | The average time to a response to a pharmacy refill request decreased from 1.57 days to 1.04 days (p <0.004). | + |
| Senholzi (2003) ¹⁹² Design: Before-after N = 349 pharmacist interventions Implementation: 11/2001 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated e-MAR: Nursing Medication Administration Record | Acute care/tertiary, 633 Beds Inpatient hospital based, Academic | Number of pharmacist interventions | The number of pharmacist interventions remained the same before and after CPOE implementation in the control unit (80 before and 84 after) In the CPOE unit the number of pharmacist interventions increased from 76 to 109, p <0.01. | + |
| Varkey (2007) ¹⁷² Design: Cross- sectional N = 4,527 prescriptions Implementation: 00/0000 Study Start: 00/1996 Study End: 00/2002 | CPOE/POE system Integrated CDSS/CDS/CCDS/re minders | Ambulatory care, Other institution based | frequency of intercepted prescription errors* | Statistically significant decrease in frequency of intercepted prescription errors among handwritten and computerized prescriptions was observed (7.4% vs. 4.9%, p = 0.0048). | + |
| Wess (2007) ¹⁹³ Design: Before-after N = 3,791 medication orders Implementation: 06/2005 Study Start: 00/0000 Study End: 00/0000 | CPOE/POE system Integrated EHR/EMR system Hospital information system | General Hospital, Inpatient hospital based, Academic | mean time from provider order entry to pharmacist verification, -community hospital, - university hospital, proportion of clarification calls placed, - community hospital, - university hospital | The mean time from provider order entry to pharmacist verification decrease for both community (152 vs. 32 minutes, p <0.0001) and university hospitals (108 vs. 50 minutes, p <0.0001) The call back percentage also decreased for both community (2.8 vs. 0.4%, RRR 86%, p <0.0001) and university hospitals (2.8% vs. 0.5%, RRR 82%, p <0.0001). | + |

Evidence Table 2. KQ1: primary process outcomes for all technologies assisting order communication (continued)

| Article Information | HIT Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|-----------------------------------|---|---------|
| Wietholter (2009) ¹⁹⁴ Design: Before-after N = 2,988 medication orders Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CPOE/POE system Integrated Pharmacy | Acute care/tertiary, 761 Beds Inpatient hospital based | mean order-processing time (min)* | The mean order-processing time before CPOE implementation was 115 minutes from prescriber composition to pharmacist verification. After CPOE implementation, the mean order-processing time was reduced to 3 minutes (p <0.0001). | + |

Evidence Table 3. KQ1: primary process outcomes for all technologies assisting dispensing

| Article Information | HIT Studied | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|---|---|---------|
| Alvarez Diaz (2009) ¹⁹⁵ Design: Observational study N = 54,169 opportunities for error (the number of medication order lines validated) Implementation: 00/0000 Study Start: 02/2008 Study End: 10/2008 | Integrated systems 3 medication dispensing systems Integrated CPOE/POE system Pharmacy | General Hospital, 1070 Beds Pharmacy Inpatient hospital based | validation errors *, filling errors* | Error rate and OR for validation errors in unit dose distribution system (UDDS) without CPOE (No-CPOE-UDDS) in relation to the UDDS with CPOE (CPOE-UDDS) was 146/13,645 (1.1) vs. 63/20,240 (0.3), (OR 0.289, 95% CI 0.215 to 0.388). For No-CPOE-UDDS in relation to automated dispensing system with CPOE (CPOE-ADS), it was 146/13,645 (1.1) vs. 83/13,932 (0.6), (OR 0.554, 95%CI 0.423 to 0.726) Error rate and OR for filling errors No-CPOE-UDDS in relation to CPOE-UDDS was 265/13,645 (1.9) vs. 345/20,240 (1.7), (OR 0.876, 95% CI 0.745 to 1.029). For No-CPOE-UDDS in relation to CPOE-ADS it was 265/13,645 (1.9) vs. 309/13,932 (2.2), (OR 1.145, 95% CI 0.97 to 1.352). | + |
| Halkin (2001) ⁶⁵ Design: Time series N = 775,186 prescriptions Implementation: 11/1997 to 00/1998 Study Start: 01/1998 Study End: 06/1999 | CDSS/CDS/CCDS/reminders Integrated Pharmacy | Pharmacy, HMO pharmacy | drug interaction rate prescriptions 90% pharmacies and 50% of physicians compared with baseline drug interaction rate prescriptions 95% pharmacies and 90% physicians compared with baseline | Dispensing of drug interaction prescriptions was reduced by 21.1% and by 67.5% in periods II and III compared with period I (OR 0.79; 95% CI 0.75 to 0.83 and OR 0.28; 95% CI 0.26 to 0.30, respectively). | + |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: ADS = Automated Dispensing System; CDS = Clinical / Computerized Decision Support ; CDSS = Clinical Decision Support System; CIT = clinical information technology; CPOE = Computerized Provider Order Entry; EHR = Electronic Health Record; e-MAR = Electronic Medication Administration Record; EMR = Electronic Medical Records; e-TAR = Electronic Treatment Authorization Request; HIT = Health Information Technology; HMO = Health Maintenance Organization; OR = Odds ratio; N = Sample Size; NS = Not specified; p = Probability; RCT = Randomized Controlled Trial; RRR = Relative Risk Reduction; UDDS = Unit Dose Drug Dispensing System; vs. = Versus

Evidence Table 3. KQ1: Primary Process outcomes for all technologies assisting dispensing (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|---|--|---------|
| Mahoney (2007) ⁹⁹ Design: Before-after N = 2,843,135 inpatient medication orders Implementation: 02/2002 Study Start: 02/2002 Study End: 06/2006 | CDSS/CDS/CCDS/ reminders CPOE/POE system, Pharmacy information system Integrated EHR/EMR system Hospital information system | General Hospital, Pediatric stand alone hospital, 966 in 2 hospitals Beds Pharmacy Inpatient hospital based Academic | rate of -drug allergy violations*, -excessive doses*, -incomplete or unclear orders*, -therapeutic duplication* | Medication errors decreased after implementation of the CIT with respect to drug allergy violations (OR 0.14, CI 0.11 to 0.17, p <0.001), excessive doses (OR 0.68, CI 0.62 to 0.74, p <0.001) and incomplete or unclear orders (0.35, CI 0.32 to 0.38, p <0.001) but no decrease in therapeutic duplications. Turnaround time between drug ordering and administration decreased from 90 minutes to 11 minutes, no stats given. The override rate also decreased (7.1 to 2.9%, RRR 59%, p = 0.001). | + |
| Murray (1999) ¹⁸⁸ Design: Cohort N = 11,102 observations of 28 pharmacists Implementation: 03/1995 Study Start: 11/1995 Study End: 01/1996 | Pharmacy information system Integrated EHR/EMR system Imaging systems Laboratory system | Acute care/tertiary, Pharmacy, Inpatient hospital based | distribution of pharmacist time on activities, functions and contacts* | The electronic guidelines and reminders were associated with the overall distribution of activities (more time discussing information and less time checking and preparing prescriptions) p <0.001; overall functions (more time advising or discussing information or problem solving and less time filling prescriptions) p <0.001, and distribution of contacts (more time with other pharmacy personnel, patients, and clinicians and less time working alone) p <0.001. | + |
| Pearce (2010) ¹⁹¹ Design: Before-after N = 332 medication refill orders Implementation: 05/2006 Study Start: 02/2006 Study End: 03/2007 | e-Pprescribing Integrated EHR/EMR system Pharmacy | Ambulatory care, Pharmacy Pharmacy chain | time to a response for refill request* | The average time to a response to a pharmacy refill request decreased from 1.57 days to 1.04 days (p <0.004). | + |

Evidence Table 3. KQ1: Primary Process outcomes for all technologies assisting dispensing (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|---|----------|
| <p>Raebel (2007)¹³⁰ Design: RCT N = 59,680 patients Implementation: 00/0000 Study Start: 05/2005 Study End: 05/2006</p> | <p>CDSS/CDS/CCDS/ reminders Pharmacy information system Integrated EHR/EMR system</p> | <p>Ambulatory care, HMO pharmacy</p> | <p>new dispensings of targeted medications*, dispensings of targeted medications considered inappropriate*</p> | <p>In the analysis of all dispensings of targeted medications, there was a significant reduction of new dispensings of at least one targeted medication (2.2% vs. 1.8%, RRR 16%, p <0.002) For dispensings of targeted medications considered inappropriate, there was also a significant reduction with the use of the alerting system (1.5% vs. 1.1%, RRR 27%, p <0.001).</p> | <p>+</p> |
| <p>Reeve (2007)¹⁹⁶ Design: RCT N = 2,396 clinical interventions by pharmacists Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Pharmacy information system</p> | <p>Pharmacy, Other, Stand alone non chain store (eg family run)</p> | <p>Rate of pharmacist interventions /100 diabetic patients-aspirin therapy for diabetic patients * The total rate of pharmacist intervention/100 patients *</p> | <p>The rate of targeted interventions for aspirin therapy for high risk diabetic patients was 0 for the control group and 4.82/100 diabetic patients (p <0.05) .Clinicians who received the prompts had a higher rate of intervening with patients overall (1.74 per 100 patients) compared with pharmacists who did not receive the prompts (mean 0.91 per 100 patients), p <0.001. When the prompts were stopped the rate of aspirin interventions fell to pre-prompt levels.</p> | <p>+</p> |
| <p>Wilson (1997)¹⁹⁷ Design: Before-after N = 00 not stated number of medications, etc Implementation: 02/1994 Study Start: 07/1993 Study End: 06/1995</p> | <p>e-Medication administration system (e-MAR, e- TAR) Integrated Formulary, Hospital information system</p> | <p>Acute care/tertiary, 362 Beds Inpatient hospital based Academic</p> | <p>Medication occurrences per admission*, Medication occurrences per patient day*, Medication occurrences per order, Medication occurrences per dose</p> | <p>Self-reported medication occurrences (errors) per admission (11% vs. 7%, RRR 39%, p <0.001), per patient day (1.4% vs. 7%, RRR 34%, p <0.001), per order (0.4% vs. 0.3%, RRR 34%, p <0.001), and per dose (0.05% vs. 0.03%, RRR 40%, p <0.001) were all significantly reduced following implementation of a shared electronic medication record for pharmacists and nurses.</p> | <p>+</p> |

Evidence Table 4. KQ1: primary process outcomes for all technologies assisting drug administration

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|---|---|---------|
| Banet (2004) ¹⁹⁸ Design: Before-after N = 55 nurses Implementation: 05/2003 Study Start: 00/0000 Study End: 00/0000 | CPOE/POE system, e-MAR, e-Medication administration system (e-MAR, e-TAR) Integrated Imaging systems Laboratory system Pharmacy | Emergency department Academic | distribution of nurses' time on activities, functions and contacts* | Time-motion study demonstrated that after implementing the information system changes, nurses spent less time (mean percent of total time) on paper documentation (17% vs. 2%, RRR 90%, p <0.05) and searching for charts (0.4% vs. 0.1%, RRR 75%, p <0.05). They spent more time using computers (10% vs. 26%, RRR -157%, p <0.05), and charting in patients rooms (0.2% vs. 2.1%, RRR -950%, p <0.05). They spent the same amount of time on documentation tasks overall (27% vs. 28%, RRR 3%, NS) and direct patient care (41% vs. 39%, RRR 4%, NS). | + |
| Climent (2008) ¹⁹⁹ Design: Cross-sectional N = 2,242 opportunities for error Implementation: 00/0000 Study Start: 05/2005 Study End: 06/2006 | 3 different drug delivery systems, e-Medication administration system (e-MAR, e-TAR), e-Rx | Acute care/tertiary, 1,500 Beds Academic | medication error rate*, medication error rate-reaching patients* | The integrated MMIT unit dose delivery system with e-Rx (DUPEA) had an error rate similar to the non-integrated unit dose system (DUTI), and the ward stock system (9.5% stock vs. 7.8% DUPEA vs. 4.7% DUTI). The error rate reaching patients with the DUPEA was lower than stock but higher than DUTI (8.1% stock vs. 5.5% DUPEA vs. 0.4% DUTI, p <0.05). | - |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: AR = Absolute Reduction; BCMA = Bar Code Medication Administration ; CCDS = Computerized Clinical Decision Support; CCU = Critical Care Unit; CDS = Clinical / Computerized Decision Support ; CDSS = Clinical Decision Support System; CI = CI; CPOE = Computerized Provider Order Entry; EHR = Electronic Health Record; e-MAR = Electronic Medication Administration Record; EMR = Electronic Medical Records; e-RX = Electronic Prescribing; e-TAR = Electronic Treatment Authorization Request; hr = Hour; ICU = Intensive Care Unit; MMIT = Medication Management Information Technology; N = Sample Size; NICU = Neonatal Intensive Care Unit; NS = Not specified; OR = OR; OSUH= Ohio State University Health System; p = Probability; POE = Provider Order Entry; RRR = Relative Risk Reduction; SD = Standard deviation; UDDS = Unit Dose Drug Dispensing System; vs. = Versus

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|--|---------|
| DeYoung (2009) ²⁰⁰ Design: Before-after N = 1,465 medication administrations in 92 patients Implementation: 01/2007 Study Start: 12/2006 Study End: 05/2007 | BCMA (BCMA) Integrated e-MAR | Critical care units (CCU, ICU, NICU) 38 in ICU, 744 in hospital Beds Academic | error rate-overall*, - excluding documentation errors*, - wrong administration time* | The medication error rate was reduced by 56% after the implementation of BCMA (19.7% vs. 8.7%, p <0.001). This rate increased to 63% when documentation orders were excluded (p <0.001). The benefit was related to a reduction associated with errors of wrong administration time. Wrong administration time errors decreased from 18.8% during preimplementation to 7.5% postimplementation (p <0.001). There were no significant differences in other error types. | + |
| Fontan (2003) ⁴⁶ Design: Cross-sectional N = 4,532 prescriptions Implementation: 00/1988 Study Start: 02/1999 Study End: 03/1999 | Computerized UDDS Integrated Hospital information system | Other specialty hospital (rehab, oncology) Pediatric stand alone hospital, 510 Beds | Prescribing error rate, Administering error rate | Errors were decreased with the use of the e-RX and computerized dispensing system compared with the hand-written prescriptions and ward distribution system. Prescribing errors were reduced from 87.9% to 10.6%, RRR 88%, p <0.00001 Administrative errors with time errors were reduced from 29.3% to 22.5%, RRR 23%, p <0.001. | + |

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|--|---------|
| Franklin (2007) ⁵⁰ Donyai (2008) ⁵¹ Barber (2007) ⁵² Franklin (2008) ⁵³ Franklin (2007) ⁵⁴ Design: Before-after N = 4,803 medication orders Implementation: 06/2003 Study Start: 00/0000 Study End: 00/0000 | Automated Dispensing Machine, e-Medication administration system (e-MAR, e-TAR) e-Rx Integrated Pharmacy | Acute care/tertiary, 28 surgery bed ward of a teaching hospital Beds Inpatient hospital based, Academic | error rate for new prescriptions*, error rate for drug administrations*, %administered <1hr ⁵³ , rate of pharmacist interventions ⁵¹ Total pharmacy time taken on study ward | The prescription error rate for new orders dropped significantly after implementation of the system (3.8% vs. 2.0%, RRR 47%, p = 0.0004) Medication administration error rate also significantly declined (8.6% vs. 4.4%, RRR 49%, p = 0.0003). ⁵³ Postintervention medication timeliness was improved (%administered <1hr, 79% vs. 89%, p <0.001). ⁵¹ The rate of pharmacist interventions declined significantly after implementation (3.0% vs. 1.9%, AR 1.1 (95% CI 0.2,2.0)). ⁵⁴ Total pharmacy time taken on study ward increased after implementation (1h 8min vs. 1h 38min, p = 0.001). Pharmacists were required to endorse fewer orders (50% vs. 21%, RRR 58%, p <0.0001) and endorsed fewer orders (55% vs. 30%, RRR 45%, p <0.0001). | + |

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|---|---|----------|
| <p>Helmons (2009)²⁰¹ Design: Before-after N = 2,353 opportunities for error Implementation: 05/2007 Study Start: 09/2007 Study End: 04/2008</p> | <p>BCMA Integrated Handheld CPOE/POE system EHR/EMR system, e-MAR, Pharmacy</p> | <p>Critical care units (CCU, ICU, NICU) 386 Beds Academic</p> | <p>error rate-surgical medical unit*, error rate-ICU*</p> | <p>The total medication administration error rates did not significantly decrease on the medical–surgical units (11% vs. 8%, RRR 23%, NS) the ICU (13% vs. 14% RRR - 7%, NS) or overall (13% vs. 14% RRR - 7%, NS) Accuracy measured by 6 indicators of accuracy reflecting error-prone process variations. Baseline medication administration accuracy higher in medical–surgical units compared with the ICUs. On the medical– surgical units, 3 accuracy indicators changed after the introduction of BCMA; improved compliance with checking patient identity after BCMA implementation was offset by more distractions and interruptions and less explanation of the medication to the patient. These 3 indicators did not change in the ICUs However, implementation of BCMA resulted in improved charting and labelling of medications administered in the ICUs.</p> | <p>-</p> |
| <p>Low (2002)²⁰² Design: Before-after N = not reported prescriptions Implementation: 03/2000 Study Start: 03/1999 Study End: 03/2001</p> | <p>BCMA Integrated Hospital information system</p> | <p>Acute care/tertiary, Pharmacy Inpatient hospital based</p> | <p>rate of errors per 1,000 doses</p> | <p>The rate of errors per 1,000 doses did not differ across the 24 month periods before and after BCMA (0.125 vs. 0.145, p = 0.6).</p> | <p>-</p> |

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|---|---------|
| Mahoney (2007) ⁹⁹ Design: Before-after N = 2,843,135 inpatient medication orders Implementation: 02/2002 Study Start: 02/2002 Study End: 06/2006 | CDSS/CDS/CCDS/reminders, CPOE/POE system, Pharmacy information system Integrated EHR/EMR system, Hospital information system | General Hospital, Pediatric stand alone hospital, 966 in 2 hospitals Beds Pharmacy Inpatient hospital based, Academic | rate of -drug allergy violations*, -excessive doses*, -incomplete or unclear orders*, -therapeutic duplication* | Medication errors decreased after implementation of the CIT with respect to drug allergy violations (OR 0.14, CI 0.11 to 0.17, p <0.001) excessive doses (OR 0.68, CI 0.62 to 0.74, p <0.001) and incomplete or unclear orders (0.35, CI 0.32 to 0.38, p <0.001) but no decrease in therapeutic duplications. Turnaround time between drug ordering and administration decreased from 90 minutes to 11 minutes, no stats given. The override rate also decreased (7.1 to 2.9%, RRR 59%, p = 0.001). | + |
| Mekhjian (2002) ¹⁸⁶ Design: Before-after N = 28,898 patients Implementation: 05/2000 Study Start: 02/2000 Study End: 01/2001 | CPOE/POE system, e-Medication administration system (e-MAR, e-TAR) Integrated Dietary system, EHR/EMR system, Imaging systems, Laboratory system | Acute care/tertiary, Other specialty hospital (rehab, oncology) Academic | medication turn-around time, proportion of verbal orders countersigned, rate of transcription errors | Combining the data showed that time from initiation of the prescription and administration was reduced after POE: mean 5:28 hours before vs. 1:51 hours after, 64% relative reduction, p <0.001. The proportion of signed verbal orders increased for both hospitals: OSUH 56.4% vs. 76%, RRR 76%, p <0.001 and James Cancer 72.8% vs. 99.0, RRR 36%, p <0.001. The volume of transcription errors was reduced after POE from 11.3% to 0%, RRR 100%, p <0.001. | + |
| Mitchell (2004) ¹⁸⁷ Design: Cross-sectional N = 4,297 prescriptions Implementation: 0/1999 Study Start: 09/2002 Study End: 12/2002 | e-Medication administration system (e-MAR, e-TAR), e-Rx Integrated Formulary, Pharmacy | Acute care/tertiary, Academic | 15 aspects of data completeness for e-MAR were sought with implementation of the e-MAR. | e-MAR was more accurate (more inclusion of important information) for nurses 9 of the 15 were statistically significantly improved including presence of dosing recommendations (30% v3 99%, RRR 230%, p <0.01), Errors detected by the pharmacist did not differ before and after implementation of the e-Rx system. Only minor errors were reduced with the system | + |

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|--|----------|
| <p>Morriss (2009)²⁰³ Design: Cohort study N = 958 patients; 92,398 doses administered Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated Pharmacy</p> | <p>Critical care units (CCU, ICU, NICU) 36 beds NICU Beds Academic</p> | <p>Medication Error*, Potential ADEs*, preventable ADEs*</p> | <p>When the BCMA system was not operative, the unadjusted medication error rates were 69.5/1,000 doses and mean 0.53 (SD 0.98)/subject/day. The unadjusted medication error rates increased in the study NICU when the BCMA system was operative to 79.7/1,000 doses and mean 0.60 (SD 0.99)/subject/day (p <0.001). The increase in medication error was associated with a 117% increase in detected wrong-time errors from 1412 before the BCMA system to 3075 when the system was operative. Significant decrease in potential ADEs [0.11 (0.47) vs. 0.033 (0.20), p <0.001], or unadjusted targeted, preventable ADEs [0.0065 (0.082) vs. 0.0032 (0.060) p <0.008] for subjects cared for in the BCMA system-equipped beds.</p> | <p>+</p> |
| <p>Paoletti (2007)²⁰⁴ Design: Before-after N = 1,868 Doses observed Implementation: 08/2003 Study Start: 00/0000 Study End: 00/0000</p> | <p>BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated Hospital information system, Pharmacy</p> | <p>General Hospital, 521 Beds</p> | <p>error rate*</p> | <p>The error rate compared between pre and postimplementation period in the three groups were: 19.6% vs. 20.6%, p = 0.762 (control); 25.3% vs. 19.2%, p = 0.065 (Intervention Group 1) and 15.6% vs. 10%, p = 0.035 (Intervention Group 2). Group 1 and 2 were noted to have different practices during baseline measurement. [unsure if this would be considered a positive trial].</p> | <p>+</p> |

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|---|--|----------|
| <p>Persell (2008)²⁰⁵ Design: RCT N = 242 patients Implementation: 00/0000 Study Start: 10/2004 Study End: 03/2005</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system</p> | <p>Ambulatory care, Academic</p> | <p>self-reported aspirin use* by all patients, self-reported aspirin use* by patients excluding long-term aspirin users and patients reporting medical contraindication</p> | <p>The control rate (reminders only) of self-reported aspirin use was not significantly different than the intervention (reminders plus clinician emails and patient phone calls) group (39% vs. 46%, p = 0.20). Excluding long-term aspirin users and patients reporting medical contraindication (30% vs. 43%, p = 0.013).</p> | <p>-</p> |
| <p>Poon (2006)²⁰⁶ Design: Before-after N = 232 observation sessions Implementation: 00/0000 Study Start: 02/2005 Study End: 10/2005</p> | <p>BCMA</p> | <p>Acute care/tertiary, 735 Beds</p> | <p>proportion time on medication administration, proportion time nurses spent on direct care</p> | <p>The proportion of time nurses spent on the major activity groups remained stable. Before BCMA implementation, nurses spent 26.5% of their time on medication administration. After BCMA implementation, this proportion remained statistically unchanged at 24.5% (RRR 8%, p = 0.22). The proportion of time nurses spent on direct care activities unrelated to medication administration remained statistically unchanged (20.1% vs. 23.7%, RRR -18%, p = 0.15).</p> | <p>-</p> |
| <p>Poon (2010)²⁰⁷ Design: Before-after N = 14041 medication administration Implementation: 04/2005 Study Start: 02/2005 Study End: 10/2005</p> | <p>BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated EHR/EMR system, Pharmacy</p> | <p>Acute care/tertiary, 735 Beds Academic</p> | <p>Non-timing errors in medication administration*, Timing errors in medication administration*, transcription error (2ndary outcome)</p> | <p>On units without the bar-code e-MAR, 776 (11.5%) non-timing medication-administration errors was observed compared to 495 (6.8%) on units with the bar-code e-MAR (p <0.001). The overall incidence of medication doses directly observed to be administered either early or late decreased from 16.7% without the bar-code e-MAR to 12.2% with its use (p = 0.001). The units without bar-code e-MAR observed 110 (6.1%) transcription errors while those with it observed no errors (p <0.001).</p> | <p>+</p> |

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|--------------------------------------|---|--|---------|
| Shirley (1999) ²⁰⁸ Design: Before-after N = 163 medication administrations Implementation: 00/0000 Study Start: 05/1997 Study End: 08/1997 | Automated drug dispensing system Integrated, Pharmacy | Acute care/tertiary, 270 Beds | proportion of medications administered as scheduled*, mean time deviation between actual and scheduled administration times*, | Before implementation of the automated dispensing system, 59% of 76 medication doses were administered as scheduled, after 77% of 87 doses were administered as scheduled (RRR -31%, p = 0.02). The mean time deviation between actual and scheduled administration times did not change significantly postimplementation (130 minutes vs. 101 minutes, p = 0.157). | - |
| Taylor (2008) ²⁰⁹ Design: Before-after N = 521 medication administrations Implementation: 07/2005 Study Start: 09/2004 Study End: 04/2006 | CPOE/POE system Integrated e-MAR, Pharmacy | Critical care units (CCU, ICU, NICU) | variance in medication administration | Medication variances were detected for 19.8% of administrations during the pre-CPOE period, compared with 11.6% with CPOE (RRR 41%, p <0.05). The reasons for medication administration variances during the pre-CPOE and CPOE were not statistically different. Overall, administration mistakes, pharmacy problems and prescribing problems accounted for 74% of all variances observed. | + |
| Wax (2007) ²¹⁰ Design: Before-after N = 14,465 patients Implementation: 02/2005 Study Start: 06/2004 Study End: 12/2005 | Anesthesia information management system (AIMS), CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Acute care/tertiary, Academic | overall compliance with antibiotic administration before surgery, noncompliance due to late administration, noncompliance due to early administration | Compliance (antibiotics 60 min before surgery) for the entire data set increased from 82.4% to 89.1% (RRR -8%, p <0.01) following the event icon implementation. Noncompliance rates decreased following implementation for late administration (15.2% vs. 8.1%, RRR 47%, p <0.01), but remained unchanged for early administration (2.4% vs. 2.8%, RRR -17%, p = 0.07). | + |

Evidence Table 4. KQ1: Primary Process outcomes for all technologies assisting drug administration (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|---|---|---------|
| Wilson (1997) ¹⁹⁷ Design: Before-after N = 00 not stated number of medications, etc Implementation: 02/1994 Study Start: 07/1993 Study End: 06/1995 | e-Medication administration system (e-MAR, e-TAR) Integrated Formulary, Hospital information system | Acute care/tertiary, 362 Beds Inpatient hospital based, Academic | Medication occurrences per admission*, Medication occurrences per patient day*, Medication occurrences per order, Medication occurrences per dose | Self-reported medication occurrences (errors) per admission (11% vs. 7%, RRR 39%, p <0.001), per patient day (1.4% vs. 7%, RRR 34%, p <0.001), per order (0.4% vs. 0.3%, RRR 34%, p <0.001), and per dose (0.05% vs. 0.03%, RRR 40%, p <0.001) were all significantly reduced following implementation of a shared electronic medication record for pharmacists and nurses. | + |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|---|---|---------|
| Abboud (2006) ¹ Design: Before-after N = 336 orders Implementation: 04/2002 Study Start: 10/2003 Study End: 03/2004 | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated CDSS/CDS/CCDS/ reminders EHR/EMR system, Formulary, Hospital information system, Imaging systems, Laboratory system, Pharmacy | Pediatric stand alone hospital, 423 Beds | Antibiotics courses with no lab order* | There were no significant differences between the baseline and the corollary order periods on courses of antibiotics associated with no laboratory monitoring [31(19.5%) vs. 31(17.5%), p = NS]. | - |
| Bertoni (2009) ¹² Design: RCT N = 8,878 patients Implementation: 00/0000 Study Start: 06/2001 Study End: 04/2006 | CDSS/CDS/CCDS/ reminders Integrated Handheld | Ambulatory care | adherence to guideline-screening*, adherence to guideline-appropriate lipid management* | There was no difference in screening rates between the CDSS-PDA group and the control. The control group had a 10.8% drop in appropriate management from baseline, while the PDA group had a 1.1% drop, p <0.01. Stable adherence was observed in the PDA intervention group, whereas a decline in guideline adherence was observed in the control group. | - |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: A1c = haemoglobin A1c; ACE = Angiotensin Converting Enzyme; ALT = Alanine Aminotransferase; AST = Aspartate Aminotransferase; CCDS = Computerized Clinical Decision Support; CCU = Critical Care Unit; CDS = Clinical / Computerized Decision Support ; CDSS = Clinical Decision Support System; CI = CI; CPOE = Computerized Provider Order Entry; DM = Diabetes Mellitus ; EHR = Electronic Health Record; EMR = Electronic Medical Records; GP = General Practitioner; HIT = Health Information Technology; HIV = Human Immunodeficiency Virus; HMO = Health Maintenance Organization; ICU = Intensive Care Unit; K = Potassium; Mg = Magnesium; N = Sample Size; NICU = Neonatal Intensive Care Unit; NPs = Nurse Practitioners; NS = Not statistically significant; NSAID = Nonsteroidal anti-inflammatory drug; OR = Odds Ratio; p = Probability; PCA = Patient-Controlled Analgesia PDA = Personal Digital Assistants ; PHR = Patient Health Record; POE = Provider Order Entry; RCT = Randomized Controlled Trial; RRR = Relative Risk Reduction; vs. = Versus; yr = Year

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|---|----------|
| <p>Bertsche (2009)²¹¹ Design: Before-after N = 100 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/ reminders Integrated Formulary</p> | <p>Acute care/tertiary, 1621 Beds Academic</p> | <p>number of patients with at least one deviation from international guidelines*</p> | <p>At discharge, the number of patients with at least one deviation from international guidelines decreased by the intervention from 37 (74%) in control group to 7 (14%) in the intervention group (p <0.001).</p> | <p>+</p> |
| <p>Chambers (2008)²¹² Design: Before-after N = 87 patients Implementation: 00/0000 Study Start: 01/2005 Study End: 06/2006</p> | <p>CDSS/CDS/CCDS/ reminders Computer-Based Antimicrobial Monitoring System Integrated EHR/EMR system, Laboratory system, Pharmacy</p> | <p>Unspecified Hospital</p> | <p>Vancomycin de-escalation rates*, Mean duration of Vancomycin therapy (days)*, Combination-Antimicrobial de-escalation rate*, Mean duration of Combination-Antimicrobial therapy (days)*</p> | <p>Vancomycin de-escalation rates significantly improved from 33% to 68% with intervention (p = 0.001). In addition, the average duration of therapy was decreased from 10.4 ± 7.3 days to 7.7 ± 2.4 days (p = 0.014). Combination-Antimicrobial de-escalation rates were not statistically improved upon (67% vs. 63%, p = 0.763). The average duration of therapy was decreased from 12.8 ± 5.5 days to 9.5 ± 2.5 days, p = 0.335).</p> | <p>+</p> |
| <p>Chisholm (2003)¹⁹ Design: Before-after N = 790 children admitted to hospital with asthma exacerbations Implementation: 10/2002 Study Start: 11/2001 Study End: 12/2003</p> | <p>CPOE/POE system Integrated Billing/administration system, EHR/EMR system, Laboratory system</p> | <p>Pediatric stand alone hospital, 323 Beds</p> | <p>systemic corticosteroids use*, metered-dose inhaler use*</p> | <p>More use was made of systemic corticosteroids (OR 5.61, 95% CI 3.46 to 9.11) and of metered-dose inhalers (OR 1.42, 95% CI 1.04 to 1.94) after implementation of standard order sets in the CPOE for asthma patients.</p> | <p>-</p> |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|-----------------------------|---|---|---------|
| Cobos (2005) ²³ Design: RCT N = 2,221 patients Implementation: 04/2000 Study Start: 04/2000 Study End: 05/2002 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | proportion of patients prescribed lipid lowering drugs (secondary) | The proportion of patients prescribed lipid lowering drugs was significantly lower in the CDSS guideline intervention group (59.1% vs. 40.8% RRR 31%, p <0.0001). | + |
| Demakis (2000) ²¹³ Design: RCT N = 12,989 patients Implementation: 00/0000 Study Start: 01/1995 Study End: 06/1996 | CDSS/CDS/CCDS/ reminders Integrated Hospital information system | Ambulatory care Academic | adherence rates for 5 medication management standards of care*, monitoring warfarin treatment; treatment of atrial fibrillation with warfarin, aspirin or ticlopidine; treatment of myocardial infarction with beta-blockers or switching NSAID therapy for gastrointestinal bleeds, pneumococcal vaccination | Adherence rates for medication management standards of care were not significantly different for 4 of the 5 medication management standards of care. There was a large effect for pneumococcal vaccination (12.7% vs. 4.3%; OR 3.26; 95% CI, 2.09 to 5.09). adherence was significantly improved for 13 standards (53.5% vs. 58.8%, OR 12.4 (95% CI 1.08 to 1.42, p = 0.002). | - |
| Evans (1990) ³⁶ Design: Before-after N = 7,656 patients Implementation: 00/0000 Study Start: 06/1985 Study End: 09/1986 | CDSS/CDS/CCDS/ reminders Hospital information system Integrated Laboratory system, Pharmacy | Unspecified Hospital | mean number of antibiotic doses per patient, proportion of patients receiving perioperative antibiotics, proportion of patients receiving antibiotics for too long | Surgical patients received an average of 19 antibiotic doses before implementation of the 'stop orders' and 13 after (p <0.001). There were non significant changes in the proportion of patients receiving perioperative antibiotics (64% vs. 66%, NS) or those receiving antibiotics for too long (40% vs. 35%, NS). | + |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|--|---------|
| Evans (1999) ²¹⁴ Design: Before-after N = 13,384 Patients Implementation: 01/1005 Study Start: 04/2005 Study End: 03/2006 | CDSS/CDS/CCDS/ reminders Integrated Pharmacy | Critical care units (CCU, ICU, NICU) 12 beds in the shock/trauma/respiratory intensive care unit. of a 520 bed academic hospital Beds Not specified, Inpatient hospital based, Academic | The mean number of days with excessive antibiotic dosing* | The mean number of days with excessive antibiotic dosing was lower after the intervention (4.7 days vs. 2.9 days, p <0.001). | + |
| Feldstein (2006) ²¹⁵ Smith (2009) ²¹⁶ Design: RCT N = 961 patients Implementation: 09/2004 Study Start: 09/2003 Study End: 01/2005 | CDSS/CDS/CCDS/ reminders Integrated Billing/administration system, EHR/EMR system, Pharmacy | Ambulatory care HMO pharmacy | rates of completing lab monitoring* | Patients in the EMR group were 2.5 times more likely than patients in the Usual Care group to complete laboratory monitoring (p <0.001), patients in the automated telephone voice message group were 4.1 times more likely (p <0.001), and patients in the pharmacy team outreach group were 6.7 times more likely (p <0.001). | + |
| Field (2009) ⁴⁰ Design: RCT N = 833 patients (10 physicians and 213,967 patient days) Implementation: 00/000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated EHR/EMR system | Long term care (nursing homes) | proportion of appropriate orders*, proportion of inappropriate drugs avoided | The proportion of appropriate antidepressant order rates for patients with renal insufficiency was higher in the CDSS group (52% vs. 63%, OR 1.2, 95% CI 1.0 to 1.4). More inappropriate drugs were avoided (15% vs. 46%, OR 2.6, 95% CI 1.4 to 5.0). Improvements were seen in frequency and missing information but not for doses in the CDSS group. | + |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|---|---|----------|
| <p>Galanter (2004)⁵⁷ Design: Before-after N = 620 patients Implementation: 00/0000 Study Start: 02/2001 Study End: 03/2002</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CDSS/CDS/CCDS/ reminders CPOE/POE system, Laboratory system</p> | <p>Acute care/tertiary, Academic</p> | <p>compliance with digoxin monitoring guidelines - synchronous alerts*, compliance with hypokalemia and hypomagnesemia treatment guidelines - synchronous alerts*, compliance with hypokalemia and hypomagnesemia treatment guidelines - asynchronous alerts*</p> | <p>Postimplementation, synchronous alerts significantly increased test ordering for digoxin levels, K levels and Mg levels at 1 hr and 24 hrs (p <0.01 for all). Supplementation of Mg at 1 hour was significantly improved, but not at 24 hrs. Supplementation of K was not improved at 1 or 24 hrs. Synchronous alerts resulted in improved compliance at 1 hr and 24 hrs for bot K and Mg supplementation (p <0.01).</p> | <p>+</p> |
| <p>Gill (2009)⁶⁰ Design: RCT N = 64,150 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 10/2006</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Up-to-date lipid test*, Lipid medication if not at goal (high risk patients only)*</p> | <p>Outcomes improved for most measures from before to 1 year after the intervention (univariate analysis). However, after controlling for confounding variables and for clustering in multilevel modeling, only up-to-date lipid testing for high-risk patients was statistically better in the intervention group as compared to the control group (adjusted OR 15.0, p <0.05). Intervention status was NS for any other analysis.</p> | <p>-</p> |
| <p>Gilutz (2009)⁶¹ Design: RCT N = 7,448 patients from 56 control and 56 intervention clinics Implementation: 00/0000 Study Start: 01/2000 Study End: 12/2003</p> | <p>CDSS/CDS/CCDS/ reminders Integrated Hospital information system, Laboratory system, Pharmacy</p> | <p>Ambulatory care Academic</p> | <p>rate of adequate monitoring, Positive treatment trend, overall up-titration rate in patients with LDL = 110 mg/dl</p> | <p>A higher rate of adequate monitoring was documented in the intervention arm (54.8% vs. 48.7%, p <0.001). Medication initiation or up-titration was recommended for patients with LDL levels above 110 mg/dl. The results showed that overall positive trends were minimally more prominent in the intervention arm (59.1% vs. 53.7%, p <0.003). Difference constitutes a higher rate of drug initiation (2.5%), up-titration (1.8%) and avoiding drug cessation (1.1%). Overall up-titration in patients with LDL = 110 mg/dl was poor, both in the intervention arm and in the control arm (8.6% vs. 7.4%, NS).</p> | <p>+</p> |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|---|---------|
| Grant (2008) ²¹⁷ Design: RCT N = 244 patients Implementation: 00/2002 Study Start: 09/2005 Study End: 03/2007 | PHR Integrated Billing/administration system, EHR/EMR system, Imaging systems, Laboratory system, Patient decision support system | General Hospital, Ambulatory care Home | Proportion of followup visits with DM related medication changes. | For the secondary outcome, significantly more followup visits included DM related medication changes in intervention patients than the control group, 15% vs. 53%, RRR 253%, p <0.001. | + |
| Hicks (2007) ⁶⁶ Design: RCT N = 1,422 patients Implementation: 00/0000 Study Start: 07/2003 Study End: 02/2005 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Other, Academic | blood pressure controlled, receiving a recommended drug class medication within 1 week of the clinic visit adjusted This study had 4 groups: usual care, CDS, NPs, and NPs+CDS. | No difference was seen across all 4 groups for blood pressure readings: Usual care vs. CDS: 45% controlled vs. 48% controlled, OR 0.96 (95% CI 0.78 to 1.19). Patients in the CDS group were more likely to have received a recommended drug class medication within 1 week of the clinic visit than patients in the usual care group: adjusted OR 1.32 (95% CI 1.09 to 1.61). | + |
| Javitt (2005) ²¹⁸ Design: RCT N = 39,462 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders Integrated Insurance | Ambulatory care | Compliance with recommendations to add-a-drug* | Physicians complied with 24% of these “add-a-drug” recommendations in the intervention group. In the control group, physicians spontaneously instituted the treatment that would have been recommended in 17% of instances in which the recommendation was triggered but not issued. This 42% relative difference in compliance was statistically significant (p = 0.007). | + |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|--|--|----------|
| <p>Koide (1999)⁸⁴ Design: Before-after N = 1,024 prescriptions for 111 patients and 68 physicians Implementation: 09/1994 Study Start: 09/1994 Study End: 09/1996</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, Hospital information system, Laboratory system</p> | <p>Acute care/tertiary, 1040 Beds Academic</p> | <p>rate of appropriate prescribing</p> | <p>Of 491 preintervention prescriptions, 127 (25.9%) were classified as appropriate because they were accompanied by a normal value of ALT or AST within 3 months. Of 533 postintervention prescriptions, 353 (66.2%) were classified as appropriate. Sudden increase occurred immediately after the start of the intervention (p <0.0001).</p> | <p>+</p> |
| <p>Kucher (2005)⁸⁹ Design: RCT N = 2,506 patients Implementation: 00/0000 Study Start: 09/2000 Study End: 01/2004</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, Hospital information system</p> | <p>Acute care/tertiary, Academic</p> | <p>received pharmacological interventions</p> | <p>More patients in the CDSS group received pharmacological interventions. (13% vs. 24%, RRR 69%, p <0.001).</p> | <p>+</p> |
| <p>Kuilboer (2006)²¹⁹ Design: RCT N = 32 primary care practices (78,926 patients of whom 9,798 had asthma or related symptoms) Implementation: 07/1998 Study Start: 07/1998 Study End: 05/1999</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>rate of prescribing for cromoglycate-12 to 39yr, rate of prescribing for cromoglycate-40-59yr</p> | <p>Of 20 potential changes in measurement rates, 8 were observed: The AsthmaCritic group had more contacts for the 12 to 39 year group (p = 0.03), more measurement of peak flow total for the 0 to 11 year group (p = 0.02), more FEV1 total peak flow ratio measurement in the 12-59 year groups (p = 0.04 and 0.009), and more measurement of FEV1 rates in the 3, 12 and older groups (p = 0.01, 0.01, and 0.016) Prescribing for cromoglycate was reduced in the 12 to 39 year and 40 to 59 year groups (12 to 39: 9.9/1000 patients vs. 4.1, p = 0.03) and (40 to 59: 9.0/1000 patients vs. 4.2, p = 0.05). Other prescribing (3 drugs or drug classes and 4 age groups) did not differ across groups.</p> | <p>+</p> |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|-----------------------------|---|--|---------|
| Lester (2005) ⁹⁴ Design: RCT N = 235 patients and 14 clinicians Implementation: 07/2003 Study Start: 07/2003 Study End: 07/2004 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care Academic | proportion of patients receiving statins*, proportion of patients receiving statins at 1 yr* | At 1 month more patients in the email group had received statins than control patients (3%, 15%, RRR 400, p <0.001). At 1 year the difference in receipt of statins had disappeared (17% vs. 25%, NS). | + |
| Lo (2009) ²²⁰ Design: RCT N = 3,673 potential alert trigger events (prescriptions) Implementation: 00/2000 Study Start: 07/2003 Study End: 01/2004 | CDSS/CDS/CCDS/ reminders Integrated CDSS/CDS/CCDS/ reminders Imaging systems | Ambulatory care Academic | proportion of events resulting in lab testing | 3,673 total events where baseline lab tests would have been advised: 1,988 events in the control group and 1,685 in the intervention group. In the control group, baseline labs were requested for 771 (39%) of the medications. In the intervention group, baseline labs were ordered by clinicians in 689 (41%) of the cases. Overall, no significant association existed between the intervention and the rate of ordering appropriate baseline laboratory tests (RRR 5%, p = 0.782). | - |
| Matheny (2008) ²²¹ Design: RCT N = 2,507 outpatient visits in 1,922 geriatric patients and 303 primary care physicians Implementation: 00/0000 Study Start: 01/2004 Study End: 06/2004 | CDSS/CDS/CCDS/ reminders Integrated CDSS/CDS/CCDS/ reminders Laboratory system | Ambulatory care Academic | rate of receiving appropriate laboratory testing within 14 days of the clinical encounter/ 10 medication-lab reminder categories | Reminders for appropriate laboratory monitoring had no impact on rates of receiving appropriate testing for creatinine, potassium, liver function, renal function, or therapeutic drug level monitoring for patients overdue for lab monitoring NSAIDs; Angiotensin Receptor Blockers ; Metformin; Potassium Supplements; Potassium Sparing Diuretics, Thiazide Diuretics; ACE Inhibitors; HMG Co-A Reductase Inhibitors; Thyroxine; (or the following therapeutic drugs combined: Carbamazepine; Cyclosporine, Phenobarbital, Phenytoin, Proc-NAPA, Valproate). | - |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|--|----------|
| <p>McDonald (1976)²²² Design: RCT N = 601 patient visits Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system</p> | <p>General Hospital, Academic</p> | <p>compliance with drug monitoring test alerts*, compliance with recommendations to change therapeutic regimens*</p> | <p>Alerts to patients overdue for drug monitoring tests resulted in an increased number of tests ordered (11% vs. 36%, RRR -227%, p <0.0001). Recommendations for changes to therapeutic regimens were followed in 28% of study events compared to 13% of control events (p <0.026).</p> | <p>+</p> |
| <p>McGregor (2006)¹⁰⁴ Design: RCT N = 4,507 patients Implementation: 00/000 Study Start: 05/2004 Study End: 08/2004</p> | <p>CDSS/CDS/CCDS/ reminders Integrated Laboratory system, Pharmacy</p> | <p>Acute care/tertiary, 648 Beds Inpatient hospital based, Academic</p> | <p>mean time spent on antimicrobial management</p> | <p>Team members spent 3.2 hours per day on management of antimicrobials with the decision support system compared with 4 hours per day without. No statistical testing was done.</p> | <p>+</p> |
| <p>McMullin (1999)¹⁰⁵ Design: Before-after N = 265 patients Implementation: 01/1996 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Laboratory system, Pharmacy</p> | <p>Acute care/tertiary, Pharmacy, Inpatient hospital based</p> | <p>rate of concomitant orders for contraindicated medications with cisapride*</p> | <p>The rate of ordering contraindicated drugs with cisapride was reduced with COPE (9% vs. 3.1%, RRR 65%, p <0.001).</p> | <p>+</p> |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|---|---|----------|
| <p>Montgomery (2000)¹⁰⁷ Design: RCT N = 552 patients Implementation: 00/0000 Study Start: 09/1996 Study End: 09/1998</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>probability of patients taking 2 drugs, probability of patients taking 3 drugs</p> | <p>Adjusted data showed that compare with those in the risk chart group alone, those with computer support had a lower probability of patients taking 2 drugs (OR 0.5, 95% CI 0.2 to 0.9) < p <0.05) or 3 drugs (OR 0.3, CI 0.1 to 0.6, p <0.05).</p> | <p>-</p> |
| <p>Niiranen (2008)¹¹³ Design: Time series N = 18,818 patient followups Implementation: 03/2005 Study Start: 04/2005 Study End: 12/2007</p> | <p>CDSS/CDS/CCDS/ reminders Integrated Laboratory system</p> | <p>Ambulatory care Home</p> | <p>proportion of patient followups assigned by nurses, year 1 to 2, proportion of patient followups assigned by nurses, year 2 to 3</p> | <p>In general, the share of patient followups assigned by nurses was similar in year 1 and 2 (56.7% vs. 55.1%, RRR 3%, NS), and increased significantly between year 2 and 3 (55.1% vs. 58.7%, RRR -7%, p <0.001).</p> | <p>+</p> |
| <p>Okon (2009)²²³ Design: Before-after N = 51,619 severe pain events Implementation: 09/2005 Study Start: 10/2005 Study End: 06/2007</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, nurse charting system</p> | <p>Acute care/tertiary, 504 Beds</p> | <p>rate of reassessment errors, time to resolution of pain events-minutes (4 time periods)</p> | <p>Aggregate delayed reassessment error postintervention rate of 35.8% compared with preintervention (56.2%, p <0.0001) for relative error reduction of 36%. Observed median time to resolution of severe pain events among all hospitalized patients decreased from 195 (T0) to 117 minutes (T1), 106 minutes (T2), and 101 minutes (T3) (all p <0.0001).</p> | <p>+</p> |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|--|----------|
| <p>Overhage (1997)¹¹⁷ Design: RCT N = 86 physicians on 6 services (services randomized) Implementation: 00/0000 Study Start: 10/1992 Study End: 04/1994</p> | <p>CDSS/CDS/CCDS/ reminders Integrated, CPOE/POE system, EHR/EMR system, Laboratory system</p> | <p>General Hospital, Academic</p> | <p>immediate compliance with corollary ordering*, 24 hour compliance*, hospital-stay compliance*</p> | <p>Intervention physicians placed corollary orders twice as often as control physicians did when measured by immediate compliance (46.3% vs. 21.9%, RRR -111%, p <0.0001). Significant differences between study and control physicians also appear in 24 hour compliance (50.4% vs. 29.0%, RRR -74%, p <0.0001) and hospital stay compliance (55.9% vs. 37.1%, RRR 51%, p <0.0001).</p> | <p>+</p> |
| <p>Palen (2006)¹²⁰ Design: RCT N = 26,586 index dispensings Implementation: 00/0000 Study Start: 11/2002 Study End: 10/2003</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CDSS/CDS/CCDS/ reminders CPOE/POE system, EHR/EMR system, Pharmacy</p> | <p>Ambulatory care</p> | <p>compliance rate</p> | <p>There was no significant difference between the control and intervention group physicians in the overall rate of compliance with ordering the recommended laboratory monitoring for patients prescribed study medications. Laboratory monitoring was performed as recommended 56.6% of the time in the intervention group compared with 57.1% of the time in the control group (p = 0.31). In cases in which a statistically significant difference was demonstrated, improved compliance favored the intervention group 71.2% vs. 62.3% (p = 0.003) for gemfibrozil; 75.7% vs. 73.9% (p = 0.05) for statins, 52.8% vs. 46% for colchicine (p = 0.05); 42.9% vs. 0% for methotrexate (p = 0.03)</p> | <p>-</p> |
| <p>Patel (2009)²²⁴ Design: Before-after N = 25,503 patients Implementation: 00/2001 Study Start: 00/2001 Study End: 00/2007</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Hospital information system</p> | <p>Acute care/tertiary, 617 Beds Academic</p> | <p>percent compliance*</p> | <p>There were no statistical differences in percent compliance for all outcomes at 2001 baseline between hospitals (p >0.05). Adherence to all outcome criteria in the 5 high-risk populations over the 6-year time frame resulted in a 119% change compared with 91% at the non-REACH® hospital (p = 0.470).</p> | <p>+</p> |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|---|--|---------|
| Quinn (2008) ¹²⁷ Design: RCT N = 30 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders Diabetes Management Tool Integrated Web-based data analytics and therapy optimization tools | Ambulatory care | Changes in medication (medication intensified) | Patients using WellDoc System were more likely to have physicians intensify diabetes medications (84.6% vs. 23.08%, p = 0.002). | + |
| Raebel (2005) ¹²⁹ Design: RCT N = 10,169 dispensings Implementation: 00/0000 Study Start: 09/2002 Study End: 12/2003 | CDSS/CDS/CCDS/ reminders Integrated Laboratory system, Pharmacy | Ambulatory care HMO pharmacy | percentage of dispensings with baseline monitoring | Recommended laboratory monitoring was completed in 74.7% (n=7,598) of dispensings at initiation of therapy. Compared to the usual care group, monitoring was higher in the intervention group (70% vs. 79%, RRR -13%, p <0.001) | + |
| Riggio (2009) ¹³³ Design: Before-after N = 100 patients with heparin induced thrombocyte-penia Implementation: 06/2005 Study Start: 03/2004 Study End: 09/2006 | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated Hospital information system | Acute care/tertiary, 728 Beds Academic | Time from platelet count criterion until heparin/enoxaparin stop*, Time from platelet count criterion until 1st HIT laboratory test drawn*, Time from platelet count criterion until direct thrombin inhibitor started* | Counter to expectations, the time (in days) taken from alert to heparin stop order was significantly higher after implementation (1.3 vs. 2.9, p = 0.04). There were no significant differences in time (in days) from alert to lab test (2.3 vs. 3.0, NS), nor time to start of treatment with direct thrombin inhibitor (19.3 vs. 15.0, NS). | - |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|---|---|---------|
| Rind (1994) ²²⁵ Design: Time series N = 562 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Formulary, Hospital information system, Laboratory system, Pharmacy | Acute care/tertiary, 504 Beds Academic | mean time interval between event and medication change* | The mean interval between the occurrence of an event and discontinuation of a medication or a change in its dosage was 97.5 hours during the control period and 75.9 hours during the intervention period, a difference of 21.6 hours (p <0.0001). | + |
| Rollman (2002) ¹³⁶ Design: RCT N = 200 Patients with documented major depression Implementation: 00/0000 Study Start: 04/1997 Study End: 12/1998 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | antidepressant prescribing rate (secondary) | Prescribing of antidepressants (continuous use of change in prescriptions) did not differ across the 3 groups at 3 or 6 months. | - |
| Rood (2005) ¹³⁷ Design: RCT N = 484 patients Implementation: 04/2001 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders Integrated Hospital information system | Critical care units (CCU, ICU, NICU) 18 Beds Academic | adherence to glucose measurement timing recommendations*, adherence to insulin dose advice* | Rate of compliance with glucose measurement timing recommendations control-intervention-control (29% vs. 38% vs. 41% with period 2 and 3 greater than period 1, p = 0.05). During the intervention period the rate for computerized group was higher than the control (36% vs. 40%, p = 0.05) Rate of compliance with insulin dose advice was higher in period 2 than 1, and decreased significantly in period 3 (56% vs. 70% vs. 42%, p = 0.05). During the intervention period the rate for computerized group was higher than the control (64% vs. 77%, p = 0.05). | + |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|---|--|---------|
| Roumie (2006) ²²⁶ Roumie (2007) ²²⁷ Design: RCT N = 871 patients Implementation: 00/0000 Study Start: 06/2004 Study End: 12/2004 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care Outpatient hospital based | prescribing changes* | No differences were seen comparing the groups who had provider education alone vs. those who had provider education and computer alerts for prescribing of any medication, changing doses, or adding medications (all data adjusted for multiple variables). | - |
| Safran (1995) ¹⁴¹ Safran (1993) ¹⁴² Design: RCT N = 349 patients with HIV Implementation: 00/0000 Study Start: 05/1992 Study End: 09/1993 | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated EHR/EMR system | Ambulatory care Academic | mean response time to alerts*, mean response times to reminders* | Physicians who got alerts responded more quickly to them (mean 52 vs. 11 days, p <0.0001). Physicians who got reminders responded more quickly to them (mean 500 vs. 114 days, p = 0.0001). | + |
| Schnipper (2008) ¹⁴³ Design: Before-after N = 30 clinicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | Antiplatelet prescribed or contraindication documented*, Beta-blocker prescribed *, Change in diabetic therapy if A1c > 7.0 * | Antiplatelet prescribed or contraindication documented improved from 3.2% in the preintervention to 31.0% in the postintervention period (p <0.001). Beta-blocker prescribed or contraindication documented was 4.2 % in the preintervention compared to 66.7% in the post period (p = 0.03). Change in diabetic therapy if A1c >7.0 was 10.7% in the pre-period and 16.9% in the post period, p = 0.11. | + |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|---|---------|
| Sequist (2005) ²²⁸ Design: RCT N = 6,243 Patients Implementation: 07/2000 Study Start: 10/2002 Study End: 04/2003 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Imaging systems, Laboratory system | Acute care/tertiary, General Hospital, Ambulatory care Community (school, community centre etc) Academic | compliance rate with Diabetes reminders*, compliance rate with Coronary Artery Disease reminders* | Diabetes reminders resulted in the recommended action in 19% in the intervention group vs. 14% in the control group. After adjusting for baseline patient and physician characteristics, patients in the intervention group were more likely than control patients to receive recommended diabetes care based on the composite outcome (OR 1.30, 95% CI 1.01 to 1.67). Coronary artery disease reminders resulted in the recommended action for overdue items in 22% in the intervention group vs. 17% in the control group. Using the composite outcome, patients in the intervention group received recommended coronary artery disease care more often than those in the control group (OR 1.25, 95% CI 1.01 to 1.55) after adjusting for baseline differences. | + |
| Shiffman (2000) ¹⁴⁷ Design: Before-after N = 9 physicians Implementation: 00/0000 Study Start: 09/1996 Study End: 10/1998 | CDSS/CDS/CCDS/ reminders Integrated Handheld | Ambulatory care | adherence rate with metered-dose inhaler/nebulization*, rate of systemic corticosteroid prescriptions* | Adherence with metered-dose inhaler/nebulization rates did not differ between control and intervention (73% vs. 91%, NS), nor did rate of prescribing systemic corticosteroids (43% vs. 57%, NS). | - |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|---|--|---------|
| Tierney (2003) ¹⁶⁶ Design: RCT N = 706 patients, 20 pharmacists, 94 physicians and 1 nurse practitioner Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Pharmacy, | Ambulatory care Outpatient hospital based, Academic | compliance with cardiac care suggestions* | Neither the physician nor the pharmacist intervention had any significant effect on whether patients' cardiac care was compliant with the suggestions (p > 0.8 across the 4 intervention groups by analysis of variance, with p > 0.7 and p > 0.4 when testing the physician and pharmacist interventions separately). | - |
| Tierney (2005) ¹⁶⁷ Design: RCT N = 706 patients Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996 | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system, Pharmacy | Ambulatory care Pharmacy, Outpatient hospital based, Academic | adherence to the care suggestions* | There were no differences between the four study groups in either adherence to the care suggestions, combined or individually (32% control, 32% physician intervention, 32% pharmacist intervention, 37% both interventions, NS). | - |
| White (1984) ²²⁹ Design: RCT N = 396 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Imaging systems, Laboratory system | Acute care/tertiary, Academic | physician actions* | Physicians were 1.22 times as likely to take action in the alert group as compared to the non-alert group (p <0.003). Actions included medication and lab monitoring changes. | + |

Evidence Table 5. KQ1: primary process outcomes for all technologies assisting monitoring (continued)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|---|--|--------------------------------|---|--|---------|
| Wrona (2007) ¹⁷⁷ Design: Observational study N = 536 PCA patients Implementation: 00/2003 Study Start: 01/2003 Study End: 03/2004 | CPOE/POE system Integrated EHR/EMR system, Imaging systems, Laboratory system | Pediatric stand alone hospital | Rates of respiratory monitoring. Rates of oxygen saturation monitoring | Compared to the control group of 'no order set', patients in the Acute Pain Team Service had a higher rate of respiratory monitoring (43% vs. 66.3%, RRR -54%, p <0.05) and oxygen saturation monitoring (86.1% vs. 98.6%, RRR -15%, p <0.05). Compared to the control group of 'no order set', patients in the prescriber initiated PCA had higher respiratory rate monitoring (43% vs. 57.8%, RRR -34%, p <0.05). No other comparisons were significant. | + |

Evidence Table 6. KQ1: Primary Process outcomes for all technologies assisting education and other aspects of medication management

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|---|--|---------|
| Agrawal (2009) ²³⁰ Design: Before-after N = 19,356 MedRecon events Implementation: 06/2006 Study Start: 08/2006 Study End: 12/2007 | CDSS/CDS/CCDS/ reminders medication reconciliation at admission, transfer and discharge Integrated CPOE/POE system, EHR/EMR system | Unspecified Hospital, 630 Beds Other, Academic | compliance with performing medication reconciliation* | On a monthly basis, clinicians performed medication reconciliation more often after the MedRec system and its reminder system were in place. Compliance improved from approximately 34% to 98% to 100%, statistically significant improvement. | + |
| Grasso (2002) ²³¹ Design: Before-after N = 200 discharge summaries Implementation: 04/2001 Study Start: 06/2000 Study End: 07/2001 | PDA use to construct discharge summaries Handheld Integrated EHR/EMR system, Pharmacy | Other specialty hospital (rehab, oncology), Inpatient hospital based | errors rate | The rate of errors in discharge summaries from a psychiatric hospital decreased after the implementation of PDAs to produce the summaries (22% vs. 8%, RRR 64%, p <0.05). | + |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: GP = General Practitioner; HIT = Health Information Technology; N= Sample Size; p = Probability; PDA = Personal Digital Assistants ; PDMW = Physician Discharge Medication Worksheet; RRR = Relative Risk Reduction; vs. = Versus

Evidence Table 6. KQ1: Primary Process outcomes for all technologies assisting education and other aspects of medication management (cont'd)

| Article Information | HIT Studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|---|--|----------|
| <p>Poole (2006)²³² Design: Before-after N = 100 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Medication Reconciliation Discharge System Integrated e-MAR, Hospital information system</p> | <p>General Hospital, Inpatient hospital based</p> | <p>frequency discrepancies *, Dose discrepancies*, Therapeutic drug duplication discrepancies*</p> | <p>The PDMW was found to be effective in reducing discrepancies in frequency and dose and reducing therapeutic drug duplication at the time of discharge. Resolution of discrepancies in frequency improved by 65% with the tool (18% vs. 76%, p <0.001). Resolution of discrepancies in dosages improved by 60% (28% vs. 82%, p <0.001), and therapeutic drug duplication was addressed in 58% more cases (p <0.001).</p> | <p>+</p> |
| <p>Quinn (2008)¹²⁷ Design: RCT N = 30 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>CDSS/CDS/CCDS/ reminders Diabetes Management Tool Integrated Web-based data analytics and therapy optimization tools</p> | <p>Ambulatory care</p> | <p>changes in medication (medication intensified)</p> | <p>Patients using WellDoc System were more likely to have physicians intensify diabetes medications (84.6% vs. 23.08%, p = 0.002).</p> | <p>+</p> |
| <p>van der Kam (2001)²³³ Design: Cohort study N = 1,149 medications Implementation: 00/0000 Study Start: 02/1998 Study End: 05/1998</p> | <p>e-Rx, Pharmacy information system Integrated Pharmacy</p> | <p>Unspecified Hospital, Pharmacy, Not specified</p> | <p>Agreement of GP and pharmacist with patient for drug reported on admission, Agreement of GP and pharmacist with patient for drug reported on 10 days after discharge</p> | <p>Agreement of GP and pharmacist with patient for drugs reported was 31% for paper-based group compared to 49% for electronic group on admission (RRR 58%, p <0,001). The figures on 10 days after discharge were 33% and 53% respectively (RRR 61%, p <0.001). Total number of drugs reported by patients on admission was 38% and 29% for paper-based and electronic groups respectively. The figures on 10 days after discharge were 38% and 28% respectively.</p> | <p>+</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|------------------------------|---|-----------------------------------|--|--|---------|
| Chan (2008) ²³⁴ Design: Survey N = 1,174 nursing homes Implementation: 00/0000 Study Start: 09/2004 Study End: 12/2004 | Administering, Dispensing | BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated EHR/EMR system, Pharmacy | Long term care (nursing homes) | factors associated with use of electronic drug dispensing systems*, factors associated with use of electronic medication administration records* | nursing home facility characteristics associated with use of electronic information systems for drug dispensing: northwest region (OR 0.45 95% CI 0.31 to 0.67), west region (OR 1.63 (95% CI 1.10 to 2.43), administrator in place 5-9 years, (0.54, 95% CI 0.37 to 0.78) and number of services offered (OR 1.23 (95% CI 1.13 to 1.34). Factors associated with medication administration records use in nursing homes included northwest (OR 0.43, 95% CI 0.27 to 0.66) and west (OR 1.85, 95% CI 1.24 to 2.75) regions, urban centers (0.7, 95% CI 0.50 to 0.97), occupancy rates of 70 to 79% (OR 0.41, 95% CI 0.23 to 0.72), administrator in place <5 years (OR 0.49, 95% CI 0.33 to 0.71) and number of services offered (OR 1.21, 95% CI 1.10 to 1.31). | - |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: ASyMS = Advanced Symptom Management System; BCMA = Bar Code Medication Administration; BPOC = Barcode-enabled Point of Care; CCDS = Computerized Clinical Decision Support; CCU = Critical Care Unit; CDS = Clinical / Computerized Decision Support; CDSS = Clinical Decision Support System; CI = CI; CPOE = Computerized Provider Order Entry; DDI = drug drug interaction; DHCp = Decentralized Hospital Computer Program; HER = Electronic Health Record; e-MAR = Electronic Medication Administration Record; EMR = Electronic Medical Records; e-RX = Electronic Prescribing; e-TAR = Electronic Treatment Authorization Request; GP = General Practitioner; HIT = Health Information Technology; ICU = Intensive Care Unit; MAS-NAS = Medication Administration System-Nurses Assessment of Satisfaction; MM = Medication Management; mos = Months; N = Sample Size; NICU = Neonatal Intensive Care Unit; NPs = Nurse Practitioners; OR = OR; p = Probability; PHR = Patient Health Record; POE = Provider Order Entry; PWS = Physician Workstation; r = Correlation Coefficient; RCT = Randomized Controlled Trial; RR = Relative Risk; RRR = Relative Risk Reduction; SES = Socioeconomic; SYW = Show Your Work (program); vs. = Versus

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|-------------|--|-------------------------------|--|---|---------|
| Ghahramani (2009) ²³⁵ Design: Survey N = 413 health professionals working in a hospital Implementation: 05/2005 Study Start: 11/2005 Study End: 00/0000 | Prescribing | CPOE/POE system | Acute care/tertiary, Academic | factors associated with user satisfaction*, Frequency of use | User satisfaction was negatively associated with age ($p < 0.0001$) and positively associated with system familiarity ($p < 0.0001$), frequency of use ($p < 0.0001$) and system characteristics ($p < 0.0001$). Frequency of use was negatively associated with age ($p < 0.001$) and training ($p = 0.002$) and was positively associated with user satisfaction ($p < 0.0001$), user friendliness ($p < 0.0001$), system familiarity ($p = 0.0002$), and system characteristics ($p < 0.0001$). | + |
| Glassman (2006) ²³⁶ Design: Survey N = 97 clinic physicians and NPs (prescribers) Implementation: 03/2000 Study Start: 12/2000 Study End: 12/2002 | Prescribing | CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Hospital information system | Ambulatory care | knowledge of DDI*, perceptions towards CPOE* | The groups differed for knowledge for 19 drug interactions, improved for 3 and decreased for 1. Knowledge did not differ by specialty. At year 1, 45% of clinicians preferred CPOE for prescriptions. By year 3 this had increased to 63%, $p < 0.001$. the other 4 reported perceptions had not changed. 8 barriers were assessed with both surveys. 6 did not differ between the 2 time periods. The perception that important alerts were missing (15% vs. 29%, $p = 0.01$) and poor visual presentation (7% vs. 21%, $p = 0.02$) differed. | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--------------------|---|--|---|---|----------|
| <p>Graumlich (2009)²³⁷ Graumlich (2009)²³⁸ Design: RCT N = 631 patients Implementation: 00/0000 Study Start: 11/2004 Study End: 01/2007</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system</p> | <p>Acute care/tertiary, 730 Beds Academic</p> | <p>patient mean score for discharge preparedness*, patient score for satisfaction with medication information, outpatient physicians perception of discharge software</p> | <p>When comparing patients assigned to discharge software vs. usual care, patient mean (SD) scores for discharge preparedness were higher (17.7 [4.1] vs. 17.2 [4.0]; p = 0.042), patient score for satisfaction with medication information were unchanged (12.3 [4.8] vs. 12.1 [4.6]; p = 0.567) and their outpatient physicians scored higher quality discharge (17.2 [3.8] vs. 16.5 [3.9]; p = 0.027). Hospital physicians found mean effort to use discharge software was more difficult than the usual care (6.5 [1.9] vs. 7.9 [2.1]; p = 0.011) and discharge software users had satisfaction (7.4 [1.4] vs. 7.9 [1.4]; p = 0.129) for usual care physicians</p> | <p>-</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---------------|---|--|---|---|---------|
| Holden (2008) ²³⁹ Design: Mixed methods N = 215 nurses Implementation: 12/2006 Study Start: 11/2006 Study End: 11/2007 | Administering | BCMA Integrated CDSS/CDS/CCDS/reminders CPOE/POE system, EHR/EMR system | Pediatric stand alone hospital, Academic | perceived ease of use*, perceived usefulness*, satisfaction with process* | The predicted process acceptance model was supported pre-BCMA (F(12,88) = 5.13, MSE = 0.67, p <0.05) and post-BCMA (F(12,61) = 5.00, MSE = 0.61, p <0.05). Perceived ease of use of the process was significantly and uniquely associated with process acceptance both pre-BCMA ($\beta = 0.28$, p <0.05) and post-BCMA ($\beta = 0.49$, p <0.05). Perceived usefulness was significantly associated with process acceptance pre-BCMA ($\beta = 0.40$, p <0.05) but not post-BCMA ($\beta = 0.18$, p = 0.10). These two process beliefs accounted for 31.3% and 32.2% of the variance in process acceptance pre- and post-BCMA, respectively. | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---------------|---|---|--|--|---------|
| Holden (2010) ²⁴⁰ Design: Survey N = 294 nurses Implementation: 07/2007 Study Start: 10/2006 Study End: 08/2007 | Administering | BCMA Integrated CPOE/POE system | Acute care/tertiary, Pediatric stand alone hospital, Academic | perceptions of accuracy, usefulness, consistency, time-efficiency, ease of performance, error likelihood, error detection likelihood | Nurses perceptions of the administration process changed at the hospital that implemented BCMA, whereas perceptions of nurses at the control hospital did not. BCMA appeared to improve the safety of the processes of matching medications to the medication administration record and checking patient identification. The accuracy, usefulness, and consistency of checking patient identification improved as well. In contrast, nurses perceptions of the usefulness, time efficiency, and ease of the documentation process decreased post-BCMA. | + |
| Hurley (2007) ²⁴¹ Design: Mixed methods N = 1,087 nurses Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Administering | BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated CPOE/POE system, EHR/EMR system, Pharmacy | Acute care/tertiary, Academic | satisfaction -The Medication Administration System-Nurses Assessment of Satisfaction Scale (MAS-NAS) | Overall scores on The Medication Administration System-Nurses Assessment of Satisfaction (MAS-NAS) Scale, significantly increased following implementation of the BCMA/e-MAR system (4.1 vs. 5.0, p <0.001). There were significant increases in each of the 3 subscales of efficacy, safety and access (p <0.001). | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|---|--|---|----------|
| <p>Johnson (2010)⁷⁵ Design: RCT N = 3,285 patients Implementation: 00/0000 Study Start: 04/2007 Study End: 08/2007</p> | <p>Prescribing, Transmission, order communication</p> | <p>CDSS/CDS/CCDS/reminders e-Rx Integrated EHR/EMR system</p> | <p>Ambulatory care, Pharmacy, Not specified, Academic</p> | <p>perceptions*</p> | <p>When asked if SYW helped avoid callbacks the majority of respondents agreed or strongly agreed (69%). Pharmacists found the allergy override information helpful (69% agree or strongly agree). A majority of pharmacists (79%) felt that information about patient's insurance eligibility was less helpful; 41% of pharmacists were neutral, 31% were in disagreement, and 7% were in strong disagreement with the statement "SYW was helpful with insurance eligibility".</p> | <p>+</p> |
| <p>Kawasumi (2006)²⁴² Design: Cross-sectional N = 28 primary care physicians caring for 4,096 patients Implementation: 00/0000 Study Start: 03/2003 Study End: 11/2003</p> | <p>Prescribing</p> | <p>drug management system, e-Rx Integrated EHR/EMR system, Insurance</p> | <p>Ambulatory care</p> | <p>rate of use of electronic medication histories, rate of use of e-Rx</p> | <p>Physicians differed in their use of electronic medication histories for patients based on their SES: 10.8% for high SES, 14.6% for middle SES (comparing middle and high, RR 1.55, 95% CI 1.15 to 2.47), and 17.9% for low SES (comparing high and low SES RR 1.70, 95%CI 1.15 to 2.47). Use of e-Rx did not differ by SES (36.1% for high, 39.0% for middle and 37.2% for low SES, NS).</p> | <p>-</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|------------------------------|---|--|---------|
| Kirking (1986) ²⁴³ Design: Survey N = 218 pharmacists Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Transmission, order communica-tion | CDSS/CDS/CCDS/ reminders Pharmacy information system Integrated Pharmacy | Pharmacy, Other | reported frequency of DDI per week, reported number of daily contacts with prescribers, proportion of contacts with prescribers related to DDI and allergies | computer users reported potential drug interactions an average of 16.1 interactions detected per week compared to 8.7 for non-users (p>0.05). As a group, computer users compared to non-users were found to have more contacts per day with prescribers (21.5 vs. 16.0, respectively, p <0.05), and a higher percentage of their reported contacts were related to interaction and allergy problems (3.9% vs. 2.8% respectively, p <0.05). | + |
| Kralewski (2008) ²⁴⁴ Design: Survey N = 93 physicians Implementation: 00/0000 Study Start: 09/2006 Study End: 10/2006 | Prescribing | e-Rx | Ambulatory care, Academic | proportion of prescriptions sent electronically | Practice-level variables explain most of the variance in the use of e-scripts by physicians, although there are significant differences in use among specialties as well. General internists have slightly lower use rates for e- Rx and pediatricians have the highest rates. Larger practices and multispecialty practices have higher use rates, and five practice culture dimensions influence these rates; two have a negative influence and three (organizational trust, adaptive, and a business orientation) have a positive influence. | |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|----------------|---|--|---|--|---------|
| Kramer (2007) ²⁴⁵ Design: Before-after N = 283 patients Implementation: 00/0000 Study Start: 09/2004 Study End: 10/2005 | reconciliation | electronic medication reconciliation system Integrated EHR/EMR system | Unspecified Hospital, 760 beds plus 102 bassinets Beds Inpatient hospital based | improved self-reported perceptions, satisfaction; self-reported perceptions of clear instructions on what medications to take, how much and how often the medications were to be taken, other instructions on taking the medication, potential side effects, and general understanding of the medications | Patients reported satisfaction with the reconciliation system (better knowledge of their discharge medications) for 5 of 5 factors. Improved self-reported perceptions of clear instructions on what medications to take (p = 0.007), how much and how often the medications were to be taken (p = 0.007), other instructions on taking the medication (p = 0.006), potential side effects (p = 0.001), and general understanding of the medications (p = 0.001). Healthcare provider Physician assistants and nurse practitioners reported that patients had clearer instructions on discharge (p = 0.01); how much, how often, and when to take their medications at home (p = 0.05); and the medication discharge process was views as being sufficient for them as care givers (p = 0.0003). | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--------------------|--|---|--|--|----------|
| <p>Lee (1996)²⁴⁶ Design: Survey N = 205 physicians and nurses Implementation: 05/1993 Study Start: 00/1993 Study End: 00/1995</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated CDSS/CDS/CCDS/ reminders</p> | <p>Acute care/tertiary, 720 Beds Academic</p> | <p>correlates of satisfaction*</p> | <p>Overall satisfaction was most strongly correlated with characteristics related to the efficiency of POE, such as impact on productivity (r = 0.69) and ease of use (r = 0.67). Characteristics related to the quality of care, such as reducing error or giving information, were less strongly correlated with overall satisfaction (r = 0.32 and r = 0.36, respectively), although these correlations were still significant.</p> | <p>+</p> |
| <p>Li (2006)²⁴⁷ Design: Qualitative N = 2 qualitative researchers (nurse and human factors psychology) Implementation: 02/2004 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated Hospital information system</p> | <p>Acute care/tertiary</p> | <p>usability The 2 researchers used heuristic methods and identified 5 major problem areas with the CPOE system. These problems centered on text presentation, too much information/too many decisions at one time, color scheme (monochromatic blue/grey with red used as accent and not to note caution or problems).</p> | <p>Problems were given to the developers who addressed them in the next redesign of the system.</p> | |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--------------------|---|---|-------------------|--|----------|
| <p>Lindenauer (2006)²⁴⁸ Design: Mixed methods N = 356 physicians at 2 hospitals. Implementation: 00/0000 Study Start: 05/2003 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Imaging systems, Pharmacy</p> | <p>Acute care/tertiary, 725 Beds Academic</p> | <p>use</p> | <p>The physician users of CPOE were categorized as being low (n = 109), intermediate (n = 88), or high (n = 141) users. Groups did not differ for use by gender, use of a computer in outpatient practice, years since graduate from medical school, practice at study institution, or total number of orders placed. Specialty was associated with use: more anesthesiologists, pediatricians, and surgeons used CPOE (p <0.0001). Physicians who trained with CPOE (p = 0.045) and those who used computers daily were more likely to use CPOE (p = 0.04). High and intermediate users were 3 times as likely to believe that the user interface of the system supported their work flow. Similarly, 19% of low users, 31% of intermediate users, and 45% of high users believed that entering orders into the system was faster than writing orders.</p> | <p>-</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---|--|--|------------------------|---|---------|
| Liu (2004) ²⁴⁹ Design: Survey N = 154 hospitalized patients Implementation: 07/2002 Study Start: 10/2002 Study End: 12/2007 | Education of patients and clinicians but not pre-professional education | Pharmacy information system Integrated Hospital information system | Acute care/tertiary, Academic | patient knowledge* | Patients reported improved drug knowledge: improved abilities to use their prescriptions, avoid adverse drug events, know contraindications and side effects of their drugs, and acquire needed information, p <0.001 for each before and after comparison. | + |
| McAlearney (2005) ²⁵⁰ Design: Observational study N = 1,010 patients Implementation: 04/2002 Study Start: 11/2001 Study End: 11/2003 | Prescribing | CPOE/POE system, order sets Integrated CDSS/CDS/CCDS/reminders Laboratory system, Pharmacy | Pediatric stand alone hospital, 328 Beds Academic | order set utilization* | Order set utilization varied significantly by condition (X ² = 339.2, p <0.001). The asthma order set use rate (88.1%) was highest, followed by appendectomy order set use (79.4%), followed by a relatively low CAP order set use rate (21.1%). | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|------------------------|---|---|---------|
| <p>McCann (2008)²⁵¹ Design: Mixed methods N = 53 patients Implementation: 0 Study Start: 03/2006 Study End: 09/2006</p> | <p>Monitoring including patient adherence and compliance</p> | <p>symptom management system Integrated Handheld</p> | <p>Ambulatory care</p> | <p>preintervention perceptions, postintervention perceptions</p> | <p>Preintervention, patients' expectations of participating were largely positive: 87% anticipated that using the ASyMS© handset would help them communicate with their doctors and nurses; 79% thought using the ASyMS© handset to record symptoms would help manage symptoms; and patients reported positive expectations about the alerting system, frequently using terms such as 'reassuring', 'excellent idea', 'confident' and 'comforting'. Patients anticipated they would find their overall experience of being involved in the study challenging (32%), rewarding (62%), educational (41%) and interesting (63%). Postintervention, patients reported positive experiences of being involved in the study, describing their experience as interesting (80%), valuable (77%) and educational (34%).</p> | |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--------------------|-------------------------------|-----------------------------|---|---|----------|
| <p>Musser (2006)²⁵² Design: RCT N = 51 anesthesiologists Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system</p> | <p>Unspecified Hospital</p> | <p>time used in CPOE sessions in seconds (text-based (OS) vs. graphical format (IF))* rate of miscellaneous orders placed per session (text-based vs. graphical format)*</p> | <p>Users predominately chose to use the IF format: used for 70% of the orders in the free-choice phase, and 17/24 (71%) of survey respondents preferred IF. OS format gained substantial support, 15 of 26 (58%) answered that they would choose to keep either both formats or the OS alone; and those users initially assigned to the IF format were more likely than their counterparts (36% vs. 21%) to prefer the OS format. Experience level (based on number of orders placed) had a small but significant (p = 0.02) correlation with preference of format, with more experienced users preferring for the OS format. According to time measurements from the usage logs, CPOE sessions in which the IF format was used averaged 27 seconds shorter (162 vs. 189 seconds, p <0.01). No statistically significant differences between IF and OS formats were found for length of stay, rate of mistakes made, or the number of orders for diagnostic tests or medications; miscellaneous orders were placed slightly more frequently (5.44 vs. 5.14 orders per session, p = 0.03) from the OS format.</p> | <p>+</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|----------------------------|--|--|--|--|---------|
| <p>Niazkhani (2009)²⁵³ Design: Before-after N = 211 nurses Implementation: 12/2001 Study Start: 09/2003 Study End: 10/2004</p> | Administering, Prescribing | CPOE/POE system Integrated EHR/EMR system, Hospital information system | Acute care/tertiary, 1237 Beds Academic | Overall mean scores of medication process attitudes* Kardex vs. TIMED, correlates of attitudes toward CPOE*. Overall mean scores of attitudes were summed across the 15 questionnaire items (relating to regarding medication process, characteristics of medication orders, registration of drug administration, learning and speed of process and managing non-stock medications) and compared pre- and post for Kardex and TIMED units. | Following implementation of CPOE, there was an increase in scores for the Kardex system (3.2 vs. 3.6, p <0.001) but not for the TIMED units (3.4 vs. 3.5, NS). Overall score with the CPOE was strongly correlated with user satisfaction (r=0.75, p <0.001), clarity of administration record (r=0.66, p <0.001), ease of the process (r=0.63, p <0.001), and clarity of the drug review form (r=0.63, p <0.001) but not with professional status, computer experience or ward. | + |
| <p>O'Morrow Snyder (2003)²⁵⁴ Design: Before-after N = 17 nurses Implementation: 07/2002 Study Start: 01/2003 Study End: 05/2003</p> | Administering | BCMA Integrated CPOE/POE system, Hospital information system, Imaging systems, Laboratory system, Pharmacy | General Hospital, 457 Beds | nurses attitudes toward BPOC | No differences in responses about attitudes toward the BPOC system before or after training and implementation took place (p >0.05) for any of the 7 factors included in the survey: patient care, charting, computer benefit, computer capability, computer characteristics, legal issues or management tools. | - |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|----------------------|--|---|---|--|----------|
| <p>Onzenoort (2008)²⁵⁵ Design: Observational study N = 15,162 drug administrations Implementation: 09/2005 Study Start: 00/0000 Study End: 00/0000</p> | <p>Administering</p> | <p>BCMA Integrated CPOE/POE system, EHR/EMR system</p> | <p>Acute care/tertiary, Academic</p> | <p>variables related to use of BCMA</p> | <p>Nurses verified the bar codes of about half of medications administered. Variables that increased the use of bar code verification were department (more in rheumatology and metabolic and infectious diseases and less in oncology, hematology, and gastroenterology and neurology and neurosurgery); oral administration (and not parenteral, inhalation, rectal or other); more with more than 46 nurses; more with nurses younger than 30 years; more with 6-7 shifts worked per month.</p> | <p>-</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--------------------|---|---|---|---|----------|
| <p>Pirnejad (2008)²⁵⁶ Pirnejad (2009)²⁵⁷ Design: Mixed methods N = 149 nurses Implementation: 09/2003 Study Start: 11/2003 Study End: 06/2007</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/reminders CPOE/POE system Integrated EHR/EMR system, Hospital information system</p> | <p>Acute care/tertiary, 1237 Beds Academic</p> | <p>nurses attitudes toward paper based system vs. CPOE*</p> | <p>²⁵⁶ The first question asked nurses whether their current medication system supported their work process and showed a statistically significant difference between nurses' attitudes in pre- and postimplementation (60.5% agreed for the paper-based system and 68.5% for the CPOE system, p = 0.048). More in the paper group said 'no' (32.9% vs. 2.7%), while fewer were unsure (3.9% vs. 28.8%).²⁵⁷ Questions about the perceived physical appearance of the prescription and administration registrations system were analyzed; The analysis showed that nurses judged CPOE system prescriptions to be significantly better than those from the paper-based system with regard to legibility (p <0.001) and completeness (p <0.001). However, there was no statistically significant difference between prescription layout in the two systems (p > 0.006).</p> | <p>+</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|-----------------------------------|--|----------|---|--|---------|
| Porteous (2003) ²⁵⁸ Design: Mixed methods N = 787 General practitioners, opinion leaders, computing experts, pharmacists, and patients Implementation: 00/0000 Study Start: 01/2000 Study End: 05/2000 | Transmission, order communication | electronic communication between physicians and pharmacists, e-Rx | Pharmacy | percentage of responders supporting electronic transfer | Responders in all three groups (68% of patients [95% CI 64% to 72%], 83% of GPs [95% CI 77% to 89%], and 87% of community pharmacists [95% CI 82% to 92%]) thought that electronic transfer of prescription related information was a good idea in principle. All groups were concerned about security and sharing confidential patient information. | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|-------------|--|---|---|---|---------|
| Rahimi (2009) ²⁵⁹ Design: Survey N = 310 healthcare providers Implementation: 00/2008 Study Start: 02/2009 Study End: 00/2009 | Prescribing | CPOE/POE system Integrated EHR/EMR system, Formulary, medication administration, Pharmacy | Unspecified Hospital, Ambulatory care | Perceptions Data analyzed physicians and nurses' experiences related to the adoption of a CPOE system, structuring the analyses according to three attributes of diffusion of innovation theory, i.e. the relative advantage of the system, its compatibility with professional values and needs, and its complexity of use. | More nurses (56.7%) than physicians (31.3%) stated that the CPOE system introduction had worked well in their clinical setting (p <0.001). Similarly, more physicians (73.9%) than nurses (50.7%) reported that they found the system not adapted to their specific professional practice (p = < 0.001). Also more physicians (25.0%) than nurses (13.4%) stated that they did want to return to the previous system (p = 0.041). Relative advantages of the CPOE system were estimated to be significantly (p <0.001) higher among nurses (39.6%) than physicians (16.5%). Physicians' agreements with the compatibility of the CPOE and with its complexity were significantly higher than the nurses (p <0.001). An important reason behind the reluctance of physicians and nurses to use the CPOE system was that the system was not adapted to their work routines. | - |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|--|--|---------|
| Rogers (1999) ²⁶⁰ Design: Before-after N = 42 practices Implementation: 10/1995 Study Start: 04/1997 Study End: 10/1997 | Prescribing | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Ambulatory care | use of the decision support system in 2 time periods | More clinicians used the decision support system in phase 2 as compared with phase 1: 9.3% vs. 27%, RRR 186, p <0.001. | + |
| Rohrig (2007) ²⁶¹ Design: Cross-sectional N = 40 physicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system | Critical care units (CCU, ICU, NICU) Academic | usability of Antibiotic Wizard* | In a survey physicians compared Antibiotic Wizard with Microsoft Word on 6 scales. 3 of the scales showed no difference in comparison (suitability for the task, conformity with user expectations, and suitability for individualization). For the other 3 scales, Antibiotic Wizard was perceived as being better than Word: self descriptiveness, controllability, and error tolerance (p <0.01 for all 3 comparisons). | + |
| Rosenbloom (2005) ²⁶² Design: RCT N = 418,739 opportunities to access an information item Implementation: 00/1995 Study Start: 04/19999 Study End: 03/2000 | Prescribing | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system | Acute care/tertiary, 609 Beds Academic | access rate for educational opportunities | Study physicians accessed educational opportunities for 278 of 240,504 (0.12%) vs. 18 of 178,235 opportunities (0.01%), RRR 1100, p <0.05. | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|---|--|--|---|--|---------|
| Ross (2004) ²⁶³ Design: RCT N = 107 patients Implementation: 00/0000 Study Start: 08/2001 Study End: 09/2002 | Monitoring including patient adherence and compliance | Patient accessible Medical Record Integrated messaging system | Acute care/tertiary, Ambulatory care, Academic | change in the self-efficacy domain of the Kansas City Cardiomyopathy Questionnaire* | A change of 7.7 was set to be minimal clinically significant difference. There was a trend towards an improvement in the intervention group, with scores of 85 at baseline, 88 at 6 mos and 91 at 12 mos (p = 0.08); but the improvement of 6 points did not reach the threshold value set as a standard for this outcome. | - |
| Rotman (1996) ²⁶⁴ Design: RCT N = 34 Physicians Implementation: 00/0000 Study Start: 07/1994 Study End: 06/1995 | Prescribing | CDSS/CDS/CCDS/reminders e-Rx Integrated Hospital information system, Laboratory system | Ambulatory care | User Satisfaction Rating* | After the physicians used the PWS, their user-satisfaction, score decreased by 0.34 Likert-scale units (approximately one half of one SD of the mean score, p = 0.008). In contrast, the mean satisfaction in the control group (DHCP) increased by 0.49 Likert-scale units (p <0.0001). Overall, the two groups diverged with a difference of 0.83 Likert-scale units between the two groups (p <0.0001). | - |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---------------------------------------|----------------------------|------------------------|--|---------|
| Rupp (2008) ²⁶⁵ Design: Survey N = 1,094 pharmacy staff Implementation: 00/0000 Study Start: 04/2006 Study End: 07/2006 | Dispensing, Transmission, order communication | e-Rx Integrated Pharmacy | Pharmacy Pharmacy chain | satisfaction with e-Rx | Pharmacy personnel reported general satisfaction but also perceived key weaknesses with electronic prescribing (e-Rx). Pharmacists, considered e-Rx technology to be significantly more positive in terms of safety, efficacy, and efficiency than pharmacy technicians. Effect on pharmacy efficiency was the most influential predictive variable for determining staff satisfaction with e-Rx; followed by communications with the physician and patient safety (final model retained were as follows: satisfaction = 0.6071+ 0.3562 efficiency + 0.2075 communications with physician + 0.1720 safety + 0.1698 relations with patient + 0.1487 effectiveness). | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--------------------|--|---|--|--|----------|
| <p>Schectman (2005)²⁶⁶ Design: Survey N = 84 physicians Implementation: 09/2003 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>e-Rx Integrated, CDSS/CDS/CCDS/ reminders</p> | <p>Ambulatory care, Academic</p> | <p>correlates of use* System utilization rate was strongly associated with perceived actual ease of use and usefulness; perceptions that their patients liked them to use the system; perceived efficiency; and feeling comfortable. Interestingly, use was not associated with satisfaction with ease of use or system capabilities since even high utilizers felt that the system should be easier to use. Utilization was also not associated with the amount of prior computer experience or with clinical practice satisfaction.</p> | <p>There was a strong association between self reported rate of recent system use and the actual number of prescriptions written in the two months prior to the survey based on electronic utilization data (r = 0.70, p <0.0001). There was no association between respondent age, gender, or level of training and utilization. There was no overall association between the mean score on the scale of attitudes toward computers' effect (beneficial vs. detrimental) on the practice of medicine and utilization of the expert system (p = 0.18). However, there was an association between prescription writing and the specific beliefs that computers enhanced the enjoyment of the practice of medicine (p = 0.04) and the quality of health care (p = 0.004).</p> | <p>+</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|----------------------|--|---|---------|
| Schmidt (2008) ²⁶⁷ Design: Cohort study N = 62 patients with CHF Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Monitoring including patient adherence and compliance | patient adherence reporting Integrated EHR/EMR system | Ambulatory care | association between self report and pill container | Clinicians could not assess patient adherence. Patients' self reports of adherence were similar to what was measured using the automated pill boxes and response to the EHRs. 51% wanted to continue after 6 months of monitoring. No changes were noted in physical health but both groups reported improved levels of mental health. p <0.001 at 2 months p <0.01 at 6 month association between self report and pill container (NS). | + |
| Shannon (2005) ²⁶⁸ Design: Cohort study N = 9 physicians Implementation: 00/1999 Study Start: 00/0000 Study End: 00/0000 | Prescribing | e-Rx Integrated, Hospital information system | Emergency department | rate of use of e-Rx* | The addition of wireless handheld computers resulted in a statistically significant increase in prescription-writing by physicians. The mean of the observed rates of prescribing was 52% during the control period and 64% during the intervention period, a 12.5% increase (SE 0.057, p = 0.03). | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--------------------|---|--|--|--|----------|
| <p>Sittig (2006)²⁶⁹ Design: Survey N = 110 PCPs Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system</p> | <p>Ambulatory care, Academic</p> | <p>factors affecting CDS system acceptance</p> | <p>Younger clinicians appreciated the system more than those who had been practicing longer in the hospital, most respondents liked the system (helped take care of patients better (3.5), work the time it takes (3.5), and reminds me of something I had forgotten (3.1)), cost. safety and health maintenance reminders were valued about the same; clinicians were more likely to look up patient information (3.9), enter orders for patients (3.8), show patient data (2.9); they wanted fewer alerts and noted that they came at an inappropriate time.</p> | |
| <p>Tan (Woan Shin Tan) (2009)²⁷⁰ Design: Survey N = 179 health care providers Implementation: 00/2006 Study Start: 10/2007 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system, e-Rx Integrated CDSS/CDS/CCDS/reminders Pharmacy</p> | <p>Ambulatory care, Other institution based</p> | <p>physician satisfaction with e-Rx*</p> | <p>85% of physicians were satisfied with the e-Rx system. Their satisfaction was associated with the ability to prescribe a new medication ($p = 0.002$) or change an existing one ($p = 0.05$), and the amount of time taken to enter prescription information ($p = 0.04$). 77% of pharmacists were satisfied with the system. their satisfaction was associated with the amount of time spend on processing standard purchases ($p = 0.05$).</p> | <p>+</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---------------|---|--------------------------------|--|--|---------|
| Tierney (1994) ²⁷¹ Design: Survey N = 212 Physicians and Medical students Implementation: 00/0000 Study Start: 04/1990 Study End: 10/1991 | Prescribing | CPOE/POE system Integrated EHR/EMR system | General Hospital | overall attitude score | Factor analysis created an overall attitude score of 16 items loading into 3 factors that explained 57% variance. The score was significantly different between groups (p = 0.0002) declining progressively from juniors (mean 47.1, SD 7.0) interns (mean 44.3, SD 7.1), and residents (40.9, 6.9). Junior students sig dif from all other groups. Gender, typing ability and computer ownership not factors. | - |
| Topps, (2005) ²⁷² Design: Mixed methods N = 313 Healthcare provider Implementation: 11/2002 Study Start: 05/2002 Study End: 06/2003 | Administering | BCMA Integrated Billing/administration system, Hospital information system, Pharmacy | Pediatric stand alone hospital | perceived effect of new system on medication errors*, perceived staff time using system* | The comparison of pre and post tests revealed that statistically significant pre vs. post differences were observed for perceived effect of new system on medication errors". Mean pre bar-code was 1.91; mean post-score-2.23 with difference between means statistically significant (F=6.55; df = 1, 308; p = 0.011); however the score was higher post bar-code than pre bar-code, indicating that the staff felt errors had not decreased as much as they thought they would. For staff time using system, mean pre bar-code was 3.55; mean postbar-code was 3.95 with difference between means statistically significant (F = 8.80; df = 1, 312; p = 0.003). | - |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--------------------|---|---|---|--|----------|
| <p>Wang (2009)²⁷³ Design: Mixed methods N = 228 physicians (139 e-Rx users and 89 non-users) Implementation: 09/2006 Study Start: 10/2006 Study End: 12/2006</p> | <p>Prescribing</p> | <p>e-Rx Integrated Stand-Alone</p> | <p>Ambulatory care</p> | <p>predictors of use of e-Rx*</p> | <p>Specialty (p = 0.05) and practice setting (smaller practices) (p = 0.002) were associated with use but not age, attitude towards computer, practice size, or use of EMRs. Performance measures were associated with volume of use of e-Rx (p <0.001) and usability issues were associated with stopping use of e-Rx (p = 0.03).</p> | |
| <p>Weiner (1999)²⁷⁴ Design: Survey N = 271 clinicians Implementation: 06/1996 Study Start: 11/1996 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated Imaging systems, Laboratory system, Pharmacy</p> | <p>Acute care/tertiary, Academic</p> | <p>perceptions of nurses and physicians towards CPOE*</p> | <p>More nurses reported the POE system easier to use than house officers and attendings (78% vs. 63% vs. 37%, p <0.03). House officers and attendings were more likely than nurses to report the use of POE decreased their time with patients (9% for nurses, 44% for house officers, and 34% for attendings, p <0.05). House officers were more likely than nurses to state that POE was associated with more tests and more errors in ordering. More nurses felt that the system improved the quality of care (56%) compared with 29% for house officers and 34% for attendings, p <0.03).</p> | <p>+</p> |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|--|-------------------------------------|--|---------|
| Weingart (2008) ²⁷⁵ Design: Cohort study N = 267 patients Implementation: 09/2000 Study Start: 04/2001 Study End: 06/2002 | Monitoring including patient adherence and compliance | patient messaging via PHR Integrated Billing/administration system, EHR/EMR system, Imaging systems, Laboratory system | Ambulatory care, Academic | patient usage of MedCheck messages* | Patients opened 79% of MedCheck messages and responded to 12%; 77% responded within 1 day. Patients often identified problems filling their prescriptions (48%), problems with drug effectiveness (12%), and medication symptoms (10%). Clinicians responded to 68% of patients' messages; 93% answered within 1 week. Clinicians often supplied or requested information (19%), or made multiple recommendations (15%). | |
| Wilson (2000) ²⁷⁶ Design: Survey N = 112 prescribers and pharmacy staff Implementation: 00/1990 Study Start: 05/1998 Study End: 06/1998 | Prescribing | CPOE/POE system Integrated Billing/administration system, Hospital information system, Imaging systems, Laboratory system, Pharmacy | Unspecified Hospital, 48 Beds Ambulatory care | correlates of satisfaction* | Overall, users were satisfied with the CHCS POE system. Satisfaction was significantly positively correlated with ratings of the POE system's impact on productivity, ease of use, effect on the quality of care, reliability, and provision of information to help providers write better orders (p <0.05). | + |

Evidence Table 7. KQ1: primary intermediate outcomes for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|-------------|--|---|--|--|---------|
| Zaidi (2006) ²⁷⁷ Design: Survey N = 115 clinicians Implementation: 00/0000 Study Start: 02/2005 Study End: 08/2005 | Prescribing | antibiotic approval program, CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Acute care/tertiary, Inpatient hospital based, Academic | clinicians' perceptions of ease of use and usefulness of a web-based antibiotic approval system* | Use of the iApprove CDSS system was negatively associated with number of years of experience (p = 0.004). Use was positively associated with self-rated computer sophistication (p = 0.03), frequency of accessing laboratory data (p = 0.012), the system was perceived to be easy to learn (p = 0.001) or easy to show others how to use the system (p = 0.014), or if they perceived the system to be integrated into daily work flow (p = 0.028), the perceived ease of finding additional information related to recommendation (p = 0.009), and ease of logging out of the system (p = 0.034). | + |

Evidence Table 8a. Summary of full economic evaluation studies

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Perspective (Time horizon) | Currency (year) Cost elements | Effective-ness measure | Intervention and alternative being evaluated | Main economic findings |
|--|-----------------------------|--|--|---|---|--|---|---|--|
| Fretheim (2006) ⁵⁵ Fretheim (2006) ⁵⁶ Norway | Cost-effectiveness analyses | Compared costs and effects of a multifaceted intervention aimed at improving prescribing of anti-hypertensive and cholesterol lowering drugs compared with usual care. | Using data from a cluster-RCT of private practices, the cost-effectiveness included the cost incurred per additional patient started on a thiazide rather than another anti-hypertensive drug. | Intervention: 73 practices with 70 included in analysis; control: 73 with 69 included in analysis | Perspective of the health care system, (1 year) | 2002 USD (used 2002 avg. exchange rate from Norwegian kroner) Development of software; training of outreach visitors; printed material; travel; cost of pharmacists doing outreach; admin costs; opportunity cost of physician time; technical support; drug expenditure; number of consultations per patient; laboratory tests | Number of patients prescribed thiazides for hypertension, number of patients that had a cardiovascular risk assessment done, number of patients who achieved treatment target goal (BP, LDL, total cholesterol) | multifaceted intervention: (1) educational outreach visits to clinics; (2) audit & feedback on current adherence to guidelines & recommendations; (3) computerized reminders to physicians during pt encounter vs. passive dissemination of guidelines through national medical journal | The cost-effectiveness of the intervention was USD\$454 per additional patient started on thiazides. |

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: ADE = adverse event; BP = blood pressure; BWH = Brigham and Women's Hospital; CDSS = computerized decision support system; CPOE = computerized physician order entry; CVR = cardiovascular risk; ESCHM = European Society of Cardiology and other societies for Hypercholesterolemia Management; GINA = Global Initiative for Asthma; LOS = length of stay; MAR = medication administration record; MOE = medication ordering entry; pADE = preventable adverse drug events; QOL = quality of Life; RCT = randomized controlled trial; SADC = system of clinical decision support; SGRQ = St. George Respiratory Questionnaire; USD = United States Dollars

Evidence Table 8a. Summary of full economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Perspective (Time horizon) | Currency (year) Cost elements | Effective-ness measure | Intervention and alternative being evaluated | Main economic findings |
|--------------------------------------|-----------------------------|---|---|---|--|--|--|---|--|
| Karnon (2008) ²⁷⁸ U.K. | Cost-utility analysis | To estimate the net benefits of interventions that aim to reduce the impact of medication errors, either through prevention or detection. | A decision tree model to describe a series of error points and subsequent error detection points in pathways through the medication process in a generic secondary care setting. Assumed an acute hospital size of 400 beds | populated model with quantitative estimates to describe the incidence and impacts of medication errors. The effective-ness of potential interventions was estimated by describing impact of interventions on error incidence and detection rates, which feed through to alter the estimated frequency of medication errors and pADEs. | Five years to represent the predicted useful life of the IT-based interventions. | U.K. sterling (2006) Monetary values were assigned to the interventions, efficiency savings, treatment of, and the health effects of pADEs. | Quality of life utility decrements associated with experiencing a pADE | CPOE vs. additional ward pharmacists vs. bar coding | The fully estimated net benefits of the three interventions are dominated by the estimated monetary valuations of the health effects of pADEs, with mean net benefits of £31.5, £27.25, and £13.1 million over a five year time horizon for CPOE, ward pharmacists and bar coding, respectively. |

Evidence Table 8a. Summary of full economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Perspective (Time horizon) | Currency (year) Cost elements | Effective-ness measure | Intervention and alternative being evaluated | Main economic findings |
|--|-----------------------------|---|---|---|--|---|--|---|--|
| Plaza, V. (2005) ²⁷⁹ Spain | Cost-effectiveness | To evaluate the cost-effectiveness of an intervention to promote the recommendation of the Global Initiative for Asthma compared with standard practice | Physicians were randomized to CDSS offering recommendations or no CDSS groups in a multicenter, prospective, pragmatic study. Eligible patients were followed for 1 year. The incremental cost-effectiveness ratio was defined as the increase in total cost per patient divided by the change in QoL score | 20 physicians (10 pulmonologists and 10 primary care physicians) & (included 198 asthmatic patients) | Societal perspective & national health system (i.e. payer) | Euros (2001) Direct (resource x unit cost, treatment costs) and indirect (time off work due to medical visits) costs for societal perspective and direct costs for payer perspective | Difference in QOL using St. Georges Respiratory Questionnaire. GRQ, healthcare resources consumed, number of medical visits, hospitalizations, asthma treatment, blood analysis, spirometry, chest radiographs | CDSS vs. no CDSS | Not clear what currency the results are presented in. Hard to decipher but it may be that from the societal perspective the intervention was dominant (less costly and more effective) and from the payer perspective the ICER was \$61/percentage point reduction in SGRQ scale |
| Rosser (1992) ¹³⁹ Canada | Cost-effectiveness analysis | To assess the effect of three computerized reminder systems on compliance with tetanus vaccination. | Prospective randomized controlled trial (4 arms). Setting: Hospital Family Medicine Centre over 1 year | 5242 randomization patients and 2369 non-randomized patients ≥ 20 years of age not in a hospital or institution | Health care practice (1 year) | CDN (1985/1986) Physician time, clerical and nurse time, stationary, stamps, prepaid envelope and clerical time, cost to set up computerized reminder system was not included. | Proportion of patients who received tetanus toxoid in the study year or who had a claim of vaccination in the previous 10 years | Computer-generated physician reminder, vs. telephone reminder to patient, vs. letter reminder to patient to recommend tetanus vaccination vs. control group | cost to practice per additional vaccination recorded was 22¢ to 43¢ for physician reminders, \$4.43 to \$5.43 for telephone reminders; and \$6.05 for the letter reminders. |

Evidence Table 8a. Summary of full economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Perspective (Time horizon) | Currency (year) Cost elements | Effectiveness measure | Intervention and alternative being evaluated | Main economic findings |
|---|-----------------------------|--|--|----------------|--|---|--|---|---|
| Wu, RC. (2007) ²⁸⁰ Canada | Cost-effectiveness analysis | To determine the potential incremental cost-effectiveness of an electronic MOE/MAR system. | Incremental cost-effectiveness analysis comparing mean effectiveness of electronic MOE/MAR vs. Standard paper ordering for prevention of ADEs. Setting: Three tertiary care teaching hospitals | N/A | Health care institution (10 years with 5% discount rate) | USD (2004) Implementation costs (software, project management, clinical team involvement and training); operating costs (support for new interface, training)) | Reduction of preventable ADEs and mortality (rates obtained by review of literature) | MOE/MAR (i.e. CPOE) compared with conventional paper-based system | Incremental costs for CPOE system vs. paper was \$12,700 per ADE averted. This value is sensitive to the ADE rate, system effectiveness of ADE reduction, system cost, and costs due to possible increase in doctor workload. |

Evidence Table 8b. Summary of partial economic evaluation studies

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-----------------------------------|-----------------------------|---|--|--|---|--|---|---|
| Chertow (2001) ¹⁸ U.S. | Cost-analysis | To determine if a system application for adjusting drug dose and frequency in patients with renal insufficiency, when merged with a computerized order entry system, improves drug prescribing and patient outcomes | Four consecutive 2-month intervals consisting of control (usual computerized order entry) alternating with intervention (computerized order entry plus decision support system) conducted in September 1997–April 1998 at a 720-bed urban tertiary care teaching hospital. | Hospitalized patients with renal insufficiency, 7,887 admissions during the 2 intervention periods (2 months each) 9,941 admissions in the 2 control periods (2 months each) | ?? assumed 1997/1998 Hospital and pharmacy charges | Rates of appropriate prescription by dose and frequency, length of stay, and changes in renal function, compared among patients with renal insufficiency | Real-time computerized decision support system for prescribing drugs in patients with renal insufficiency. During intervention periods, the adjusted dose list, default dose amount, and default frequency were displayed to the order-entry user and a notation was provided that adjustments had been made based on renal insufficiency. During control periods, these recommended adjustments were not revealed to the order-entry user, and the unadjusted parameters were displayed. | There were no significant differences in estimated hospital and pharmacy costs. USD\$4,881 vs. USD\$4,968 in total costs and USD\$168 vs. USD\$166 for the intervention and the control groups, respectively LOS was shorter for the intervention group 4.3 days vs. 4.5 days, p = 0.009 even after adjusting for sex, age and DRG there remained a significant difference p = 0.002. |

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations = ADE = adverse drug event; AMI = acute myocardial infarction; CC = care considerations; CDSS=computerized decision support system; CHF = congestive heart failure; CMS = Center for Medicare and Medicaid Services; COPD = Chronic Obstructive Pulmonary Disease; CPOE=computerized physician order entry; CPR computer-based patient record; CVR = cardiovascular risk; DRG = diagnosis related group; EMR = electronic medical record; ER = emergency room; ESCHM = European Society of Cardiology and other societies for Hypercholesterolemia Management; HF=heart failure; HMO = health maintenance organization; ICU=intensive care unit; IHD=ischemic heart disease; IQR = *interquartile range* JCAHO = Joint Commission for Accreditation of Healthcare Organizations; LOS = length of stay; pmpm = per member per month; POE = physician order entry; QALY = Quality Adjusted Life Year; QOL = quality of life; RCT = randomized controlled trial; Rx = treatment; SD = standard deviation; USD = United States Dollars

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|--|-----------------------------|--|--|---|---|---------------------------|---|--|
| Barenfanger (2001) ²⁸⁷ U.S. | Cost-analysis | To assess the impact of improved interventions facilitated by (i) a computer software program which electronically notifies pharmacists of potential problems with a patient's antimicrobial therapy, and (ii) the education of pharmacists making interventions and notification of the medical staff of the program. | Quasi RCT (2 arm) of hospitalized patients prospective study in a 450-bed community teaching hospital over a 5 month time period | Patients: (i) infected with a bacterial isolate with no order for antimicro-bial therapy, (ii) infected with bacteria resistant to current antimicrobial therapy, (iii) on therapy not tested, and (iv) on antimicro-bial therapy but from whom no sample for culture had been taken. Analysis A: 24 patients in control group, 52 patients study group; Analysis B&C: (DRG) matched samples study group: 188, control group: 190 | ?? assumed 1998/1999 Total costs, fixed costs (overhead) variable direct (pharmacy costs, supplies, lab tests, radiology tests) fixed indirect costs | Mortality, length of stay | Compared patients whose microbiologic data were processed in the normal manual manner in the pharmacy to patients whose microbiological data were processed using the computer software, TheraTrac 2, a computer software program which electronically links susceptibility testing results immediately to the pharmacy and alerts pharmacists of potential interventions | Analysis A: study group had average total standard cost of \$21,189 per patient; control group had average total standard cost of \$51,790 per patient, a decrease of \$30,601 per patient in study group (p = 0.41) Analysis B (DRG-matched patients for whom susceptibility testing was done): study group had average total standard cost of \$13,294 per patient; control group had average total standard cost of \$18,601 per patient, a decrease of \$5,308 per patient in study group (p = 0.008). Analysis C:(severity adjustment) study group had average total standard cost of \$13,294 per patient; severity-adjusted control group had average total standard cost of \$16,106 per patient, a decrease of \$2,812 per patient in study group (no statistical analyses performed) By using these severity-adjusted data (that the data management team relies on), estimated variable cost savings annually from the improvement of interventions is \$2,932,000 (2,000 in patients for whom |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evalua- tion | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-----------------------------|--|--------------------|-----------------------------------|-------------------|----------------------------------|-------------------|--|---|
| | | | | | | | | <p>susceptibility testing is done X \$1,466). If the list price of TheraTrac 2 (\$44,500) is subtracted from the expected annual cost savings from the use of our program to improve interventions (\$2,932,000), the resulting savings (\$2,887,500) is still substantial in the first year. The present study demonstrates the financial benefits of improved interventions involving antimicrobial agents, namely, statistically significant differences in lengths of stay, total costs, variable costs, and radiology costs.</p> |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-----------------------------------|-----------------------------|---|--|--|---|---|---|--|
| Chisolm (2006) ¹⁹ U.S. | Cost analysis | To assess the relationship between use of a computerized order set within a CPOE and processes of care pediatric asthma treatment | Before/after. 'Pre-set' patients: those admitted prior to order set implementation; 'no set': those admitted after implementation when asthma order set not used; 'set' patients admitted after implementation and the order set was used. Inpatient pediatric teaching hospital | Asthma patients between the age of 2 and 20 years admitted to hospital between November 2001 and November 2003 (excluded those admitted to ICU). N=790 (261 'pre-set'; 63 'no set'; 466 'set' cases) | USD (year not stated) Length of stay, total inpatient charges and pharmacy charges | Use of systemic corticosteroids, use of pulse oximetry, and use of metered-dose inhalers. | Computerized order set within a CPOE system before and after implementation of the asthma order set | No significant difference in costs or lengths of stay among the three groups. Total charges were \$3,620, \$3,567, \$3,759; pharmacy charges were \$416, \$373, \$429; and LOS was 1.94, 1.93 and 1.77 for the 'no set', 'pre-set', and 'set' groups respectively. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-------------------------------------|-----------------------------|--|--|--|---|---|--|--|
| Cobos (2005) ²³ Spain | cost analysis | To assess the cost and effectiveness of a CDSS based on recommendations of the ESCHM in comparison with usual care for patients with hypercholesterolaemia | A multi-centre cluster-randomized, unblinded, pragmatic trial. (Primary care) Perspective not stated (1 year time horizon) | Patients with hypercholesterolaemia, which was defined as total cholesterol concentrations >200 mg/dL. Patients were excluded if they had triglyceride concentrations >400 mg/dL or were participating in another study. 44 practices, 2,221 patients (1,161 usual care, 1,060 CDSS) | Euros (2002) Direct costs only: physician visits, lab analyses, lipid-lowering drugs | Achievement of LDL-C reduction goals in patients with CVR of >20% over 10 yrs or keeping it <20% when patient baseline was <20% | CDSS vs. usual care | The treatment costs were €214,683 in the usual care group and €125,569 in the intervention group. The total costs were €264,658 in the usual care group and €170,061 in the intervention group. The adjusted means of the treatment costs per patient were €237 in the usual care group and €178 in the intervention group. The difference was €59 (95% CI: €34 to €83; p <0.0001). The adjusted means of the total costs per patient were €283 in the usual care group and €223 in the intervention group. The difference was €60 (95% CI: €33 to €86; p = 0.001). The CDSS did not alter the effectiveness of usual care but induced considerable savings. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|----------------------------------|-----------------------------|--|---|--|--|---|--|---|
| Evans (1992) ²⁸³ U.S. | quasi cost analysis | To use a hospital information system to help identify ADEs and to create a database of ADEs to prevent specific types of ADE | Pre-post design using a computerized ADE surveillance system vs. a control group with no ADEs | Hospital-ized patients | USD (not stated) Hospitalization costs | Reduction in ADEs and LOS | Computerized surveillance with physician notification only of verified ADEs if classified as severe or life-threatening vs. physician immediately notified of all ADEs when they were verified. Either the clinical pharmacist or ADE study nurse contacted the prescribing physician and recommended a change in drug or dosage vs. a control population of patients who received drugs but did not have ADEs | The average cost of hospitalization was \$38,007 for patients with severe ADEs compared to \$22,474 (p <0.002) for patients with moderate ADEs and \$6,320 for patients without ADEs. |
| Evans (1998) ³⁵ U.S. | Cost analysis | To evaluate a CDSS to improve the use of and reduce the cost of antibiotics | Prospective study in a 12 bed Shock/Trauma/Respiratory ICU. (before/after) 12 months | 398 patients in intervention (divided into those who got the recommended treatment and those who did not); 766 patients in control. # of physicians not stated | USD (1995) Cost of antibiotics and cost of hospitalization | # of ADEs, # of days of excessive antibiotic dosage, LOS, and mortality | Antibiotics ordered using CDSS by physicians during the study period compared to the control period | The cost of anti-infective agents was \$102 vs. \$340 and \$427 (p <0.001) and the total cost of hospitalization was \$26,315 vs. \$35,283 and \$44,865, (p <0.001). for control, regimen followed, and regimen overridden, respectively. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|----------------------------------|-----------------------------|--|---|--|-----------------------------------|---|--|--|
| Evans (1999) ²¹⁴ U.S. | Cost analysis | To examine the effect of a computer-assisted antibiotic dose monitor used to reduce the number of days that patients receive excessive dosages of antibiotics and the number of ADEs secondary to antibiotics. | Descriptive epidemiologic study of a two-year preintervention period and one-year intervention period. 12 month intervention period | All patients ≥18 years, admitted to Hospital from April 1 1993 to March 31 1996, who received ≥1 of 5 targeted antibiotics who had a serum creatinine or a urine creatinine clearance test result before antibiotic therapy, and who were never admitted or transferred to the ICU. # of physicians not stated | USD (1996) Cost of antibiotics | # of ADEs, # of days of excessive antibiotic dosage | Antibiotics ordered using CDSS by physicians during the study period compared to the control period | The intervention group had a lower and at a lower mean cost (\$80.62 vs. \$92.96; p <0.02) of antibiotics than patients during the preintervention period. |
| Evans (1994) ³⁷ U.S. | Cost analysis | To evaluate a CDSS to assist physicians in the selection of appropriate empiric antibiotics | Two-stage random-selection study (tertiary, private hospital and major teaching centre associated with a university). 12 month time frame | 28 physicians, 482 cultures | USD (1994) Cost of antibiotics | Computer-suggested antibiotics with results of susceptibility tests of cultures and antibiotics selected by physician | Antibiotics ordered using CDSS by randomized physicians were then compared between crossover periods of antibiotic consultant use. | The average cost for 24 hours of therapy for the computer-suggested antibiotics was \$41.08 per patient, compared with an average of \$51.93 (p <0.001) for the antibiotics actually prescribed by physicians. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-----------------------------------|-----------------------------|--|---|---|---|--|---|--|
| Evans (1995) ²⁸⁴ U.S. | Cost analysis | To evaluate a CDSS to improve the use of and reduce the cost of antibiotics | A 7-month pilot was compared with 12-months previous in a 12-bed Shock/Trauma/Respiratory ICU. 7 months | 588 orders for antibiotics, # of physicians not stated | USD (1994) Cost of antibiotics | # of ADEs and LOS | Antibiotics ordered using CDSS by physicians during the study period compared to the control period | The mean cost of antibiotics was \$87.03 (p <0.04) less per patient during the study period as compared to the control period. |
| Javitt (2005) ²¹⁸ U.S. | Cost analysis | To demonstrate the potential effect of deploying a sentinel system that scans administrative claims information and clinical data to detect and mitigate errors in care and deviations from best medical practices | RCT, members of an HMO were randomly assigned to an intervention or a control group. Care considerations (CC) generated by the CDSS for subjects in the intervention group were relayed to treating physicians, and those for the control group were deferred to study end. | Intervention and control group members consisted of all health plan enrollees who were between the ages of 12 and 64 years and had incurred at least 1 physician claim or 1 pharmacy claim in the 12 months before enrollment | USD (not specified) Total charges, in-patient charges; out-patient charges; Rx charges; professional charges | CCs generated by group; physician compliance with recommendation; and hospital utilization | CDSS tool that produces an electronic record from administrative data and runs it through a set of decision rules identifies "issues" and sends a CC message to the physician in the intervention group but nothing sent to the control group until the end of the study. | Charges for those whose recommendations were communicated were \$77.91 per member per month (pmpm) lower and paid claims were \$68.08 pmpm lower than controls compared with the baseline (p = 0.003 for both). Paid claims for the entire intervention group (with or without recommendations) were \$8.07 pmpm lower than those for the entire control group. In contrast, the intervention cost \$1.00 pmpm, suggesting an 8-fold return on investment. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|----------------------------------|-----------------------------|---|---|---|--|-------------------------------|--|---|
| Javitt (2008) ⁷⁴ U.S. | Cost analysis | To determine whether a CDSS tool improves quality of care and the effect of the intervention on average charges per member per month. | RCT, participants randomized to study group had the software turned on. Software was not turned on for patients in the control group until the 1-year experiment was over. Conducted in a large HMO | Patients all had medical charges in the previous year, all patients <65 yrs n = 19,719 intervention group, n = 19,792 control group | USD (2001) Total charges, in-patient charges; out-patient charges; Rx charges; professional charges | Rate at which CC are resolved | CDSS tool that produces an electronic record from billing records, lab feeds and pharmacies then runs the record through a set of decision rules, identifies "issues" and sends a CC message, 3 levels of CCs; level 1 contains potentially life-threatening situations, level 2 might have an important effect on clinical outcomes, level 3 are preventative care issues. All CCs reviewed by doctors employed by software company. HMOs medical director received level 1 messages and called the appropriate physician. Level 2 and 3 were received by an HMO nurse who then decided whether to send message to physician. Data collected in control group but CCs turned off. | The intervention reduced the average of total charges in the study group by 6.1%; average charge for the control group (\$327.54 vs. \$352.31 pmpm) |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|------------------------------------|------------------------------------|---|---|--|---|--|---|---|
| Kaushal (2006) ²⁸⁶ U.S. | Cost analysis | To assess the costs and benefits associated with the implementation of a CPOE and CDSS system over 10 years (1993-2002) | Cost and benefit estimates of a hospital CPOE system in a 720-adult bed, tertiary care academic hospital. With 7% discounting | Patients admitted to the hospital over the 10 year timeframe | USD (2002) Capital and operational costs, drug costs, hospital costs | Reductions in ADEs, LOS, proportion of appropriate prescriptions, laboratory & radiology tests (some measures from the literature) | CPOE with graduated CDSS over 10 years compared to estimates of what it might have been like without the CPOE | \$11.8 million to develop, implement, and operate CPOE; over 10 yrs, the system saved the hospital \$28.5 million. It took over 5 years to realize a net benefit and over 7 years to realize an operating budget benefit. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|---|-----------------------------|---|---|--|--|---|---|---|
| Macdonald (2002) ²⁸⁵ Canada | Cost analysis | Evaluation of the safety and potential cost savings of a computerized, laboratory-based program to manage inpatient warfarin thromboprophylaxis after major joint arthroplasty. | A consecutive-case study of adults admitted over a 54-month period (July 1994–December 1998) in a tertiary care orthopedic institution compared with Patients who underwent similar procedures in the 18-month period before the program was introduced (<1994) served as historical controls. These patients received the identical loading doses of warfarin and were individually managed by staff surgeons or internists. | Patients requiring joint arthroplasty who had no recent episodes of thromboembolic disease, no mechanical heart valve, atrial fibrillation, severe liver disease or baseline inter-national normalized ratio [INR] greater than 1.3 (n = 4,729, intervention vs. n = 279, control) | CAD (year not stated) Pharmacy and comparative nursing care costs associated with the program | Test results maintained within the desired therapeutic range (INR 2.0–3.0), clinically severe bleeding episodes, readmission rates, clinically symptomatic and venographically proven episodes of venous thrombosis or pulmonary embolism | Major joint arthroplasty with warfarin therapy administered through the computerized program compared with an historical control group Patients who underwent similar procedures in the 18-month period before the program was introduced served as historical controls. These patients received the identical loading doses of warfarin and were individually managed by staff surgeons or internists. | The potential savings per patient would be 11 minutes of nursing time or \$5.50/patient daily for a total annual figure, based on 10,152 patient days per yr of \$55,836. NOTE: The cost estimates and potential cost savings are speculative and are meant to be illustrative and not conclusive in nature. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-------------------------------------|-----------------------------|--|--|--|--|--|---|---|
| McGregor (2006) ¹⁰⁴ U.S. | Cost-analysis | To evaluate the effectiveness and cost effectiveness of a web-based, computerized CDSS for the management of antimicrobial utilization | RCT (2 arm), in-patients a 648-bed tertiary care, academic hospital over a 3 month period | n = 4,507 (n = 2,237 intervention arm & n = 2,270 control) | USD (2004) Hospital antimicrobial costs (primary outcome) | Mortality, LOS, frequency of tests for C. difficile, time spent managing antimicrobial utilization | Antimicrobial utilization was managed by an existing antimicrobial management team (AMT) using the system in the intervention arm and without the system in the control arm. The system was developed to alert the AMT of potentially inadequate antimicrobial therapy. This is a "back-end" or post-prescription review. | Hospital antimicrobial expenditures were \$285,812 for intervention vs. \$370,006 in the control arm, for a savings of \$84,194 (23%), or \$37.64 per patient |
| McMullin (2004) ²⁸¹ U.S. | Cost analysis | To evaluate the impact on prescription costs of a computerized decision support system (CDSS) | Retrospective cohort study (before-after) using pharmacy claims database in primary care. Clinicians using CDSS were matched to controls with 6 month followup | 19 physicians in each group | USD (not stated) New prescription costs | NIL | CDSS that provides evidence-based recommendations to clinicians during the electronic prescribing process before and after implementation | Average cost for intervention group per new prescription \$4.16 lower (p = 0.02); for new and refilled prescriptions \$4.99 lower (p = 0.01). The 6 month savings from new prescriptions and their refills were estimated to be \$3,450 (95% CI, \$1,030 to \$5,863) per clinician. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-------------------------------------|-----------------------------|---|---|-----------------------------------|---|---|--|--|
| McMullin (2005) ²⁸² U.S. | Cost analysis | To evaluate the impact on prescription costs of a computerized decision support system (CDSS) | Retrospective cohort (before-after) study using pharmacy claims database in primary care. Clinicians using CDSS matched to controls. 12 months (extension of the 6-month study described above) | 19 physicians in each group | USD (not stated) New and existing prescription costs | NIL | CDSS that provides evidence-based recommendations to clinicians during the electronic prescribing process before and after implementation | The average cost per new prescription decreased by \$1.00 (-2.4%) in the intervention group while it increased by \$3.75 (9.0%) in the control group. The 12 month savings on new prescriptions were \$109,897 |
| Mekhjian (2002) ¹⁸⁶ U.S. | Cost-analysis | To evaluate the benefits of computerized physician order entry (POE) and electronic medication administration record (e-MAR) on the delivery of health care | Cohort of inpatient nursing units in an academic health system (3 sites Cancer hospital, state hospital, rehab centre), before-and-after POE The study comprised before-and-after comparisons between phase 1, preimplementation of POE (pre-POE) and phase 2, postimplementation of POE (post-POE) and, within phase 2, a comparison of POE and the combination of POE plus e-MAR | Cohort of inpatient nursing units | USD (2002) Total costs per patient | LOS, medication turn-around time, radiology turn-around time, laboratory test turn-around time, medication transcription errors | phase 1, preimplementation of POE (pre-POE) and phase 2, postimplementation of POE (post-POE) and, within phase 2, a comparison of POE and the combination of POE plus e-MAR | State hospital total costs for the heart transplant service (pre-POE, \$5,264; post-POE, \$4,871; p = 0.013) and organ transplant service (pre-POE, \$8,382; post-POE, \$7,711; p = 0.043) showed a statistically significant decrease, whereas costs for general surgery (pre-POE, \$4,995; post-POE, \$5,567; p = 0.008) showed a statistically significant increase. There were no statistically significant changes in other services. Cancer: services such as surgical oncology (pre-POE, \$6,087; post-POE, \$5,631; p = 0.008) and neurology/ neurosurgery (pre-POE, \$5,600; post-POE, \$5,125; p = 0.045) showed statistically |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|------------------------------------|-----------------------------|--|--|--|---|---|---|---|
| | | | for a period of 10 to 12 months across all services in the respective hospitals. | | | | | significant reductions in total costs, whereas the POE, \$5,821; p <0.001) showed a statistically significant increase in total costs and thoracic surgery (pre=POE, \$5,181; post=POE, \$5,946; p = 0.055) showed a nonsignificant increase. When all the services were combined, severity adjusted total cost per admission did not change significantly in either state (pre-POE, \$5,697; post-POE, \$5,661; p = 0.687) or in the cancer hospital (pre-POE, \$6,427; post-POE, \$6,518; p = 0.502). |
| Mullett (2001) ¹⁰⁹ U.S. | Cost-analysis | To evaluate the impact of a pediatric anti-infective CDSS. | Cohort, patients in a 26-bed pediatric intensive care unit in an academic 232-bed hospital 6-month pre- vs. postimplementation | N=1758 (809 control, 949 intervention) | USD (1999) Hospital costs, anti-infective drug charges | Number of anti-infective drugs used, total doses used, LOS, mortality | CDSS vs. pre-CDSS (all patient care orders from the physicians were handwritten. Antibiotic and other medication orders typically were interpreted by the clerk and rewritten onto the bedside medication administration record. Carbon copies of the handwritten order were physically sent to the pharmacy and read by a pharmacist, who entered the order via the keyboard into the HELP system's pharmacy module. | no difference in hospital costs \$28,257.67 (control) vs. \$25,032.11 (intervention) no difference in mean anti-infective cost/patient \$274.79 (control) vs. \$289.60 (intervention) |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-------------------------------------|-----------------------------|---|--|-----------------------------|---------------------------------------|----------------|--|---|
| Ornstein (1999) ²⁸⁸ U.S. | Cost-analysis | To determine the impact of displaying prescription cost information in a computer-based patient record (CPR) system on decreasing drug costs by family physicians | During a 6-month period, cost information was not displayed; during the subsequent 6-month intervention period, costs were displayed at the time of prescribing. Academic family practice setting. | 10 physicians, 36 residents | USD (1995/1996) Prescription costs | Nil | CPR system that displays drug cost information at time of prescription order compared to no cost information being displayed during the control period | This study failed to detect an impact of CPR-based prescription drug cost information on overall drug costs to patients among family physicians in an academic family medicine ambulatory clinical practice. The mean (SD) cost per prescription in the control period was \$21.83 (\$27.00), and in the intervention period was \$22.03 (\$28.12), (p = 0.61). The mean (SD) cost/contact control \$12.49 (\$29.35) vs. intervention \$13.03 (\$30.06) (p = 0.12). |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|--|-----------------------------|--|--|--|---|--|---|---|
| Paul (2006) ¹²³ Israel, Germany, Italy | Cost analysis | Cohort study: the aim was to compare CDSS advice with physician performance for antibiotic treatment and antibiotic costs. In the RCT, the goal was to assess whether the CDSS improved physician performance and patient-related outcomes | Prospective cohort study comparing a CDSS for antibiotic treatment advice to physician's treatment followed by a multicentre, cluster randomized trial comparing wards using the CDSS vs. antibiotic monitoring without the CDSS. (Cohort-6-month time period between 2002/2003 in each of the 3 countries; RCT-6-month period in 2004) 3 university affiliated primary and tertiary hospitals (Israel, Germany, Italy) | Patients suspected of harboring bacterial infections in 3 university affiliated primary and tertiary hospitals (Israel, Germany, Italy) Cohort: 1,203 patients RCT: 2,326 patients | Euros (2002/ 2004) Antibiotic costs including: (1) direct drug & administration, (2) ADE (rates from the literature and assigned costs in hospital days and QALYs), (3) ecological costs (patient costs, probability of infection and antibiotic failure; costs to eco-system for loss of antibiotic efficacy, penalty cost for drugs of last resort (antibiotic costs, including costs related to future antibiotic resistance) | Appropriate antibiotic treatment, mortality, LOS | CDSS recommends treatment by highlighting the 3 top-ranked antibiotic regimens, with the highest cost-benefit difference, including no antibiotic treatment wards using the CDSS vs. antibiotic monitoring without CDSS). CDSS advises antibiotic therapy for inpatients using data available at the time of empirical antibiotic treatment. | COHORT: All cost components, except those related to expected adverse events, were significantly lower for the treatments suggested by the CDSS compared with those used by physicians. Total antibiotic costs were €289 lower per patient for CDSS compared with physicians, a relative decrease of 48%. RCT: the use of the CDSS resulted in significantly lower antibiotic costs in intervention vs. control wards, the difference originating from lower ecological costs in intervention wards in Israel and Italy. Direct antibiotic costs, as well as costs incurred by observed adverse events, were similar -mean total antibiotic costs per patient €623.2 (control) vs. €565.4 (intervention) p = 0.007 Total projected costs for the appropriate CDSS regimens were lower than physician's treatment by €262 per patient, a relative decrease of 44%, with the reduction originating mainly from lower ecological costs |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|------------------------------------|-----------------------------|--|--|---|---|---|---|---|
| Piontek (2010) ²⁹⁰ U.S. | Cost analysis | The effects of an adverse-ADE alert system on cost and quality outcomes in community hospitals were evaluated. | Retrospective observational study evaluated the effects of an ADE alert system in seven hospitals in a health network. Outcomes after, and one year before, the deployment of an ADE alert system were evaluated. Inpatients in two hospitals without any computerized ADE alert system constituted the control group. Administrative data were gathered for patients from these facilities for the same time frames as for the pre-implementation and post-implementation groups. | All inpatients admitted to one of seven hospitals in a health network | USD (not indicated) Primary outcomes evaluated included pharmacy department costs, variable drug costs. Secondary outcomes included total hospitalization costs | Primary outcomes included mortality rates. Secondary outcomes included LOS, rate of readmission, and case-mix index | Pre-post ADE alert system. Four distinct groups were evaluated: (1) preimplementation of the ADE alert system (internal control group), (2) postimplementation group, (3) external control group matching internal control time frame, and (4) external control group matching ADE postimplementation time frame. | Statistically significant decreases were observed in average pharmacy department costs per patient (\$867 vs. \$826, p <0.001) from preimplementation to post-implementation. In contrast, the external control group had a significant increase in pharmacy department costs (\$734 vs. \$797, p = 0.029). Drug costs decreased significantly from baseline (\$360 vs. \$337, p <0.001) in the study group. Conversely, there were significant increases in drug costs in the external control group (\$401 vs. \$429, p = 0.029). |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|----------------------------------|-----------------------------|--|--|---|--|--|--|--|
| Stone (2009) ¹⁵⁸ U.S. | Cost-analysis | Presentation of the implementation of a CPOE system in the management of surgical patients | Retrospective and prospective analyses of patient-safety measures 6 months pre- and 6 months post-CPOE institution, respectively. Inpatients of a multispecialty hospital academic surgical practice | Paper only provides the number of surgical procedures pre and post: 6,815 procedures in the pre period and 5,963 in the first post 6 month and 6,106 in the second 6 months postimple-mentation | USD (2007/2008) Personnel requirements (efficiencies) and capital costs of implementation | Patient safety, medication errors, order implementation time | CPOE compared to no CPOE | -total capital costs for implementation \$2.9 million and operating costs of \$2.3 million -decrease in the number of unit secretaries (clarified orders and transcribed the orders to a required format); -savings of \$445,500 (personnel changes occurred as a consequence of work-load redistribution). Considerable gains in efficiency were noted, which included the time necessary to have orders accessible to nursing, radiology, and laboratory. This gain in efficiency will likely result in long-term cost savings and increased quality of care. Additionally, personnel needs were reduced, which subsequently resulted in additional financial benefit for our institution. |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|------------------------------------|-----------------------------|--|--|---|---|---|--|---|
| Tierney (2005) ¹⁶⁷ U.S. | Cost analysis | To assess whether guideline-based care suggestions delivered via physicians' and pharmacists' computer (CDSS) workstations could improve the outpatient management and outcomes among patients with asthma or COPD | 1-year, 2x2 factorial, 4 arm RCT, academic general internal medical practice in a hospital | 246 physicians and 20 outpatient pharmacists randomized (706 patients included) | USD (1994-1996) Total health care charges (Outpatient charges + inpatient charges) | Adherence to treatment guidelines, QOL, patient satisfaction with physician & pharmacists, ER visits, hospitalizations. | Care recommendations provided electronically to physicians, pharmacists, both physician & pharmacist vs. no care recommendations/ intervention | Patients in the group receiving only the physician intervention had significantly elevated total health care charges, possibly because of just a small number of extremely high-cost hospitalizations costs Control (n=80):\$5,800 (SD: 8,536) Physician only (n=81):\$8,006 (SD: \$18,720) Pharmacist only (n=80):\$5,333 (SD:\$9,400) Both physician & pharmacist (n=82):\$5,652 (SD: \$10,579) |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|------------------------------------|-----------------------------|--|--|--|---|---|---|---|
| Tierney (2003) ¹⁶⁶ U.S. | Cost analysis | To assess the effects of an established EMR system containing a CPOE with a guideline-based CDSS for managing patients with IHD and chronic HF | 1-year, 2x2 factorial, 4 arm RCT, academic, primary care group practice (targeting physicians and pharmacists) | 11 full time & 9 part-time outpatient pharmacists; ?? physicians; 32 practice sessions (706 patients included) | USD (1994-1996) Total health care charges (Outpatient charges + inpatient charges) | Adherence to recommendation, health-related QOL, exacerbation of heart disease, patient satisfaction with physician and pharmacist, medication compliance, satisfaction with care, physician attitude toward intervention | Evidence-based cardiac care recommendations displayed electronically to physicians, pharmacists, physician & pharmacists vs. no recommendations for enrolled patients | No difference in total costs across groups Costs: Control (n=181): \$7,025 (SD \$17,024) Physician only (n=197): \$6,302 (SD 10,928) Pharmacist only (n=158): \$7,387 (SD: \$13,206) Both physician and pharmacist (n=170): \$7,639 (SD:\$16,921) |

Evidence Table 8b. Summary of partial economic evaluation studies (continued)

| Author (year) Country | Type of economic evaluation | Study objective | Study design (include setting) | Population (n) | Currency (year) Cost elements | Effect measure | Intervention and alternative being evaluated | Main economic findings |
|-------------------------------------|-----------------------------|--|--|--|--|---------------------------|--|--|
| Tierney (1993) ²⁸⁹ U.S. | Cost analysis | To assess the effects on health care resource utilization of a network of microcomputer workstations for writing all inpatient orders (CPOE) that encourage cost effective ordering. Aim of increasing cost consciousness and reducing costs | RCT in an inpatient internal medicine service of an urban public hospital over 6 months | 6 medical services were randomly assigned to intervention or control: 5,219 patients (1,859 intervention from 22 teams & 3,360 controls from 46 teams) | USD (1990/1991) Inpatient charges (bed, tests and drugs) | LOS, time in motion | Microcomputer workstations, linked to a comprehensive EMR system for all inpatient order vs. hand-written orders | Total charges per admission were significantly less (mean difference: \$887, 12.7% reduction) for intervention teams than for control teams, with similar differences in bed charges, test charges, and drug charges. Hospital stays for intervention admissions were 0.89 day (10.5%) shorter than for controls (p = 0.11). This would amount to more than \$3 million in charges annually for that hospital's medicine service |
| Weingart (2009) ²⁹¹ U.S. | Cost analysis | To understand the potential benefits of medication safety alerts in ambulatory care using a (CPOE) | A multifaceted study from January 1 through June 30 2006. An expert panel reviewed a sample of common drug interaction alerts, estimating the likelihood and severity of ADEs associated with each alert, the likely injury to the patient, and the health care utilization required to address each ADE | 279, 476 alerted prescriptions written by 2,321 ambulatory care clinicians | USD (2006) Hospitalization, emergency room visit, office visit, filled prescription | ADEs and related injuries | Potential benefit of electronic prescribing with decision support based on expert panel estimates | Alerts potentially resulted in a cost savings of \$402,619 (IQR, \$141,012 to \$1,012,386). Drug alerts have the potential to prevent harm and reduce health care costs |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|-------------|----------------------------------|--|----------------------------|--|---------|
| Ali (2005) ⁴ Design: Before-after N = 91 patients Implementation: 02/2000 Study Start: 05/2000 Study End: 05/2002 | Prescribing | CPOE/POE system | Critical care units (CCU, ICU, NICU) 25 Beds Academic | LOS - Secondary Outcome | LOS did not differ between patients cared for with the initial CPOE vs. the modified CPOE (9.9 days vs. 9.0 days, NS) | - |

The HIT system studied is in **bold**, followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: A1c = hemoglobin A1c; ADE = Adverse Drug Event; AHR = Airway Hyper-responsiveness; aPTT = Activated Partial Thromboplastin Time; AQLQ = Asthma Quality of Life Questionnaire; ARR = Adjusted Relative Risk; BMI = Body Mass Index; BG = blood glucose; BP = Blood Pressure; CAGES = computer assisted guideline enhancement system; CCDS = Computerized Clinical Decision Support; CDS = Clinical / Computerized Decision Support; CDSS = Clinical Decision Support System; CI = Confidence Interval; CPOE = Computerized Provider Order Entry; DBP = Diastolic Blood Pressure; DVT = Deep Vein Thrombosis; EHR = Electronic Health Record; e-MAR = Electronic Medication Administration Record; EMR = Electronic Medical Records; e-TAR = Electronic Treatment Authorization Request; FEV1 = Forced Expiratory Volume in the first second.; GHb = Glycohemoglobin; GP = General Practitioner; HbA1c = Glycated hemoglobin; HF = Hear Failure; HIT = Health Information Technology; ICU = Intensive Care Unit; INR = International Normalized Ratio; Kg/m² = Kilogram per square metre; LDL = Low density Lipoprotein; LOS = Length of Stay; mg/dL = milligrams per decilitre; micro-mol/L = micro moles per litre; mL/min = millilitre per minute; MM = Medication Management; MMC = Montefiore Medical Center; mmHg = millimeter of mercury; mmol/l = millimoles per litre; N or n = Sample Size; NS = Not Statistically Significant; OR = OR; OSUH = Ohio State University Health System; p = Probability; PADEs = Potential Adverse Drug Events; PANSS = Positive and Negative Syndrome Scale PCA = Patient-Controlled Analgesia; PE = Pulmonary Embolism; PHR = Patient Health Record; POE = Provider Order Entry; PRISM = Pediatric Risk of Mortality; QoL = Quality of Life; RCT = Randomized Controlled Trial; RRR = Relative Risk Reduction; SBP = Systolic Blood Pressure; SD = Standard Deviation; SF-36 = Short Form 36; SICU = Surgical Intensive Care Unit; vs. or vs. = Versus; VTE = Venous thromboembolism

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|---|---|---|---------|
| Balcezak (2000) ²⁹² Design: Cohort study N = 747 Patients Implementation: 00/0000 Study Start: 10/1996 Study End: 01/1997 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system | Acute care/tertiary, 900 Beds Academic | aPPT exceeding therapeutic threshold by 24 hours, aPPT exceeding therapeutic threshold by 48 hours, aPPT within therapeutic threshold by 48 hours | Use of the nomogram was 10% (low). More patients who received the nomogram ordering exceeding the therapeutic threshold by 24 hours and by 48 hours were 79% vs. 56% (p <0.001), and 88% and 66% (p <0.001) respectively. More patients achieved a therapeutic range by 24 hours and 48 hours with the weight-based nomogram compared with physician-guided dosing were 47% vs. 39% (p = 0.027), and 69% and 52% (p = 0.019) respectively. Use of the nomogram also had a higher rate of being within the therapeutic range by 48 hours (69% vs. 52%, RRR 25%, p = 0.02). | + |
| Barenfanger (2001) ²⁸⁷ Design: Cohort study N = 450 patients Implementation: 00/0000 Study Start: 10/1998 Study End: 02/1999 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system, Pharmacy | Acute care/tertiary, 450 Beds Pharmacy, Inpatient hospital based, Academic | mortality rate, average length of stay | In matched patient analysis, the study patients did not differ for mortality (10% vs. 11%, p = 0.7). The control group patients stayed longer in the hospital (13.7 vs. 11.0 days, p = 0.04). | + |
| Baroletti (2008) ²⁹³ Design: Cohort study N = 9,527 patients Implementation: 00/000 Study Start: 01/2004 Study End: 07/2006 | Prescribing | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system | Unspecified Hospital | symptomatic VTE or PE* | The primary end point of symptomatic DVT or PE at 90 days occurred in 5.1% of patients in the cohort group and 4.9% of patients in the historical alert group, respectively, p = 0.82 | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|--|--|---------|
| Beccaro (2006) ²⁹⁴ Design: Before-after N = 2,533 patients Implementation: 11/2003 Study Start: 10/2002 Study End: 12/2004 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Hospital information system, Imaging systems, Laboratory system | Pediatric stand alone hospital, 270 Beds | unadjusted total mortality | Introduction of the CPOE system was not associated with changes in mortality rates measured 13 months after implementation. The unadjusted mortality rates before implementation was 4.2% and after 3.4%, RRR 18%, (95% CI -21% to 45%), NS. No mortality difference was seen either for transfer patients (7.8% before and 9.6% after, RRR 34%, 95% CI -47% to 71%, NS) or for children with congenital cardiovascular disease (4.4% before and 2.6% after, RRR 41%, 95% CI -63% to 79%, NS). | - |
| Boord (2007) ²⁹⁵ Design: Before-after N = 351 ICU patients Implementation: 11/2004 Study Start: 10/2004 Study End: 01/2005 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders, CPOE/POE system, e-Rx Integrated EHR/EMR system | Acute care/tertiary, Critical care units (CCU, ICU, NICU) 21 Beds Academic | percentage of patients within ideal glucose range*, time spent in ideal glucose range (minutes)* | Patients were studied for 5 days in the SICU. The percentage of patients with their blood glucose in the ideal range increased with the CPOE insulin protocol (29.3% vs. 37.7%, RRR -29%, p = 0.006). Patients who were cared for under CPOE/CDSS also spent more time on average within normal glucose levels across all 5 days (mean difference 116 minutes, p = 0.029). | + |
| Chabot (2003) ²⁹⁶ Design: Cohort study N = 100 people with hypertension Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated Pharmacy Computerized Prescription Management System | Pharmacy, Other | Mean BP (mm Hg), Adherence, Controlled BP (Measure based on Recommendation before 1999) or in 1999 | The groups did not differ for blood pressure: mean SBP: 139 vs. 141, p = 0.747; DBP: 78 vs. 78, p = 0.357 or for adherence based on pharmacy recorded: 93% vs. 98%, p = 0.643 or self reported data: 83% vs. 68%, p = 0.085 or rates of controlled BP (recommendations before 1999) 81% vs. 78%, p = 0.684 (recommendations in 1999) 44% vs. 54%, p = 0.3 | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|--|--|---------|
| Chen (2010) ²⁹⁷ Design: Case control N = 200 patients Implementation: 00/0000 Study Start: 08/2003 Study End: 00/0000 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system | Unspecified Hospital | LDL-C goal, | Among the first 200 consecutive patients followed up more than 1 year, 65% reached the LDL-C goal in 1 year. For those whose treatment followed CAGES, 74% reached the LDL-C goal. For those whose treatment was entered without CAGES, 57% reached the LDL-C goal. The OR is 2.1 (1.2, 3.8, 95% CI) (p = 0.022), patients whose treatment followed CAGES were twice as likely to reduce their LDL-C | + |
| Chertow (2001) ¹⁸ Design: Time series N = 19,982 admissions Implementation: 00/0000 Study Start: 09/1997 Study End: 04/1998 | Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated Hospital information system Imaging systems | Acute care/tertiary, 720 Beds Academic | length of stay, percentage of patients with a decline of creatinine clearance >10 mL/min | Length of stay decreased with CPOE/CDSS (mean 4.5 days vs. 4.3 days, p = 0.009). No changes in renal function were observed | + |
| Chisholm (2003) ¹⁹ Design: Before-after N = 790 children admitted to hospital with asthma exacerbations Implementation: 10/2002 Study Start: 11/2001 Study End: 12/2003 | Monitoring including patient adherence and compliance, Prescribing | CPOE/POE system Integrated Billing/administration system, EHR/EMR system, Laboratory system | Pediatric stand alone hospital, 323 Beds | length of stay | No difference was seen in LOS (1.8 vs. 1.9 days) NS | - |
| Cobos (2005) ²³ Design: RCT N = 2,221 patients Implementation: 04/2000 Study Start: 04/2000 Study End: 05/2002 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | proportion of patients achieving successful lipoprotein-cholesterol goals or cardiovascular risk reassessment* | Effectiveness was defined as success or failure for patients achieving either their LDL cholesterol goal or a reassessment of their cardiovascular risk maintained at <20%. The proportion of patients achieving success in the intention to treat analysis was similar between usual care and intervention groups (50.5% vs. 54%, NS). | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|--|---|---------|
| Cook (2004) ²⁹⁸ Design: Time series N = 334 paired visits Implementation: 11/2001 Study Start: 11/2001 Study End: 05/2002 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Handheld | Ambulatory care | fasting glucose levels*, random glucose*, A1c* | For 117 paired visits using fasting glucose for insulin adjustment, paired fasting glucose levels decreased from 220 ± 85 to 149 ± 61 mg/dL (p <0.0001). For 103 paired visits where random glucose was used for dosing, random glucose decreased from 249 ± 93 to 168 ± 69 mg/dL (p <0.0001). For 114 paired visits using A1c for insulin adjustment, A1c levels improved from 10.4 ± 2.9% to 7.9 ± 2.0% (p <0.0001). | + |
| Evans (1995) ²⁸⁴ Design: Before-after N = 962 patients Implementation: 07/1994 Study Start: 07/1993 Study End: 02/1995 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated Hospital information system | Critical care units (CCU, ICU, NICU) | ADE rate, Length of Stay | The rate of ADE did not differ before and after implementation (2.4% vs. 0.,9%, NS). The length of stay in the unit did not differ (mean 6.2 vs. 5.8 days, NS). | - |
| Evans (1998) ³⁵ Design: Before-after N = 1,681 patients Implementation: 00/0000 Study Start: 07/1992 Study End: 06/1995 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Imaging systems, Laboratory system | Acute care/tertiary, 520 Beds Academic | ADE* | ADE rate decreased significantly following the implementation of the reminder (3.7% vs. 1%, RRR 73%, p = 0.018) | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|---|---|---|---|---|---------|
| Evans (1992) ²⁸³ Design: Before-after N = 45,544 patients (May '89-Apr '91) Implementation: 05/1989 Study Start: 05/1989 Study End: 00/1991 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/reminders surveillance system Integrated EHR/EMR system, Hospital information system, Laboratory system, Pharmacy | Unspecified Hospital | rate of type B ADEs, rate of severe ADEs, length of stay | Rate of type B ADEs declined following the use of the surveillance system and the addition of computer alerts (15% vs. 1.4%, p <0.001). The rate of severe ADEs with early notification of physicians to all ADEs as soon as they were verified reduced from 7.6% to 2.2% (p <0.001). The average length of stay for patients with type B ADEs to hospital-administered drugs was 17 days compared to 14 days (p <0.013) for patients with type A ADEs and only five days for the control patients that did not have ADEs. The average length of hospitalization for patients with severe ADEs was 20 days compared to 13 days for patients with moderate (p <0.024). | + |
| Evans (1994) ²⁹⁹ Design: Time series N = 1,865 ADEs Implementation: 00/1980s Study Start: 05/1989 Study End: 04/1992 | Monitoring including patient adherence and compliance | Health Information System Integrated Laboratory system, Pharmacy | Acute care/tertiary, 520 Beds Academic | ADE-known drug allergies*, ADE-rapid antibiotic administration rates* | The ADE surveillance system identified drug allergy and rapids antibiotic administration rates as areas of concern. In year 2 and 3, when clinicians were alerted to all ADEs and had in service for antibiotic admin rates, the number of known drug allergy ADEs declined to 0 (p <0.002) and there was a significant decrease in the number of ADEs related to antibiotic administration rates (p <0.01). | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|---|--|---|---------|
| Evans (1999) ²¹⁴ Design: Before-after N = 13,384 Patients Implementation: 01/1005 Study Start: 04/2005 Study End: 03/2006 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated Pharmacy | Critical care units (CCU, ICU, NICU), 12 beds in the shock/trauma/respir atory ICU. of a 520 bed academic hospital Beds Not specified, Inpatient hospital based, Academic | Rates of adverse drug events* | The rate of adverse drug events related to 5 antibiotics was lower in patients who were followed with the drug monitoring system (0.9% vs. 0.3%, RRR 67%, p <0.001). | + |
| Fiumara (2010) ³⁰⁰ Design: Cohort study N = 880 patients Implementation: 00/0000 Study Start: 07/2006 Study End: 05/2008 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system, Hospital information system | Unspecified Hospital | Symptomatic DVT or PE at 90 days*, PE at 90 days, DVT at 90 days, Death at 90 days, Major haemorrhage at 30 days | There was no significant difference in symptomatic 90-day VTE rates between the two cohorts (2.8% for the one-screen vs. 2.2% for the three-screen, p = 0.55). PE at 90 days was 1.1% vs. 0.9%, p = 0.25. DVT at 90 days was 1.1% vs. 1.9%, p = 0.14. Death at 90 days was less frequent among patients in the one-screen alert cohort than the three-screen alert cohort (14.6% vs. 22.2%, p = 0.004). The frequency of major haemorrhage was similar in both alert cohorts (1.3% vs. 1.8%, p = 0.51). | - |
| Frances (2001) ⁴⁸ Design: RCT N = 63 physicians and 730 patients Implementation: 00/0000 Study Start: 03/1997 Study End: 06/1997 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Pharmacy | Ambulatory care | LDL level <100 mg/dL* | The proportion of patients with a level of LDL cholesterol in the desired range (< 100 mg/dL) Did not improve cholesterol management in patients (73.2 % vs. 71.0%, p = 0.512) with CAD. | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|---|---------|
| Garthwaite (2004) ³⁰¹ Design: Cohort study N = 939 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Acute care/tertiary, Ambulatory care, Academic | Serum cholesterol, creatinine kinase, alkaline phosphatase levels, creatinine clearance, cyclosporine dose, cyclosporine trough levels, DBP, SBP | Serum cholesterol*, creatinine kinase, alkaline phosphatase levels*, creatinine clearance*, cyclosporine doses and trough levels, lipid-lowering drugs, and dia*- and SBP measurements were compared between baseline and 6 months. *indicates significant improvements. | + |
| Gill (2009) ⁶⁰ Design: RCT N = 64,150 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 10/2006 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/reminders Integrated EHR/EMR system | Ambulatory care | Lipids at goal* | After controlling for confounding variables and for clustering in multilevel modeling, the proportion of patients with lipids at goal was not significantly different between control and intervention groups. | - |
| Gilutz (2009) ⁶¹ Design: RCT N = 7,448 patients from 56 control and 56 intervention clinics Implementation: 00/0000 Study Start: 01/2000 Study End: 12/2003 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/reminders Integrated Hospital information system, Laboratory system, Pharmacy | Ambulatory care, Academic | LDL level reduction* | In the group of patients with initial LDL levels above 120 mg/dl, a significant decrease in LDL levels was observed in the two groups, which was minimally more pronounced in the intervention arm from 145.5 ± 22.3 mg/dl to 121.9 ± 34.2, mg/dl, 16.2% reduction than in the control arm from 145.8 ± 22.9 to 124.3 ± 34.6, 14.8% reduction; (p <0.02). | + |
| Grant (2008) ²¹⁷ Design: RCT N = 244 patients Implementation: 00/2002 Study Start: 09/2005 Study End: 03/2007 | Monitoring including patient adherence and compliance | PHR Integrated Billing/administration system, EHR/EMR system, Imaging systems, Laboratory system, Patient decision support system | General Hospital, Ambulatory care, Home | HbA1c levels* | For the primary outcome, study participants had relatively good glycemic control (mean HbA1c levels) at baseline with modest improvement over the study period that did not differ by treatment arm (7.1% vs. 7.2%, p = 0.45), with nearly three-quarters of all patients at goal (73% vs. 68% among control patients; p = 0.53). | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|-------------|--|--|--|---|---------|
| <p>Graumlich (2009)²³⁷ Graumlich (2009)²³⁸ Design: RCT N = 631 patients Implementation: 00/0000 Study Start: 11/2004 Study End: 01/2007</p> | Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system | Acute care/tertiary, 730 Beds Academic | Readmitted within 6 months*, emergency department visit within 6 months, adverse events within 1 month | When comparing patients assigned to discharge software vs. usual care, there was no difference in hospital readmission within 6 months (37.0% vs. 37.8%; OR 0.005 [95% CI, - 0.074 to 0.065]; p = 0.894), emergency department visit within 6 months (35.4% vs. 40.6%; OR 0.052 [95% CI, -0.115 to 0.011]; p = 0.108), or adverse events within 1 month (7.3% vs. 7.3%; OR 0.003 [95% CI, -0.037 to 0.043]; p = 0.884) | - |
| <p>Gurwitz (2008)³⁰² Design: RCT N = 1,118 residents Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system Laboratory system | Long term care (nursing homes) | ADE rates per 100 resident months*, Preventable ADE targeted by alerts | ADE rates per 100 resident months were similar for control and intervention units (10.4 vs. 10.8, NS). The same was found for the rate of preventable ADEs per 100 resident months (3.9 vs. 4.0, NS). Of the 152 preventable events on the intervention units, 59 (38.8%) might have been prevented as a result of one or more of the alerts. Of the 126 preventable events identified on the control units, 56 (44.4%) might have been prevented as a result of one or more of the alerts. NS. | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|---|--|---------|
| Han (2005) ³⁰³ Design: Before-after N = 1,942 patients Implementation: 10/2002 Study Start: 10/2001 Study End: 03/2003 | Dispensing, Transmission, order communication | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated Hospital information system | Acute care/tertiary, Pediatric stand alone hospital, Academic | mortality rate | The unadjusted mortality rate after implementation was 3.9%. A step-wise regression analysis was done using 19 variables. For the model that was adjusted for PRISM score 7 factors including use of CPOE were associated with mortality: shock, Glasgow Coma scale score, surgical referral, prematurity, cardiovascular problems, and PRISM score. The OR for mortality for the presence of CPOE is 3.71, 95% CI 2.13 to 6.46. Post CPOE mortality affected children and ICU admission most severely. In the primary regression model that adjusted for PRISM score, shock was highly associated with increased odds of mortality (OR: 6.24; 95% CI:2.94 to 13.26), followed by CPOE (OR: 3.71; 95% CI:2.13 to 6.46) and severe coma (OR: 3.43; 95% CI: 1.88 to 6.25). | + |
| Hetlevik (1999) ³⁰⁴ Design: RCT N = 1,998 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated EHR/EMR system | Ambulatory care | SBP mmHg, DBP mmHg, Serum cholesterol mmol/l, BMI kg/m2 | The groups did not differ for BP, cholesterol levels or BMI: SBP was 155.6 vs. 156.8 mmHG (95 % CI -0.6 to 3.0) between the control and the intervention group. DBP was 89.8 vs. 88.8 mmHg (95% CI -1.9 to -0.2). Serum cholesterol was 6.57 mmol/l vs. 6.64 mmol/l (95% CI -0.1 to 0.2) between the two groups. BMI was 27.7 kg/m2 vs. 27.8 kg/m2 (95% CI - 0.4 to 0.07). | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|---|--|---------|
| Holdsworth (2007) ³⁰⁵ Design: Before-after N = 2,407 patient admissions Implementation: 00/0000 Study Start: 00/0000 Study End: Oct 2004 | Administering, Prescribing | CPOE/POE system Integrated EHR/EMR system, Pharmacy | Acute care/tertiary, 50 Beds | Total ADEs per 100 admissions* Preventable ADEs per 100 admissions*, Potential ADEs per 100 admission*, | Patients were classified as having an ADE, a preventable ADE or potential ADE. All rates of ADEs were reduced after implementation of the CPOE system. Total ADEs per 100 admissions: 6.3 vs. 3.1 RRR 37%, 95% CI 0.05 to 0.57. Preventable ADEs per 100 admissions: 3.8 vs. 2.2, RRR 44%, 95% CI 0.09 to 0.66. Potential ADEs per 100 admissions: 7.9 vs. 2.2, RRR 63%, 95% CI 0.45 to 0.75. | + |
| Holman (1996) ³⁰⁶ Design: RCT N = 5 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Administering, Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated Handheld | Home | Pre-prandial blood glucose levels* | Pre-prandial blood glucose levels* were significantly less during the 'advice on' period compared to the 'advice off' period (7.5 vs. 8.9 mmol/l, p = 0.015) | + |
| Hwang (2002) ⁷⁰ Design: Time series N = 171 patients Implementation: 10/1999 Study Start: 06/1999 Study End: 05/2000 | Prescribing | CPOE/POE system Integrated Hospital information system Imaging system | Acute care/tertiary, 1,000 plus Beds Academic | LOS (mean number of days) | LOS in mean number of days decreased over the three time periods (11.4 vs. 10.3 vs. 8.2), with a significant reduction before implementation to 6 months postimplementation (p = 0.049). | + |
| Janssen (2009) ³⁰⁷ Design: Observational study N = 522 patients Implementation: 00/0000 Study Start: 10/2000 Study End: 04/2002 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system | Ambulatory care | PANSS Positive score*, PANSS Negative score* | Pronounced group-effect was found when comparing the Düsseldorf group using the Decision-Support System and the control group (Munich 1) providing treatment-as-usual, 14.1 (6.5) vs. 13.8 (6.7), p = 0.004) with respect to positive symptoms. No group effects were apparent concerning negative symptoms. The interaction effect of time 9 group was significant with regard to the negative score (p <0.039) and the positive score (p <0.001) (Figs. 1, 2). | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|---|---|---------|
| Javitt (2005) ²¹⁸ Design: RCT N = 39,462 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated Insurance | Ambulatory care | Admissions per 1,000 persons* | Among those in both groups who triggered recommendations, there were 19% fewer hospital admissions in the intervention group compared with the control group (213.8 ± 5.7 vs. 264.6 ± 5.7, p <0.001). | + |
| Keene (2007) ³⁰⁸ Design: Before-after N = 1,291 patients Implementation: 00/2001 Study Start: 09/2000 Study End: 02/2003 | Prescribing | CPOE/POE system Integrated EHR/EMR system, Laboratory system, Pharmacy | Critical care units (CCU, ICU, NICU), Academic | mortality* | Overall, 29 (3.16%) patients admitted during the pre-CPOE period and nine (2.41%) patients admitted in the post- CPOE period died under MMC care (p = 0.466). | - |
| Kucher (2005) ⁸⁹ Design: RCT N = 2,506 patients Implementation: 00/0000 Study Start: 09/2000 Study End: 01/2004 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, Hospital information system | Acute care/tertiary, Academic | Clinically diagnosed DVT at 90 Days, Clinically Diagnosed PE at 90 days | The primary end point for clinically diagnosed DVT at 90 days occurred in 103 (8.2%) in the control group as compared with 61 patients (4.9%) in the intervention group (RRR 40%, p = 0.001). For clinically diagnosed PE at 90 days the numbers were 35 (2.8%) in the control group as compared with 14 (1.1%) in the intervention group (RRR 61%, p = 0.004). The groups did not differ for proximal- or distal DVT, DVT of the arms, death, or hemorrhage. | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|--|--|---|---------|
| Lecumberri (2008) ⁹¹ Design: Time series N = 19,338 patients Implementation: 09/2005 Study Start: 01/2005 Study End: 06/2007 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated hospital guidelines | Unspecified Hospital, Academic | rate of VTE during hospitalization*, control to year 1, rate of VTE during hospitalization*, year 1 to year 2 | A non-significant reduction of VTE during hospitalization was achieved. Compared with the first semester of 2005, before implementing the computer-alert program, the overall rate of VTE during hospitalization was reduced from 3.26/1,000 (21 episodes in 6,441 patients) to 1.74/1,000 patients, (relative reduction 46.6%) in 2006. During the first semester of 2007, the rate of VTE during hospitalization was 1.67/1,000. OR: 0.53, 95% CI 0.25 to 1.10 and OR: 0.51, 95% CI 0.24 to 1.05 during the first semesters of 2006 and 2007 respectively, the impact being significant (p <0.05) among medical patients in 2007, OR: 0.36, 95%CI 0.12 to 0.98. | - |
| Lesprit (2009) ⁹³ Design: Observational study N = 932 prescriptions Implementation: 11/2006 Study Start: 11/2006 Study End: 10/2007 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Laboratory system | Acute care/tertiary, 960 Beds Academic | mortality rate*, median LOS*, readmission rate* | Clinical outcomes mortality (5.6% vs. 4.1%, RRR 27%, p = 0.348) and readmission (15.2% vs. 15.4%, RRR -1%, p = 0.936) were similar between intervention and non-intervention patients, LOS was significantly longer for intervention patients (15 days vs. 19 days, p = 0.011). | - |
| Lester (2005) ⁹⁴ Design: RCT N = 235 patients and 14 clinicians Implementation: 07/2003 Study Start: 07/2003 Study End: 07/2004 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care, Academic | LDL cholesterol | For the first assessment of LDL levels, the email group had lower levels of LDL cholesterol (138 vs. 119 mg/dL, p = 0.004). At the end of the study both groups had decreased their cholesterol levels and the difference between them was no longer seen (129 vs. 111 mg/dL, p = 0.055). | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|--|---------|
| Liu (2008) ⁹⁷ Design: Time series N = 858 patients Implementation: 00/1989 Study Start: 01/2005 Study End: 12/2006 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system | Acute care/tertiary | post-operative wound infection rates-clean procedures, post- operative wound infection rates- clean-contaminated procedures | The post-operative wound infection rate did not change significantly among 3 groups. In clean procedures, the post-operative wound infection rates were 0.63, 0.72 and 0.71% in group 1, group 2 and group 3, respectively (p = 0.995). In clean-contaminated procedures, the postoperative wound infection rates were 8.5%, 12.0%, and 9.4% in group 1, group 2, and group 3, respectively (p = 0.736). | - |
| Macdonald (2002) ²⁸⁵ Design: Before-after N = 5,008 patients Implementation: 07/1994 Study Start: 07/1994 Study End: 12/1998 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated Laboratory system | Acute care/tertiary, Academic | proportion of time within normal INR range*, readmission rates* | Patients in the computer group spent more time with INRs in the normal range (52% vs. 62%, p <0.05). No difference in readmission rates were found (3.8% vs. 3.0%, p = 0.9). | - |
| Madaras-Kelly (2006) ⁹⁸ Design: Time series N = not reported Implementation: 00/0000 Study Start: 07/2001 Study End: 06/2004 | Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated Hospital information system | Acute care/tertiary, 87 Beds | nosocomial infections rates* | All nosocomial infections decrease after the implementation of computer prompts, requirement for justification, and suggestion of alternate antibiotics beyond fluoroquinolones (1.37 cases/100 patient days vs. 0.62 cases, p = 0.02) | + |
| McGregor (2006) ¹⁰⁴ Design: RCT N = 4,507 patients Implementation: 00/000 Study Start: 05/2004 Study End: 08/2004 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated Laboratory system, Pharmacy | Acute care/tertiary, 648 Beds Inpatient hospital based, Academic | mortality, Length of stay | The groups did not differ from mortality. All patients 3.0% vs. 3.3%, p = 0.6) or for those patients who got alerts (8.2% vs. 7.8%, p = 0.5). Length of stay did not differ. All patients: 4.0 days vs. 3.8, p = 0.4 and 5 vs. 4 days for patients with alerts, p = 0.6 | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|----------------------------|--|---------|
| Meigs (2003) ³⁰⁹ Design: RCT N = 598 patients Implementation: 05/1998 Study Start: 05/1997 Study End: 04/1999 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated Laboratory system | Ambulatory care | HbA1c levels* | The intervention had a modest but nonsignificant benefit on glycemic control; HbA1c levels tended to improve in the intervention group (change -0.23) and worsen in the control group (change +0.14). p = 0.09 | - |
| Mekhjian (2002) ¹⁸⁶ Design: Before-after N = 28,898 patients Implementation: 05/2000 Study Start: 02/2000 Study End: 01/2001 | Administering, Transmission, order communication | CPOE/POE system, e- Medication administration system (e-MAR, e-TAR) Integrated Die-TARy system, EHR/EMR system, Imaging systems, Laboratory system | Acute care/tertiary, Other specialty hospital (rehab, oncology) Academic | severity-adjusted LOS * | After POE and e-MAR, severity-adjusted length of stay was reduced in OSUH (3.9 to 3.7 days, p = 0.002) but not James Cancer (3.7 to 3.6 days, NS) | + |
| Miskulin (2009) ¹⁰⁶ Design: Cohort study N = 8,941 patients Implementation: 00/2005 Study Start: 11/2005 Study End: 04/2006 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | Hb levels * | In the model adjusted for only center, average Hb levels were 11.8 ± 0.2 (SE) g/dL in patients treated using manual dosing and 0.11 ± 0.04 (SE) g/dL lower (p <0.001) in those treated with CDS. | + |
| Montgomery (2000) ¹⁰⁷ Design: RCT N = 552 patients Implementation: 00/0000 Study Start: 09/1996 Study End: 09/1998 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | SBP, DBP | SBP and DBP was not reduced in the CDSS group (SBP 153. vs. 153 mmHg) (DBP 85 vs. 85 mmHg) compared to the usual care group (EMR alone) | - |
| Murray (2004) ³¹⁰ Design: RCT N = 712 patients Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Pharmacy | Ambulatory care, Pharmacy, Outpatient hospital based, Academic | SF-36 QoL* | No intergroup differences were found for the primary endpoint the SF-36 QoL* scale (Table 3). No analysis presented. | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|---|---------------------------------------|---|---|---------|
| Niiranen (2008) ¹¹³ Design: Time series N = 18,818 patient followups Implementation: 03/2005 Study Start: 04/2005 Study End: 12/2007 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated Laboratory system | Ambulatory care, Home | proportion of patient followups with patients within p-INR, year 1 to 2 | The share of patient followups with patients within p-INR target range was significantly lower in year 2 than year 1 (67.1% vs. 63.1%, RRR 6%, p <0.001), then was constant from year 2 to year 3 (63.1% vs. 63.1%, RRR 0%, NS) | - |
| Novis (2010) ¹¹⁴ Design: Before-after N = 800 patients Implementation: 08/2007 Study Start: 03/2007 Study End: 03/2008 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Acute care/tertiary | postoperative bleeds, 30-, 60-, and 90- day DVT rates | Despite the increase in preoperative prophylaxis administration, there was no significant change in postoperative bleeds, with the rate of confirmed bleeds actually decreasing from 4% to 3% after implementation of the risk assessment (p = 0.34; NS). Over the course of the study, there was a trend toward decreased DVT events. The 30-, 60-, and 90-day DVT rates prior to implementation were 1.5%, 1.8%, and 2.0% respectively. After implementation, the 30-, 60-, and 90-day DVT rates were 0.3%, 0.5%, 1.3% respectively. This represents an overall 80% decrease in the 30-day rate of DVT and a 36% decrease in the 90-day rate of DVT, NS (p <0.12, p <0.58 respectively). There were no confirmed PE events at 90 days postoperation in this study population. | - |
| Oliven (2005) ¹¹⁵ Design: Cross-sectional N = 1,350 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated Drug order database, EHR/EMR system, Hospital information system, Laboratory system | Acute care/tertiary, 88 Beds Academic | Hospital stays | The average hospital stay was significantly shorter in department with CDOE than the department where prescriptions were handwritten and transcribed (6.9 ± 6.2 vs. 8.9 ± 7.9, p <0.001). | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|--|---|---------|
| Overhage (1997) ¹¹⁷ Design: RCT N = 86 physicians on 6 services (services randomized) Implementation: 00/0000 Study Start: 10/1992 Study End: 04/1994 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system, Laboratory system | General Hospital, Academic | LOS | LOS was not different for intervention patients compared with control patients (8.12 vs. 7.62, a difference of -0.5 days, 95% CI -0.17 to 1.19; p = 0.94). | - |
| Peterson (2005) ¹²⁴ Design: Cohort study N = 7,456 Medication orders Implementation: 00/0000 Study Start: 10/2001 Study End: 05/2002 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system | Acute care/ tertiary, Critical care units (CCU, ICU, NICU), 720 Beds Academic | LOS per 100 patient days, Fall per 100 patients days, Altered mental status per 100 patient days | There was no difference in the LOS between control and intervention period. (4 days for both, p = 0.43) or rate of altered mental status/100 patient days (21% vs. 22%, p = 0.17). The rate of falls was reduced in the CPOE group (0.64 falls/100 patient days for control vs. 0.28/100 patient days for the CPOE group, p = 0.001.) | - |
| Pielmeier (2010) ³¹¹ Design: Before-after N = 10 patients Implementation: 00/0000 Study Start: 02/2009 Study End: 03/2009 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Critical care units (CCU, ICU, NICU) | Mean log-normal BG ± standard deviation (mmol/L) | Hypoglycemia (blood glucose [BG] $b3.5\text{ mmol/L}$) was not observed. Mean log-normal BG ± standard deviation was reduced from 8.6 ± 2.4 mmol/L preintervention to 7.0 ± 1.1 mmol/L during the Glucosafe intervention (p <0.01). Mean log-normal BG ± standard deviation was reduced from 7.0 ± 1.1 mmol/L Glucosafe intervention to 7.4 ± 1.5 mmol/L during the intervention (p <0.03) | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|---|--|--|---------|
| Piontek (2010) ²⁹⁰ Design: Before-after N = 229,463 patients Implementation: 05/2001 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Laboratory system, Pharmacy | Unspecified Hospital, For Study group: bed sizes ranged from 58 to 303 (Indiana), 237 to 442 (Ohio), and 371 (California).For external control group: 460 (Maryland) and 365 (Idaho) Beds Inpatient hospital based | Mortality rates | Simple mortality rates exhibited no statistically significant changes in either the study group(3.86% vs. 3.87%, p <=0.999) or the control groups (2.99% vs. 2.88%, p = 0.963).However, severity-adjusted mortality rates decreased significantly only in the study group (1.049% vs. 0.975%, p <0.001). | - |
| Plaza (2005) ²⁷⁹ Design: RCT N = 198 patients Implementation: 03/2000 Study Start: 10/1999 Study End: 02/2001 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated Handheld | Ambulatory care | QoL-St George's Respiratory Questionnaire* | Scores on the St George's Respiratory Questionnaire were significantly lower for intervention patients (34.1 vs. 27.3, p = 0.002, difference 6.8 (95% CI 2.5 to 11.1). | + |
| Quinn (2008) ¹²⁷ Design: RCT N = 30 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Education of patients and clinicians but not pre- professional education, Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Daibetes Management Tool Integrated Web-based data analytics and therapy optimization tools | Ambulatory care | Average decrease in A1c values | Average decrease in A1c for intervention patients was 2.03% compared to .68% for control patients (p <0.04) | + |
| Ralston (2009) ³¹² Design: RCT N = 83 patients Implementation: 00/0000 Study Start: 08/2002 Study End: 05/2004 | Monitoring including patient adherence and compliance | Patient accessible Medical Record Integrated CDSS/CDS/CCDS/ reminders EHR/EMR system, Personal health records systems | Ambulatory care, Academic | Absolute change in GHb* | Absolute change in GHb declined significantly in the intervention group compared with the usual care group (0.2 vs. -0.9, change -0.7%; p = 0.01) at 12 months after adjusting for age, sex, and baseline GHb. | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|---|--|---------|
| Rasmussen (2005) ¹³² Design: RCT N = 253 patients Implementation: 00/0000 Study Start: 00/2001 Study End: 00/0000 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated Internet based electronic diary | Ambulatory care (clinic, doctors office etc) Academic | Improved Symptoms, Improved Quality of Life (AQLQ), Improved lung function (FEV1>=300 mL), Improved Airway hyperresponsiveness (AHR) | The treatment and monitoring with the Internet-based management tool lead to significant improvement in the Internet group regarding: Improved Asthma symptoms: Internet vs. specialist: OR 2.64 (95% CI 1.43 to 4.88), p = 0.002 Internet vs. GP: OR 3.26 (95% CI 1.71 to 6.19); p <0.001 2) Improved QOL: Internet vs. specialist: OR 2.21 (95% CI 1.09 to 4.47), p = 0.03 Internet vs. GP: OR 2.10 (95% CI 1.02 to 4.31), p = 0.04 3) Lung function: Internet vs. specialist: OR 3.26 (95% CI 1.50 to 7.11), p = 0.002 Internet vs. GP: OR 4.86 (95% CI 1.97-11.94), p <0.001 4)Airway responsiveness: Internet vs. Specialist: OR 1.26 (95% CI 0.57-2.79), p = NS Internet vs. GP: OR 3.06 (95% CI 1.13 to 8.31), p = 0.02 | + |
| Rind (1994) ²²⁵ Design: Time series N = 562 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Formulary, Hospital information system, Laboratory system, Pharmacy | Acute care/tertiary, 504 Beds Academic | difference in change in creatinine levels at 3 days*, difference in change of creatinine level at 7 days*, serious renal impairment | For medical service patients with changes in renal function, more patients had serious renal impairment in the control group compared (7.5% vs. 3.4%, p = 0.034). Difference in changes in creatinine levels at 3 days (14 mmol/L, p = 0.007) and 7 days (26 mmol/L, p <0.05) favored alerts medication event showed significant decreases for patients in the. | + |
| Rohrig (2008) ¹³⁵ Design: Before-after N = 156 patients Implementation: 00/1999 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system EHR/EMR system | Critical care units (CCU, ICU, NICU) 14 bed unit Beds Academic | Delta-SOFA, length of stay ICU (hours) duration of ventilation (hours) | Delta-SOFA decreased from 1.9% in the pre-period to 1.4% in the post-period, p = 0.23; length of stay (ICU) hours decreased from 472 to 337, p = 0.07; duration of ventilation hours decreased from 254 to 178, p = 0.07. | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|---|--|--|---|---------|
| Rollman (2002) ¹³⁶ Design: RCT N = 200 Patients with documented major depression Implementation: 00/0000 Study Start: 04/1997 Study End: 12/1998 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care | mean depression scores* | All groups improved their mean depression scores at 3 and 6 months. However, the groups did not differ from each other in mean scores at 3 or 6 months. | - |
| Roumie (2006) ²²⁶ Roumie (2007) ²²⁷ Design: RCT N = 871 patients Implementation: 00/0000 Study Start: 06/2004 Study End: 12/2004 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system | Ambulatory care, Outpatient hospital based | the proportion of patients achieving goal blood pressure | Patients of providers who were randomly assigned to the patient education group had better blood pressure control (138/75 mm Hg) than those in the provider education and alert or provider education alone groups (146/76 mm Hg and 145/78 mm Hg, respectively). More patients in the patient education group had a systolic blood pressure of 140 mm Hg or less, compared with those in the provider education or provider education and alert groups RR 1.31 (95% CI, 1.06 to 1.62) p = 0.012. The proportion achieving goal blood pressure differed in the 3 groups: 107/255 (42.0%) vs. 148/362 (40.9%) vs. 213/358 (59.5%) (p = 0.003) in the provider education; provider education and alert; and provider education, alert, and patient education groups, respectively. | - |
| Safran (1995) ¹⁴¹ Safran (1993) ¹⁴² Design: RCT N = 349 patients with HIV Implementation: 00/0000 Study Start: 05/1992 Study End: 09/1993 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated EHR/EMR system | Ambulatory care, Academic | rate of hospitalizations, rate of mortality | Patients in the control group had a higher rate of hospitalizations than those in the intervention group (44% vs. 35%, RRR 20%, p = 0.04). No significant difference in mortality rate (p = 0.18) | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|------------------------|--|---|----------|
| <p>Schmidt (2008)²⁶⁷ Design: Cohort study N = 62 patients with CHF Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Monitoring including patient adherence and compliance</p> | <p>patient adherence reporting Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Self-reported compliance, physical and mental health status in regard to health-related quality of life</p> | <p>The concordance between self-reported compliance (Yes/No) and telematic compliance monitoring was high; patients of the study group, who reported as noncompliant, highly significantly showed lower compliance scores, measured across a 2 month monitoring period with the telematic approach (T = 9.71, p <0.001). The same effect was true to the 6 month period (T = 3.51, p <0.01). Pre-post comparisons with respect to both physical and mental health status in regard to health-related quality of life showed significant differences between baseline and 1-month followup (T = -3.09, p ≤0.01), as well as baseline and 6-month followup (T = 1.81, p = 0.05). However, there were neither significant increases nor decreases between 1-month followup and 6-month followup. The changes from baseline to 1-month followup were stronger with respect to mental health than to physical health. Changes were insignificant in the control group.</p> | <p>-</p> |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--|---|---|---------|
| Schnipper (2009) ³¹³ Design: RCT N = 322 patients Implementation: 00/0000 Study Start: 05/2006 Study End: 06/2006 | Medication Reconciliation | medication reconciliation application Integrated CPOE/POE system | Acute care/tertiary Academic | PADE rate per patient*, PADE rate per patient per hospital, rate of hospital readmission or emergency department visit within 30 days | Among 160 control patients, there were 230 PADEs (1.44 per patient), while among 162 intervention patients there were 170 PADEs (1.05 per patient) (ARR, 0.72; 95% CI, 0.52 to 0.99). A significant benefit was found at hospital 1 (ARR, 0.60; 95% CI, 0.38 to 0.97) but not at hospital 2 (ARR, 0.87; 95% CI, 0.57 to 1.32) (p = 0.32 for test of effect modification). Hospitals differed in the extent of integration of the medication reconciliation tool into computerized provider order entry applications at discharge. The rate of hospital readmission or emergency department visit within 30 days was 20% in the intervention arm and 24% in the usual care arm (clustered OR, 0.76; 95% CI, 0.43 to 1.35). | + |
| Schnipper (2009) ³¹⁴ Design: Before-after N = 169 patients Implementation: 00/2006 Study Start: 07/2005 Study End: 06/2006 | Monitoring including patient adherence and compliance | CPOE/POE system Integrated EHR/EMR system, Hospital information system, Laboratory system, Pharmacy | Acute care/tertiary, Academic | Mean percent glucose readings 60–180 mg/dL per patient* | Mean percent of glucose readings between 60 and 180 mg/dL per patient, was 59.1% in the preintervention period and 64.7% in the postintervention (p = 0.13 in unadjusted analysis). When adjusted for A1c, admission glucose, and insulin use prior to admission, the adjusted absolute difference in the percent of glucose readings within range was 9.7% (95% CI [CI], 0.6%-18.8%; p = 0.04) | + |
| Sintchenko (2005) ¹⁵² Design: Before-after N = not reported n/a Implementation: 10/2002 Study Start: 04/2002 Study End: 03/2003 | Prescribing | CDSS/CDS/CCDS/reminders Integrated Laboratory system | Acute care/tertiary, Critical care units (CCU, ICU, NICU) 800 (18 bed ICU) Beds Academic | LOS, mean number of days*, mortality rate* | LOS decreased significantly from a mean of 7.12 days to 6.22 days (p = 0.02). Mortality rate was not different before and after the intervention (11.5% vs. 13.2%, NS) | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|--|--|--|---|--|---------|
| Steele (2005) ¹⁵⁷ Design: Before-after N = 54,206 patient visits Implementation: 12/2002 Study Start: 08/2002 Study End: 04/2003 | Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system EHR/EMR system, Laboratory system | Ambulatory care | “definite” or “probable” ADE rate | There was a non-statistically significant difference towards less “definite” or “probable” adverse drug events defined by Naranjo scoring (10.3% at baseline vs. 4.3% during postintervention, p = 0.23). | - |
| Takada (2003) ³¹⁵ Design: Before-after N = 374 patients Implementation: 00/1995 Study Start: 01/1998 Study End: 06/2002 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system | Other specialty hospital (rehab, oncology), 650 Beds | testing for renal function at 1 month*, 6 months*, decreased renal function at 1 month*, 6 months* | Introduction of the CDSS was not associated with testing for renal function at 1 month (43.8 vs. 48.3, p = 0.46 NS) or 6 months (85.3% vs. 84.5%, p = 0.84) or for having decreased renal function at 1 month (3.1% vs. 3.4%, p = 0.86) or at 6 months (5.4% vs. 5.1%, p = 0.92). NS at each stage. | - |
| Tierney (2003) ¹⁶⁶ Design: RCT N = 706 patients, 20 pharmacists, 94 physicians and 1 nurse practitioner Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Pharmacy | Ambulatory care, Outpatient hospital based, Academic | QOL SF-36*, heart failure exacerbation* | Across the 4 groups (physician intervention, pharmacist intervention, both interventions, and controls) the SF-36 (8 subscales), or for Heart Failure exacerbation (4 subscales), and emergency department visits or hospitalizations (all or related to HF) were NS. | - |
| Tierney (2005) ¹⁶⁷ Design: RCT N = 706 patients Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996 | Monitoring including patient adherence and compliance Prescribing | CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system, Pharmacy | Ambulatory care, Pharmacy, Outpatient hospital based Academic | QOL SF-36, Chronic Respiratory Disease Questionnaire, hospitalizations (Control vs. Physician intervention vs. Pharmacist Intervention vs. Both Intervention). | No significant change in QOL measures. Hospitalization was measured and not affected. | - |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|---|--|--|--|--|---|---------|
| Upperman (2005) ¹⁶⁸ Design: Before-after N = Not reported ADE/1,000 doses Implementation: 00/2002 Study Start: 01/2002 Study End: 00/0000 | Prescribing | CPOE/POE system Integrated EHR/EMR system | Acute care/tertiary, Pediatric stand alone hospital, Academic | rate of harmful ADEs | After implementation of CPOE the rate of harmful ADEs decreased (0.05/1,000 doses vs. 0.03/1,000 doses, p = 0.05). | + |
| Vartak (2009) ³¹⁶ Design: Before-after N = 41,865 patients Implementation: 00/2005 Study Start: 10/2004 Study End: 07/2005 | Prescribing | CPOE/POE system, EHR and Emergency Room (ER) event tracking system Integrated CPOE/POE system, EHR/EMR system, Hospital information system, Imaging systems, Laboratory system, Pharmacy | Emergency department, 193 Beds | Mean LOS*, number of patients treated in ED* | Although the system was designed to enhance efficiency, the mean (LOS) increased significantly from 116.8 minutes during the preimplementation period to 134.2 minutes during the postimplementation period (p <0.0001). The volume of patients treated in the ED however decreased significantly (p <0.0001) from preimplementation (n = 22,936) to postimplementation (n = 18,929). | - |
| Weingart (2008) ²⁷⁵ Design: Cohort study N = 267 patients Implementation: 09/2000 Study Start: 04/2001 Study End: 06/2002 | Monitoring including patient adherence and compliance | patient messaging via PHR Integrated Billing/administration system, EHR/EMR system, Imaging systems, Laboratory system | Ambulatory care, Academic | ADE rate | Patients experienced 21 total ADEs; responders reported significantly more ADEs electronically (13%) than non-responders (3%) RRR-333%, p = 0.01. | + |
| Wrona (2007) ¹⁷⁷ Design: Observational study N = 536 PCA patients Implementation: 00/2003 Study Start: 01/2003 Study End: 03/2004 | Monitoring including patient adherence and compliance, Prescribing | CPOE/POE system Integrated EHR/EMR system, Imaging systems, Laboratory system | Pediatric stand alone hospital | Occurrences of low respiratory rate, low oxygen saturation rate | Occurrences of low respiratory rate and low oxygen saturation were compared between 'no order set' and each of the two order sets groups (3.3% vs. 4.3% vs. 9.9%); the Acute Pain Service order set group had significantly higher rate of low respiratory rate (3.3% vs. 9.9%, - 200%, p <0.05). No significant differences were found in the number of cases in which low oxygen saturation was recognized (13.4% vs. 20.9% vs. 14.6%). | + |

Evidence Table 9. KQ1: primary clinical outcomes for all technologies assisting all medication phases (continued)

| Article Information | MM Phase(s) | HIT Studied Integrated System | Settings | Outcomes Measured | Results | Outcome |
|--|-------------|---|--------------------------------|--|--|---------|
| Yu (2009) ³¹⁷ Design: Case control N = 22,665 patients Implementation: 00/0000 Study Start: 10/2005 Study End: 09/2006 | Prescribing | CPOE/POE system Integrated Imaging systems, Laboratory system, Pharmacy | Pediatric stand alone hospital | Reportable ADE, no CPOE vs. some CPOE* | Univariate conditional logistic regression analysis showed that the lack of CPOE in hospitals was associated with increased risk of ADE. Specifically, after controlling for co-morbidities, the odds of experiencing a reportable ADE were 42% higher for hospitals without CPOE compared with those with CPOE, after adjusting for the number of co-morbidities. OR of experiencing a reportable ADE, no CPOE vs. some CPOE 1.42 (95% CI 1.28 to 1.57) | + |
| Zanetti (2003) ¹⁸⁰ Design: RCT N = 273 patients Implementation: 00/0000 Study Start: 03/2000 Study End: 06/ 2000 | Prescribing | CDSS/CDS/CCDS/reminders Integrated Hospital information system | Acute care/tertiary, Academic | rate of infection | The rate of infection was similar in both groups (4% in alarm plus reminder group vs. 6% in the control, p = 0.4) and both were lower than before the study (p = 0.2) | - |

Evidence Table 10. KQ1: primary qualitative outcomes for all technologies across phases

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|-------------|--|-------------------------------|---|---|
| Agostini (2007) ³¹⁹ Design: Qualitative N = 36 house officers most of whom were PGY1 Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CDSS/CDS/CCDS/reminders Integrated CPOE/POE system EHR/EMR system | Acute care/tertiary, Academic | Benefits and barrier themes were identified. Benefits include awareness of patient safety risks (delirium, falls, and general patient safety risks), usefulness of computer technology, and value of educational content of the reminder (geriatrics pharmacology review and nonpharmacologic treatment options). Barriers were related to demands of reading the reminder, role of clinical experience, and information content of the reminder. | Both barriers and benefits of computer-based reminders were identified by house officers dealing with the elderly patients with insomnia. |
| Ahearn (2003) ³²⁰ Design: Qualitative N = 22 general practitioners Implementation: 00/0000 Study Start: 04/2002 Study End: 05/2002 | Prescribing | e-Rx | Ambulatory care | 7 main themes emerging from the focus groups; (1) reaction to prompts; (2) concerns and potential problems re: comprehensiveness and accuracy of alerts; (3) effects on prescribing behaviour; (4) need for training; (5) helpful CDSS features e.g. sensitivity settings, alerts in red, etc; (6) suggested improvements; and (7) attitudes to evidence-based guidelines. | GPs believed that important interactions may be missed because of desensitization from too many alerts (which also intrude on workflow); that interaction alerts need to be severity graded and only significant ones should appear; and that improved computer-user interface design could enhance the usefulness of the decision support systems. |

The HIT system studied is in **bold**, followed by the systems that it was integrated with.

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: ADE = Adverse Drug Event; AMDs = Automated Medication Dispensing Systems; BCMA = Bar Code Medication Administration ; CCDS = Computerized Clinical Decision Support; CDS = Clinical / Computerized Decision Support ; CDSS = Clinical Decision Support System; CIT = Clinical Information Technology; CPOE = Computerized Provider Order Entry; CR = computer reminder; ED = Emergency Department; EDI = Electronic Data Interchange ; EHR = Electronic Health Record; e-MAR = Electronic Medication Administration Record; EMR = Electronic Medical Records; EPA = Electronic Prescribing and Administration System; e-RX = Electronic Prescribing; e-TAR = Electronic Treatment Authorization Request; GPs = General Practitioners; HIT = Health Information Technology; ICT = Information and Communication Technology; MICU = Medical Intensive Care Unit; MM = Medication Management; N = sample size; OTC = Over the counter; PA = Physician Assistants; PGY1 = First Year Postgraduate; POE = Provider Order Entry

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|---|--|---|--|---|
| Arar (2005) ³²¹ Design: Observational study N = 50 clinical encounters with patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Education of patients and clinicians but not pre-professional education | e-Rx Integrated EHR/EMR system, Imaging systems, Laboratory system | Ambulatory care | Direct observation and content analysis showed that the EMR/e-Rx facilitated communication with respect to the process of care that included checking active and inactive prescriptions and new and refill prescriptions, names of medication, and other medication themes (mail-order issues, adherence, self regulation, alternate OTC issues). | The EMR improved communication between physicians and patients in relation to medication issues. |
| Ash (2004) ³²² Ash (2003) ³²³ Sittig (2005) ³²⁴ Ash (1999) ³²⁵ Ash (2000) ³²⁶ Ash (2003) ³²⁷ Ash (2001) ³²⁸ Design: Qualitative N = 58 physicians, nurses, administrators, IT professionals Implementation: 1966 onwards Study Start: 00/1998 Study End: 00/2003 | Prescribing | CPOE/POE system | Acute care/tertiary, General Hospital, Academic | ³²⁴ Negative emotional responses were more prevalent than positive or neutral. ³²³ Four high-level themes were identified: (1) organizational issues such as collaboration, pride, culture, power, politics, and control; (2) clinical and professional issues involving adaptation to local practices, preferences, and policies; (3) technical/ implementation issues, including usability, time, training and support; (4) issues related to the organization of information and knowledge, such as system rigidity and integration. Relevant differences between teaching and nonteaching hospitals include extent of collaboration, staff longevity, and organizational missions. ³²² Themes included: language and misunderstandings, context matters and it affects the way of doing things, benefits and tradeoffs, 'contrasts, conflicts and contradictions', collaboration and trust, customization and organization of information, defining boundaries of CPOE, ongoing nature of implementation. ³²⁷ Explores the theme of leaders and bridgers-administrative; clinical; | ³²⁴ Designers need to recognize that CPOE features and implementation strategies can increase negative emotions and impact success of implementation. Positive feedback might alleviate some of the problems. ³²³ An organizational culture characterized by collaboration and trust and an ongoing process that includes active clinician engagement in adaptation of the technology were important elements in successful implementation of physician order entry at the institutions that we studied. ³²² Publication of the results of these iterative inquiries served to promote a realization that implementation of CPOE is not easy and that the negatives must be weighed against the positives. ³²⁷ Understanding multiple perspectives should be undertaken, with insights used to form strategic implementation plans. ³²⁵ house officers felt that CPOE assists patient care but may undermine education; it works best when tailored to fit local and |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--|-----------------------------------|---------------|--|---|
| | | | | bridgers/support staff; skills and training. ³²⁶ Physicians, admin and IT have different perspectives of the technical and organizational aspects of CPOE; the multiple perspectives model was used to offer structure to the results. ³²⁵ Themes relating to housestaff perceptions of CPOE included education; benefits; problems; feelings about; implementation strategies and the future of CPOE. | individual workflow; implementation strategies should include mechanisms for engaging housestaff in decision process. |
| Avery (2005) ³²⁹ Design: Survey N = 21 experts (Delphi panel members) Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Monitoring including patient adherence and compliance, Prescribing | e-Rx | Not specified | Key Themes: (1) importance of computerized alerts; (2) need to minimize spurious alerts; (3) making it difficult to override critically important alerts; (4) having audit trails of such overrides; (5) support for safe repeat prescribing; (6) effective computer-user interface; (7) importance of call and recall; (8) need to be able to run safety reports. | The high level of agreement among the expert panel members indicates clear themes and priorities that need to be addressed in any further improvement of safety features in primary care computing systems. |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|----------------------|---|--|---|--|
| <p>Banet (2004)¹⁹⁸ Design: Before-after N = 55 nurses Implementation: 05/2003 Study Start: 00/0000 Study End: 00/0000</p> | <p>Administering</p> | <p>CPOE/POE system, e-MAR, e-Medication administration system (e-MAR, e-TAR) Integrated Imaging systems, Laboratory system, Pharmacy</p> | <p>ED, Academic</p> | <p>For the open-ended question on ease of CPOE documentation, responses fell into the following themes: improvements in the clarity of orders, system helps organize and time their tasks, positive responses about efficiency and standardization of documentation provided by templates, general improvement in ED processes, decreased number of verbal orders and time searching for charts. For the open ended question for suggestions for improvements, themes included: additional terms and phrases for templates, process issues not affected by the ED application, complaints regarding technical problems with the system, suggestions for additional functionality, comments about the medication order icon on the tracking board.</p> | <p>The findings from this study indicate that users perceived no change in the total amount of time spent on documentation, a perception that was corroborated by the results of the time-motion studies. Nurses also perceived that certain processes, such as laboratory and radiology tests, were accomplished more efficiently after the implementation.</p> |
| <p>Bastholm Rahmner (2004)³³⁰ Design: Qualitative N = 21 physicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>e-Rx Integrated CDSS/CDS/CCDS/ reminders Pharmacy</p> | <p>General Hospital</p> | <p>4 categories for possibilities and obstacles. (1) possibilities related to access to patient drug history (which is not met by the new system), increased pharmacological knowledge from alerts etc., access to information more readily and time saved; (2) obstacles centered around technical problems given current problems with the EMR and too frequent alerts, computer shortages within the ED, altering routines and habits and the resulting diminishing patient contact since they need to leave the consulting room to enter the prescriptions;</p> | <p>Gaining access to patient drug history enables physicians to carry out work in a professional way. Alerts and producer-independent drug information are valuable in reducing workload. However, technical prerequisites form the base for a successful implementation. Time must be given to adapt to new ways of working.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|---------------|--|--|--|---|
| Beuscart-Zephir, (2010) ³³¹ Design: Qualitative N = Not Specified Nurse Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Administering | CPOE/POE system Integrated EHR/EMR system | Acute care/ tertiary, 3,000 Beds Academic | (3) standard ethnographic methods were used to support the analysis of the current work system and work situations, coupled with cognitive task analysis methods and documents review; (4) usability inspection (heuristic evaluation) and both in-lab (simulated tasks) and on-site (real tasks) usability tests were performed for the evaluation of the CPOE candidate. The study focused on the nurses' tasks of preparing and administering oral route drugs to the patients, with a particular attention to the nurses' needs in terms of information necessary to efficiently and safely support their tasks. | The analysis of the work situations identified different work organizations and procedures across the hospital's departments. The most important differences concerned the doctor–nurse communications and cooperation modes and the procedures for preparing and administering the medications. The assessment of the medication CPOE functions uncovered a number of usability problems, including severe ones which could be impossible to detect. |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--------------------|---|---|--|---|
| <p>Boonstra (2004)³³² Design: Qualitative N = 36 GPs Implementation: 00/2001 Study Start: 00/2001 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/ reminders e-Rx Integrated CDSS/CDS/CCDS/ reminders EHR/EMR system, Formulary, Insurance</p> | <p>Ambulatory care</p> | <p>Five factors related to the perceived advantages and disadvantages of the system. (1) system: usability issues and features of the system viewed as positive by some (user friendly, integrated) and negative by others (unfamiliar with the disease codes system, lack of flexibility, lack of computer resources); (2) finance: though the software was free, the government reaped the economic benefits of using it and the GPs were required to keep their EHR systems up to date; (3) system in consultation process: some felt it was more efficient during consultation and led to better quality; others felt it took longer and took away from patient focus; (4) cultural factors: users tended to have a culture of professional quality, non-users tended to focus on human relations; (5) policy environment: helps doctors become more cost conscious, but benefit only for insurers, and focused solely on costs.</p> | <p>Designing a system that met the diverse needs of users more satisfactorily, in being more compatible with their diverse cultures, may have encouraged wider and more creative use, and thus achieved more savings than the present arrangements have achieved.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|----------------------|-----------------------------------|----------------------------|---|---|
| <p>Buhrer (2008)³³³ Design: Qualitative N = 67 Nurses Implementation: 11/2007 Study Start: 10/2007 Study End: 12/2007</p> | <p>Administering</p> | <p>BCMA</p> | <p>Acute care/tertiary</p> | <p>Pre-BCMA: (1) scheduled medication passes take longer (35/35); (2) system is overwhelming (25/35); (3) the system would direct nurses' attention away from patients (23/35); (4) nurses expected system to improve patient safety (30/35). Post-BCMA: (1) liked working with the system (19/32); (2) BCMA improves safety (28/32); (3) overwhelmed at beginning of the implementation (12/32); (4) more focused on system than the patient and found this annoying (20/32); (5) would like to switch back to the previous, paper-based system. Negative attitude: (1) computer carts: too heavy and too big and some without storage drawer; (2) scanners: too few wireless scanners; (3) batteries: unreliable power indicators and weak batteries; (4) lost orders: sometimes disappeared from the medication schedule, causing confusion; (5) documentation: required launching a separate cumbersome application. Positive attitude: (1) organization: nurses found BCMA system's scheduling function helpful; (2) carts: some use cart as a "portable desk".</p> | <p>Implementation of BCMA into the active process of medication administration was a significant source of negative attitudes in nurses. Qualitative examination of users' attitudes (negative and positive) toward specific attributes can result in improved design of both technology and implementation strategies.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--|---|--|---|--|
| <p>Campbell (2009)³³⁴ Design: Qualitative N = 32 semi-structured interviews=43 hours; 400 hours of observation shadowing 95 clinical providers Implementation: 00/0000 Study Start: 08/2004 Study End: 04/2005</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated EHR/EMR system</p> | <p>Acute care/tertiary, General Hospital, 340 (Wishard); 893 (Mass.); 150 (Faulkner); 725 (Brigham); 238 (Alamance) Beds Academic</p> | <p>Themes: CPOE systems can affect clinical work by: (1) introducing or exposing human/computer interaction problems; (2) altering the pace, sequencing, and dynamics of clinical activities; (3) providing only partial support for the work activities of all types of clinical personnel; (4) reducing clinical situation awareness; (5) poorly reflecting organizational policy and procedure.</p> | <p>CPOE systems are tools intended to support and improve the delivery of care, and are not solutions for all problems related to clinical practice. Workflow issues resulting from CPOE can be mitigated by iteratively altering both clinical workflow and the CPOE system until a satisfactory fit is achieved.</p> |
| <p>Cross (2009)³³⁵ Design: Qualitative N = 10 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Monitoring including patient adherence and compliance</p> | <p>CDSS/CDS/CCDS/reminders home automated tele-management Integrated electronic messaging system, Personal health records systems</p> | <p>Ambulatory care, Home, Academic</p> | <p>Patients' perceptions: (1) constant communication: assist them in monitoring the symptoms of disease, both from a medical provider and a patient perspective; (2) use of computer was not difficult; (3) improved safety; (4) keep the patient and provider up to date on changes in symptoms. Analysis of the responses were sorted into three topic areas: (1) user attitudes about the interface; (2) user attitudes about the content of self-testing; (3) user attitudes about the self-testing process.</p> | <p>Pilot testing of a tele-management system customized for UC revealed a high level of acceptance and interest among patients. The results suggest that implementation of a tele-management system will be feasible on a long-term basis with only minor modifications.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--|---|---|--|---|
| <p>Crosson (2007)³³⁶ Design: Qualitative N = 47 clinicians Implementation: 10/2006 Study Start: 03/2006 Study End: 11/2006</p> | <p>Prescribing, Transmission, order communication</p> | <p>e-Rx, e-Transmission of the prescription to/from doctor to pharmacy Handheld Integrated, EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Practices which successfully implemented the e-Rx system exhibited greater familiarity with the capabilities of the systems and had more realistic expectations of the benefits. Physicians in these practices tended to have positive attitudes about and previous experiences with e-Rx or EMR, participation in continuing education courses relating to e-Rx, and plans for the future use of other HIT. Physicians in the 3 practices where the programs were successfully installed but unevenly implemented had high expectations about the ease of implementation, but at the same time reported concerns about how e-Rx might affect their clinical independence or undermine their authority with patients. Prescribers and staff members in the 2 practices that successfully installed, but then discontinued use of the program exhibited very little advance knowledge of program functions or the potential effect on prescription workflow. Two practices failed to install e-Rx; physicians and support staff in these practices expected that e-Rx would lead to greater efficiency and safety but, at the same time, had little specific knowledge of program functionality.</p> | <p>Practice leaders should plan implementation carefully, ensuring that practice members prepare for the effective integration of e-Rx technology into clinical workflow.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|-------------|--|--|---|---|
| Feldstein (2004) ³³⁷ Design: Qualitative N = 20 Clinicians Implementation: 00/1996 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated CDSS/CDS/CCDS/ reminders EHR/EMR system | Ambulatory care | The study found some common theme with respect to prescribers' frustrations with CPOE systems: (1) alerts that contained low-priority information; (2) intrusive alerts presented at the wrong time in the workflow; (3) difficult-to-interpret alerts; (4) delays caused by the alert; (5) redundant and repetitive alerts. | Although alerts may slow the work process, busy clinicians generally find them helpful. Safety alerts need to be concise and relevant, have clear action steps, and provide options for users with different experience levels and work styles. Health care decisionmakers should prioritize safety-related alerts and educational programs to facilitate the implementation of CDSS at CPOE. |
| Fernando (2009) ³³⁸ Design: Qualitative N = 9 ED specialists and registrars Implementation: 01/2006 Study Start: 05/2006 Study End: 12/2006 | Prescribing | CPOE/POE system Integrated EHR/EMR system, Imaging systems, Laboratory system | Acute care/tertiary, 66 ED beds | Three major issues emerged from the findings: (1) the implementation of the new system was accompanied by major shifts in ED work responsibilities and tasks; (2) the appearance of dysfunctional consequences related to the excess time it took to electronically order and the usability of some features of the new system; (3) doctors' concerns that their views and opinions about design and implementation of the new system had not been adequately addressed | The implementation of electronic ordering has important implications for ED functioning and the delivery of patient care. The complexity of the ED makes it vulnerable to disruption caused by inadequate system design and ineffective channels of communication across the hospital. |
| Fields (2007) ³³⁹ Design: Qualitative N = 17 Health care providers Implementation: 08/2006 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CPOE/POE system Integrated Hospital information system | Acute care/tertiary, Critical care units (CCU, ICU, NICU) 19 beds in the MICU Beds | Four themes were suggested: (1) ease of use; (2) speed; (3) trust; (4) hopefulness. Participants valued CPOE potential and were hopeful that future systems would be easy to use, decrease error potential, be more customizable for individual users, and contain concise physician order sets to foster medication order safety. | Participants valued CPOE potential although they commented on improvements and challenges. |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|---|--|--|--|--|
| Franklin (2007) ⁵⁰ Donyai (2008) ⁵¹ Barber (2007) ⁵² Franklin (2008) ⁵³ Franklin (2007) ⁵⁴ Design: Before-after N = 4,803 medication orders Implementation: 06/2003 Study Start: 00/0000 Study End: 00/0000 | Administering, Prescribing | Automated Dispensing Machine, e-Medication administration system (e-MAR, e-TAR) e-Rx Integrated Pharmacy | Acute care/tertiary, 28 surgery bed ward of a teaching hospital Beds Inpatient hospital based, Academic | The system was successfully implemented on the ward, and remained in operation for over 2 years. Many of the technical components of the system initially showed problems, but evolved with increased functionality and improved performance. Attitudes to the system in the early stages were mixed. Over time, staff attitudes changed to become more balanced and the potential benefits of the system became clearer to most. The system structured the work of staff, sometimes unexpectedly. | This theory-led evaluation offers valuable insights into a critical contemporary policy area. Technical systems are never perfect, and they require time and effort to become embedded into any particular clinical context. The effectiveness of ICT changes and develops over time, have quite different effects in different settings. For this reason a sophisticated evaluation framework is necessary. |
| Georgiou (2009) ³⁴⁰ Design: Qualitative N = 50 hospital employees Implementation: 00/0000 Study Start: 01/2006 Study End: 03/2006 | Prescribing, Transmission, order communication | CPOE/POE system Integrated EHR/EMR system, e-MAR | Acute care/tertiary, Not specified, Inpatient hospital based, Academic | The 20 recurring themes were grouped into 4 major constructs: Will it help?, Will it work?, Will it impair existing interaction?, and Will we cope? | The hospital employees had major concerns before implementation of a CPOE system. The elucidation and understanding of these concerns and worries can help to inform and strengthen implementation strategies. |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|--|--|---|--|--|
| <p>Graham (2008)³⁴¹ Design: Qualitative N = 7 physicians Implementation: 00/0000 Study Start: 00/2006 Study End: 00/2007</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system, Formulary</p> | <p>Emergency department, Academic</p> | <p>Coding categories for identifying usability problems from the analysis of video-based data included: (1) interface problems; (2) content problems; (3) slips and mistakes. From 56 recorded sessions, a total of 422 events were recorded. The events were further grouped into seven main categories: (1) negative comments; (2) positive comments; (3) neutral comments; (4) application events; (5) problems; (6) slips; (7) mistakes.</p> | <p>This study provides a framework for evaluating CDSS applications in a clinical environment and has identified specific areas for improvement in the applications utilized. A number of interface issues that could lead directly to adverse medical events that were identified raises concerns about the potential for similar undocumented problems in other clinical applications currently in use or being developed for implementation. Application of usability engineering principles can help identify interface problems that may lead to medical adverse events, and need to be incorporated early in the design phase to ensure that such problems can be corrected while there is still time and it is cost effective to do so.</p> |
| <p>Grossman (2007)³⁴² Design: Qualitative N = 26 organizations Implementation: 00/0000 Study Start: 11/2005 Study End: 03/2006</p> | <p>Prescribing, Transmission, order communication</p> | <p>e-Rx, e-Transmission- of the prescription to/from doctor to pharmacy Integrated Stand-Alone, EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Qualitative data were narratively analyzed from 44 telephone interviews with 26 medical practices, 21 with e-Rx.</p> | <p>Barriers were reported related to maintaining complete lists of patients and their medications, use of CDSS, getting patient-specific formulary data, and EDI. Factors associated with these issues related to product limitations, external implementation challenges (e.g., communication with pharmacists and vendor support), and physician preferences on specific product features.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|----------------------|---|---------------------------------------|---|---|
| <p>Holden (2010)³¹⁸ Design: Qualitative N = 20 attending Physicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system, EMR Integrated EHR/EMR system</p> | <p>Acute care/tertiary, 400+ Beds</p> | <p>Behavioral beliefs: (1) performance outcomes; (2) productivity and efficiency outcomes; (3) patient outcomes; (4) financial, organizational, and other outcomes; (5) affective outcomes; External Normative beliefs; Control beliefs: controllability; self-efficacy.</p> | <p>EMR and CPOE were commonly believed to both improve and worsen the ease and quality of personal performance, productivity and efficiency, and patient outcomes. Physicians felt encouraged by employers and others to use the systems but also had personal role-related and moral concerns about doing so. Perceived facilitators and barriers were numerous and had their sources in all aspects of the work system.</p> |
| <p>Hurley (2007)²⁴¹ Design: Mixed methods N = 1,087 nurses Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Administering</p> | <p>BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated CPOE/POE system, EHR/EMR system, Pharmacy</p> | <p>Acute care/tertiary, Academic</p> | <p>Interview questions followed the same subscales as the satisfaction scale. Nurses found the new system more time consuming but acknowledged that the extra time was wisely spent to assure verification. They viewed saving time on handwritten, paper-based medication sheets transcribing as a positive change. They felt there was an increased sense of safety for the patients and the nurses and that the system helped with the 5 rights. In terms of access, they appreciated greater access to medications and information (policies, guidelines, drug resources, patient files, etc.), but felt there were still some delays in getting medications from pharmacy.</p> | <p>A medication administration system that nurses view as being effective, by promoting efficacy, safety, and easy access, will support their nursing practice. Results of this study can give confidence to nurse executives that nurses can be satisfied with technology to make medication administration safer and more efficient and provide easier access to system components.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|--|--|---|--|---|
| <p>Johansson (2010)³⁴³ Design: Mixed methods N = 15 home-care nurses Implementation: 00/2007 Study Start: 12/2007 Study End: 3/2008</p> | <p>Administering</p> | <p>BCMA, CDSS/CDS/CCDS/ reminders Integrated Handheld, Drug reference software</p> | <p>Home, Academic</p> | <p>Medication profiles: Most of the nurses had access to the mobile information and the possibility to obtain a profile of the patients' medication regarding drug–drug interactions, therapeutic duplications and warnings for drugs unsuitable for elderly. Usability: The nurses discussed that it was a time consuming learning threshold, but once used to the LIFe-reader®s' functions, they were regarded as fast. The nurses experienced that the keyboard of the LIFe-reader® was too small and not suited to the Swedish language, and that the pen was not easy to use. Usefulness: The nurses believed that it would be different to use the PDA once they started to work as district nurses. Some nurses thought they would have used the LIFereader ® in a different way if they could have had it for a longer time or if they knew they could have kept it. The drug reference text in the LIFe-reader® had the highest priority but there was also a potential for more functions and features</p> | <p>We found that the LIFe-reader® has the potential to be a useful and user-friendly MDSS for nurses in home care when obtaining profiles of the patients' medication regarding drug–drug interactions, therapeutic duplications and warnings for drugs unsuitable for elderly patients.</p> |
| <p>Johnson (2010)⁷⁵ Design: RCT N = 3,285 patients Implementation: 00/0000 Study Start: 04/2007 Study End: 08/2007</p> | <p>Prescribing, Transmission, order communication</p> | <p>CDSS/CDS/CCDS/ reminders e-Rx Integrated EHR/EMR system</p> | <p>Ambulatory care, Pharmacy, Not specified, Academic</p> | <p>Improving communication between prescribers and dispensers; Decreases callbacks in some cases; Pediatric dosing information helps check for potential errors; Increases callbacks in some cases; Need more information to be included in annotations; New SYW feature request</p> | <p>Comments suggested that SYW increased callbacks where necessary and decreased them in other situations, but did not contribute to unnecessary callbacks. These findings support the continued and potentially expanded use of SYW by e-Rx systems to enhance communication with pharmacists.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|---------------|--|---|--|---|
| Kazemi (2008) ³⁴⁴ Design: Qualitative N = 19 physicians Implementation: 00/0000 Study Start: 12/2006 Study End: 01/2007 | Prescribing | CPOE/POE system Integrated Hospital information system | General Hospital, 234 Beds Academic | 3 themes emerged on current prescription process: (1) decision-making errors; (2) transcription errors; (3) over confidence errors. 3 themes were identified in the expected benefits category: (1) confidentiality issues; (2) reduction of medication errors; (3) educational benefits. 4 themes emerged in the perceived obstacles category: (1) high cost; (2) social and cultural barriers; (3) data entry time; (4) problems with technical support. | Prescription patterns in Iranian teaching hospitals are physician centered, top-down with possibility for medication errors. Although barriers exist towards implementation of CPOE, there is a general willingness among the physicians to use such a system if it provides significant benefit. |
| Koppel (2005) ³⁴⁵ Design: Mixed methods N = 291 health care providers Implementation: 00/1997 Study Start: 00/2002 Study End: 00/2003 | Prescribing | CPOE/POE system Integrated nurses medication lists, Pharmacy | Acute care/tertiary, 750 Beds Academic | Identified 22 previously unexplored medication-error sources that users report to be facilitated by CPOE. We group these as: (1) information errors generated by fragmentation of data and failure to integrate the hospital's several computer and information systems; (2) human-machine interface flaws reflecting machine rules that do not correspond to work organization or usual behaviors. | A leading CPOE system often facilitated medication error risks, with many reported to occur frequently. As CPOE systems are implemented, clinicians and hospitals must attend to errors that these systems cause in addition to errors that they prevent. |
| Koppel (2008) ³⁴⁶ Design: Mixed methods N = 14,2203 medication administrations Implementation: 12/ 2001 Study Start: 00/2003 Study End: 00/2006 | Administering | BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated Hospital information system | Acute care/tertiary, 1,399 Beds Academic | 15 workarounds falling into 3 categories were identified: omission of process steps (7 workarounds), steps performed out of sequence (1 workaround) and unauthorized process steps (7 workarounds). The probable causes and potential errors for each workaround were determined. Probable causes included technology, task, organizational, patient and environmental related causes. | BCMA systems are intended to advance medication safety, our data reveal that integrating BCMAs within real-world clinical workflows requires critical attention to ensure that technology safety features are used as intended and that systems are designed to support this use. Compliance with patient safety protocols is best achieved by configuring BCMAs for efficient as well as safe patient care. Repeated examinations and corrections of BCMA actual uses are needed to optimize their role in preventing medication errors. |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|---|---|---|---|--|
| <p>Krall (2002)³⁴⁷ Design: Qualitative N = 16 clinicians (physicians, PAs and nurses) in 3 focus groups Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Monitoring including patient adherence and compliance, Prescribing</p> | <p>Outpatient EMR Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>5 themes were identified from the focus group data. (1) efficiency: alerts and reminders being efficient and not wasting time; (2) usefulness: alerts being useful and appropriate; (3) Information content: about timely, rich, and accessible information; (4) user interface: important for smooth and efficient work and provision of valuable information quickly and accurately; (5) workflow: issues related to the information being available when needed. Note that considerable emotion was associated with alerts and reminders (criticism, embarrassment, guilt, frustration, annoyance, and anger).</p> | <p>the clinicians provided considerable feedback on the usefulness and usability of alerts and reminders in EMRs.</p> |
| <p>Lai (2007)³⁴⁸ Design: Mixed methods N = 15 pharmacists Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Dispensing, Transmission, order communication</p> | <p>CPOE/POE system</p> | <p>Unspecified Hospital, Pharmacy, Inpatient hospital based</p> | <p>1) patient safety: pharmacy leaders all believed CPOE would improve patient safety, allergy, dosing and interaction alerts. Some expressed concern that poor design/implementation could lead to increased errors; 2) pharmacy practice: most believed the system would lead to improved efficiencies facilitating more time spent with patients; 3) pharmacy profession: most felt CPOE would improve working relationships with physicians and nurses by facilitating new collaborations The scaling analysis found that pharmacy leaders of community, academia, and hospitals had different experience and/or opinions regarding the impact of CPOE.</p> | <p>Most pharmacy leaders held positive opinions regarding the impact of CPOE on the pharmacy practice and the profession, with varying concerns regarding its impact on practice and safety.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--------------------|---|----------------------------|---|--|
| <p>Lapane (2008)⁹⁰ Design: Mixed methods N = 276 primary care prescribers and their staff Implementation: 00/2003 Study Start: 04/2006 Study End: 08/2006</p> | <p>Prescribing</p> | <p>e-Rx</p> | <p>Ambulatory care</p> | <p>An open-ended approach was used to elicit information about the benefits and drawbacks of e-Rx. 15 different parent nodes were defined. Attention focused on 2 parent nodes, impact on clinical practice and software features. Physicians found the drug allergy alerts useful. For drug-drug interactions, they found these beneficial to patient safety. Many of the interaction alerts were however ignored and many were viewed as too trivial or unnecessary. Physicians suggested that alerts be provided for current medication only and for them to be less sensitive, more sensible, possibly having a personal setting for severity levels.</p> | <p>Prescribers believe that refinements to the drug alerting systems are necessary to reduce common overriding of alerts. In addition to honing the specificity of the alerts and permitting prescribers to set the severity threshold for alerts, prescribers recommend having the drug alert algorithms run against current medication regimens.</p> |
| <p>Li (2006)²⁴⁷ Design: Qualitative N = 2 qualitative researchers (nurse and human factors psychology) Implementation: 02/2004 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated Hospital information system</p> | <p>Acute care/tertiary</p> | <p>The 2 researchers used heuristic methods and identified 5 major problem areas with the CPOE system. These problems centered on text presentation, too much information/too many decisions at one time, color scheme (monochromatic blue/grey with red used as accent and not to note caution or problems). Problems were given to the developers who addressed them in the next redesign of the system.</p> | <p>The 5 problem areas that were identified were given to the developers who addressed them in the next redesign of the system.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--|---|---|---|---|
| <p>McAlearney (2006)³⁴⁹ Design: Qualitative N = 71 Healthcare providers (Physicians) Implementation: 00/0000 Study Start: 04/2002 Study End: 05/2005</p> | <p>Prescribing</p> | <p>Computerized order sets & hand held computers, CPOE/POE system Integrated EHR/EMR system</p> | <p>Pediatric stand alone hospital, Ambulatory care, Other, Academic</p> | <p>2 major themes emerged: (1) Can it work? Physicians expressed concerns about: (a) appropriateness of physician-directed CIT as a solution for medical errors; (b) current technical capabilities; (c) level of technical support for CIT solutions; (d) introduction of new errors. (2) At what cost to the medical profession? Physicians were concerned about the time efficiency and workload redistribution associated with the introduction of CIT.</p> | <p>The study concluded that health care organization attempting to promote physician use of CIT should consider physician's perspectives about technology adoption and use to address their concerns, reduce skepticism, and increase the likelihood of implementation success.</p> |
| <p>McCann (2008)²⁵¹ Design: Mixed methods N = 53 patients Implementation: 0 Study Start: 03/2006 Study End: 09/2006</p> | <p>Monitoring including patient adherence and compliance</p> | <p>symptom management system Integrated Handheld</p> | <p>Ambulatory care</p> | <p>1) training and familiarization of the handset: patients felt that the training was adequate and the handset was straightforward and easy to use; 2) length of data collection: patients felt that entering data twice a day for 14 days was acceptable; 3) daily routine: the system did not appear to impact on patients' daily routines as it was incorporated into their day in a variety of ways; 4) symptoms: patients often felt that the six symptoms that were recorded on the handset were adequate, although some patients did indicate that they would have liked the opportunity to report other symptoms; 5) the alerting facility: overall, patients were happy with the alerting facility of the system, and the real-time, quick response rate of the data collected.</p> | <p>The results of this study indicate that patients with breast, lung and colorectal cancer had positive perceptions and experiences of using ASyMS© to monitor and manage chemotherapy related toxicity.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|-------------|--|--|--|--|
| Motulsky (2008) ³⁵⁰ Design: Qualitative N = 12 community pharmacists Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Prescribing | e-Rx Integrated Insurance, Personal health records systems | Pharmacy, Other | The model of the effects of e-Rx on professionalization of community pharmacists had 7 themes: (1) increased analytical capacity; (2) greater dissemination of knowledge; (3) better integration of process tasks; (4) increased process automation; (5) elimination of intermediaries; (6) increased tracking capability; (7) greater informational capability. The main effects of the e-Rx were analytical capacity of the pharmacists and physician and dissemination of knowledge, integration of process tasks, process automation, facilitates interpretation of prescriptions, improves relevance and meaningfulness of interaction and improves quality of information transmitted. | e-Rx has tremendous capacity to change and improve pharmacists professional work and interactions. |
| Nanji, (2009) ³⁵¹ Design: Qualitative N = 10 pharmacy staff Implementation: 11/2003 Study Start: 00/0000 Study End: 00/0000 | Dispensing | Barcoding-dispensing Integrated Barcoding system, Pharmacy | Acute care/tertiary, 750 Beds Pharmacy, Inpatient hospital based, Academic | 3 barrier themes: (1) process (training requirements and process flow issues); (2) technology (hardware, software, and the role of vendors); (3) resistance (communication issues, changing roles, and negative perceptions about technology). | Bar code scanning system implementation is a difficult process with several barriers involving processes, technology and organizational resistance. Adequate training, continuous improvement, and adaptation of workflow to address one's own needs mitigated process barriers. Ongoing vendor involvement, acknowledgment of technology limitations, and attempts to address them were crucial in overcoming technology barriers. Staff resistance was addressed through clear communication, identifying champions, emphasizing new information provided by the system, and facilitating collaboration. |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|----------------------|---|---------------------------------------|---|---|
| <p>Novak (2008)³⁵² Design: Qualitative N = 50 hours of observations Implementation: 00/2007 Study Start: 00/0000 Study End: 00/0000</p> | <p>Administering</p> | <p>BCMA, e-Medication administration system (e-MAR, e-TAR) Integrated CPOE/POE system</p> | <p>Acute care/tertiary</p> | <p>For prior to BCMA implementation: Themes from the analytical coding were organized according to the nurses' practice goals-the familiar "Five Rights" of medication: Right Patient, Right Drugs, Right Dose, Right Time, and Right Way. For after BCMA implementation: In addition to the "Five Rights" of medication another theme emerged, namely, "New Articulation Work" and describes support and problem resolution strategies employed as nurses developed new coordination mechanisms.</p> | <p>The implementation of new information technology in the clinical setting can be disruptive to existing patterns of articulation work, or work that coordinates the activities of people across time and space. Implementation teams must familiarize themselves with articulation work and support users in developing new ways of coordinating with colleagues on other shifts and in remote physical spaces.</p> |
| <p>Novek (2000)³⁵³ Design: Mixed methods N = 124 Health care providers (mostly nurses and pharmacists) Implementation: 05/1997 Study Start: 02/1998 Study End: 00/0000</p> | <p>Dispensing</p> | <p>AMDs Integrated Pharmacy</p> | <p>Long term care (nursing homes)</p> | <p>Distrust, resistance, miscommunication, unrealistic expectations, speed and scale of implementation, concurrent changes, inadequate support, and social factors.</p> | <p>Nurses were generally distrustful of the AMDs and skeptical that it reduced medication errors.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|----------------------|---|---|--|--|
| <p>O'Grady (2006)³⁵⁴ Design: Qualitative N = 20 Patients Implementation: 06/2003 Study Start: 04/2003 Study End: 02/2004</p> | <p>Prescribing</p> | <p>e-Medication administration system (e-MAR, e-TAR) e-Rx Integrated Barcoding system, Ward-based automated dispensing system</p> | <p>Unspecified Hospital, 28 on the general surgery ward Beds</p> | <p>Themes (1) pre-EPA views: attitude about paper-based system was generally positive; (2) anticipated advantages of EPA before its introduction (save time, improve accuracy, and decrease mistakes); (3) the new system was expected to save time and be efficient (flexibility, comparisons with old system). (4) Concerns were shown over time, loss of personal touch, and not understanding the system; (5) advantage for staff when language is not English; (6) error reduction; (7) pre- EPA: inherent mistrust for computer systems; (8) post EPA: perceived disadvantages of the paper-based systems; (9) post EPA: perceived extra time needed if nursing staff had to check the drugs prescribed on the computer.</p> | <p>Patients generally had a good understanding of how paper-based system had worked and majority had safety concerns with it. Anticipated advantages were mostly about increased efficiency and reduced time. On balance, inpatients seemed neither for nor against EPA.</p> |
| <p>Patterson (2002)³⁵⁵ Design: Qualitative N = 33 nurses--7 before BCMA and 26 after Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Administering</p> | <p>BCMA Integrated EHR/EMR system</p> | <p>Acute care/tertiary, Other specialty hospital (rehab, oncology) 784 in the 4 settings Beds Long term care (nursing homes)</p> | <p>6 unanticipated side effects were noted: (1) confusion by automated removal of medications by BCMA; (2) degraded coordination between nurses and physicians; (3) dropping activities to reduce workload during busy periods; (4) increased prioritization of monitored activities during goal conflicts; (5) decreased ability to deviate from routine sequences; (6) to reduce workload wristbands were not scanned and medication scanning was delayed.</p> | <p>Unanticipated adverse effects happen and nurses find solutions to cope with workloads.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--|---|--|--|---|
| Patterson (2004) ³⁵⁶ Design: Qualitative N = 46 clinicians working in 6 primary care clinics Implementation: 00/0000 Study Start: 10/2001 Study End: 10/2002 | Monitoring including patient adherence and compliance, Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated Hospital information system, Laboratory system, Pharmacy | Ambulatory care | 7 barriers were identified, some of which were not on the original list: 1) workload; 2) time to document; 3) reminder did not apply; 4) inapplicability to the situation; 5) training lacks; 6) quality of provider- patient interaction; 7) use of paper forms. | Barriers exist. 17 recommendations were made to improve the situation: 9 related to design, 4 to the organization, and 1 each to team and role design, individual attitudes, patient and situation specific context, and interactions with other systems making issues redundant. |
| Pirnejad (2008) ²⁵⁶ Pirnejad (2009) ²⁵⁷ Design: Mixed methods N = 149 nurses Implementation: 09/2003 Study Start: 11/2003 Study End: 06/2007 | Prescribing | CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated EHR/EMR system, Hospital information system | Acute care/tertiary, 1237 Beds Academic | ²⁵⁶ The coding scheme included differentiation between those features that were considered supportive from features that were considered non- supportive to nurses' and physicians' medication work. Many of the paper- based system's non-supportive features were improved by the CPOE system. And, more useful features such as safety alerts and the possibility for physicians to prescribe electronically from everywhere in the hospital greatly benefited the prescription phase and improved the medication process. Nevertheless, nurses and physicians listed many non-supportive features of the CPOE system as well. ²⁵⁷ Workflow impediments from the perspective of physicians and nurses are described. The care providers devised compensatory work-arounds due to interoperabilities in the CPOE system. | ²⁵⁶ It is clear that moving from the paper based to the CPOE system had positive and negative impacts on nurses' and physicians' medication work. In our study, many of the CPOE system's non- supportive features were listed because the system damaged the synchronization and feedback mechanisms between nurses and physicians. ²⁵⁷ The interviews revealed that both nurses and physicians considered the system to be an improvement in their medication work compared to the old paper-based system. They complained about problems in coordination and collaboration. Problems forced them to develop informal rules and work methods to adapt the system in a way that it met their work requirements. |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|--|---|------------------------|---|---|
| <p>Ruiz (2010)³⁵⁷ Design: Qualitative N = 19 primary care practitioners Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/ reminders Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>Our data analysis elicited a number of themes, of which six are most relevant to the two areas of our inquiry. (1) pain as part of growing old; (2) concerns about using pain medications; (3) waiting times for pain clinic; (4) value of ancillary services; (5) poor training in pain management; (6) value of CPRS as a support tool.</p> | <p>The findings of this study clearly point to the need for a more systematic and solid understanding of the competencies of primary care practitioners in managing chronic nonmalignant pain in elderly veteran patients. While various types of support have been made available to primary care providers, competency-based training targeted on the elderly population must occur to facilitate the assessment and treatment of such pain. Particular attention must be given to the role of the EMR system as a source of clinical decision support complementary to and reinforcing competency-based training approaches.</p> |
| <p>Saleem (2005)³⁵⁸ Design: Qualitative N = 90 Healthcare providers Implementation: 00/0000 Study Start: 01/2004 Study End: 06/2004</p> | <p>Monitoring including patient adherence and compliance</p> | <p>CDSS/CDS/CCDS/ reminders Integrated CPOE/POE system, EHR/EMR system, Laboratory system</p> | <p>Ambulatory care</p> | <p>Five barriers, four of which have related subcategories, and four facilitators, organized by three themes: (1) organizational; (2) workflow; (3) computer interface. Barriers: (a) Lack of coordination between nurses and providers; (b) Using the reminders while not with the patient, impairing data acquisition and/or implementation of recommended actions; (c) Workload; (d) Lack of CR flexibility; (e) Poor interface usability. Facilitators: (a) Limiting the number of reminders at a site; (b) Strategic location of the computer workstations; (c) Integration of reminders into workflow; (d) Ability to document system problems and receive prompt administrator feedback.</p> | <p>Barriers might explain some of the variability in the use of CRs. These barriers may be difficult to overcome but some strategies may increase user acceptance and therefore the effectiveness of the CRs. These include explicitly assigning responsibility for each CR to nurses or providers, improving visibility of positive results from CRs in the electronic medical record, creating a feedback mechanism about CR use, and limiting the overall number of CRs.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|---|----------------------|---|---|--|--|
| <p>Schoville (2009)³⁵⁹ Design: Qualitative N = 58 nurses Implementation: 09/2007 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/ reminders CPOE/POE system Integrated EHR/EMR system, Imaging systems, Laboratory system, Pharmacy</p> | <p>Acute care/tertiary, Pediatric stand alone hospital, Pediatric Hospital: 184 beds; Women's Hospital: 40 Beds Academic</p> | <p>There were 5 types of work-arounds and artifacts identified by both nursing leadership and staff nurses: (1) workflow timing of events; (2) communication changes; (3) system problems; (4) learning curve of the CPOE system; (5) patient safety</p> | <p>Although CPOE is considered a technical solution to prevent or reduce errors and enhance communication among caregivers, errors could result because of the redundancy in documentation between the paper record and the EMR, systems not interfacing with one another, and multiple screens needing to be viewed to find information about the patient. It was verified that multiple variables affect a successful transition to an electronic order entry system and that workarounds and artifacts were used.</p> |
| <p>Topps, (2005)²⁷² Design: Mixed methods N = 313 Healthcare provider Implementation: 11/2002 Study Start: 05/2002 Study End: 06/2003</p> | <p>Administering</p> | <p>BCMA Integrated Billing/administration system Hospital information system, Pharmacy</p> | <p>Pediatric stand alone hospital</p> | <p>Qualitative</p> | <p>Themes derived from the pre-survey indicated that medications would be given in a timely manner with less error, but may result in an increase in time with increase in safety along with more reported errors, but fewer errors in administering actual medications (near misses). The surveys collected post-implementation indicated that the staff felt there were fewer medication errors with a smoother administration of medication; however, it was perceived that more time was spent administering medications taking time away from patient care.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--------------------|--|----------------------------------|--|---|
| <p>Varonen (2008)³⁶⁰ Design: Qualitative N = 39 physicians Implementation: 00/0000 Study Start: 10/2005 Study End: 12/2005</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/ reminders</p> | <p>Ambulatory care, Academic</p> | <p>Facilitating factors: Flexibility of the system; (tailoring the selection of topics or patients for reminders and possibility to switch off the system); Reliability; Reliable knowledge base and trust in the developers of the system; Simplicity and ease of use; Concise reminders that facilitate and help work processes; Adequate budgeting; Concise and tailored education for the use of CDSS barriers. In all groups, repeatedly: experience of imperfect health care information systems; Threats to doctor–patient relationship: the computer should not have the leading role in the encounter; Obscured responsibilities; loss of own reasoning and clinical autonomy; Knowledge management: too much information or erroneous information; Resistance towards change; Issues of compatibility and updating, problems with several poorly interacting computer programs</p> | <p>Finnish physicians interviewed in this qualitative study had positive attitudes towards implementation of CDSS provided that they have some control over the system. They expected flexibility, individual tailoring and reliability of the CDSS. The high level of computerized practices and wide use of electronic guidelines have paved the way for the CDSS in Finland.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--------------------|--|---|---|--|
| <p>Vaziri (2009)³⁶¹ Design: Qualitative N = >30 informaticians, academic clinicians, pharmacists, clinicians with an IT (information technology) interest, human factor/user experience consultants and medical and non- medical commercial IT vendors, as well as members of the National Health Service (NHS) national programme for IT development team Implementation: 00/000 Study Start: 00/2008 Study End: 00/2008</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/ reminders e-Rx Integrated EHR/EMR system</p> | <p>Ambulatory care</p> | <p>End-users (principally GPs) at the workshop reported that prescribing alerts were more often a source of frustration more than of help. Delegates reported concerns about the current prescribing support prompts, primarily the low specificity of the pop-ups, which were too numerous, often unhelpful and therefore ignored. Information overload may have a negative impact on cognitive performance.</p> | <p>Prescribing errors remain a major source of unnecessary morbidity and mortality and current systems do not appear to have significantly reduced this problem; nor has the extensive literature about how to reduce unnecessary alerts been taken into account. We need a new and more rational basis for the selection and presentation of alerts that would help, not hinder, the clinician's performance.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|----------------------|--|---------------------------------------|---|---|
| <p>Vogelsmeier (2008)³⁶² Design: Qualitative N = 88 nursing home staff Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Administering</p> | <p>e-Medication administration system (e-MAR, e-TAR) Integrated EHR/EMR system</p> | <p>Long term care (nursing homes)</p> | <p>Workarounds fell into 2 categories, relating to the technology itself and organizational processes. They occurred at new medication order entry, communication with the pharmacy and administration. The technology introduced intentional blocks (safety features such as excessive dose blocking, dual documentation and ADE monitoring) that lead to workarounds. Unintentional blocks leading to workarounds included wireless speed and printing each order on a separate page. Organization process blocks leading to workarounds included double checking of preparation and administration documents and limited resources such as fax machines.</p> | <p>As new technologies are introduced, continued monitoring to identify work flow is needed so appropriate changes can be made to address the underlying problems that create work flow blocks ultimately leading to potential workarounds. Additionally, as technology is implemented, organizational processes that will interface with the technology must be carefully re-engineered to reduce the unintended consequences of change.</p> |
| <p>Weingart (2009)³⁶³ Design: Qualitative N = 25 health care providers Implementation: 00/0000 Study Start: 00/2007 Study End: 00/2007</p> | <p>Prescribing</p> | <p>e-Rx Handheld Integrated, CDSS/CDS/CCDS/reminders Formulary, Pharmacy</p> | <p>Ambulatory care</p> | <p>problematic features: list management for patients; creating medication lists; poor recording of allergy information; awkward prescription writing leading to work-arounds; problematic alerts leading to alert fatigue</p> | <p>Front-line clinicians find many features of the e-Rx system burdensome. The value of e-Rx alerts is diminished by the quantity of irrelevant and inappropriate alerts. e-Rx triggers a variety of clinician behaviors (other than terminating or changing a prescription) that may improve patient safety.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--------------------|---|--|--|--|
| <p>Weir (1994)³⁶⁴ Design: Qualitative N = 40 hospital staff (admin, physicians, support staff etc) Implementation: 03/1993 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated Pharmacy</p> | <p>Unspecified Hospital</p> | <p>A survey requesting a list of 6-10 factors facilitating and 6-10 most significant barriers from staff at 3 hospitals with successful implementation and 3 with unsuccessful implementations of CPOE was analyzed using a modified Delphi technique. Fourteen facilitating factors and 14 barriers were identified. Several categories differentiated the two hospital groups. Significantly more people from the successful hospital group reported supportive administration and supportive heads of medical sections; direct involvement of physicians, mandatory implementation, adequate training, and sufficient hardware facilitated success. In terms of barriers, only inadequate hardware and lack of ability to easily do patient transfer and advance admission orders (medical records package) differentiated the two groups and in both cases the item was mentioned more frequently by the successful hospitals.</p> | <p>These findings support the notion that the changes involved in instituting a physician order entry system are system wide and involve individual as well as organizational factors.</p> |
| <p>weir (2007)³⁶⁵ Design: Qualitative N = 88 interviews Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated Hospital information system</p> | <p>Ambulatory care</p> | <p>Tasks were related to organization, assigning, determination, educating, scheduling, tracking, overview, correlating documenting, reminding, handing off, prioritizing, accepting, communicating, conforming, and informing. Task components were related to cueing, status, timing, communication, ownership, and linkage. Goals were associated with relevance screening, ensuring accuracy, minimizing memory load, and negotiating responsibility.</p> | <p>User creates strategies to learn how to effectively deal with new systems and processes, information overload must be carefully managed, and communication is vital and is often affected by new systems.</p> |

Evidence Table 10. KQ1: Primary qualitative outcomes for all technologies across phases (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results | Conclusions |
|--|--------------------|---|-----------------------------|--|---|
| <p>Wentzer (2007)³⁶⁶ Design: Qualitative N = 6 clinicians(2 physicians and 4 nurses) Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated EHR/EMR system</p> | <p>Acute care/tertiary</p> | <p>The study started with 3 relations (physician and patient interacting, physician and nurse coordination work, and the patients further route and medication path). Themes centered on transformation of the prescription and drug order during physician rounds, transformation of the drug order and dispensing with the CPOE system (user rights, inflexibilities and displacements with the use of CPOE, going back to the paper system, unified and inflexible CPOE medication model), transformation of continuing medication with the system (discharge, withdrawal or discontinuous patient routes, and new tasks and demands on the clinicians with the CPOE system).</p> | <p>CPOE system did not meet naive and early expectations. Some adverse effects of the CPOE system were noted.</p> |
| <p>Zhan (2006)¹⁸¹ Design: Mixed methods N = 138,922 number of errors/100,000 doses Implementation: 00/0000 Study Start: 01/2003 Study End: 12/2003</p> | <p>Prescribing</p> | <p>CPOE/POE system</p> | <p>Unspecified Hospital</p> | <p>Some of the themes taken from the CPOE-related error descriptions included: faulty computer interface, CPOE design failures, especially lack of connection with other parallel systems, inadequacy of decision support and human errors occurring in interactions with the computer.</p> | <p>A national, voluntary medication error-reporting database cannot be used to determine the effectiveness of a CPOE system in reducing medication errors because of the variability in the level of underreporting from different institutions. However, it may provide valuable and useful information on the specific types of errors related to CPOE systems.</p> |

Evidence Table 11. KQ1: primary composite/population level outcomes for all technologies assisting education or other phase

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Outcomes Measured | Results | Outcome |
|--|---|---|--------------------------------|-------------------|---|---------|
| Holbrook (2009) ³⁶⁷ Design: RCT N = 511 patients Implementation: 00/0000 Study Start: 00/2002 Study End: 12/2003 | Monitoring including patient adherence and compliance | CDSS/CDS/CCDS/ reminders Integrated Laboratory system | Ambulatory care, Home | Composite | A shared electronic decision-support system to support the primary care of diabetes improved the process of care and some clinical markers of the quality of diabetes care. | + |
| Yu (2009) ³¹⁷ Design: Case control N = 22,665 patients Implementation: 00/0000 Study Start: 10/2005 Study End: 09/2006 | Prescribing | CPOE/POE system Integrated Imaging systems, Laboratory system, Pharmacy | Pediatric stand alone hospital | Population Level | Using actual reportable ADEs from a relatively large number of pediatric hospitals, the study found significant beneficial associations between reportable ADE and CPOE implementation. | + |

The HIT system studied is in **bold**; followed by the systems that it was integrated with. The outcome column indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: ADE = Adverse Drug Event; CCDS = Computerized Clinical Decision Support; CDS = Clinical / Computerized Decision Support; CDSS = Clinical Decision Support System; CPOE = Computerized Provider Order Entry; HIT = Health Information Technology; N = Sample Size; POE = Provider Order Entry; RCT = Randomized Controlled Trial

Evidence Table 12. KQ1: adverse effects measured for all technologies assisting all phases of medication management

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results |
|--|--------------|---|---|---|
| Ash (2007) ³⁶⁸ Design: Observational study N = 95 clinicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CPOE/POE system Integrated CDSS/CDS/CCDS/reminders EHR/EMR system | Acute care/tertiary, General Hospital, Academic | 47 examples of unintended consequences of CDS systems were observed. Thematic analysis showed 2 major patterns: generation by the content or presentation of the information. Issues related to content centered around elimination or shifting of human roles, currency of the CDS content or wrong or misleading CDS content. Issues related to the presentation centered around rigidity of the system, alert fatigue, sources of potential errors. |
| Ash (2007) ³⁶⁹ Design: Survey N = 176 hospitals Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CPOE/POE system | Acute care/tertiary | The preliminary qualitative study identified major types of UAC of CPOE. ³⁷⁰ The survey results verified the existence of eight UAC asked about at most of the 176 hospitals with CPOE. All types of UAC are widespread although two of them, power shifts and new kinds of errors, were not considered as important as the others (more/new work, workflow, system demands, communication, emotions, dependence on technology). |
| Ash (2007) ³⁷⁰ Design: Qualitative N = 4 outpatient clinics Implementation: 00/1997 Study Start: 04/2003 Study End: 10/2003 | Prescribing, | CPOE/POE system | Ambulatory care, Not specified | Using the diffusion of innovations model, unintended sociotechnical consequences of CPOE were analyzed along 3 classifications: desirable vs. undesirable; direct vs. indirect; and anticipated vs. unanticipated for a total of 8 possible combinations. Examples of each combination were found in the narrative data. There were error and security concerns, and issues related to alerts, workflow, ergonomics, interpersonal relations, and reimplementation. |

The HIT system studied is in **bold**, followed by the systems that it was integrated with.

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: BCMA= Bar Code Medication Administration, CCDS= Computerized Clinical Decision Support, CDS= Clinical / Computerized Decision Support, CDSS= Clinical Decision Support System, CI=confidence Interval CPOE= Computerized Provider Order Entry, EHR= Electronic Health Record, e-MAR= Electronic Medication Administration Record, EMR= Electronic Medical Records, e-TAR= Electronic Treatment Authorization Request, HIT= Health Information Technology, HMO= Health Maintenance Organization, ICU= Intensive Care Unit, MM= Medication Management, MMC= Montefiore Medical Center, N= Sample Size, p = probability, POE= Provider Order Entry, RCT= Randomized Controlled Trial, RRR Relative Risk Reduction, UACs= Unintended Adverse Consequences, vs.= versus, WA= Workarounds

Evidence Table 12. KQ1: adverse effects measured for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results |
|---|--------------------|---|---|---|
| <p>Campbell (2006)³⁷¹ Campbell (2006)³⁷² Ash (2006)³⁷³ Campbell (2009)³⁷⁴ Design: Qualitative N = 95 clinicians (various) Implementation: 00/0000 Study Start: 09/2004 Study End: 04/2005</p> | <p>Prescribing</p> | <p>CPOE/POE system</p> | <p>Acute care/tertiary, General Hospital, Pharmacy, Inpatient hospital based, Academic</p> | <p>³⁷¹ UACs fell into nine major categories (in order of decreasing frequency): (1) more work for clinicians; (2) unfavorable workflow issues; (3) never ending system demands; (4) problems related to paper persistence; (5) untoward changes in communication patterns and practices; (6) negative emotions; (7) generation of new kinds of errors; (8) unexpected changes in the power structure; and (9) overdependence on the technology. Clinical decision support features introduced many of these unintended consequences.³⁷⁴ CPOE systems can affect clinical work by: (1) introducing or exposing human/computer interaction problems; (2) altering the pace, sequencing, and dynamics of clinical activities; (3) providing only partial support for the work activities of all types of clinical personnel; (4) reducing clinical situation awareness; and (5) poorly reflecting organizational policy and procedure.³⁷² Careful analysis of overdependence on technology data revealed 3 themes: (1) system downtime can create chaos when there are insufficient backup systems in place; (2) users have false expectations regarding data accuracy and processing; and 3) some clinicians cannot work efficiently without computerized systems.³⁷³ CPOE enables shifts in power related to work redistribution and safety initiatives and causes a perceived loss of control and autonomy by clinicians.</p> |
| <p>Campbell (2009)³³⁴ Design: Qualitative N = 32 semi-structured interviews=43 hours; 400 hours of observation shadowing 95 clinical providers Implementation: 00/0000 Study Start: 08/2004 Study End: 04/2005</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated EHR/EMR system</p> | <p>Acute care/tertiary, General Hospital, 340 (Wishard); 893 (Massachusetts); 150 (Faulkner); 725 (Brigham); 238 (Alamance) Beds Academic</p> | <p>To identify and describe unintended adverse consequences related to clinical workflow when implementing or using computerized provider order entry (CPOE) systems we analyzed qualitative data from field observations and formal interviews gathered over a three-year period at five hospitals in three organizations. Five multidisciplinary researchers worked together to identify themes related to the impacts of CPOE systems on clinical workflow.</p> |

Evidence Table 12. KQ1: adverse effects measured for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results |
|--|--|--|--|--|
| Han (2005) ³⁰³ Design: Before-after N = 1,942 patients Implementation: 10/2002 Study Start: 10/2001 Study End: 03/2003 | Dispensing, Transmission, order communication | CDSS/CDS/CCDS/reminders CPOE/POE system Integrated Hospital information system | Acute care/tertiary, Pediatric stand alone hospital, Academic | Mortality increased among patients transported in to the tertiary care pediatric center following the implementation of the CPOE system (2.8% vs. 6.6%, RRR -135%, p<0.001) as a result of delays in entering and processing orders and changes to workflow and communication among staff. |
| Keene (2007) ³⁰⁸ Design: Before-after N = 1,291 patients Implementation: 00/2001 Study Start: 09/2000 Study End: 02/2003 | Prescribing | CPOE/POE system Integrated EHR/EMR system, Laboratory system, Pharmacy | Critical care units (CCU, ICU, NICU) Academic | The initiation of CPOE for the pediatric critically ill at MMC took place without the increase in mortality reported during a similar initiation period by Han and colleagues. Careful preparation, unit-by-unit tailoring, and extensive technical support may have improved the results at MMC. |
| Koppel (2005) ³⁴⁵ Design: Mixed methods N = 291 health care providers Implementation: 00/1997 Study Start: 00/2002 Study End: 00/2003 | Prescribing | CPOE/POE system Integrated nurses medication lists, Pharmacy | Acute care/tertiary, 750 Beds Academic | a widely used CPOE system facilitated 22 types of medication error risks. Examples include fragmented CPOE displays that prevent a coherent view of patients' medications, pharmacy inventory displays mistaken for dosage guidelines, ignored antibiotic renewal notices placed on paper charts rather than in the CPOE system, separation of functions that facilitate double dosing and incompatible orders, and inflexible ordering formats generating wrong orders. Three quarters of the house staff reported observing each of these error risks, indicating that they occur weekly or more often. Use of multiple qualitative and survey methods identified and quantified error risks not previously considered, offering many opportunities for error reduction. |
| Lin (2008) ⁹⁵ Design: Time series N = 1,123 high severity order checks Implementation: 00/1997 Study Start: 01/2001 Study End: 01/2006 | Prescribing | CDSS/CDS/CCDS/reminders CPOE/POE system Integrated CPOE/POE system, EHR/EMR system | Acute care/tertiary, General Hospital, 444 Beds Ambulatory care, Long term care (nursing homes) | There were 215 high severity order checks in 2001 (0.5% of orders) and 908 in 2006 (2.5% of orders). Rate of overrides for drug-drug checks remained the same between 2001 and 2006 (88% vs. 87%, NS). Rate of overrides for drug-allergy order checks increased significantly from 2001 to 2006 (69% vs. 81%, RRR - 17%, p<0.005). Override rates remain high and drug-allergy override rates increased. |

Evidence Table 12. KQ1: adverse effects measured for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results |
|---|---|--|--|---|
| <p>Patterson (2002)³⁵⁵ Design: Qualitative N = 33 nurses--7 before BCMA and 26 after Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Administering</p> | <p>Barcoding-medication administering Integrated EHR/EMR system</p> | <p>Acute care/tertiary, Other specialty hospital (rehab, oncology) 784 in the 4 settings Beds Long term care (nursing homes)</p> | <p>6 unanticipated side effects were noted: confusion by automated removal of medications by BCMA, degraded coordination between nurses and physicians, dropping activities to reduce workload during busy periods. Increased prioritization of monitored activities during goal conflicts, decreased ability to deviate from routine sequences. To reduce workload, wristbands were not scanned and medication scanning was delayed.</p> |
| <p>Raebel (2007)¹³¹ Design: RCT N = 11,100 women Implementation: 00/0000 Study Start: 01/2003 Study End: 04/2003</p> | <p>Prescribing</p> | <p>CDSS/CDS/CCDS/reminders Integrated Hospital information system, Pharmacy</p> | <p>Ambulatory care, HMO pharmacy</p> | <p>Although the study intervention was successful at decreasing the proportion of pregnant women with contraindicated drug dispensings, the study intervention was stopped after 4 of the planned 12 months. The 2 predominant factors contributing to the decision to end the intervention were the false-positive alerts resulting from misidentification of medications as contraindicated in pregnancy by the pharmacy information system and misidentification of pregnancy related to delayed transfer of diagnosis information.</p> |
| <p>Santell (2009)³⁷⁵ Design: Observational study N = 90,876 potential errors Implementation: 00/1998 Study Start: 07/2001 Study End: 12/2005</p> | <p>Administering, Dispensing, Monitoring including patient adherence and compliance, Prescribing, Transmission, order communication</p> | <p>CPOE/POE system, Medication-error reporting system</p> | <p>Acute care/tertiary, Academic</p> | <p>The analysis of the national database focused on errors by non-prescribers resulting from CPOE. Errors generally occurred during dispensing (50.9%) and transcribing or documenting (42.5%). Errors tended to be improper dose or quantity (32.5%), omissions (22.2%), or unauthorized/wrong drug (14.35). Causes of errors included performance deficits (59.1%), inaccurate transcriptions (30.0%), procedure or protocol not followed (21.7%), documentation (19.5) and communication (18.3%). 62.2% of errors did not reach the patient. Fairly similar patterns were observed at University of Pittsburgh Medical Center.</p> |

Evidence Table 12. KQ1: adverse effects measured for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results |
|--|--------------------|---|---|--|
| <p>Shulman (2005)¹⁵⁰ Design: Time series N = 3,465 prescriptions over 4 time points Implementation: 04/2002 Study Start: 09/2001 Study End: 12/2002</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated Hospital information system</p> | <p>Critical care units (CCU, ICU, NICU) 22 (in the ICU) Beds Academic</p> | <p>Three intercepted errors with CPOE could have caused permanent harm or death if they had been administered as prescribed. These intercepted errors were not administered to the patient because either the pharmacist intercepted the prescription before administration or the nurse recognized the error.</p> |
| <p>Singh (2009)³⁷⁶ Design: Cross-sectional N = 997 prescriptions Implementation: 00/1998 Study Start: 10/2007 Study End: 01/2008</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated CDSS/CDS/CCDS/reminders EHR/EMR system, Pharmacy</p> | <p>Acute care/tertiary, Pharmacy, Inpatient hospital based</p> | <p>Of 55,992 new prescriptions, 532 (0.95%) were reported to contain inconsistent communication (control prescriptions = 465), a rate comparable to that obtained from the unreported group. Drug dosage was the most common inconsistent element among both groups. Certain medications were more likely associated with errors, as was the inpatient setting (OR 3.30; 95% CI 2.18 to 5.00) and surgical subspecialty (OR 2.45; 95% CI 1.57 to 3.82). About 20% of errors could have resulted in moderate to severe harm, for which significant independent predictors were found. Despite standardization of data entry, inconsistent communication in CPOE poses a significant risk to safety.</p> |
| <p>Spencer (2005)¹⁵⁶ Design: Before-after N = 5,063 medication errors Implementation: 10/2002 Study Start: 01/2002 Study End: 05/2003</p> | <p>Prescribing</p> | <p>CPOE/POE system Integrated</p> | <p>Acute care/tertiary, 688 Beds Academic</p> | <p>A total of 23 reported errors were attributable to errors in computerized order entry after implementation of CPOE. Of these, 6 occurred during the calendar month when CPOE was implemented on the given unit and are therefore not accounted for in Table 3. All of these errors were classified as minor, with 14 (61%) constituting only potential errors. Twenty-one errors in computerized order entry (91%) were of severity category 1 or lower.</p> |

Evidence Table 12. KQ1: adverse effects measured for all technologies assisting all phases of medication management (continued)

| Article Information | MM Phase(s) | HIT studied Integrated systems | Settings | Results |
|---|---------------|--|--------------------------------|---|
| Vogelsmeier (2008) ³⁶² Design: Qualitative N = 88 nursing home staff Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Administering | e-Medication administration system (e-MAR, e-TAR) Integrated EHR/EMR system | Long term care (nursing homes) | WA presented in two distinct patterns: WA related to blocks introduced by technology and WA related to organizational processes that had not been reengineered to integrate effectively with the implementation of technology. WA resulted as nursing home staff attempted to individually problem solve how to overcome a work flow block rather than seeking to identify and understand the underlying cause of the work flow block. WA frequently circumvented the built in security features of the system. |
| Wentzer (2007) ³⁶⁶ Design: Qualitative N = 6 clinicians(2 physicians and 4 nurses) Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000 | Prescribing | CPOE/POE system Integrated EHR/EMR system | Acute care/tertiary | The study started with 3 relations (physician and patient interacting, physician and nurse coordination work, and the patients further route and medication path). Themes centered on transformation of the prescription and drug order during physician rounds, transformation of the drug order and dispensing with the CPOE system (user rights, inflexibilities and displacements with the use of CPOE, going back to the paper system, unified and inflexible CPOE medication model), transformation of continuing medication with the system (discharge, withdrawal or discontinuous patient routes, and new tasks and demands on the clinicians with the CPOE system). |
| Zhan (2006) ¹⁸¹ Design: Mixed methods N = 138,922 number of errors/100000 doses Implementation: 00/0000 Study Start: 01/2003 Study End: 12/2003 | Prescribing | CPOE/POE system | Unspecified Hospital | There were 7,029 CPOE related errors reported from May to December 2003. Most were potential errors, about 4.7 reached patients, 0.1% inflicted temporary harm. Error types, causes and contributing factors were further described. |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality

| PICOM [^] | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|--|-------------------------|----------------|-----------------------------------|-----------------------|
| An RCT of hospitalized patients with coronary artery diseases. To asses whether a computerized alert identifying patients with elevated troponin I levels to pharmacists who mediated academic detailing was effective in increasing adherence to secondary prevention guidelines for coronary artery disease. 895 patients were considered eligible for the study and 216 discharge physicians were involved. | Bailey (2007) ⁷ Design: RCT N = 853 patients Implementation: 00/0000 Study Start: 02/2000 Study End: 05/2001 | Acute care/ tertiary | Academic | 16 months | 3 |

[^]PICOM = description of patient, intervention, comparator, outcome and method

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: AMT = Antimicrobial Management Team; ATP III = Third Adult Treatment Panel; BMD = Bone Mineral Density; BP = Blood Pressure; CAD = Coronary Artery Disease; CCU = Critical Care Unit; CDS = Clinical / Computerized Decision Support; CDSS = Clinical Decision Support System; CIS = Clinical Information System; COPD = Chronic Obstructive Pulmonary Disease; CPOE = Computerized Provider Order Entry; CRS = Computer Reminder System; DCGP = Dutch College of General Practitioners; DHCP = Decentralized Hospital Computer System; DMA = Disease Management Application; DS = Decision Support; DSS = Decision Support System; DVT = Deep Vein Thrombosis; ED = Emergency Department; HER = Electronic Health Record; EMR = Electronic Medical Records; FEV1 = Forced Expiratory Volume in the first second; FU = Followup; GINA = Global Initiative for Asthma; GMC = General Medical Clinic; GPs = General Practitioners; HbA1c = Glycated hemoglobin; HCP = Health Care Provider; HF = Heart Failure; HIV = Human Immunodeficiency Virus; HMO = Health Maintenance Organization; ICU = Intensive Care Unit; IHD = Ischemic Heart Disease; JNC-7 = Seventh Report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; LDL = Lowdensity Lipoprotein; MDI = Microbiologically Documented Infections; N = Sample Size; NAEP = National Asthma Education Program; NICU = Neonatal Intensive Care Unit; NSAID = Nonsteroidal anti-inflammatory drug; OA = Oral Anticoagulant; PCC = Pediatric Care Center; PE = Pulmonary Embolism; PEF = Peak Expiratory Flow; PICOM = description of patient, intervention, comparator, outcome and method; POC = Point of Care; PRISM = Prescription in Ischaemic Stroke Management; PWS = Physician Workstation; RAMQ = Régie de l'assurance maladie du Québec; RCT = Randomized Controlled Trial; RRR = Relative Risk Reduction; SADC = System of Clinical Decision Support; SGRQ = St. Georges Respiratory Questionnaire; SOC = Standards of Care; SYW = show your work; TIA = Transient Ischemic Attack; USFDA = United States Food and Drug Administration; VA = Veterans Affairs; VTE = Venous thromboembolism

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|------------------|--|-----------------------------------|-----------------------|
| <p>A prospective cluster-randomized trial was conducted in 12 primary care sites within the Children’s Hospital of Philadelphia over a 1-year period. Practices were stratified for analysis according to whether the site was urban or suburban. A clinical decision support (CDS) embedded in an electronic health record (EHR) to improve clinician adherence to the NAEPP guidelines for asthma management was assessed in the intervention group. Control group had passive access to the same asthma management tools. Proportion of children with persistent asthma with at least 1 prescription for a controller medication in each time period; with persistent asthma with an up-to-date asthma care plan filed in the previous year; with documentation of spirometry performed were measured and compared. To balance practices with previous asthma education or involvement in resident teaching and patient characteristics, the practices were stratified according to site (urban UP or suburban SP) in blocks of 2. Therefore, 4 clusters of practices were compared in the analysis: 2 control UPs, 2 intervention UPs, 4 control SPs, and 4 intervention SPs. 19,450 children with asthma were included in the analysis.</p> | <p>Bell (2010)¹¹ Design: RCT N = 19,450 patients Implementation: 00/0000 Study Start: 04/2007 Study End: 04/2008</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>12 months</p> | <p>4</p> |
| <p>An RCT in a university-based resident clinic on 59 internal medicine residents using standardized patients to assess the impact of a PDA-based CDSS for safe prescribing of NSAIDs. Prescriptions were judged as safe or unsafe. The main outcome measure was the differential change in unsafe prescribing of NSAIDs for the intervention versus the control group. Both groups received PDAs with DS rule sets, the intervention group receiving the NSAID set following a 6 month baseline assessment period.</p> | <p>Berner (2006)¹² Design: RCT N = 59 internal medicine residents Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>8 months</p> | <p>5</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|---|------------------|--|-----------------------------------|-----------------------|
| <p>29 practices received the Third Adult Treatment Panel (ATP III) intervention and 32 receiving an alternative intervention focused on the Seventh Report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC-7). The ATP III providers received a personal digital assistant providing the Framingham risk scores and ATP III–recommended treatment. All practices received copies of each clinical practice guideline, an introductory lecture, 1 performance feedback report, and 4 visits for intervention specific academic detailing. Data were abstracted at 61 practices from random samples of medical records of patients treated from June 1 2001, through May 31 2003 (baseline), and from May 1 2004, through April 30 2006 (follow-up). Effect on screening of lipid levels and appropriate management of lipid level test results were compared for 8,878 patients.</p> | <p>Bertoni (2009)¹⁴ Design: RCT N = 8,878 patients Implementation: 00/0000 Study Start: 06/2001 Study End: 04/2006</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>24 months</p> | <p>5</p> |
| <p>This RCT was to determine the effectiveness of three different prompts, 92 providers from 5 intervention primary care clinics were randomized to receive one of the three prompts and compared to 6 control clinics. Prompts included letters sent to patients about lipid therapy prior to their scheduled visit, a progress note message within the computerized patient record system notifications area and a computerized reminder screen within the specific patient chart during the patient’s visit.</p> | <p>Bloomfield (2005)¹⁵ Design: RCT N = 9,105 patients Implementation: 04/2002 Study Start: 10/2001 Study End: 10/2003</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>2</p> |
| <p>A RCT in a pediatric care clinic of a point of care (POC) evidenced based message system with real time evidence to providers based on their prescribing practice for otitis media. Compared change in prescribing behavior of the intervention and control providers before and after implementation of the message pop-up. Prescribing behavior change was measured as the change in the proportion of prescriptions of antibiotics for less than 10 days duration from baseline. The study included 38 providers and 1,339 visits for acute otitis media.</p> | <p>Christakis (2001)²³ Design: RCT N = 38 providers Implementation: 00/0000 Study Start: 03/0000 Study End: 05/0000</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>8 months</p> | <p>4</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|--|------------------|--|-----------------------------------|-----------------------|
| <p>This study was conducted to assess the impact in effectiveness and direct costs of a CDSS to assist physicians in the implementation of an adapted version of the recommendations of the European Society of Cardiology and other societies for Hypercholesterolemia Management (ESCHM). 2,221 patients were included from 42 practices in the cluster-randomized controlled trial comparing lipid profile, cardiovascular risk, use of lipid lowering drugs and costs.</p> | <p>Cobos (2005)²⁵ Design: RCT N = 2,221 patients Implementation: 04/2000 Study Start: 04/2000 Study End: 05/2002</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>5</p> |
| <p>This was a cluster randomized clinical trial of provider behavior change. Prescribing behavior was measured in both the intervention and control groups before and after the introduction in the intervention group of a pop-up DSS alert providing evidence at the time of electronic prescribing. The conditions included in the intervention were acute otitis media, allergic rhinitis, sinusitis, constipation, pharyngitis, croup, urticaria, and bronchiolitis. In this study the unit of intervention was the provider. This study was conducted at two clinical sites. One was the Pediatric Care Center (n = 36 Health Care Providers), an outpatient teaching clinic for pediatric residents and a clinical practice site staffed by full-time pediatric providers. The other site was Skagit Pediatrics (n = 8 HCP), a primary care pediatric clinic serving a rural and semi-urban patient mix.</p> | <p>Davis (2007)³⁰ Design: RCT N = 44 health care providers Implementation: 11/1999 Study Start: 11/1999 Study End: 12/2003</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>18 to 50 months</p> | <p>6</p> |
| <p>To examine whether a computerized reminder system operating in multiple Veterans Affairs (VA) ambulatory care clinics improves resident physician compliance with standards of ambulatory care an RCT was undertaken. A total of 275 resident physicians at 12 VA medical centers were randomly assigned in firms or half-day clinic blocks to either a reminder group (n=132) or a control group (n=143). During a 17-month study period (January 31 1995 to June 30 1996), the residents cared for 12,989 unique patients for whom at least 1 of the studied standards of care (SOC) was applicable. Compliance with 13 SOC, were compared, 5 relating to medication management. The reminders were presented to intervention residents in the electronic chart in the examination room and a paper copy was put into the patient paper chart with the standard health summaries printed at each clinic visit. Control residents continued to receive the health summaries.</p> | <p>Demakis (2000)²¹³ Design: RCT N = 12,989 patients Implementation: 00/0000 Study Start: 01/1995 Study End: 06/1996</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>17 months</p> | <p>2</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|-----------------------------|-----------------|-----------------------------------|-----------------------|
| <p>An RCT of all hospitalized patients during an 18-month period, to determine the effects of computerized reminders vs. no reminders on the rates of compliance with 4 preventative therapies; national guidelines for the use of pneumococcal vaccination, influenza vaccination, prophylactic enteric-coated aspirin for cardiovascular disease and prophylactic subcutaneous heparin to reduce the risk of thromboembolic events. The reminder system identified 3,416 patients (53.6%) as eligible for preventive measures.</p> | <p>Dexter (2001)³³ Design: RCT N = 3,416 patients Implementation: 00/0000 Study Start: 05/1997 Study End: 10/1998</p> | <p>General Hospital</p> | <p>Academic</p> | <p>18 months</p> | <p>4</p> |
| <p>To determine the effects of computerized physician standing orders compared with physician reminders on inpatient vaccination rates. Randomized trial of 3,777 general medicine patients discharged from 1 of 6 study wards during a 14-month period (November 1, 1998, through December 31, 1999) composed of 2 overlapping influenza seasons at an urban public teaching hospital. The intervention was the use of the hospital CPOE to either, for patients with standing orders, automatically produce vaccine orders directed to nurses at the time of patient discharge or, for patients with reminders, provide reminders to physicians that included vaccine orders during routine order entry sessions. The main outcome measure was vaccine administration.</p> | <p>Dexter (2004)³⁴ Design: RCT N = 1,677 patients Implementation: 11/1998 Study Start: 11/1998 Study End: 12/1999</p> | <p>General Hospital</p> | <p>Academic</p> | <p>3 days</p> | <p>4</p> |
| <p>A two-stage, random-selection study to develop and evaluate appropriate empiric antibiotics in tertiary care Hospital in Salt Lake City. Antibiotics suggested by the antibiotic consultant with 482 associated antibiotic susceptibility results and the concurrent antibiotics ordered by physicians were compared. The antibiotics ordered by randomized physicians were then compared between crossover periods of antibiotic consultant use.</p> | <p>Evans (1994)³⁹ Design: RCT N = 482 cultures Implementation: 00/000 Study Start: 07/1990 Study End: 01/1991</p> | <p>Acute care/ tertiary</p> | | <p>9 months</p> | <p>2</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|--|------------------|------------------------|-----------------------------------|-----------------------|
| <p>This study evaluated interventions to improve laboratory monitoring at initiation of medication therapy. This cluster-randomized trial compared 3 interventions to usual care for 10 medications in 15 primary care clinics in a health maintenance organization with an electronic medical record system. Eligible patients, identified from electronic databases, had not received recommended laboratory monitoring within 5 days after new dispensing of a study medication. Interventions were an electronic medical record reminder to the prescribing health care professional, an automated voice message to the patient, and a pharmacy team outreach to the patient. Primary outcome was completion of all recommended baseline laboratory monitoring. 961 patients were included. Direct HMO costs were calculated (repeat testing, extra visits, intervention costs) using trial data and external sources.</p> | <p>Feldstein (2006)²¹⁵ Smith (2009)²¹⁶ Design: RCT N = 961 patients Implementation: 09/2004 Study Start: 09/2003 Study End: 01/2005</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>4.5 months</p> | <p>7</p> |
| <p>An interrupted time series RCT was performed at 15 primary care clinics including 239 primary care providers and 9,910 patients taking warfarin. EMR alerts and group academic detailing were implemented to reduce the co-prescribing of warfarin and 5 interacting medications. Physicians could continue with the prescription, change the medication or select from options presented. The academic detailing included group educational session. The unit of randomization was the primary care clinic; the unit of intervention was the primary care provider; and the unit of analysis was time (study month). The primary outcome was the "interacting prescription rate," defined as the number of co-prescriptions of warfarin-interacting medications per 10,000 warfarin users per month. The effect of the interventions was evaluated using an interrupted time series design, analyzed with segmented regression models that control for pre-intervention trends.</p> | <p>Feldstein (2006)⁴⁰ Design: RCT N = 9,910 patients with 239 care providers in 15 primary care clinics Implementation: 12/2002 Study Start: 01/2000 Study End: 08/2004</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>6</p> |
| <p>A 3-arm RCT of women needing osteoporosis care after a fracture. Women were allocated to usual care, reminder letters and EMR notes alone or reminders plus patient education and related information. Outcomes were obtaining BMD measurement or starting medication or both by the end of the 6-month trial. Care recommendations were based on guidelines.</p> | <p>Feldstein (2006)⁴¹ Design: RCT N = 311 women Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>6 months</p> | <p>5</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|---|------------------|-------------------------------------|-----------------------------------|-----------------------|
| <p>Cluster RCT of 22 long stay units in long term care settings was done to determine if CPOE with CDSS improved prescribing for antidepressants in patients with renal insufficiency (94 alerts related to 62 drugs). Alerts centered on maximum daily doses or frequencies, medications to be avoided and missing values for creatinine clearance. Outcomes were the proportion of alerts that lead to appropriate drug orders and rates of inappropriate drugs avoided. 10 physicians and 833 patients (213,967 patient days) were studied.</p> | <p>Field (2009)⁴² Design: RCT N = 833 patients (10 physicians and 213,967 patient days) Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Long term care</p> | <p>12 months</p> | <p>5</p> |
| <p>This study was conducted at the first 20 practices in the Children's Hospital of Philadelphia Pediatric Research Consortium that implemented the ambulatory EHR EpicCare. A prospective, 20-primary care site, cluster-randomized, decision-support trial between October 1, 2006, and March 31 2007 was conducted. At intervention sites, electronic health record-based clinical alerts for influenza vaccine appeared at all office visits for children between 5 and 19 years of age with asthma who were due for vaccine. For each site, captured opportunities for influenza vaccination and influenza vaccination rates were compared with those for the same period in the previous year.</p> | <p>Fiks (2009)⁴³ Design: RCT N = 22,586 patients Implementation: 00/0000 Study Start: 10/2006 Study End: 05/2007</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>6 months</p> | <p>3</p> |
| <p>A RCT among general practitioners (GPs) in Italy. An electronic reminder was put in a standard software system for patient data management to remind GPs to prescribe anti-platelet drugs in diabetic patients who were at high risk of developing cardiovascular disease. A letter summarizing the beneficial effects of anti-platelet drugs in such type of patients were given to both the intervention and the control group. Patients were classified into 3 risk groups. Data for patients receiving anti-platelet drug treatment in the control and the intervention group at the baseline and at the follow-up among the three risk groups were analyzed. 300 GPs and 15,343 high-risk diabetic patients were involved in the study.</p> | <p>Filippi (2003)⁴⁴ Design: RCT N = 15,343 patients Implementation: 00/0000 Study Start: 05/2001 Study End: 11/2001</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>7 months</p> | <p>4</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|--|------------------|--|-----------------------------------|-----------------------|
| <p>A clustered RCT was conducted at general practices in Norway. the Winmed electronic medical record system was used by the practices. Computer based DS and reminders were implemented based on evidence based guidelines for urinary tract infection or sore throat. Changes in rates of ordering of antibiotics were compared between the intervention and the control group for sore throat and urinary tract infection.</p> | <p>Flottorp, (2002)⁴⁷ Design: RCT N = 26,826 Consultation Implementation: 00/0000 Study Start: 01/2000 Study End: 01/2001</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>4.5 months</p> | <p>2</p> |
| <p>To evaluate the effectiveness of computerized prescribing alerts, with or without physician-led group educational sessions, compared to usual care, to reduce the prescribing of heavily marketed hypnotic medications. 14 internal medicine practice sites were randomly allocated to receive usual care, computerized prescribing alerts alone, or alerts plus group educational sessions. Proportion of heavily marketed hypnotics prescribed before and after the implementation of computerized alerts and educational sessions were compared. Usual care included an alert of the copayment tier of the medication; the computer alerts recommended generic brands; group education sessions were held at 4 sites and an educational information packet was sent to all internal medicine clinicians from those sites.</p> | <p>Fortuna (2009)⁴⁸ Design: RCT N = 257 clinicians Implementation: 00/1997 Study Start: 03/2006 Study End: 03/2008</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>12 months</p> | <p>6</p> |
| <p>An RCT was conducted to determine whether the combination of a computer- generated and written reminder system provided during patient visits could increase patient receipt of aspirin, beta-blockers, and cholesterol-lowering agents in patients with CAD. Physicians were randomly assigned to either a control group or an intervention group. The intervention group received computerized and written reminders for their patients with coronary artery disease, whereas those assigned to the control group were not contacted. Proportion of patients who had an active prescription for aspirin; the proportion of patients with myocardial infarction who had an active beta-blocker prescription; the proportion of patients receiving a cholesterol-lowering agent; and the proportion of patients with a level of low-density lipoprotein (LDL) cholesterol in the desired range (< 100 mg/dL) were evaluated and compared between the control and intervention group. 730 patients and 63 physicians were involved in the study. Sample size adjusted for clustering.</p> | <p>Frances (2001)⁴⁹ Design: RCT N = 63 physicians and 730 patients Implementation: 00/0000 Study Start: 03/1997 Study End: 06/1997</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>3</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|------------------|------------------------|-----------------------------------|-----------------------|
| <p>A quasi-randomized trial within a primary clinic with 10 physicians looking at the effectiveness of in consultation computer reminders about 12 outstanding preventive care activities. Patients were the unit of randomization; 5,118 in the intervention group and 5,389 in the control group. Reminders appeared on the medical record screen and pertained to 4 vaccine reminders and 8 non-medication related preventive care recommendations.</p> | <p>Frank (2004)⁵⁰ Design: RCT N = 10,507 patients Implementation: 00/0000 Study Start: 03/1998 Study End: 03/1999</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>3</p> |
| <p>This cluster RCT is a complex set of multifaceted interventions to improve care and an economic evaluation (cost minimization and cost effectiveness analysis). The project sought to study passive vs. tailored interventions to improve management of cardiovascular risk factors according to guideline based care. The control group was usual care and the intervention group received an educational outreach visit by a pharmacist with audit and feedback, and computerized reminders linked to the EMR. The main outcomes were first time prescriptions for hypertension where thiazides were prescribed, patients assessed for cardiovascular risk before prescribing anti hypertensive or cholesterol-lowering agents, and patients treated for hypertension or high levels of cholesterol for 3 or more months who had achieved recommended treatment goals. Cost minimization framework was adopted, costs of intervention were set against reduced treatment costs. Net annual cost and cost per additional patient being started on thiazides.</p> | <p>Fretheim (2006)⁵⁶ Fretheim (2006)⁵⁷ Design: RCT N = 139 practices and 501 physicians Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>5</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|------------------|-------------------------------------|-----------------------------------|-----------------------|
| <p>A cluster RCT was conducted in a national network of primary care offices that use the Centricity EMR. The study examined the impact of an EMR-based intervention for lipid management incorporating nationally recognized guidelines (specifically the ATP-III guidelines) into the EMR. Prompts were generated at the point of care and included 3 pages: screening, assessment and management information. The 3 main outcome variables compared: proportion of patients tested adequately for hyperlipidemia, the proportion of patients whose most recent low-density lipoprotein cholesterol (LDL-C) was at goal (<100 for high-risk patients, <130 for moderate-risk patients, and <160 for low-risk patients), and the proportion of high-risk patients with an LDLC \geq130 who were prescribed lipid-lowering medications. Univariate (McNemar) and multivariate analysis (accounting for clustering) were performed. Results were presented with patients stratified by risk groups. A total of 105 physicians from 25 practices and 64,150 patients were included in the study.</p> | <p>Gill (2009)⁶¹ Design: RCT N = 64,150 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 10/2006</p> | <p>NA</p> | <p>ambulatory care</p> | <p>12 months</p> | <p>3</p> |
| <p>A cluster RCT was conducted in primary care clinics in Israel comparing the intervention with standard care. The pilot study was to evaluate the feasibility of a CDSS mainly on secondary prevention measure outcomes in patients with CAD and dyslipidemia followed by primary care physicians. In the intervention arm, a written reminder with patient tailored recommendations was mailed to the primary care physicians and nurses. The recommendations were based on the last 6 months data for new patients, and 4 months for patients in periodic follow-up. Rate of adequate monitoring, positive treatment trend, overall uptitration rate in patients with LDL = 110 mg/dl and LDL Levels were compared in between the control and intervention arms. 7,448 patients were included in the study. The intervention clinics included 204 general practitioners and 396 nurses.</p> | <p>Gilutz (2009)⁶² Design: RCT N = 7,448 patients from 56 control and 56 intervention clinics Implementation: 00/0000 Study Start: 01/2000 Study End: 12/2003</p> | <p>NA</p> | <p>ambulatory care Academic</p> | <p>21 months</p> | <p>4</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|---------------------------------|-----------------------|-----------------------------------|-----------------------|
| <p>A cluster randomized trial was conducted to assess the value of a discharge software application of CPOE in a 730-bed, tertiary care teaching hospital in central Illinois. The study compared the benefits of a CPOE with discharge software with usual handwritten discharge care for patients at high risk for repeat admission. Software features included required fields, pick lists, standard drug doses, alerts, reminders, and online reference information. The software prompted the discharging physician to enter pending tests and order tests after discharge. Hospital physicians used the software on the day of discharge and automatically generated 4 discharge documents. Proportion of patients readmitted at least once within 6 months of index hospitalization, emergency visits within 6 months and adverse events within 1 month were measured and compared. Perceptions about discharge from the perspective of patients, outpatient physicians and hospital physicians were examined by interview and survey. 631 patients, 70 hospital physicians and 496 outpatient physicians were involved in the study.</p> | <p>Graumlich (2009)²³⁷ Graumlich (2009)²³⁸ Design: RCT N = 631 patients Implementation: 00/0000 Study Start: 11/2004 Study End: 01/2007</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>6 months</p> | <p>2</p> |
| <p>A cluster-randomized controlled trial involving 1,118 long-term care residents at 29 resident care units in 2 facilities. The resident care units, each with computerized provider order entry, were randomized to having a clinical DS system (intervention units) or not (control units). Alerts in the form of warning messages appeared in the CPOE of intervention units. The number of adverse drug events, severity of events, and whether the events were preventable were measured in this study.</p> | <p>Gurwitz (2008)³⁰² Design: RCT N = 29 units randomized containing 1,118 patients Implementation: 00/0000 Study Start: 00/2000 Study End: 00/2000</p> | <p>NA</p> | <p>Long term care</p> | <p>6.3 Months</p> | <p>5</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|--|------------------|--|-----------------------------------|-----------------------|
| <p>RCT at general practice in Sør- and Nord-Trøndelag counties in Norway. The CDSS with clinical guidelines for treatment of hypertension was implemented as an external computer program, accessible from the main computerized record system. Health centers in intervention group had the CDSS. The CDSS guided the doctors in diagnostics, history and additional tests taking, physical examination, and treatments. Both doctors and assistants could use the CDSS, however, some parts of the CDSS were reserved for the doctors only. Doctors in control group followed their ordinary procedures for patients with hypertension. Group differences in level of systolic and diastolic BP, serum cholesterol, body mass index, and risk score for myocardial hypertension in general practice by use of a computer-based clinical infarction were calculated and compared. 53 doctors participated.</p> | <p>Hetlevik (1999)³⁰⁴ Design: RCT N = 1,998 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>18 months</p> | <p>4</p> |
| <p>A cluster RCT of 2,027 racially diverse adults receiving hypertension care in 14 primary care practices. To examine the effectiveness of computerized DS (CDS) designed to improve hypertension care and outcomes. Intervention arm was 18 mo of the physicians receiving CDS for each hypertensive patient compared to usual care without computerized support (control). Assessed prescribing of guideline recommended drug therapy and levels of BP control for patients and examined if the effects of the intervention differed by patients' race/ethnicity using interaction terms.</p> | <p>Hicks (2007)⁶⁷ Design: RCT N = 1,422 patients Implementation: 00/0000 Study Start: 07/2003 Study End: 02/2005</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>18 months</p> | <p>4</p> |
| <p>Randomized trial of 511 adult patients with type 2 diabetes receiving either usual care or intervention involving shared access by patient and primary care provider to a Web-based diabetes tracker. The tracker interfaced with the providers EMR and a phone reminder system which sent monthly reminders for medications, labs or doctor visits. The primary outcome measure was a process composite score.</p> | <p>Holbrook (2009)³⁶⁷ Design: RCT N = 511 patients Implementation: 00/0000 Study Start: 00/2002 Study End: 12/2003</p> | <p>NA</p> | <p>Ambulatory care Home</p> | <p>6 months</p> | <p>7</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|------------------|------------------------|-----------------------------------|-----------------------|
| <p>Cross-over randomized trial of 6 type 1 diabetic patients using a hand held insulin regimen optimizer. Patients had a 1 week run-in period using the tool as an electronic log book for their glucose measures. They then underwent 2 consecutive 3 week study periods, with and without the computerized insulin dose advice switched on. The clinical DS was suggesting optimum insulin dose based on patients entered data. Blood glucose was the primary outcome.</p> | <p>Holman (1996)³⁰⁶ Design: RCT N = 6 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Home</p> | <p>1.2 months</p> | <p>3</p> |
| <p>An RCT was conducted to demonstrate the potential effect of deploying a sentinel system that scans administrative claims information and clinical data to detect and mitigate errors in care and deviations from best medical practices. The study was performed among the commercially insured population of a university-affiliated managed care plan. The system relayed all triggered recommendations to intervention physicians (those for control group were deferred until the end of the study). Compliance with recommendations, hospital admissions and attendant cost were measured and compared between control and intervention groups. A total of 39,462 subjects were initially enrolled in the study. Charges were also compared.</p> | <p>Javitt (2005)²¹⁸ Design: RCT N = 39,462 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>6</p> |
| <p>A randomized trial of the effect of a DS tool designed to detect and help physicians from one HMO correct "missteps". Study group had "Care consideration" alerts given to the physicians (n =19,716) and the control group did not (n = 19,792). The DS tool used data from billing, lab and pharmacy to detect care considerations that fell into 3 recommendation categories: stop a drug, do a test, start a drug along a gradient of severity from severe, moderate, least severe. Severe alerts were phoned in to the medical director who called the physician, moderate and least severe alerts were processed through HMO nursing staff and passed on at their discretion. Alert resolution rates and costs were assessed.</p> | <p>Javitt (2008)⁷⁵ Design: RCT N = 39,508 patients Implementation: 01/2001 Study Start: 01/2001 Study End: 12/2001</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>6</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|--|--|---|-----------------------------------|-----------------------|
| <p>A randomized controlled double-blinded study was conducted at 3 pharmacies affiliated with an academic center to assess the impact of a system called Show Your Work (SYW) on pharmacy callbacks. The SYW system automatically annotates e-prescriptions by appending alerts and override comments to e-Prescriptions generated by an e-Prescribing system. This process adds notes below each medication order to describe any DS warnings that were displayed at the time of prescribing, any overrides to drug alerts provided by the prescriber during the session, and any dose calculations for pediatric prescriptions. Prescriptions were either printed or faxed to pharmacies throughout Tennessee. Pharmacy callback rates were measured and compared between those with "SYW off" and "SYW on" at 3 affiliated pharmacies; pharmacists' perceptions of the system were also studied using a qualitative and quantitative survey.</p> | <p>Johnson (2010)⁷⁶ Design: RCT N = 3,285 patients Implementation: 00/0000 Study Start: 04/2007 Study End: 08/2007</p> | <p>NA</p> | <p>Pharmacies Academic</p> | <p>4 months</p> | <p>7</p> |
| <p>A randomized trial assessing whether off-line data analysis, instead of event monitoring, was a viable method for initiating a clinical quality alert. A cohort of patients eligible for an alert was identified by off-line data analysis and a flag was set in their ambulatory Electronic Medical Records. One hundred clinicians were randomly assigned either to a control group or to a group that received the alert when viewing the electronic medical record of eligible patients. A low dose aspirin therapy alert was selected to test the feasibility. Comparisons were made on the proportion of patients no longer eligible for alert at end of month.</p> | <p>Krall (2004)⁸⁹ Design: RCT N = 1,076 patients Implementation: 00/1994 Study Start: 01/2000 Study End: 02/2000</p> | <p>Acute care/ tertiary</p> | | <p>1 month</p> | <p>3</p> |
| <p>An RCT at the Brigham and Women's Hospital in Boston, MA. A computer program linked to the patient database identified consecutive hospitalized patients at increased risk of VTE using 8 common risk factors identified from the EHR. Physicians in the intervention group were alerted of a patient's risk of VTE - they were required to acknowledge the alert and then withhold or order prophylaxis. Number of patients with DVT or PE and death were compared between at 90 days. Of the 2,506 patients studied, 2,361 were followed up beyond the index hospitalization.</p> | <p>Kucher (2005)⁹⁰ Design: RCT N = 2,506 patients Implementation: 00/0000 Study Start: 09/2000 Study End: 01/2004</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>3 months</p> | <p>6</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|--|------------------|--|-----------------------------------|-----------------------|
| <p>Cluster RCT of 32 Dutch General Practices. All had EMRs and half were given Asthma Critic which suggested actions for patients with asthma. Reminders were based on the Dutch College of General Practitioners asthma and COPD treatment guidelines. Outcomes were 3 patient (contact frequency, peak-flow, FEV1) and 5 prescribing (cromoglycate, deproprine, antihistamines, and oral bronchodilators).</p> | <p>Kuilboer (2006)²¹⁹ Design: RCT N = 32 primary care practices (78,926 patients of whom 9,798 had asthma or related symptoms) Implementation: 07/1998 Study Start: 07/1998 Study End: 05/1999</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>5 months</p> | <p>4</p> |
| <p>This RCT recruited physicians within an academic primary care practice who used EHR for majority of visits, then randomized adults (without their consent) identified by EHR review as having CAD or CAD risks and high levels of LDL cholesterol. Physicians received 1 email per intervention patient facilitating statin prescription and monitoring. Outcomes were changes in statin prescription, and cholesterol levels across times during the 1-year trial.</p> | <p>Lester (2005)⁹⁵ Design: RCT N = 235 patients and 14 clinicians Implementation: 07/2003 Study Start: 07/2003 Study End: 07/2004</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>12 months</p> | <p>6</p> |
| <p>A cluster RCT was conducted at 27 primary care clinics (matched pairs by size for all but 1 practice) associated with Partners HealthCare that uses internally developed full featured EHR called LMR. ARI Smart Form is an LMR module that is launched from the notes page of the EHR only when a physician triggers the module. Among many features, it provides DS in antibiotic prescribing, and antibiotic choices generation of diagnostic appropriate order set. Rate of antibiotic prescribing to patients with ARI (acute respiratory infection) were compared between the control and intervention groups. 111,820 patients and 443 clinicians were involved in the study. Analysis was adjusted for clustering.</p> | <p>Linder (2009)⁹⁷ Design: RCT N = 111,820 patients Implementation: 00/0000 Study Start: 11/2005 Study End: 05/2006</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>6 months</p> | <p>4</p> |
| <p>A randomized, controlled trial of 22 primary care clinics using either the existing system of no lab monitoring alerts or noninterruptive, on-screen recommendations for baseline laboratory tests when prescribing new medications. Prescribers did not need to respond to the alert. Differences in the proportion of visits resulting in lab testing within 14 days were analyzed. The clinics included 366 physicians, 2,765 patients and 3,673 events requiring lab monitoring test orders.</p> | <p>Lo (2009)²²⁰ Design: RCT N = 3,673 potential alert trigger events (prescriptions) Implementation: 00/2000 Study Start: 07/2003 Study End: 01/2004</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>0.5 months</p> | <p>3</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|--|--|--|-----------------------------------|-----------------------|
| <p>Cluster-RCT with an incomplete block design in the south of the Netherlands to assess the effect on drug-prescribing behavior of implementing prescribing guidelines by means of a reactive computer reminder system (CRS). 25 GPs (7 GP practices) received reminders about antibiotics and asthma/COPD prescriptions, 28 GPs (7 GP practices) received reminders about cholesterol prescriptions. Prescription guidelines were integrated into the computerized GP information system. Both performance indicators and prescription volumes were calculated as the main outcome measures.</p> | <p>Martens (2007)¹⁰¹ Design: RCT N = 77 physicians (GPs) Implementation: 04/2004 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>4</p> |
| <p>An RCT trial of electronic clinical reminders for primary care physicians to improve lab monitoring for maintenance therapy of potassium, creatinine, liver function, thyroid function and therapeutic drug levels. Reminders were generated if patients were on a target medication for at least 365 days with no record of a relevant lab test within the previous 365 days. Compliance rates were compared with usual care.</p> | <p>Matheny (2008)²²¹ Design: RCT N = 2,507 outpatient visits in 1,922 geriatric patients and 303 primary care physicians Implementation: 00/0000 Study Start: 01/2004 Study End: 06/2004</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>6 months</p> | <p>3</p> |
| <p>Prospective randomized study of diabetic patients at an adult diabetes clinic at Wishard Memorial Hospital to assess the impact of computer reminders to clinicians about a) out-of-date test results and b) specific changes in therapeutics. Each patient visit (n = 794 visits by 257 patients) was regarded as an independent event during the 8 month trial. Computer reminders consisted of paper reports printed for each patient encounter.</p> | <p>McDonald (1976)²²² Design: RCT N = 601 patient visits by 226 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>8 months</p> | <p>1</p> |
| <p>An RCT in Maryland design to test if AMTs improve hospitalized patient care with respect to costs, mortality, length of stay, or time spent managing antimicrobial utilization. Patients were randomized to DS (DS) for the AMT or usual care in all wards according to their chart number (odd/even). The reminder system was within the pharmacy information system. No EMR or CPOE were available.</p> | <p>McGregor (2006)¹⁰⁵ Design: RCT N = 4,507 patients Implementation: 00/000 Study Start: 05/2004 Study End: 08/2004</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>3 months</p> | <p>3</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|--|------------------|------------------------|-----------------------------------|-----------------------|
| <p>This study was intended to test effects of a web-based DS tool, the diabetes Disease Management Application (DMA), developed to improve evidence-based management of type 2 diabetes. A group RCT of 12 intervention and 14 control staff providers and 307 intervention and 291 control patients with type 2 diabetes was conducted in a hospital-based internal medicine clinic. Providers were randomly assigned from May 1998 through April 1999 to have access to the DMA (intervention) or not to have access (control). The web-based DMA is not an involuntary reminder system, but needs to be actively opened; it displays interactive patient-specific clinical data, treatment advice, and links to other web-based care resources. We compared patients in the intervention and control groups for changes in processes and outcomes of care from the year preceding the study through the year of the study by intention-to-treat analysis. Power analysis performed for change in HbA1c levels which is abstracted as the primary outcome.</p> | <p>Meigs (2003)³⁰⁹ Design: RCT N = 598 patients Implementation: 05/1998 Study Start: 05/1997 Study End: 04/1999</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>4</p> |
| <p>Cluster RCT of 27 general practices in Avon, U.K. which included 614 patients between 60 and 79 years with high BP. The trial was to investigate the effect of a computer based clinical DS system and a risk chart on absolute cardiovascular risk, BP, and prescribing of cardiovascular drugs in hypertensive patients. Interventions: Patients got CDSS system plus cardiovascular risk chart, cardiovascular risk chart alone, or usual care.</p> | <p>Montgomery (2000)¹⁰⁸ Design: RCT N = 552 patients Implementation: 00/0000 Study Start: 09/1996 Study End: 09/1998</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>5</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|-------------------------|--|-----------------------------------|-----------------------|
| <p>Randomized controlled trial with a 2 x 2 factorial design of physician and pharmacist CDSS alert interventions was conducted at a large, inner city, academic, internal medicine practice. The primary venues for this study were the general medicine practice and the Wishard Memorial Hospital outpatient pharmacy. The study assessed the effects of evidence-based treatment suggestions for hypertension made to physicians and pharmacists using a comprehensive electronic medical record system. Using data from patients' electronic medical records from Regenstrief medical records system, (RMRS) and data entered by physicians after patient visit. The computer-based ordering system generated care suggestions for both intervention and control groups; All hypertension care suggestions for intervention patients were displayed as "suggested orders" on physicians' workstations when they wrote orders after patient visits. For this study pharmacist intervention included the same care suggestions as generated for physicians and were displayed on the PIRS. There were 4 groups: control, physician intervention, pharmacy intervention and both interventions. QoL was the primary endpoint. 712 patients were included in the study.</p> | <p>Murray (2004)³¹⁰ Design: RCT N = 712 patients Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>12 months</p> | <p>5</p> |
| <p>To determine if computer reminders increase the provision of inpatient preventive care for 22 preventive care actions, 8 of which related to medication issues. Randomized, controlled trial on the general medicine inpatient service of an urban, university-affiliated public hospital. Study subjects were 78 house staff rotating on the 6 general medicine services. The intervention was reminders to physicians printed on daily rounds reports about preventive care for which their patients were eligible, and suggested orders for preventive care provided through the physicians' workstations. The preventive care guidelines were derived from the US Preventive Care Task Force recommendations. Compliance with preventive care guidelines and house staff attitudes toward providing preventive care to hospitalized patients were the main outcome measures. Reminders were generated for a total of 4,649 preventive care measures.</p> | <p>Overhage (1996)¹¹⁷ Design: RCT N = 24 practice teams Implementation: 10/1991 Study Start: 10/1992 Study End: 03/1993</p> | <p>General Hospital</p> | <p>Academic</p> | <p>6 months</p> | <p>6</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|-----------------------------|------------------------|-----------------------------------|-----------------------|
| <p>A RCT to assess if automated, guideline-based reminders to physicians, provided on-screen as they wrote orders, could impact compliance rates with 87 corollary (associated monitoring) orders. During the 6-month trial, reminders about corollary orders were presented to 48 intervention physicians and withheld from 41 control physicians in a general medicine public teaching hospital. All physicians had access to the guidelines, intervention physicians received the onscreen reminders that they could easily accept, reject or modify; for control physicians the computer tracked the number of time corollary orders would have been triggered. Compliance rates were compared immediately (at the time of the trigger order), at 24 hours post trigger order and within hospital stay compliance rates. In all there were 7,394 trigger orders and 11,404 suggestions for corollary orders.</p> | <p>Overhage (1997)¹¹⁸ Design: RCT N = 86 physicians on 6 services (services randomized) Implementation: 00/0000 Study Start: 10/1992 Study End: 04/1994</p> | <p>General Hospital</p> | <p>Academic</p> | <p>6 months</p> | <p>3</p> |
| <p>An RCT of 207 HMO primary care physicians who either received or did not receive drug laboratory monitoring alerts form within the CPOE system. The intervention group used the same CPOE and had the same electronic medication list as controls with additional information recommending specific lab monitoring for 25 select medications provided as nonintrusive reminders on the ordering screen. Compliance with guidelines for lab monitoring was compared between the groups, rates among the different drugs were also compared.</p> | <p>Palen (2006)¹²¹ Design: RCT N = 26,586 (index dispensing) patients Implementation: 00/0000 Study Start: 11/2002 Study End: 10/2003</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>3</p> |
| <p>Cohort study comparing TREAT's advice, CDSS to physician's treatment followed by a cluster RCT comparing wards using TREAT (intervention) versus antibiotic monitoring without TREAT (control). Patients had suspected harboring bacterial infections in 3 hospitals (Israel, Germany and Italy) 2,326 patients. The primary outcome, appropriate antibiotic treatment, was assessed among patients with MDI. Length of hospital stay, adverse events, mortality and antibiotic costs, including costs related to future antibiotic resistance, were compared for all patients.</p> | <p>Paul (2006)¹²⁴ Design: RCT N = 3,529 patients in the RCT and 1,203 in the cohort study Implementation: 00/0000 Study Start: 05/2004 Study End: 11/2004</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>6 months</p> | <p>7</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|---|--|-----------------------------------|-----------------------|
| <p>In a cluster-randomized design, 19 physicians caring for 334 eligible patients at least 40 years of age were randomized. All clinicians received computerized reminders through their EMR at office visits. Intervention physicians also received e-mails asking whether aspirin was indicated for each patient. If so, patients received a mailing and nurse telephone call addressing aspirin. The primary outcome was self reported regular aspirin use in 242 patients.</p> | <p>Persell (2008)²⁰⁵ Design: RCT N = 242 Patients Implementation: 00/0000 Study Start: 10/2004 Study End: 03/2005</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>6 months</p> | <p>4</p> |
| <p>A randomized trial to determine physicians' response rates of guided dosing for hospitalized patients 65 and older. Designed to assess if a guided dosing system delivering advice to physicians about appropriate initial dosing for a minority of drug orders as well as discouraged prescription of contraindicated drugs affects their compliance with guidelines. Dosing is to reduce falls in the elderly. 9,111 study-related orders by 778 providers were entered for 2,981 patients.</p> | <p>Peterson (2007)¹²⁶ Design: RCT N = 9,111 medication orders Implementation: 00/0000 Study Start: 12/2005 Study End: 08/2006</p> | <p>Acute care/ tertiary, Critical care units, Emergency department</p> | <p>Academic</p> | <p>9 months</p> | <p>3</p> |
| <p>Multicenter, prospective, pragmatic, with randomization of groups (clusters) designed to determine the cost-effectiveness of an intervention to promote the recommendations of the Global Initiative for Asthma (GINA). Group of 10 pulmonologists and 10 primary care physicians (who recruited 98 and 100 patients with persistent asthma respectively) were randomized to intervention and control. The intervention consisted of providing physicians with a hand-held clinical decision support system (SADC) that offered recommendations based on the GINA PLUS nurse trainers to assist patients. Doctors in the control group had handheld but did not have the SADC or nurses. Effectiveness was determined by measuring the quality of life through the St. Georges Respiratory Questionnaire (SGRQ). Costs were calculated from the consumption of resources registration for 12 months and determined the cost effectiveness of intervention by an incremental analysis.</p> | <p>Plaza (2005)²⁸⁰ Design: RCT N = 20 physicians+B44 Implementation: 03/2000 Study Start: 10/1999 Study End: 02/2001</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12.3 months</p> | <p>4</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|---|-----------------------------|------------------------|-----------------------------------|-----------------------|
| <p>A clustered RCT by Prescription in Ischaemic Stroke Management (PRISM) Study Group. Hospitals were randomized CDSS or control. Baseline clinical data were entered via an automated telephone data entry system as soon as possible after the hospital admission (inpatients) or clinic appointment (outpatients) of a study patient. The CDSS estimated individual acute stroke or TIA annual risks of recurrent ischaemic stroke, haemorrhagic stroke, myocardial infarction, other ischaemic vascular events and other haemorrhagic complications associated with each possible antiplatelet or anticoagulant therapy. Relative risk reduction (RRR) in ischaemic and haemorrhagic vascular events was compared. The information provided by the CDSS enabled informed prescribing decisions.</p> | <p>Prescription in Ischaemic Stroke Management (PRISM) Study Group (2003)¹²⁷ Design: RCT N = 1,640 Patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Unspecified Hospital</p> | | <p>6 months</p> | <p>4</p> |
| <p>A non-blinded pilot RCT was conducted in Maryland to assess the impact on A1c of a cell phone based diabetes management software system used with web data analytics and therapy optimization tools. Study patients received a Bluetooth enable blood glucose meter, a cell phone and WellDoc's proprietary diabetes management software, Diabetes Manager. Blood glucose reading are automatically sent to the cell phone and the phone-based software CDSS is initiated providing real time feedback. Patient is then prompted to enter insulin dosage and on hitting "OK" data is sent to the WellDoc server. Patient data were analysed by automated algorithms and by the research team. Average decrease of A1c and physicians change of medication were measured and compared between the groups. 30 patients were enrolled in the study.</p> | <p>Quinn (2008)¹²⁸ Design: RCT N = 30 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>3 months</p> | <p>4</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|------------------|------------------------|-----------------------------------|-----------------------|
| <p>An RCT was conducted on 400,000 HMO members 18 years and older who were assigned to the intervention group (alerts) or usual care group. The objective of the study was to determine whether computerized alerts were effective at increasing the percentage of ambulatory patients with laboratory monitoring at initiation of drug therapy. The primary outcome measure was the percentage of drug dispensing with baseline laboratory monitoring. Alerts were triggered by a dispensing of one of 15 target drug or drug classes. The alert was sent electronically to the Clinical Pharmacy Call Center daily if lab tests were not completed. This team of pharmacists contacted patients by phone to remind them their test was due or to order the tests if the physician did not do so. The intervention therefore had 2 stages; the alerting of the pharmacist by the computer and the phone follow-up by the pharmacist. 10,169 dispensing were included; the primary outcome was the percentage of drug dispensing with baseline lab monitoring.</p> | <p>Raebel (2005)¹³⁰ Design: RCT N = 9,565 patients, 10,169 dispensing Implementation: 00/0000 Study Start: 09/2002 Study End: 12/2003</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>15 months</p> | <p>6</p> |
| <p>Randomized trial of HMO patients 65 and over prescribed one of a newly targeted group of 11 potentially inappropriate medications. To determine if a computerized pharmacy alert system plus collaboration between health care professionals can affect the rate of potentially inappropriate medication dispensed in elderly patients. During this 1-year study, 1,187 patients (2.0% of 59,680 included) were newly dispensed one or more of the 11 medications. An alert generated in the pharmacy system prevented printing of the label until a pharmacist intervened by contacting prescribing clinicians by phone.</p> | <p>Raebel (2007)¹³¹ Design: RCT N = 59,680 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>5</p> |
| <p>Randomized trial of a computerized alert for pharmacists when pregnant patients were prescribed USFDA category D or X medications (evidence for fetal risk) compared to usual care. Measured by the proportion of pregnant women dispensed a category D or X medication and the total number of first dispensing of targeted medications. Alerts were sent to pharmacists who had to review prescription and contact prescriber before the prescription label would print. The alerts were generated from the integration of administrative and EHR data with the pharmacy system. Patients included 11,100 potentially pregnant women, HMO members, between 18 to 50 years randomized to intervention or usual care.</p> | <p>Raebel (2007)¹³² Design: RCT N = 11,100 women Implementation: 00/0000 Study Start: 01/2003 Study End: 04/2003</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>4 months</p> | <p>7</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|---|-----------------------------|------------------------|-----------------------------------|-----------------------|
| <p>This 3 arm cluster RCT randomized 200 US primary care physicians to 3 groups. Patients were in groups according to their physician: 78 patient's physicians received active reminders on treatment advice for patients with depression, 77 patient's physicians were told of the patient's depression, and 71 patient's physicians received usual notes. Notes were produced from an EMR (reminder system CDS). CDS based on AHCPR guidelines on depression.</p> | <p>Rollman (2002)¹³⁷ Design: RCT N = 200 Patients with documented major depression Implementation: 00/0000 Study Start: 04/1997 Study End: 12/1998</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>6 months</p> | <p>3</p> |
| <p>A randomized, controlled trial with an off-on-off design whereby a glucose regulation guideline was implemented in an intensive care unit in paper form during the first study period. During the second period, the guideline was randomly applied in either paper or computerized form. In the third period, the guideline was available only in paper form. Periods were 6, 10 and 4 weeks respectively. For the computer intervention each noncompliant event triggered a pop-up window to appear on top of the active CIS screen, alerting clinical staff members. This window appeared on bedside workstations and at any workstation where the patient's record was activated. 484 patients from an 18-bed ICU were included, 120 during the intervention period, attended by 11 intensivists and 93 nurses. The two guideline-related outcome measures consisted of compliance with: (a) glucose measurement timing recommendations and (b) insulin dose advice.</p> | <p>Rood (2005)¹³⁸ Design: RCT N = 484 patients Implementation: 04/2001 Study Start: 00/0000 Study End: 00/0000</p> | <p>Critical care units</p> | <p>Academic</p> | <p>5 months</p> | <p>6</p> |
| <p>To evaluate whether displaying context sensitive links to infrequently accessed educational materials and patient information via the user interface of an inpatient computerized care provider order entry (CPOE) system would affect access rates to the materials. The CPOE of Vanderbilt University Hospital included "baseline" clinical DS advice for safety and quality. Authors augmented this with 7 new, primarily educational DS features. A prospective, RCT compared clinicians' utilization rates for the new materials via two interfaces. Control subjects could access study-related DS from a menu in the standard CPOE interface. Intervention subjects received active notification when study-related DS was available through context sensitive, visibly highlighted, selectable hyperlinks. Rates of opportunities to access and utilization of study-related DS materials from April 1999 through March 2000.</p> | <p>Rosenbloom (2005)²⁶² Design: RCT N = 418,739 opportunities to access an information item Implementation: 00/1995 Study Start: 04/1999 Study End: 03/2000</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>12 months</p> | <p>4</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|------------------|--|-----------------------------------|-----------------------|
| <p>An RCT at 6 teaching practices of the Ottawa Civic Hospital Family Medicine. With the use of a standard randomization computer program these patients were assigned to the control group, the physician reminder group, the telephone reminder group or the letter reminder group. For patients in the physician reminder group a computer-generated reminder to ask the patient about tetanus vaccination was included on the routinely printed encounter form used for billing purposes. Proportion of patients who received tetanus toxoid during the study year or who had a claim of vaccination in the previous 10 years. 8,069 patients participated in the study. Costs were also assessed.</p> | <p>Rosser (1992)² Design: RCT N = 8,069 patients Implementation: 00/0000 Study Start: 04/1985 Study End: 03/1986</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>12 months</p> | <p>4</p> |
| <p>A randomized controlled trial was conducted at General Medical Clinic (GMC) at the Veterans Affairs (VA) Palo Alto Health Care System. The study chose two-period parallel design with the study subjects randomly divided into two groups the Physician Workstation (PWS) group and the Decentralized Hospital Computer Program (DHCP), group. The PWS system contained features designed to reduce prescription-drug costs and to reduce the number of adverse drug interactions. The PWS system provided alerts about potential adverse drug interactions. User Satisfaction rating was measured and compared. 34 physicians were involved in the study.</p> | <p>Rotman (1996)²⁶⁴ Design: RCT N = 34 Physicians Implementation: 00/0000 Study Start: 07/1994 Study End: 06/1995</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>2</p> |
| <p>Cluster randomized, controlled trial to improve Blood pressure within 2 hospital-based and 8 community-based clinics in the Veterans Affairs Tennessee Valley Healthcare System. There were 1,341 veterans with essential hypertension cared for by 182 providers compared across 3 study arms for 6 months. Providers were randomly assigned to receive an e-mail with a Web-based link to the 7th Report of the Joint National Committee on the Prevention, Detection, Evaluation and Treatment of High Blood Pressure guidelines (provider education); provider education and a patient-specific hypertension computerized alert (provider education and alert); or provider education, hypertension alert, and patient education, in which patients were sent a letter advocating drug adherence, lifestyle modification, and conversations with providers (patient education).</p> | <p>Roumie (2006)²²⁶ Roumie (2007)²²⁷ Design: RCT N = 871 patients Implementation: 00/0000 Study Start: 06/2004 Study End: 12/2004</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>6 months</p> | <p>6</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|---|--|--|-----------------------------------|-----------------------|
| <p>A controlled clinical trial (cluster randomization) among physicians and nurse practitioners caring for 349 patients (191 I and 158 C) infected with the human immuno-deficiency virus (HIV). Intervention was a very poorly described 'knowledge-based medical record ' which was an integration of the on-line patient record, rule-based DS, and full-text information retrieval into a clinical workstation. Main outcome was time to implementation of clinical alerts with secondary review of and improved quality of care. In the 18 month trial, 191 patients were treated by 70 physicians and nurse practitioners assigned to the intervention group, and 158 patients were treated by 66 physicians and nurse practitioners assigned to the control group.</p> | <p>Safran (1995)¹⁴¹ Safran (1993)¹⁴² Design: RCT - cluster N = 126 physicians, 10 nurse practitioners Implementation: 00/0000 Study Start: 05/1992 Study End: 09/1993</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>18 months</p> | <p>4</p> |
| <p>A RCT at Partners HealthCare System which is an integrated health care network consisting of outpatient clinics, community hospitals, and 2 academic teaching hospitals (Brigham and Women's Hospital and Massachusetts General Hospital) in Boston. Internally developed ambulatory EMR system integrated with patient specific electronic clinical reminder (recommendation for diabetes care and coronary artery disease[CAD]) allowed physicians to maintain patient problem, medication, and allergy lists and view laboratory results. Physicians also used the system to enter patient notes and medication prescriptions. Each time a clinician opened a patient chart within the system, the algorithm for all reminders determined whether the patient had received care in accordance with the recommended practice guidelines. Diabetes and CAD reminders resulting in recommended action were compared. 194 primary care physicians and 6,243 patients.</p> | <p>Sequist (2005)²²⁸ Design: RCT N = 6,243 Patients Implementation: 07/2000 Study Start: 10/2002 Study End: 04/2003</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>6 months</p> | <p>3</p> |
| <p>RCT comparing the prescribing of vancomycin by hospital physicians receiving real time computerized guidelines during the physician order entry process vs. no computerized guidelines during physician order entry. Measures of vancomycin prescribing were the number of orders, duration of the therapy and number of days per course of treatment. 396 physicians and 1,798 patients in a tertiary-care teaching hospital were studied.</p> | <p>Shojania (1998)¹⁴⁸ Design: RCT N = 396 physicians Implementation: 00/0000 Study Start: 06/1996 Study End: 03/1997</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>9 months</p> | <p>5</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|--|------------------|------------------------|-----------------------------------|-----------------------|
| <p>A cluster RCT in Quebec to test whether CDS in primary care EMR would reduce inappropriate prescribing. Physicians in the CDS group obtained information on each patient by downloading updates of dispensed prescriptions from the Régie de l'assurance maladie du Québec (RAMQ) drug-insurance program. All retail pharmacies had a data link to the RAMQ for online prescription adjudication, which provided a daily update of all prescriptions dispensed for each patient. These data were integrated into the patient's HR and categorized as having been prescribed by the study physician or by another physician. Alerts identified 159 clinically relevant prescribing problems in the elderly, a list established previously by expert consensus. The alerts appeared when the electronic chart was opened, when prescription-record updates were downloaded from the RAMQ, and when current health problems and prescriptions were recorded by the physician in the chart. Each alert identified the nature of the problem and possible consequences and suggested alternative therapy in accordance with the expert consensus. The primary outcomes were initiation and discontinuation rates of the 159 prescription-related problems. 107 physicians participated.</p> | <p>Tamblyn (2003)¹⁵⁹ Design: RCT N = 12,560 Patients Implementation: 00/0000 Study Start: 01/1997 Study End: 02/1998</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>13 months</p> | <p>2</p> |
| <p>An RCT to determine if a cardiovascular medication tracking and nonadherence alert system, incorporated into a computerized health record system, would increase drug profile review by primary care physicians, increase the likelihood of therapy change, and improve adherence with antihypertensive and lipid-lowering drugs. There were 2,293 primary care patients prescribed lipid-lowering or antihypertensive drugs by 59 physicians who were randomized to the adherence tracking and alert system or active medication list alone to determine if the intervention increased drug profile review, changes in cardiovascular drug treatment, and refill adherence in the first 6 months.</p> | <p>Tamblyn (2010)¹⁶⁰ Design: RCT N = 2,293 patients Implementation: 00/0000 Study Start: 04/2006 Study End: 00/0000</p> | <p>NA</p> | <p>ambulatory care</p> | <p>6 months</p> | <p>7</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|---|---|--|--|-----------------------------------|-----------------------|
| <p>An RCT at the Wishard Memorial Hospital is a tax-supported, 450-bed, university-affiliated, urban, public hospital located on the Indiana University Medical Center campus. Physicians were randomized. The intervention was CDSS designed to reduce prescribing of potentially inappropriate medications for older adults. CDSS was provided only when a physician in the intervention group prescribed a targeted inappropriate medication for a patient aged 65 and older who was being discharged from the Emergency department (ED). Proportion of ED visits by seniors with an inappropriate medication was measured. 5,162 patients and 63 physicians were involved in the study.</p> | <p>Terrell (2009)¹⁶⁴ Design: RCT N = 5,162 Patients Implementation: 00/0000 Study Start: 01/2005 Study End: 07/2007</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>30 months</p> | <p>7</p> |
| <p>An RCT was conducted at the Wishard Memorial Hospital, a tax-supported, 450-bed, university-affiliated, urban, public hospital. DS advised against use of nine potentially inappropriate medications and recommended safer substitute therapies. DS was provided only when a physician in the intervention group attempted to prescribe a targeted inappropriate medication for a patient aged 65 and older who was being discharged from the ED. The primary outcome measured was the proportion of ED visits by older adults that resulted in one or more prescriptions for a targeted inappropriate medication. The secondary outcome of interest examined was the proportion of all prescribed medications that were potentially inappropriate. 5,162 patients and 63 physicians were involved in the study.</p> | <p>Terrell (2009)¹⁶⁵ Design: RCT N = 63 physicians had 5,162 patient visits Implementation: 00/0000 Study Start: 01/2005 Study End: 07/2007</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>30 months</p> | <p>8</p> |
| <p>An RCT at an Inner-city academic general internal medicine practice to assess the effects of guideline-based care suggestions for asthma and CPOD delivered to physicians when writing orders on computer workstations. 246 physicians general internists, and 20 outpatient pharmacists were randomized to Care suggestions concerning drugs and monitoring. This 2 X 2 factorial randomization of practice sessions and pharmacists resulted in four groups of patients: physician intervention, pharmacist intervention, both interventions, and controls. Enrolled 706 of their primary care patients with HF or IHD. Outcomes were adherence to preventive care guidelines. 4 groups studied: control, pharmacist, physician or both health professionals. CPOE and CDSS intervention for physicians, pharmacist prompts were to provide education on several computer identified issues.</p> | <p>Tierney (2003)¹⁶⁶ Design: RCT N = 706 patients, 20 pharmacists, 94 physicians and 1 nurse practitioner Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>12 months</p> | <p>4</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|--|--|-----------------------------------|-----------------------|
| <p>An RCT at an Inner-city, academic, general internal medicine practice to assess the effects of guideline-based care (NAEP Expert Panel Report and Canadian Thoracic Society) suggestions for asthma and CPOE delivered to physicians when writing orders on computer workstations. 246 physicians general internists, and 20 outpatient pharmacists were randomized to Care suggestions concerning drugs and monitoring. This 2 X 2 factorial randomization of practice sessions and pharmacists resulted in four groups of patients: physician intervention, pharmacist intervention, both interventions, and controls. Enrolled 706 of their primary care patients with asthma or COPD. Outcomes were adherence to preventive care guidelines. 4 groups studied: control, pharmacist, physician or both health professionals. CPOE and CDSS intervention for physicians, pharmacist prompts were to provide education on several computer identified issues.</p> | <p>Tierney (2005)¹⁶⁷ Design: RCT N = 706 patients Implementation: 00/0000 Study Start: 01/1994 Study End: 05/1996</p> | <p>NA</p> | <p>Ambulatory care Academic</p> | <p>12 months</p> | <p>4</p> |
| <p>A cluster RCT in practices using ELIAS EHR system in the Netherlands were invited to participate in the study. Non-commercial home grown CDSS for lipid management based on recommendations from the guideline of Dutch College of General Practitioner (DCGP) was developed and integrated with the EHR system. Practices were randomly assigned to 3 arms of the study: control arm, and 2 intervention arms (an on-demand arm and an alerting arm). Each practice was subsequently assigned by simple random allocation to CDSS alerting, CDSS on demand, or control groups for the complete study period. The CDSS analyzed and interpret the patient data in the EHR, generating patient-specific guideline recommendations for preventative activities. Data on patients requiring treatment and patient treated based on the two intervention arms were measured and compared. 87,866 patients participated in the study. 77 physicians completed the study.</p> | <p>Van Wyk (2007)¹⁷¹ Design: RCT N = 87,860 Patients Implementation: 00/0000 Study Start: 05/2004 Study End: 00/0000</p> | <p>NA</p> | <p>Ambulatory care</p> | <p>12 months</p> | <p>3</p> |
| <p>A randomized trial was undertaken at LDS Hospital to assess the effects of digoxin alert reports generated by nightly review of the patient database, lab data and electrocardiographic findings. Reports were printed in the nursing division and placed in patient charts. 396 patients were randomly assigned to alert and non-alert groups over 3 months. Rate of physician actions were compared.</p> | <p>White (1984)²²⁹ Design: RCT N = 396 patients Implementation: 00/0000 Study Start: 00/0000 Study End: 00/0000</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>3 months</p> | <p>7</p> |

Evidence Table 13. KQ7: integrated CDSS study characteristics: setting and quality (continued)

| PICOM | Article Information | Type of Hospital | Other Settings | Length of Follow-up (mean months) | Summary Methods Score |
|--|---|-----------------------------|-----------------|-----------------------------------|-----------------------|
| <p>RCT to assess the impact of an automated intraoperative alert to redose prophylactic antibiotics in prolonged cardiac operations. All patients were randomized to an audible and visual reminder on the operating room computer console at 225 minutes after the administration of preoperative antibiotics (reminder group, n =137) or control (n =136).</p> | <p>Zanetti (2003)¹⁸⁰ Design: RCT N = 273 patients having cardiac surgery Implementation: 00/0000 Study Start: 03/2000 Study End: 06/ 2000</p> | <p>Acute care/ tertiary</p> | <p>Academic</p> | <p>1 week</p> | <p>6</p> |

Evidence Table 14. KQ7: integrated CDSS study characteristics: participants and interventions

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--------------------------------|--------------------------------|---------------------------|--------------|---------------------|------------------|--|-------------------------|---------------|
| Field (2009) ⁴² | Meds, Prescrip, Orders | NA | NA | renal Insufficiency | anti-depressants | alerts related to medication prescribing for residents with renal insufficiency were displayed to prescribers in the intervention units and hidden but tracked in control units. | monitoring, prescribing | Comm |
| Holbrook (2009) ³⁶⁷ | Patients | NA | UnDiff | diabetes | No | usual care of vascular risk patients | monitoring | NR |
| Lo (2009) ²²⁰ | Patients | NA | NR | No | No | usual care | monitoring | NR |
| Gurwitz (2008) ³⁰² | Patients | NA | 65+ | No | No | usual care - units already had CPOE | prescribing | HG |

*indicates outcomes noted as being the primary outcome by the paper's authors

Abbreviations: ACE = Angiotensin Converting Enzyme; AD = Academic Detailing; Adol = Adolescents; AMTs = Antimicrobial Management Teams; APAP = acetaminophen; ARB = Angiotensin Receptor Blocker; ARI = acute respiratory infection; CAD = Coronary Artery Disease; CDS = Clinical / Computerized Decision Support; Comm = Commercial; COPD = Chronic Obstructive Pulmonary Disease; CPOE = Computerized Provider Order Entry; CVD = Cardiovascular Disease; DMA = Disease Management Application; ED = Emergency Department; ESCHM = European Society of Cardiology and other societies for Hypercholesterolemia Management; GI = Gastrointestinal; GINA = Global Initiative for Asthma; GP = General Practitioner; HG = Homegrown; HIT = Health Information Technology; HMG Co-A = 3-hydroxy-3-methylglutaryl-coenzyme A; Meds = medications; MI = Myocardial Infarction; MM = Medication Management; NA = Not Applicable; NR = not reported; NSAIDs = Nonsteroidal anti-inflammatory drugs; Prescrip = prescriptions; PRISM = Prescription in Ischemic Stroke Management; UnDiff = undifferentiated, SYW = show your work; UTI = Urinary Tract Infection

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--------------------------------|--------------------------------|---------------------------|-------------------|-----------------|---|--|--------------------|---------------|
| Matheny (2008) ²²¹ | Outpatient visits | Primary Care | 65+ | multiple | Metformin; Potassium Supplement; Potassium Sparing Diuretic, Thiazide Diuretic; Angiotensin Converting Enzyme Inhibitor; HMG Co-A Reductase Inhibitor; Thyroxine; Carbamazapine; Cyclosporine, Phenobarbital, Phenytoin, Proc-NAPA, Valproate | usual care - clinics were randomized so that physicians received either usual care or electronic reminders at the time of office visits focused on potassium, creatinine, liver function, thyroid function, and therapeutic drug levels. | monitoring | NR |
| Javitt (2008) ⁷⁵ | Patients | Physicians UnDiff | ages 13 to 44 | no | no | usual care- no care consideration with decision support tool | prescribing | Comm |
| Hicks (2007) ⁶⁷ | Patients | NA | 19 to 64, and 65+ | hypertension | antihypertensive drugs | usual care without decision support | monitoring | NR |
| Martens (2007) ¹⁰¹ | Health Care Providers | Primary Care GP | NA | asthma | antibiotics and cholesterol-lowering drugs | reminder about 2 different types of prescriptions. All GPs were blind to the fact that they only received a specific subset of all available prescribing reminders and that they were analysed on certain prescribing behavior as controls | prescribing | NR |
| Peterson (2007) ¹²⁶ | Meds, Prescrip, Orders | Hospitalists | NA | No | No | recommended drugs | prescribing | HG |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--|--------------------------------|---------------------------|----------------|--------------------------------------|--|---|-------------------------|---------------|
| Raebel (2007) ¹³² | Patients | NA | 19 to 64 years | pregnancy | 21 drugs that were pregnancy risk category D (contraindicated with fetal risk although some therapeutic benefit) and X (evidence of fetal risk and no therapeutic benefit) | usual care | prescribing | Comm HG |
| Bailey (2007) ⁷ | Patients | NA | UnDiff | heart disease | ACE inhibitors, statins, aspirin and B-blockers | Usual care- Acute MI Patients in the control group received standard care | prescribing | NR |
| Paul (2006) ¹²⁴ | Patients | NA | 19- 64 years | No | antibiotics | Control group antibiotic monitoring without CDS | prescribing | HG |
| Fretheim (2006) ⁵⁶ Fretheim (2006) ⁵⁷ | Patients | Primary Care GP | 45-64 and 65+ | hypertension or hypercholesterolemia | thiazides | Usual care | Prescribing | NR |
| Feldstein (2006) ²¹⁵ Smith (2009) ²¹⁶ | Patients | NA | 19-64 and 65+ | No | 10 study medications or medication classes; ACE/ARB, Allopurinol, Carbamazepine, Diuretic, Metformin, Phenytoin, Pioglitazone, Potassium | Usual care | monitoring | NR |
| Kuilboer (2006) ²¹⁹ | Patients | NA | UnDiff | asthma | cromoglycate, deproprine, antihistamines, and oral bronchodilators | Usual care | Monitoring, Prescribing | NR |
| Roumie (2006) ²²⁶ Roumie (2007) ²²⁷ | Patients | NA | 19-64 and 65+ | hypertension | antihypertensive medications | Not clear - it was a Multiple Intervention comparison | prescribing | Comm |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--------------------------------|--------------------------------|--------------------------------------|----------------|-----------------------|--|--|-------------------------|---------------|
| McGregor (2006) ¹⁰⁵ | Patients | NA | 19- 64 and 65+ | No | 23 restricted antimicrobials | Control was without the system in the control arm - antimicrobial management teams | prescribing | Comm |
| Palen (2006) ¹²¹ | Meds, Prescrip, Orders | NA | NA | No | 25 specific drugs requiring lab monitoring according to guidelines, within the following classes: ACE inhibitors (2), Angiotension II receptor blocker (1), antiarrhythmic (1), antiinfective agents (2), antigout (2), cholesterol-lowering (5), diuretics (5), hyperglycemics (2), metabolic (2), neurological (3) | Control group - Did NOT receive drug laboratory monitoring alerts within the CPOE system. | Monitoring, Prescribing | Comm |
| Feldstein (2006) ⁴⁰ | Meds, Prescrip, Orders | NA | NA | No | Warfarin | Control - All 15 clinics received electronic medical record alerts for the coprescription of warfarin and 5 interacting medications. | prescribing | NR |
| Feldstein (2006) ⁴¹ | Patients | NA | 45-64 and 65+ | probably osteoporosis | No | Usual care- Control was no provider reminder or patient education, just usual care. | prescribing | NR |
| Berner (2006) ¹² | Health Care Providers | Primary Care GP and Other Physicians | NA | No | NSAIDS | Control group did not receive the rule for GI risk assessment. | prescribing | Comm HG |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|----------------------------------|--------------------------------|---------------------------|---------------|-------------------------|---|--|-------------------------|---------------|
| Lester (2005) ⁹⁵ | Patients | NA | 19-64 and 65+ | high cholesterol levels | statins | Control - Physicians did NOT receive a visit-independent disease management tool which was initiated by an email with CDS and facilitated "one-click" order writing. | Monitoring, Prescribing | HG |
| Raebel (2005) ¹³⁰ | Meds, Prescrip, Orders | NA | NA | No | 15 drugs/drug classes requiring lab monitoring | Pharmacists were alerted to missing laboratory test information only for intervention patients. Pharmacists were not provided information about laboratory monitoring for patients in the usual-care group. | Monitoring, Prescribing | HG |
| Bloomfield (2005) ¹⁵ | Patients | NA | NR | heart disease | lipid modifying therapy: fibrates (such as gemfibrozil), statins, bile acid binding resins, or niacin | 6 control clinic did NOT receive prompts. | prescribing | NR |
| Rosenbloom (2005) ²⁶² | Health Care Providers | Hospitalists | NA | No | No | Control subjects could access study-related decision support from a menu in the standard CPOE interface but they DID NOT receive active notification when study-related decision support was available through context sensitive, visibly highlighted, selectable hyperlinks | Prescribing | Comm |
| Tierney (2005) ¹⁶⁷ | Patients | Hospitalists | 19-64 and 65+ | asthma and COPD | No | 4 comparison groups 1 was control - usual care | prescribing | HG |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--------------------------------|--------------------------------|---------------------------|----------------|------------------------|--|--|-------------------------|---------------|
| Kucher (2005) ⁹⁰ | Patients | NA | 19-64 and 65+ | venous thromboembolism | No | Control group - no alert given | Prescribing | HG |
| Tierney (2003) ¹⁶⁶ | Patients | NA | 19-64 and 65+ | heart disease | No | control - No Evidence-based cardiac care suggestions | Prescribing | HG |
| Rollman (2002) ¹³⁷ | Health Care Providers | Primary Care GP | 18- 64 years | depression | antidepressants | Usual pt care for depression | Monitoring, Prescribing | Comm |
| Krall (2004) ⁸⁹ | Patients | NA | UnDiff | No | aspirin | Control - no alert for aspirin | prescribing | Comm |
| Filippi (2003) ⁴⁴ | Patients | NA | 19-64 and 65+ | diabetes | antiplatelet drugs | Control - No electronic reminder only a letter summarizing the beneficial effects of antiplatelet drugs | prescribing | NR |
| Zanetti (2003) ¹⁸⁰ | Patients | NA | 19- 64 and 65+ | heart disease | antibiotics - Cefazolin | Control - No audible and visual reminder on the operating room computer console at 225 minutes in surgery | prescribing | HG |
| Flottorp, (2002) ⁴⁷ | Consults | NA | NA | sore throat UTI | antibiotics | 72 practices received interventions to implement guidelines for urinary tract infection and 70 practices received interventions to implement guidelines for sore throat, serving as controls for each other. | prescribing | NR |
| Dexter (2001) ³³ | Patients | NA | NR | No | pneumococcal vaccination, influenza vaccination, prophylactic enteric coated aspirin and prophylactic subcutaneous heparin | Control - no preventive care reminders | prescribing | NR |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--|--------------------------------|---|---------------------|----------------------|------------------------------|---|---------------------------|---------------|
| Christakis (2001) ²³ | Health Care Providers | Primary Care GP, Hospitalists, Other Physicians | NA | acute otitis media | antibiotics for otitis media | Control - Providers did not receive real time evidence-based prompts on their prescribing practice for otitis media. | prescribing | HG |
| Montgomery (2000) ¹⁰⁸ | Patients | NA | 60-79 years | hypertension | No | Usual care for patients with Hypertension | monitoring | Comm HG |
| Shojania (1998) ¹⁴⁸ | Health Care Providers | Hospitalists | NA | No | vancomycin | Control physicians encountered no guidelines screens only the usual computer prompt to renew or discontinue the order after 72 hours of therapy | prescribing | HG |
| Overhage (1997) ¹¹⁸ | Health Care Providers | Hospitalists, Other Physicians | 19-64 and 65+ years | No | No | Control - reminders about corollary orders were withheld | monitoring | HG |
| Holman (1996) ³⁰⁶ | Patients | NA | 19- 64 years | diabetes | insulin | Control - patients were their own control - had device but dose support turned off (could still enter glucoses) | Administering, Monitoring | HG |
| Safran (1995) ¹⁴¹ Safran (1993) ¹⁴² | Patients | Physicians, Nurse Practitioners | NR | HIV | No | Control - alerts were not visible to control group | Monitoring, Prescribing | HG |
| Evans (1994) ³⁹ | antibiotic cultures | NA | NA | No | antibiotics | Control -No computerized antibiotic consultant. Two-stage random-selection study. antibiotics ordered compared between crossover periods. | Prescribing | HG |
| Cobos (2005) ²⁵ | Patients | NA | 19-64 and 65+ | Hypercholesterolemia | No | Usual Care | Prescribing | NR |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--------------------------------|--------------------------------|---------------------------|---------------|--|-------------------------------------|--|-------------------------|---------------|
| Davis (2007) ³⁰ | Health Care Providers | Primary Care GPs | NA | acute otitis media, allergic rhinitis, sinusitis, constipation, pharyngitis, croup, urticaria, and bronchiolitis | No | Control Group Provider - Did NOT have point-of-care evidence-based prescription writer and decision support system. | Prescribing | HG |
| Dexter (2004) ³⁴ | Patients | NA | NR | No | influenza and pneumococcal vaccines | Comparison of computerized physician standing orders compared with physician reminders for inpatient vaccinations. | prescribing | NR |
| Fiks (2009) ⁴³ | Patients | NA | 2 to 18 years | asthma | Influenza vaccine | Control sites had no electronic health record-based clinical alerts for influenza vaccine | prescribing | Comm |
| Hetlevik (1999) ³⁰⁴ | Patients | NA | 45-64 and 65+ | hypertension | No | Control - No CDS, doctors in control group were supposed to follow their ordinary procedures in the treatment of patients with hypertension. | Prescribing | NR |
| Rood (2005) ¹³⁸ | Patients | NA | UnDiff | No | No | Control was glucose regulation guideline in an intensive care unit in paper form. Paper form was implemented in the first and third period of the study. | Monitoring, Prescribing | Comm |
| Rosser (1992) ² | Patients | NA | 19-64 and 65+ | No | Tetanus Toxoid | Control – no reminder | Prescribing | NR |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--|--------------------------------|---------------------------|---------------|------------------|------------------------------|---|-------------------------|---------------|
| Sequist (2005) ²²⁸ | Patients | NA | 45-64 and 65+ | diabetes CAD | No | Control was usual care for diabetics and CAD pts, no guideline recommendations. | monitoring | HG |
| Van Wyk (2007) ¹⁷¹ | Patients | NA | 19-64 and 65+ | high cholesterol | No | Usual care | prescribing | HG |
| Persell (2008) ²⁰⁵ | Patients | NA | 19-64 and 65+ | diabetes | aspirin | Control group received electronic Reminder only for aspirin | Administering | NR |
| Terrell (2009) ¹⁶⁴ | Patients | NA | 65+ | No | No | Control - No Decision support that advised against use of potentially inappropriate medications in ED visit | prescribing | NR |
| Prescription in Ischemic Stroke Management (PRISM) Study Group (2003) ¹²⁷ | Patients | NA | 45-64 and 65+ | ischemic stroke | antiplatelets anticoagulants | Usual care | prescribing | NR |
| Tamblyn (2003) ¹⁵⁹ | Patients | NA | 65+ | No | No | Control no CDS | prescribing | NR |
| Rotman (1996) ²⁶⁴ | Health Care Providers | Physicians UnDiff | NA | No | No | Control - not using Physician Workstation | Prescribing | HG |
| Bertoni (2009) ¹⁴ | Health Care Providers | Primary Care GPs | NA | No | lipid lowering therapy | Control - Alternative Intervention for High BP treatment | Monitoring, Prescribing | NR |
| Demakis (2000) ²¹³ | Health Care Providers | Other Physicians | NA | No | No | Control - No reminder on standards of ambulatory care. | monitoring | NR |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--------------------------------|--------------------------------|---|--------------|-----------------|--|--|-------------------------|---------------|
| Fortuna (2009) ⁴⁸ | Health Care Providers | Primary care GPs, Specialist, Nurses, Mid level practitioners | NA | No | hypnotics: Ambien CR [®] (zolpidem tartrate extended release), Lunesta [®] (eszopiclone), Sonata [®] (zaleplon), and Rozerem [®] (ramelteon) | Control usual care - received an alert stating only the copayment tier of the medication. NO computerized alerts or educational sessions for hypnotic prescribing. | prescribing | Comm HG |
| Frank (2004) ⁵⁰ | Patients | NA | UnDiff | No | No | Control - usual care, no reminders of the 12 preselected preventive care activities. | prescribing | NR |
| Meigs (2003) ³⁰⁹ | Patients | NA | NR | diabetes | No | Control providers continued their usual care practices during the intervention and did not have access to the CDS DMA. | Monitoring, | NR |
| Overhage (1996) ¹¹⁷ | Health Care Providers | Physicians UnDiff | NA | No | No | Control - no electronic preventative care guideline reminders | prescribing | HG |
| White (1984) ²²⁹ | Patients | NA | UnDiff | No | digoxin | Control - no alert for digoxin intoxication | monitoring | HG |
| McDonald (1976) ²²² | Patients | NA | UnDiff | diabetes | No | Control - without computer reminders/ suggestions | monitoring | HG |
| Plaza (2005) ²⁸⁰ | Patients | Primary Care GPs, Respirologist | 14 years + | asthma | No | Access to handheld but no nurse trainers of Global Initiative for Asthma guideline advice. | Monitoring, Prescribing | Comm (likely) |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|------------------------------|--------------------------------|---------------------------|--|-----------------|---|--|----------------------------|---------------|
| Raebel (2007) ¹³¹ | Patients | NA | Geriatric (65 plus) | No | Drugs inappropriate for the elderly - amitriptyline, chlorthalidone, chlorpropamide, diazepam, doxepin, flurazepam, aspirin in combination with hydrocodone or oxycodone, ketorolac, oral meperidine, and piroxicam | Usual Care – Pharmacists did not receive the medication alerts generated by the pharmacy information management system for elderly patients newly prescribed a potentially inappropriate medication. | Dispensing, Prescribing | HG |
| Murray (2004) ³¹⁰ | Patients | NA | Geriatric (65 plus) Middle age (45 to 64) | hypertension | antihypertensive agents were angiotensin-converting enzyme (ACE) inhibitors, b-blockers, calcium channel blockers, oral clonidine and topical patch, diuretics, and other less commonly prescribed drugs such as methyldopa and reserpine | Control - Neither physician nor pharmacist received hypertension care suggestions for patients | Prescribing | HG |
| Frances (2001) ⁴⁹ | Patients | NA | Geriatric (65 plus) | heart diseases | aspirin, Beta-blockers, and cholesterol-lowering agents | Usual care | Prescribing | NR |
| Javitt (2005) ²¹⁸ | Patients | NA | Adol (13 to 18) Adults (19 to 44), Middle age (45 to 64) | No | No | Control Group - The system relayed all triggered recommendations to intervention physicians - those for control group were deferred until the end of the study | Monitoring and Prescribing | HG |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|--|--------------------------------|---------------------------|---|-----------------|--------------|--|------------------------------------|---------------|
| Quinn (2008) ¹²⁸ | Patients | NA | Adol (13 to 18) Adults (19 to 44) Middle age (45 to 64) | diabetes | No | Usual out patient care - Controls received one touch ultra BG meters, Testing strips and lancets. Faxed or called in results. Intervention pts -bluetooth enable blood glucose meter, a cell phone and WellDoc's proprietary diabetes management software, Diabetes Manager, automatically sent to the cell phone and the phone-based software CDSS is initiated providing real time feedback. | Monitoring, Prescribing, Education | Comm |
| Linder (2009) ⁹⁷ | Patients | NA | Middle age (45 to 64) | ARI | antibiotics | Usual care | Prescribing | HG |
| Bell (2010) ¹¹ | Patients | NA | Adol (13 to 18) Children (2 to 12) | asthma | No | Control group had passive access to the same asthma management tools. | Prescribing | Comm |
| Graumlich (2009) ²³⁷ Graumlich (2009) ²³⁸ | Patients | NA | Adol (13 to 18) Adults (19 to 44) Geriatric (65 plus) Middle age (45 to 64) | No | No | Usual Care - usual handwritten discharge care for patients at high risk for repeat admission. | Prescribing | NR |

Evidence Table 14. KQ7: integrated CDSS Study Characteristics: participants and interventions (continued)

| Author (year) | Primary unit of study analysis | Health Care Provider type | Patient type | Disease Specify | Drug specify | Control Group Type | MM PHASE(S) Target | Nature of HIT |
|-------------------------------|--------------------------------|---------------------------|---|---------------------|---|--|--|---------------|
| Terrell (2009) ¹⁶⁵ | Patients | NA | Geriatric (65 plus) | No | Promethazine Diphenhydramine Diazepam Propoxyphene with APAP Hydroxyzine Amitriptyline Cyclobenzaprine Clonidine Indomethacin | Usual care no DS - . Physicians in the control group did not receive the decision support, but the computer system tracked their prescribing. | Prescribing | NR |
| Johnson (2010) ⁷⁶ | Meds, Prescrip, Orders | NA | NA | No | No | Control - Each day, SYW output across the enterprise was turned "on" or "off" randomly for all e-prescriptions. Three pharmacies, blinded to SYW status, submitted callback logs each day. | Prescribing, Transmission, order communication | NR |
| Gilutz (2009) ⁶² | Patients | NA | Adults (19 to 44) Geriatric (65 plus) Middle age (45 to 64) | CAD dyslipidemia | No | Usual Care | Monitoring, Prescribing | NR |
| Tamblyn (2010) ¹⁶⁰ | Patients | NA | Geriatric (65 plus) Middle age (45 to 64) | CVD | anti hypertensive lipid lowering therapy | Usual care received medication list alone | Monitoring, Prescribing | NR |
| Gill (2009) ⁶¹ | Patients | NA | Adults (19 to 44) Geriatric (65 plus) Middle age (45 to 64) | hyper-lipidemia | No | Usual Care did not have the disease management tool. | Monitoring, Prescribing | Comm |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|----------------------------|---------------------------|------------------------------|-----------------|---|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Bailey (2007) ⁷ | CDSS/ CDS/ CCDS/ reminder | Hospital information system | 853 patients | compliance rates - pts discharged on a full-complement regimen of secondary prevention medications*, compliance rates- ACE inhibitor*, compliance rates- statins*, compliance rates-aspirin, compliance rates-beta-blockers | When individual drug class exclusions were considered, compliance rates increased for pts discharged on a full-complement regimen of secondary prevention medications (70.3% vs. 83.6%, RRR -19% , p<.001). Compliance rates for ACE inhibitor (83.6 vs. 89.9, RRR - 8%, p = 0.01) and statin use (89.3 vs. 94.2%, RRR 5%, p = 0.02) were significantly higher, while rates for aspirin (96.5% vs. 96.4%, RRR 0%, p = 0.95) and beta-blockers (91.8% vs. 95.9%, RRR -5%, p = 0.08) remained the same. | + | No | NA | NA |

The outcome columns (+/-) indicates whether at least 50% of the relevant outcomes abstracted were positively impacted by the MMIT (+) or not (-).

*indicates outcomes noted as being the primary outcome by the paper's authors

A1c = hemoglobin A1c; ACE= Angiotensin Converting Enzyme; ADE= Adverse Drug Event; AHR= Airway Hyperresponsiveness; AQLQ= Asthma Quality of Life Questionnaire; BMD= Bone Mineral Density; BMI= Body Mass Index; BP = Blood Pressure; CCDS= Computerized Clinical Decision Support; CCs= Care Considerations; CDS= Clinical / Computerized Decision Support ; CDSS= Clinical Decision Support System; CI= CI; COPD= Chronic Obstructive Pulmonary Disease; CPOE= Computerized Provider Order Entry; DBp = Diastolic Blood Pressure; DHCp = Decentralized Hospital Computer Program; DVT= Deep Vein Thrombosis; ED= Emergency Department; EHR= Electronic Health Record; EMR= Electronic Medical Records; e-RX= Electronic Prescribing; FEV1= Forced Expiratory Volume in the first second.; GPs= General Practitioners; HbA1c= Glycated hemoglobin; HMG Co-A= 3-hydroxy-3-methylglutaryl-coenzyme A; INR= International Normalized Ratio; LDL= Lowdensity Lipoprotein; MCID= Minimal Clinically Important Difference; MM= Medication Management; mmHg= millimeter of mercury; mmol/l= millimoles per litre; MMR= Measles, Mumps and Rubella; NPs= Nurse Practitioners; NS= Not specified; NSAID= Nonsteroidal anti-inflammatory drug; OR= OR; p = Probability; PDA= Personal Digital Assistants ; POE= Provider Order Entry; QoL= Quality of Life; RCT= Randomized Controlled Trial; RR= Relative Risk; RRR Relative Risk Reduction; SBP = Systolic Blood Pressure; SD= Standard Deviation; SF-36= Short Form 36; UTI= Urinary Tract Infection; vs.= Versus; yr= Year

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-----------------------------|---------------------------|------------------------------|--------------------------------|--|--|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Bell (2010) ¹¹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 19,450 patients | proportion of children with asthma having at least 1 prescription for controller medication*, proportion of children with asthma having an up-to-date asthma care plan*, proportion of children with asthma having spirometry performed* | Increases in the number of prescriptions for controller medications, over time, was 6% greater (p = 0.006) and 3% greater for spirometry (p = 0.04) in the intervention urban practices. Filing an up-to-date asthma care plan improved 14% (p = 0.03) and spirometry improved 6% (p = 0.003) in the suburban practices with the intervention | + | No | NA | NA |
| Berner (2006) ¹² | CDSS/ CDS/ CCDS/ reminder | Handheld | 59 internal medicine residents | proportion of unsafe NSAID prescribing per physician at follow-up | The proportion of cases per physician with unsafe NSAID prescriptions were similar at baseline for control (0.29) and intervention residents (0.27). At follow-up, the rates were statistically different, with lower proportions for intervention residents after adjustment for baseline rates (0.45 control vs. 0.23 intervention, p<0.05). Control group prescribing degraded over time while the intervention group was stable. | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|---------------------------------|---------------------------|------------------------------|-----------------|---|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Bertoni (2009) ¹⁴ | CDSS/ CDS/ CCDS/ reminder | Handheld | 8,878 patients | adherence to guideline-screening*, adherence to guideline-appropriate lipid management* | There was no difference in screening rates between the CDSS-PDA group and the control. The control group had a 10.8% drop in appropriate management from baseline, while the PDA group had a 1.1% drop, p<0.01. Stable adherence was observed in the PDA intervention group, whereas a decline in guideline adherence was observed in the control group. | - | No | NA | NA |
| Bloomfield (2005) ¹⁵ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 9,105 patients | rate of prescription -lipid therapy (before-after), rate of prescription -lipid therapy among prompt groups | Rate of lipid therapy prescriptions after implementation of the prompts: intervention clinics (8.3%) control (39.1%), RRR -371, p<0.0001 Prescription rates: 40.7% for progress notes, 36.9% for patient letters, 39.4% for reminders (p = 0.60, NS). Alternative logistic regression analysis: significant interaction between group and site, indicating that the efficacy of prompts differed by site. | + | No | NA | NA |
| Christakis (2001) ²³ | CDSS/ CDS/ CCDS/ reminder | online prescription writer | 38 providers | change in the frequency of antibiotic prescription* | Providers in the intervention arm had a 44% change in the frequency with which they prescribed antibiotics for <10 days, whereas providers in the control arm had a 10% change. Change in behavior was significantly related to the intervention, although both groups improved (p<0.01). | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-------------------------------|---------------------------|--------------------------------|--------------------------------|---|--|-----|--|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Cobos (2005) ²⁵ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 2,221 patients | proportion of patients prescribed lipid lowering drugs (secondary) | The proportion of patients prescribed lipid lowering drugs was significantly lower in the CDSS guideline intervention group (59.1% vs. 40.8%, RRR 31%, p<0.0001). | + | Proportion of patients achieving successful lipoprotein-cholesterol goals or cardiovascular risk reassessment* | The proportion of patients achieving success in the ITT analysis was similar between usual care and intervention groups (50.5% vs. 54%, NS). | - |
| Davis (2007) ³⁰ | e-prescribe CDSS | CPOE/POE system EHR/EMR system | 44 health care providers | changed physician behavior in accordance with the intervention message screens* | Prescribing behavior in accordance with the evidence improved only marginally, by 1% in control group and 4% in the intervention group (absolute difference 3%, 95% CI 1% to 15%). | + | No | NA | NA |
| Demakis (2000) ²¹³ | CDSS/ CDS/ CCDS/ reminder | Hospital information system | 12,989 patients 275 physicians | adherence rates for 5 medication management standards of care* | Adherence rates for medication management standards of care were NS different for monitoring warfarin treatment; treatment of atrial fibrillation with warfarin, aspirin or ticlopidine; treatment of myocardial infarction with beta-blockers or switching NSAID therapy for gastrointestinal bleeds. There was a large effect for pneumococcal vaccination (12.7% vs. 4.3%; OR, 3.26; 95% CI, 2.09 to 5.09). Overall, for 13 standards including non-medicinal preventive care actions, adherence was significantly improved (53.5% vs. 58.8%, OR 12.4; 95% CI 1.08 to 1.42, p = 0.002). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--|---|--|--------------------------------|---|--|-----|--|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Dexter (2001) ³³ | CDSS/ CDS/ CCDS/ reminder | Imaging systems, Pharmacy | 3,416 patients | proportion compliance - pneumococcal vaccination* , proportion compliance - influenza vaccination* , proportion compliance - subcutaneous heparin , proportion compliance -aspirin at discharge | The use of the reminders led to a higher ordering rate all 4 preventive therapies for eligible patients; pneumococcal vaccination (0.8% vs. 35.8%, RRR -4375%, p<0.001) Influenza vaccination (1.0% vs. 51.4%, RRR -5040%, p<0.001) subcutaneous heparin (18.9% vs. 32.2%, RRR -70%, p<0.001) aspirin at discharge (27.6% vs. 36.4%, RRR -32%, p<0.001). | + | No | NA | NA |
| Dexter (2004) ³⁴ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system | 1,677 patients | rate of receipt of vaccination - influenza*, rate of receipt of vaccination - pneumococcal* | Pts in the standing order group received both vaccinations more often than patients in the pop-up reminder group; for the influenza vaccine 30% reminder vs. 42% standing order, p < 0.001; for the pneumococcal vaccine 51% vs. 31%, p < 0.001. | + | Length of stay | The median length of stay for the first hospitalization were identical between the 2 study groups. | NA |
| Evans (1994) ³⁹ | CDSS/ CDS/ CCDS/ reminder, CPOE/ POE system | EHR/EMR system, Laboratory system | 482 cultures from 451 patients | rate of prescribing antibiotics to which all of the isolated pathogens were susceptible | The computer group had a higher rate of prescribing antibiotics to which all of the isolated pathogens were susceptible (77% vs. 94%, RRR 22%, p< 0.001). | + | No | NA | NA |
| Feldstein (2006) ²¹⁵ Smith (2009) ²¹⁶ | CDSS/ CDS/ CCDS/ reminder | Billing/ administration system, EHR/EMR system, Pharmacy | 961 patients | rates of completing lab monitoring* | Pts in the EMR group were 2.5 times more likely than patients in the Usual Care group to complete laboratory monitoring (p< 0.001), patients in the Automated telephone Voice Message group were 4.1 times more likely (p< 0.001), and patients in the pharmacy team outreach group were 6.7 times more likely (p< 0.001). | + | Chronic Disease score, also number of patients with abnormal lab results, so needed some actual lab results though not reported. | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|-----------------------------------|---|--|---|-----|--|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Feldstein (2006) ⁴⁰ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, Laboratory system | 9,910 patients with 239 care providers in 15 primary care clinics | interacting prescription rate (/10,000 warfarin users/month) slope for interacting prescription rate | Overall interacting prescription rate decreased immediately after the alerts were implemented, with an estimated reduction of 329.7 interacting prescriptions per 10,000 warfarin users in the first month (p = 0.002). The alerts also significantly changed the trend in the interacting prescription rate, with a preintervention increasing rate of 1.1 and a postintervention decreasing rate of 21.3 (slope change, - 22.4; p = 0.01). Academic detailing did not have an effect on interacting prescription rates. | + | No | NA | NA |
| Feldstein (2006) ⁴¹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, Laboratory system | 311 women | rate of completion of BMD or medication for osteoporosis | The control group had fewer women who had BMD completer or medication for osteoporosis compared with the reminder and reminder plus education groups (5.9% control, 51.5% reminders, and 33% reminders and education, p< 0.01 for both comparisons with control RRR for reminders alone 690% and RRR for reminders and education 460%). The same pattern was evidence for medication only (5.0% for control, 27.7% for reminders and 20.2% for reminders plus education; p< 0.01 for comparisons with control. | + | Yes - BMD, Charlson Comorbidity Index, patients weight and a satisfaction questionnaire at baseline and 6 months | BMD test and Charlson Comorbidity Index and actual weight - results not reported. Satisfaction questionnaire showed no significant differences. | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|------------------------------|---|-----------------------------------|---|--|--|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Field (2009) ⁴² | CDSS/ CDS/ CCDS/ reminder, CPOE/ POE system | EHR/EMR system | 833 patients 10 physicians 213,967 patient days | proportion of appropriate orders*, proportion of inappropriate drugs avoided | The proportion of appropriate antidepressant order rates for patients with renal insufficiency was higher in the CDSS group (52% vs. 63%, OR 1.2, 95% CI 1.0 to 1.4). More inappropriate drugs were avoided (15% vs. 46%, OR 2.6, CI 1.4 to 5.0). Improvements were seen in frequency and missing information but not for doses in the CDSS group. | + | No | NA | NA |
| Fiks (2009) ⁴³ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system EHR/EMR system | 22,586 patients | rates of up-to-date influenza vaccination*, rates of captured opportunities for vaccination* | Rates of up-to-date influenza vaccination increased from 44.2% to 48.2% at control sites and from 45.0% to 53.0% at intervention sites, a 4.0% (95% CI -1.3% to 9.1%) NS. Overall rates of captured opportunities for vaccination increased 3.8%, from 12.3% to 16.1%, at control sites and 4.8%, from 14.4% to 19.2%, at intervention sites, a difference of 1% (95% CI -2.4% to 4.9%). | - | No | NA | NA |
| Filippi (2003) ⁴⁴ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system EHR/EMR system | 15,343 patients | antiplatelet drug treatment | Number of treated patients was significantly increased in the intervention group (OR 1.99, 95% CI 1.79 to 2.22) | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|------------------------------|---|--|--|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Flottorp, (2002) ⁴⁷ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 26,826 Consults Actual- 18,106 patients, 113 practices completed. | use of antibiotics for sore throat, use of antibiotics for UTI | Pts in the sore throat group were 3% less likely to receive antibiotics after the intervention (49.5% vs. 43.8%, p = 0.032). Those for UTI were 43.4% vs. 46.3%, p = 0.639. Women with symptoms of UTI in the intervention group were 5.1% less likely to have a laboratory test ordered (55% vs. 49.8%, p = 0.046). For the sore throat, the numbers were 39.7% vs. 42.0%, p = 0.638. The absolute increase in the proportion of telephone consults for sore throat was 1.2% greater in the control group than in the intervention group (14.1% vs. 12.9%, p = 0.128). The proportion decreased for UTI (18.9% vs. 19.8%, p = 0.874). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|------------------------------|---------------------------|------------------------------|--------------------------------|--|--|-----|-----------------------|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Fortuna (2009) ⁴⁸ | CDSS/ CDS/ CCDS/ reminder | eRx | 257 clinicians | relative risk of prescribing heavily marketed medications* | The relative risk of prescribing heavily marketed medications in the alert-group during the intervention period was less than in the usual-care group (RRR 0.74; 95% CI 0.57 to 0.96; p = 0.02). The RR of prescribing heavily marketed hypnotics in the alert-plus-education group was less than in the usual-care group (RRR 0.74; 95% CI 0.58 to 0.97, p = 0.03). The prescribing of heavily marketed medications was similar in the alert-only group and the alert-plus-education group (RRR 1.02; 95% CI 0.80 to 1.29; p = 0.90). | + | No | NA | NA |
| Frances (2001) ⁴⁹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system Pharmacy | 63 physicians and 730 patients | receiving aspirin*, History of MI and receiving beta-blocker*, Receiving cholesterol-lowering agent* | 1. the proportion of patients who had an active prescription for aspirin 37.9% vs. 35.1%, RRR 7%, p = 0.440, NS; 2. the proportion of patients with MI who had an active beta-blocker prescription 22.2% vs. 33.3%, RRR -50%, p = 0.465, NS; 3. the proportion of patients receiving a cholesterol-lowering agent 73.2 % vs. 71.0%, RRR -15% p = 0.512. | - | LDL level <100 mg/dL* | The proportion of patients with a level of LDL cholesterol in the desired range (< 100 mg/dL) Did not improve cholesterol management in patients (73.2 % vs. 71.0%, p = 0.512) with CAD. | - |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--|---------------------------|------------------------------|----------------------------------|--|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Frank (2004) ⁵⁰ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 10,507 patients 10 physicians | proportion of opportunities taken for preventive activity* | Reminders did not improve adherence to MMR and flu vaccinations, but there was a significant increase in tetanus immunization (1.5% vs. 2.8%, RR 1.89, 95% CI 1.59 to 2.25). and pneumococcal immunization rates (1.6% vs. 2.8%, RR 1.70, 95% CI 1.10 to 2.62). Two of 8 non-medication related preventive care recommendations were significantly improved as well. | + | No | NA | NA |
| Fretheim (2006) ⁵⁶ Fretheim (2006) ⁵⁷ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 139 practices 501 physicians | thiazides prescription rates*, rates of cardiovascular risk assessment , proportion of patients achieving treatment goal at 3 months | Prescribing of thiazides increased in the reminders + group (11% vs. 15%, RRR 54%, p < 0.001, RR 1.94 95% CI 1.49 to 2.49). The groups did not differ for cardiovascular risk assessment (RR 1.04, 95% CI 0.60 to 1.71) or proportion that achieved treatment goal at 3 months (RR 0.98, 95% CI 0.93 to 1.02). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-----------------------------|------------------------------------|--|--|---|---|-----|----------------------|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Gill (2009) ⁶¹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 64,150 patients | up-to-date lipid test*, Lipid medication if not at goal (high risk patients only)* | Outcomes improved for most measures from before to 1 year after the intervention (univariate analysis). However, after controlling for confounding variables and for clustering in multilevel modeling, only up-to-date lipid testing for high-risk patients was statistically better in the intervention group as compared to the control group (ARR 15.0, p<0.05). Intervention status was NS for any other analysis. | - | Lipids at goal* | After controlling for confounding variables and for clustering in multilevel modeling, the proportion of patients with lipids at goal was NS between control and intervention groups. | - |
| Gilutz (2009) ⁶² | CDSS/ CDS/ CCDS/ reminder | Hospital information system, Laboratory system, Pharmacy | 7,448 patients from 56 control and 56 intervention clinics | rate of adequate monitoring, positive treatment trend, overall up titration rate in patients with LDL = 110 mg/dl | A higher rate of adequate monitoring was documented in the intervention arm (54.8% vs. 48.7%, p<0.001). Medication initiation or up-titration was recommended for patients with LDL levels above 110 mg/dl. The results showed that overall positive trends were minimally more prominent in the intervention arm (59.1% vs. 53.7%, p< 0.003). This difference constitutes a higher rate of drug initiation (2.5%), up-titration (1.8%) and avoiding drug cessation (1.1%). However, overall up titration in patients with LDL = 110 mg/dl was poor, both in the intervention arm and in the control arm (8.6% vs. 7.4%, NS). | + | LDL level reduction* | In the group of patients with initial LDL levels above 120 mg/dl a significant decrease in LDL levels was observed in the two groups, which was minimally more pronounced in the intervention arm (from 145.5 ±22.3 mg/dl to 121.9 ± 34.2, mg/dl, 16.2% reduction) than in the control arm (from 145.8 ± 22.9 to 124.3 ± 34.6, 14.8% reduction; p< 0.02). | + |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--|---|------------------------------|-----------------|--|---|-----|---|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Graumlich (2009) ²³⁷ Graumlich (2009) ²³⁸ | CDSS/ CDS/ CCDS/ reminder CPOE/ POE system | CPOE/POE system | 631 patients | patient mean score for discharge preparedness*, patient score for satisfaction with medication information, outpatient physicians perception of discharge software | When comparing patients assigned to discharge software vs. usual care, patient mean (standard deviation [SD]) scores for discharge preparedness were higher (17.7 [4.1] vs. 17.2 [4.0]; p = 0.042), patient score for satisfaction with medication information were unchanged (12.3 [4.8] vs. 12.1 [4.6]; p = 0.567). and their outpatient physicians scored higher quality discharge (17.2 [3.8] vs. 16.5 [3.9]; p = 0.027). Hospital physicians found mean effort to use discharge software was more difficult than the usual care (6.5 [1.9] vs. 7.9 [2.1]; p = 0.011) and discharge software users had mean (SD) satisfaction 7.4 (1.4) vs. 7.9 (1.4) for usual care physicians; p = 0.129. | - | Readmitted within 6 months*, emergency department visit within 6 months, adverse events within 1 month. | When comparing patients assigned to discharge software vs. usual care, there was no difference in hospital readmission within 6 months (37.0% vs. 37.8%; coefficient 0.005 95% CI - 0.074 to 0.065 p = 0.894), emergency department visit within 6 months (35.4% vs. 40.6%; coefficient 0.052, 95% CI -0.115 to 0.011; p = 0.108), or adverse events within 1 month (7.3% vs. 7.3%; coefficient 0.003, 95% CI -0.037 to 0.043, p = 0.884). | - |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---|--|---------------------------|-------------------|---------|-----|--|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Gurwitz (2008) ³⁰² | CDSS/ CDS/ CCDS/ reminder | Integrated CPOE/ POE system, Laboratory system | 1,118 residents/ patients | No | NA | NA | ADE rates per 100 resident months* | similar for control and intervention units (10.4 vs. 10.8, NS). The same was found for the rate of preventable ADEs per 100 resident months (3.9 vs. 4.0, NS). Hospitalization measured. | - |
| Hetlevik (1999) ³⁰⁴ | CDSS/ CDS/ CCDS/ reminder CPOE/ POE system | EHR/EMR system | 1,998 patients | NA | NA | NA | *SBP mmHg (SD), DBP mmHg (SD), Serum Cholesterol mmol/l (SD), BMI kg/m2 (SD) | The groups did not differ for BP, cholesterol levels or BMI: SBP was 155.6 vs. 156.8 mmHG (95 % CI -0.6 to 3.0) between the control and the intervention group. DBP was 89.8 vs. 88.8 mmHg (95% CI -1.9 to -0.2). Serum cholesterol was 6.57 mmol/l vs. 6.64 mmol/l (95% CI -0.1 to 0.2) between the two groups. BMI was 27.7 kg/m2 vs. 27.8 kg/m2 (95% CI -0.4 to 0.07). | - |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|--|-----------------|---|---|-----|---|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Hicks (2007) ⁶⁷ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 1,422 patients | BP controlled, receiving a recommended drug class medication within 1 week of the clinic visit adjusted | This study had 4 groups: usual care, CDS, NPs, and NPs+CDS. No difference was seen across all 4 groups for BP readings: Usual care vs. CDS: 45% controlled vs. 48% controlled, OR 0.96 (95% CI 0.78 to 1.19). Pts in the CDS group were more likely to have received a recommended drug class medication within 1 week of the clinic visit than patients in the usual care group: adjusted OR 1.32 (95% CI 1.09 to 1.61). | - | Study endpoints included BP control and mean SBP or DBP at the outcome visit. | Adjusting for patients' demographic and clinical variables, the number of prior visits, and levels of baseline BP control, there were no differences between intervention groups in the odds of outcome BP control. | NA |
| Holbrook (2009) ³⁶⁷ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, Laboratory system, Personal health records systems | 511 patients | 1) Composite* 2) Adequacy of monitoring A1c, BP, lipids, foot, eye | 1) A shared electronic decision-support system to support the primary care of diabetes improved the process of care and some clinical markers of the quality of diabetes care; 2) Improvement in monitoring was seen significantly more in the intervention group than in the control group. Number of visits to the primary care provider (as recommended) increased significantly more in the intervention group than in the control group (difference of 0.66, 95% CI 0.37 to 1.02, p<0.001). 3) Satisfaction, ease of use, usefulness, preference for paper vs. computer. | + | A1c, BP, cholesterol, urine albumin, foot, eye | no significant change reported | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|------------------------------|---------------------------|------------------------------|-----------------|--|---|-----|--|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Holman (1996) ³⁰⁶ | CDSS/ CDS/ CCDS/ reminder | Handheld, Stand-Alone | 5 patients | NA | NA | NA | Glycemic control NR, therefore includes Pre-prandial blood glucose levels, A1c and fructosamine* | Pre-prandial blood glucose levels* were significantly less during the 'advice on' period compared to the 'advice off' period (7.5 vs. 8.9 mmol/l, p = 0.015) but A1c and fructosamine not changed | - |
| Javitt (2005) ²¹⁸ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 39,462 patients | compliance with recommendations to add-a-drug* | Physicians complied with 24% of these "add-a-drug" recommendations in the intervention group. In the control group, physicians spontaneously instituted the treatment that would have been recommended in 17% of instances in which the recommendation was triggered but not issued. This 42% relative difference in compliance was statistically significant (P = .007). | - | Admissions per 1,000 persons* | Among those in both groups who triggered recommendations, there were 19% fewer hospital admissions in the intervention group compared with the control group (213.8 ± 5.7 vs. 264.6 ± 5.7, p< .001). | + |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|------------------------------|--|--------------------------------------|-----------------|--|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Javitt (2008) ⁷⁵ | CDSS/ CDS/ CCDS/ reminder | Pharmacy, Laboratory system, billing | 39,508 patients | resolution rate-add a drug alert*, resolution rate-stop a drug*, resolution rate -do a test* | Resolution rate for “add a drug” CCs was 8.6 % higher in the study group than the control group (p <0.05). There was, however, no significant difference in the resolution rates for “stop a drug” CCs (change -6%, NS). Resolution rates for “do a test” CCs were 5.8% higher in the study group, p <0.05. | + | No | NA | NA |
| Johnson (2010) ⁷⁶ | CDSS/ CDS/ CCDS/ reminder e-prescribe | EHR/EMR system | 3,285 patients | rate of call backs generated* perceptions* | There was no significant difference in the call back rates between the “SYW off” and the “SYW on “ periods (0.4% vs. 0.45%; p = 0.47) Other Outcomes: perceptions* of Show your work were mostly positive trends, in the questionnaire. | - | No | NA | NA |
| Krall (2004) ⁸⁹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 1,076 patients | proportion of patients no longer eligible for alerts at the end of the month* | Following implementation of the alert, more patients were ‘no longer eligible for alerts at the end of the month’ (25.8% pre vs. 54.3% post, RRR - 103%, p <0.001). | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-----------------------------|---------------------------|---------------------------------|-----------------|-------------------|---------|-----|--------------------|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Kucher (2005) ⁹⁰ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, CPOE/POE system | 2,506 patients | No | NA | NA | *DVT, PE, bleeding | Clinically diagnosed DVT at 90 days Control: 103 (8.2%) Intervention: 61 patients (4.9%) (RRR 40%, p = 0.001); Clinically diagnosed PE at 90 days Control: 35 (2.8%) Intervention: 14 (1.1%) (RRR 61%, p = 0.004). The groups did not differ for proximal- or distal DVT, DVT of the arms, death, or hemorrhage. | + |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|--|------------------------------|---|---|--|-----|------------------------------|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Kuilboer (2006) ²¹⁹ | CDSS/ CDS/ CCDS/ reminder CPOE/ POE system | EHR/EMR system | 32 primary care practices (78,926 patients of whom 9,798 had asthma or related symptoms) Actual: 156,772 patients and 40 GPs | rate of prescribing for cromoglycate | Prescribing for cromoglycate was reduced in the 12 to 39 year and 40 to 59 year groups (12 to 39: 9.9/1,000 patients vs. 4.1, p = 0.03) and (40 to 59: 9.0/1,000 patients vs. 4.2, p = 0.05). Other prescribing (3 drugs or drug classes and 4 age groups) did not differ across groups. | - | Peak flow measurements | Of 20 potential changes, 8 were observed: The AsthmaCritic group had more contacts for the 12 to 39 year group (p = 0.03), more measurement of peak flow total for the 0 to 11 year group (p = 0.02), more FEV1 total peak flow ratio measurement in the 12-59 year groups (p = 0.04 and 0.009), and more measurement of FEV1 rates in the 3, 12 and older groups (p = 0.01, 0.01, and 0.016). | - |
| Lester (2005) ⁹⁵ | CDSS/ CDS/ CCDS/ reminder, email message of high levels of LDL | EHR/EMR system | 235 patients 14 clinicians | proportion of patients with changes in statin prescriptions at 1 and 12 months* | At 1 month more patients in the email group had received statins than control patients (15.3%, vs. 2%, p = 0.001). At 1 year the difference in receipt of statins had disappeared (24.6% vs. 17.1%, p = 0.14). | - | LDL cholesterol also process | LDL cholesterol did not differ between group | - |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-----------------------------|---------------------------|------------------------------|--|---|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Linder (2009) ⁹⁷ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 111,820 patients, 443 physicians within 27 practices | rate of antibiotic prescribing to patients with ARI * | In the intent-to-intervene analysis, clinicians prescribed antibiotics to 43% of patients with ARI diagnoses in control clinic compared to 39% in the intervention clinic (OR. 0.8; 95% CI, 0.6 to 1.2, p = 0.30). The ARI Smart Form did not significantly reduce overall antibiotic prescribing. The smart form was used by 33% of intervention clinicians (86/262) at least once. For the as-used analysis, appropriate antibiotic prescribing rate was 88% (n = 990 visits) | - | No | NA | NA |
| Lo (2009) ²²⁰ | CDSS/ CDS/ CCDS/ reminder | Imaging systems | 3,673 potential alert trigger events Actual: 2,765 patient 366 providers | clinic, doctors office, etc. | 3,673 total events where baseline lab tests would have been advised: 1,988 events in the control group and 1,685 in the intervention group. control group: baseline labs requested for 771 (39%) of the medications. intervention group: baseline labs ordered by clinicians for 689 (41%) of the cases. No significant association existed between the intervention and the rate of ordering appropriate baseline laboratory tests (RRR 5%, p = 0.782, NS). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-------------------------------|---------------------------|------------------------------|---------------------|--|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Martens (2007) ¹⁰¹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 77 physicians (GPs) | quinolone prescriptions , inhaled corticosteroids for newly diagnosed COPD in patients > 40 yr , first choice drugs for sore throats | GPs got reminders to either stop prescribing drugs or to prescribe a specific first-line drug. No differences were seen for either group to prescribe a drug. No differences were found for those in the cholesterol reminder group. GPs in the antibiotics, asthma and COPD group showed changes in 3 of 8 drug categories. Outcome measures were for sum scores for drug volume: lower scores were improvements in prescribing. Reminder physicians prescribed fewer quinolones (4.6 (95% CI 2.8 to 8.1)) vs. (1.5 (95% CI 0.8 to 2.2)); fewer inhaled corticosteroids for COPD in newly diagnosed patients >40 yrs (0.5 (95% CI 0.3 to 0.9)) vs. (0.0 (95% CI 0 to 0.1), p = 0.00); and better first choice drugs for sore throats (0.8 (95% CI 0.3 to 2.4) vs. (0.2 (95% CI 0.0 to 0.4), p = 0.03). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|------------------------------|---|---|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Matheny (2008) ²²¹ | CDSS/ CDS/ CCDS/ reminder | Laboratory system | 2,507 outpatient visits in 1,922 geriatric patients 303 primary care physicians | rate of receiving appropriate laboratory testing within 14 days of the clinical encounter/ 10 medication lab reminder categories. | Reminders for appropriate laboratory monitoring had no impact on rates of receiving appropriate testing for creatinine, potassium, liver function, renal function, or therapeutic drug level monitoring for patients overdue for lab monitoring NSAIDs; Angiotensin Receptor Blockers; Metformin; Potassium Supplements; Potassium Sparing Diuretics, Thiazide Diuretics; Angiotensin Converting Enzyme Inhibitors; HMG Co-A Reductase Inhibitors; Thyroxine; (or the following therapeutic drugs combined: Carbamazepine; Cyclosporine, Phenobarbital, Phenytoin, Proc-NAPA, Valproate). | - | No | NA | NA |
| McDonald (1976) ²²² | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 601 patient visits by 226 patients | compliance with drug monitoring test alerts*, compliance with recommendations to change therapeutic regimens* | Alerts to patients overdue for drug monitoring tests resulted in an increased number of tests ordered (11% vs. 36%, RRR -227%, p< 0.0001). Recommendations for changes to therapeutic regimens were followed in 28% of study events compared to 13% of control events (p < 0.026). | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|-----------------------------------|------------------------------------|--|--|-----|---------------------------------|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| McGregor (2006) ¹⁰⁵ | CDSS/ CDS/ CCDS/ reminder | Pharmacy | 4,507 patients | mean time spent on antimicrobial management: | Team members spent 3.2 hours per day on management of antimicrobials with the decision support system compared with 4 hours per day without. Not statistical testing was done. | - | Yes - mortality, Length of stay | Mortality: NS All patients 3.0% vs. 3.3%, p = 0.6 or for those patients who got alerts 8.2% vs. 7.8%, p = 0.5. Length of stay: All patients: 4.0 days vs. 3.8, p = 0.04 and 5 vs. 4 days for patients with alerts, p = 0.6 (NS) | - |
| Meigs (2003) ³⁰⁹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, Laboratory system | 598 patients 26 staff providers | NA | NA | NA | HbA1c levels* | The intervention had a modest benefit on glycemic control; HbA1c levels tended to improve in the intervention group (change - 0.23) and worsen in the control group (change +0.14) NS | - |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|----------------------------------|---------------------------|--|------------------------------------|---|---|-----|---|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Montgomery (2000) ¹⁰⁸ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 552 patients | probability of patients taking 2 drugs , probability of patients taking 3 drugs | Adjusted data showed that compare with those in the risk chart group alone, those with computer support had a lower probability of patients taking 2 drugs (OR 0.5, 95% CI 0.2 to 0.9) p< 0.05) or 3 drugs (OR 0.3, 95% CI 0.1 to 0.6, p< 0.05). | - | *a five year cardiovascular risk >10%, SBP, DBP | no difference between groups with cardiovascular risk reduced below 10%. SBP and DBP were not reduced in the CDSS group (SBP 153 vs. 153 mmHg) (DBP 85 vs. 85 mmHg) | - |
| Murray (2004) ³¹⁰ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system Pharmacy | 712 patients | NA | NA | NA | SF-36 QoL* | No intergroup differences were found for the primary endpoint the SF-36 QoL* scale (Table 3). No analysis presented. | - |
| Overhage (1996) ¹¹⁷ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system EHR/EMR system, Hospital information system, Laboratory system, Pharmacy | 24 practice teams - 78 house staff | rates of compliance with preventive care recommendations* | Overall, control teams complied with 24% of the reminders compared with 23% for intervention teams (P = 0.78). When preventive care measures were analyzed individually, 2 significant differences were seen in compliance (24-hour urine protein and angiotensin-converting enzyme [ACE] inhibitor) between control and intervention teams. These were assumed to be due to chance with multiple testing and because they were in the opposite directions. | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|--|-----------------------------|---|--|-----|-------------------|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Overhage (1997) ¹¹⁸ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system EHR/EMR system Laboratory system | 86 physicians on 6 services | immediate compliance with corollary ordering*, 24 hour compliance*, hospital-stay compliance* | Intervention physicians ordered the corollary orders required by the guidelines twice as often as control physicians did when measured by immediate compliance (46.3% vs. 21.9%, RRR - 111%, p < 0.0001). Significant differences between study and control physicians also appear in 24 hour compliance (50.4% vs. 29.0%, RRR -74%, p < 0.0001) and hospital-stay compliance (55.9% vs. 37.1%, RRR 51%, p < 0.0001). | + | Length of stay | Length of stay was not different for intervention patients compared with control patients (8.12 days vs. 7.62 days, a difference of -0.5 days, 95% CI 0.17 to 1.19; p = 0.94). | - |
| Palen (2006) ¹²¹ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system, Pharmacy | 26,586 patients | compliance rate | No significant differences between group physicians in the overall rate of compliance with ordering recommended lab monitoring for patients prescribed study meds. Lab monitoring Intervention: 56.6% Control: 57.1% (p = .31). Improved compliance: Gemfibrozil 71.2% vs. 62.3% (p = .003); Statins 75.7% vs. 73.9% (p = .05), Colchicine 52.8% vs. 46% (p = 0.05); Methotrexate 42.9% vs. 0% (p = 0.03). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|---------------------------------|---|--|--|-----|--|------------------------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Paul (2006) ¹²⁴ | CDSS/ CDS/ CCDS/ reminder | Hospital information system | 3,529 patients in the RCT and 1,203 in the cohort study | appropriate antibiotic prescribing increased | Appropriate antibiotic prescribing increased for both intention to treat analyzes (64.5% vs. 72.7%, RRR 13%, p< 0,05) and for per protocol analyzes (64.5% vs. 85.1%, RRR 32%, p< 0.05). The cohort study showed similar increases in improved prescribing (57% vs. 70%, p< 0.001) | + | Yes -(secondary outcomes) - duration of stay, duration of fever, or 30-day mortality, adverse events, costs. | Mortality not affected | NA |
| Persell (2008) ²⁰⁵ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 242 patients | self-reported aspirin use* | the control rate (reminders only) of self-reported aspirin use was NS different than the intervention (reminders plus clinician emails and patient phone calls) group (39% vs. 46%, p = 0.20) | - | NA | NA | NA |
| Peterson (2007) ¹²⁶ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system, EHR/EMR system | 9,111 medication orders Actual : 778 providers 2,981 patients | ratio between prescribed and recommended doses | Ratio between the prescribed dose and recommended dose showed that compared to controls the intervention group (reminders) received lower doses (3.0 vs. 2.5, p< 0.001). | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--|---------------------------|------------------------------|-----------------|--|---|-----|--|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Plaza (2005) ²⁸⁰ | CDSS/ CDS/ CCDS/ reminder | Handheld | 198 patients | NA | NA | NA | QoL-St George's Respiratory Questionnaire* | Scores on the St George's Respiratory Questionnaire were significantly lower for intervention patients (34.1 vs. 27.3, p = 0.002, difference 6.8 (95% CI 2.5 to 11.1). % patients reaching MCID of decrease by 4 points was 65.3% I vs. 41.0% C | + |
| Prescription in Ischemic Stroke Management (PRISM) Study Group (2003) ¹²⁷ | CDSS/ CDS/ CCDS/ reminder | Hospital information system | 1,640 Pts | relative risk reduction (RRR) in ischemic and hemorrhagic vascular events* | For each patient, the CDSS was used to calculate the relative risk reduction (RRR) in ischemic and hemorrhagic vascular events which was achieved by the actual therapy prescribed vs. the option of 'no antiplatelet or anticoagulant therapy'. Estimated RRR(%) for the control and intervention in the first phase was 16.7 (13.2–23.7) vs. 16.3 (15.2–21.2) (NS different). For the second phase it was 16.3 (13.1–23.8) vs. 16.7 (13.5–22.9) (NS different). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|------------------------------|---|------------------------------|--|---|--|-----|---|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Quinn (2008) ¹²⁸ | CDSS/ CDS/ CCDS/ reminder | Glucose Meter | 30 patients | changes in medication (medication intensified) | Process Monitoring - Pts using WDS were more likely to have physicians intensify diabetes medications (84.6% vs. 23.08%, p = 0.002). | + | Average decrease in Alc values * | Average decrease in A1c for intervention patients was 2.03% compared to .68% for control patients (p<.04) | + |
| Raebel (2005) ¹³⁰ | CDSS/ CDS/ CCDS/ reminder | Laboratory system, Pharmacy | 9,565 patients with 10,169 dispensings | percentage of dispensings with baseline monitoring* | Recommended laboratory monitoring was completed in 74.7% (n= 7,598) of dispensings at initiation of therapy. Compared to the usual care group, monitoring was higher in the intervention group (70% vs. 79%, RRR - 13%, p<0.001). | + | Lab tests - alanine aminotransferase /aspartate aminotransferase ; CBC; TSH | no results given in this paper, reported directly to care provider only | NA |
| Raebel (2007) ¹³¹ | CDSS/ CDS/ CCDS/ reminder pharma info system | EHR/EMR system | 59,680 patients | new dispensing of targeted medications* | In the analysis of all dispensing of targeted medications, there was a significant reduction of new dispensing of at least one targeted medication (2.2% vs. 1.8%, RRR 16%, p<0.002). For dispensing of targeted medications considered inappropriate, there was also a significant reduction with the use of the alerting system (1.5% vs. 1.1%, RRR 27%, p<0.001). | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-------------------------------|---------------------------|---------------------------------------|--|---|---|-----|-------------------------|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Raebel (2007) ¹³² | CDSS/ CDS/ CCDS/ reminder | Hospital information system, Pharmacy | 11,100 women | proportion of pregnant women dispensed a category D or X medication*, total number of first dispensings of targeted medications | The alerts resulted in a significant 47% reduction in the proportion of pregnant patients receiving category D or X drugs (p<0.001). Intervention patients received 238 dispensings of unique targeted medications and usual care patients received 361 dispensings (p = 0.03). The study was stopped primarily due to 2 false-positive alert types: Misidentification of medications as contraindicated in pregnancy by the pharmacy information system and misidentification of pregnancy related to delayed transfer of diagnosis information. | + | No | NA | NA |
| Rollman (2002) ¹³⁷ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 200 Pts with documented major depression | antidepressant prescribing rate (secondary) | Prescribing of antidepressants (continuous use of change in prescriptions) did not differ across the 3 groups at 3 or 6 months. | - | Mean depression scores* | All groups improved their mean depression scores at 3 and 6 months. However the groups did not differ from each other in mean scores at 3 or 6 months. | - |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|----------------------------------|--|--------------------------------|--|---|---|-----|----------------------|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Rood (2005) ¹³⁸ | CDSS/ CDS/ CCDS/ reminder | Hospital information system | 484 patients | adherence to glucose measurement timing recommendations*, adherence to insulin dose advice* | Rate of compliance with glucose measurement timing recommendations control-intervention-control (29% vs. 38% vs. 41% with period 2 and 3 greater than period 1, p = 0.05). During the intervention period the rate for computerized group was higher than the control (36% vs. 40%, p = 0.05) Rate of compliance with insulin dose advice was higher in period 2 than 1, and then decreased significantly in period 3 (56% vs. 70% vs. 42%, p = 0.05). During the intervention period the rate for computerized group was higher than the control (64% vs. 77%, p = 0.05) | + | Glucose measurements | Glucose measurement actual levels NR. Measured the number of times the levels fell within normal range. | NA |
| Rosenbloom (2005) ²⁶² | CDSS/ CDS/ CCDS/ reminder CPOE/ POE system | CPOE/POE system EHR/EMR system | 418,739 opportunities to access an information item 147 house staff | access rate for educational opportunities | Study physicians accessed educational opportunities for 278 of 240,504 (0.12%) vs. 18 of 178,35 opportunities (0.01%), RRR 1100, p< 0.05. | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|------------------------------|---|--|-----------------|-------------------------------------|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Rosser (1992) ² | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 8,069 patients | rate of tetanus toxoid vaccination* | The rates of tetanus toxin given were 3.2% in control, 22.8% in physician reminder, 24% in telephone reminder, and 30.6% in the letter reminder. The differences in the recorded vaccination rate between the randomized control group and the three reminder groups are as follows: 19.6% in the physician reminder group (95% CI 17.1 to 22.2, p < 0.00001), 20.8% in the telephone reminder group (95% CI 18.3 to 23.5, p < 0.00001) and 27.4% in the letter group (95% CI 24.8 to 30.2, p < 0.00001). | + | No | NA | NA |
| Rotman (1996) ²⁶⁴ | CDSS/ CDS/ CCDS/ reminder e-prescribe | Hospital information system, Laboratory system | 34 Physicians | user Satisfaction Rating* | After the physicians used the PWS, their user-satisfaction, score decreased by 0.34 Likert-scale units (approximately one half of one standard deviation of the mean score, p = 0.008). In contrast, the mean satisfaction in the control group (DHCP) increased by 0.49 Likert-scale units (p < 0.0001). Overall, the two groups diverged with a difference of 0.83 Likert-scale units between the two groups (p < 0.0001). | - | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--|---------------------------|------------------------------|-----------------|----------------------|--|-----|---|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Roumie (2006) ²²⁶ Roumie (2007) ²²⁷ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 871 patients | prescribing changes* | No differences were seen comparing the groups who had provider education alone vs. those who had provider education and computer alerts for prescribing of any medication, changing doses, or adding medications (all data adjusted for multiple variables). | - | Yes -proportion of patients with controlled hypertension* | Pts of providers randomly assigned to the patient education group had better BP control (138/75 mm Hg) than those in the provider education (146/76 mm Hg) and alert or provider education alone (145/78 mm Hg). The patient education group had a SBP of 140 mm Hg or less compared with those in the provider education or provider education and alert groups ARR 1.31 (95% CI, 1.06 to 1.62; p = 0.012). | - |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--|--|--|-----------------------|--|--|-----|---|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Safran (1995) ¹⁴¹ Safran (1993) ¹⁴² | CDSS/ CDS/ CCDS/ reminder, CPOE/ POE system | EHR/EMR system | 349 patients with HIV | mean response time to alerts* | Physicians who got alerts responded more quickly to them (mean 52 vs. 11 days, $p < 0.0001$). Physicians who got reminders responded more quickly to them (mean 500 vs. 114 days, $p = 0.0001$). | + | Rate of emergency visits, rate of non-primary care visits, hospitalizations, deaths | Pts in the control group had a higher rate of hospitalizations than those in the intervention group (44% vs. 35%, RRR 20%, $p = 0.04$) but other outcomes negative. Mortality not affected. | - |
| Sequist (2005) ²²⁸ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, Imaging systems, Laboratory system | 6,243 Pts | compliance rate with Diabetes reminders* , compliance rate with Coronary Artery Disease reminders* | Diabetes reminders resulted in the recommended action in 19% of patients in the intervention group vs. 14% of patients in the control group. After adjusting for baseline patient and physician characteristics, patients in the intervention group were more likely than control patients to receive recommended diabetes care based on the composite outcome (OR] 1.30, 95% CI 1.01 to 1.67). CAD reminders resulted in the recommended action for overdue items: Intervention 22% Control: 17% Pts in the intervention group received recommended CAD care more often than those in the control group (OR 1.25, 95% CI 1.01 to 1.55) after adjusting for baseline differences. | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|--------------------------------|---------------------------|------------------------------|---------------------------|--|---|-----|-------------------|---|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Shojania (1998) ¹⁴⁸ | CDSS/ CDS/ CCDS/ reminder | Imaging systems Pharmacy | 396 physicians | number of vancomycin orders/ prescriber*, mean duration of treatment prescribed per physician*, mean number of days of vancomycin per course of treatment* | The total number of orders for vancomycin for physicians in the control group was higher than in the intervention group (16.7 vs. 11.3 orders per physician, p = 0.04). Physicians in the intervention group prescribed vancomycin for 36% fewer days than physicians in the control group (26.5 vs. 41.2, p = 0.05). The number of days of vancomycin per course of treatment was also lower for the physicians in the intervention group, mean of 1.8 d vs. 2.0 for the control group (p = 0.05). | + | Length of stay | There was no significant differences between the groups with respect to the average length of stay. | |
| Tamblyn (2003) ¹⁵⁹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 12,560 Pts 107 physicians | rate of initiation of inappropriate drugs per 1,000 visits, Rate of discontinuation of inappropriate drugs per 1,000 | During the study the number of new potentially inappropriate prescriptions per 1000 visits was lower (52.2 v 43.8) in the CDS group than in the control group (RR 0.82, 95% CI, 0.69 to 0.98). The rate of discontinuation of inappropriate drugs per 1,000 was not different: 67.4 vs. 71.4, RR (95% CI 1.06, 0.089 to 1.26) | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-------------------------------|---------------------------|-------------------------------------|----------------------------|--|---|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Tamblyn (2010) ¹⁶⁰ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, Insurance databases | 2,293 patients | rate of drug profile review, Changes in therapy | Process Monitoring - Significant increase in drug profile review in the intervention compared to the control group (44.5% vs. 35.5%;p<0.001). There was NS difference between the intervention and control group in the proportion of patients who had increases in therapy (28.5% vs. 29.1%; OR, 0.98; p = 0.86). | + | No | NA | NA |
| Terrell (2009) ¹⁶⁴ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system | 5,162 Pts 63 physicians | proportion of ED visits by seniors with an inappropriate medication*, proportion of medications that were potentially inappropriate was also reduced | The decision support reduced the proportion of ED discharges that resulted in potentially inappropriate prescriptions (3.9% vs. 2.6%; p = 0.02; OR = 0.55, 95% CI 0.34 to 0.89). The proportion of medications that were potentially inappropriate was also reduced, from 5.4% to 3.4% (p = 0.006; OR 0.59, 95% CI 0.41 to 0.85). | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-------------------------------|------------------------------------|---------------------------------|--|---|--|-----|-------------------|---------|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Terrell (2009) ¹⁶⁵ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system, CPOE/POE system | 63 physicians had 5,162 patient visits | visits with an inappropriate medication prescription* , prescriptions that were inappropriate | Primary Outcome: Decision support significantly reduced the proportion of ED discharges that resulted in a potentially inappropriate prescription (3.9% vs. 2.6%; p = 5.02; OR 50.55, 95% CI 50.34 to 0.89). This difference represents an ARR of 1.3% (95% CI 50.4 to 2.3%). Secondary Outcome: When analyzed as a percentage of all medications prescribed by physician subjects, the proportion of medications that were potentially inappropriate was significantly reduced, from 5.4% to 3.4% (p = 5.006; OR 50.59, 95% CI 50.41 to 0.85), with an ARR 2.0% (95% CI 50.7 to 3.3%). | + | No | NA | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-------------------------------|---------------------------|--|---|------------------------------------|---|-----|--|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Tierney (2003) ¹⁶⁶ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 706 patients, 20 pharmacists, 94 physicians, 1 nurse practitioner | NA | NA | NA | Quality of Life - SF-36*, heart failure exacerbations* | Across the 4 groups (physician intervention, pharmacist intervention, both interventions, and controls) no differences were seen for the SF-36 (8 subscales), or for Heart Failure exacerbations (4 subscales) and emergency department visits or hospitalizations (all or related to HF). | - |
| Tierney (2005) ¹⁶⁷ | CDSS/ CDS/ CCDS/ reminder | CPOE/POE system EHR/EMR system, Pharmacy | 706 patients | adherence to the care suggestions* | There were no differences between the four study groups in either adherence to the care suggestions, combined or individually (32% control, 32% physician intervention, 32% pharmacist intervention, 37% both interventions, NS). | - | Quality of Life - SF-36, Chronic Respiratory Disease Questionnaire, hospitalizations | Change in Quality of life measures (NS). Hospitalization was measured and not affected. | NA |

Evidence Table 15. KQ7: integrated CDSS study characteristics: results (continued)

| Author (year) | MM System studied | Systems CDSS Integrated with | Number Analyzed | Process | | | Clinical | | |
|-------------------------------|---------------------------|------------------------------|-------------------------------------|---|---|-----|-------------------|--|-----|
| | | | | Outcomes Measured | Results | +/- | Outcomes measured | results | +/- |
| Van Wyk (2007) ¹⁷¹ | CDSS/ CDS/ CCDS/ reminder | EHR/EMR system | 87,860 Pts 77 physicians | percentage of patient treated | Of the patients requiring treatment, 66% were treated in alerting arm, 40% in on-demand arm, and 36% in control arm. After adjustment for differences between arms, likelihood of being treated was 40% higher in alerting arm (adjusted RR 1.40; 95% CI 1.15 to 1.70) and 19% higher (NS) in on-demand arm in comparison to the control arm (adjusted RR 1.19; 95% CI 0.94 to 1.50). A similar pattern was shown for the need for screening within the 3 groups. | + | No | NA | NA |
| White (1984) ²²⁹ | CDSS/ CDS/ CCDS/ reminder | Imaging systems | 396 patients | physician actions* | Physicians were 1.22 times as likely to take action in the alert group as compared to the non-alert group (p < 0.003). Actions included medication and lab monitoring changes. | + | No | NA | NA |
| Zanetti (2003) ¹⁸⁰ | CDSS/ CDS/ CCDS/ reminder | Hospital information system | 273 patients having cardiac surgery | more patients in the alarm plus reminder group received appropriate redosing of antibiotics after > 240 minutes in surgery. | More patients in the alarm plus reminder group received appropriate redosing of antibiotics after > 240 minutes in surgery (adjusted OR 3.31, 95% CI 1.97 to 5.56, p< 0.0001). | + | Rate of infection | Rate of infection Intervention: 4% Control: 6% (p = 0.4) lower than before the study (p = 0.2) | - |

Evidence Table 16. Article references for studies across the phases of medication management (and education and reconciliation) by research design

| Design | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|---------------|---|---|---------------------------------------|--|--|-----------|-----------------------|
| Cohort | 87,106,121,124,138,169,268,281,292,293,297,300,317 | 188,190 | 188,190 | 203 | 267,275,287,292,296,301 | | 233 |
| Observational | 16,26,27,29,44,59,81,91,93,97,113,128,135,144,146,153,155,158,162,176,235,244,253,278,375-377 1,3,50,67,73,77,78,85,99,140,151,172-174,177,242,277,280,295,305,368,369 17,19,34,58,72,80,82,108,110,115,119,150,152,154,157,168,236,266,269,286,294,329 2,4,6,11,14,15,18,20,25,33,43,46,57,65,69,70,83,84,105,109,111,156,163,274,276,315 22,35,36,63,64,71,95,98,112,122,133,145,147,161,175,179,246,260,261,270,284,288,298,316 7,30,62,76,79,86,92,101-103,114,134,143,149,170,178,250,259,271,290,291,307,308 | 65,99,172,182-187,189,191-194,243,265,303,375 | 65,99,191,195,197,234,265,278,303,375 | 46,50,99,186,187,197-202,204,206-210,234,240,253-255,278,280,305,375 | 1,19,57,84,105,113,177,214,223,225,283,285,286,295,299,315,329,375 36,133,143,147,211,212,224,261,298,311,314 | 249,341 | 230-232,245 |

Evidence Table 16. Article references for studies across the phases of medication management (and education and reconciliation) by research design (continued)

| Design | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|---------------|--|----------------------------|-------------------|--|---|------------------|------------------------------|
| Qualitative | 90,181,248,256,273,319,320,322, 330,336,337,339,340,342,344, 347,349,350,356,360,363,364, 366,370,371 247,318,332,334,338,341,345, 354,357,359,361,365 | 258,336,340,342,348 | 348,351,353 | 239,241,272,331,333,343, 346,352,355,362 | 251,335,347,356,358 | | |
| RCT | 5,10,13,24,38- 40,55,66,74,89,94,100,104,120, 123,125,129- 132,166,167,262,302 8,21,23,28,31,32,37,42,45,48,88, 107,117,136,148,180,306,310 9,12,41,47,49,60,61,75,96,116, 118,126,127,137,139,141,159, 160,164,165,171,237,252,264, 289,304 | 75 | 130,196 | 205,306 | 23,40,66,89,94,104, 107,117,120,129,132, 136,166,167,215,217, 219- 221,226,263,306,312, 367 12,60,61,127,137, 141,213,218,222,228, 229,279,309 | 127 | 313 |

Evidence Table 17. Article references for studies across settings for the phases of medication management (and reconciliation and education)

| Setting | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|--|--|------------------------|------------|---------------|---|-----------|-----------------------|
| Ambulatory care | 10,13,29,38,39,55,67,73,74,82,90,94, 100,113,120,129- 132,136,140,154,157,162,166,167, 172,236,242,244,266,269,273,281, 336,337,342,349,360,370 15,20,21,23,28,41,42,45,48,83,87,95, 107,118,139,147,161,171,260,270, 276,298,304,310,320,347,356,363 9,12,30,47,49,60- 62,75,92,96,106,127,141,143,145, 159,160,259,264,291,307,332,357, 361,365 | 75,172,185,191,336,342 | 130,191 | 205 | 23,94,107,113,120,129, 132,136,166,167,215, 217,219- 221,226,263,275,301, 312,347,367 12,60,61,127,141,143, 147,213,218,228,251, 267,279,298,309,335, 356,358 | 127,321 | |
| Community (school, community centre etc) | | | | | 228 | | |
| Home | 113,306 | | | 306,343 | 113,217,306,335,367 | | |

Evidence Table 17. Article references for studies across settings for the phases of medication management (and reconciliation and education) (continued)

| Setting | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|--------------------------------|---|--|---|---|---|-----------|-----------------------|
| Hospital | 16,26,27,59,73,78,81,85,91,93,97, 121,125,128,135,144,146,151,153, 155,158,174,176,177,235,253,256, 277,278,293,305,319,339,340,344, 349,366,368,375,376 1,3,5,17,19,24,34,50,58,72,77,80,82, 99,104,108,115,119,123,150,152,173, 181,248,262,268,280,286,294,295, 369,371 2,4,6,8,11,14,18,25,31,33,35,43,46, 57,63,69,70,84,88,89,105,109,110, 117,122,124,148,156,163,168,180, 274,276,288,292,315,322,330 7,22,32,36,37,64,71,95,98,112,116, 126,133,137,149,164,169,175,179, 246,247,250,252,261,271,284,289, 308,316,345,354,364 6,79,86,101- 103,114,134,165,170,178,237,259, 290,297,300,317,318,334,338,341, 359 | 99,186-189,192- 194,303,340,348,375 | 99,188,195,197,278,303, 348,351,375 | 46,50,99,186,198- 200,202,204,208- 210,241,253,255,278,28 0,305,346,355,375 187,197,201,203,206, 207,239,240,254,272, 331,333,352 | 1,19,57,84,89,104,105, 117,177,214,217,223, 225,263,283,285- 287,292,295,299,301, 315,375 36,133,137,211,212, 222,224,228,229,261, 311,314 | 249 | 230-233,245,313 |
| Long term care (nursing homes) | 40,95,302,377 | | 234,353 | 234,355,362 | 40 | | |
| Pharmacy | 65,75,99,105,111,167,173,310,350, 371,376 | 65,75,99,182- 184,188,190,191,243,258,265, 348 | 65,99,188,190,191,195, 196,265,348,351 | 99,202 | 105,167,287,296 | | 233 |

Evidence Table 18. Article references for outcomes studies evaluating clinicians across the medication management for phases, education, and reconciliation

| Provider | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|--------------------------------------|---|-------------------------|---------------------|--|-------------|-----------|-----------------------|
| Primary care physicians | 10,12,21,28,47,55,65,67,100,118,140,161,236,242,244,260,269,273,320,332,337,356,357,360,361,363 | 65,258 | 65 | | 12,356,358 | 321 | |
| Specialists | 47,147,169,236,246,248,252,268,273,338,360 | | | | 147 | | |
| Hospitalists | 21,58,64,117,125,148,152,167,235,246,262,277,319,330,338,345,349,368 | | | | 117,167 | | |
| Other Physicians | 7,10,21,117,235,271,349 | | | | 117,213 | | |
| Physicians undifferentiated | 58,90,116,138,143,149,248,256,259-261,264,266,274,276,318,322,334,338-341,344,345,347,364 | 340 | | 355 | 143,261,347 | | 245 |
| Nurses | 47,50,67,235,246,253,256,259,260,274,322,334,339,340,345,347,359,364,366,371 | 340 | | 50,198,206,239-241,253,254,272,331,333,343,346,352,355,362 | 347,358 | | 245 |
| Mid level practitioners (PA, NP, MW) | 47,67,236,337,347,357 | | | | 347,358 | | |
| Pharmacists | 167,276,277,322,334,339,340,345,350,361,370,371,377 | 188,243,258,265,340,348 | 188,196,265,348,351 | 272,355 | 167 | | 245 |
| Other health professionals | 90,152,260,270,271,276,334,339,361,371 | | 351,353 | 272,346,352,362 | 358 | | |
| Hospital administrators | 322,334,340,364 | 340 | | 346 | | | |

Evidence Table 19. Article references for patients studied by phase of medication management and education and reconciliation

| Patient | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|------------------------|--|---------------------|------------|---------------|--|-------------|-----------------------|
| Infants (0 to 2 years) | 25,62,70,79,176,294 | 303 | 303 | 203 | 219 | | |
| Children (2 to 12) | 9,19,41,62,70,176,177,250,294,305,308 | 303 | 303 | 305 | 19,177,219 | | |
| Adolescents (13 to 18) | 9,19,41,70,74,86,102,127,132,176,177,237,250,294,295 | | | | 19,127,132,177,211,218,219,295,335 | 127 | 245 |
| Adults (19 to 44) | 8,16,17,23,42,58,66,70,74,83,85,89,94,104,108,119,123,131-133,135,167,177,180,293,295 60,61,86,102,103,106,114,127,139,170,171,237,250,307 | | | | 23,60,61,66,89,94,104,127,132,133,167,177,211,215,218,219,226,263,275,295,296,311,335 | 127,249 | 245 |
| Middle age (45 to 64) | 8,16,17,26,35,39,42,55,58,66,70,74,83,85,89,91,94,104,107,108,119,123,131,132,135,136,166,167,180,292,293,295,310,315 23,60,61,86,92,96,102,106,112,114,126,127,133,139,145,160,170,171,237,300,306,307 | 189 | | 306 | 66,89,94,104,107,132,136,166,167,214,215,217,219,225,226,263,275,287,292,295,296,301,306,315 23,60,61,127,133,211,212,218,228,251,267,311,314,335 | 127,249,321 | 245 |
| Geriatric (65 plus) | 3,8,16,17,26,39,42,48,55,65,66,70,83,85,89,91,94,104,107,108,115,117,119,123,130,135,166,167,180,242,292,293,295,302,310,315 23,60,61,64,92,102,106,112,114,126,133,139,159-161,164,165,170,171,175,237,297,300,304 | 65,189 | 65,130 | 205 | 23,66,89,94,104,107,117,166,167,215,217,219,221,225,226,275,287,292,295,296,312,315 60,61,133,211,212,228,267,311,314,335 | 249,321 | 232,245,313 |
| Undifferentiated | 5,22,33,36,49,88,137,289,298 | | | | 36,137,222,229,279,285,298,367 | | |

Evidence Table 20. Main health IT studied by medication management phase and education and reconciliation

| MMIT System | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|-----------------------|--|------------------------------|---------------|--|---|-----------|-----------------------|
| CDSS/ reminders | 3,5,16,26,29,40,59,66,74,78,81,85,91,93,97, 99,100,113,121,125,128,130,131,135,144, 146,155,162,176,256,277,293,295,302,319, 360,377 1,10,11,13,17,20,38,39,55,58,69,72,82,88,89, 94,104,108,115,119,120,123,124,129,132, 136,140,152,154,157,166,167,173,236,248, 262,269,281,294,337 2,6,14,15,18,21,31,33,35- 37,42,43,45,48,57,63,65,83,84,87,105,107, 109,117,122,148,163,180,260,284,288,292, 306,310,315 9,12,22,23,28,32,41,47,49,64,79,95,96,98, 102,103,112,114,116,126,127,133,137,139, 141,145,147,159,161,164,169,171,175,261, 264,297,298,300,304,332,356,361 30,60- 62,75,76,86,92,101,106,134,143,160,165, 170,178,237,290,307,341,357,359 | 65,75,99,183,185,192,243,303 | 65,99,130,303 | 99,205,210,306,343 | 1,36,40,57,66,84,89, 94,104,105,107,113, 117,120,129,132,136, 166,167,214,215,219- 221,223,225,226,283, 285,287,292,295,296, 301,306,315,367 12,23,60,61,127,133, 137,141,143,147,211- 213,218,222,224,228, 229,261,279,298,309, 311,335,356,358 | 127 | 230 |
| Barcoding- dispensing | | | 351 | | | | |
| BCMA | 278 | | 234,278 | 200- 204,206,207,234,239- 241,254,255,272,278, 333,343,346,352,355 | | | |

Evidence Table 20. Main health IT studied by medication management phase and education and reconciliation (continued)

| MMIT System | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|---|---|--|--------------------|--|--|-----------|-----------------------|
| CPOE/POE system | 1,19,24,27,34,40,77,80,99,128,153,155,158,172-174,177,235,248,253,256,278,286,295,305,339,340,344,349,366,368-371,375,376 2,4,8,17,18,20,25,70,71,95,98,109,115,118,119,133,150,156,168,181,236,246,270,274,276,304,322,337,356,364 7,30,79,141,149,170,175,179,237,247,250,252,259,271,289,308,316-318,334,338,345,359,365 | 99,172,183,186,189,192-194,303,340,348,375 | 99,278,303,348,375 | 99,186,198,209,253,278,305,331,375 | 1,19,40,133,141,177,286,295,314,356,375 | | |
| e-Rx | 44,50,67,73,75,90,111,138,151,242,244,264,266,268,270,273,280,281,288,291,295,320,329,330,332,336,342,350,354,361,363 | 75,182,184,187,190,191,258,265,336,342 | 190,191,265 | 50,187,199,280 | 295,329 | 321 | 233 |
| e-Transmission of the prescription to/from doctor to pharmacy | 336,342 | 182,336,342 | | | | | |
| Pharmacy information system | 99,130 | 99,188,243 | 99,130,188,196 | 99 | | 249 | 233 |
| e-Medication administration system (e-MAR, e-TAR) | 50,354 | 186,187 | 197,234 | 50,186,187,197-199,203,204,207,234,241,346,352,362 | | | |
| Other | 36,46,50,110,127,242,250,277,316,318,347,349,375 | 258,375 | 195,353,375 | 46,50,198,199,208,210,375 | 36,127,212,217,251,263,267,275,283,299,312,335,347,375 | 127 | 230-232,245,313 |

Evidence Table 21. Article references for articles where the primary technology being studied was integrated with the various health IT systems

| Integrated System | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|--------------------------|---|---|------------------------------------|--|--|------------------|------------------------------|
| EHR/EMR system | 26,29,40,66,67,73,85,93,97,99,100, 125,130,135,144,151,162,176,177, 242,253,256,277,295,305,319,336, 340,342,349,366,368,376,377 1,2,13- 15,19,20,34,38,39,42,45,48,55,69, 80,87,88,94,105,107,108,115,119, 120,124,136,154,157,166- 168,288,292,294,310,337,347 9,23,35,37,41,47,49,71,95,96,114, 116,117,141,145,148,159,161,171, 175,260,271,289,300,304,308,316, 332 30,60,62,75,86,92,103,106,143,160, 165,259,290,318,334,338,341,357, 359,361 | 75,99,185,186,188,189,191, 193,336,340,342 | 99,130,188,191,234 | 99,186,201,205,207,210, 234,239,241,253,255, 305,331,355,362 | 1,19,40,66,94,105, 107,117,120,136,166, 167,177,215,217,219, 225,226,275,283,287, 292,295,301,312,347 23,60,141,143,212, 222,224,228,229,267, 311,314,358 | 321 | 230,231,245 |
| Formulary | 1,3,8,44,77,138,259,280,332,341, 363 | 187 | 197 | 187,197,280 | 1,211,225 | | |
| Pharmacy | 1,25,31,36,48,50,65,71,73,74,77, 104,105,111,120,128,129,131,146, 148,151,162,166,167,173,248,274, 276,280,305,310,330,356,364,376 61,112,116,170,178,250,259,270, 290,308,316,317,345,359,363 | 65,184,185,187,190,191,194 ,243,265 | 65,190,191,195,234, 265,351,353 | 50,187,198,201,203,204, 207- 209,234,241,254,272, 280,305 | 1,36,61,104,105,120, 129,166,167,212,214, 215,225,283,287,299, 314,356 | | 231,233 |

Evidence Table 21. Article references for articles where the primary technology being studied was integrated with the various health IT systems (continued)

| Integrated System | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|---------------------------------|--|----------------------------|-------------------|------------------------------------|--|------------------|------------------------------|
| CPOE/POE system | 3,11,14,24,33,38,42,43,57,58,69,78, 81,82,84,89,97,108,117,119-121,124,125,135,148,154,155,157, 162,167,262,292,293,302,315,319, 377 28,32,41,64,76,95,101-103,116,118,134,164,165,178,261, 297,300,307,316 | | 195 | 201,239-241,254,255,352 | 57,84,89,117,120, 167,223,261,287,292, 315,358 | | 230,313 |
| Hospital information system | 1,5,6,8,16,18,22,24,33,46,59,70,72, 77,83,84,89,99,109,110,115,122, 131,150,163,180,236,253,256,268, 269,276,284,286,294,339,344 7,61,76,79,98,116,126,133,134,137, 149,247,264,300,316,356,365 | 99,183,193,303 | 99,197,303 | 46,99,197,202,204,253,2 54,272,346 | 1,61,84,89,133,137, 213,224,225,283,286, 314,356 | 249 | 232 |
| Laboratory system | 1,2,5,19,24,34,39,57,74,84,93,104, 105,110,113,115,129,146,152,157, 162,163,176,177,276,294,302,377 35-37,61,96,112,116,117,148,175,250, 264,274,289,290,308,316,317,338, 356,359 | 186,188,189 | 188 | 186,198,254 | 1,19,36,57,61,84,104, 105,113,117,129,177, 212,217,221,225,228, 229,275,283,285,299, 309,314,356,358,367 | 321 | |
| Imaging systems | 1,2,18,25,35,70,77,96,112,118,148, 163,175,177,248,274,276,294,316, 317,338,359 | 186,188 | 188 | 186,198,254 | 1,177,217,220,228, 229,275 | 321 | |
| CDSS/reminders | 1,14,27,38,57,71,119,120,128,154, 172,174,246,250,266,270,289,291, 330,332,337,363,368,376 | 172 | | 239 | 1,57,120,220,221,312 | | |
| Billing/administration system | 6,8,19,24,74,112,118,276 | | | 272 | 19,215,217,275 | | |
| Insurance | 44,160,242,332,350 | | | | 218 | | |
| Personal health records systems | 350 | | | | 312,335 | | |
| Patient decision support system | | | | | 217 | | |
| Barcoding system | 170,354 | | 351 | | | | |

Evidence Table 21. Article references for articles where the primary technology being studied was integrated with the various health IT systems (continued)

| Integrated System | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|--------------------------|--|----------------------------|-------------------|-------------------------|--------------------------------|------------------|------------------------------|
| Not specified | 4,10,12,17,63,90,123,139,140,147,153,156,158,179,181,235,237,244,252,273,278,281,298,306,320,322,329,360,369-371,375 | 182,258,348,375 | 196,278,348,375 | 199,206,278,306,333,375 | 12,147,251,279,298,306,329,375 | | |
| Other | 14,21,47,73,91,115,127,132,160,169,178,259,340,345,354 | 186,192,340 | | 186,200,201,209,343 | 127,132,223,263,296,335 | 127 | 232 |

Evidence Table 22. Study designs used in studies measuring intermediate outcomes across the phases for medication management

| Design | Prescribing | Order Communication | Dispensing | Administering | Monitoring | Education | Reconciliation/ Other |
|---------------------------|---|----------------------------|-------------------|----------------------|-------------------|------------------|------------------------------|
| RCT | 75,237,252,262,264 | 75 | | | 263 | | |
| Cohort | 268 | | | | 267,275 | | |
| Observational | 235,236,242,244,246,250,253,259-251,266,269-271,274,276,277 | 243,265 | 234,265 | 234,240,253-255 | 261 | 249 | 245 |
| Qualitative Mixed Methods | 247,248,256,273 | 258 | | 239,241,272 | 251 | | |

Appendix C References

1. Abboud PA, Ancheta R, McKibben M, et al. Impact of workflow-integrated corollary orders on aminoglycoside monitoring in children. *Health Informatics Journal* 2006;12(3):187-98.
2. Achtmeyer CE, Payne TH, Anawalt BD. Computer order entry system decreased use of sliding scale insulin regimens. *Methods Inf Med* 2002;41(4):277-81.
3. Agostini J, Shang Y, Inouye S. Use of a coputer-based reminder to improve sedative-hypnotic prescribing in older hospitalized patients. *J Am Geriatr Soc* 2007;55:43-8.
4. Ali NA, Mekhjian HS, Kuehn PL, et al. Specificity of computerized physician order entry has a significant effect on the efficiency of workflow for critically ill patients. *Crit Care Med* 2005;33(1):110-4.
5. Bailey TC, Noirot LA, Blickensderfer A, et al. An intervention to improve secondary prevention of coronary heart disease. *Arch Intern Med* 2007;167(6):586-90.
6. Bates DW, Teich JM, Lee J, et al. The impact of computerized physician order entry on medication error prevention. *J Am Med Inform Assoc* 1999;6(4):313-21.
7. Bates D, Boyle D, Teich J. Impact of computerized physician order entry on physician time. *Proceedings of the AMIA Symposium* 1994;996
8. Bates DW, Leape LL, Cullen DJ, et al. Effect of computerized physician order entry and a team intervention on prevention of serious medication errors.[see comment]. *JAMA* 1998;280(15):1311-6.
9. Bell LM, Grundmeier R, Localio R, et al. Electronic health record-based decision support to improve asthma care: A cluster-randomized trial. *Pediatrics* 2010;125(4):e770-e777
10. Berner ES, Houston TK, Ray MN, et al. Improving ambulatory prescribing safety with a handheld decision support system: a randomized controlled trial. *J Am Med Inform Assoc* 2006;13(2):171-9.
11. Bernstein SL, Whitaker D, Winograd J, et al. An electronic chart prompt to decrease proprietary antibiotic prescription to self-pay patients. *Acad Emerg Med* 2005;12(3):225-31.
12. Bertoni A, Bonds D, Chen H, et al. Impact of a multifaceted intervention on cholesterol management in primary care practices. *Arch Intern Med* 2009;169(7):678-86.
13. Bloomfield HE, Nelson DB, van Ryn M, et al. A trial of education, prompts, and opinion leaders to improve prescription of lipid modifying therapy by primary care physicians for patients with ischemic heart disease. *Qual Safe Health Care* 2005;14(4):258-63.
14. Bogucki B, Jacobs BR, Hingle J, et al. Computerized reminders reduce the use of medications during shortages. *J Am Med Inform Assoc* 2004;11(4):278-80.
15. Bouaud J, Seroussi B, Antoine EC, et al. A before-after study using OncoDoc, a guideline-based decision support-system on breast cancer management: impact upon physician prescribing behaviour. *Stud Health Technol Inform* 2001;84(Pt.1):1-4.
16. Buising KL, Thursky KA, Black JF, et al. Improving antibiotic prescribing for adults with community acquired pneumonia: Does a computerised decision support system achieve more than academic detailing alone?--A time series analysis. *BMC Med Inform Decis Mak* 2008;8:35
17. Butler J, Speroff T, Arbogast PG, et al. Improved compliance with quality measures at hospital discharge with a computerized physician order entry system. *Am Heart J* 2006;151(3):643-53.
18. Chertow GM, Lee J, Kuperman GJ, et al. Guided medication dosing for inpatients with renal insufficiency. *JAMA* 2001;286(22):2839-44.
19. Chisolm DJ, McAlearney AS, Veneris S, et al. The role of computerized order sets in pediatric inpatient asthma treatment. *Pediatric Allergy & Immunology* 2006;17(3):199-206.

20. Choi SS, Jazayeri DG, Mitnick CD, et al. Implementation and initial evaluation of a Web-based nurse order entry system for multidrug-resistant tuberculosis patients in Peru. *Stud Health Technol Inform* 2004;107(Pt:1):1-6.
21. Christakis DA, Zimmerman FJ, Wright JA, et al. A randomized controlled trial of point-of-care evidence to improve the antibiotic prescribing practices for otitis media in children. *Pediatrics* 2001;107(2):e15
22. Clancy CM, Gelfman D, Poses RM. A strategy to improve the utilization of pneumococcal vaccine. *J Gen Intern Med* 1992;7(1):14-8.
23. Cobos A, Cobos A, Vilaseca J, et al. Cost effectiveness of a clinical decision support system based on the recommendations of the European Society of Cardiology and other societies for the management of hypercholesterolemia: Report of a cluster-randomized trial. *Disease Management and Health Outcomes* 2005;13(6):421-32.
24. Colpaert K, Claus B, Somers A, et al. Impact of computerized physician order entry on medication prescription errors in the intensive care unit: a controlled cross-sectional trial. *Critical Care (London, England)* 2006;10(1):R21
25. Cordero L, Kuehn L, Kumar RR, et al. Impact of computerized physician order entry on clinical practice in a newborn intensive care unit. *J Perinatol* 2004;24(2):88-93.
26. Cote GA, Rice JP, Bulsiewicz W, et al. Use of physician education and computer alert to improve targeted use of gastroprotection among NSAID users. *Am J Gastroenterol* 2008;103(5):1097-103.
27. Cunningham TR, Geller ES, Clarke SW. Impact of electronic prescribing in a hospital setting: a process-focused evaluation. *Int J Med Inf* 2008;77(8):546-54.
28. Davis RL, Wright J, Chalmers F, et al. A cluster randomized clinical trial to improve prescribing patterns in ambulatory pediatrics. *PLoS Clinical Trials* 2007;2(5):e25
29. de Jong JD, Groenewegen PP, Spreeuwenberg P, et al. Do decision support systems influence variation in prescription? *BMC Health Serv Res* 2009;9:9-20.
30. Devine EB, Wilson-Norton JL, Lawless NM, et al. The impact of an ambulatory CPOE system on medication errors. *AMIA* 2008;928
31. Dexter PR, Perkins S, Overhage JM, et al. A computerized reminder system to increase the use of preventive care for hospitalized patients. *N Engl J Med* 2001;345(13):965-70.
32. Dexter PR, Maharry K, Jones K, et al. Inpatient computer-based standing orders vs. physician reminders to increase influenza and pneumococcal vaccination rates: A randomized trial. *JAMA* 2004;292(19):2366-71.
33. Durieux P, Nizard R, Ravaud P, et al. A clinical decision support system for prevention of venous thromboembolism: effect on physician behavior. *JAMA* 2000;283(21):2816-21.
34. Eslami S, Abu-Hanna A, de Keizer NF, et al. Errors associated with applying decision support by suggesting default doses for aminoglycosides. *Drug Saf* 2006;29(9):803-9.
35. Evans RS, Pestotnik SL, Classen DC, et al. A computer-assisted management program for antibiotics and other anti-infective agents.[see comment]. *N Engl J Med* 1998;338(4):232-8.
36. Evans RS, Pestotnik SL, Burke JP, et al. Reducing the duration of prophylactic antibiotic use through computer monitoring of surgical patients. *DICP* 1990;24(4):351-4.
37. Evans RS, Classen DC, Pestotnik SL, et al. Improving empiric antibiotic selection using computer decision support. *Arch Intern Med* 1994;154(8):878-84.
38. Feldstein AC, Smith DH, Perrin N, et al. Reducing warfarin medication interactions: an interrupted time series evaluation. *Arch Intern Med* 2006;166(9):1009-15.
39. Feldstein A, Elmer PJ, Smith DH, et al. Electronic medical record reminder improves osteoporosis management after a fracture: a randomized, controlled trial. *J Am Geriatr Soc* 2006;54(3):450-7.

40. Field TS, Rochon P, Lee M, et al. Computerized clinical decision support during medication ordering for long-term care residents with renal insufficiency. *J Am Med Inform Assoc* 2009;16(4):480-5.
41. Fiks AG, Hunter K, Localio R, et al. Impact of electronic health record-based alerts on influenza vaccination for children with asthma. *Pediatrics* 2009;124(1):159-69.
42. Filippi A, Sabatini A, Badioli L, et al. Effects of an automated electronic reminder in changing the antiplatelet drug-prescribing behavior among Italian general practitioners in diabetic patients: an intervention trial. *Diabetes Care* 2003;26(5):1497-500.
43. Fischer MA, Solomon DH, Teich JM, et al. Conversion from intravenous to oral medications: assessment of a computerized intervention for hospitalized patients. *Arch Intern Med* 2003;163(21):2585-9.
44. Fischer MA, Vogeli C, Stedman M, et al. Effect of electronic prescribing with formulary decision support on medication use and cost. *Arch Intern Med* 2008;168(22):2433-9.
45. Flottorp S, Oxman AD, Havelsrud K, et al. Cluster randomised controlled trial of tailored interventions to improve the management of urinary tract infections in women and sore throat. *BMJ* 2002;325(7360):367
46. Fontan JE, Maneglier V, Nguyen VX, et al. Medication errors in hospitals: computerized unit dose drug dispensing system versus ward stock distribution system. *Pharm World Sci* 2003;25(3):112-7.
47. Fortuna R, Zhang F, Ross-Degnan D, et al. Reducing the prescribing of heavily marketed medications: A randomized controlled trial. *J Gen Intern Med* 2009;24(8):897-903.
48. Frances CD, Alperin P, Adler JS, et al. Does a fixed physician reminder system improve the care of patients with coronary artery disease? A randomized controlled trial. *West J Med* 2001;175(3):165-6.
49. Frank O, Litt J, Beilby J. Opportunistic electronic reminders: Improving performance of preventive care in general practice. *Aust Fam Physician* 2004;33(1/2):87-90.
50. Franklin BD, O'Grady K, Donyai P, et al. The impact of a closed-loop electronic prescribing and administration system on prescribing errors, administration errors and staff time: a before-and-after study. *Qual Safe Health Care* 2007;16(4):279-84.
51. Donyai P, O'Grady K, Jacklin A, et al. The effects of electronic prescribing on the quality of prescribing. *Br J Clin Pharmacol* 2008;65(2):230-7.
52. Barber N, Cornford T, Klecun E. Qualitative evaluation of an electronic prescribing and administration system. *Qual Safe Health Care* 2007;16(4):271-8.
53. Franklin BD, Jacklin A, Barber N. The impact of an electronic prescribing and administration system on the safety and quality of medication administration. *Int J Pharm Pract* 2008;16(6):375-9.
54. Franklin BD, O'Grady K, Donyai P, et al. The impact of a closed-loop electronic prescribing and automated dispensing system on the ward pharmacist's time and activities. *Int J Pharm Pract* 2007;15(2):133-9.
55. Fretheim A, Aaserud M, Oxman AD. Rational prescribing in primary care (RaPP): economic evaluation of an intervention to improve professional practice. *PLoS Medicine* 2006;3(6):e216
56. Fretheim A, Oxman AD, Havelsrud K, et al. Rational prescribing in primary care (RaPP): a cluster randomized trial of a tailored intervention. *PLoS Medicine / Public Library of Science* 2006;3(6):e134
57. Galanter WL, Polikaitis A, Didomenico RJ. A trial of automated safety alerts for inpatient digoxin use with computerized physician order entry. *J Am Med Inform Assoc* 2004;11(4):270-7.
58. Galanter WL, Didomenico RJ, Polikaitis A. A trial of automated decision support alerts for contraindicated medications using computerized physician order entry. *J Am Med Inform Assoc* 2005;12(3):269-74.
59. Gerard MN, Trick WE, Das K, et al. Use of clinical decision support to increase influenza vaccination: multi-year evolution of the system. *J Am Med Inform Assoc* 2008;15(6):776-9.

60. Gill JM, Chen YX, Glutting JJ, et al. Impact of decision support in electronic medical records on lipid management in primary care. *Popul Health Manag* 2009;12(5):221-6.
61. Gilutz H, Novack L, Shvartzman P, et al. Computerized community cholesterol control (4C): meeting the challenge of secondary prevention. *The Israel Medical Association journal : IMAJ* 2009;11(1):23-9.
62. Ginzburg R, Barr WB, Harris M, et al. Effect of a weight-based prescribing method within an electronic health record on prescribing errors. *Am J Health Syst Pharm* 2009;66(22):2037-41.
63. Goethe JW, Schwartz HI, Szarek BL. Physician compliance with practice guidelines. *Conn Med* 1997;61(9):553-8.
64. Griffey RT. Guided medication dosing for elderly emergency department patients using a real-time, computerized decision support tool. *Ann Emerg Med* 2009;54(3):265
65. Halkin H, Katzir I, Kurman I, et al. Preventing drug interactions by online prescription screening in community pharmacies and medical practices. *Clinical Pharmacology & Therapeutics* 2001;69(4):260-5.
66. Hicks LS, Sequist TD, Ayanian JZ, et al. Impact of computerized decision support on blood pressure management and control: a randomized controlled trial. *J Gen Intern Med* 2008;23(4):429-41.
67. Hollingworth W, Devine EB, Hansen RN, et al. The impact of e-prescribing on prescriber and staff time in ambulatory care clinics: a time motion study. *J Am Med Inform Assoc* 2007;14(6):722-30.
68. Devine EB, Hollingworth W, Hansen RN, et al. Electronic prescribing at the point of care: A time-motion study in the primary care setting. *Health Serv Res* 2010;45(1):152-71.
69. Hulgán T, Rosenbloom ST, Hargrove F, et al. Oral quinolones in hospitalized patients: an evaluation of a computerized decision support intervention. *J Intern Med* 2004;256(4):349-57.
70. Hwang JI, Park HA, Bakken S. Impact of a physician's order entry (POE) system on physicians' ordering patterns and patient length of stay. *Int J Med Inf* 2002;65(3):213-23.
71. Igboechi CA, Ng C, Yang C, et al. Impact of computerized prescriber order entry on medication errors at an acute tertiary care hospital. *Hosp Pharm* 2003;38(3):227-31.
72. St Jacques P, Sanders N, Patel N, et al. Improving timely surgical antibiotic prophylaxis redosing administration using computerized record prompts. *Surgical Infections* 2005;6(2):215-21.
73. Jani YH, Ghaleb MA, Marks SD, et al. Electronic prescribing reduced prescribing errors in a pediatric renal outpatient clinic. *J Pediatr* 2008;152(2):214-8.
74. Javitt JC, Rebitzer JB, Reisman L. Information technology and medical missteps: evidence from a randomized trial. *J Health Econ* 2008;27(3):585-602.
75. Johnson KB, Ho Y-X, Cala CM, et al. Showing Your Work: Impact of annotating electronic prescriptions with decision support results. *J Biomed Inform* 2010;43(2):321-5.
76. Kadmon G, Bron-Harlev E, Nahum E, et al. Computerized order entry with limited decision support to prevent prescription errors in a PICU. *Pediatrics* 2009;124(3):935-40.
77. Kaplan JM, Ancheta R, Jacobs BR, et al. Inpatient verbal orders and the impact of computerized provider order entry. *J Pediatr* 2006;149(4):461-7.
78. Karson AS, Campbell EJ, Panagou CM, et al. Using computerized provider order entry application to improve compliance with co-signature of verbal orders. *AMIA Proceedings* 2007;1004
79. Kazemi A, Fors UG, Tofighi S, et al. Physician order entry or nurse order entry? Comparison of two implementation strategies for a computerized order entry system aimed at reducing dosing medication errors. *J Med Internet Res* 2010;12(1):e5

80. Kim GR, Chen AR, Arceci RJ, et al. Error reduction in pediatric chemotherapy: computerized order entry and failure modes and effects analysis. *Archives of Pediatrics & Adolescent Medicine* 2006;160(5):495-8.
81. Kim JY, Sohn JW, Park DW, et al. Control of extended-spectrum β -lactamase-producing *Klebsiella pneumoniae* using a computer-assisted management program to restrict third-generation cephalosporin use. *J Antimicrob Chemother* 2008;62(2):416-21.
82. Kirk RC, Li-Meng GD, Packia J, et al. Computer calculated dose in paediatric prescribing. *Drug Saf* 2005;28(9):817-24.
83. Kitahata MM, Dillingham PW, Chaiyakunapruk N, et al. Electronic human immunodeficiency virus (HIV) clinical reminder system improves adherence to practice guidelines among the University of Washington HIV Study Cohort. *Clin Infect Dis* 2003;36(6):803-11.
84. Koide D, Ohe K, Ross-Degnan D, et al. Computerized reminders to monitor liver function to improve the use of etretinate. *Int J Med Inf* 2000;57(1):11-9.
85. Kooij FO, Klok T, Hollmann MW, et al. Decision support increases guideline adherence for prescribing postoperative nausea and vomiting prophylaxis. *Anesth Analg* 2008;106(3):893-8.
86. Kooij FO, Klok T, Hollmann MW, et al. Automated reminders increase adherence to guidelines for administration of prophylaxis for postoperative nausea and vomiting. *Eur J Anaesthesiol* 2010;27(2):187-91.
87. Kralj B, Iverson D, Hotz K, et al. The impact of computerized clinical reminders on physician prescribing behavior: evidence from community oncology practice. *Am J Med Qual* 2003;18(5):197-203.
88. Krall MA, Traunweiser K, Towery W. Effectiveness of an electronic medical record clinical quality alert prepared by off-line data analysis. *Stud Health Technol Inform* 2004;107(Pt:1):1-9.
89. Kucher N, Koo S, Quiroz R, et al. Electronic alerts to prevent venous thromboembolism among hospitalized patients. *N Engl J Med* 2005;352(10):969-77.
90. Lapane KL, Waring ME, Schneider KL, et al. A mixed method study of the merits of e-prescribing drug alerts in primary care. *J Gen Intern Med* 2008;23(4):442-6.
91. Lecumberri R, Marques M, Diaz-Navarraz MT, et al. Maintained effectiveness of an electronic alert system to prevent venous thromboembolism among hospitalized patients. *Thrombosis & Haemostasis* 2008;100(4):699-704.
92. Ledwich LJ, Harrington TM, Ayoub WT, et al. Improved influenza and pneumococcal vaccination in rheumatology patients taking immunosuppressants using an electronic health record best practice alert. *Arthritis Rheum* 2009;61(11):1505-10.
93. Lesprit P, Duong T, Girou E, et al. Impact of a computer-generated alert system prompting review of antibiotic use in hospitals. *J Antimicrob Chemother* 2009;63(5):1058-63.
94. Lester WT, Grant RW, Barnett GO, et al. Randomized controlled trial of an informatics-based intervention to increase statin prescription for secondary prevention of coronary disease. *J Gen Intern Med* 2006;21(1):22-9.
95. Lin C, Payne T, Nichol W, et al. Evaluating Clinical Decision Support Systems: Monitoring CPOE Order Check Override Rates in the Department of Veterans Affairs' Computerized Patient Record System. *J Am Med Inform Assoc* 2008;15(5):620-6.
96. Linder JA, Schnipper JL, Tsurikova R, et al. Documentation-based clinical decision support to improve antibiotic prescribing for acute respiratory infections in primary care: A cluster randomised controlled trial. *Inform Prim Care* 2009;17(4):231-40.
97. Liu SA, Chiu YT, Lin WD, et al. Using information technology to reduce the inappropriate use of surgical prophylactic antibiotic. *Eur Arch Otorhinolaryngol* 2008;265(9):1109-12.
98. Madaras-Kelly KJ, Remington RE, Lewis PG, et al. Evaluation of an intervention designed to decrease the rate of nosocomial methicillin-resistant *Staphylococcus aureus* infection by encouraging decreased fluoroquinolone use. *INFECT CONTROL HOSP EPIDEMIOL* 2006;27(2):155-69.

99. Mahoney CD, Berard-Collins CM, Coleman R, et al. Effects of an integrated clinical information system on medication safety in a multi-hospital setting. *Am J Health Syst Pharm* 2007;64(18):1969-77.
100. Martens JD, van der WT, Severens JL, et al. The effect of computer reminders on GPs' prescribing behaviour: a cluster-randomised trial. *Int J Med Inf* 2007;76(Suppl 3):S403-S416
101. Mattison ML, Afonso KA, Ngo LH, et al. Preventing Potentially Inappropriate Medication Use in Hospitalized Older Patients With a Computerized Provider Order Entry Warning System. *Arch Intern Med* 2010;170(15):1331-6.
102. Maynard GA, Morris TA, Jenkins IH, et al. Optimizing prevention of hospital-acquired venous thromboembolism (VTE): prospective validation of a VTE risk assessment model. *Journal of Hospital Medicine (Online)* 2010;5(1):10-8.
103. McCluggage L, Lee K, Potter T, et al. Implementation and evaluation of vancomycin nomogram guidelines in a computerized prescriber-order-entry system. *Am J Health Syst Pharm* 2010;67(1):70-5.
104. McGregor JC, Weekes E, Forrest GN, et al. Impact of a computerized clinical decision support system on reducing inappropriate antimicrobial use: a randomized controlled trial. *J Am Med Inform Assoc* 2006;13(4):378-84.
105. McMullin ST, Reichley RM, Watson LA, et al. Impact of a Web-based clinical information system on cisapride drug interactions and patient safety.[see comment]. *Arch Intern Med* 1999;159(17):2077-82.
106. Miskulin DC, Weiner DE, Tighiouart H, et al. Computerized decision support for EPO dosing in hemodialysis patients. *Am J Kidney Dis* 2009;54(6):1081-8.
107. Montgomery AA, Fahey T, Peters TJ, et al. Evaluation of computer based clinical decision support system and risk chart for management of hypertension in primary care: randomised controlled trial.[see comment]. *BMJ* 2000;320(7236):686-90.
108. Morrison RS, Meier DE, Fischberg D, et al. Improving the management of pain in hospitalized adults. *Arch Intern Med* 2006;166(9):1033-9.
109. Mullett CJ, Evans RS, Christenson JC, et al. Development and impact of a computerized pediatric antiinfective decision support program. *Pediatrics* 2001;108(4):e75
110. Nash IS, Rojas M, Hebert P, et al. Reducing excessive medication administration in hospitalized adults with renal dysfunction. *Am J Med Qual* 2005;20(2):64-9.
111. Newby DA, Fryer JL, Henry DA. Effect of computerised prescribing on use of antibiotics. *Med J Aust* 2003;178(5):210-3.
112. Niemi K, Geary S, Quinn B, et al. Implementation and evaluation of electronic clinical decision support for compliance with pneumonia and heart failure quality indicators. *Am J Health Syst Pharm* 2009;66(4):389-97.
113. Niiranen ST, Yli-Hietanen JM. Analysis of computer-supported oral anticoagulant treatment follow-up workflow. *Conference Proceedings: 2008*;4330-2.
114. Novis SJ, Havelka GE, Ostrowski D, et al. Prevention of thromboembolic events in surgical patients through the creation and implementation of a computerized risk assessment program. *J Vasc Surg* 2010;51(3):648-54.
115. Oliven A, Michalake I, Zalman D, et al. Prevention of prescription errors by computerized, on-line surveillance of drug order entry. *Int J Med Inf* 2005;74(5):377-86.
116. Overhage J, Tierney W, McDonald C. Computer reminders to implement preventive care guidelines for hospitalized patients. *Arch Intern Med* 1996;156(14):1551-6.
117. Overhage JM, Tierney WM, Zhou XH, et al. A randomized trial of "corollary orders" to prevent errors of omission. *J Am Med Inform Assoc* 1997;4(5):364-75.
118. Overhage JM, Perkins S, Tierney WM, et al. Controlled trial of direct physician order entry: Effects on physicians' time utilization in ambulatory primary care internal medicine practices. *J Am Med Inform Assoc* 2001;8(4):361-9.

119. Ozdas A, Speroff T, Waitman LR, et al. Integrating “best of care” protocols into clinicians’ workflow via care provider order entry: impact on quality-of-care indicators for acute myocardial infarction. *J Am Med Inform Assoc* 2006;13(2):188-96.
120. Palen TE, Raebel M, Lyons E, et al. Evaluation of laboratory monitoring alerts within a computerized physician order entry system for medication orders. *Am J Manag Care* 2006;12(7):389-95.
121. Paterno MD, Maviglia SM, Gorman PN, et al. Tiering drug-drug interaction alerts by severity increases compliance rates. *J Am Med Inform Assoc* 2009;16(1):40-6.
122. Patterson R. A computerized reminder for prophylaxis of deep vein thrombosis in surgical patients. *Proceedings / AMIA 1998;Annual Symposium.*:573-6.
123. Paul M, Andreassen S, Tacconelli E, et al. Improving empirical antibiotic treatment using TREAT, a computerized decision support system: cluster randomized trial. *J Antimicrob Chemother* 2006;58(6):1238-45.
124. Peterson JF, Kuperman GJ, Shek C, et al. Guided prescription of psychotropic medications for geriatric inpatients. *Arch Intern Med* 2005;165(7):802-7.
125. Peterson JF, Rosenbaum BP, Waitman LR, et al. Physicians’ response to guided geriatric dosing: initial results from a randomized trial. *Stud Health Technol Inform* 2007;129(Pt:2):2-40.
126. Weir CJ, Lees KR, MacWalter RS, et al. Cluster-randomized, controlled trial of computer-based decision support for selecting long-term anti-thrombotic therapy after acute ischaemic stroke. *QJM: An International Journal of Medicine* 2003;96(2):143-53.
127. Quinn C, Clough S, Minor J, et al. WellDoc mobile diabetes management randomized controlled trial: Change in clinical and behavioral outcomes and patient and physician satisfaction. *Diabetes Technol Ther* 2008;10(3):160-8.
128. Quinzler R, Schmitt SP, Pritsch M, et al. Substantial reduction of inappropriate tablet splitting with computerised decision support: a prospective intervention study assessing potential benefit and harm. *BMC Med Inform Decis Mak* 2009;9:30-5.
129. Raebel MA, Lyons EE, Chester EA, et al. Improving laboratory monitoring at initiation of drug therapy in ambulatory care: a randomized trial. *Arch Intern Med* 2005;165(20):2395-401.
130. Raebel MA, Charles J, Dugan J, et al. Randomized trial to improve prescribing safety in ambulatory elderly patients. *J Am Geriatr Soc* 2007;55(7):977-85.
131. Raebel MA, Carroll NM, Kelleher JA, et al. Randomized trial to improve prescribing safety during pregnancy. *J Am Med Inform Assoc* 2007;14(4):440-50.
132. Rasmussen LM, Phanareth K, Nolte H, et al. Internet-based monitoring of asthma: a long-term, randomized clinical study of 300 asthmatic subjects. *Journal of Allergy & Clinical Immunology* 2005;115(6):1137-42.
133. Riggio JM, Cooper M, Leiby B, et al. Effectiveness of a clinical decision support system to identify heparin induced thrombocytopenia. *J Thromb Thrombolysis* 2009;28(2):124-31.
134. Riggio JM, Sorokin R, Moxey ED, et al. Effectiveness of a clinical-decision-support system in improving compliance with cardiac-care quality measures and supporting resident training. *Acad Med* 2009;84(12):1719-26.
135. Rohrig R, Niczko EJ, Beutefuhr H, et al. Examination of computer assisted prescribing of an initial calculated antibiotic treatment. *Stud Health Technol Inform* 2008;136:63-8.
136. Rollman BL, Hanusa BH, Lowe HJ, et al. A randomized trial using computerized decision support to improve treatment of major depression in primary care. *J Gen Intern Med* 2002;17(7):493-503.
137. Rood E, Bosman R, van der Spoel J, et al. Use of a computerized guideline for glucose regulation in the intensive care unit improved both guideline adherence and glucose regulation. *J Am Med Inform Assoc* 2005;12(2):172-80.

138. Ross SM, Papshev D, Murphy EL, et al. Effects of electronic prescribing on formulary compliance and generic drug utilization in the ambulatory care setting: a retrospective analysis of administrative claims data. *J Manag Care Pharm* 2005;11(5):410-5.
139. Rosser WW, Hutchison B, McDowell I, et al. Use of reminders to increase compliance with tetanus booster vaccination. *Can Med Assoc J* 1992;146(6):911-7.
140. Rubin MA, Bateman K, Donnelly S, et al. Use of a personal digital assistant for managing antibiotic prescribing for outpatient respiratory tract infections in rural communities. *J Am Med Inform Assoc* 2006;13(6):627-34.
141. Safran C, Rind D, Davis R, et al. Guidelines for management of HIV infection with computer-based patient's record. *The Lancet* 1995;346:341-6.
142. Safran C, Rind DM, Davis RM, et al. An electronic medical record that helps care for patients with HIV infection. Proceedings - the Annual Symposium on Computer Applications in Medical Care 1993;224-8.
143. Schnipper JL, McColgan KE, Linder JA, et al. Improving management of chronic diseases with documentation-based clinical decision support: results of a pilot study. *AMIA 2008;Annual:Symposium*
144. Scotton DW, Wierman H, Coughlan A, et al. Assessing the appropriate use of metformin in an inpatient setting and the effectiveness of two pharmacy-based measures to improve guideline adherence. *Qual Manag Health Care* 2009;18(1):71-6.
145. Segarra-Newnham M. Tracking vaccination rates among HIV-positive patients with a computerized reminder system. *Hosp Pharm* 2003;38(8):758-62.
146. Sellier E, Colombet I, Sabatier B, et al. Effect of alerts for drug dosage adjustment in inpatients with renal insufficiency. *J Am Med Inform Assoc* 2009;16(2):203-10.
147. Shiffman RN, Brandt C, Liaw Y, et al. A guideline implementation system using handheld computers for office management of asthma: Effects on adherence and patient outcomes. *Pediatrics* 2000;105(4):767-73.
148. Shojania KG, Yokoe D, Platt R, et al. Reducing vancomycin use utilizing a computer guideline: results of a randomized controlled trial.[see comment]. *J Am Med Inform Assoc* 1998;5(6):554-62.
149. Bates D, Shu K, Narasimhan D, et al. Comparing time spent writing orders on paper and physician computer order entry. *Proceedings of the AMIA Symposium* 2000;965
150. Shulman R, Singer M, Goldstone J, et al. Medication errors: a prospective cohort study of hand-written and computerised physician order entry in the intensive care unit. *Critical Care (London, England)* 2005;9(5):R516-R521
151. Delgado SE, Soler VM, Perez Menendez-Conde C, et al. Prescription errors after the implementation of an electronic prescribing system. *Farm Hosp* 2007;31(4):223-30.
152. Sintchenko V, Iredell JR, Gilbert GL, et al. Handheld computer-based decision support reduces patient length of stay and antibiotic prescribing in critical care. *J Am Med Inform Assoc* 2005;12(4):398-402.
153. Small MD, Barrett A, Price GM. The impact of computerized prescribing on error rate in a department of Oncology/Hematology. *Journal of Oncology Pharmacy Practice* 2008;14(4):181-7.
154. Smith DH, Perrin N, Feldstein A, et al. The impact of prescribing safety alerts for elderly persons in an electronic medical record: an interrupted time series evaluation. *Arch Intern Med* 2006;166(10):1098-104.
155. Sobieraj DM. Development and implementation of a program to assess medical patients' need for venous thromboembolism prophylaxis. *Am J Health Syst Pharm* 2008;65(18):1755-60.
156. Spencer DC, Leininger A, Daniels R, et al. Effect of a computerized prescriber-order-entry system on reported medication errors. *Am J Health Syst Pharm* 2005;62(4):416-9.
157. Steele AW, Eisert S, Witter J, et al. The effect of automated alerts on provider ordering behavior in an outpatient setting. *PLoS Medicine* 2005;2(9):e255

158. Stone WM, Smith BE, Shaft JD, et al. Impact of a computerized physician order-entry system. *J Am Coll Surg* 2009;208(5):960-7.
159. Tamblyn R, Huang A, Perreault R, et al. The medical office of the 21st century (MOXXI): Effectiveness of computerized decision-making support in reducing inappropriate prescribing in primary care. *Can Med Assoc J* 2003;169(6):549-56.
160. Tamblyn R, Reidel K, Huang A, et al. Increasing the detection and response to adherence problems with cardiovascular medication in primary care through computerized drug management systems: a randomized controlled trial. *Med Decis Making* 2010;30(2):176-88.
161. Tang PC, LaRosa M, Newcomb C, et al. Measuring the effects of reminders for outpatient influenza immunizations at the point of clinical opportunity. *J Am Med Inform Assoc* 1999;6(2):115-21.
162. Tang MB, Tan ES, Tian EA, et al. Electronic e-isotretinoin prescription chart: improving physicians' adherence to isotretinoin prescription guidelines. *Australas J Dermatol* 2009;50(2):107-12.
163. Teich JM, Merchia PR, Schmiz JL, et al. Effects of computerized physician order entry on prescribing practices. *Arch Intern Med* 2000;160(18):2741-7.
164. Terrell KM, Perkins AJ, Dexter PR, et al. Computerized decision support to reduce potentially inappropriate prescribing to older emergency department patients: A randomized, controlled trial. *J Am Geriatr Soc* 2009;57(8):1388-94.
165. Terrell KM, Perkins AJ, Dexter PR, et al. Computerized decision support to reduce potentially inappropriate prescribing to older emergency department patients: a randomized, controlled trial. *J Am Geriatr Soc* 2009;57(8):1388-94.
166. Tierney WM, Overhage JM, Murray MD, et al. Effects of computerized guidelines for managing heart disease in primary care. *J Gen Intern Med* 2003;18(12):967-76.
167. Tierney WM, Overhage JM, Murray MD, et al. Can computer-generated evidence-based care suggestions enhance evidence-based management of asthma and chronic obstructive pulmonary disease? A randomized, controlled trial. *Health Serv Res* 2005;40(2):477-97.
168. Upperman JS, Staley P, Friend K, et al. The impact of hospitalwide computerized physician order entry on medical errors in a pediatric hospital. *J Pediatr Surg* 2005;40(1):57-9.
169. Uttaro T, Finnerty M, White T, et al. Reduction of concurrent antipsychotic prescribing practices through the use of PSYCKES. *Adm Policy Ment Health* 2007;34(1):57-61.
170. van Doormaal JE, van den Bemt PM, Zaal RJ, et al. The influence that electronic prescribing has on medication errors and preventable adverse drug events: an interrupted time-series study. *J Am Med Inform Assoc* 2009;16(6):816-25.
171. Van Wyk JT, van Wijk M, Sturkenboom M, et al. Electronic alerts versus on-demand decision support to improve dyslipidemia treatment: A cluster randomized controlled trial. *Circulation* 2008;117(3):371-8.
172. Varkey P, Aponte P, Swanton C, et al. The effect of computerized physician-order entry on outpatient prescription errors. *Manag Care Interface* 2007;20(3):53-7.
173. Voeffray M, Pannatier A, Stupp R, et al. Effect of computerisation on the quality and safety of chemotherapy prescription. *Qual Safe Health Care* 2006;15(6):418-21.
174. Walsh KE, Landrigan CP, Adams WG, et al. Effect of computer order entry on prevention of serious medication errors in hospitalized children. *Pediatrics* 2008;121(3):e421-e427.
175. Were MC, Abernathy G, Hui SL, et al. Using computerized provider order entry and clinical decision support to improve referring physicians' implementation of consultants' medical recommendations. *J Am Med Inform Assoc* 2009;16(2):196-202.

176. Wilkes JJ, Zaoutis TE, Keren R, et al. Treatment with oseltamivir in children hospitalized with community-acquired, laboratory-confirmed influenza: review of five seasons and evaluation of an electronic reminder. *Journal of Hospital Medicine (Online)* 2009;4(3):171-8.
177. Wrona S, Chisolm DJ, Powers M, et al. Improving processes of care in patient-controlled analgesia: the impact of computerized order sets and acute pain service patient management. *Paediatr Anaesth* 2007;17(11):1083-9.
178. Xamplas RC, Itokazu GS, Glowacki RC, et al. Implementation of an extended-infusion piperacillin-tazobactam program at an urban teaching hospital. *Am J Health Syst Pharm* 2010;67(8):622-8.
179. Yu FB, Menachemi N, Berner ES, et al. Full implementation of computerized physician order entry and medication-related quality outcomes: A study of 3364 hospitals. *Am J Med Qual* 2009;24(4):278-86.
180. Zanetti G, Flanagan HL, Jr., Cohn LH, et al. Improvement of intraoperative antibiotic prophylaxis in prolonged cardiac surgery by automated alerts in the operating room. *INFECT CONTROL HOSP EPIDEMIOL* 2003;24(1):13-6.
181. Zhan C, Hicks RW, Blanchette CM, et al. Potential benefits and problems with computerized prescriber order entry: analysis of a voluntary medication error-reporting database. *Am J Health Syst Pharm* 2006;63(4):353-8.
182. Astrand B, Montelius E, Petersson G, et al. Assessment of ePrescription quality: an observational study at three mail-order pharmacies. *BMC Med Inform Decis Mak* 2009;9:8
183. Beer J. Physician order entry: A mixed blessing to pharmacy? *Journal of Oncology Pharmacy Practice* 2002;8(4):
184. Ekedahl A, Mansson N. Unclaimed prescriptions after automated prescription transmittals to pharmacies. *Pharm World Sci* 2004;26(1):26-31.
185. Humphries TL, Carroll N, Chester EA, et al. Evaluation of an electronic critical drug interaction program coupled with active pharmacist intervention. *Ann Pharmacother* 2007;41(12):1979-85.
186. Mekhjian HS, Kumar RR, Kuehn L, et al. Immediate benefits realized following implementation of physician order entry at an academic medical center. *J Am Med Inform Assoc* 2002;9(5):529-39.
187. Mitchell D, Usher J, Gray S, et al. Evaluation and audit of a pilot of electronic prescribing and drug administration. *Journal on Information Technology in Healthcare* 2004;2(1):19-29.
188. Murray MD, Loos B, Tu W, et al. Work patterns of ambulatory care pharmacists with access to electronic guideline-based treatment suggestions. *Am J Health Syst Pharm* 1999;56(3):225-32.
189. Nam HS, Han SW, Ahn SH, et al. Improved time intervals by implementation of computerized physician order entry-based stroke team approach. *Cerebrovasc Dis* 2007;23(4):289-93.
190. Nilsson JLG, Backstrom S, Sundstrom J. Electronically transferred prescriptions: Picked up faster than paper prescriptions. *Int J Pharm Pract* 2007;15(2):157-8.
191. Pearce DD, Opperman JM. Electronic medical record reduces HIV medication refill response time and emergency refills in a Latino community clinic. *Int J STD AIDS* 2010;21(3):184-6.
192. Senholzi C, Gottlieb J. Pharmacist interventions after implementation of computerized prescriber order entry. *Am J Health Syst Pharm* 2003;60(18):1880-2.
193. Wess ML, Embi PJ, Besier JL, et al. Effect of a Computerized Provider Order Entry (CPOE) System on medication orders at a community hospital and university hospital. *AMIA Proceedings* 2007;Oct 11:796-800.
194. Wietholter J, Sitterson S, Allison S. Effects of computerized prescriber order entry on pharmacy order-processing time. *Am J Health Syst Pharm* 2009;66(15):1394-8.

195. Alvarez Diaz AM, Delgado SE, Perez Menendez-Conde C, et al. New technologies applied to the medication-dispensing process, error analysis and contributing factors. [Spanish]. *Farm Hosp* 2010;34(2):59-67.
196. Reeve JF, Tenni PC, Peterson GM. An electronic prompt in dispensing software to promote clinical interventions by community pharmacists: a randomized controlled trial. *Br J Clin Pharmacol* 2008;65(3):377-85.
197. Wilson AL, Hill JJ, Wilson RG, et al. Computerized medication administration records decrease medication occurrences. *Pharm Pract Manag Q* 1997;17(1):17-29.
198. Banet GA, Jeffe DB, Williams JA, et al. Effects of implementing computerized practitioner order entry and nursing documentation on nursing workflow in an emergency department. *J Healthc Inf Manag* 2006;20(2):45-54.
199. Climent C, Font-Noguera I, Poveda Andres JL, et al. Medication errors in a tertiary hospital with three different drug delivery systems. *Farm Hosp* 2008;32(1):18-24.
200. DeYoung JL, Vanderkooi ME, Barletta JF. Effect of bar-code-assisted medication administration on medication error rates in an adult medical intensive care unit. *Am J Health Syst Pharm* 2009;66(12):1110-5.
201. Helmons PJ, Wargel LN, Daniels CE. Effect of bar-code-assisted medication administration on medication administration errors and accuracy in multiple patient care areas. *Am J Health Syst Pharm* 2009;66(13):1202-10.
202. Low DK, Belcher JV. Reporting medication errors through computerized medication administration. *Comput Inform Nurs* 2002;20(5):178-83.
203. Morriss F, Abramowitz P, Nelson S, et al. Effectiveness of a Barcode Medication Administration System in Reducing Preventable Adverse Drug Events in a Neonatal Intensive Care Unit: A Prospective Cohort Study. *J Pediatr* 2009;154(3):363-8.
204. Paoletti RD, Suess TM, Lesko MG, et al. Using bar-code technology and medication observation methodology for safer medication administration. *Am J Health Syst Pharm* 2007;64(5):536-43.
205. Persell SD, Denecke-Dattalo TA, Dunham DP, et al. Evidence-based medicine. Patient-directed intervention versus clinician reminders alone to improve aspirin use in diabetes: A cluster randomized trial. *Jt Comm J Qual Patient Saf* 2008;34(2):98-105.
206. Poon EG, Keohane CA, Bane A, et al. Impact of barcode medication administration technology on how nurses spend their time providing patient care. *J Nurs Adm* 2006;38(12):541-9.
207. Poon EG, Keohane CA, Yoon CS, et al. Effect of bar-code technology on the safety of medication administration. *N Engl J Med* 2010;362(18):1698-707.
208. Shirley KL. Effect of an automated dispensing system on medication administration time.[erratum appears in *Am J Health Syst Pharm* 1999 Oct 15;56(20):2142]. *Am J Health Syst Pharm* 1999;56(15):1542-5.
209. Taylor JA, Loan LA, Kamara J, et al. Medication administration variances before and after implementation of computerized physician order entry in a neonatal intensive care unit. *Pediatrics* 2008;121(1):123-8.
210. Wax DB, Beilin Y, Levin M, et al. The effect of an interactive visual reminder in an anesthesia information management system on timeliness of prophylactic antibiotic administration. *Anesth Analg* 2007;104(6):1462-6.
211. Bertsche T, Askoxylakis V, Habl G, et al. Multidisciplinary pain management based on a computerized clinical decision support system in cancer pain patients. *Pain* 2009;147(1-3):20-8.
212. Chambers RM, Wilson JW, Estes LL. Computer-based monitoring as a tool for antimicrobial de-escalation. *Hosp Pharm* 2008;43(3):199-205.

213. DEmakis J, Beauchamp C, Cull W, et al. Improving residents' compliance with standards of ambulatory care: Results from the VA cooperative study on computerized reminders. *JAMA* 2000;284(11):1411-6.
214. Evans RS, Pestotnik SL, Classen DC, et al. Evaluation of a computer-assisted antibiotic-dose monitor. *Ann Pharmacother* 1999;33(10):1026-31.
215. Feldstein AC, Smith DH, Perrin N, et al. Improved therapeutic monitoring with several interventions: a randomized trial. *Arch Intern Med* 2006;166(17):1848-54.
216. Smith DH, Feldstein AC, Perrin NA, et al. Improving laboratory monitoring of medications: an economic analysis alongside a clinical trial. *Am J Manag Care* 2009;15(5):281-9.
217. Grant RW, Wald JS, Schnipper JL, et al. Practice-linked online personal health records for type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med* 2008;168(16):1776-82.
218. Javitt J, Steinberg G, Locke T, et al. Using a claims data-based sentinel system to improve compliance with clinical guidelines: Results of a randomized prospective study. *Am J Manag Care* 2005;11:93-102.
219. Kuilboer MM, van Wijk MA, Mosseveld M, et al. Computed critiquing integrated into daily clinical practice affects physicians' behavior--a randomized clinical trial with AsthmaCritic. *Methods Inf Med* 2006;45(4):447-54.
220. Lo HG, Matheny ME, Seger DL, et al. Impact of non-interruptive medication laboratory monitoring alerts in ambulatory care. *J Am Med Inform Assoc* 2009;16(1):66-71.
221. Matheny ME, Sequist TD, Seger AC, et al. A randomized trial of electronic clinical reminders to improve medication laboratory monitoring. *J Am Med Inform Assoc* 2008;15(4):424-9.
222. McDonald C. Use of a computer to detect and respond to clinical events: Its effect on clinical behavior. *Ann Intern Med* 1976;84(2):162-7.
223. Okon TR, Lutz PS, Liang H. Improved pain resolution in hospitalized patients through targeting of pain mismanagement as medical error. *Journal of Pain & Symptom Management* 2009;37(6):1039-49.
224. Patel PV, Gilski D, Morrison J. Improving outcomes in high-risk populations using REACH: An inpatient cardiac risk reduction program. *Critical Pathways in Cardiology* 2009;8(3):112-8.
225. Rind DM, Safran C, Phillips RS, et al. Effect of computer-based alerts on the treatment and outcomes of hospitalized patients. *Arch Intern Med* 1994;154(13):1511-7.
226. Roumie CL, Elasy TA, Greevy R, et al. Improving blood pressure control through provider education, provider alerts, and patient education: a cluster randomized trial. *Ann Intern Med* 2006;145(3):165-75.
227. Roumie CL, Elasy TA, Wallston KA, et al. Clinical inertia: a common barrier to changing provider prescribing behavior. *Jt Comm J Qual Patient Saf* 2007;33(5):277-85.
228. Sequist TD, Gandhi T, Karson A, et al. A randomized trial of electronic clinical reminders to improve quality of care for diabetes and coronary artery disease. *J Am Med Inform Assoc* 2005;12(4):431-7.
229. White K, Lindsay A, Pryor T, et al. Application of a computerized medical decision-making process to the problem of digoxin intoxication. *J Am Coll Cardiol* 1984;4(3):571-6.
230. Agrawal A, Wu WY. National patient safety goals. Reducing medication errors and improving systems reliability using an electronic medication reconciliation system. *Jt Comm J Qual Patient Saf* 2009;35(2):106-14.
231. Grasso BC, Genest R, Yung K, et al. Reducing errors in discharge medication lists by using personal digital assistants. *Psychiatr Serv* 2002;53(10):1325-6.
232. Poole D, Chainakul J, Pearson M, et al. JHQ 177 medication reconciliation: A necessity in promoting a safe hospital discharge. *J Healthc Qual* 2006;28(3):12-9.

233. van der Kam WJ, Meyboom dJ, Tromp TF, et al. Effects of electronic communication between the GP and the pharmacist. The quality of medication data on admission and after discharge. *Fam Pract* 2001;18(6):605-9.
234. Chan S. Factors Associated With the Use of Electronic Information Systems for Drug Dispensing and Medication Administration Records in Nursing Homes. *J Am Med Dir Assoc* 2008;9(6):414-21.
235. Ghahramani N, Lendel I, Haque R, et al. User satisfaction with computerized order entry system and its effect on workplace level of stress. *J Med Syst* 2009;33(3):199-205.
236. Glassman PA, Belperio P, Simon B, et al. Exposure to automated drug alerts over time: effects on clinicians' knowledge and perceptions. *Med Care* 2006;44(3):250-6.
237. Graumlich JF, Novotny NL, Stephen NG, et al. Patient readmissions, emergency visits, and adverse events after software-assisted discharge from hospital: cluster randomized trial. *Journal of Hospital Medicine (Online)* 2009;4(7):E11-E19
238. Graumlich JF, Novotny NL, Nace GS, et al. Patient and physician perceptions after software-assisted hospital discharge: Cluster randomized trial. *Journal of Hospital Medicine* 2009;4(6):356-63.
239. Holden R J, Scanlon M C, Brown R L and others. What is IT? New conceptualizations and measures of pediatric nurses' acceptance of bar-coded medication administration information technology. In Santa Monica, CA, U.S.: Human Factors and Ergonomics Society; 2008. p.768-72.
240. Holden RJ, Brown RL, Alper SJ, et al. That's nice, but what does IT do? Evaluating the impact of bar coded medication administration by measuring changes in the process of care. *International Journal of Industrial Ergonomics* 2010;(accepted):
241. Hurley AC, Bane A, Fotakis S, et al. Nurses' satisfaction with medication administration point-of-care technology. *J Nurs Adm* 2007;37(7-8):343-9.
242. Kawasumi Y, Tamblyn R, Platt R, et al. Evaluation of the use of an integrated drug information system by primary care physicians for vulnerable population. *Int J Med Inf* 2008;77(2):98-106.
243. Kirking D, Thomas J, Ascione F, et al. Detecting and preventing adverse drug interactions: The potential contribution of computers in pharmacies. *Soc Sci Med* 1986;22(1):1-8.
244. Krlewski JE, Dowd BE, Cole-Adeniyi T, et al. Factors influencing physician use of clinical electronic information technologies after adoption by their medical group practices. *Health Care Manage Rev* 2008;33(4):361-7.
245. Kramer JS, Hopkins PJ, Rosendale JC, et al. Implementation of an electronic system for medication reconciliation. *Am J Health Syst Pharm* 2007;64(4):404-22.
246. Lee F, Teich JM, Spurr CD, et al. Implementation of physician order entry: user satisfaction and self-reported usage patterns. *J Am Med Inform Assoc* 1996;3(1):42-55.
247. Li Q, Douglas S, Hundt A et al. A heuristic usability evaluation of a computerized provider order entry (CPOE) technology. *Proceedings of the IEA2006 Congress*, 2006. Available at: http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGETS_0_204532_0_0_18/Lietal-2006-IEA-CPOEUsability.pdf.
248. Lindenauer PK, Ling D, Pekow PS, et al. Physician characteristics, attitudes, and use of computerized order entry. *Journal of Hospital Medicine (Online)* 2006;1(4):221-30.
249. Liu CT, Yeh YT, Chiang IJ, et al. Development and evaluation of an integrated pharmaceutical education system. *Int J Med Inf* 2004;73(4):383-9.
250. McAlearney A, Chisolm D, Veneris S, et al. Utilization of evidence-based computerized order sets in pediatrics. *Int J Med Inf* 2006;75:501-12.

251. McCann L, Maguire R, Miller M, et al. Patients' perceptions and experiences of using a mobile phone-based advanced symptom management system (ASyMS) to monitor and manage chemotherapy related toxicity. *European Journal of Cancer Care* 2009;18(2):156-64.
252. Musser R, Tcheng J. Quantitative and qualitative comparison of text-based and graphical user interfaces for computerized provider order entry. *Proceedings of the AMIA Symposium* 2006;1041
253. Niazkhani Z, van der SH, Pirnejad H, et al. Same system, different outcomes: comparing the transitions from two paper-based systems to the same computerized physician order entry system. *Int J Med Inf* 2009;78(3):170-81.
254. O'Morrow SD. Nurses' attitudes toward use of an automated medication management system. *Medical University of South Carolina - College of Health Professions* 2003; D H A 2000;14(2):1-13.
255. van Onzenoort HA, van de PA, Kessels AG, et al. Factors influencing bar-code verification by nurses during medication administration in a Dutch hospital. *Am J Health Syst Pharm* 2008;65(7):644-8.
256. Pirnejad H, Niazkhani Z, van der SH, et al. Impact of a computerized physician order entry system on nurse-physician collaboration in the medication process. *Int J Med Inf* 2008;77(11):735-44.
257. Pirnejad H, Niazkhani Z, van der SH, et al. Evaluation of the impact of a CPOE system on nurse-physician communication--a mixed method study. *Methods Inf Med* 2009;48(4):350-60.
258. Porteous T, Bond C, Robertson R, et al. Electronic transfer of prescription-related information: comparing views of patients, general practitioners, and pharmacists. *Br J Gen Pract* 2003;53(488):204-9.
259. Rahimi B, Timpka T, Vimarlund V, et al. Organization-wide adoption of computerized provider order entry systems: a study based on diffusion of innovations theory. *BMC Med Inform Decis Mak* 2009;9:52
260. Rogers J, Jain NL, Hayes GM. Evaluation of an implementation of PRODIGY phase two. *AMIA Proceedings* 1999;604-8.
261. Rohrig R, Beutefuhr H, Hartmann B, et al. Summative software evaluation of a therapeutic guideline assistance system for empiric antimicrobial therapy in ICU. *J Clin Monit Comput* 2007;21(4):203-10.
262. Rosenbloom ST, Geissbuhler AJ, Dupont WD, et al. Effect of CPOE user interface design on user-initiated access to educational and patient information during clinical care. *J Am Med Inform Assoc* 2005;12(4):458-73.
263. Ross SE, Moore LA, Earnest MA, et al. Providing a web-based online medical record with electronic communication capabilities to patients with congestive heart failure: randomized trial. *J Med Internet Res* 2004;6(2):e12
264. Owens DK, McDonald TW, Brown BW, et al. A randomized controlled trial of a computer-based physician workstation in an outpatient setting. Implementation barriers to outcome evaluation. *J Am Med Inform Assoc* 1996;3(5):340-8.
265. Rupp MT, Warholak TL. Evaluation of e-prescribing in chain community pharmacy: best-practice recommendations. *J Am Pharm Assoc* 2008;48(3):364-70.
266. Schectman JM, Schorling JB, Nadkarni MM, et al. Determinants of physician use of an ambulatory prescription expert system. *Int J Med Inf* 2005;74(9):711-7.
267. Schmidt S, Sheikzadeh S, Beil B, et al. Acceptance of telemonitoring to enhance medication compliance in patients with chronic heart failure. *Telemedicine and e-Health* 2008;14(5):426-33.
268. Shannon T, Feied C, Smith M, et al. Wireless handheld computers and voluntary utilization of computerized prescribing systems in the emergency department. *J Emerg Med* 2006;31(3):309-15.
269. Sittig DF, Krall MA, Dykstra RH, et al. A survey of factors affecting clinician acceptance of clinical decision support. *BMC Med Inform Decis Mak* 2006;6:6

270. Tan WS, Phang JS, Tan LK. Evaluating user satisfaction with an electronic prescription system in a primary care group. *Annals of the Academy of Medicine Singapore* 2009;38(6):494-597.
271. Tierney W, Overhage J, McDonald C, et al. Medical students' and housestaff's opinions of computerized order-writing. *Acad Med* 1994;69(5):386-9.
272. Topps C, Lopez L, Messmer PR, et al. Perceptions of pediatric nurses toward bar-code point-of-care medication administration. *Nurs Adm Q* 2005;29(1):102-7.
273. Wang CJ, Patel MH, Schueth AJ, et al. Perceptions of standards-based electronic prescribing systems as implemented in outpatient primary care: a physician survey. *J Am Med Inform Assoc* 2009;16(4):493-502.
274. Weiner M, Gress T, Thiemann DR, et al. Contrasting views of physicians and nurses about an inpatient computer-based provider order-entry system. *J Am Med Inform Assoc* 1999;6(3):234-44.
275. Weingart SN, Hamrick HE, Tutkus S, et al. Medication safety messages for patients via the web portal: the MedCheck intervention. *Int J Med Inf* 2008;77(3):161-8.
276. Wilson JP, Bulatao PT, Rascati KL. Satisfaction with a computerized practitioner order-entry system at two military health care facilities. *Am J Health Syst Pharm* 2000;57(23):2188-95.
277. Zaidi ST, Marriott JL, Nation RL. The role of perceptions of clinicians in their adoption of a web-based antibiotic approval system: do perceptions translate into actions? *Int J Med Inf* 2008;77(1):33-40.
278. Karnon J, McIntosh A, Dean J, et al. Modelling the expected net benefits of interventions to reduce the burden of medication errors. *J Health Serv Res Policy* 2008;13(2):85-91.
279. Plaza V, Cobas A, Ignacio-Garcia J, et al. Cost-effectiveness of an intervention based on the recommendations of the Global Initiative for Asthma (GINA) by a computerized clinical decision support: A trial with randomization of physicians. *Med Clin (Barc)* 2005;124(6):201-6.
280. Wu RC, Laporte A, Ungar WJ. Cost-effectiveness of an electronic medication ordering and administration system in reducing adverse drug events. *J Eval Clin Pract* 2007;13(3):440-8.
281. McMullin ST, Lonergan TP, Rynearson CS, et al. Impact of an evidence-based computerized decision support system on primary care prescription costs. *Annals of Family Medicine* 2004;2(5):494-8.
282. McMullin ST, Lonergan TP, Rynearson CS. Twelve-month drug cost savings related to use of an electronic prescribing system with integrated decision support in primary care. *J Manag Care Pharm* 2005;11(4):322-32.
283. Evans RS, Pestotnik SL, Classen DC, et al. Prevention of adverse drug events through computerized surveillance. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1992;437-41.
284. Evans RS, Classen DC, Pestotnik SL, et al. A decision support tool for antibiotic therapy. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1995;651-5.
285. Macdonald D, Bhalla P, Cass W, et al. Computerized management of oral anticoagulant therapy: experience in major joint arthroplasty. *Can J Surg* 2002;45(1):47-52.
286. Kaushal R, Jha AK, Franz C, et al. Return on investment for a computerized physician order entry system. *J Am Med Inform Assoc* 2006;13(3):261-6.
287. Barenfanger J, Short MA, Groesch AA. Improved antimicrobial interventions have benefits. *J Clin Microbiol* 2001;39(8):2823-8.
288. Ornstein SM, MacFarlane LL, Jenkins RG, et al. Medication cost information in a computer-based patient record system. Impact on prescribing in a family medicine clinical practice. *Arch Fam Med* 1999;8(2):118-21.
289. Tierney W, Miller M, Overhage J, et al. Physician inpatient order writing on microcomputer workstations: effects on resource utilization. *JAMA* 1993;269(3):379-83.

290. Piontek F, Kohli R, Conlon P, et al. Effects of an adverse-drug-event alert system on cost and quality outcomes in community hospitals. *Am J Health Syst Pharm* 2010;67(8):613-20.
291. Weingart SN, Simchowitz B, Padolsky H, et al. An empirical model to estimate the potential impact of medication safety alerts on patient safety, health care utilization, and cost in ambulatory care. *Arch Intern Med* 2009;169(16):1465-73.
292. Balczak TJ, Krumholz HM, Getnick GS, et al. Utilization and effectiveness of a weight-based heparin nomogram at a large academic medical center. *Am J Manag Care* 2000;6(3):329-38.
293. Baroletti S, Munz K, Sonis J, et al. Electronic alerts for hospitalized high-VTE risk patients not receiving prophylaxis: a cohort study. *J Thromb Thrombolysis* 2008;25(2):146-50.
294. Del Beccaro MA, Jeffries HE, Eisenberg MA, et al. Computerized provider order entry implementation: no association with increased mortality rates in an intensive care unit. *Pediatrics* 2006;118(1):290-5.
295. Boord JB, Sharifi M, Greevy RA, et al. Computer-based insulin infusion protocol improves glycemia control over manual protocol. *J Am Med Inform Assoc* 2007;14(3):278-87.
296. Chabot I, Moisan J, Gregoire JP, et al. Pharmacist intervention program for control of hypertension. *Ann Pharmacother* 2003;37(9):1186-93.
297. Chen C, Chen K, Hsu CY, et al. A guideline-based decision support for pharmacological treatment can improve the quality of hyperlipidemia management. *Comput Methods Programs Biomed* 2010;97(3):280-5.
298. Cook CB, Mann LJ, King EC, et al. Management of insulin therapy in urban diabetes patients is facilitated by use of an intelligent dosing system. *Diabetes Technology and Therapeutics* 2004;6(3):326-35.
299. Evans RS, Pestotnik SL, Classen DC, et al. Preventing adverse drug events in hospitalized patients. *Ann Pharmacother* 1994;28(4):523-7.
300. Fiumara K, Piovella C, Hurwitz S, et al. Multi-screen electronic alerts to augment venous thromboembolism prophylaxis. *Thrombosis & Haemostasis* 2010;103(2):312-7.
301. Garthwaite EA, Will EJ, Bartlett C, et al. Patient-specific prompts in the cholesterol management of renal transplant outpatients: results and analysis of underperformance. *Transplantation* 2004;78(7):1042-7.
302. Gurwitz JH, Field TS, Rochon P, et al. Effect of computerized provider order entry with clinical decision support on adverse drug events in the long-term care setting. *J Am Geriatr Soc* 2008;56(12):2225-33.
303. Han YY, Carcillo JA, Venkataraman ST, et al. Unexpected increased mortality after implementation of a commercially sold computerized physician order entry system. *Pediatrics* 2005;116(6):1506-12.
304. Hetlevik I, Holmen J, Kruger O. Implementing clinical guidelines in the treatment of hypertension in general practice: Evaluation of patient outcome related to implementation of a computer-based clinical decision support system. *Scand J Prim Health Care* 1999;17(1):35-40.
305. Holdsworth MT, Fichtl RE, Raisch DW, et al. Impact of computerized prescriber order entry on the incidence of adverse drug events in pediatric inpatients. *Pediatrics* 2007;120(5):1058-66.
306. Holman RR, Smale AD, Pemberton E, et al. Randomized controlled pilot trial of a hand-held patient-oriented, insulin regimen optimizer. *Med Inform (Lond)* 1996;21(4):317-26.
307. Janssen B, Ludwig S, Eustermann H, et al. Improving outpatient treatment in schizophrenia: effects of computerized guideline implementation--results of a multicenter-study within the German research network on schizophrenia. *Eur Arch Psychiatry Clin Neurosci* 2010;260(1):51-7.
308. Keene A, Ashton L, Shure D, et al. Mortality before and after initiation of a computerized physician order entry system in a critically ill pediatric population. *Pediatric Critical Care Medicine* 2007;8(3):268-71.

309. Meigs J, Cagliero E, Dubey A, et al. A controlled trial of web-based diabetes disease management. *Diabetes Care* 2003;26(3):750-7.
310. Murray MD, Harris LE, Overhage JM, et al. Failure of computerized treatment suggestions to improve health outcomes of outpatients with uncomplicated hypertension: results of a randomized controlled trial. *Pharmacotherapy* 2004;24(3):324-37.
311. Pielmeier U, Andreassen S, Juliussen B, et al. The Glucosafe system for tight glycemic control in critical care: A pilot evaluation study. *J Crit Care* 2010;25(1):97-104.
312. Ralston JD, Hirsch IB, Hoath J, et al. Web-based collaborative care for type 2 diabetes: a pilot randomized trial. *Diabetes Care* 2009;32(2):234-9.
313. Schnipper JL, Hamann C, Ndumele CD, et al. Effect of an electronic medication reconciliation application and process redesign on potential adverse drug events: a cluster-randomized trial. *Arch Intern Med* 2009;169(8):771-80.
314. Schnipper JL, Ndumele CD, Liang CL, et al. Effects of a subcutaneous insulin protocol, clinical education, and computerized order set on the quality of inpatient management of hyperglycemia: results of a clinical trial. *Journal of Hospital Medicine (Online)* 2009;4(1):16-27.
315. Takada M, Demizu M, Shibakawa M. Physicians' prescribing attitudes to combined therapy with statins and fibrates. *J Clin Pharm Therapeut* 2003;28(6):445-50.
316. Vartak S, Crandall DK, Brokel JM, et al. Professional practice and innovation: transformation of emergency department processes of care with EHR, CPOE, and ER event tracking systems. *Health Information Management Journal* 2010;38(2):27-32.
317. Yu F, Salas M, Kim YI, et al. The relationship between computerized physician order entry and pediatric adverse drug events: a nested matched case-control study. *Pharmacoepidemiology & Drug Safety* 2009;18(8):751-5.
318. Holden RJ. Physicians' beliefs about using EMR and CPOE: In pursuit of a contextualized understanding of health it use behavior. *Int J Med Inf* 2010;79(2):71-80.
319. Agostini JV, Concato J, Inouye SK. Improving sedative-hypnotic prescribing in older hospitalized patients: provider-perceived benefits and barriers of a computer-based reminder. *J Gen Intern Med* 2008;23(Suppl 1):32-6.
320. Ahearn MD, Kerr SJ. General practitioners' perceptions of the pharmaceutical decision-support tools in their prescribing software. *Med J Aust* 2003;179(1):34-7.
321. Arar NH, Wen L, McGrath J, et al. Communicating about medications during primary care outpatient visits: the role of electronic medical records. *Inform Prim Care* 2005;13(1):13-22.
322. Ash JS, Sittig DF, Seshadri V, et al. Adding insight: a qualitative cross-site study of physician order entry. *Stud Health Technol Inform* 2004;107(Pt:2):2-7.
323. Ash JS, Gorman PN, Lavelle M, et al. A cross-site qualitative study of physician order entry. *J Am Med Inform Assoc* 2003;10(2):188-200.
324. Sittig DF, Krall M, Kaalaas-Sittig J, et al. Emotional aspects of computer-based provider order entry: A qualitative study. *J Am Med Inform Assoc* 2005;12(5):561-7.
325. Ash J, Gorman P, Hersh W, et al. Perceptions of house officers who use physician order entry. *Proceedings of the AMIA Symposium* 1999;471-5.
326. Ash J, Gorman P, Lavelle M, et al. Multiple perspectives on physician order entry. *Proceedings of the AMIA Symposium* 2000;27-31.
327. Ash J, Stavri P, Dykstra R, et al. Implementing computerized physician order entry: the importance of special people. *Int J Med Inf* 2003;69:235-50.
328. Ash J, Gorman P, Lavelle M, et al. Investigating physician order entry in the field: Lessons learned in a multi-center study. *Medinfo* 2001;

329. Avery AJ, Savelyich BS, Sheikh A, et al. Identifying and establishing consensus on the most important safety features of GP computer systems: e-Delphi study. *Inform Prim Care* 2005;13(1):3-12.
330. Bastholm RP, Andersen-Karlsson E, Arnhjort T, et al. Physicians' perceptions of possibilities and obstacles prior to implementing a computerised drug prescribing support system. *Int J Health Care Qual Assur Inc Leadersh Health Serv* 2004;17(4-5):173-9.
331. Beuscart-Zephir M-C, Pelayo S, Bernonville S. Example of a Human Factors Engineering approach to a medication administration work system: Potential impact on patient safety. *Int J Med Inf* 2010;79(4):e43-e57
332. Boonstra A, Boddy D, Fischbacher M. The limited acceptance of an electronic prescription system by general practitioners: Reasons and practical implications. *New Tech Work Employ* 2004;19(2):128-44.
333. Buhner ED, Novak LL, Lorenzi NM. Implementing barcode medication administration: nurses' attitudes. *AMIA* 2008;Annual:1072
334. Campbell EM, Guappone KP, Sittig DF, et al. Computerized provider order entry adoption: implications for clinical workflow. *J Gen Intern Med* 2009;24(1):21-6.
335. Cross RK, Cheevers N, Finkelstein J. Home telemanagement for patients with ulcerative colitis (UC HAT). *Dig Dis Sci* 2009;54(11):2463-72.
336. Crosson JC, Isaacson N, Lancaster D, et al. Variation in electronic prescribing implementation among twelve ambulatory practices. *J Gen Intern Med* 2008;23(4):364-71.
337. Feldstein A, Simon SR, Schneider J, et al. How to design computerized alerts to safe prescribing practices. *Jt Comm J Qual Patient Saf* 2004;30(11):602-13.
338. Fernando S, Georgiou A, Holdgate A, et al. Challenges associated with electronic ordering in the emergency department: a study of doctors' experiences. *Emergency Medicine Australasia* 2009;21(5):373-8.
339. Fields W, Snyder R. Community hospital CPOE system implementation: The lived experience of multi-disciplinary healthcare team members. *AMIA Proceedings* 2007;954
340. Georgiou A, Ampt A, Creswick N, et al. Computerized Provider Order Entry--what are health professionals concerned about? A qualitative study in an Australian hospital. *Int J Med Inf* 2009;78(1):60-70.
341. Graham TA, Kushniruk AW, Bullard MJ, et al. How usability of a web-based clinical decision support system has the potential to contribute to adverse medical events. *AMIA* 2008;Annual:257-61.
342. Grossman JM, Gerland A, Reed MC, et al. Physicians' experiences using commercial e-prescribing systems. *Health Aff (Millwood)* 2007;26(3):w393-w404
343. Johansson PE, Petersson GI, Nilsson GC. Personal digital assistant with a barcode reader-A medical decision support system for nurses in home care. *Int J Med Inf* 2010;79(4):232-42.
344. Kazemi A, Ellenius J, Tofighi S, et al. CPOE in Iran--a viable prospect? Physicians' opinions on using CPOE in an Iranian teaching hospital. *Int J Med Inf* 2009;78(3):199-207.
345. Koppel R, Metlay J, Cohen A, et al. Role of computerized physician order entry systems in facilitating medication errors. *JAMA* 2005;293(10):1197-203.
346. Koppel R, Wetterneck T, Telles JL, et al. Workarounds to barcode medication administration systems: their occurrences, causes, and threats to patient safety. *J Am Med Inform Assoc* 2008;15(4):408-23.
347. Krall MA, Sittig DF. Clinician's assessments of outpatient electronic medical record alert and reminder usability and usefulness requirements. *AMIA Proceedings* 2002;400-4.
348. Lai JS, Yokoyama G, Louie C, et al. Impact of computerized prescriber order entry (CPOE) on clinical pharmacy practice: A hypothesis-generating study. *Hosp Pharm* 2007;42(10):931-8.

349. McAlearney AS, Chisolm DJ, Schweikhart S, et al. The story behind the story: physician skepticism about relying on clinical information technologies to reduce medical errors. *Int J Med Inf* 2007;76(11-12):836-42.
350. Motulsky A, Winslade N, Tamblyn R, et al. The impact of electronic prescribing on the professionalization of community pharmacists: a qualitative study of pharmacists' perception. *Journal of Pharmacy & Pharmaceutical Sciences* 2008;11(1):131-46.
351. Nanji KC, Cina J, Patel N, et al. Overcoming barriers to the implementation of a pharmacy bar code scanning system for medication dispensing: a case study. *J Am Med Inform Assoc* 2009;16(5):645-50.
352. Novak LL, Lorenzi NM. Barcode medication administration: supporting transitions in articulation work. *AMIA* 2008;Annual:515-9.
353. Novek J, Bettess S, Burke K, et al. Nurses' perceptions of the reliability of an automated medication dispensing system. *J Nurs Care Qual* 2000;14(2):1-13.
354. O'Grady K, Donyai P, Franklin BD. Patients' views about an electronic prescribing and drug administration system in secondary care. *BJHC & IM* 2006;23(7):15-8.
355. Patterson ES, Cook RI, Render ML. Improving patient safety by identifying side effects from introducing bar coding in medication administration. *J Am Med Inform Assoc* 2002;9(5):540-53.
356. Patterson ES, Nguyen A, Halloran J, et al. Human Factors Barriers to the Effective Use of Ten HIV Clinical Reminders. *J Am Med Inform Assoc* 2004;11(1):50-9.
357. Ruiz JG, Qadri SS, Nader S, et al. Primary care management of chronic nonmalignant pain in veterans: a qualitative study. *EDUC GERONTOL* 2010;36(5):372-93.
358. Saleem JJ, Patterson ES, Militello L, et al. Exploring barriers and facilitators to the use of computerized clinical reminders. *J Am Med Inform Assoc* 2005;12(4):438-47.
359. Schoville RR. Work-arounds and artifacts during transition to a computer physician order entry: what they are and what they mean. *J Nurs Care Qual* 2009;24(4):316-24.
360. Varonen H, Kortteisto T, Kaila M, et al. What may help or hinder the implementation of computerized decision support systems (CDSSs): a focus group study with physicians. *Fam Pract* 2008;25(3):162-7.
361. Vaziri A, Connor E, Shepherd I, et al. Are we setting about improving the safety of computerised prescribing in the right way? A workshop report. *Inform Prim Care* 2009;17(3):175-82.
362. Vogelsmeier AA, Halbesleben JR, Scott-Cawiezell JR. Technology implementation and workarounds in the nursing home. *J Am Med Inform Assoc* 2008;15(1):114-9.
363. Weingart SN, Massagli M, Cyrulik A, et al. Assessing the value of electronic prescribing in ambulatory care: A focus group study. *Int J Med Inf* 2009;78(9):571-8.
364. Weir C, Lincoln M, Roscoe D, et al. Dimensions associated with successful implementation of a hospital based integrated order entry system. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1994;653-7.
365. Weir CR, Nebeker JJR, Hicken BL, et al. A cognitive task analysis of information management strategies in a computerized provider order entry environment. *J Am Med Inform Assoc* 2007;14(1):65-75.
366. Wentzer HS, Bottger U, Boye N. Unintended transformations of clinical relations with a computerized physician order entry system. *Int J Med Inf* 2007;76(Suppl 3):S453-S461
367. Holbrook A, Thabane L, Keshavjee K, et al. Individualized electronic decision support and reminders to improve diabetes care in the community: COMPETE II randomized trial. *CMAJ Canadian Medical Association Journal* 2009;181(1-2):37-44.
368. Ash JS, Sittig DF, Campbell EM, et al. Some unintended consequences of clinical decision support systems. *AMIA Proceedings* 2007;26-30.

369. Ash JS, Sittig DF, Poon EG, et al. The extent and importance of unintended consequences related to computerized provider order entry. *J Am Med Inform Assoc* 2007;14(4):415-23.
370. Ash JS, Sittig DF, Dykstra RH, et al. Categorizing the unintended sociotechnical consequences of computerized provider order entry. *Int J Med Inf* 2007;76(Suppl 1):S1-S7
371. Campbell EM, Sittig DF, Ash JS, et al. Types of unintended consequences related to computerized provider order entry. *J Am Med Inform Assoc* 2006;13(5):547-56.
372. Campbell EM, Sittig DF, Guappone KP, et al. Overdependence on technology: an unintended adverse consequence of computerized provider order entry. *AMIA* 2007;94-8.
373. Ash JS, Sittig DF, Campbell E, et al. An unintended consequence of CPOE implementation: shifts in power, control, and autonomy. *AMIA* 2006;11-5.
374. Campbell EM. Computerized provider order entry adoption: Implications for clinical workflow. *J Gen Intern Med* 2009;24(1):21-6.
375. Santell JP, Kowiatek JG, Weber RJ, et al. Medication errors resulting from computer entry by nonprescribers. *Am J Health Syst Pharm* 2009;66(9):843-53.
376. Singh H, Mani S, Espadas D, et al. Prescription errors and outcomes related to inconsistent information transmitted through computerized order entry: a prospective study. *Arch Intern Med* 2009;169(10):982-9.
377. Field TS, Rochon P, Lee M, et al. Costs associated with developing and implementing a computerized clinical decision support system for medication dosing for patients with renal insufficiency in the long-term care setting. *J Am Med Inform Assoc* 2008;15(4):466-72.

Appendix D. Technical Expert Panel and Peer Reviewers

Technical Expert Panel

David Bates, M.D., M.Sc

Director, The Center for Patient Safety Research and Practice Partners
Brigham and Women's Hospital
Boston, MA

Chris Gibbons, M.D., M.P.H.

Assistant Professor
Associate Director, Johns Hopkins Urban Health Institute
Johns Hopkins Bloomberg School of Public Health
Baltimore, MD

Joseph Hanlon, Pharm.D, M.S.

Professor of Medicine
Health Scientist
University of Pittsburgh
Pittsburgh, PA

Kevin Johnson, M.D., M.S.

Associate Professor
Department of Pediatrics
Vanderbilt University Medical Center
Nashville, TN

Bimla Schwarz, M.D., M.S.

Assistant Professor of Medicine
Assistant Professor of Obstetrics, Gynecology and Reproductive Science
Assistant Professor of Epidemiology
University of Pittsburgh
Pittsburgh, PA

Doug Bell, Ph.D., M.D.

Research Scientist, RAND Corporation;
Associate Professor, Department of Medicine
University of California
Los Angeles CA

Gordon Schiff, M.D.

Associate Professor of Medicine, Harvard Medical School
Clinical Researcher, Division of General Medicine
Brigham and Women's Hospital
Center for Patient Safety Research and Practice
Boston, MA

Jerry Gurwitz, M.D.

Executive Director, Meyers Primary Care Institute
The Doctor John Meyers Professor of Primary Care Medicine
University of Massachusetts
Boston, MA

Joy Grossman, Ph.D.

Senior Health Researcher
Center for Studying Health System Change
Washington, DC

Dennis Tribble, Pharm.D

Chief Technology Officer
ForHealth Technologies, Inc.
Daytona Beach, FL

John Poikonen, Pharm.D

Director of Clinical Informatics
UMass Memorial Medical Center
Worcester, MA

Kenneth Boockvar, M.D., M.S.

Associate Professor, Brookdale Department of Geriatrics and Adult Development Mount Sinai
School of Medicine
New York, NY

Reviewers

Anne Bobb, B.S.Pharm.

Department of Quality and Clinical Informatics
Northwestern Memorial Hospital
Chicago, IL

Elizabeth Chrischilles, M.S. Ph.D.

Director, Health Effectiveness Research Center
University of Iowa
Iowa City, IA

Allen J. Flynn, Pharm.D
Medication Lead CPOE
University of Michigan Hospitals
Ann Arbor, MI

Joy Grossman, Ph.D.
Senior Health Researcher
Center for Studying Health System Change
Washington, DC

Joseph Hanlon, Pharm.D, M.S.
Professor of Medicine
Health Scientist
University of Pittsburgh

Kevin Johnson, M.D., M.S.
Associate Professor
Department of Pediatrics
Vanderbilt University Medical Center

Kevin Marvin, R.Ph., M.S.
Consultant
Medication Systems Informatics
Burlington, VT

John Poikonen, Pharm.D
Director of Clinical Informatics
UMass Memorial Medical Center
Worcester, MA

Bimla Schwarz, M.D., M.S.
Assistant Professor of Medicine
Assistant Professor of Obstetrics, Gynecology and Reproductive Science
Assistant Professor of Epidemiology
University of Pittsburgh

Dennis Tribble, Pharm.D
Chief Technology Officer
ForHealth Technologies, Inc.
Daytona Beach, FL

Appendix E. Excluded Studies

Organized security. To address compliance and IT security, Piedmont Healthcare chose to establish baseline metrics for its security risk. Health Manag Technol 1918;30(12):14-6, 18. OVID EMBASE.

Exclude - Not a Primary Study

New device stems loss of drug charges in ER. Drug Topics 1982;134:25 Database: IPA.

Exclude - Not a Primary Study

Bar-coded unit doses: Coming to a hospital near you. Pharm Times 1983;70(12):11

Database: IPA.

Exclude - Not a Primary Study

E-scrips could save billions. Pharm Times 1983;70(6):16 Database: IPA.

Exclude - Not a Primary Study

Bars and stripes forever? ... Well, maybe .. Hosp Purch Manage 1985;10(Dec):3-8. Database: IPA.

Exclude - Not MMIT

Pharmacy microcomputer software exchange. Hosp Pharm 1986;21(Apr):384-5. Database: IPA.

Exclude - Not MMIT

GPs favor computer prescribing. Pharmaceutical Journal 1991;235(Nov 23):678 Database: IPA.

Exclude - No Outcomes of Interest

Medical informatics needs assessment. 1993.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_846504_0_0_18/medinds3.doc Grey Lit.

Exclude - Not a Primary Study

Technology in pharmacy: Friend or foe? Computertalk for the Pharmacist 1993;14(Nov-Dec):16-23. Database: IPA.

Exclude - Not a Primary Study

Automated decentralized pharmacy dispensing systems. Health Devices 1996;25(12):452-73.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ciba-Geigy program warns doctors. Pharmaceutical Journal 1996;236(Feb 1):134 Database: IPA.

Exclude - No Outcomes of Interest

Computer scripts favoured--survey. Pharmaceutical Journal 1996;236(Mar 8):300 Database: IPA.

Exclude - No Outcomes of Interest

Equipment management guide. Improving the drug distribution process--do you need an automated decentralized pharmacy dispensing system? Health Devices 1996;25(12):441-51. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Top-priority actions for preventing adverse drug events in hospitals: Recommendations of an expert panel. Am J Health Syst Pharm 1996;53(Apr 1):747-51. Database: IPA.

Exclude - Not MMIT

Driving change: How pharmacists are innovating to meet real needs. Computertalk for the Pharmacist 1997;17(Jul-Aug):20-33. Database: IPA.

Exclude - Not a Primary Study

Physician order entry system cuts error rate, improves path compliance, tracks data. Health Care Cost Reengineering Rep 1997;2(10):152-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Pick of the crop: Vendor profiles. Computertalk for the Pharmacist 1997;17(Mar-Apr):10-81. Database: IPA.

Exclude - Not a Primary Study

Proceedings of the 1997 9th Annual Quest for Quality and Productivity in Health Services. In St.Louis, MO, USA: IIE; 1997. 1998023929188

Database: Compendex.

Exclude - Not MMIT

State of the chains: Special report. Computertalk for the Pharmacist 1997;17(Nov-Dec):19-27. Database: IPA.

Exclude - Not a Primary Study

Even in best-run centers, preventable medication errors lengthen LOS, increase costs. Health Care Cost Reengineering Rep 1998;3(10):149-51. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Electronic point-of-care prescribing: Selling the benefits, identifying initial steps for success. Formulary 1999;34(1): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Antibiotic-related ADEs plummet and pharmacy costs shrink with computer-aided decision support. Clinical Resource Management 2000;1(10):151-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Can an automatic alert system eliminate medication errors? Hospital system gives it a go. Clinical Resource Management 2000;1(10):154-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chains speak out on technology. Computertalk for the Pharmacist 2000;20(Jul-Aug):14-25. Database: IPA.

Exclude - Not a Primary Study

Electronic prescribing reduces Rx costs, boosts generic utilization for physician group. Formulary 2000;35(1): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Healtheon/WebMD: Connecting health care. Computertalk for the Pharmacist 2000;20(Jul-Aug):26-9. Database: IPA.

Exclude - Not a Primary Study

Patient safety/medication safety: the impact of computerized physician order entry on medication error prevention in hospitalized patients (project). The Netherlands Organisation for Health Research and Development (ZonMw) 2000; <http://www.zonmw.nl/> Grey Lit saved abstract -describes a research study not yet complete.

Exclude - Not a Primary Study

Bar code labeling standards proposed for drug product packaging. Am J Health Syst Pharm 2001;58(Sep 1):1582 Database: IPA.

Exclude - Not a Primary Study

Computerized physician order entry. Institute for Clinical Systems Improvement; 2001. <http://www.icsi.org/index.asp> Grey Lit.

Exclude - Not a Primary Study

Computerized provider order entry systems. Health Devices 2001;30(9-10):323-59.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Making health care safer: A critical analysis of patient safety practices. AHRQ; 2001. saved as NYAM AHRQ.pdf

Grey Lit.

Exclude - Not a Primary Study

RxHUB venture concerns independents: NCPA calls on FTC, Congress to investigate. American Journal of Clinical Dermatology 2001;123(Jun):13-6. Database: IPA.

Exclude - Not MMIT

Technologies expedite distribution within hospital setting. Healthcare Distributors 2001;53(Feb-Mar):30-1. Database: IPA.

Exclude - Not MMIT

Bar coding landscape. Healthcare Information and Management Systems Society; 2002. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=29279> Grey Lit.

Exclude - Not a Primary Study

Can e-prescribing reduce drug costs under capitation? Capitation Manag Rep 2002;9(4):58-62. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Computers. Good medicine: e-prescribing. Fam Pract Manag 2002;9(9):63-4. Database: CINAHL.

Exclude - Not MMIT

CPOE - An executive's guide. Hospitals & Health Networks 2002;76(6):41-3. Database: IPA.

Exclude - Not a Primary Study

Executive preview: The value of CPOE in ambulatory settings. 2002.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=29895> Grey Lit.
Exclude - Not a Primary Study

What's hot in the chain drug market - A report on the technology agenda based on a survey of nearly 40 large and small chains. Computertalk for the Pharmacist 2002;22(4):14-8.
Database: IPA.
Exclude - Not a Primary Study

What's on the technology agenda in 2002? A sector analysis. Computertalk for the Pharmacist 2002;22(1):17-8. Database: IPA.
Exclude - Not a Primary Study

Advanced technologies to lower healthcare costs and improve quality. Massachusetts Technology Collaborative Board of Directors; 2003.
http://www.masstech.org/ehealth/STATFinal9_24.pdf Grey Lit.
Exclude - Not a Primary Study

Bar-coded medication administration (BCMA) systems. Future promise, present challenges. Health Devices 2003;32(10):373-81. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Barcodes. Computertalk for the Pharmacist 2003;23(6):22-6. Database: IPA.
Exclude - Not a Primary Study

Computerized physician order entry. Institute for Clinical Systems Improvement (ICSI); 2003. Grey Lit.
Exclude - Not MMIT

CPOE cuts time needed to deliver meds, X-rays. Healthc Benchmarks Qual Improv 2003;10(4):44-5. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Drug bar code regulation proposed. WHO Drug Information 2003;17(1):17 Database: IPA.
Exclude - Not MMIT

E-Rx is coming and we're ready. America's Pharmacist 2003;125(4):16-21. Database: IPA.
Exclude - Not a Primary Study

National survey of physicians and quality of care. 2003.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_865229_0_0_18/2003_physician_survey.pdf Grey Lit.
Exclude - Not a Primary Study

Report: CPOE adoption a long-term process. Healthc Benchmarks Qual Improv 2003;10(9):105-7. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Scanning medication barcodes improves accuracy at Lehigh Valley Hospital. Performance Improvement Advisor 2003;7(10):132-4. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Special chain market report. Computertalk for the Pharmacist 2003;23(4):19-20. Database: IPA.

Exclude - Not a Primary Study

The cost of computing. Mark Health Serv 2003;23(2):5 Database: BSC.

Exclude - Not MMIT

The technology agenda for 2003. Computertalk for the Pharmacist 2003;23(1):20-2.

Database: IPA.

Exclude - Not a Primary Study

Use these tools to comply with patient safety goals. ED Manag 2003;15(3):27-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

'Most wired' hospitals widen gap over others. Healthc Benchmarks Qual Improv 2004;11(10):116-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

2004 HIMSS nursing informatics survey. Healthcare Information and Management Systems Society; 2004. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67708> Grey Lit.

Exclude - Not MMIT

Bar code labeling of unit dose packages recommended. Am J Hosp Pharm 2004;44(Jan):16

Database: IPA.

Exclude - Not MMIT

Code compliance. Manufacturing Chemist 2004;75(8):35-6. 8202114

Database: Inspec.

Exclude - Not a Primary Study

Computer feedback promotes generic drug prescribing. Am Pharm 2004; Database: IPA.

Exclude - No Outcomes of Interest

e-Commerce enabled dispensing. Pharmacy Review 2004;27(3): Database: IPA.

Exclude - Not a Primary Study

E-commerce in pharmacy: Opportunity or threat? Acta Farmaceutica Bonaerense 2004;9(3):8-9. Database: IPA.

Exclude - Not MMIT

e-Health in Massachusetts. 2004. http://www.masstech.org/ehealth/11_05/exsumbest.pdf Grey Lit.

Exclude - No Outcomes of Interest

e-prescriptions: Surescripts making a difference? Computertalk for the Pharmacist 2004;24(3):16 Database: IPA.

Exclude - Not a Primary Study

E-scripts to provide new loyalty opportunities for pharmacy. Australian Journal of Pharmacy 2004;83(982):54 Database: IPA.

Exclude - Not MMIT

Electronic prescribing: Planning and implementation to achieve success and maximize value. eHealth Initiative; 2004.

[http://ehr.medigent.com/assets/collaborate/2005/04/02/ElectronicPrescribing_CCBH Annual Meeting_2004_.pdf](http://ehr.medigent.com/assets/collaborate/2005/04/02/ElectronicPrescribing_CCBH%20AnnualMeeting_2004_.pdf) Grey Lit.

Exclude - Not a Primary Study

Electronic prescription tested in Portalegre. Farmacia Portuguesa 2004;27: Database: IPA.

Exclude - Not MMIT

FDA issues final bar coding rule for drugs and blood. Healthc Financ Manage 2004;58(4):12 Database: BSC.

Exclude - Not MMIT

HIMSS ambulatory technology survey. Healthcare Information and Management Systems Society; 2004. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=42882> Grey Lit.

Exclude - Not MMIT

Leapfrog patient-safety standards a stretch for most hospitals. Care Management 2004;10(2):39 Database: CINAHL.

Exclude - Not a Primary Study

Longitudinal records enable instant QI changes. Healthc Benchmarks Qual Improv 2004;11(12):136-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Massive NHS IT project goes under the microscope. IEE Review 2004;50(9):15 Database: BSC.

Exclude - Not MMIT

Medicare reform at the 1-year mark. Pharmacy Today (Washington D C) 2004;10(12):22 Database: IPA.

Exclude - Not a Primary Study

Philadelphia hosts pharma innovations. Manufacturing Chemist 2004;69(May):33 Database: IPA.

Exclude - Not MMIT

Physician order entry perception study: FOCUS: Leading physician order entry -- clinical information system vendors. ElectronicHealthcare 2004;1(3):60-3. Database: CINAHL.

Exclude - Not MMIT

Prescribing connectivity. America's Pharmacist 2004;126(6):30-1. Database: IPA.

Exclude - Not a Primary Study

Recommendations to reduce Rx errors. American Druggist 2004;213(Oct):18 Database: IPA.

Exclude - Not a Primary Study

Senior manager survey addresses CPOE barriers. Healthc Financ Manage 2004;58(8):21 Database: BSC.

Exclude - Not MMIT

The EU's prescription for EHR success. Information Management Journal 2004;38(4):24
<http://search.ebscohost.com/login.aspx?direct=true&db=lxh&AN=13791750&site=ehost-live>
Database: LISTA.

Exclude - Not MMIT

The HIT report from KLAS. Medication administration 2004: FOCUS: Medication administration application performance. HEALTHC Q 2004;8(2):107-10. Database: CINAHL.

Exclude - Not a Primary Study

Today's priority. Computertalk for the Pharmacist 2004;24(4):26-8. Database: IPA.

Exclude - Not a Primary Study

Treatment plan: High tech transfusion. Case statement for implementation of CPOE in all Massachusetts hospitals. Westborough, MA: Massachusetts Technology Collaborative; 2004.
http://www.masstech.org/ehealth/health_final_4.0.pdf Grey Lit.

Exclude - Not a Primary Study

Where we are headed with technology - What industry executives have to say. Computertalk for the Pharmacist 2004;24(1):20-4. Database: IPA.

Exclude - Not a Primary Study

"As directed" is never acceptable for prescription directions and frequency. America's Pharmacist 2005;127(3):16 Database: IPA.

Exclude - Not a Primary Study

At a tipping point: Transforming medicine with health information technology a guide for consumers. MedStar Health e-Health Initiative; 2005.

http://ehr.medigent.com/assets/collaborate/2005/04/14/April12MedStar_final.pdf Grey Lit.

Exclude - Not a Primary Study

Bar coding helps hospitals reduce medication errors. Healthc Benchmarks Qual Improv 2005;12(5):49-52. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

HIMSS analysis of CPOE study: Implemented correctly, CPOE can help reduce medical errors. Healthcare Purchasing News 2005;

http://findarticles.com/p/articles/mi_m0BPC/is_4_29/ai_n13650601/ Database: BSC.

Exclude - Not MMIT

HIMSS AutoID and bar coding task force reports on bar coding for automated patient identification. Healthcare Information and Management Systems Society; 2005.

<http://www.himss.org/asp/ContentRedirector.asp?ContentID=65519> Grey Lit.

Exclude - Not MMIT

Hospitals still error-prone: medical mistakes continue even in highly computerized facilities. Industrial Engineer 2005; <http://www.allbusiness.com/specialty-businesses/476352-1.html>

Database: BSC.

Exclude - Not MMIT

Leapfrog group releases hospital quality and safety survey. Quality Progress 2005;38(1):16 Database: BSC.

Exclude - Not MMIT

Medical records institute's seventh annual survey of electronic health record: Trends and usage for 2005. 2005.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_846502_0_0_18/ehrsurvey05.pdf Grey Lit.

Exclude - Not MMIT

Medicare program: E-prescribing and the prescription drug program. Department of Health and Human Services; 2005. <http://www.cms.hhs.gov/EPrescribing/Downloads/E-Prescribingfinalruleonfoundationstandards.pdf> Grey Lit.

Exclude - Not a Primary Study

Medication safety issue brief: Bar code implementation strategies. Hosp Health Netw 2005;79(7): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

NHIN and RHIOs: Getting started - Organizing models for regional health information organizations. Manatt; 2005.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67597> Grey Lit.

Exclude - Not a Primary Study

Strategies to reduce medication errors with reference to older adults. Aust Nurs J 2005;14(4):26-9. Database: CINAHL.

Exclude - Not a Primary Study

Study shows limited use of electronic medical records. Biomed Instrum Technol 2005;39(5): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Study: CPOE facilitates 22 types of medication errors. Healthc Benchmarks Qual Improv 2005;12(5):52-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tech trends. America's Pharmacist 2005;127(1):19-21. Database: IPA.

Exclude - Not a Primary Study

A primer on the new physician self-referral exceptions and anti-kickback safe harbors for electronic prescribing and electronic health records technology. 2006.

http://www.himss.org/content/files/Stark_Anti-kickback_exceptions.PDF Grey Lit.

Exclude - Not a Primary Study

Assessing the economics of EMR adoption and successful implementation in physician small practice settings. Moshman Associates Inc., Booz Allen Hamilton; 2006.

http://library.ahima.org/xpedio/groups/public/documents/government/bok1_035685.pdf Grey Lit.

Exclude - Not a Primary Study

Comparing EMR adoption to care outcomes at UHC hospitals, including Davies award winners, using HIMSS analytics' EMR adoption model scores. Healthcare Information and Management Systems Society; 2006. <http://www.himssanalytics.org/PDFFiles/UHC25.pdf> Grey Lit.

Exclude - Not MMIT

CPOE study shows drop in hospital errors. Healthc Benchmarks Qual Improv 2006;13(5):54-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Electronic health records overview. National Institutes of Health; 2006.

<http://www.ncrr.nih.gov/publications/informatics/EHR.pdf> Grey Lit.

Exclude - Not a Primary Study

Health information technology promotion act of 2006. 2006.

http://www.himss.org/Content/files/HR4157_Report.pdf Grey Lit.

Exclude - Not a Primary Study

Missouri current status working group survey tool. 2006.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_874957_0_0_18/Missouri.attachment.php.doc Grey Lit.

Exclude - Not a Primary Study

Ontario's medication management system - Transformation wanted! Emergis Inc, Courtyard Group Ltd.; 2006. Grey Lit.

Exclude - Not a Primary Study

Regulatory changes affecting health information exchanges. Deloitte Center for Health Solutions; 2006.

http://www.deloitte.com/dtt/cda/doc/content/us_chs_RegulatoryPaper_HIE010906.pdf Grey Lit.

Exclude - Not a Primary Study

Saving lives, reducing costs: Computerized physician order entry lessons learned in community hospitals. Massachusetts Technology Collaborative; 2006.

http://www.masstech.org/ehealth/CPOE_lessonslearned.pdf Grey Lit.

Exclude - Not a Primary Study

Testing times. Brand 2006;5(5):42-8. 20063710110044

Database: Compendex.

Exclude - Not MMIT

University health network achieves over 85% CPOE with Misys CPR, improves accuracy and saves time. HEALTHC Q 2006;10:Spec-6 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

White paper. Meru Networks; 2006.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67695> Grey Lit.

Exclude - Not a Primary Study

2007 HIMSS nursing informatics survey. Healthcare Information and Management Systems Society; 2007. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67709> Grey Lit.

Exclude - Not MMIT

Anon, editor. 2007 Joint workshop on high confidence medical devices, software, and systems and medical device plug-and-play interoperability. In Anon, editor. Cambridge, MA, United states: Institute of Electrical and Electronics Engineers Computer Society; 2007. 20083211433252

Database: Compendex.

Exclude - Not MMIT

2007 Physician quality reporting initiative (PQRI): Module one. 2007.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=66661> Grey Lit.

Exclude - Not MMIT

2007 State legislation on health information exchanges and networks. National Conference of State Legislatures; 2007.

http://www.ncsl.org/programs/health/forum/hitch/HIE_networks.htm Grey Lit.

Exclude - Not MMIT

A vision of health care from and framework for the state alliance for e-health. 2007.

<http://www.nga.org/Files/pdf/0701EHEALTHFRAMEWORK.PDF> Grey Lit.

Exclude - Not a Primary Study

Barriers to equitable access to quality health information with emphasis on developing countries. The Rockefeller Foundation; 2007. http://www.ehealth-connection.org/files/conf-materials/Barriers%20to%20Equitable%20Access_0.pdf Grey Lit.

Exclude - Not MMIT

Cutting the cord: Wireless connectivity for healthcare. Healthcare Information and Management Systems Society; 2007.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68119> Grey Lit.

Exclude - Not a Primary Study

Designing inherent safety into electronic medication order entry systems. Healthcare Information and Management Systems Society; 2007.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67110> Grey Lit.

Exclude - Not a Primary Study

Determining health informatics workforce needs in developing countries. The Rockefeller Foundation; 2007. http://www.ehealth-connection.org/files/conf-materials/Determining%20Health%20Informatics_0.pdf Grey Lit.

Exclude - Not MMIT

eHealth policy: The road to the new digital divide? The Rockefeller Foundation; 2007.

http://www.ehealth-connection.org/files/conf-materials/eHealth%20Policy%20-%20The%20Road%20to%20the%20New%20Digital%20Divide_0.pdf Grey Lit.

Exclude - Not MMIT

EHR implementation in ambulatory care. Healthcare Information and Management Systems Society; 2007. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67208> Grey Lit.

Exclude - Not a Primary Study

Endorsing the adoption of electronic medical records and health information exchange systems that improve the quality, safety and value of health care. Rhode Island; 2007. <http://www.rilin.state.ri.us/billtext07/senatetext07/s1085.pdf> Grey Lit.

Exclude - Not a Primary Study

Evaluations of the impact of eHealth technologies in developing countries: A systematic review. The Rockefeller Foundation; 2007. http://www.ehealth-connection.org/files/conf-materials/Evaluation%20of%20the%20Impact%20of%20eHealth%20Technologies_0.pdf Grey Lit.

Exclude - Not a Primary Study

Health information exchange: From start up to sustainability. eHealth Initiative; 2007. http://ehr.medigent.com/assets/collaborate/2007/07/10/Health_Information_Exchange-Start_Up_to_Sustainability_Full_Report_07.09.2007001.pdf Grey Lit.

Exclude - Not a Primary Study

Health information technology champions (HITCh). National Conference of State Legislatures; 2007. <http://www.ncsl.org/programs/health/forum/hitch/> Grey Lit.

Exclude - Not MMIT

Health information technology champions (HITCh): 2007 Enacted legislation on health information technology. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/programs/health/forum/Hitch/enacted.htm> Grey Lit.

Exclude - Not MMIT

Health information technology champions (HITCh): 2007 State legislation on health information technology financing. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/programs/health/forum/Hitch/finance.htm> Grey Lit.

Exclude - Not MMIT

HIMSS/GSA national e-authentication project whitepaper. Healthcare Information and Management Systems Society; 2007.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67206> Grey Lit.

Exclude - Not a Primary Study

HIT and telemedicine: Part II of a three part series on the state alliance for e-health. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/print/health/forum/Telemedicine.pdf> Grey Lit.

Exclude - Not a Primary Study

House Bill 134: An act relative to electronic prescribing for prescription drugs. 2007.

<http://www.gencourt.state.nh.us/legislation/2007/HB0134.html> Grey Lit.

Exclude - Not a Primary Study

Interoperability standards for health information systems. The Rockefeller Foundation; 2007.

<http://www.ehealth-connection.org/files/conf-materials/Interoperability%20Standards%20for%20Health%20Information%20Systems.pdf> Grey Lit.

Exclude - No Outcomes of Interest

Interoperable digital identity management in the electronic exchange of health information. Signatures and Authentication for Everyone; 2007.

<http://toolkit.ehealthinitiative.org/Interoperable%20Digital%20ID%20Managment%20Report.pdf> Grey Lit.

Exclude - Not a Primary Study

Leapfrog Group releases its first 'top hospitals' list. Healthc Benchmarks Qual Improv 2007;14(1):1-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Major health information exchange legislation: 2007 Comparison of five state proposals. National Conference of State Legislatures; 2007.

http://www.ncsl.org/programs/health/forum/hitch/five_state.htm Grey Lit.

Exclude - Not MMIT

Medication management detailed use case. U.S. Department of Health and Human Services; 2007. Grey Lit.

Exclude - Not a Primary Study

Mobile services evolution. The Rockefeller Foundation; 2007. <http://www.ehealth-connection.org/files/conf-materials/Mobile%20Services%20Evolution.pdf> Grey Lit.

Exclude - Not MMIT

NCSL health information technology champions (HITCh) policy partnership. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/programs/health/forum/Hitch/scopingmeet0806b.htm> Grey Lit.

Exclude - Not MMIT

NCSL project HITCh phase I report. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/programs/health/forum/hitch/hitchreport.htm> Grey Lit.

Exclude - Not MMIT

North star state illuminates one way toward e-health. National Conference of State Legislatures; 2007. <http://www.ncsl.org/programs/health/shn/2007/sn501a.htm> Grey Lit.

Exclude - Not MMIT

Pharmacy informatics task force: Managing a barcoded medication inventory. Healthcare Information and Management Systems Society; 2007.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67111> Grey Lit.

Exclude - Not a Primary Study

Phase II HITCh work plan. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/programs/health/forum/Hitch/phaseIIplan.htm> Grey Lit.

Exclude - Not MMIT

Regional health information exchanges (RHIOs) & health information exchanges (HIEs) overview. Healthcare Information and Management Systems Society (HIMSS); 2007.

http://www.himss.org/content/files/RHIO/RHIO_HIE_11_10_07.pdf Grey Lit.

Exclude - Not a Primary Study

Registries for evaluating patient outcomes: A user's guide. AHRQ; 2007.
http://library.ahima.org/xpedio/groups/public/documents/government/bok1_037334.pdf Grey Lit.

Exclude - Not a Primary Study

Report from the health information protection taskforce to the state alliance for e-health. US Department of Health and Human Services; 2007.

<http://www.nga.org/Files/pdf/0708EHEALTHREPORT.PDF> Grey Lit.

Exclude - Not a Primary Study

Second report from the health care practice taskforce to the state alliance for e-health. US Department of Health and Human Services; 2007.

<http://www.nga.org/files/pdf/0710EHEALTHHCPREPORT.PDF> Grey Lit.

Exclude - Not a Primary Study

Security standards: Implementation for the small provider. 2007.

http://library.ahima.org/xpedio/groups/public/documents/government/bok1_037819.pdf Grey Lit.

Exclude - Not a Primary Study

State level health information exchange initiative. Development workbook: A guide to key issues, options and strategies. Foundation of Research and Education of the American Health Information Management Association; 2007.

http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_038398.pdf Grey Lit.

Exclude - Not a Primary Study

States turning to electronic health records to save money, improve quality. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/programs/health/shn/2007/484a.htm> Grey Lit.

Exclude - Not MMIT

Study shows efficacy of bar-code usage. *Healthc Benchmarks Qual Improv* 2007;14(1):7-8.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Taming the information beast: States consider how to restrain costs, protect patients. National Conference of State Legislatures; 2007.

<http://www.ncsl.org/programs/health/shn/2007/sn496a.htm> Grey Lit.

Exclude - Not MMIT

The ideal barcode point of care system for the pharmacy informaticist. Healthcare Information and Management Systems Society; 2007.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67112> Grey Lit.

Exclude - Not a Primary Study

US must adopt and implement ICD-10-CM and ICD-10-PCS: Immediate action to upgrade medical code set standards needed. American Health Information Management Association; 2007. http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_034662.hcsp Grey Lit.

Exclude - Not a Primary Study

2008 National survey on patient flow challenges and technologies. Statcon; 2008.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68501> Grey Lit.
Exclude - Not MMIT

A consumer's guide to e-prescribing: Understanding the benefits of e-prescribing. How it works, and what you can do. Washington, DC: The Centre for Improving Medication Management; 2008.
http://www.thecimm.org/PDF/eHI_CIMM_Consumer_Guide_to_ePrescribing.pdf Grey Lit.
Exclude - Not a Primary Study

A guide for health care payers to improve the medication management process: Value creation, market feasibility, and implementation opportunities through technology innovations. The Center for Improving Medication Management; 2008 Oct 6.
http://www.ehealthinitiative.org/assets/Documents/eHI_CIMM_Guide_for_Payers_Final.pdf Grey Lit.
Exclude - Not a Primary Study

Accelerating progress: Using health information technology and electronic health information exchange to improve care. National Governors Association; 2008.
<http://www.nga.org/Files/pdf/0809EHEALTHREPORT.PDF> Grey Lit.
Exclude - Not a Primary Study

Authentication of consumers. Connecting for Health Common Framework; 2008.
http://library.ahima.org/xpedio/groups/public/documents/external/bok1_039087.pdf Grey Lit.
Exclude - Not a Primary Study

BLUES project list of refined metrics. 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_869061_0_0_18/Karen_Fox_IQHIT_Q3_Outcome_Measures%20or_BLUES_Project.pdf Grey Lit.
Exclude - Not a Primary Study

Clinical practice guidelines ensure consistent quality cancer care. ASCO News & Forum 2008;3(4):22-3.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010176644&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=3174&accno=2010176644>
EBSCO CINAHL.
Exclude - Not a Primary Study

Clinician usability study: Workflow and clinician satisfaction improvement for physician CPOE and nursing eMAR using Cerner Millennium, PowerChart and the Motion C5. 2008.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68609> Grey Lit.
Exclude - Not a Primary Study

Computerized Physician Order Entry (CPOE). SO: Journal of clinical pharmacology 2008;
<http://www.mrw.interscience.wiley.com/cochrane/clhta/articles/HTA-32008100294/frame.html> Database: Cochrane.
Exclude - Not MMIT

Consumers as network participants. Connecting for Health Common Framework; 2008.
http://library.ahima.org/xpedio/groups/public/documents/external/bok1_039076.pdf Grey Lit.
Exclude - Not a Primary Study

Decisionmaker brief: Computerized provider order entry (CPOE). AHRQ; 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_831190_0_0_18/08-0093_cpoe.pdf Grey Lit.

Exclude - Not a Primary Study

Decisionmaker brief: Bar-coded medication administration (BCMA). AHRQ; 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_831403_0_0_18/08-0085_bcma.pdf Grey Lit.

Exclude - Not a Primary Study

Defining key health information technology terms. The National Alliance for Health Information Technology; 2008. Grey Lit.

Exclude - Not a Primary Study

Dental recommendations for preventing complications in patients with chronic conditions: Health partners research foundation eDent study. 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_869058_0_0_18/James_Fricton_IQHIT_Q1_HP_Recommendations_for_Preventing_Complications_for_Chronic_Illnesses.pdf Grey Lit.

Exclude - Not a Primary Study

Drug & device news. Medical Malpractice Law & Strategy 2008;25(12):7
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010060816&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1256&accno=2010060816>
EBSCO CINAHL.

Exclude - Not a Primary Study

E-prescribing: Missives from the front. National Conference of State Legislatures; 2008.
<http://www.ncsl.org/programs/health/shn/2008/sn508c.htm> Grey Lit.

Exclude - Not MMIT

Efficiency of hospital information management computer based program in pharmacy service at inpatient ward. Thai Journal of Hospital Pharmacy 2008;18(3):278-89. Database: IPA.

Exclude - Unable to Retrieve Foreign

Electronic health records: A global perspective. Healthcare Information and Management Systems Society; 2008. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68372> Grey Lit.

Exclude - Not a Primary Study

Electronic prescribing: Becoming mainstream practice. A collaborative report from the ehealth initiative and the center for improving medication management. eHealth Initiative; 2008. <http://nyam.waldo.kohalibrary.com/cgi-bin/koha/opac-detail.pl?biblionumber=250993> saved as NYAM Centre for Improving MM.pdf

Grey Lit.

Exclude - Not a Primary Study

Electronic prescribing in ambulatory care: Taming a rapidly evolving market. American Medical Informatics Association; 2008.

<http://www.amia.org/pubs/proceedings/symposia/2001/D010001400.pdf> Grey Lit.

Exclude - Not MMIT

Electronic prescribing: Becoming mainstream practice. A collaborative report from the ehealth initiative and the center for improving medication management. The Centre for Improving Medication Management; 2008.

http://www.thecimm.org.libaccess.lib.mcmaster.ca/PDF/eHI_CIMM_ePrescribing_Report_6-10-08.pdf Grey Lit.

Exclude - Not a Primary Study

Employee/Staff pre-go-live expectations/perceptions clinical information systems survey (Dr. only). 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_835019_0_0_18/CIS%20Expectations%20-%20Experiences%20Survey%20-%20post%20go-live%20-%20Shortened%20for%20MD%20-%20oct%2006.doc Grey Lit.

Exclude - Not a Primary Study

Employee/Staff pre-go-live expectations/perceptions clinical information systems survey. 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_835114_0_0_18/CIS%20Expectations%20-%20Experiences%20Survey%20-%20pre-%20go-live%20-%20Long%20Version%20for%20all%20clinical%20staff.doc Grey Lit.

Exclude - Not a Primary Study

ESP-VAERS case identification and reporting protocol. 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_869021_0_0_18/Lazarus_ESP_VAERS_algorithm_v1_2007_11_05_1.pdf Grey Lit.

Exclude - Not a Primary Study

Evidence on the costs and benefits of health information technology. Congress of the United States, Congressional Budget Office; 2008. <http://www.cbo.gov/ftpdocs/91xx/doc9168/05-20-HealthIT.pdf> Grey Lit.

Exclude - Not a Primary Study

FDA bar code rule overview and implementation. 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_226702_0_0_18/FDA-BCMA-AHRQ2007.pdf Grey Lit.

Exclude - Not MMIT

Health care at the crossroads: Guiding principles for the development of the hospital of the future. The Joint Commission; 2008.

http://www.jointcommission.org/NR/rdonlyres/1C9A7079-7A29-4658-B80D-A7DF8771309B/0/Hospital_Future.pdf Grey Lit.

Exclude - Not a Primary Study

Health information technology survey for physicians, physician assistants and nurse practitioners. 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_835112_0_0_18/Provider%20Survey-%20Final%20Version%20sent%20out%20Fall%202006.doc Grey Lit.

Exclude - Not a Primary Study

Health information technology: 2007 and 2008 state legislation. National Conference of State Legislatures; 2008. http://www.ncsl.org/print/health/forum/HIT_Enacted.pdf Grey Lit.

Exclude - Not a Primary Study

Health IT: Support for effective use of electronic prescribing. AHRQ; 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_834454_0_0_18/08-PFS015_eRx.pdf Grey Lit.

Exclude - Not a Primary Study

Healthcare information and management systems society briefing February 26, 2008. U.S. Department of Health and Human Services; 2008.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67797> Grey Lit.

Exclude - Not a Primary Study

Heparin overdose scare in 14 babies at Texas hospital: could CPOE have prevented this tragic error? Healthc Benchmarks Qual Improv 2008;15(9):89-90.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010009888&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010009888

EBSCO CINAHL.

Exclude - Not a Primary Study

High-tech medical records. National Conference of State Legislatures; 2008.

http://www.ncsl.org/magazine/articles/2008/08SLJune08_MedicalRecords.htm Grey Lit.

Exclude - Not MMIT

Identifying, categorizing, and evaluating health care efficiency measures. AHRQ; 2008.

http://library.ahima.org/xpedio/groups/public/documents/government/bok1_038923.pdf Grey Lit.

Exclude - Not MMIT

In brief. Nurs Times 2008;104(48):6

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010140327&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=275&accno=2010140327 EBSCO

CINAHL.

Exclude - Not a Primary Study

Industry watch. HHS secretary supports ePrescribing incentives. Health Manag Technol 2008;29(9):8

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010034903&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010034903 EBSCO

CINAHL.

Exclude - Not a Primary Study

Interview questions prior to EHR implementation. AHRQ; 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_898126_0_0_18/Pohl_Qual_Interview_EHR.pdf Grey Lit.

Exclude - Not a Primary Study

IQ Health survey for clinicians. 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_878096_0_0_18/IQHealth_Clinicians.pdf Grey Lit.

Exclude - Not a Primary Study

Measures of impact for BLUES project. 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_869062_0_0_18/Karen_Fox_IQHIT_Q4_Measures_of_Impact_for_BLUES_Project.pdf Grey Lit.

Exclude - Not a Primary Study

Medicare's practical guide to the e-prescribing incentive program. U.S. Department of Health and Human Services; 2008. <http://www.cms.hhs.gov/partnerships/downloads/11399.pdf> Grey Lit.

Exclude - Not a Primary Study

Medicare program; Standards for e-prescribing under medicare part D and identification of backward compatible version of adopted standard for e-prescribing and the medicare prescription drug program (version 8.1). 2008.

<http://edocket.access.gpo.gov/libaccess.lib.mcmaster.ca/2008/pdf/08-1094.pdf> Grey Lit.

Exclude - Not a Primary Study

Medication adherence and compliance patient letter. 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_909117_0_0_18/James_Veline_IQHIT_Q6_Adherence_and_Compliance_Patient_Letter.pdf Grey Lit.

Exclude - Not a Primary Study

New computer network helps EDs to reduce redundant test orders: observers see significant savings, benefits in patient safety. *ED Manag* 2008;20(12):133-4.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010128228&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2083&accno=2010128228

EBSCO CINAHL.

Exclude - Not a Primary Study

New resource encourages physicians to embrace electronic prescribing. *Am Fam Physician* 2008;78(10):1138 Database: CINAHL.

Exclude - Not a Primary Study

New Technology. *Healthcare Purchasing News* 2008;32(12):44

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=48343155&site=ehost-live&scope=site> EBSCO-BusinessSourceComplete.

Exclude - Not a Primary Study

ONCHIT to Fund Assessment of Medical Identity Theft Problem. *hfm (Healthcare Financial Management)* 2008;62(7):12

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=33142219&site=ehost-live&scope=site> EBSCO-BusinessSourceComplete.

Exclude - Not a Primary Study

Partnerships advance e-Health records. *Information Management Journal* 2008;42(3):10 Database: BSC.

Exclude - Not MMIT

PCIP Evaluation patient survey. NYC Health; 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_887509_0_0_18/PCIP_Evaluation_Patient_Survey.pdf Grey Lit.

Exclude - Not a Primary Study

PCIP evaluation provider survey. AHRQ; 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/P...8_887510_0_0_18/PCIP_Evaluation_Provider_Survey.pdf Grey Lit.

Exclude - Not MMIT

PCIP Post-EHR implementation survey of providers. NYC Health; 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_887511_0_0_18/PCIP_Post-EHR_Implementation_Survey_Providers.pdf Grey Lit.

Exclude - Not a Primary Study

Privacy surveillance in healthcare. FairWarning, Inc; 2008.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68522> Grey Lit.

Exclude - Not a Primary Study

Provider experiences with and perceptions of current patients' use of email communication with their doctor. 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_835121_0_0_18/Provider%20experiences%20and%20perception%20survey%20PES%202_4_08.doc Grey Lit.

Exclude - Not a Primary Study

Physician HIT survey. Rhode Island Department of Health; 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_835119_0_0_18/Rhode%20Island%20Physician%20HIT%20Survey_061108.doc Grey Lit.

Exclude - Not a Primary Study

Quebec health record result chains. 2008.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_840769_0_0_18/Quebec%20Health%20Record%20Result%20Chains_v3.pdf Grey Lit.

Exclude - Not a Primary Study

Recommendations for standardized consumer consent policies and procedures for RHIOs in New York to advance interoperable health information exchange to improve care. New York Statewide Collaboration Process (SCP), New York Health Information Security and Privacy Collaboration (HISPC); 2008.

http://www.nyehealth.org/files/File_Repository16/pdf/Consent_White_Paper_20081125.pdf Grey Lit.

Exclude - Not a Primary Study

Regional news: South. Mod Healthc 2008;38(36):20
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010060050&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010060050> EBSCO CINAHL.

Exclude - Not a Primary Study

Reversing the trend of resistant infections: cutting costs with automated screening. Healthc Benchmarks Qual Improv 2008;15(11):115-7.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010085880&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010085880> EBSCO CINAHL.

Exclude - Not a Primary Study

Safely implementing health information and converging technologies. The Joint Commission; 2008.

http://www.jointcommission.org/SentinelEvents/SentinelEventAlert/sea_42.htm Grey Lit.

Exclude - Not a Primary Study

State level health information exchange. Final report part I: Roles in ensuring governance and advancing interoperability. Foundation of Research and Education of American Health Information Management Association; 2008.

http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_040348.pdf Grey Lit.

Exclude - Not MMIT

State level health information exchange. Final report part II: Coordinating policies that impact the access, use and control of health information. Foundation of Research and Education of American Health Information Management Association; 2008.

http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_040349.pdf Grey Lit.

Exclude - Not MMIT

Survey of health center use of electronic health information. NACHC; 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_835625_0_0_18/HIT%20Survey%2007_16_08_forNORC.doc Grey Lit.

Exclude - Not a Primary Study

The importance of enterprise integration: A work product of the HIMSS enterprise integration task force. Healthcare Information and Management Systems Society; 2008.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67643> Grey Lit.

Exclude - Not a Primary Study

The ONC-coordinated federal health IT strategic plan: 2008-2012. Department of Health & Human Services; 2008.

http://library.ahima.org/xpedio/groups/public/documents/government/bok1_038569.pdf Grey Lit.

Exclude - Not a Primary Study

The ONC-coordinated federal health IT strategic plan: 2008-2012. Department of Health and Human Services; 2008. Grey Lit.

Exclude - Not a Primary Study

The state of health information technology in California: Use among hospitals and long term care facilities. California HealthCare Foundation; 2008.

<http://www.chcf.org/topics/view.cfm?itemID=133656> Grey Lit.

Exclude - Not MMIT

The technology factor: is it our friend or our foe? Healthc Benchmarks Qual Improv 2008;15(12):130-1.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010099482&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010099482>

EBSCO CINAHL.

Exclude - Not a Primary Study

Tissue welding technology maximizes efficiency, cuts costs, improves outcomes. Healthcare Purchasing News 2008;32(4):62

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=31566938&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Top nine health industry issues in 2009: outside forces will disrupt the industry. Mod Healthc 2008;38(51):1-8.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010155635&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010155635 EBSCO CINAHL>.

Exclude - Not a Primary Study

Transcription turnaround time for common document types. Perspectives in Health Information Management, Summer 2008, 2008.

http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_039584.pdf Grey Lit.

Exclude - Not a Primary Study

Understanding the benefits of e-prescribing: How does it work? What can you do? The Center for Improving Medication Management; 2008.

http://www.ehealthinitiative.org/assets/Documents/One_Page_Consumer_Pamphlet_Final.pdf Grey Lit.

Exclude - Not a Primary Study

Understanding the benefits of e-prescribing: How does it work? What can you do? Washington, DC: The Centre for Improving Medication Management; 2008.

http://www.thecimm.org/PDF/One_Page_Consumer_Pamphlet.pdf Grey Lit.

Exclude - Not MMIT

1st ACM SIGMM International Workshop on Media Studies and Implementations that Help Improving Access to Disabled Users, MSIADU'09, Co-located with the 2009 ACM International Conf. Multimedia, MM'09. In Beijing, China: Association for Computing Machinery; 2009. p. ACM.Engineering Village Compendex and Inspec.

Exclude - Not a Primary Study

AHRQ's ambulatory safety and quality program: Health IT portfolio. AHRQ; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_872385_0_0_18/09-P001.pdf Grey Lit.

Exclude - Not a Primary Study

Assessment of body mass index screening of elementary school children - Florida, 2007-2008. MMWR: Morbidity & Mortality Weekly Report 2009;58(17):460-3.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010281489&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1581&accno=2010281489 EBSCO CINAHL>.

Exclude - Not a Primary Study

Bar-code-assisted medication administration shows inconsistent results. AACN Bold Voices 2009;1(5):11

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010458100&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=4140&accno=2010458100 EBSCO CINAHL.

Exclude - Not a Primary Study

Break down these barriers to medication safety. Healthc Benchmarks Qual Improv 2009;16(6):66-7. PMID:19472681 OVID MEDLINE.

Exclude - Not a Primary Study

Bulletin board. CPOE systems roll out slowly. JAHIMA2009;80(8):14

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010396443&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=205&accno=2010396443 EBSCO CINAHL.

Exclude - Not a Primary Study

Change in storage and administration helps hospital: use of prepared bar-coded labels improves nursing work flow, drives scan rates. Briefings on Patient Safety 2009;10(9):6-7.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010381061&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2276&accno=2010381061 EBSCO CINAHL.

Exclude - Not a Primary Study

Clinical clips. Nursing Spectrum -- DC, Maryland & Virginia Edition 2009;19(11):24

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010398330&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=3087&accno=2010398330 EBSCO CINAHL.

Exclude - Not a Primary Study

Clinical decision support (CDS) fact sheet. Healthcare Information and Management Systems Society; 2009. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=69332> Grey Lit.

Exclude - Not a Primary Study

Closing the loop in medication management: Why an integrated, closed-loop solution is vital for hospitals. Microsoft; 2009. Grey Lit.

Exclude - Not a Primary Study

Compliance with verbal orders standards poor. Healthc Benchmarks Qual Improv 2009;16(8):90-1. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Computerized prescribing in the ambulatory environment. eHealth Initiative; 2009.

<http://www.ehealthinitiative.org/initiatives/erx> Grey Lit.

Exclude - Not a Primary Study

Computerized support to avoid inappropriate prescribing to seniors. Brown University Geriatric Psychopharmacology Update 2009;13(11):4

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010457399&site=ehost-live> EBSCO CINAHL.

Exclude - Unable to Retrieve

Consumer engagement in developing electronic health information systems. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_888520_0_0_18/09-0081-EF.pdf Grey Lit.

Exclude - Not a Primary Study

E-prescribing saves costs and improves patient safety. CMA Today 2009;42(3):13
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010470643&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2589&accno=2010470643
EBSCO CINAHL.

Exclude - Not a Primary Study

Education may not be enough. ED Manag 2009;21(10):113-4.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010428168&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2083&accno=2010428168
EBSCO CINAHL.

Exclude - Not a Primary Study

Electronic prescribing: Toward maximum value and rapid adoption. eHealth Initiative; 2009.
<http://www.ehealthinitiative.org/assets/documents/eHIFullReport-ElectronicPrescribing2004.pdf> Grey Lit.

Exclude - Not MMIT

Electronic prescribing. H&HN: Hospitals & Health Networks 2009;83(5):46
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010313760&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1774&accno=2010313760
EBSCO CINAHL.

Exclude - Not a Primary Study

EMR white paper. 2009.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68426> Grey Lit.

Exclude - Not a Primary Study

EMRs: a 'plaintiff's dream'? Reduce your risks... emergency medical record. ED Legal Letter 2009;20(7):77
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010386681&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2160&accno=2010386681
EBSCO CINAHL.

Exclude - Not a Primary Study

Ensuring safer prescribing for children: AAP members funded to launch the STEPSTools project. 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_869063_0_0_18/Kevin%20Johnson_IQHIT_Q3_AAP-COCIT_Newsletter_Submission.pdf Grey Lit.

Exclude - Not a Primary Study

Errors associated with new technology: Proactively addressing potential medication safety issues. Joint Commission Perspectives on Patient Safety 2009;7(5):5-6. Database: CINAHL.
Exclude - No Outcomes of Interest

Health information exchange economic sustainability panel: Final report. Chicago: NORC; 2009. Grey Lit
Grey Lit.
Exclude - Not MMIT

Health plan, physicians collaborate to improve patient care. Case Management Advisor 2009;20(10):109-12.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010412936&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=570&accno=2010412936 EBSCO CINAHL.
Exclude - Not a Primary Study

Healthcare information technology standards home page. Healthcare Information Technology Standards; 2009. <http://www.hitsp.org/default.aspx> Grey Lit.
Exclude - No Outcomes of Interest

HIMSS 2009 informatics nurse impact survey. Healthcare Information and Management Systems Society; 2009. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=69103> Grey Lit.
Exclude - Not MMIT

HITSP library of standards documents and other materials. Healthcare Information Technology Standards; 2009. <http://www.hitsp.org/default.aspx?show=library> Grey Lit.
Exclude - Not MMIT

Impact of health IT on nurses' time spent on direct patient care. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_877556_0_0_18/Impact%20of%20Health%20IT%20on%20nurses.pdf Grey Lit.
Exclude - Not a Primary Study

In brief. Nurs Times 2009;105(6):7
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010241318&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=275&accno=2010241318 EBSCO CINAHL.
Exclude - Not a Primary Study

In case you haven't heard.. Alcoholism & Drug Abuse Weekly 2009;21(10):8
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010222895&site=ehost-live> EBSCO CINAHL.
Exclude - Not a Primary Study

Industry watch. Health Manag Technol 2009;30(1):6-7.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010174561&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010174561 EBSCO CINAHL.
Exclude - Not a Primary Study

Industry watch. Health Manag Technol 2009;30(5):8
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010316305&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010316305 EBSCO CINAHL.
Exclude - Not a Primary Study

International conference on eHealth, telemedicine, and social medicine. In Cancun, Mexico: Inst. of Elec. and Elec. Eng. Computer Society; 2009. 20091512030440
Database: Compendex.
Exclude - Not MMIT

Lab's HIE solution connects LIS to EMR and HIS. MLO Med Lab Obs 2009;41(12):30
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010512946&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=246&accno=2010512946 EBSCO CINAHL.
Exclude - Not a Primary Study

Lessons learned from a journey to EMR. Health Manag Technol 2009;30(11):24-7.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010470330&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010470330 EBSCO CINAHL.
Exclude - Not a Primary Study

Look what's happening @ HCI Online. Healthc Inform 2009;26(5):6
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010375190&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1239&accno=2010375190 EBSCO CINAHL.
Exclude - Not a Primary Study

Medication management and health IT in ambulatory care. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_891066_0_0_18/09-0098.pdf Grey Lit.
Exclude - Not a Primary Study

Medication turnaround time in the inpatient setting. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/P...0_1248_868887_0_0_18/Medication_Turnaround_Time.pdf Grey Lit.
Exclude - Not a Primary Study

Member profile. ACE member educates on copy/paste. AHIMA Advantage 2009;13(7):1
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010484018&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2010&accno=2010484018 EBSCO CINAHL.
Exclude - Not a Primary Study

National/international technology guide white paper. Healthcare Information and Management Systems Society; 2009.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=69152> Grey Lit.
Exclude - Not a Primary Study

News. CIN COMPUT INFORM NURS 2009;27(1):1-6.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010253850&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2306&accno=2010253850 EBSCO CINAHL.
Exclude - Not a Primary Study

News & advocacy. AORN Connections 2009;7(5):4-5.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010299377&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2592&accno=2010299377
EBSCO CINAHL.

Exclude - Not a Primary Study

News & advocacy [corrected] [published erratum appears in AORN CONNECT 2009 Nov-Dec;7(11):5]. AORN Connections 2009;7(10):4
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010438882&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2592&accno=2010438882
EBSCO CINAHL.

Exclude - Not a Primary Study

Patient safety Q&A. Briefings on Patient Safety 2009;10(12):12
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010495516&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2276&accno=2010495516
EBSCO CINAHL.

Exclude - Not a Primary Study

Patient satisfaction upon discharge improved: hospitalists, PCPs use communication software. Healthc Benchmarks Qual Improv 2009;16(10):113-4.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010400603&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010400603
EBSCO CINAHL.

Exclude - Not a Primary Study

Patient satisfaction upon discharge improved. Hosp Case Manag 2009;17(11):170
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010443643&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=842&accno=2010443643 EBSCO CINAHL.

Exclude - Not a Primary Study

Percentage of alerts or reminders that resulted in desired action. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_868888_0_0_18/Percent_of_Alerts.pdf Grey Lit.

Exclude - Not a Primary Study

Percentage of orders entered by authorized providers using CPOE. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_868889_0_0_18/Percent_of_Orders_Entered.pdf Grey Lit.

Exclude - Not a Primary Study

Percentage of verbal orders. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_868890_0_0_18/Percent_of_Verbal_Orders.pdf Grey Lit.

Exclude - Not a Primary Study

Prescribing patterns of cost-effective drugs. AHRQ; 2009.
http://healthit.ahrq.gov.libaccess.lib.mcmaster.ca/portal/server.pt/gateway/PTARGS_0_1248_868891_0_0_18/Prescribing_Patterns_of_Cost_Effective.pdf Grey Lit.

Exclude - Not a Primary Study

Prescribing patterns of preferred or formulary medications. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_868892_0_0_18/Prescribing_Patterns_of_Preferred.pdf Grey Lit.

Exclude - Not a Primary Study

Profiles of progress: State health IT initiatives. 2009.

<http://www.nascio.org/publications/documents/NASCIO-ProfilesInProgress3.pdf> Grey Lit.

Exclude - Not MMIT

Project HealthDesign: Rethinking the power and potential of personal health records. Round One Final Report. Madison, WI: Robert Wood Johnson Foundation; 2009.

<http://www.projecthealthdesign.org/media/file/Round%20One%20PHD%20Final%20Report6.17.09.pdf> Grey Lit.

Exclude - Not a Primary Study

Public governance models for a sustainable health information exchange industry. 2009.

<http://www.nga.org/Files/pdf/0902EHEALTHHIEREPORT.PDF> Grey Lit.

Exclude - Not a Primary Study

Reduction in hospital-acquired complications and infections. AHRQ; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_898613_0_0_18/09-0097.pdf Grey Lit.

Exclude - Not a Primary Study

Scribes, EMR please docs, save \$600,000. ED Manag 2009;21(10):117-8. PMID:20162997
OVID MEDLINE.

Exclude - Not a Primary Study

Seek out aids to risk prevention. ED Manag 2009;21(10):115-6.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010428178&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2083&accno=2010428178

EBSCO CINAHL.

Exclude - Not a Primary Study

SonoSite supports transplant charity. Operating Theatre Journal 2009;(230):16

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010499557&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2766&accno=2010499557

EBSCO CINAHL.

Exclude - Not a Primary Study

SonoSites MICROMAXX system is the right choice for regional nerve blocks. Operating Theatre Journal 2009;(223):8

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010258627&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2766&accno=2010258627

EBSCO CINAHL.

Exclude - Not a Primary Study

Standards on verbal orders rank high among common compliance problems. *ED Manag* 2009;1-2.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010310231&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2083&accno=2010310231
EBSCO CINAHL.

Exclude - Not a Primary Study

Study finds benefits in 'paperless' hospitals: 'patients appear safer and hospital bottom lines may improve'. *American Dental Association News* 2009;40(6):21

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010239079&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2360&accno=2010239079
EBSCO CINAHL.

Exclude - Not a Primary Study

Study says E-prescribing systems boost efficiency. *Same Day Surg* 2009;33(6):63-4.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010300618&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=508&accno=2010300618 EBSCO CINAHL.

Exclude - Not a Primary Study

Technological advances in adherence interventions are not expected panacea. Electronic messaging can't do it all. *AIDS Alert* 2009;24(12):136-7. PMID:20063474 OVID MEDLINE.

Exclude - Not a Primary Study

Technology, government incentives drive electronic prescribing. *Ocular Surgery News* 2009;27(24):22-3.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010509277&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2294&accno=2010509277
EBSCO CINAHL.

Exclude - Not a Primary Study

Technology: bar codes reduce errors... sometimes. *Nursing (Lond)* 2009;39(10):29

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010430992&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=280&accno=2010430992 EBSCO CINAHL.

Exclude - Not a Primary Study

The observatory. New tests. *MLO Med Lab Obs* 2009;41(4):8

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010262281&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=246&accno=2010262281 EBSCO CINAHL.

Exclude - Not a Primary Study

The week in nursing. *Nurs Times* 2009;105(46):6-7.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010493848&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=275&accno=2010493848 EBSCO CINAHL.

Exclude - Not a Primary Study

Universal patient floor increases patient flow, decreases handoff, improves patient safety. Briefings on Patient Safety 2009;10(10):5-6.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010451799&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2276&accno=2010451799
EBSCO CINAHL.

Exclude - Not a Primary Study

Welcome to CPOE.org. Oregon Health & Science University; 2009.

<http://www.ohsu.edu/academic/dmice/research/cpoe/index.php> Grey Lit.

Exclude - Not MMIT

2010 quality standards. Remington Report 2010;18(2):46

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010607676&site=ehost-live> EBSCO CINAHL.

Exclude - Not a Primary Study

An independent's experience with e-prescribing. America's Pharmacist 2010;127(7):10
Database: IPA.

Exclude - Not a Primary Study

Are you ready for a post-reform world? BioPharm International 2010;23(5):66 OVID
EMBASE.

Exclude - Not a Primary Study

Bulletin board. PQRI and e-prescribing changes for 2010... Physician Quality Reporting Initiative. JAHIMA2010;81(1):14-5.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010520184&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=205&accno=2010520184 EBSCO CINAHL.

Exclude - Not a Primary Study

By the numbers. Largest CPOE systems. Mod Healthc 2010;40(16):34

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010630523&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010630523 EBSCO CINAHL.

Exclude - Not a Primary Study

CE test. Professional Case Management 2010;15(2):68-9.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010621803&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=3340&accno=2010621803
EBSCO CINAHL.

Exclude - Not a Primary Study

Changes to the Physician Quality Reporting Initiative (PQRI) and the Electronic Prescribing Incentive Program. Remington Report 2010;18(2):33

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010607669&site=ehost-live> EBSCO CINAHL.

Exclude - Not a Primary Study

Clinical digest. Better drug adherence after transplant in children sent texts. Nurs Stand 2010;24(19):17

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010536220&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=530&accno=2010536220 EBSCO CINAHL.

Exclude - Not a Primary Study

Clinical news. Informatics. CVS Caremark announces e-prescribing agreement with Allscripts. J Pharm Technol 2010;26(2):89-90.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010636887&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=967&accno=2010636887 EBSCO CINAHL.

Exclude - Not a Primary Study

Clinical news. Substance abuse. Prescription monitoring programs reveal frequency of “doctor shopping”. J Pharm Technol 2010;26(2):91

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010636890&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=967&accno=2010636890 EBSCO CINAHL.

Exclude - Not a Primary Study

Clinical rounds. Communication: mixed reviews for electronic technology. Nursing (Lond) 2010;40(3):20-1.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010583781&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=280&accno=2010583781 EBSCO CINAHL.

Exclude - Not a Primary Study

CMS overview of e-prescribing. U.S. Department of Health & Human Services; 2010.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68716> Grey Lit.

Exclude - Not a Primary Study

Consider human factors engineering when designing your patient safety projects. Briefings on Patient Safety 2010;11(4):1-3.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010615826&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2276&accno=2010615826 EBSCO CINAHL.

Exclude - Not a Primary Study

Drug & device news. Medical Malpractice Law & Strategy 2010;27(7):7

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010627441&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1256&accno=2010627441 EBSCO CINAHL.

Exclude - Not a Primary Study

Federal government’s paperless requirements come with many Practical benefits. Caribbean Business 2010;38(20):B47

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=51302858&site=ehost-live&scope=site> EBSCO-BusinessSourceComplete.

Exclude - Not a Primary Study

Follow-up calls help patients adhere to treatment plan, avoid readmissions. Hosp Case Manag 2010;18(1):7-8.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010518297&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=842&accno=2010518297 EBSCO CINAHL.

Exclude - Not a Primary Study

FYI. MLA News 2010;50(4):4-5.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010646873&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1269&accno=2010646873 EBSCO CINAHL.

Exclude - Not a Primary Study

Health system sets 'zero errors' as its goal for patient safety, quality. Healthc Benchmarks Qual Improv 2010;17(4):37-41.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010621649&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010621649 EBSCO CINAHL.

Exclude - Not a Primary Study

Individuals and Employer Requirements CQ Weekly 2010;68(15):916

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=49546225&site=ehost-live&scope=site> EBSCO-BusinessSourceComplete.

Exclude - Not a Primary Study

Industry resources on e-prescribing. Healthcare Information and Management Systems Society; 2010. http://www.himss.org/ASP/topics_FocusDynamic.asp?faid=279 Grey Lit.

Exclude - Not MMIT

Kaiser/VA/DoD partnership piloting a nationwide EHR network: early testing encouraging; improvements in quality anticipated. Healthc Benchmarks Qual Improv 2010;17(2):13-5.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010540294&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010540294 EBSCO CINAHL.

Exclude - Not a Primary Study

Late news. Mod Healthc 2010;40(7):4

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010557907&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010557907 EBSCO CINAHL.

Exclude - Not a Primary Study

Med mal news. Medical Malpractice Law & Strategy 2010;27(4):5

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010570031&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1256&accno=2010570031 EBSCO CINAHL.

Exclude - Not a Primary Study

New products. Pharmacy practice. J Pharm Technol 2010;26(1):39
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010583124&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=967&accno=2010583124 EBSCO CINAHL.

Exclude - Not a Primary Study

Pioneers in healthcare IT: Wolters Kluwer Health. Increasing ROI from EMR/CPOE: trusted clinical content and advanced decision support drive outcomes. Health Manag Technol 2010;31(2):28

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010585790&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010585790 EBSCO CINAHL.

Exclude - Not a Primary Study

POE Publications. Oregon Health & Science University; 2010.

<http://www.ohsu.edu/academic/dmice/research/cpoe/publications.php> Grey Lit.

Exclude - Not MMIT

QI efforts lead to success in VTE prophylaxis. Healthc Benchmarks Qual Improv 2010;17(6):63-5.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010657663&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010657663 EBSCO CINAHL.

Exclude - Not a Primary Study

Standardize order sets for improved care. Health Manag Technol 2010;31(2):38-9.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010565553&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010565553 EBSCO CINAHL.

Exclude - Not a Primary Study

State e-health activities in 2007: Findings from a state survey. The Commonwealth Fund; 2010.

http://www.commonwealthfund.org/publications/publications_show.htm?doc_id=669309 Grey Lit.

Exclude - Not MMIT

The electronic healthcare network accreditation commission (EHNAC) homepage. 2010.

<http://www.ehnac.org/> Grey Lit.

Exclude - Not MMIT

VA strengthens critical patient-safety procedure. Health Manag Technol 2010;31(4):28-9.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010624789&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010624789 EBSCO CINAHL.

Exclude - Not a Primary Study

What works. To err is human: bar codes help streamline medication administration. *Healthcare Purchasing News* 2010;34(3):66
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010586602&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2373&accno=2010586602
 EBSCO CINAHL.
 Exclude - Not a Primary Study

Aarnio E, Raitoharju R. Patient's medication information and e-Health development in Finland: A case study of a Finnish primary care organization. In Reading, UK: Academic Publishing Limited; 2008. p.17-23. *Engineering Village Compendex and Inspec*.
 Exclude - No Outcomes of Interest

Aarts J, Doorewaard H, Berg M. Understanding implementation: The case of a computerized physician order entry system in a large Dutch university medical center. *J Am Med Inform Assoc* 2004;11(3):207-16. Database: CINAHL.
 Exclude - Not a Primary Study

Aarts J, Berg M. Same systems, different outcomes: Comparing the implementation of computerized physician order entry in two dutch hospitals. *Methods Inf Med* 2006;45:53-61.
 Exclude - No Outcomes of Interest

Aarts J, van der SH. CPOE, alerts and workflow: taking stock of ten years research at Erasmus MC. *Studies in Health Technology & Informatics* 2009;148:165-9. PMID:19745247
 OVID MEDLINE.
 Exclude - No Outcomes of Interest

Abarca J, Colon LR, Wang VS, et al. Evaluation of the performance of drug-drug interaction screening software in community and hospital pharmacies. *J Manag Care Pharm* 2006;12(5):383-9. Database: Ovid MEDLINE(R).
 Exclude - Not MMIT

Abbrecht P, O'Leary T, Behrendt D. Evaluation of a computer-assisted method for individualized anticoagulation: Retrospective and prospective studies with a pharmacodynamic model. *Clinical Pharmacology & Therapeutics* 1982;32(1):129-36.
 Exclude - Not a Primary Study

Abdo YM. Designing a patient care medication and recording system that uses bar code technology. *Comput Nurs* 1992;10(3):116-20. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Abernethy AP, Herndon JE, Wheeler JL, et al. Feasibility and acceptability to patients of a longitudinal system for evaluating cancer-related symptoms and quality of life: pilot study of an e/Tablet data-collection system in academic oncology. *Journal of Pain & Symptom Management* 2009;37(6):1027-38.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010309332&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=463&accno=2010309332 EBSCO CINAHL.
 Exclude - Not MMIT

Abraham JA, Stapinski C. Safe medication practices compared at a teaching hospital and a community hospital. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
 Exclude - Not a Primary Study

Abrams H, Carr D. The human factor: unexpected benefits of a CPOE and electronic medication management implementation at the University Health Network. HEALTHC Q 2005;8:Spec-8 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Abston,K.C. Using the electronic medical record to predict the pharmacological management of acute myocardial infarction The University of UtahEditor. 1999. Grey Lit.

Exclude - No Outcomes of Interest

Acharyulu G. RFID in the healthcare supply chain: Improving performance through greater visibility. ICFAI Journal of Management Research 2007;6(11):32-45. Database: BSC.

Exclude - Not a Primary Study

Adams R, Ruffin R, Smith B, et al. Problems and some solutions in adapting clinical practice guidelines for asthma patient management into a computerised management system. The Western region asthma pilot project (Wrapp). Informatics in Healthcare Australia 1998;7(1):16-21. Database: CINAHL.

Exclude - No Outcomes of Interest

Adamson R, Principe P, Shah M. Implementation of a computerized physician order entry (CPOE)-based protocol to improve the care of patients with community-acquired pneumonia. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.

Exclude - Not a Primary Study

Adenot I. The pharmaceutical record. Ann Pharm Fr 2007;65(5):325-30. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Adhikari N, Lapinsky SE. Medical informatics in the intensive care unit: overview of technology assessment. J Crit Care 2003;18(1):41-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Adler KG. E-prescribing: why the fuss?[see comment. Fam Pract Manag 2009;16(1):22-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Adler K. How to successfully navigate your EHR implementation: These clues can help you avoid the pitfalls you'll encounter on your EHR journey. Fam Pract Manag 2007;14(2):33-9. <http://www.aafp.org/fpm/20070200/33howt.html> Grey Lit.

Exclude - Not a Primary Study

Adwan KM, Ragonese D. Implementation of computerized chemotherapy provider order entry in veterans affairs medical center and its role in a three tier check system. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Adwan KM, Ragonese D. Implementation of computerized chemotherapy provider order in veterans affairs medical center and its role in the three tier check system. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Agarwal R and Khuntia J. Personal health information management and the design of consumer health information technology: Background report. AHRQ; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_907458_0_0_18/09-0075-EF.pdf Grey Lit.

Exclude - Not a Primary Study

Agno W, Turpie G. A randomized comparison of a computer-based dosing program with a manual system to monitor oral anticoagulant therapy. *Thromb Res* 1998;91:237-40. Exclude - Not MMIT

Agno W, Johnson J, Nowacki B, et al. A computer generated induction system for hospitalized patients starting on oral anticoagulant therapy. *Thrombosis & Haemostasis* 2000;83(6):849-52. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Agha Z, Schapira RM, Maker AH. Cost effectiveness of telemedicine for the delivery of outpatient pulmonary care to a rural population. *Telemedicine Journal and e-Health* 2002;8(3):281-91. Grey Lit.

Exclude - Not MMIT

Agrawal A, Glasser AR. Barcode medication. Administration implementation in an acute care hospital and lessons learned. *J Healthc Inf Manag* 2009;23(4):24-9. PMID:19894483 OVID MEDLINE.

Exclude - No Outcomes of Interest

Agrawal A. Medication errors: Prevention using information technology systems. *Br J Clin Pharmacol* 2009;67(6): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Ague T. A dynamic duo: Barcoding and RFID prevent errors. *Healthcare Information and Management Systems Society*; 2007.

<http://www.himss.org/asp/ContentRedirector.asp?ContentID=66795> Grey Lit.

Exclude - Not a Primary Study

Aguila A, Valenzuela P. Experience with electronic files in a university neonatology unit. *Rev Med Chil* 2005;133(2):241-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ahlfeldt H, Johansson B, Linnarsson R, et al. Experiences from the use of data-driven decision support in different environments. *Computers in Biology & Medicine* 1994;24(5):397-404. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ahmad A, Rucker D, Teater P. Implementing end-to-end computerized physician order entry. *Healthcare Information and Management Systems Society*; 2002.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=12284> Grey Lit.

Exclude - Not a Primary Study

Ain KB, Pucino F, Drass JA, et al. Comparison of a restrictive and nonrestrictive levothyroxine formulary system. *ASHP Annual Meeting* 1992;49: Database: IPA.

Exclude - Not a Primary Study

Akiyama M, Kondo T. Risk management and measuring productivity with POAS--point of act system. *Studies in Health Technology & Informatics* 2007;129(Pt:1):1-12. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Akl EA. A Decision Aid for COPD patients considering inhaled steroid therapy: Development and before and after pilot testing. *BMC Med Inform Decis Mak* 2007;7: Database: Embase Sept 22-09.

Exclude - Not MMIT

Al sheikh YT, Haug PJ, Wong A, et al. Using a diabetes data mart in individualizing diabetes management. *AMIA 2008;Annual:Symposium* PMID:18999166 OVID MEDLINE.

Exclude - No Outcomes of Interest

Alamo J M R, Wong J, Babbitt R and others. MISS: Medicine information support system in the smart home environment. In Berlin, Germany: Springer-Verlag; 2008. p.185-99.10159677

Database: Inspec.

Exclude - Not a Primary Study

Alamo JMR, Hen I, Babbitt R, et al. Support for Medication Safety and Compliance in Smart Home Environments. *International Journal of Advanced Pervasive and Ubiquitous Computing* 2009;1(3):42-60. Engineering Village Compendex and Inspec.

Exclude - Unable to Retrieve

Alamo J M R, Wong J, Babbitt R and others. Using web services for medication management in a smart home environment. In 2007; Tours, France) SUBJECT(S) Identifier Ubiquitous computing; Home automation; Self-help devices for people with disabilities; Medical telematics; Smart homes; Health telematics; ICOST Note(s) Includes bibliographical references and index.: 2009. p.265-8.Grey Lit.

Exclude - Not a Primary Study

Alan EL. Error potential of automated systems. *ASHP Annual Meeting* 1995;52: Database: IPA.

Exclude - Not a Primary Study

Alanon Pardo AM, Salazar BM, Calleja Hernandez MA. Dose adjustment of antibiotics in patients with renal failure. [Spanish]. *Atencion Farmaceutica* 2010;12(1):33-8. OVID EMBASE.

Exclude - Unable to Retrieve

Alapetite A. Speech recognition for the anaesthesia record during crisis scenarios. *Int J Med Inf* 2008;77(7):448-60. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Alasaarela E, Oliver NS. Wireless solutions for managing diabetes: A review and future prospects. [Review] [59 refs]. *Technology & Health Care* 2009;17(5):353-67. PMID:20051615 OVID MEDLINE.

Exclude - No Outcomes of Interest

Albers N. [Computerized prescription service for pediatric intensive care]. [German]. Monatsschrift Kinderheilkunde 1993;Organ der Deutschen Gesellschaft für Kinderheilkunde. 141(10):814-7. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Albisser AM. Six generations of the insulin dosage computer: a new clinical device for diabetes self-management through specialized centres. Hormone & Metabolic Research - Supplement 1990;24:140-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Albisser AM, Harris RI, Sakkal S, et al. Diabetes intervention in the information age.[erratum appears in Med Inf (Lond) 1997 Apr-Jun;22(2):205]. Med Inform (Lond) 1996;21(4):297-316. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Albisser AM. Clinical studies with home glucose clamping. Ann Endocrinol (Paris) 2001;62(1:Pt:1):t-8 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Albisser AM, Harris R, I, Albisser JB, et al. The impact of initiatives in education, self-management training, and computer-assisted self-care on outcomes in diabetes disease management. Diabetes Technology and Therapeutics 2001;3(4):571-9. Grey Lit.

Exclude - Not MMIT

Albisser A, Wright C, Sakkal S. Averting iatrogenic hypoglycemia through glucose prediction in clinical practice: Progress towards a new procedure in diabetes. Diabetes Res Clin Pract 2007;76:207-14. Exclude - No Outcomes of Interest

Aldrich A, Nelson M. Integration of a pharmacy based anticoagulation parameter into a medical group quality scorecard. Ashp Summer Meeting 2009;65: Database: IPA.

Exclude - Not a Primary Study

Alexander GL, Wakefield DS. Information technology sophistication in nursing homes. J AM MED DIR ASSOC 2009;10(6):398-407.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010418002&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2207&accno=2010418002
EBSCO CINAHL.

Exclude - Not MMIT

Ali M, Vaccine Safety DG, Do CG, et al. The use of a computerized database to monitor vaccine safety in Vietn. Bull World Health Organ 2005;83(8):604-10. Database: IPA.

Exclude - Not MMIT

Ali NA. Implementing CPOE in a medical intensive care unit: A physician's experience. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Allain H. Drug distribution system in hospitals. Therapie 2002;57(4): Database: Embase Sept 22-09.

Exclude - Not MMIT

Allan EL, Barker KN. Review of research on automated systems and medication errors. Ashp Midyear Clinical Meeting 1995;30: Database: IPA.

Exclude - Not a Primary Study

Allen C, Manyika P, Jazayeri D, et al. Rapid deployment of electronic medical records for ARV rollout in rural Rwanda. AMIA 2006;840 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Allen J, Ferguson G, Blaylock N, et al. Chester: towards a personal medication advisor. Journal of Biomedical Informatics 2006;39(5):500-13. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Allen SI. Prescription-writing with a PC. Comput Methods Programs Biomed 1986;22(1):127-35. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Allenet B, Schmitt D, Leger S, et al. Acceptability of computerized physician's drug prescription in Grenoble Teaching Hospital, France: Survey of 44 prescribers. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.

Exclude - Not a Primary Study

Allig J. Electronic transmission of drug-ordering from wards to clinical pharmacies by intra- or internet. Krankenhauspharmazie 2004;25(6): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Almeida-Hunt M, Coll RE, Kinney SD, et al. Improving medication safety in a community hospital: Identifying medication errors/near misses and developing policies for high-risk medications. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.

Exclude - Not a Primary Study

Almond M, Gordon K, Kent JR, et al. The effect of the controlled entry of electronic prescribing and medicines administration on the quality of prescribing, safety and success of administration on an acute medical ward. BJHC & IM 2002;19(2):41-2. Database: CINAHL.

Exclude - No Outcomes of Interest

Alonso Lopez FA, Iturrioz A, I, Molina IA, et al. [An analysis of long-term computerized prescriptions for those over and under 65 at a health center]. [Spanish]. Aten Primaria 1996;17(9):555-8. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Altersberger IE, Chan SY. Carbamazepine interactions. ASHP Annual Meeting 1996;53: Database: IPA.

Exclude - Not a Primary Study

Alvarez RL, Martin Conde JA, Alberdi LA, et al. Assessing an automated dispensation system in the emergency department of a level III-hospital. Farmacia Hospitalaria 2003;27(2):72-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Amarasingham R. Clinical information technologies and inpatient outcomes a multiple hospital study. Arch Intern Med 2009;169(2):108-14. Database: Embase Sept 22-09.

Exclude - Not MMIT

Amatayakul M, Cohen M. First communication, then automation. *Healthc Financ Manage* 2004;58(5):102-4. Database: BSC.

Exclude - Not a Primary Study

Ambrosiadou BV, Goulis DG, Pappas C. Clinical evaluation of the DIABETES expert system for decision support by multiple regimen insulin dose adjustment. *Computer Methods & Programs in Biomedicine* 1996;49(1):105-15. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

American Academy of Pediatrics Council on Clinical Information Technology, Gerstle RS. Electronic prescribing systems in pediatrics: the rationale and functionality requirements. *Pediatrics* 2007;119(6):1229-31. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

American Hospital Association, American Society of Health-System Pharmacists, Hospitals & Health Networks. Medication safety issue brief. Using automation to reduce errors. Part 2. *Hospitals & Health Networks* 2001;75(2):33-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

American Hospital Association, American Society of Health-System Pharmacists, Hospitals and Health Networks. Medication Safety Issue Brief. Bar Code implementation strategies. *Hospitals & Health Networks* 2005;79(7):65-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Amin AN, Deitelzweig SB. Optimizing the prevention of venous thromboembolism: recent quality initiatives and strategies to drive improvement. *Jt Comm J Qual Patient Saf* 2009;35(11):558-64. PMID:19947332 OVID MEDLINE.

Exclude - Not Primary Study

Ammenwerth E, Talmon J, Ash J, et al. Impact of CPOE on mortality rates - Contradictory findings, important messages. *Methods Inf Med* 2006;45:586-94. Exclude - Not a Primary Study

Ammenwerth E. The Effect of Electronic Prescribing on Medication Errors and Adverse Drug Events: A Systematic Review. *J Am Med Inform Assoc* 2008;15(5):585-600. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Ammenwerth E, Schnell-Inderst P, Siebert U. Vision and challenges of Evidence-Based Health Informatics: A case study of a CPOE meta-analysis. *Int J Med Inf* 2010;79(4):e83-e88 OVID EMBASE.

Exclude - No Outcomes of Interest

Ampt A, Westbrook JI. Measuring nurses' time in medication related tasks prior to the implementation of an electronic medication management system. *Studies in Health Technology & Informatics* 2007;130:157-67. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Amsden DB. Rules-based clinical alerting systems: A pharmacy opportunity. *Ashp Midyear Clinical Meeting* 2002;37. Database: IPA.

Exclude - Not a Primary Study

Andersen LS, Kjeldsen LJ, Haugbolle LS. Suitability of the Personal Electronic Medication profile for estimation of medication compliance. *Ugeskr Laeger* 2009;171(11):899-903. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Anderson HJ. Avoiding 'alert fatigue'. *Health Data Manag* 2009;17(10):42 PMID:19845096 OVID MEDLINE.

Exclude - Not a Primary Study

Anderson JG. Computer-based ambulatory information systems: recent developments. *J Ambulatory Care Manage* 2000;23(2):53-63. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Anderson JG, Jay SJ, Anderson M, et al. Evaluating the capability of information technology to prevent adverse drug events: a computer simulation approach. *J Am Med Inform Assoc* 2002;9(5):479-90. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Anderson JG. Evaluation in health informatics: computer simulation. *Computers in Biology & Medicine* 2002;32(3):151-64. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Anderson JG. A framework for considering business models. *Studies in Health Technology & Informatics* 2003;92:3-11. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Anderson JG. Information technology for detecting medication errors and adverse drug events. *Expert Opinion on Drug Safety* 2004;3(5):449-55. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Anderson JG, Ramanujam R, Hensel D, et al. The need for organizational change in patient safety initiatives. *Int J Med Inf* 2006;75(12):809-17. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Anderson JG. Social, ethical and legal barriers to e-health. *Int J Med Inf* 2007;76(5-6):480-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Anderson J, Jay S, Perry J, et al. Informal communication networks and change in physicians' practice behavior. *International Sociological Association* 1990;27:127-32.

Database: Sociological Abstracts.

Exclude - Not MMIT

Anderson KJ, Malone PM. Electronic prescriptions in pharmacy. *Am J Health Syst Pharm* 1999;56(Jul 1):1351-3. Database: IPA.

Exclude - Not a Primary Study

Anderson M, Baker M, Bell R, et al. The business case for patient safety. *HEALTHC Q* 2004;10(Spec):20-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Anderson SA, Jensen RJ. Multidisciplinary approach to decreasing the occurrence of medication errors and variances. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.
Exclude - Not a Primary Study

Andrade AS, McGruder HF, Wu AW, et al. A programmable prompting device improves adherence to highly active antiretroviral therapy in HIV-infected subjects with memory impairment. *Clin Infect Dis* 2005;41(6):875-82. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Andrade SE, Kahler KH, Frech F, et al. Methods for evaluation of medication adherence and persistence using automated databases. *Pharmacoepidemiology & Drug Safety* 2006;15(8):565-74. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Andreoni M, Beretta D, Carzaniga A, et al. A web-based family health record supporting genealogical clinical anamnesis. *AMIA* 2007;862 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Androulidakis A, Nielsen AD, Prentza A, et al. A distributed environment for the integration of multiple high-performance decision support systems into clinical workflow. *Technology & Health Care* 2006;14(3):157-70. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Angelo LB, Christensen DB, Ferreri SP. Does workflow automation make a difference in patient counseling? *Computertalk for the Pharmacist* 2002;22(5):16-7. Database: IPA.
Exclude - Not a Primary Study

Angelo LB, Christensen DB, Ferreri SP. Impact of community pharmacy automation on workflow, workload, and patient interaction. *Journal of the American Pharmacists Association: JAPhA* 2005;45(2):138-44. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Anhoj J, Nielsen L. Quantitative and qualitative usage data of an internet-based asthma monitoring tool. *J Med Internet Res* 2004;6(3):e23 Database: PsycINFO.
Exclude - Not MMIT

Anhoj J, Moldrup C. Feasibility of collecting diary data from asthma patients through mobile phones and SMS (short message service): response rate analysis and focus group evaluation from a pilot study. *J Med Internet Res* 2004;6(4):e42 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Anogianakis G, Goulis D, Vakalis D. Computer-aided prescription--a prototype system. *Studies in Health Technology & Informatics* 1997;43 Pt A:272-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Ansari M, Shlipak MG, Heidenreich PA, et al. Improving guideline adherence: a randomized trial evaluating strategies to increase beta-blocker use in heart failure. *Circulation* 2003;107(22):2799-804. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Anton BB, Schafer JJ, Micenko A, et al. Clinical decision support. How CDS tools impact patient care outcomes. *J Healthc Inf Manag* 2009;23(1):39-45. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Apostolopoulos A. Application of health informatics in the education of diabetic patients for the improvement of self-management and reporting to specialists. *Journal on Information Technology in Healthcare* 2008;5(6): Database: Embase Sept 22-09.

Exclude - Not MMIT

Appleby DH, Cullison TE, Edgerton WL. Reducing lost drug charges by monitoring computer generated interdepartmental transfers. *Am J Hosp Pharm* 1983;40(Jun):982-3. Database: IPA.

Exclude - Not MMIT

Arensman W, Price N, Pickens K. Automated interpretation of medical prescription text. *AMIA Annual Symposium* 2008;865 PMID:18998831 OVID MEDLINE.

Exclude - No Outcomes of Interest

Arlia EA, Mahoney CD, Berard CM, et al. Evaluation of the effect of computerized physician order entry (CPOE) on actual and prevented medication errors: lessons learned. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Arlicot N, Pourrat X, Bourgoin-Herard H, et al. Are antibiotic drugs well prescribed in case of renal insufficiency? A retrospective study. *Ren Fail* 2007;29(8):1055-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Armijo D, McDonnell C, Werner K. Electronic health record usability: Evaluation and use case framework. *AHRQ*; 2009.
[http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_907504_0_0_18/09\(10\)-0091-1-EF.pdf](http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_907504_0_0_18/09(10)-0091-1-EF.pdf) Grey Lit.

Exclude - Not a Primary Study

Armijo D, McDonnell C, Werner K. Electronic health record usability: Interface design considerations. *AHRQ*; 2009.
[http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_907505_0_0_18/09\(10\)-0091-2-EF.pdf](http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_907505_0_0_18/09(10)-0091-2-EF.pdf) Grey Lit.

Exclude - Not a Primary Study

Armour CL, Smith L, Krass I. Community pharmacy, disease state management, and adherence to medication: A review. *Disease Management & Health Outcomes* 2008;16(4):245-54. Database: CINAHL.

Exclude - Not MMIT

Armstrong I, Cox MA. Horus meets Nightingale in the modern age: How nursing communicates with pharmacy in HCIT era. *Studies in Health Technology & Informatics* 2006;122:585-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Armstrong K S, Davis J P, Bonnell R D and others. A model and prototype for using intelligent software agents to monitor patient adherence to a medication regimen. In Philadelphia; PA: 1998. p. 82. Grey Lit.

Exclude - No Outcomes of Interest

Arnow P, Chamberlin W. Debugging computer-assisted antibiotic prescribing. *Lancet* 1995;345(8944):207-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Aronsky D, Haug PJ. An integrated decision support system for diagnosing and managing patients with community-acquired pneumonia. *Proceedings / AMIA 1999;Annual Symposium.*:197-201. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Arotaritei D, Toma CM, Turnea M, et al. MIDAS intelligent platform for medical services, support for decision optimization in virtual medical communities. *Rev Med Chir Soc Med Nat Iasi* 2008;112(2):538-41. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Artinian NT, Harden JK, Kronenberg MW, et al. Pilot study of a Web-based compliance monitoring device for patients with congestive heart failure. *Heart & Lung* 2003;32(4):226-33. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Asakura S, Yamashita K, Takano Y, et al. Computerized checking system for drug interactions developed in Oita Medical University Hospital. *Japanese Journal of Hospital Pharmacy* 1997;23(5):467-77. Database: IPA.

Exclude - Unable to Retrieve Foreign

Asare AL, Huda H, Klimczak JC, et al. Integrating molecular diagnostic and flow cytometric reporting for improved longitudinal monitoring of HIV patients. *Proceedings / AMIA 1998;Annual Symposium.*:952-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Asaro PV, Boxerman SB. Effects of computerized provider order entry and nursing documentation on workflow. *Acad Emerg Med* 2008;15(10):908-15.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010386514&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2265&accno=2010386514 EBSCO CINAHL.

Exclude - Not MMIT

Ash JS, Gorman PN, Seshadri V, et al. Computerized physician order entry in U.S. hospitals: results of a 2002 survey. *J Am Med Inform Assoc* 2004;11(2):95-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ash JS. The unintended consequences of computerized provider order entry: Findings from a mixed methods exploration. *Int J Med Inf* 2009;78(Suppl 1):S69-76. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Ash J, Fournier L, Stavri P, et al. Principles for a successful computerized physician order entry implementation. *Proceedings of the AMIA Symposium 2003*;36-40. Exclude - Not a Primary Study

Ash J, Guappone K. Qualitative evaluation of health information exchange efforts. *Journal of Biomedical Informatics* 2007;40(6 Suppl 1):S33-S39

http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PJ6GKF-2&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=9&_fmt=full&_orig=browse&_srch=doc-

[info\(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume\)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=acc3d191ff4e6cb10597cfe86f226552](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PJ6GKF-2&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=9&_fmt=full&_orig=browse&_srch=doc-info(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=acc3d191ff4e6cb10597cfe86f226552) Grey Lit.

Exclude - Not a Primary Study

Aspinall S, Sevick MA, Donohue J, et al. Medication errors in older adults: a review of recent publications. *American Journal Geriatric Pharmacotherapy* 2007;5(1):75-84.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Assimakopoulos N. A medication management system. *Applied Mathematics and Computation* 1987;21(1):73-91. 2870781

Database: Inspec.

Exclude - No Outcomes of Interest

Astheimer H, Gortitz I, Bachert A. 10 years of computer-aided therapy planning in a department of pediatric hematology and oncology. *Klin Padiatr* 2003;215(6):345-51.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Atkins PM, Barone LD. Case study on measuring and improving the medication use system. *Jt Comm J Qual Improv* 1999;25(Jan):40-9. Database: IPA.

Exclude - Not MMIT

Atkinson V, Andrews JD. Effect of computer generated prompts on physician prescribing of multiple daily doses. *Can J Hosp Pharm* 1987;40(Jun):91-3. Database: IPA.

Exclude - Not MMIT

Atkinson WL, Frey D. Integration of a medication management model into outcome-based quality improvement: A pilot program in a rural proprietary home Healthcare agency. *Home Health Care Serv Q* 2005;24(1/2):29-45. Database: IPA.

Exclude - Not MMIT

Atreja A, Buck M, Jain A, et al. Drug-age alerting for outpatient geriatric prescriptions: a joint study using interoperable drug standards. *AMIA* 2005;886 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Audet AM, Doty MM, Peugh J, et al. Information technologies: when will they make it into physicians' black bags? *Medgenmed [Computer File]: Medscape General Medicine* 2004;6(4):2 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Austin G, Klasko S, Leaver WB. The art of health IT transformation. *Mod Healthc* 2009;5-16.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010470399&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010470399 EBSCO CINAHL.
Exclude - Not a Primary Study

Autenrieth-Ansorge L. Use of electronic data processing (EDP) in the pharmacy for communication. *Krankenhauspharmazie* 1993;14(Jan):29-31. Database: IPA.
Exclude - Unable to Retrieve

Avery AJ. Safer medicines management in primary care. *Br J Gen Pract* 2002;52(Suppl):S17-S22 Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Avery AJ, Savelyich BS, Sheikh A, et al. Improving general practice computer systems for patient safety: qualitative study of key stakeholders. *Quality & Safety in Health Care* 2007;16(1):28-33. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Avorn J. Drug use in academic medical centres in an era of ferment. *Pharmacoeconomics* 1996;10(Suppl 2):124-9. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Avorn J. Balancing the cost and value of medications: the dilemma facing clinicians. *Pharmacoeconomics* 2002;20(Suppl 3):67-72. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Aydin CE. Professional agendas and computer-based patient records: negotiating for control. *Top Health Inf Manage* 1994;15(1):41-51. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Aylett M. Computerised repeat prescriptions: simple system. *British Medical Journal Clinical Research Ed* 1985;290(6475):1115-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Ayshford H. The need to know [pharma manufacturers]. *Manufacturing Chemist* 2009;80(4):24-5. Engineering Village Compendex and Inspec.
Exclude - Not a Primary Study

Azagra LR, Aguye BA. Computer-aided prescribing in Spain. *Aten Primaria* 2005;35(9):457-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Aziz S, Crabtree D, Meollentin D, et al. The role of clinical informatics technician in improving clinical pharmacy program. *Ashp Summer Meeting* 2003;60: Database: IPA.
Exclude - Not a Primary Study

Babamoto EN, Uchida KM, Eames JL, et al. Tools to evaluate and implement the use of technology to decrease medication errors. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Badham J, Seah LP, Kidd M, et al. National electronic medication management systems. *Studies in Health Technology & Informatics* 2001;84(2):1524 Grey Lit.
Exclude - Not a Primary Study

Baer DM, Kotschi ML. On-line computer pharmacokinetics program: lessons learned from its failure. *Clin Chem* 1995;41(4):491-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Bagheri H, Michel F, Lapeyre-Mestre M, et al. Detection and incidence of drug-induced liver injuries in hospital: a prospective analysis from laboratory signals. *Br J Clin Pharmacol* 2000;50(5):479-84. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bailey FA. Palliative care intervention for choice and use of opioids in the last hours of life. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences* 2008;63(9):949-50. Database: Embase Sept 22-09.
Exclude - Not MMIT

Bailey M, Ferro K. Innovative drug formulary management through computer-assisted protocols. *J Manag Care Pharm* 1998;4(May-Jun):246-7. Database: IPA.
Exclude - Not MMIT

Bailey TC, Troy MS. Using information systems technology to improve antibiotic prescribing. *Crit Care Med* 2001;29(4:Suppl):N87-91. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Bailey TC, Noirot LA, Gage BF, et al. Improving adherence to coronary heart disease secondary prevention medication guidelines at a community hospital. *AMIA* 2006;850 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bails D, Clayton K, Roy K, et al. Implementing online medication reconciliation at a large academic medical center. *Jt Comm J Qual Patient Saf* 2008;34(9):499-508. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Baird TK, Broekemeier RL, Anderson MW. Effectiveness of a computer-supported refill reminder system. *Am J Hosp Pharm* 1984;41(11):2395-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bajaj AN, Karandikar SM, Baichwal MR. Computer--aided drug interaction retrieval programme. *J Assoc Physicians India* 1994;42(9):706-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Baker GR, Norton P. Making patients safer! Reducing error in Canadian healthcare. *Healthcarepapers* 2001;2(1):10-31. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Baker V, Copping M. A pilot study exploring the clinical benefits when using a Mobile Clinical Assistant, the Motion C5 in medical wards. *Studies in Health Technology & Informatics* 2009;146:121-5. PMID:19592820 OVID MEDLINE.
Exclude - No Outcomes of Interest

Bakken K. Local monitoring center for clozapine therapy: Quality assurance of drug treatment in a group of psychiatric patients. *Tidsskr Nor Laegeforen* 1998;118(7): Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Balaguer Santamaria JA, Fernandez Ballart JD, Escribano SJ. Usefulness of a software package to reduce medication errors in neonatal care. *An Esp Pediatr* 2001;55(6):541-5. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Balas EA, Jaffrey F, Kuperman GJ, et al. Electronic communication with patients: Evaluation of distance medicine technology. *JAMA* 1997;278(2):152-9. Grey Lit.

Exclude - Not MMIT

Balas EA, Krishna S, Kretschmer RA, et al. Computerized knowledge management in diabetes care. *Med Care* 2004;42(6):610-21. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Balfour DC, III, Evans S, Januska J, et al. Health information technology--results from a roundtable discussion. *J Manag Care Pharm* 2009;15(1:Suppl:A):S10-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Balka E. Technology, governance and patient safety: Systems issues in technology and patient safety. *Int J Med Inf* 2007;76(Suppl 1):S35-47. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Balka E. Who is in charge of patient safety? Work practice, work processes and utopian views of automatic drug dispensing systems. *Int J Med Inf* 2007;76(Suppl 1):S48-57. Database: Embase Sept 22-09.

Exclude - Not MMIT

Balkrishnan R, Foss CE, Pawaskar M, et al. Monitoring for medication errors in outpatient settings. *Journal of Dermatological Treatment* 2009;20(4):229-32. PMID:19085201 OVID MEDLINE.

Exclude - No Outcomes of Interest

Ball FR. DEA issues proposed rule on e-prescribing. *J Med Pract Manage* 2009;24(4):243 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ball MJ. Enabling technologies promise to revitalize the role of nursing in an era of patient safety. *Int J Med Inf* 2002;69(1):29-38. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Ball MJ, Douglas JV. IT, patient safety, and quality care. *J Healthc Inf Manag* 2002;16(1):28-33. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ball MJ, Douglas JV. Redefining and improving patient safety. *Methods Inf Med* 2002;41(4):271-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ball MJ, Garets DE, Handler TJ. Leveraging information technology towards enhancing patient care and a culture of safety in the US. *Methods Inf Med* 2003;42(5):503-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ballentine AJ, Kinnaird D, Wilson JP. Prescription errors occur despite computerized prescriber order entry. *Am J Health Syst Pharm* 2003;60(7):708-9. Database: IPA.

Exclude - Not a Primary Study

Balon D. Design of a computer program for automatic capture of adverse drug interaction and contraindication data detected during prescription labelling. *Int J Pharm Pract* 1997;5(2): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Balust J. Can anesthesia information management systems improve quality in the surgical suite? *Current Opinion in Anaesthesiology* 2009;22(2):215-22. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Bambauer KZ, Adams AS, Zhang F, et al. Physician alerts to increase antidepressant adherence: fax or fiction? *Arch Intern Med* 2006;166(5):498-504. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Banahan BF, Bentley JP. GIGO, NINO, and compliance-enhancement programs. *Computertalk for the Pharmacist* 1996;16(Jul-Aug):22-3. Database: IPA.

Exclude - Not a Primary Study

Barber N. Hospital computer prescribing: opportunity or threat? *Pharmaceutical Journal* 1989;244(Jun 30):786 Database: IPA.

Exclude - Not MMIT

Barber N. Designing information technology to support prescribing decision making. *Quality & Safety in Health Care* 2004;13(6):450-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Barclay S. Signature authentication can restore confidence. *Pharmaceutical Technology* 2007;31(6): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Bard M. E-prescribing cuts costs and reduces medical errors. *Managed Healthcare* 2009;12(5):46 Database: IPA.

Exclude - Not a Primary Study

Bargren M, Lu DF. An evaluation process for an electronic bar code medication administration information system in an acute care unit. *Urol Nurs* 391;29(5):355-67. PMID:19863043 OVID MEDLINE.

Exclude - Not MMIT

Barham C. The National Programme for Information Technology in the NHS. *Anaesthesia and Intensive Care Medicine* 2007;11(12):507-8. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Barker A. Electronic prescribing improves patient safety - An audit. Hospital Pharmacist 2007;14(7):225 Database: Embase Sept 22-09.

Exclude - Not MMIT

Barker KN, Pearson RE, Hepler CD, et al. Effect of an automated bedside dispensing machine on medication errors. Am J Hosp Pharm 1984;41(7):1352-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Barker KN, Allan EL. Research on drug-use-system errors. Am J Health Syst Pharm 1995;52(4):400-3. Database: IPA.

Exclude - Not a Primary Study

Barker KN, Felkey BG, Flynn EA, et al. White paper on automation in pharmacy. Consultant Pharmacist 1998;13(3):256-93. Database: IPA.

Exclude - Not a Primary Study

Barlas S. Surf's up! - New electronic health records initiative creates waves. P & T 2004;29(9):540 Database: IPA.

Exclude - Not MMIT

Barlas S. E-prescribing standards: A prescription for paralysis? Medicare excludes RxHub formulary Protocols. P & T 2005;30(12):693 Database: IPA.

Exclude - Not a Primary Study

Barlow RD. Unleashing the potential of RFID, RTLS. (cover story). Healthcare Purchasing News 2008;32(8):10-1.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=33362699&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Barlow, Rick Dana. Bar Coding: Hurdles at the starting block. Healthcare Purchasing News 2010 Mar34(3):16-21. 2010 Mar. KSR Publishing.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=48633301&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Barlow S, Johnson J, Steck J. The economic effect of implementing an EMR in an outpatient clinical setting. J Healthc Inf Manag 2004;18(1):46-51.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=60711> Grey Lit.

Exclude - Not MMIT

Barnes GD. Anticoagulation: Where we are and where we need to go. J Thromb Thrombolysis 2009;28(2):220-3. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Barnes PM, Magee MJ, Parnes JG, et al. Reengineering controlled substance distribution in a hospital pharmacy. ASHP Annual Meeting 1996;53: Database: IPA.

Exclude - Not a Primary Study

Barnsteiner JH. Medication reconciliation. Journal of Infusion Nursing 2005;28: Database: Embase Sept 22-09.

Exclude - Not MMIT

Baron RJ, Fabens EL, Schiffman M, et al. Electronic health records: just around the corner? Or over the cliff? *Ann Intern Med* 2005;143(3):222-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Baron RJ, Fabens EL, Schiffman M, et al. Perspective. Electronic health records: Just around the corner? Or over the cliff? *Ann Intern Med* 2005;143(3):222-6. Database: CINAHL.
Exclude - Not MMIT

Baronet G. Electronic medical prescription in a psychiatry unit. *Atencion Farmaceutica* 2008;10(2): Database: Embase Sept 22-09.
Exclude - Unable to Retrieve Foreign

Barret K, Watson A. Physician perspectives on a pilot prescription monitoring program. *J PAIN PALLIAT CARE PHARMACOTHER* 2005;19(3):5-13. Database: PsycINFO.
Exclude - Not MMIT

Barrett JS, Mondick JT, Narayan M, et al. Integration of modeling and simulation into hospital-based decision support systems guiding pediatric pharmacotherapy. *BMC Med Inform Decis Mak* 2008;8:6 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Barrett K, Watson A. Physician perspectives on a pilot prescription monitoring program. *J PAIN PALLIAT CARE PHARMACOTHER* 2005;19(3):5-13. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Barringer C, Williams JM, McCrirrick A, et al. Regional anaesthesia and propofol sedation for carotid endarterectomy. *ANZ Journal of Surgery* 2005;75(7):546-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Barrios AC. The comprehensive management of anticoagulation: Ochsner Coumadin Clinic. *Ochsner Journal* 2002;4(1):37-40. Database: Embase Sept 22-09.
Exclude - Not MMIT

Barrons R. Evaluation of personal digital assistant software for drug interactions. *Am J Health Syst Pharm* 2004;61(4):380-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Barrueco N. Computerized prescriber order entry with decision support: Drug interactions. *Acta Pediatr Esp* 2008;66(6): Database: Embase Sept 22-09.
Exclude - Unable to Retrieve Foreign

Bartelink IH. Error reduction by computerised physician order entry in hospitals. *Pharm Weekbl* 2003;138(20): Database: Embase Sept 22-09.
Exclude - Unable to Retrieve Foreign

Barton C. Contraindicated medication use among patients in a memory disorders clinic. *American Journal Geriatric Pharmacotherapy* 2008;6(3):147-52. Database: Embase Sept 22-09.
Exclude - Not MMIT

Bassi J, Lau F, Bardal S. Use of information technology in medication reconciliation: A scoping review. *Ann Pharmacother* 2010;44(5):885-97. OVID EMBASE.
Exclude - Not a Primary Study

Batenburg R, Van den BE. Pharmacy information systems: the experience and user satisfaction within a chain of Dutch pharmacies. *International Journal of Electronic Healthcare* 2008;4(2):119-31. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bates DW, Boyle DL, Vander Vliet MB, et al. Relationship between medication errors and adverse drug events. *J Gen Intern Med* 1995;10(4):199-205. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bates DW. Medication errors. How common are they and what can be done to prevent them? *Drug Saf* 1996;15(5):303-10. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bates DW, Makary MA, Teich JM, et al. Asking residents about adverse events in a computer dialogue: how accurate are they? *Jt Comm J Qual Improv* 1998;24(4):197-202. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bates DW. Frequency, consequences and prevention of adverse drug events. *J Qual Clin Pract* 1999;19(1):13-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Bates DW, Pappius E, Kuperman GJ, et al. Using information systems to measure and improve quality. *Int J Med Inf* 1999;53(2-3):115-24. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Bates DW. Medication safety issues in the new millennium. *ASHP Annual Meeting* 2000;57: Database: IPA.
Exclude - Not a Primary Study

Bates DW. Using information technology to reduce rates of medication errors in hospitals. *Br Med J* 2000;320(7237): Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Bates DW. Using information technology to reduce rates of medication errors in hospitals. *Br Med J* 2000;320(Mar 18):788-91. Database: IPA.
Exclude - Not a Primary Study

Bates DW, Cohen M, Leape LL, et al. Reducing the frequency of errors in medicine using information technology. *J Am Med Inform Assoc* 2001;8(4):299-308. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Bates DW. John M. Eisenberg Patient Safety Awards. Research: David W. Bates, MD, MSc, Brigham and Women's Hospital. Interview by Steven Berman. *Jt Comm J Qual Improv* 2002;28(12):651-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Bates DW, Evans RS, Murff H, et al. Detecting adverse events using information technology. *J Am Med Inform Assoc* 2003;10(2):115-28. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bates DW. Using information technology to screen for adverse drug events. *Am J Health Syst Pharm* 2003;59(23):2317-9. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Bates DW. Preventing medication errors: a summary. *Am J Health Syst Pharm* 2007;64(14:Suppl:9):S3-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bates D and Gawande A. Improving safety with information technology. 2007. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=63038> Grey Lit.

Exclude - Not a Primary Study

Bauman JL, Didomenico RJ, Galanter WL. Mechanisms, manifestations, and management of digoxin toxicity in the modern era. *American Journal of Cardiovascular Drugs* 2006;6(2):77-86. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Beard R, Candlish C. Does electronic prescribing contribute to clinical governance? *BJHC & IM* 2005;21(4):27-9. Database: CINAHL.

Exclude - No Outcomes of Interest

Beaucaire G. How to improve the prescription of antibiotics in hospitals? Training, organization, and evaluation: Evaluation and action. *Medecine et Maladies Infectieuses* 2003;33(Suppl 1):93-104. Database: Embase Sept 22-09.

Exclude - Not MMIT

Beaulieu JE, McGaughey A, Matulewicz A, et al. Utilizing information technology on patient care rounds to enhance pharmacist effectiveness. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Bedouch P, Allenet B, Labarere J, et al. Diffusion of pharmacist interventions within the framework of clinical pharmacy activity in the clinical ward. *Therapie* 2005;60(5):515-22. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Bedouch P, Baudrant M, Detavernier M, et al. Drug supply chain safety in hospitals: current data and experience of the Grenoble university hospital. *Ann Pharm Fr* 2009;67(1):3-15.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Beers MH, Fingold SF, Ouslander JG. A computerized system for identifying and informing physicians about problematic drug use in nursing homes. *J Med Syst* 1992;16(6):237-45.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Begg A. Computerized decision support system is key to anticoagulation service. *Guidelines in Practice* 2009;8(2):35-6. Database: CINAHL.

Exclude - No Outcomes of Interest

Begg E, Atkinson H, Jeffery G, et al. Individualised aminoglycoside dosage based on pharmacokinetic analysis is superior to dosage based on physician intuition at achieving target plasma drug concentrations. *Br J Clin Pharmacol* 1989;28:137-41. Exclude - Not MMIT

Bekker H, Thornton JG, Airey CM, et al. Informed decision making: An annotated bibliography and systematic review. *Health Technol Assess* 1999;3(1):1-156. <http://www.hta.ac.uk/project.asp?PjtId=973> Grey Lit. Exclude - Not MMIT

Belgamwar RB. Use of hospital prescribing data to monitor the implementation of clinical guidelines. *Pharmacy Education* 2005;5(3-4):219-21. Database: Embase Sept 22-09. Exclude - No Outcomes of Interest

Bell DS, Cretin S, Marken RS, et al. A conceptual framework for evaluating outpatient electronic prescribing systems based on their functional capabilities. *J Am Med Inform Assoc* 2004;11(1):60-70. Database: Ovid MEDLINE(R). Exclude - Not a Primary Study

Bell DS, Friedman MA. E-prescribing and the medicare modernization act of 2003. *Health Aff (Millwood)* 2005;24(5):1159-69. Database: Ovid MEDLINE(R). Exclude - Not a Primary Study

Bell JS, Vaananen M, Ovaskainen H, et al. Providing patient care in community pharmacies: practice and research in Finland. *Ann Pharmacother* 2007;41(6):1039-46. Database: Ovid MEDLINE(R). Exclude - Not MMIT

Bellazzi R, Arcelloni M, Ferrari P, et al. Management of patients with diabetes through information technology: tools for monitoring and control of the patients' metabolic behavior. *Diabetes Technol Ther* 2004;6(5):567-78. Database: Ovid MEDLINE(R). Exclude - Not MMIT

Benjamin DM. Reducing medication errors and increasing patient safety: case studies in clinical pharmacology. *J Clin Pharmacol* 2003;43(7):768-83. Database: Ovid MEDLINE(R). Exclude - Not a Primary Study

Bennett J W, Glasziou P P. Computerised support of medication management: A systematic review. In 2002; Melbourne, Australia): 2002. p.16-7. Grey Lit. Exclude - Not a Primary Study

Bennett JW, Glasziou P, Del Mar C, et al. A computerised prescribing decision support system to improve patient adherence with prescribing. A randomised controlled trial. *Aust Fam Physician* 2003;32(8):667-71. Database: Ovid MEDLINE(R). Exclude - Not MMIT

Bennett JW, Glasziou PP. Computerised reminders and feedback in medication management: a systematic review of randomised controlled trials. *Med J Aust* 2003;178(5):217-22. Database: Ovid MEDLINE(R). Exclude - Not a Primary Study

Bennett SW, Scott AC. Computer assisted customized antimicrobial dosages. *Am J Hosp Pharm* 1980;37(4):523-9. Database: IPA.

Exclude - Not a Primary Study

Benrimoj SI, Thornton PD, Langford JH. Review of drug distribution systems. Part 2. Automated dispensing for unit dose systems. *Australian Journal of Hospital Pharmacy* 1995;25(3):230-5. Database: IPA.

Exclude - Not MMIT

Benson H. Florida health information technology strategy. National Conference of State Legislatures; 2007. <http://www.ncsl.org/programs/health/shn/2006/sn480c.htm> Grey Lit.

Exclude - Not MMIT

Benson VM. Dyslipidemia treatment guidelines and management systems in Kaiser Permanente. *Am J Cardiol* 1997;80(8 Suppl2):85H-8H. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Berard CM, Mahoney CD, Welch DW, et al. Computer software for pharmacy oncology services. *Am J Health Syst Pharm* 1996;53(7):752-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Berdine HJ, Nesbit SA. Equianalgesic dosing of opioids. *J PAIN PALLIAT CARE PHARMACOTHER* 2006;20(4):79-84. Database: IPA.

Exclude - Not a Primary Study

Berend ME. Computer-assisted TKA: greater precision, doubtful clinical efficacy: affirms. *Orthopedics* 2009;32(9):678

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010413390&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1212&accno=2010413390 EBSCO CINAHL.

Exclude - Not a Primary Study

Berg KM, Arnsten JH. Practical and conceptual challenges in measuring antiretroviral adherence. *Journal of Acquired Immune Deficiency Syndromes: JAIDS* 2006;43(Suppl 1):S79-87. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Berger RG, Kichak JP. Computerized physician order entry: helpful or harmful?[see comment. *J Am Med Inform Assoc* 2004;11(2):100-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bergeron BP. Medical errors: computers are no panacea. *J Med Pract Manage* 2005;21(1):31-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bergman A, Olsson J, Carlsten A, et al. Evaluation of the quality of drug therapy among elderly patients in nursing homes. *Scand J Prim Health Care* 2007;25(1):9-14. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bergman SJ, Slain D, Petros KO. Evaluation of a newly implemented once-daily aminoglycoside dosing and monitoring program. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Bergman U. Use of antibiotics at hospitals in Stockholm: A benchmarking project using internet. *Pharmacoepidemiology and Drug Safety* 2004;13(7):465-71. Database: Embase Sept 22-09.

Exclude - Not MMIT

Berkenstadt H, Yusim Y, Katznelson R, et al. A novel point-of-care information system reduces anaesthesiologists' errors while managing case scenarios. *Eur J Anaesthesiol* 2006;23(3):239-50. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bermejo T, Hidalgo F, Beltran-Huertas Y, et al. Economical impact of an automated dispensing system in the emergency ward of a general hospital. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Bermejo T, Hidalgo F, Beltran-Huertas Y, et al. Opinion survey about the correct use of an automated dispensing machine in the emergency ward of a general hospital. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Bermejo VT, Alvarez Diaz AM, Delgado SE, et al. Analysis of failures detected during the medication-dispensing process and their contributing factors. [Spanish]. *Trauma* 2009;20(3):194-9. OVID EMBASE.

Exclude - Unable to Retrieve

Bernard D, Vanhee D, Blaton V. Implementation of an integrated instrument control and data management system for point of care blood gas testing. *Clin Chim Acta* 2001;307(1-2):169-73. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bernardi GM. Barcode: Contribution to the rationalization in the assembly process. *Krankenhauspharmazie* 1990;11(Jun):246-50. Database: IPA.

Exclude - Unable to Retrieve

Berner ES, Kasiraman RK, Yu F, et al. Data quality in the outpatient setting: impact on clinical decision support systems. *AMIA* 2005;41-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Berner ES. Solutions in the non-peer-reviewed literature for reducing medication errors. *Journal of Pharmaceutical Finance, Economics and Policy* 2007;15(3): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Berner E. Clinical decision support systems: State of the art. *AHRQ*; 2009. http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_874024_0_0_18/09-0069-EF.pdf Grey Lit.

Exclude - Not a Primary Study

Berney DR, Palmgren GN, Stowers AD. Use of telepharmacy technology offers potential for improved financial management. [Abstract] Ashp Midyear Clinical Meeting 2001;36:

Reference ID: 27791

Notes: 38-12948

English

Abstract of meeting presentation. Journal article

200111

Bernstein E L. The use of real time clinical decision support in integrated medication therapy. In Newton, MA, USA: Med. Records Inst; 1997. p.228-32.5787582

Database: Inspec.

Exclude - No Outcomes of Interest

Berry D. Bar coding rule good news for hospital pharmacy. Pharmacy Today (Washington D C) 2004;10(4):24 Database: IPA.

Exclude - Not a Primary Study

Berry D, Gillie T, Banbury S. What do patients want to know: An empirical approach to explanation generation and validation. Expert Systems with Applications 1995;8(4):419-28.

1995292724265

Database: Compendex.

Exclude - Not MMIT

Bertsche T, Kaltschmidt J, Haefeli WE. Patient safety based on computer-assisted drug therapy. Electronic check-up of the patient. Internist 2009;50(6):748-56. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bertsche T, Pfaff J, Schiller P, et al. Prevention of adverse drug reactions in intensive care patients by personal intervention based on an electronic clinical decision support system. Intensive Care Med 2010;36(4):665-72. OVID EMBASE.

Exclude - Not MMIT

Besier J. Organizational behavior in a complex system. A pilot study: The impact of a change model on health care practice improvement. DAI 2004;65(1): Database: Sociological Abstracts.

Exclude - Not MMIT

Best V. Medicare's e-prescribing incentive program. An introduction. Adv Nurse Pract 2009;17(5):49-50. PMID:20014717 OVID MEDLINE.

Exclude - Not a Primary Study

Beuscart-Zephir MC, Pelayo S, Anceaux F, et al. Cognitive analysis of physicians and nurses cooperation in the medication ordering and administration process. Int J Med Inf 2007;76(Suppl 1):S65-77. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bhatt AD. Drug-related problems and adverse drug events: negligence, litigation and prevention. J Assoc Physicians India 1999;47(7):715-20. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bhattacharjee A, Hikmet N. Physicians' resistance toward healthcare information technology: A theoretical model and empirical test. *European Journal of Information Systems* 2007;16(6):725-37. Database: BSC.

Exclude - No Outcomes of Interest

Bickham P. Improving patient safety and compliance with Joint Commission on Accreditation of Healthcare Organizations (JCAHO) medication management standards in radiology. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Bindoff I, Tenni P, Kang B and others. Intelligent decision support for medication review. In Berlin, Germany: Springer; 2006. p.120-31.9253598

Database: Inspec.

Exclude - No Outcomes of Interest

Bindoff IK, Tenni PC, Peterson GM, et al. Development of an intelligent decision support system for medication review. *J Clin Pharm Ther* 2007;32(1):81-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Binnersley J, Woodcock A, Wallace L M and others. Establishing user requirements for a patient held electronic record system in the United Kingdom. In San Antonio, TX, United states: Human Factors an Ergonomics Society Inc.; 2009. p.714-7. Engineering Village Compendex and Inspec.

Exclude - Unable to Retrieve

Binns P. The impact of the electronic health record on patient safety: an Alberta perspective. *Healthcarepapers* 1982;5(3):47-51. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Birdsey GH, Weeks GR, Bortoletto DA, et al. Pharmacist-initiated electronic discharge prescribing for cardiology patients. *Journal of Pharmacy Practice & Research* 2005;35(4):287-90. Database: IPA.

Exclude - Not MMIT

Bishop SC, Rodriguez EM, Cebula MW. Conversion of conventional human insulin vials to analog insulin pens in a community hospital. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Bjeldbak-Olesen I, Jacklin A. E-health solutions in hospital pharmacy. *EJHP Practice* 2008;14(1):22 OVID EMBASE.

Exclude - Not a Primary Study

Bjornstad GC, Farr F, Vernon P, et al. Automated patient care documentation: what's in it for us? An expert system emergency drug card printout. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1995;17-21. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Blackledge CG, Jr., Veltri MA, Matlin C, et al. Patient safety in emergency situations: A Web-based pediatric arrest medication calculator. *Journal for Healthcare Quality: Promoting Excellence in Healthcare* 2006;28(2):27-31. Database: CINAHL.

Exclude - Not a Primary Study

Blake RT, Massey AP, Bala H, et al. Driving health IT implementation success: Insights from The Christ Hospital. *Bus Horiz* 2010;53(2):131-8.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=47956940&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Blakey SA, Longley JM, Mullis SR, et al. Implementation and evaluation of a pharmacy-based computer-assisted antimicrobial surveillance service. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Blank D. Biometric devices: Are you ready for them? *Drug Topics* 2002;146(7):66 Database: IPA.

Exclude - Not a Primary Study

Blendon RJ, Schoen C, Donelan K, et al. Physicians' views on quality of care: a five-country comparison. *Health Aff (Millwood)* 2001;20(3):233-43. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bloom MZ. Connecting physicians to pharmacies and patients. *J Manag Care Pharm* 1998;4(Sep-Oct):473-4. Database: IPA.

Exclude - Not a Primary Study

Bloomfield EL. The anesthesia information management system for electronic documentation: What are we waiting for? *Journal of Anesthesia* 2008;22(4):404-11.

Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Bluestein D, Brantley C, Barnes-Eley M, et al. Measuring international normalized ratios in long-term care: a comparison of commercial laboratory and point-of-care device results. *J AM MED DIR ASSOC* 2007;8(6):404-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Blum BI, Johns CJ, McColligan EE, et al. Low cost ambulatory medical information system. *J Clin Eng* 1979;4(4):372-7. 1565930

Database: Inspec.

Exclude - Not MMIT

Blumenfeld B. Computerized physician order entry of medications and fluids in a neonatal intensive care unit (NICU). *Clinical Informatics Research & Development*; 2002.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=30562> Grey Lit.

Exclude - Not a Primary Study

Bobb AM, Liebovitz DM, Christensen KL, et al. Clinical decision support to avoid adverse drug events with anticoagulants. *AMIA* 2007;874 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bobb AM, Payne TH, Gross PA. Viewpoint: Controversies surrounding use of order sets for clinical decision support in computerized provider order entry. *J Am Med Inform Assoc* 2007;14(1):41-7. Database: CINAHL.

Exclude - Not a Primary Study

Bobb AM, Boecler LA. Optimization and maintenance. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Bochicchio GV, Smit PA, Moore R, et al. Pilot study of a web-based antibiotic decision management guide. *J Am Coll Surg* 2006;202(3):459-67. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Boecler LA, Bobb AM. Implementation of CPOE. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Boerner CM. View HIPAA Breaches Affecting 500 or More Individuals Online. *Journal of Health Care Compliance* 2010;12(3):31-68.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=50320776&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Bogner M. There is more to error in healthcare than the care provider. In Orlando, FL, United states: Human Factors an Ergonomics Society Inc.; 2005. p.952-4.20082311293334

Database: Compendex.

Exclude - Not MMIT

Bojke C, Nestor G. What determines GPs' use of a clinical decision support software system: An analysis from PRODIGY Phase II. *Health Informatics Journal* 2000;6(3):147-55.

Database: CINAHL.

Exclude - Not a Primary Study

Bollen C, Warren J, Whenan G. Introduction of electronic prescribing in an aged care facility. *Aust Fam Physician* 2005;34(4):283-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bomba D, Land T. The feasibility of implementing an electronic prescribing decision support system: a case study of an Australian public hospital. *Aust Health Rev* 2006;30(3):380-8.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bond CA, Raehl CL. 2006 national clinical pharmacy services survey: clinical pharmacy services, collaborative drug management, medication errors, and pharmacy technology. *Pharmacotherapy* 2008;28(1):1-13. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bondmass M. The effect of home monitoring and telemanagement on blood pressure control among African Americans. *Telemed J* 2000;6(1):15-23. Database: Embase Sept 22-09.

Exclude - Not MMIT

Bonnevie B, Jensen BA. Prescribing and dispensing drugs in Denmark. Frequency of and intervention against errors in documentation and dispensing of drugs. *Ugeskr Laeger* 2002;164(40):4656-9. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Bordowitz R. Electronic health records: A primer. *Laboratory Medicine* 2008;39(5):301-7. Database: CINAHL.

Exclude - Not a Primary Study

Bornhauser M, Schmidt M, Ehninger U, et al. Computer-based quality control in high-dose chemotherapy and bone marrow transplantation. *Bone Marrow Transplant* 1998;21(5):505-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Borsi JD, Klepp O, Moe PJ. PharmCalc: program for the calculation of clinical pharmacokinetic parameters of methotrexate. *Cancer Chemotherapy & Pharmacology* 1988;22(4):339-43. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Borycki EM, Kushniruk A, Key E, et al. Toward an integrated simulation approach for predicting and preventing technology-induced errors in healthcare: implications for healthcare decision-makers. *HEALTHC Q* 2009;12(Spec.):90-6. PMID:19667784 OVID MEDLINE.

Exclude - Not a Primary Study

Bossen C. Test the artefact--develop the organization. The implementation of an electronic medication plan. *Int J Med Inf* 2007;76(1):13-21. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bosworth HB, Olsen MK, Oddone EZ. Improving blood pressure control by tailored feedback to patients and clinicians. *Am Heart J* 2005;149(5):795-803. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bosworth K, Gustafson D. CHESS: Providing decision support for reducing health risk behavior and improving access to health services. *Interfaces* 1991;21(3):93-104. Database: BSC.

Exclude - Not a Primary Study

Bot H, Brouwer J, Dijk W A and others. Brewer: An integrated on-line coding system for diagnoses and medication. In 1994; (Bethesda; MD): 1994. p.185-8. Grey Lit.

Exclude - Not a Primary Study

Bott OJ. HIS modelling and simulation based cost-benefit analysis of a telemedical system for closed-loop diabetes therapy. *Int J Med Inf* 2007;76: Database: Embase Sept 22-09.

Exclude - Not MMIT

Bottiger Y, Laine K, Andersson ML, et al. SFINX-a drug-drug interaction database designed for clinical decision support systems. *Eur J Clin Pharmacol* 2009;65(6):627-33. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bouam S, Girou E, Brun-Buisson C, et al. Development of a Web-based clinical information system for surveillance of multiresistant organisms and nosocomial infections. *AMIA* 1999;1999(1999):696-700. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bouchand F, Thomas A, Zerhouni L, et al. Pharmacists' interventions before and after prescription computerization in an internal medicine department. *Presse Med* 2007;36(3p1):410-8. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Bouhaddou O, Warner H. An interactive patient information and education system (Medical HouseCall) based on a physician expert system (Iliad). *Medinfo* 1995;8 Pt 2:1181-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bouhaddou O, Warnekar P, Parrish F, et al. Exchange of computable patient data between the Department of Veterans Affairs (VA) and the Department of Defense (DoD): terminology mediation strategy. *J Am Med Inform Assoc* 2008;15(2):174-83. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Boukhors Y. The use of information technology for the management of intensive insulin therapy in type 1 diabetes mellitus. *Diabetes Metab* 2004;29(6):619-27. Database: Embase Sept 22-09.

Exclude - Not MMIT

Bourget P, Barath V, Guntz JP, et al. Pharmaceutical traceability integrated with the patient file. Development of a computerized hospital application. *Pathol Biol* 2001;49(8):624-33. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bourka A, Kaliontzoglou A, Polemi D, et al. Enriching healthcare applications with cryptographic mechanisms and XML- based security services. *Technology & Health Care* 2003;11(1):61-76. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bourret JA. Major medication management issues in an integrated health system. *ASHP Annual Meeting* 1996;53: Database: IPA.

Exclude - Not a Primary Study

Bourret JA, Demers RF, Wordell D, et al. Medication use review process and information systems utilized for oncology chemotherapy quality improvement. *Pharm Pract Manag Q* 1996;16(1):1-17. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Boustani M, Munger S, Beck R, et al. A gero-informatics tool to enhance the care of hospitalized older adults with cognitive impairment. *Clinical Interventions In Aging* 2007;2(2):247-53. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bowman L, Carlstedt BC, Black CD. Detection of adverse drug reactions (ADRs) in hospitalized patients using computerized physician online order entry (OLOE). Ashp Midyear Clinical Meeting 1992;27: Database: IPA.
Exclude - Not a Primary Study

Bowser A. Brigham and women's embarks on integrated bar-code system. Pharmacy Practice News 2004;31(11):32 Database: IPA.
Exclude - Not a Primary Study

Boxer M. Impact of a real-time peer review audit on patient management in a radiation oncology department. Journal of Medical Imaging and Radiation Oncology 2009;53(4):405-11. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Boyd M. Bar coding for patients streamlines drug treatment. New Zealand Pharmacy 1993;13(3):22-3. Database: IPA.
Exclude - No Outcomes of Interest

Brailer DJ. Translating ideals for health information technology into practice. Health Aff (Millwood) 2004;318-20. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Bramble JD, Galt KA, Siracuse MV, et al. The relationship between physician practice characteristics and physician adoption of electronic health records. Health Care Manage Rev 2010;35(1):55-64.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010531792&site=ehost-live> EBSCO CINAHL.
Exclude - Not MMIT

Brandell R, Shelton M. Bar code research: Pharmacokinetic services and satellite pharmacies. ASHP Annual Meeting 1990;47: Database: IPA.
Exclude - Not a Primary Study

Brandell R, Taylor M, Meyer G, et al. Development of on-line drug specific information screens to improve the quality of medication use. Ashp Midyear Clinical Meeting 1993;28: Database: IPA.
Exclude - Not a Primary Study

Brandenburg HC, Vermeij P. Optically readable code in pharmacy. Pharm Weekbl 1987;122:505-9. Database: IPA.
Exclude - Unable to Retrieve Foreign

Branger PJ, van 't HA, van der Wouden JC, et al. Shared care for diabetes: supporting communication between primary and secondary care. Studies in Health Technology & Informatics 1998;52 Pt 1:412-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Branham CN. Washington update. Ocular Surgery News 2008;26(23):38
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010181329&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2294&accno=2010181329
EBSCO CINAHL.
Exclude - Not a Primary Study

Branham CN. Washington update. Ocular Surgery News 2009;27(9):46
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010283453&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2294&accno=2010283453
EBSCO CINAHL.

Exclude - Not a Primary Study

Branowicki P, O'Neill JB, Dwyer JL, et al. Improving complex medication systems: an interdisciplinary approach. J Nurs Adm 2003;33(4):199-200. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bransgrove LL. Use of a computerized histamine-2 receptor antagonist DUE in ambulatory care patients. Ashp Midyear Clinical Meeting 1995;30: Database: IPA.

Exclude - Not a Primary Study

Braswell A, Duggar S. The new look of bedside technology. The point-of-care evolution drives providers to rethink nursing workflow and medication management. Nurs Manag (Harrow) 2006;37(10):14-32. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Brater C. Cross-check PC. M D Computing 1988;5(2):49-50. 3161705
Database: Inspec.

Exclude - No Outcomes of Interest

Brennan S, Spours A. Barriers to the successful and timely implementation of electronic prescribing and medicines administration. BJHC & IM 2001;17(8):22-5. Database: CINAHL.

Exclude - No Outcomes of Interest

Brennan S, Spours A. Electronic prescribing and medicines administration: Are we overcoming the barriers to success? BJHC & IM 2004;20(4):19-22. Database: CINAHL.

Exclude - No Outcomes of Interest

Brenner G. Special applications in health telematics: electronic prescription/electronic patient file/digital archiving. Z Arztl Fortbild Qualitatssich 2001;95(9):646-51. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Brenninkmeijer R. Application of SOJA and InforMatrix in practice: Interactive web and workshop tools. Expert Opinion on Pharmacotherapy 2007;8(Suppl 1):S49-55. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Brenninkmeijer R. InforMatrix as an alternative tool in rational and transparent drug-decision making. Expert Opinion on Pharmacotherapy 2007;8(Suppl 1):S31-6. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Brenninkmeijer RF. InforMatrix. Pharm Weekbl 1994;129(47-48): Database: Embase Sept 22-09.

Exclude - Not MMIT

Bressan N, Castro A, Bras S, et al. Synchronization software for automation in anesthesia. Conference Proceedings: 2007;2007(2007):5298-301. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Briggs B. I.T. helps battle 'The big C'. Health Data Manag 2004;12(5):56-8. Database: IPA.
Exclude - Not MMIT

Bright,T. Development and evaluation of an ontology for guiding appropriate antibiotic prescribing Columbia UniversityEditor. 2009. Grey Lit.
Exclude - Theses

Brill CW, Forster K, Keil W. Patients' mailbox and electronic prescription. Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz 2005;48(7):732-5. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Briscoe AP, Pancerella MB, Pleskunas G. Perfect order initiative. Pharmaceutical Executive 1984;17(7):82-6. Database: IPA.
Exclude - Not MMIT

Brixey J. Development of a screen interface for a medication error database. In 2001. Washington, DC: 2001. p. 867.Grey Lit.
Exclude - Not a Primary Study

Brock TP, Smith SR. Using digital videos displayed on personal digital assistants (PDAs) to enhance patient education in clinical settings. Int J Med Inf 2007;76(11-12):829-35. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Brockenbrough G. CMS-sponsored national conference held to advance the use of e-prescribing. Orthopedics Today 2008;28(11):32-3. Database: CINAHL.
Exclude - Not a Primary Study

Brockenbrough G. CMS-sponsored national conference held to advance the use of e-prescribing. Orthopedics Today 2008;28(11):32-3.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010098902&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1950&accno=2010098902
EBSCO CINAHL.
Exclude - Not a Primary Study

Brockenbrough G. New Medicare Part D e-prescribing standards for April 2009 issued by CMS. Orthopedics Today 2008;28(6):36-7. Database: CINAHL.
Exclude - Not a Primary Study

Brodts-Suggs J, Masulli T, Burke S. Scanning the horizon: A health system upgrades its bar coding and patient auto-identification for improving patient care enterprisewide. Health Manag Technol 2007;28(9):24-6. 9736006
Database: Inspec.
Exclude - Not a Primary Study

Broekema WJ. Automated drug-dispensing system in a general psychiatric hospital surpassing unit dosage. Pharm Weekbl 1990;125:1326-32. Database: IPA.
Exclude - Unable to Retrieve Foreign

Bronzino JD, Morelli RA, Goethe JW. OVERSEER: a prototype expert system for monitoring drug treatment in the psychiatric clinic. *IEEE Trans Biomed Eng* 1989;36(5):533-40. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Brooks P, Sonnenschein C. E-prescribing: where health information and patient care intersect. *J Healthc Inf Manag* 2010;24(2):53-9. PMID:20397335 OVID MEDLINE.

Exclude - Not a Primary Study

Brouwers JR. Drug distribution in hospitalized care: Past, present and future. *Ziekenhuisfarmacie* 1995;11(2):53-8. Database: IPA.

Exclude - Unable to Retrieve Foreign

Broverman C, Kapusnik-Uner J, Shalaby J, et al. A concept-based medication vocabulary: an essential requirement for pharmacy decision support. *Pharm Pract Manag Q* 1998;18(1):1-20. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Broverman C. Overcoming obstacles to medication decision support at point-of-care: Interim report on standardization efforts. In Philadelphia; PA): 1998. p. 39. Grey Lit.

Exclude - Unable to Retrieve

Broverman CA, Clyman JI, Schlesinger JM, et al. Clinical decision support for physician order-entry: design challenges. *AMIA Annual Fall Symposium* 1996;572-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Broverman CA, Schlesinger JM, Sperzel WD, et al. The future of knowledge-based components in the electronic health record. *Studies in Health Technology & Informatics* 1998;52(Pt 1):457-61. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Brown B, Patel H, Machaj D, et al. Implementation of barcode medication administration at VA Chicago health care system: an into-disciplinary performance improvement project focused on patient safety. *Ashp Summer Meeting* 2003;60. Database: IPA.

Exclude - Not a Primary Study

Brown BT. Re-engineering a VA outpatient pharmacy using automation reduces patient waiting times. *ASHP Annual Meeting* 1999;56. Database: IPA.

Exclude - Not a Primary Study

Brown B. 25 Steps to Meaningful Use. *Journal of Health Care Compliance* 2010;12(3):33-69.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=50320777&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Brown JS, Kulldorff M, Chan KA, et al. Early detection of adverse drug events within population-based health networks: Application of sequential testing methods. *Pharmacoepidemiology and Drug Safety* 2007;16(12):1275-84. Database: Embase Sept 22-09.

Exclude - Not MMIT

Brown L and Frye J. Is CPOE safe for patients? Society for Health Systems Conference and Expo, Orlando, FL, United states: Institute of Industrial Engineers; 2008. 20091612039256 Database: Compendex.

Exclude - Not MMIT

Brown V. Reconcilable differences: A Washington healthcare enterprise works collaboratively to create a comprehensive medication reconciliation solution. Health Manag Technol 2008;29(1):12-6. 9816589

Database: Inspec.

Exclude - Not a Primary Study

Brudieu E, Grain F, Bosson JL, et al. Pharmacist's interventions evaluation on computerized prescriptions. Journal de Pharmacie Clinique 1999;18(3):227-32. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Bruegman-May K, Murphy M, Pemble K et al. Bar coded eMAR: The SynergyHealth St. Joseph's experience (Goin' to the Bar-Code). 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_226703_0_0_18/StJosephs-BCMA-AHRQ.pdf Grey Lit.

Exclude - Not MMIT

Bu D, Pan E, Walker J, et al. Benefits of information technology-enabled diabetes management. Diabetes Care 2007;30(5):1137-42. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bucher HC, Rickenbach M, Young J, et al. Randomized trial of a computerized coronary heart disease risk assessment tool in HIV-infected patients receiving combination antiretroviral therapy. Antiviral Therapy 2010;15(1):31-40. PMID:20167989 OVID MEDLINE.

Exclude - Not MMIT

Buck MD, Atreja A, Brunker CP, et al. Potentially inappropriate medication prescribing in outpatient practices: prevalence and patient characteristics based on electronic health records. American Journal Geriatric Pharmacotherapy 2009;7(2):84-92. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Buck,M.D. Development and impact of computerized clinical decision support alerts on prescribing for elderly outpatients The University of Utah. 2008. Grey Lit preview pdf saved-don't have full-text.

Exclude - Theses

Buckles CL, Howard E. Cost effectiveness of an automated dispensing and storage system. Ashp Midyear Clinical Meeting 1992;27: Database: IPA.

Exclude - Not a Primary Study

Buckley B. Cedars-sinai suspends CPOE system. Pharmacy Practice News 2003;30(4):26 Database: IPA.

Exclude - Not a Primary Study

Buckley B. Can bar-coding efforts keep pace with JCAHO? Pharmacy Practice News 2004;31(7): Database: IPA.

Exclude - Not a Primary Study

Buffington DE, Lampasona V, Chandler MH. Computers in pharmacokinetics. Choosing software for clinical decision making. Clin Pharmacokinet 1993;25(3):205-16. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Buising KL, Thursky KA, Robertson MB, et al. Electronic antibiotic stewardship--reduced consumption of broad-spectrum antibiotics using a computerized antimicrobial approval system in a hospital setting. J Antimicrob Chemother 2008;62(3):608-16. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Bukowsky SS. Redesign for the future, now! ASHP Annual Meeting 1998;55: Database: IPA.

Exclude - Not a Primary Study

Bukunt S, Hunter C, Perkins S, et al. El Camino Hospital: using health information technology to promote patient safety. Jt Comm J Qual Patient Saf 2005;31(10):561-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Bullock-Palmer RP, Weiss S, Hyman C. Innovative approaches to increase deep vein thrombosis prophylaxis rate resulting in a decrease in hospital-acquired deep vein thrombosis at a tertiary-care teaching hospital. Journal of Hospital Medicine 2008;3(2):148-55. Database: Embase Sept 22-09.

Exclude - Not MMIT

Bunn RJ, Taylor D, Holmes N, et al. Use of computerized patient medication profiles as an aid to prescribing, prescribing review and dispensing. Pharmaceutical Journal 1990;244:690-1. Database: IPA.

Exclude - Unable to Retrieve

Burgdorf AW, Phillips EM. Evaluation of a computerized physician order entry (CPOE) scheme for deep vein thrombosis (DVT) prophylaxis in a tertiary-care teaching hospital. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.

Exclude - Not a Primary Study

Burke DE, Wang BBL, Wan TTH, et al. Exploring hospitals' adoption of information technology. J Med Syst 2004;26(4):349-55. Database: Embase Sept 22-09.

Exclude - Not MMIT

Burke J. Reducing rx drug abuse through electronic monitoring. Pharm Times 1981;68(3):46 Database: IPA.

Exclude - Not a Primary Study

Burke J. Bar coding thwarts illegitimate drug use. Pharm Times 1984;71(12):84 Database: IPA.

Exclude - Not a Primary Study

Burke J. Rx abuse in rural America. Pharm Times 2009;75(11):62 OVID EMBASE.
Exclude - Not a Primary Study

Burke JP. Antibiotic resistance - Squeezing the balloon? JAMA 1998;280(14):1270-1.
Database: IPA.
Exclude - Not a Primary Study

Burke JP. Rational approaches to combating resistance. Int J Clin Pract 2002;(125):29-36.
Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Burkle T, Prokosch HU, Hussak G, et al. Knowledge based functions for routine use at a German university hospital setting: the issue of fine tuning. Proceedings/AMIA Annual Fall Symposium 1997;61-5. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Burns LL. Impact of an automated antibiotic refrigerator and an online medication administration record on timeliness of postoperative antibiotics on a spinal surgical unit. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Burton ME, Ash CL, Hill DP, et al. A controlled trial of the cost benefit of computerized bayesian aminoglycoside administration. Clinical Pharmacology & Therapeutics 1991;49(6):685-94. Exclude - Not MMIT

Burton SS, Duffus PR, Williams A. Exploration of the role of the clinical pharmacist in general practice medicine. Pharmaceutical Journal 1995;254(6824):91-3. Database: IPA.
Exclude - Not MMIT

Bury J, Hurt C, Roy A, et al. A quantitative and qualitative evaluation of LISA, a decision support system for chemotherapy dosing in childhood acute lymphoblastic leukaemia. Studies in Health Technology & Informatics 2004;107(Pt:1):197-201. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Bury J, Hurt C, Roy A, et al. LISA: a web-based decision-support system for trial management of childhood acute lymphoblastic leukaemia. Br J Haematol 2005;129(6):746-54. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Bussieres JF, Lebel D, Erickson L, et al. Evaluation of the ability of pharmacists to extract specific data from their pharmacy information management system. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.
Exclude - Not a Primary Study

Bussieres JF, Lebel D. Use of barcodes in the drug circuit of health establishments. Le Pharmactuel 2009;42(2):131-7. Database: IPA.
Exclude - No Outcomes of Interest

Bussmann H, Wester CW, Ndwapi N, et al. Hybrid data capture for monitoring patients on highly active antiretroviral therapy (HAART) in urban Botswana. Bull World Health Organ 2006;84(2):127-31. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Butler SF, Budman SH, Licari A, et al. National Addictions Vigilance Intervention and Prevention Program (NAVIPPRO): A real-time, product-specific, public health surveillance system for monitoring prescription drug abuse. *Pharmacoepidemiology and Drug Safety* 2008;17(12):1142-54. Database: Embase Sept 22-09.

Exclude - Not MMIT

Button E, Keaton P. Glycemic control after coronary bypass graft: using intravenous insulin regulated by a computerized system. *Crit Care Nurs Clin North Am* 2006;18(2):257-65. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Butts JD, Brookfield WP, Dempsey JH, et al. Use of bar coding to reduce the time required for relational database entry of data from an antibiotic order form. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Buurma H, De Smet PA, Kruijtbosch M, et al. Disease and intolerability documentation in electronic patient records. *Ann Pharmacother* 2005;39(10):1640-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Buurma,H. Clinical risk management in community pharmacy University Utrecht. 2008. Database: Embase Sept 22-09.

Exclude - Not MMIT

Buyschaert M, Jacques D, Donckier J, et al. Effect of computer-assisted insulin delivery on glycemic control of type I diabetic patients: a preliminary experience. *Acta Clin Belg* 1989;44(3):169-73. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Byrne CM, Mercincavage LM, Pan EC, et al. The value from investments in health information technology at the U.S. Department of Veterans Affairs. *Health Aff (Millwood)* 2010;29(4):629-38.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010629263&site=ehost-live>; EBSCO CINAHL.

Exclude - Not MMIT

Cabeza BJ, Salmeron GA, Vallejo R, I, et al. Technological innovation management in the implementation of computer-assisted prescription. Structure, process and results. [Spanish]. *Atencion Farmaceutica* 2009;11(4):221-31. OVID EMBASE.

Exclude - Unable to Retrieve

Cabie ED, Lee L, Hughes S. Implementation of a pediatric IV admixture service using standardization. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Cabrera RC, Del Prado Llergo JR, Panadero Ruz D, et al. Computer programme for dispensation to cystic fibrosis out-patients. *Farmacia Hospitalaria* 1992;16(Suppl 1):59-62. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Caesar BR, Hutchinson B. Reducing medication errors by using applied technology. *Nursing (Lond)* 2006;36(8):24-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cafazzo JA, Trbovich PL, Cassano-Piche A, et al. Human factors perspectives on a systemic approach to ensuring a safer medication delivery process. *HEALTHC Q* 2009;12(Spec):70-4. PMID:19667781 OVID MEDLINE.

Exclude - Not a Primary Study

Cafiero AC. Reducing medication errors in a long-term care setting. *Annals of Long-Term Care* 2003;11(2):29-35. Database: Embase Sept 22-09.

Exclude - Not MMIT

Cain C. AHRQ and CMS Pilot projects of ePrescribing standards. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Caiola K, Caliendo GC, Meyer J, et al. Continuity of care between outpatient and inpatient settings in hemodialysis patients. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Calabrese VS, Matthewson JR, Coblio NA, et al. Promoting the cognitive skills of the pharmacist: Use of telepharmacy. *Ashp Midyear Clinical Meeting* 1999;34: Database: IPA.

Exclude - Not a Primary Study

Calabretto J P, Warren J, Darzanos K and others. Building common ground for communication between patients and community pharmacists with an Internet medicine cabinet. In Los Alamitos, CA, USA: IEEE Comput. Soc; 2002. p. 161.7205349 Database: Inspec.

Exclude - Not MMIT

Calabretto JP, Warren J, Bird L. Pharmacy decision support: Where is it? A systematic literature review. *International Journal of Pharmacy Practice* 2005;13(3):157-63. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Calabretto J P, Banerjee D, Warren J and others. XForms and XML events to support decisions about medication management. In Washington, DC): 2005. p. 909. Grey Lit.

Exclude - Not a Primary Study

Calabretto J P, Warren J, Bird L. Missing information for decision making in medication management. In 1913; (Melbourne, Australia) Harrogate, England): Health Informatics Society of Australia; 2006. p.161-70.

<http://search.informit.com.au/documentSummary;dn=994875327326711;res=IELHEA> Grey Lit.

Exclude - No Outcomes of Interest

Calabretto JP, Swatman PMC. Towards a working document to support medication management. *Current Perspectives in Healthcare Computing* 2007;161-70. Grey Lit.

Exclude - No Outcomes of Interest

Caldwell R. Alerting staff to medication errors: Hospitals can save \$90,000 a year averting extended stays. *Health Manag Technol* 2000;21(8):52 6698266

Database: Inspec.

Exclude - Not a Primary Study

Caldwell RD, White SJ. Lessons learned from CPOE implementation (The Stanford experience): Practical applications vs. practice model changes. *Ashp Midyear Clinical Meeting* 1979;38: Database: IPA.

Exclude - Not a Primary Study

Caldwell RD, Teng D, Wei JC. Use of an automated dispensing device system to assist in patient care monitoring activities. *Ashp Midyear Clinical Meeting* 1997;32: Database: IPA.

Exclude - Not a Primary Study

Caldwell RD. Physician order entry: Stanford experience. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Caldwell RD. Physician electronic order entry one year later - Continual learning. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Caldwell RD, Follain DP. Implementation of computerized physician order entry: Pharmacy perspectives. *California Journal of Health-System Pharmacy* 2004;16(1):4-6. Database: IPA.

Exclude - Not a Primary Study

Caliendo G, Meyer J, Cheng J. Computerized prescriber order entry (CPOE) facilitated restriction in improving nesiritide utilization. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Caliendo GC, Aggour T, Friedman TS, et al. JCAHO periodic performance review: A tool to improve medication safety. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Callahan CL, Manthey C, Neagos G. Putting sedation, analgesia and neuromuscular blockade guidelines to work for the adult ICU patient. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Database: IPA.

Exclude - Not a Primary Study

Callahan JJ, Donati EJ, Makooi-Morehead M. Integrating pharmacist interventions with computerized order entry. *ASHP Annual Meeting* 1992;49: Database: IPA.

Exclude - Not a Primary Study

Callen J, Braithwaite J, Westbrook J. Differences in doctors' and nurses' assessments of hospital culture and their views about computerised order entry systems. *Studies in Health Technology & Informatics* 2008;136:15-20. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Callen JL, Braithwaite J, Westbrook JI. Cultures in hospitals and their influence on attitudes to, and satisfaction with, the use of clinical information systems. *Soc Sci Med* 2007;65(3):635-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Callen JL, Braithwaite J, Westbrook JI. Contextual implementation model: a framework for assisting clinical information system implementations. *J Am Med Inform Assoc* 2008;15(2):255-62. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Calman N, Kitson K, Hauser D. Using information technology to improve health quality and safety in community health centers. *Progress in Community Health Partnerships* 2007;1(1):83-8. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=66732> Grey Lit.

Exclude - Not a Primary Study

Cals JW, Hopstaken RM, Le Doux PH, et al. Dose timing and patient compliance with two antibiotic treatment regimens for lower respiratory tract infections in primary care. *Int J Antimicrob Agents* 2008;31(6):531-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Calvano C, Vacovsky M. E-prescriptions: ready for prime time? Getting the regulations in place. *Computertalk for the Pharmacist* 2000;20(Sep-Oct):26-8. Database: IPA.

Exclude - Not a Primary Study

Calvo Alcantara MJ, Inesta GA. [The impact of an intervention strategy in the prescription of generic drugs in a primary care area]. [Spanish]. *Aten Primaria* 1999;23(7):419-24. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Canizares NJ, Diaz R, Gurriss LA, et al. Microcomputer decision support system for intensive care. *Studies in Health Technology & Informatics* 1998;52(Pt 1):534-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cansdale WC, Craven KH, Fey DW, et al. Implementation of a pharmacy-based, computer-assisted, concurrent antibiotic review service in a 688 bed acute tertiary care community hospital. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Cantor MN, Feldman HJ, Triola MM. Using trigger phrases to detect adverse drug reactions in ambulatory care notes. *Quality & Safety in Health Care* 2007;16(2):132-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Cantraine F. Computer driven i.v. injection systems. State of the art, future developments. *Acta Anaesthesiol Belg* 1988;39(4):257-66. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cantrill JA. The first 18 months of primary care groups: Pharmaceutical implications. *Pharmaceutical Journal* 2002;268(7190):407-9. Database: Embase Sept 22-09.

Exclude - Not MMIT

Capdevila C. Program of personalized system of dosing. *Circular Farmaceutica* 2002;60(4):20-2. Database: IPA.

Exclude - Unable to Retrieve Foreign

Carayon P, Wettemeck TB, Sobande F, et al. Assessing nurse interaction with medication administration technologies: the development of observation methodologies. *Work with Computer Systems* 2004;319-24. Grey Lit.

Exclude - Not MMIT

Cardona JM. More than drug reference tools: Using PDA's to improve workflow. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Carlstedt BC, Lukes AL, Overhage JM, et al. Use of a physician order-entry system to implement pharmacy department policies. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Carlstedt BC, Murray MD. Doctor of Pharmacy Clerkship Student Interaction with a Physician Order-Entry System. *American Association of Colleges of Pharmacy Annual Meeting* 2002;95(Jul):2 Database: IPA.

Exclude - Not MMIT

Carmenates J, Keith MR. Impact of automation on pharmacist interventions and medication errors in a correctional health care system. *Am J Health Syst Pharm* 2001;58(9):779-83. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Carmichael JM, Alvarez A, Chaput R, et al. Establishment and outcomes of a model primary care pharmacy service system. *Am J Health Syst Pharm* 2004;61(5):472-82. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Carnicero J, Blanco O, Mateos M. The application of information and communication technologies to clinical activity: Electronic health and clinical records. *Pharmaceuticals Policy & Law* 2006;8(1):69-82. Database: BSC.

Exclude - Not a Primary Study

Carnwath T. Switching on the work force: Computerizing a drug team. *Br J Addict* 1991;86(1):33-5. Database: PsycINFO.

Exclude - Not a Primary Study

Carol R. E-prescribing effects: Pilot project studies E-prescribing standards in long-term care. *Journal of the American Health Information Management Association* 2007;78(1):36-8. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Carpenter JD, Gorman PN. What's so special about medications: a pharmacist's observations from the POE study. *Proceedings / AMIA* 2001;95-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Carpenter JD, Gorman PN. Using medication list--problem list mismatches as markers of potential error. *Proceedings / AMIA* 2002;106-10. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Carpenter JD. Preparing for the CPOE environment: Pharmacists wanted! Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Carper J. Effect of automation on errors. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.

Exclude - Not a Primary Study

Carper JL, Barker KN, Flynn EA, et al. Methodology for determining how pharmacy automation decisions are made. Ashp Midyear Clinical Meeting 1998;33: Database: IPA.

Exclude - Not a Primary Study

Carper JL, Barker KN, Flynn EA, et al. Study of the factors that determine pharmacy automation purchasing decisions. Ashp Midyear Clinical Meeting 1998;33: Database: IPA.

Exclude - Not a Primary Study

Carpi V. When consistency counts. Health Manag Technol 2008;29(2):60-1.

http://findarticles.com/p/articles/mi_m0DUD/is_2_29/ai_n24265690/9984198

Database: Inspec.

Exclude - Not a Primary Study

Carr D, Dimitrakakis J. Explore all-encompassing electronic health records. Nurs Manag (Harrow) 2003;(Suppl):24-32. Database: BSC.

Exclude - Not a Primary Study

Carroll AE, Christakis DA. Pediatricians' use of and attitudes about personal digital assistants. Pediatrics 2004;113(2):238-42. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Carroll AE, Tarczy-Hornoch P, O'Reilly E, et al. The effect of point-of-care personal digital assistant use on resident documentation discrepancies. Pediatrics 2004;113(3):450-4.

Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Carroll CA. Technology and best Practices - How to measure the impact on patient safety. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Carroll MD, Wood PM, Temples JV, et al. Redesigning the order entry process to provide time for clinical training and pharmaceutical care. Ashp Midyear Clinical Meeting 1994;29: Database: IPA.

Database: IPA.

Exclude - Not a Primary Study

Carswell JL, DiPiro CV, Gomez TA, et al. Evaluation of turnaround time for medication order processing with use of a novel scanning system. Hosp Pharm 2006;41(3):249-53.

Database: Embase Sept 22-09.

Exclude - Not MMIT

Carter BL, Taylor JW, Becker A. Evaluation of three dosage-prediction methods for initial in-hospital stabilization of warfarin therapy. Clin Pharm 1987;6(1):37-45. Exclude - Not

MMIT

Carter BL, Barr W, Rock W, et al. Warfarin dosage predictions assisted by the analog computer. *Ther Drug Monit* 1988;10(1):69-73. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Carter BL, Farris KB, Abramowitz PW, et al. The Iowa Continuity of Care study: Background and methods. *Am J Health Syst Pharm* 2008;65(17):1631-42. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Carvalho MV, Boucherle D, Allenet B, et al. Economic impact of computerized prescriber order entry (CPOE) on drug costs. *Ashp Summer Meeting* 2005;62: Database: IPA.
Exclude - Not a Primary Study

Cash J. Computer alert fatigue analysis in a pediatric pharmacy information system. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Cassiani SH. Medication errors: prevention strategies [Portuguese]. *Rev Bras Enferm* 2000;53(3):424-30. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Cassiani SH, Gimenes FR, Freire CC. Evaluation of electronic drug prescriptions at a university hospital [Portuguese]. *Rev Bras Enferm* 2002;55(5):509-13. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Cassiani SH, Freire CC, Gimenes FR. Electronic medical prescription at a university hospital: writing failures and users' opinions. *Rev Esc Enferm USP* 2003;37(4):51-60. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Cassiani SHDB. Errors in medication: preventive strategies [Portuguese]. *Rev Bras Enferm* 2001;53(3):424-30. Database: CINAHL.
Exclude - Not MMIT

Cassiani SHDB, Gimenes FRE, Freire CC. Evaluation of computerized orders in an university hospital [Portuguese]. *Rev Bras Enferm* 2002;55(5):509-13. Database: CINAHL.
Exclude - Unable to Retrieve Foreign

Casteuble T. HDMA Foundation report surveys industry's barcode readiness. *Healthcare Distributors* 2003;55(4):15-7. Database: IPA.
Exclude - Not a Primary Study

Castine SD, Dolan C, Ardolino M, et al. Implementation of a physician order entry system: One hospital's experience. *ASHP Annual Meeting* 1996;53: Database: IPA.
Exclude - Not a Primary Study

Cataldo R, Jr. Nonprogramming relational database primer: a case study--pharmacist intervention audits. *Top Hosp Pharm Manage* 1991;11(3):29-51. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Caudill-Slosberg M, Weeks WB. Case study: identifying potential problems at the human/technical interface in complex clinical systems. *Am J Med Qual* 2005;20(6):353-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cavalcanti AB, Silva E, Pereira AJ, et al. A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *J Crit Care* 2009;24(3):371-8. Database: Embase Sept 22-09.

Exclude - Not MMIT

Cavallo J. One-click prescribing. *Community Oncology* 2005;2(3):229-30. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Cavan DA, Hejlesen OK, Hovorka R, et al. Preliminary experience of the DIAS computer model in providing insulin dose advice to patients with insulin dependent diabetes. *Computer Methods & Programs in Biomedicine* 1998;56(2):157-64. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Cavell GF, Hughes DK. Does computerized prescribing improve the accuracy of drug administration? *Pharmaceutical Journal* 1997;259:782-4. Database: IPA.

Exclude - Unable to Retrieve

Celeste R, Cusack BA. EPCglobal standards in the pharmaceutical industry: Toward a safe and secure supply chain. *Journal of Pharmacy Practice* 2006;19(4):244-9. Database: IPA.

Exclude - Not a Primary Study

Celler BG, Lovell NH, Basilakis J et al. Home telecare for chronic disease management. New South Wales Univ Sydney (Australia) School of Biomedical Engineering; 2001.

<http://handle.dtic.mil/100.2/ADA409950> 2002507266223

Database: Compendex.

Exclude - Not MMIT

Celler BG, Lovell NH, Basilakis J. Using information technology to improve the management of chronic disease. *Med J Aust* 2003;179(5):242-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Centers for Medicare & Medicaid Services (CMS). Medicare program; e-prescribing and the prescription drug program. Final rule. *Fed Regist* 2005;70(214):67567-95. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Centers for Medicare & Medicaid Services (CMS). Medicare program; physicians referrals to health care entities with which they have financial relationships; exceptions for certain electronic prescribing and electronic health records arrangements. Final rule. *Fed Regist* 2006;71(152):45139-71. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cepelewicz BB, Sastow GS. Electronic medical records: for many, a voyage into the unknown. *Medical Malpractice Law & Strategy* 2009;26(12):1-4.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010425075&site=ehost-live>; EBSCO CINAHL.

Exclude - Not a Primary Study

Cervi PL, Everitt AS. Automatic voice mail for delivering computer-generated anticoagulant dose advice to patients. *J Telemed Telecare* 2002;8(5):259-63. 7495974
Database: Inspec.

Exclude - Not MMIT

Cescon DW, Etchells E. Barcoded medication administration: a last line of defense. *JAMA* 2008;299(18):2200-2. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cesta A, Bajcar JM, Ong SW, et al. The EMITT study: development and evaluation of a medication information transfer tool. *Ann Pharmacother* 2006;40(6):1074-81. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Chaffee BW, Townsend KA, Benner T, et al. Pharmacy database for tracking drug costs and utilization. *Am J Health Syst Pharm* 2000;57(7):669-76. Database: IPA.

Exclude - Not a Primary Study

Chaffee BW. Developing and assessing requirements for clinical decision support. *Am J Health Syst Pharm* 2003;60(18):1875-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chaffee BW, Bonasso J. Strategies for pharmacy integration and pharmacy information system interfaces, Part 2: Scope of work and technical aspects of interfaces. *Am J Health Syst Pharm* 2004;61(5):506-14. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chaffee BW, Bonasso J. Strategies for pharmacy integration and pharmacy information system interfaces, Part 1: History and pharmacy integration options. *Am J Health Syst Pharm* 2004;61(5):502-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chaiken BP, Christian CE, Johnson L. Quality and efficiency successes leveraging IT and new processes. *J Healthc Inf Manag* 2007;21(1):48-53. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chamberlain MA. Vertically integrated pharmacy department. *Am J Health Syst Pharm* 1998;55(7):669-75. Database: IPA.

Exclude - Not a Primary Study

Champness A. Wireless computers lead the emancipation of medical workers. *Managed Healthcare* 2004;7(10):44-5. Database: IPA.

Exclude - Not a Primary Study

Chan A, Martins S, Coleman R et al. Post-fielding surveillance of a guideline-based decision support system. *AHRQ*: 2005. Grey Lit.

Exclude - No Outcomes of Interest

Chan AL, Tsai YF. Computerized system for reminding prescribers to monitor warfarin therapy. *Am J Health Syst Pharm* 2004;61(14):1436-7. Database: IPA.
Exclude - Not a Primary Study

Chan AL, Wang HY, Leung HW. Incorporation of a gentamicin dosage calculator into a computerized prescriber-order-entry system. *Am J Health Syst Pharm* 2006;63(14):1344-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Chan D. Cost-savings benefit of automation at a long term acute care hospital. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
Exclude - Not a Primary Study

Chan DS, Callahan CW, Hatch-Pigott VB, et al. Internet-based home monitoring and education of children with asthma is comparable to ideal office-based care: results of a 1-year asthma in-home monitoring trial. *Pediatrics* 2007;119(3):569-78. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Chan JT. Computerisation of accident and emergency departments in Hong Kong. *Hong Kong Medical Journal* 2000;J..(3):276-82. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Chan SL. Scanning and image processing system (SIPS) for medication ordering. *Journal of End User Computing* 2001;13(3):14-9. 2001276567360
Database: Compendex.
Exclude - Not a Primary Study

Chan SL. *Advanced topics in end user computing Vol.1*, Hershey, PA, USA:Idea Group Publishing;2002. End-user directed requirements-a case in medication ordering. 7438298
Database: Inspec.
Exclude - Not a Primary Study.

Chan W. Increasing the success of physician order entry through human factors engineering. *J Healthc Inf Manag* 2002;16(1):71-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Chandler MH. Introduction to clinical pharmacy software. *Ashp Midyear Clinical Meeting* 1991;26: Database: IPA.
Exclude - Not a Primary Study

Chang F. One-on-one with Florence Chang, SVP & CIO, MultiCare Health System. As MultiCare rolls out more applications and expands its award-winning EHR system, data storage is becoming a high priority. Interview by Kate Huvane Gamble. *Healthcare informatics : the business magazine for information and communication systems* 2009;26(12):22, 24-2, 25. OVID EMBASE.
Exclude - Not a Primary Study

Chang K, Kirkpatrick J, Criswell M, et al. Using Pocket PC and simulation to prepare students for safe and effective medication administration... 8th Annual International Nursing Simulation/Learning Resource Centers Conference, 10 June 2009 - 13 June 2009. *Clinical Simulation in Nursing* 2009;5(3):e134-e135

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010493651&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=3970&accno=2010493651

EBSCO CINAHL.

Exclude - Unable to Retrieve

Chant C, Wilson G, Friedrich JO. Validation of an insulin infusion nomogram for intensive glucose control in critically ill patients. *Pharmacotherapy* 2005;25(3):352-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Chapman SR. Developing decision support systems for nurse prescribers. *Nurse Prescribing* 2009;5(6):251-5. Database: CINAHL.

Exclude - No Outcomes of Interest

Chapman SR. Making decisions about prescribing in diabetes. *Practice Nursing* 2009;19(3):130-1. Database: CINAHL.

Exclude - Not MMIT

Chappel D, Fernandes V. Improving the coverage of neonatal BCG vaccination. *J Public Health Med* 1996;18(3):308-12. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Chast F, Brandely ML, Bardin C. The benefits of computerized anticancer drug distribution. *Bull Acad Natl Med* 2005;189(8):1721-32. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chatellier G, Colombet I, Degoulet P. An overview of the effect of computer-assisted management of anticoagulant therapy on the quality of anticoagulation. *Int J Med Inf* 1998;49(3):311-20. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chatellier G, Colombet I, Degoulet P. Computer-adjusted dosage of anticoagulant therapy improves the quality of anticoagulation. *Studies in Health Technology & Informatics* 1998;52 Pt 2:819-23. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chatman C. How Cloud Computing is Changing the Face of Health Care Information Technology. *Journal of Health Care Compliance* 2010;12(3):37-70.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=50320779&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Chaudhry B, Wang J, Wu S, et al. Systematic review: impact of health information technology on quality, efficiency, and costs of medical care. *Ann Intern Med* 2006;144(10):742-52. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chaudhry,B. The effect and adoption of electronic health records: A systematic review and national survey of physicians in the United States University of California, Los Angeles. 2007. Grey Lit.

Exclude - Not MMIT

Chedoe I, Molendijk HA, Dittrich ST, et al. Incidence and nature of medication errors in neonatal intensive care with strategies to improve safety: a review of the current literature. *Drug Saf* 2007;30(6):503-13. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chen CF, Yao ML, Wang SC, et al. A computerized antiinfective management program for antibiotics and other anti-infective agents in MMH. *International Pharmaceutical Federation World Congress* 2004;62:111 Database: IPA.

Exclude - No Outcomes of Interest

Chen DM, Wong KM, Boro MS. Outpatient provider order entry: Implementation process and a comparison to handwritten prescriptions. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Chen J, Shabot MM, LoBue M. A real time interface between a computerized physician order entry system and the computerized ICU medication administration record. *AMIA* 2003;810 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chen JL, Ng A, Yaghdjian V, et al. Pharmacy intervention reduced excessive antibiotic monitoring in the intensive care units. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Chen LM, Rein MS, Bates DW. Costs of quality improvement: a survey of four acute care hospitals. *Jt Comm J Qual Patient Saf* 2009;35(11):544-50. PMID:19947330 OVID MEDLINE.

Exclude - Not MMIT

Chen P, Tanasijevic MJ, Schoenenberger RA, et al. A computer-based intervention for improving the appropriateness of antiepileptic drug level monitoring. *Am J Clin Pathol* 2003;119(3):432-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Chen WC, Chang P, Chen LF. Developing a camera-phone-based drug barcode reader and support system. *AMIA* 2006;882 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chen WC. Effectiveness evaluation of bar code implementation in hospital -- an illustration of out-patient pharmacy information system. *AMIA* 2008;899 PMID:18998929 OVID MEDLINE.

Exclude - No Outcomes of Interest

Chen Y, Hsieh M, Wang C and others. RFID-based intelligent systems for home-healthcare. In Piscataway, NJ, USA: IEEE; 2007. 9529109

Database: Inspec.

Exclude - Not a Primary Study

Chen Y, Liu C, Wang C and others. RFID and IPv6-enabled ubiquitous medication error and compliance monitoring system. In Taipei, Taiwan: Inst. of Elec. and Elec. Eng. Computer Society; 2007. p.105-8.20074010849348

Database: Compendex.

Exclude - Not a Primary Study

Chester MI, Zilz DA. Effects of bar coding on a pharmacy stock replenishment system. Ashp Midyear Clinical Meeting 1988;23: Database: IPA.

Exclude - Not a Primary Study

Cheung LK, Gibson GA, Bourret JA. Outcome-oriented quality assurance program for the clinical pharmacokinetics monitoring service. ASHP Annual Meeting 1992;49: Database: IPA.

Exclude - Not a Primary Study

Cheung PP, Hauser WF. Automated unit dose dispensing in a state hospital. California Journal of Hospital Pharmacy 2008;3(Jul):8 Database: IPA.

Exclude - Not a Primary Study

Cheverry C, Crepeaux R, Le Verge R. Help to therapeutics by a daily computer monitoring of potential drug interactions. Journal de Pharmacie Clinique 1982;1(1):49-61. Database: IPA.

Exclude - Unable to Retrieve Foreign

Cheverst K, Clarke K, Dewsbury G, et al. Designing assistive technologies for medication regimes in care settings. Universal Access in the Information Society 2003;2(3):235-42. 8396669

Database: Inspec.

Exclude - Not a Primary Study

Chi J. Physician order entry: Is it something for you? Hospital Pharmacist Report 2004;14(Feb):20 Database: IPA.

Exclude - Not a Primary Study

Chiarelli F, Tumini S, Morgese G, et al. Controlled study in diabetic children comparing insulin-dosage adjustment by manual and computer algorithms. Diabetes Care 1990;13(10):1080-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Chiarelli F, Di Ricco L, Catino M, et al. Modern management of childhood diabetes: a role for computerized devices? Acta Paediatr Jpn 1998;40(4):299-302. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chien YW, Lin SS. Optimisation of treatment by applying programmable rate-controlled drug delivery technology. Clin Pharmacokinet 2002;41(15):1267-99. Database: IPA.

Exclude - Not a Primary Study

Childs SA, Poikonen J. Converting from manual to computerized medication administration records. *Am J Hosp Pharm* 1993;50(7):1420-2. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chin HL, Krall MA. Successful implementation of a comprehensive computer-based patient record system in Kaiser Permanente Northwest: strategy and experience. *Eff Clin Pract* 1998;1(2):51-60. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chiu CC, Lee S, Boro MS. Implementation of online drug use criteria to facilitate nonformulary drug requests. *Ashp Midyear Clinical Meeting* 2001;36. Database: IPA.

Exclude - Not a Primary Study

Chiu MF, Leone DM, Timoney J, et al. Implementation of maximum dose information of the physician order entry system. *Ashp Midyear Clinical Meeting* 1999;34. Database: IPA.

Exclude - Not a Primary Study

Chizzali-Bonfadin C, Adlassnig KP, Koller W. MONI: an intelligent database and monitoring system for surveillance of nosocomial infections. *Medinfo* 1995;8p2:1684

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Choi H, McCarthy MW. Assessment of adherence to a tiotropium therapeutic interchange. *Ashp Midyear Clinical Meeting* 2006;41. Database: IPA.

Exclude - Not a Primary Study

Chou LS, Fichtl RE, Jordan F. Development of an antineoplastic therapy ordering process in a computerized physician order entry (CPOE) system. *Ashp Midyear Clinical Meeting* 2007;42. Database: IPA.

Exclude - Not a Primary Study

Chow CL, Chim TS. Pharmacy formulary maintenance: online approach. *ASHP Annual Meeting* 1993;50. Database: IPA.

Exclude - Not a Primary Study

Chrischilles EA, Fulda TR, Byrns PJ, et al. The role of pharmacy computer systems in preventing medication errors. *J Am Pharm Assoc (Wash)* 2002;42(3):439-48. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Christensen A, Christrup LL, Fabricius PE, et al. Survey of patient and physician assessment of a compliance reminder device in the treatment of hypertension. *Blood Press* 2009;18(5):280-5. OVID MEDLINE.

Exclude - Not MMIT

Christensen CM, Hwang J. A Disruptive Solution for Health Care. *BusinessWeek Online* 2009;22

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=36806151&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Christie P, Robinson H. Using a communication framework at handover to boost patient outcomes. *Nurs Times* 2009;105(47):13-5. PMID:20063616 OVID MEDLINE.

Exclude - Not a Primary Study

Chrymko MM, Scherpbier HJ. Computerized pharmacokinetics on an institutional clinical information system. *Pharm Pract Manag Q* 1995;15(3):36-43. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chu JJ. Preventing postoperative venous thromboembolism. *AJN, American Journal of Nursing* 2004;104: Database: IPA.

Exclude - Not a Primary Study

Chu S. ePrescription: road map from wired to wireless point-of-care order entry. In Piscataway, NJ, USA: IEEE; 2004. p.26-33.8127582

Database: Inspec.

Exclude - Not a Primary Study

Chu S. Non-intrusive guideline-based electronic disease management programme: Principles and evaluation of a pilot. *Journal on Information Technology in Healthcare* 2004;2(4):263-80. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Chua P. Reducing drug packing errors using bar code technology and visual aids. *Ashp Summer Meeting* 2009;65(Jun): Database: IPA.

Exclude - Not a Primary Study

Chuang CT. An efficient fault-tolerant order entry management information system based on special distributed client/server architecture. *Health Serv Manage Res* 1998;11(4):255-64.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chuang CT. An efficient fault-tolerant out-patient order entry system based on special distributed client/server architecture. *Med Inform (Lond)* 1998;23(2):145-57. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Chuang M-H. Computerized medication prescribing errors in hospital settings. *Tzu Chi Medical Journal* 2003;15(5): Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Chuang Y, Tan R, Wu M, et al. An empirical study of a computerized management system in ambulatory services at Chang Gung Memorial Hospital in Taiwan. *International Journal of Technology Management* 1995;74-92. 5109989

Database: Inspec.

Exclude - Unable to Retrieve

Chuang Y, Tan R, Wu M, et al. Managing information technology: an empirical study of a computerized management system in ambulatory services at Chang Gung Memorial Hospital in Taiwan. *International Journal of Computer Applications in Technology* 1996;9(4):181-92. 5447679

Database: Inspec.

Exclude - Unable to Retrieve

Chung E, Caliendo G, Paxos P, et al. The impact of pharmacy services in a neurosurgical intensive care unit. *Ashp Midyear Clinical Meeting 2002*;37: Database: IPA.
Exclude - Not a Primary Study

Chung K, Choi YB, Moon S. Toward efficient medication error reduction: error-reducing information management systems. *J Med Syst 2003*;27(6):553-60. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Churchill WW. The glory and chaos of bar-coding. *Ashp Midyear Clinical Meeting 2004*;39: Database: IPA.
Exclude - Not a Primary Study

Churchill WW. Evaluation and selection of IV “smart pump” technology. *Ashp Midyear Clinical Meeting 2005*;40: Database: IPA.
Exclude - Not a Primary Study

Cimino J J, Bright T J, Li J. Medication reconciliation using natural language processing and controlled terminologies. In *MEDINFO*; 2007. p.679-83. Grey Lit.
Exclude - Not MMIT

Cina J, Fanikos J, Mitton P, et al. Medication errors in a pharmacy-based bar-code-repackaging center. *Am J Health Syst Pharm 2006*;63(2):165-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Cioffi M, Shaw A, Robitaille P, et al. Impact of barcoding on the incidence of adverse drug events. *Ashp Midyear Clinical Meeting 1996*;31: Database: IPA.
Exclude - Not a Primary Study

Ciotti V. Great-grandfather of CPOE. *Healthc Inform 2009*;26(9):56
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010429786&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1239&accno=2010429786
EBSCO CINAHL.
Exclude - Not a Primary Study

Cipriano PF. A nursing perspective on bedside scanning systems. *Hosp Pharm 2003*;38(11):S14-S15 Database: IPA.
Exclude - Unable to Retrieve

Claes N, Buntinx F, Vijgen J, et al. The Belgian improvement study on oral anticoagulation therapy: A randomized clinical trial. *Eur Heart J 2005*;26:2159-65. Exclude - Not MMIT

Claes N, Moeremans K, Frank B, et al. Estimating the cost-effectiveness of quality-improving interventions in oral anticoagulation management within general practice. *Value in Health 2006*;9(6):369-76. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Clancy TR, Delaney C, Segre A, et al. Predicting the impact of an electronic health record on practice patterns using computational modeling and simulation. *AMIA Annu Symp Proc 2007*;145-9. Database: CINAHL.
Exclude - No Outcomes of Interest

Clark C. Information technology in action. Hospital Pharmacist 2002;9(4): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Clark I, Peters M. Mobile real-time computer links in a hospital environment. BJHC & IM 1997;14(6):27-9. Database: CINAHL.

Exclude - Not MMIT

Clark T. Computerized physician order entry prescribing events impeded by pharmacists. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.

Exclude - Not a Primary Study

Clark T. CPOE-medication system strategy. Ashp Summer Meeting 2002;59: Database: IPA.

Exclude - Not a Primary Study

Clark T. Review of CPOE systems and the EMR. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Classen D. Leapfrog computerized physician order entry (CPOE) and electronic health record (EHR) evaluation tools: Project overview and discussion. AHRQ; 2007. http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_219414_0_0_18/AHRQ%20Webcast%2011207.pdf Grey Lit.

Exclude - Not a Primary Study

Classen DC, Pestotnik SL, Evans RS, et al. Computerized surveillance of adverse drug events in hospital patients. JAMA 1991;266(20):2847-51. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Classen DC, Metzger J. Improving medication safety: the measurement conundrum and where to start. Int J Qual Health Care 2003;15(suppl 7):i41-i47 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Classen DC, Avery AJ, Bates DW. Evaluation and certification of computerized provider order entry systems. J Am Med Inform Assoc 2007;14(1):48-55. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Clauson KA, Marsh WA, Polen HH, et al. Clinical decision support tools: analysis of online drug information databases. BMC Med Inform Decis Mak 2007;7:7 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Clemmer TP. Computers in the ICU: where we started and where we are now. J Crit Care 2004;19(4):201-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cleveringa FGW. Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. Diabetes Care 2009;31(12):2273-5. Database: Embase Sept 22-09.

Exclude - Not MMIT

Clifton-Koeppel R. What nurses can do right now to reduce medication errors in the neonatal intensive care unit. *Newborn & Infant Nursing Reviews* 2008;8(2):72-82. Database: CINAHL.

Exclude - Not a Primary Study

Climent C. Impact of implementation a unit-dose system and computer-assisted prescribing on medication errors. *Atencion Farmaceutica* 2008;10(5): Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Cloud-Buckner J, Gallimore J J, Wong P K. Issues in alerting: Medication order entry in real practice. In Heraklion, Greece: 2008. p.703-7. Grey Lit.

Exclude - No Outcomes of Interest

Coddington DC, Moore KD. Leading IDSs heed the call to invest in IT. *Healthc Financ Manage* 2002;56(2):36-40. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Coe F, Norton E, Oparil S, et al. Treatment of hypertension by computer and physician - A prospective controlled study. *J Chronic Dis* 1977;30:81-92. Exclude - Not MMIT

Cohen B, Flinn M, Koontz J, et al. Robotic system: Centralized dispensing model. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Cohen H. Be a watchdog. *Nurs Manag (Harrow)* 2000;31(12):53-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cohen H. Protecting patients from harm: reduce the risks of high-alert drugs. *Nursing (Lond)* 2001;37(9):49-55. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cohen M. Are you ready to take a giant leap forward? Leapfrog: New initiative to help employee benefit plans adopt patient safety standards. *Hosp Pharm* 2000;35(9):907

Database: IPA.

Exclude - Not MMIT

Cohen MR. Drug product characteristics that foster drug-use-system errors. *Am J Health Syst Pharm* 1995;52:395-9. Database: IPA.

Exclude - Not MMIT

Cohen MR. Medication errors associated with automated dispensing modules. *Hosp Pharm* 1996;31(8):930-3. Database: IPA.

Exclude - Not MMIT

Cohen MR. Errors with Hespan IV bags. *Hosp Pharm* 1998;33(Apr):393-4. Database: IPA.

Exclude - Not MMIT

Cohen MR. Update on health care purchaser initiatives to improve patient safety. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Cohen MR. Don't lose track of orders during clarification. *Hosp Pharm* 2003;38(12):1120-1.
Database: IPA.
Exclude - Not MMIT

Cohen MR. Medication errors. Bar code blues: consider the source. *Nursing (Lond)* 2009;39(12):15
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010506137&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=280&accno=2010506137 EBSCO CINAHL.
Exclude - Not a Primary Study

Cohen MR. Medication errors. *Nursing (Lond)* 2010;40(2):17
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010550602&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=280&accno=2010550602 EBSCO CINAHL.
Exclude - Not a Primary Study

Cohen MR. Medication errors. Fixing electronic errors: allergic to change. *Nursing (Lond)* 2010;40(5):17
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010643453&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=280&accno=2010643453 EBSCO CINAHL.
Exclude - Not a Primary Study

Cohen MR. Medication errors. New alert system: attention-getter. *Nursing (Lond)* 2010;40(4):20
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010612421&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=280&accno=2010612421 EBSCO CINAHL.
Exclude - Not a Primary Study

Cohen S N, Armstrong M F, Briggs R L and others. Computer-based monitoring and reporting of drug interactions. In Amsterdam, Netherlands: North-Holland; 1974. p.889-94.1006962
Database: Inspec.
Exclude - No Outcomes of Interest

Cohn KH, Berman J, Chaiken B, et al. Engaging physicians to adopt healthcare information technology. *J Healthc Manag* 2009;54(5):291-300. OVID EMBASE.
Exclude - No Outcomes of Interest

Coiera E, Westbrook J, Wyatt J. The safety and quality of decision support systems. *Yearbook of Medical Informatics* 2006;(1):20-5. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Col NF. Using Internet technologies to improve and simplify counseling about menopause: The WISDOM website. *Maturitas* 2007;57(1):95-9. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Colella SJ. Quantifying the impact of a clinical pharmacy program to support investments in automation technology. *Ashp Midyear Clinical Meeting* 1991;26. Database: IPA.
Exclude - Not a Primary Study

Coleman B. Hospital pharmacy staff attitudes towards automated dispensing before and after implementation. *Hospital Pharmacist* 2004;11(6):248-51. Database: IPA.
Exclude - Not MMIT

Coleman H, Ashcraft K. Genelex Corporation. *Pharmacogenomics* 2008;9(4):469-75.
Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Coleman JJ, Ferner RE, Evans SJ. Monitoring for adverse drug reactions. *Br J Clin Pharmacol* 2006;61(4):371-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Coleman RL, Sanderson IC, Lubarsky DA. Anesthesia information management systems as a cost containment tool. *CRNA: the Clinical Forum for Nurse Anesthetists* 1997;8(2):77-83.
Database: CINAHL.
Exclude - Not a Primary Study

Coleman RW. Translation and interpretation: the hidden processes and problems revealed by computerized physician order entry systems. *J Crit Care* 2004;19(4):279-82. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Colen HB, Neuenschwander M, Neef C, et al. Using automated dispensing machines to improve medication safety. *Ejhp Science* 2006;12(5):71-3. Database: IPA.
Exclude - Not a Primary Study

Coley SC, Johnson VB, Unachukwu EO, et al. Implementation of injectable potassium chloride policy. *Ashp Midyear Clinical Meeting* 1997;32. Database: IPA.
Exclude - Not a Primary Study

Collin S, Reeves BC, Hendy J, et al. Implementation of computerised physician order entry (CPOE) and picture archiving and communication systems (PACS) in the NHS: quantitative before and after study. *BMJ: British Medical Journal* 2008;337(7670):622-5.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010073124&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=694&accno=2010073124 EBSCO CINAHL.
Exclude - Not MMIT

Collins CD, Pedersen CA, Schneider PJ, et al. The effect of a medication rules engine in Computerized Prescriber Order Entry (CPOE) on appropriate prescribing of Amphotericin B Lipid Complex (ABLC). *Ashp Midyear Clinical Meeting* 2003;38. Database: IPA.
Exclude - Not a Primary Study

Collins PS. Negotiating for clinical IT dollars: lessons learned. *Nurs Adm Q* 2007;31(4):300-3. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Collura J, Clark T, Grider L, et al. Quantification and evaluation of pharmacist computer medication order entry errors. *ASHP Annual Meeting* 2000;57. Database: IPA.
Exclude - Not a Primary Study

Colombet I, Bura-Riviere A, Chatila R, et al. Personalized versus non-personalized computerized decision support system to increase therapeutic quality control of oral anticoagulant therapy: an alternating time series analysis. *BMC Health Serv Res* 2004;4(1):27 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Colon-Emeric CS, Schmader KE, Twersky J, et al. Development and pilot testing of computerized order entry algorithms for geriatric problems in nursing homes. *J Am Geriatr Soc* 2009;57(9):1644-53. OVID EMBASE.

Exclude - Not MMIT

Colpaert K, Decruyenaere J. Computerized physician order entry in critical care. *Best Practice & Research* 2009;23(1):27-38. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Colvin JR. Microcomputer-controlled administration of vasodilators following cardiac surgery: Technical considerations. *J Cardiothorac Anesth* 1989;3(1):10-5. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Colvin L, Slack-Smith L, Stanley FJ, et al. Pharmacovigilance in pregnancy using population-based linked datasets. *Pharmacoepidemiology & Drug Safety* 2009;18(3):211-25. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Condella F, Lawlor H. Developing a pharmacy application using IBM's Patient Care System (PCS). *Hosp Pharm* 1987;22(4):341-53. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Conklin MH, Culley EJ, O'Donnell J. Case study of the effects of office-based generic drug sampling on antibiotic drug costs and first-line antibiotic prescribing ratios. *J Manag Care Pharm* 2009;15(1):55-61. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Conlin PR, Alexis G. E-mail system for obtaining approval to use a restricted agent. *Am J Health Syst Pharm* 1999;56(24):2535-7. Database: IPA.

Exclude - Not a Primary Study

Conn J. Enormous push for e-Rx: stimulus act gives vendors extra motivation to adopt. *Mod Healthc* 2009;39(29):14-5.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010353838&site=ehost-live>; EBSCO CINAHL.

Exclude - Not a Primary Study

Conn J. Patient safety, IT center stage. *Mod Healthc* 2009;39(41):18

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010441025&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvic?jid=772&accno=2010441025 EBSCO CINAHL.

Exclude - Not a Primary Study

Conn J. Privacy, please: IT, Congress urge HHS to add harm threshold. Mod Healthc 2009;39(44):12
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010469629&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010469629 EBSCO CINAHL.

Exclude - Not a Primary Study

Conn J. Safe and secure? Data encryption just one option under security law. Mod Healthc 2009;39(19):30-1.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010281692&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010281692 EBSCO CINAHL.

Exclude - Not a Primary Study

Conn J. Time and money: providers have issues with stimulus act's deadlines, definitions for IT. Mod Healthc 2009;39(27):14-5.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010326372&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010326372 EBSCO CINAHL.

Exclude - Not a Primary Study

Conn J. Looser IT subsidy rules? Panel advises relaxing requirements to qualify. Mod Healthc 2010;40(8):16
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010569743&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010569743 EBSCO CINAHL.

Exclude - Not a Primary Study

Conn J. Meaningful obtuse? Experts say proposed rule might be too ambitious. Mod Healthc 2010;40(2):8-9.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010524527&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010524527 EBSCO CINAHL.

Exclude - Not a Primary Study

Conn J. PHRs, portals both popular and not so popular. Mod Healthc 2010;40(9):34
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010571972&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010571972 EBSCO CINAHL.

Exclude - Not a Primary Study

Connors GP, Hays DP. Emergency department drug orders: does drug storage location make a difference? Ann Emerg Med 2007;50(4):414-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Conroy S, Sweis D, Planner C, et al. Interventions to reduce dosing errors in children: a systematic review of the literature. Drug Saf 2007;30(12):1111-25. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cook A, Schattner P, Pleteshner C. The experiences of one divisional group of GPs in introducing computers into clinical practice. *Aust Fam Physician* 1999;28(9):971-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Cook RL, Larrabee SE. Medication error: Problem to be solved. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.
Exclude - Not a Primary Study

Cooley TW, Szeto K, Churchill W, et al. Development and implementation of bar code technology in the medication use process of a new web-based CPOE-Pharmacy-EMAR system. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Cooper-DeHoff R, Handberg E, Heissenberg C, et al. Electronic prescribing via the internet for a coronary artery disease and hypertension megatrial. *Clin Cardiol* 2001;24(S5):V14-V16 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Coopmans VC. Certified registered nurse anesthetist performance and perceptions: Use of a handheld, computerized, decision making aid during critical events in a high-fidelity human simulation environment. *Virginia Commonwealth University* 2005; Database: CINAHL.
Exclude - Theses

Coopmans VC, Biddle C. CRNA performance using a handheld, computerized, decision-making aid during critical events in a simulated environment: a methodologic inquiry. *AANA J* 2008;76(1):29-35. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Copen J. Systematic review on PDA clinical application implementation and lessons learned. *Journal on Information Technology in Healthcare* 2008;6(2):114-28. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Corbo TE, Kleban S, Ewen EF. Implementation of computerized rules without computerized prescriber order entry (CPOE). *Ashp Summer Meeting* 2004;61: Database: IPA.
Exclude - Not a Primary Study

Corbo TE, Kleban S, Watson B, et al. We've got your back: Use of computerized decision support to minimize the risk of spinal hematoma. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Corcoran DK, Anderson RA. Smart cards: Nurses at the forefront of radical advances in health care. *Policy* 2001;2(1):39-46. Database: CINAHL.
Exclude - Not MMIT

Cordingley JJ, Vlasselaers D, Dormand NC, et al. Intensive insulin therapy: enhanced Model Predictive Control algorithm versus standard care. *Intensive Care Med* 2009;35(1):123-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Cordioli AV. SETA: A computer program aiding pharmacological treatment of affective disorders. *Revista ABP-APAL* 1995;16(2): Database: PsycINFO.

Exclude - Not MMIT

Corey C and Grossman J. Clinical information technology adoption varies across physician specialties. Center for Studying Health System Change; 2007.

<http://www.hschange.com/CONTENT/945/?words=electronic+prescribing> Grey Lit.

Exclude - Not MMIT

Corley ST. Electronic prescribing: a review of costs and benefits. *Top Health Inf Manage* 2003;24(1):29-38. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Corte Garcia JJ, Espin FM, Rojo EJ. Inappropriate prescribing in hospitalized elderly patients according to the beers criteria. [Spanish]. *Atencion Farmaceutica* 2009;11(4):261-6. OVID EMBASE.

Exclude - Unable to Retrieve

Cote J, Godin G, Garcia PR, et al. Program development for enhancing adherence to antiretroviral therapy among persons living with HIV. *Aids Patient Care STDS* 2008;22(12):965-75. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cotter CM. Computerized physician order entry from a chief information officer perspective. *J Crit Care* 2004;19(4):283-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cotter SA, Malone R. Improving pharmacist-documented interventions at a community hospital through education and computer generated medication warnings. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Coulter DM. Privacy issues and the monitoring of sumatriptan in the New Zealand Intensive Medicines Monitoring Programme. *Pharmacoepidemiology and Drug Safety* 2001;10(7):663-7. Database: Embase Sept 22-09.

Exclude - Not MMIT

Council on Clinical Information Technology. Pediatric aspects of inpatient health information technology systems. *Pediatrics* 2008;122(6):e1287-e1296 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Courtenay M, Carey N. Completeness of prescriptions issued to dermatology patients. *Nurse Prescribing* 2009;6(9):391-6. Database: CINAHL.

Exclude - Not MMIT

Coussaert EJ, Cantraine FR. Management software for a universal device communication controller: application to monitoring and computerized infusions. *Int J Clin Monit Comput* 1996;13(4):225-33. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cox PM, Jr., D'Amato S, Tillotson DJ. Reducing medication errors. *Am J Med Qual* 2001;16(3):81-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Cox S, Wilcock P, Young J. Improving the repeat prescribing process in a busy general practice. A study using continuous quality improvement methodology. *Qual Health Care* 1999;8(2):119-25. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Coyle GA. Bar code medication administration: Five deal breakers, pearls and perils. *Hosp Pharm* 2003;38(11):S20-S21 Database: IPA.
Exclude - Not MMIT

Coyle GA, Heinen M. Evolution of BCMA within the Department of Veterans Affairs. *Nurs Adm Q* 2005;29(1):32-8. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Craghead RM, Wartski DM. Effect of computerized prescribing on noncompliance rates. *Ashp Midyear Clinical Meeting* 1988;23: Database: IPA.
Exclude - Not a Primary Study

Craghead RM, Wartski DM. Evaluative study of unclaimed prescriptions. *ASHP Annual Meeting* 1989;46(10):310-2. Database: IPA.
Exclude - Not a Primary Study

Craig A. Electronic prescribing. *Adv Nurse Pract* 2007;15(4):22 PMID:19998953 OVID MEDLINE.
Exclude - Not a Primary Study

Craig JA, Eves GB. Minimizing drug misuse among elders: a proposal. *Public Health Rep* 1987;102(1):86-90. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Craig RP. Online adjudication of third party prescription claims: Technological revolution. *California Pharmacist* 1989;37(Oct):45-7. Database: IPA.
Exclude - Not a Primary Study

Crain LJ. Multidisciplinary approach to implementing physician order entry. *Ashp Midyear Clinical Meeting* 1999;34: Database: IPA.
Exclude - Not a Primary Study

Cramer J. Medicine partnerships. *Heart* 2003;89(Suppl 2):19-21. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Crane J, Crane FG. Preventing medication errors in hospitals through a systems approach and technological innovation: a prescription for 2010. *Hosp Top* 2006;84(4):3-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Crawford SY. ASHP national survey of hospital-based pharmaceutical services--1990. *Am J Hosp Pharm* 1990;47(12):2665-95. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Crawford SY, Myers CE. ASHP national survey of hospital-based pharmaceutical services--1992. *Am J Hosp Pharm* 1993;50(7):1371-404. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Crawford S, Grussing P, Clark T, et al. Staff attitudes about the use of robots in pharmacy before implementation of a robotic dispensing system. *Am J Health Syst Pharm* 1998;55(18):1907-14. Exclude - Not MMIT

Creasman GL, Dervin DA, Widdowson NF, et al. Barcode medication administration (BCMA): Quality management and medication error evaluation following implementation. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.
Exclude - Not a Primary Study

Cresswell KM. Adverse drug events in the elderly. *Br Med Bull* 2007;83(1):259-74. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Cresswell KM, Sheikh A. Information technology-based approaches to reducing repeat drug exposure in patients with known drug allergies. *Journal of Allergy & Clinical Immunology* 2008;121(5):1112-7. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Creswick N, Westbrook JI. The medication advice-seeking network of staff in an Australian hospital renal ward. *Studies in Health Technology & Informatics* 2007;130:217-31. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Creus N, Masso J, Nigorra M, et al. Computerized chemotherapy physician order entry to prevent medication errors. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Crisp A, Schomberg R. Assessment of the treatment of excessive anticoagulation in internal medicine patients. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.
Exclude - Not a Primary Study

Crittenden W. Drug cost savings associated with repackaging for use in an automated dispensing system: retrospective analysis. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Cronk JD. Digital scanning and consolidated entry of medication orders in a multihospital health system. *Am J Health Syst Pharm* 2002;59(8):731-3. Database: CINAHL.
Exclude - Not a Primary Study

Cross RK. Acceptance of telemanagement is high in patients with inflammatory bowel disease. *J Clin Gastroenterol* 2006;40(3): Database: Embase Sept 22-09.
Exclude - Not MMIT

Crossno CL. Using CPOE to improve communication, safety, and policy compliance when ordering pediatric chemotherapy. *Hosp Pharm* 2007;42(4):368-73. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Crosswhite R, Beckham SH, Gray P, et al. Using a multidisciplinary automated discharge summary process to improve information management across the system. *Am J Manag Care* 1997;3(3):473-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Crouch MA, Gramling R. Family history of coronary heart disease: evidence-based applications. *Prim Care* 2005;32(4):995-1010. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Crowther MA. Oral anticoagulant initiation: rationale for the use of warfarin dosing nomograms. *Seminars in Vascular Medicine* 2003;vasc.(3):255-60. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cruz-Correia R, Fonseca J, Lima L, et al. Web-based or paper-based self management tools for asthma - Patients' opinions and quality of data in a randomised crossover study. *Stud Health Technol Inform* 2007;127:178-89. Database: Embase Sept 22-09.

Exclude - Not MMIT

Cruz-Correia R, Fonseca J, Lima L, et al. Web-based or paper-based self-management tools for asthma--patients' opinions and quality of data in a randomized crossover study. *Studies in Health Technology & Informatics* 2007;127:178-89. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Cuellar Monreal MJ, Planells HC, Hernandez Fernandez de las Rojas MD, et al. Designing a module for the prevention of hypersensitivity reactions in an assisted electronic prescription system. *Farmacia Hospitalaria* 2005;HOSP..(4):241-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cummings J, Bush P, Smith D, et al. Bar-coding medication administration overview and consensus recommendations. *Am J Health Syst Pharm* 2005;62(24):2626-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Cunningham MC, Basile SB, Timmons VL, et al. Categorizing errors on the medication administration record. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Curioso WH, Kurth AE, Cabello R, et al. Usability evaluation of Personal Digital Assistants (PDAs) to support HIV treatment adherence and safer sex behavior in Peru. *AMIA Annual Symposium Proceedings* 2008; PMID:18999317 OVID MEDLINE.

Exclude - Not MMIT

Curtis C, Ford NG. Paperless electronic prescribing and medicines administration in a district general hospital. *Pharmaceutical Journal* 1997;259:734-5. Database: IPA.

Exclude - Unable to Retrieve

Curtis C, Marriott J, Langley C. Development of a prescribing indicator for objective quantification of antibiotic usage in secondary care. *J Antimicrob Chemother* 2004;54(2):529-33. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Cusack CM. Electronic health records and electronic prescribing: promise and pitfalls. *Obstetrics & Gynecology Clinics of North America* 2008;35(1):63-79. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Cusack CM, Byrne CM, Hook JM et al. Health information technology evaluation toolkit: 2009 Update. 09-0083-EF. AHRQ Publication; 2009.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_875888_0_0_18/09_0083_EF.pdf Grey Lit.
 Exclude - Not a Primary Study

Czech TR. Effective adverse drug reaction reporting as a measure of the impact of the pharmacist in a divorced setting. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
 Exclude - Not a Primary Study

D'Arcy Y. Using technology to help alleviate pain. *Nurs Manag (Harrow)* 2004;35(11):45-6. Database: BSC.
 Exclude - Not a Primary Study

Da Silva RV, Rivkin S. Increasing efficiency through point-of-prescribing technology. *Drug Benefit Trends* 2000;12(9):45-9. <http://www.medscape.com/viewarticle/409957> Database: Embase Sept 22-09.
 Exclude - Not MMIT

Dace LN. Bar codes in the hospital pharmacy. *Computertalk for the Pharmacist* 1990;10(May-Jun):6-7. Database: IPA.
 Exclude - Not a Primary Study

Dailey TL, Kozakiewicz JM. Failure mode and effects analysis for the medication use process of chemotherapy regimens. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
 Exclude - Not a Primary Study

Dailey TL. Evaluation of surgical antimicrobial prophylaxis (SAP) guidelines in a 511 bed community teaching hospital. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
 Exclude - Not a Primary Study

Dale J, Caramlau I, Docherty A, et al. Telecare motivational interviewing for diabetes patient education and support: A randomised controlled trial based in primary care comparing nurse and peer supporter delivery. *The Cochrane Central Register of Controlled Trials* 2007;8(18): <http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/305/CN-00641305/frame.html> Database: Cochrane.
 Exclude - Not MMIT

Dalen E, Wideroe TE, Hetlevik I, et al. Blood pressure records in general practice-- expectations and reality. *Tidsskr Nor Laegeforen* 2001;121(2):158-61. Database: Ovid MEDLINE(R).
 Exclude - Unable to Retrieve Foreign

Dalton MF, Klipfel L, Carmichael K. Safety issues: Use of continuous subcutaneous insulin infusion (CSII) pumps in hospitalized patients. *Hosp Pharm* 2006;41(10):956-69. Database: Embase Sept 22-09.
 Exclude - Not MMIT

Danekas P, Borchert KC, Kroeger R, et al. Implementation of medication reconciliation with an electronic medical record. *ASHP Midyear Clinical Meeting* 2005;40: Database: IPA.
Exclude - Not a Primary Study

Daniels BR, Thielke TS, Skibinski KA. Point of care systems, closing the medication use loop. *ASHP Annual Meeting* 1995;52: Database: IPA.
Exclude - Not a Primary Study

Daniels SD, Hood JC, Jones LJ, et al. Incorporation of enteral supplements into pharmacy drug/drug interaction program to detect potential therapeutic problems. *ASHP Midyear Clinical Meeting* 1991;26: Database: IPA.
Exclude - Not a Primary Study

Danielson, E. Evaluating the impact of implementing clinical information systems The University of Utah Editor. 2006. Grey Lit.
Exclude - Not MMIT

Danne T, von Schutz W, Lange K, et al. Current practice of insulin pump therapy in children and adolescents - The Hannover recipe. *Pediatric Diabetes* 2006;7(Suppl. 4):25-31. Database: Embase Sept 22-09.
Exclude - Not MMIT

Darby PW. Bar code technology in inventory control. *Can J Hosp Pharm* 1988;41(6):313-6. Database: IPA.
Exclude - Not MMIT

Das BP, Rauniar GP, Bhattacharya SK. Medical errors challenges for the health professionals: need of Pharmacovigilance to prevent. *JNMA Journal of the Nepal Medical Association* 2006;45(162):273-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Dasta JF. Computers in critical care: opportunities and challenges. *DICP* 1990;24(11):1084-92. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Dasta JF, Greer ML, Speedie SM. Computers in healthcare: overview and bibliography. *Ann Pharmacother* 1992;26(1):109-17. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Dasta JF, Reardon G. Survey of ACCP members regarding use of computers and information processing. *Pharmacotherapy* 1992;12(6):468-74. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Davenport TH, Glaser J. Just-in-time delivery comes to knowledge management. *Harv Bus Rev* 2002;80(7):107-11. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Davey P. The potential role of computerized decision support systems to improve empirical antibiotic prescribing. *J Antimicrob Chemother* 2006;58(6):1105-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

David Y. Telehealth: Current practices and future directions. In Philadelphia, PA, USA: Society of Photo-Optical Instrumentation Engineers; 1996. p.2-8.1996263115939
Database: Compendex.
Exclude - Not MMIT

Davidson PC, Steed RD, Bode BW. Glucomander: a computer-directed intravenous insulin system shown to be safe, simple, and effective in 120,618 h of operation. Diabetes Care 2005;28(10):2418-23. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Davidson SM, Heineke J. Toward an effective strategy for the diffusion and use of clinical information systems. J Am Med Inform Assoc 2007;14(3):361-7. Database: IPA.
Exclude - Not a Primary Study

Davies A. Making the connections: The role of read codes as a linkage mechanism for drug codes. BJHC & IM 1996;13(1):18-20. Database: CINAHL.
Exclude - Not MMIT

Davies EA. Multidisciplinary approach to implementing physician order entry: Physician overview. Ashp Midyear Clinical Meeting 1999;34: Database: IPA.
Exclude - Not a Primary Study

Davis AS, Wingfield CL, Forrester CW, et al. Streamlining outpatient drug therapy at a Veterans Affairs Medical Center. Am J Health Syst Pharm 1997;54(23):2719-20. Database: IPA.
Exclude - Not MMIT

Davis E. Electronic transfer of prescriptions in primary care. BJHC & IM 2003;17(8):29-31. Database: CINAHL.
Exclude - Not a Primary Study

Davis KL, Berensen NM, MacFall HM, et al. Retrospective evaluation of propofol usage and selected adverse events in adult patients. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.
Exclude - Not a Primary Study

Davis L, Lemanowicz GS, Young JA, et al. Third generation decentralization. Ashp Midyear Clinical Meeting 1990;25: Database: IPA.
Exclude - Not a Primary Study

Davis NM. Concept of physician direct computer order entry needs pharmacists' support. Hosp Pharm 1996;31(7):774-7. Database: IPA.
Exclude - No Outcomes of Interest

Davis NM. Lack or failure of the safety net as a cause of medication errors. Hosp Pharm 1997;32(2):143-4. Database: IPA.
Exclude - No Outcomes of Interest

Davis RB. Enhancing productivity in an integrated environment. Comput Healthc 1988;9(5):45-8. 3160944
Database: Inspec.
Exclude - Not MMIT

Davis RG, Hepfinger CA, Sauer KA, et al. Retrospective evaluation of medication appropriateness and clinical pharmacist drug therapy recommendations for home-based primary care veterans. *American Journal Geriatric Pharmacotherapy* 2007;5(1):40-7. Database: Embase Sept 22-09.
Exclude - Not MMIT

Davis RL, Black S, Vadheim C, et al. Immunization tracking systems: experience of the CDC Vaccine Safety Datalink sites. *HMO Pract* 1997;11(1):13-7. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Davis RPM. Northern Ireland's electronic prescribing and eligibility system. *BJHC & IM* 2007;24(1):12-4. Database: CINAHL.
Exclude - Not a Primary Study

Davison J, Holden J. Clinical pharmacist teaching rounds: The first step to caseload management. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Davydov L, Smith LG, Caliendo GC, et al. Analysis of clinical intervention documentation by dispensing pharmacists. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Davydov L, Caliendo GC, Smith LG, et al. Analysis of clinical intervention documentation by dispensing pharmacists in a teaching hospital. *Hosp Pharm* 2003;38(4):346-50. Database: Embase Sept 22-09.
Exclude - Not MMIT

Dawe U, Warnock-Matheron A, Ross S. Mapping the future of hospital information systems: priorities for nursing applications. *Comput Nurs* 1993;11(2):61-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Dawodu T, Churchill W, Cooley T, et al. Development of a NICU web-based physician order-entry system in a major tertiary care teaching hospital. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Dawson J, Kushniruk A. Improving the efficiency and accuracy of a tablet PC interface for computerized provider order entry through usability evaluation and provision of data entry strategies. *Studies in Health Technology & Informatics* 2009;143:447-52. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Daye K, Lashman D, Pandiani J. Medication history display and evaluation. In Washington; DC): American Medical Informatics Association; 1994. p. 993. Grey Lit.
Exclude - Not a Primary Study

Dayton CS, Ferguson J, Hornick D, et al. Evaluation of an Internet-based decision-support system for applying the ATS/CDC guidelines for tuberculosis preventive therapy. *Med Decis Making* 2000;20(1):1-6. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

de Bittner MR, Michocki R. Establishing a pharmaceutical care database. *J Am Pharm Assoc (Wash)* 1996;NS36(1):60-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

de Bruijn N, Lucas P, Schurink K, et al. A probabilistic approach to improved antibiotic therapy. *Studies in Health Technology & Informatics* 1999;68:690-5. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

De Carolis B, de Rosis F, Grasso F, et al. Generating recipient-centered explanations about drug prescription. *Artif Intell Med* 1996;8(2):123-45. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

de Clippele F. The law on e-Health: draft proposal for an electronic medical prescription. *Acta Chir Belg* 2005;105(5):450-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

de Gier H. The impact of automation - Decision support systems (BPP-4-2). *International Pharmaceutical Federation World Congress* 2004;62:11 Database: IPA.

Exclude - No Outcomes of Interest

de Gier JJ, Lastdrager CJ, Rutten GWH, et al. Decision support for quality assurance in medication surveillance. *Pharm Weekbl* 1996;131(1):14-21. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

de Gier JJ. The Electronic Pharmaceutical Dossier: an effective aid to documenting pharmaceutical care data. *Pharmacy World & Science* 1996;18(6):241-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

De Konin GHP, Egberts ACG, Lenderink AW. How frequently and in what amount: Medication breaks by transition between hospital and community care. *Pharm Weekbl* 2000;135(29):1071-5. Database: IPA.

Exclude - Unable to Retrieve Foreign

de Lemos ML, O'Brien RK. Development of a pharmacist-coordinated system of chemotherapy protocols in an integrated healthcare delivery organization. *Journal of Oncology Pharmacy Practice* 2002;8(2-3):39-48. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

de Lusignan S, Robinson J. Information systems in primary care. *Practice Nurse* 2006;31(3):16-20. Database: BSC.

Exclude - Not MMIT

de Lusignan S, van Vlymen J, Hague N, et al. Using computers to identify non-compliant people at increased risk of osteoporotic fractures in general practice: a cross-sectional study. *Osteoporos Int* 2006;17(12):1808-14. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

De Rijdt T. Robotisation and automatisatie, a new way of dispensing medicines. *Ashp Midyear Clinical Meeting* 2007;42. Database: IPA.

Exclude - Not a Primary Study

De Roos AM. National commitment: Up-to-date medicines list to save lives. *EJHP Practice* 2009;15(1):63 OVID EMBASE.
Exclude - Not a Primary Study

de Wet C, Bowie P. The preliminary development and testing of a global trigger tool to detect error and patient harm in primary-care records. *Postgrad Med J* 2009;85(1002):176-80. PMID:19417164 OVID MEDLINE.
Exclude - Not MMIT

de Zegher I, Milstein C, Sene B, et al. OPADE: development of an European computerized drug prescription system. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1993;144-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

de Zegher I, Venot A, Milstein C, et al. OPADE: optimization of drug prescription using advanced informatics. *Computer Methods & Programs in Biomedicine* 1994;45(1-2):131-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

de Zegher I, Milstein C, Sene B, et al. Prescription guidelines in OPADE: what are they, how are they used? *Studies in Health Technology & Informatics* 1995;16:199-205. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Dean, B.S. Creating a healthcare culture of patient safety: A retrospective analysis of change readiness associated with the implementation of computerized provider order entry 2006. Database: CINAHL.
Exclude - Theses

Dean NL. An extra dose of safety: Installation of a bar-coding system drives an entire workflow redesign at a non-profit hospital and healthcare network. *Health Manag Technol* 2007;28(4):30-4. 9422074
Database: Inspec.
Exclude - Not a Primary Study

DeBenedette V. Planning and education: Keys to drug bar-coding. *Drug Topics* 2008;148(22):HSE43 American Medical Informatics Association Database: IPA.
Exclude - Not a Primary Study

Debold P. Electronic prescription and medication documentation. In Amsterdam, Netherlands: IOS Press; 1997. p.248-54.6522644
Database: Inspec.
Exclude - No Outcomes of Interest

Decker E. Database programs: Powerful pharmacy tools. *Drug Topics* 1996;136:28
Database: IPA.
Exclude - Not a Primary Study

DeClariss J W. D-ATM, a working example of health care interoperability: from dirt path to gravel road. In Piscataway, NJ, USA: IEEE; 2009. p.4643-5. Engineering Village Compendex and Inspec.
Exclude - No Outcomes of Interest

deClifford JM, Lam SS, Leung BK. Evaluation of a pharmacist-initiated E-script transcription service for discharged patients. *Journal of Pharmacy Practice and Research* 2009;39(1):39-42. Database: Embase Sept 22-09.

Exclude - Not MMIT

DeFrancesco MS. Adapting to the changing environment: Redefining the annual visit. *Obstet Gynecol Surv* 2007;62(8):491-3. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Defreitas AJ, Waters JP, Ferraro M, et al. Pen to keyboard: A multidisciplinary approach to transitioning from a paper to an electronic medication administration record. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Degli Esposti E, Berto P, Buda S, et al. The Pandora Project: results of the pilot study. *Am J Hypertens* 1999;12(8 Pt 1):790-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Degnan D, Merryfield D, Hultgren S. Reaching out to clinicians: implementation of a computerized alert system. *J Healthc Qual* 2004;26(6):26-30. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Del Fiol G, Rocha BH, Kuperman GJ, et al. Comparison of two knowledge bases on the detection of drug-drug interactions. *Proceedings of the AMIA Symposium* 2000;171-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Del Fiol G, Rocha BH, Nohama P. Design, implementation and evaluation of a clinical decision support system to prevent adverse drug events. *Studies in Health Technology & Informatics* 2000;77:740-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Del Fiol G, Rocha RA, Bradshaw RL, et al. An XML model that enables the development of complex order sets by clinical experts. *IEEE Transactions on Information Technology in Biomedicine* 2005;9(2):216-28. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Del Fiol,G. Context-aware prediction of clinician information needs using “infobuttons.” *The University of Utah Editor*. 2008. Grey Lit.

Exclude - Not MMIT

Del Fiol G. Effectiveness of Topic-specific Infobuttons: A Randomized Controlled Trial. *J Am Med Inform Assoc* 2008;15(6):752-9. Database: Embase Sept 22-09.

Exclude - Not MMIT

Del Fiol G, Haug PJ. Classification models for the prediction of clinicians' information needs. *Journal of Biomedical Informatics* 2009;42(1):82-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Delaney BC. Potential for improving patient safety by computerized decision support systems. *Fam Pract* 2008;25(3):137-8. Database: PsycINFO.

Exclude - Not a Primary Study

Delgado Sanchez O, Escriva Torralva A, Vilanova Bolto M, et al. Comparative study of errors in electronic versus manual prescription [Spanish]. *Farmacia Hospitalaria* 2005;29(4):228-35. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Demiris G, Rantz MJ, Aud MA, et al. Older adults' attitudes towards and perceptions of 'smart home' technologies: A pilot study. *Med Inform Internet Med* 2004;29(2):87-94. 8216504

Database: Inspec.

Exclude - Not MMIT

Demkjaer K, Johansen I, Bernstein K. Third generation electronic pharmacy communications. Recommendations based on ten years' experience. *Studies in Health Technology & Informatics* 1999;68:278-82. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Dempsey C, Budzyna T, Madden S. Improve processes with perioperative EHR. *Health Manag Technol* 2009;30(10):32-3.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010443447&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010443447 EBSCO CINAHL.

Exclude - Not a Primary Study

denheim K. Health HIT and medicaid: Opportunities for states. Part I of a three part series on the state alliance for e-health. *National Conference of State Legislatures*; 2007.

http://www.ncsl.org/print/health/forum/HIT_and_Medicaid.pdf Grey Lit.

Exclude - Not a Primary Study

Dennis L. Stand-alone e-prescribing: Ready or not? *Healthcare Information and Management Systems Society*; 2007. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=67209> Grey Lit.

Exclude - Not a Primary Study

Deshimaru M, Tsuruta S, Togashi H, et al. Drug information system using distributed data processing system based on UNIX. *Japanese Journal of Hospital Pharmacy* 1995;21(6):511-8. Database: IPA.

Exclude - Not MMIT

Desouza KC, Thomas D, Zhang Y. Information integrity in healthcare enterprises: Strategies for mitigation of medical errors. *International Journal of Healthcare Technology and Management* 2004;6(2):241-55. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

DesRoches CM, Campbell EG, Vogeli C, et al. Electronic health records' limited successes suggest more targeted uses. *Health Aff (Millwood)* 2010;29(4):639-46.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010629264&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1759&accno=2010629264 EBSCO CINAHL.

Exclude - Not MMIT

Dete H, May T, Legras B, et al. A computer based medical consultation for antibiotherapy useful for medical practitioner. *Medecine et Maladies Infectieuses* 1988;18(10):420-6. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Deutsch T, Roudsari AV, Leicester HJ, et al. UTOPIA: a consultation system for visit-by-visit diabetes management. *Med Inform (Lond)* 1996;21(4):345-58. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Devamalar PMB, Bai V, Srivatsa S. The new telemedicine paradigm: Fully automated real time web-centric expert system to support diabetes diagnosis. *International Journal of Healthcare Technology and Management* 2008;9(5-6):526-39. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Devers KJ, Liu G. Leapfrog patient-safety standards are a stretch for most hospitals. *Issue Brief/Center for Studying Health System Change* 2004;(77):1-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Devine,E. Evaluating the impact of an ambulatory computerized provider order entry system on outcomes in a community-based multispecialty health system. University of Washington Editor. 2008. Grey Lit.
Exclude - Theses

Devine EB, Wilson-Norton JL, Lawless NM, et al. Characterization of prescribing errors in an internal medicine clinic. *Am J Health Syst Pharm* 2007;64(10):1062-70. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Dexheimer JW, Talbot TR, Sanders DL, et al. Prompting clinicians about preventive care measures: A systematic review of randomized controlled trials. *J Am Med Inform Assoc* 2008;15(3):311-20. Database: IPA.
Exclude - Not a Primary Study

DeYoung JL, Vanderkooi ME, Barletta JF. Effect of barcode medication administration (BCMA) on medication administration error rates in an adult medical intensive care unit (MICU). *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

DiCenzo R, Ferren DM, Resch SJ. Use of data driven decisions to allocate pharmacy resources. *Ashp Midyear Clinical Meeting* 1995;30: Database: IPA.
Exclude - Not a Primary Study

Dickinson JC, Warshaw G, Gehlbach S, et al. Improving hypertension control: Impact of computer feedback and physician education. *Med Care* 1981;19(8):843-54. Database: Embase Sept 22-09.
Exclude - Not MMIT

Diefes RS. Evaluating commercial PCOE systems for safe medication practice. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Dietzel GTW. What do we expect from ehealth and telemedicine? Telematics for integrated/seamless care - Developments in Germany and Europe. *Journal fur Kardiologie* 2003;10(7-8):314-7. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

DiFrancesco M, Andrews T. Alamance Regional Medical Center improves patient safety with CPOE. *J Healthc Inf Manag* 2004;18(1):18-23. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

DiFrancesco MC, DiFrancesco MC. CPOE in a community hospital setting: a nurse's perspective. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Dimick C. HIM breakup: Changing times pull HIM and coding apart. *Journal of the American Health Information Management Association* 2009;80(3):32-6. OVID EMBASE.

Exclude - Not a Primary Study

Dinh AK. Securing portable devices. *Journal of the American Health Information Management Association* 2009;80(1):56-7. OVID EMBASE.

Exclude - Not a Primary Study

Dinklage K, Davidow S, White SJ, et al. Adaption of bar code technology to an existing controlled drug record-keeping system. *Ashp Midyear Clinical Meeting* 1988;23: Database: IPA.

Exclude - Not a Primary Study

Dinning C, Branowicki P, O'Neill JB, et al. Chemotherapy error reduction: a multidisciplinary approach to create templated order sets. *J Pediatr Oncol Nurs* 2005;22(1):20-30. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

DiTusa L, Luzier AB, Jarosz DE, et al. Treatment of hypertension in a managed care setting. *Am J Manag Care* 2001;7(5):520-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Divine HS, Millheim ET, Adams JS. Clinic pharmacist's use of prescription assistance programs to impact patient compliance. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Dixon B and Samarth A. Innovations in using health IT for chronic disease management: Findings from the AHRQ health IT portfolio. AHRQ; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_840586_0_0_18/09-0029-EF_cdm.pdf Grey Lit.

Exclude - Not a Primary Study

Dixon B and Zafar A. Inpatient computerized provider order entry (CPOE): Findings from the AHRQ health IT portfolio. AHRQ; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_846328_0_0_18/09-0031-EF_cpoe.pdf Grey Lit.

Exclude - Not a Primary Study

Dixon JF, Wielgosz C, Pires ML. Health care research and improvement. Description and outcomes of a custom Web-based patient occurrence reporting system developed for Baylor University Medical Center and other system entities. *Baylor University Medical Center Proceedings* 2002;15(2):199-202. Database: CINAHL.
Exclude - Not MMIT

Dixon NM, Shofer M. Struggling to invent high-reliability organizations in health care settings: Insights from the field. *Health Serv Res* 2006;41(4p2):1618-32. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Dixon P. Medical identity theft: The information crime that can kill you. *World Privacy Forum*; 2006. http://www.worldprivacyforum.org/pdf/wpf_medicalidtheft2006.pdf Grey Lit.
Exclude - Not a Primary Study

Do N, Peterson C, Marcum D and others. Reduction of clinic telephone consultation workload through a novel process using physician extenders and computer-based medication refill algorithms. In 2000. Los Angeles, CA): American Medical Informatics Association; 2000. p. 995. Grey Lit.
Exclude - Not MMIT

Dobscha SK, Winterbottom L, Snodgrass L. Reducing drug costs at a Veterans Affairs hospital by increasing market-share of generic fluoxetine. *Community Ment Health J* 2007;43(1):75-84. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Dodge J. Single focus. *Health Data Manag* 2010;18(3):82-5. OVID EMBASE.
Exclude - Not a Primary Study

Dodson G. Computers and controlled drugs. *American Druggist* 1985;192(Aug):122-3. Database: IPA.
Exclude - Not a Primary Study

Doebbeling B, Chou A, Tierney M. Priorities and strategies for the implementation of integrated informatics and communications technology to improve evidence-based practice. *J Gen Intern Med* 2006;21(Suppl 2):S50-S57 Database: PsycINFO.
Exclude - Not a Primary Study

Doherty JA, Reichley RM, Noirot LA, et al. Monitoring pharmacy expert system performance using statistical process control methodology. *AMIA* 2003;205-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Doherty M. How point of sale works in pharmacy. *Irish Pharmaceutical Union Review* 2004;15(Apr):130 Database: IPA.
Exclude - Not MMIT

Dohnalek LJ, Cusaac L, Westcott J, et al. The code to safer transfusions. *Nurs Manag (Harrow)* 2004;35(6):33-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Dolan JG. Medical decision making using the analytic hierarchy process: choice of initial antimicrobial therapy for acute pyelonephritis. *Med Decis Making* 1989;9(1):51-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Donald JB. On line prescribing by computer. *British Medical Journal Clinical Research Ed* 1986;292(6525):937-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Donald JB. Prescribing costs when computers are used to issue all prescriptions. *Br Med J* 1989;298(6690):28-30. Database: Embase Sept 22-09.

Exclude - Not MMIT

Donnelly K. SNOMED-CT: The advanced terminology and coding system for eHealth. *Studies in Health Technology & Informatics* 2006;121:279-90. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Donohoe T. England's Electronic Prescription Service: Taking off now on the implementation runway. *BJHC & IM* 2006;23(2):19-21. Database: CINAHL.

Exclude - Not a Primary Study

Donohoe T. Delivering England's electronic prescriptions service. *BJHC & IM* 2009;22(6):15-6. Database: CINAHL.

Exclude - Not MMIT

Donohue P. Computer-generated prescriptions: A path for the future or a necessity for the present? *All Ireland Journal of Nursing & Midwifery* 2003;2(6):10-1. Database: CINAHL.

Exclude - No Outcomes of Interest

Donovan M J, Zielstorff R D, Mauldin T and others. Using COSTAR to assist nurses in hypertension screening and education. In Silver Spring, MD, USA: IEEE Comput. Soc. Press; 1984. p.487-90.2288283

Database: Inspec.

Exclude - Not a Primary Study

Donze A, Wolf M. Safety in the NICU: preventing medication errors with computerized provider order entry. *Nursing for Women's Health* 2007;11(6):612-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Exclude - Not a Primary Study

Doolan DF, Bates DW. Computerized physician order entry systems in hospitals: mandates and incentives. *Health Aff (Millwood)* 2002;21(4):180-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Dorenfest S. New approaches for the new millenium: IT use in ambulatory clinics. *Healthcare Information and Management Systems Society*; 2000.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=62792> Grey Lit.

Exclude - Not a Primary Study

Dorfschmid., Cornelia M. Why RAT-STATS and Sampling Are Hot. *Journal of Health Care Compliance* 2010;12(3):41-6.
<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=50320781&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.
Exclude - Not a Primary Study

Dorr D, Bonner LM, Cohen AN, et al. Informatics systems to promote improved care for chronic illness: A literature review. *J Am Med Inform Assoc* 2007;14(2):156-63. Database: IPA.
Exclude - Not a Primary Study

Douglas D W, Martin R H, Sherman B W. Automated control of life-threatening cardiac arrhythmias. In 1. USA: 1973. p. 50.544859
Database: Inspec.
Exclude - No Outcomes of Interest

Douglas GP, Deula RA, Connor SE. The Lilongwe Central Hospital Patient Management Information System: a success in computer-based order entry where one might least expect it. *AMIA 2003*;833 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Douglas J, Larrabee S. Bring barcoding to the bedside. *Nurs Manag (Harrow)* 2003;34(5):36-40. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Douglas W. Wireless-EMR provides end-to-end solution with measurable benefits in ambulatory setting. *Healthcare Information and Management Systems Society*; 2002.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=62802> Grey Lit.
Exclude - Not a Primary Study

Dovey SM, Tilyard MW. The computer research network of the Royal New Zealand College of General Practitioners: an approach to general practice research in New Zealand. *Br J Gen Pract* 1996;46(413):749-52. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Dowding D, Mitchell N, Randell R, et al. Nurses' use of computerised clinical decision support systems: A case site analysis. *J Clin Nurs* 2009;18(8):1159-67. Database: PsycINFO.
Exclude - Not MMIT

Doyle MD. Impact of the bar code medication administration (BCMA) system on medication administration errors. University of Arizona 2005; Ph D 2009; Database: CINAHL.
Exclude - Theses

Dracker P. Use of the Pyxis Medstation in a community hospital. *California Journal of Hospital Pharmacy* 1991;3(Jul):10-1. Database: IPA.
Exclude - Not a Primary Study

Drew RH. Information technology for optimizing the management of infectious diseases. *Am J Health Syst Pharm* 2006;63(10):957-65. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Duckers M, Faber M, Cruijsberg J, et al. Safety and risk management interventions in Hospitals: A systematic review of the literature. *Med Care Res Rev* 2009;66(6 SUPPL.):90S-119S. OVID EMBASE.

Exclude - Not MMIT

Dudde R, Vering T, Piechotta G, et al. Computer-aided continuous drug infusion: setup and test of a mobile closed-loop system for the continuous automated infusion of insulin. *IEEE Transactions on Information Technology in Biomedicine* 2006;10(2):395-402. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Dufour JC, Fieschi D, Fieschi M. Coupling computer-interpretable guidelines with a drug-database through a web-based system--The PRESGUID project. *BMC Med Inform Decis Mak* 2004;4:2 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Dugal B. Automating the delivery of healthcare using RFID tracking systems. *Healthcare Information and Management Systems Society*; 2007.

<http://www.himss.org/asp/ContentRedirector.asp?ContentID=66798> Grey Lit.

Exclude - Not a Primary Study

Duke KP. Implementation and impact of an electronic medication reconciliation program in a community hospital. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Dullabh P and Molfino M. Liability coverage for regional health information organizations: Lessons from the AHRQ-funded state and regional demonstration projects in health information technology and other community efforts. *AHRQ*; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_874953_0_0_18/09-0071-EF.pdf Grey Lit.

Exclude - Not a Primary Study

Duman M B, Jones K I. PharmAssist: A multimedia, multiethnic system for medication counselling. In 1996; Harrogate) SUBJECT(S) Identifier healthcare computing; HC Note(s) Also known as HC96: 1996. p.51-9. Grey Lit.

Exclude - Not a Primary Study

Dumay AC, Freriks G. Quality management issues for medical ICT. *Studies in Health Technology & Informatics* 2004;103:93-100. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Dumitru D, Boecler L. Implementing computerized physician order entry in the perioperative setting. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Dunbar PJ, Madigan D, Grohskopf LA, et al. A two-way messaging system to enhance antiretroviral adherence. *J Am Med Inform Assoc* 2003;10(1):11-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Dunham DP, Makoul G. Improving medication reconciliation in the 21st century. *Current Drug Safety* 2008;3(3):227-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Duran M. Drug-related interventions made through a computerized prescription order entry system in an Internal Medicine Unit. *Rev Clin Esp* 2009;209(6):270-8. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Durieux P, Trinquart L, Colombet I, et al. Computerized advice on drug dosage to improve prescribing practice. *Cochrane Database Syst Rev* 2008;(3):CD002894 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Duszynski KM, Beilby JJ, Marley JE, et al. Privacy considerations in the context of an Australian observational database. *Pharmacoepidemiology & Drug Safety* 2001;10(7):587-94. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Dwyer JL, Vitale K, Billett AL. Development and implementation of a physician order entry system for pediatric chemotherapy orders. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Dydek GJ, Blue PW. Application of a hospital pharmacy computer system to nuclear pharmacy. *Hosp Pharm* 1989;24(5):380-2. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Dydensborg CB, Krabbe T. Compliance of hypertension patients estimated by means of website tool. *Ugeskr Laeger* 2007;169(50):4347-50. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Dykes P. Clinical practice guideline adherence before and after implementation of the HEARTFELT (HEART Failure Effectiveness & Leadership Team) Intervention. *J Cardiovasc Nurs* 2005;20(5): Database: PsycINFO.

Exclude - Not MMIT

Dykstra R. Computerized physician order entry and communication: Reciprocal impacts. *ElectronicHealthcare* 2003;2(3):36-42. Database: CINAHL.

Exclude - Not MMIT

East T, Heermann L, Bradshaw R, et al. Efficacy of computerized decision support for mechanical ventilation: Results of a prospective multi-center randomized trial. *Proceedings of the AMIA Symposium* 1999;251-5. Exclude - Not MMIT

Eastham JH. A screening tool to identify look-alike/sound-alike medication storage risks in an automated dispensing cabinet. *California Journal of Health-System Pharmacy* 2006;21(2):6 Database: IPA.

Exclude - Not a Primary Study

Ebrahimi V, Riou C, Seroussi B, et al. Design of a decision support system for chronic diseases coupling generic therapeutic algorithms with guideline-based specific rules. *Studies in Health Technology & Informatics* 2006;124:483-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Eccles M, McColl E, Steen N, et al. Effect of computerised evidence based guidelines on management of asthma and angina in adults in primary care: cluster randomised controlled trial. *BMJ* 2002;325(7370):941 Parent Article 35569. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Eckel SF. User satisfaction with an intravenous medication safety system. *Am J Health Syst Pharm* 2006;63(15):1419-23. Database: Embase Sept 22-09.

Exclude - Not MMIT

ECRI. Patient-controlled analgesic infusion pumps. *Health Devices* 2006;35(1):5-35.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Edelstein SA. Proposed rules for electronic medical records and e-prescribing: do they go far enough. *Manag Care Interface* 2006;19(3):47-51. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Eden A, Grach M, Goldik Z, et al. The implementation of an anesthesia information management system. *Eur J Anaesthesiol* 2006;23(10):882-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Edlin M. Standardized models needed for e-prescribing adoption - Early collaborative launches show promising results, but creating the infrastructure is easier said than done. *Managed Healthcare Executive* 2003;13(12):33-4. Database: IPA.

Exclude - Not a Primary Study

Edwards DR. Computerized prescriber order entry - Tips and strategies. *Journal Michigan Pharmacist* 2004;42(3):24-7. Database: IPA.

Exclude - Not MMIT

Edwards E, Gosney M, Vellodi C, et al. GPs' responses to computer-generated prescription warnings. *British Journal of Pharmaceutical Practice* 1989;11(Jul):232-4. Database: IPA.

Exclude - Unable to Retrieve

Edwards KA, Ehrenzweig BA. Application of bar coding to a hospital top-up service. *Pharmaceutical Journal* 1987;238:234-5. Database: IPA.

Exclude - Unable to Retrieve

Edwards M, Moczygemba J. Reducing medical errors through better documentation. *Health Care Manager* 2004;23(4):329-33. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Effken JA, Brewer B, Patil A, et al. Using OrgAhead, a computational modeling program, to improve patient care unit safety and quality outcomes. *Int J Med Inf* 2005;74(7-8):605-13.

Database: Embase Sept 22-09.

Exclude - Not MMIT

Ehlert DA. Improving the safety of medication administration using low-tech and high-tech approaches. ASHP Annual Meeting 2001;58: Database: IPA.

Exclude - Not a Primary Study

Eichelberger B. Implementing a clinical information system in a managed care setting: Building clinical decision support systems to profile physician practice and develop guidelines in a staff-model HMO. Am J Health Syst Pharm 1997;54:1510-5. Database: IPA.

Exclude - Not a Primary Study

Eidem L, Bond J, Rosendale J, et al. Evaluation of a point-of-care medication bedside bar code scanning system in a tertiary care teaching hospital. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Einbinder JS, Rury C, Safran C. Outcomes research using the electronic patient record: Beth Israel Hospital's experience with anticoagulation. Proceedings - the Annual Symposium on Computer Applications in Medical Care 1995;819-23. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Eisenwine JE, Tourville JF, Hetey SK, et al. Information technology overview in a pediatric teaching health-system: A pharmacy systems perspective. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Ekedahl A, Oskarsson V, Sundberg B, et al. Impact of postal and telephone reminders on pick-up rates of unclaimed e-prescriptions. Pharmacy World & Science 2008;30(5):503-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Eklund B, Joustra-Enquist I. Sustains--direct access for the patient to the medical record over the Internet. Studies in Health Technology & Informatics 2004;100:182-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

El Aneed A, Alaghebandan R, Gladney N, et al. Prescription drug abuse and methods of diversion: The potential role of a pharmacy network. J SUBST USE 2009;14(2):75-83. Database: CINAHL.

Exclude - Not a Primary Study

El Kareh R, Gandhi TK, Poon EG, et al. Trends in primary care clinician perceptions of a new electronic health record. J Gen Intern Med 2009;24(4):464-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

El Rouby S, Rinehart K, Zucker ML, et al. Hand-held personal digital assistant program for the HEMOCHRON RxDx heparin and protamine dosing system. J Extra Corpor Technol 2003;35(3):212-7. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Elganzouri ES, Standish CA, Androwich I. Medication Administration Time Study (MATS): nursing staff performance of medication administration. *J Nurs Adm* 2009;39(5):204-10. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Elliott WJ. Compliance strategies. [Review] [70 refs]. *Current Opinion in Nephrology & Hypertension* 1994;3(3):271-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ellrodt G, Glasener R, Cadorette B, et al. Multidisciplinary rounds (MDR): an implementation system for sustained improvement in the American Heart Association's Get With The Guidelines program. *Critical Pathways in Cardiology: A Journal of Evidence-Based Medicine* 2007;pathw.(3):106-16. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Elson R. Best practices in clinical decision support and CPOE. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Elson R, Neuenschwander M. Debating key issues in technology and informatics - Computerized provider order entry versus bar code medication administration. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Elson RB, Connelly DP. Computerized decision support systems in primary care. *Primary Care* 2009;22(2):365-84. Database: CINAHL.

Exclude - No Outcomes of Interest

Emanuel JE. Implementing decentralized pharmacy order entry in critical care units following implementation of computerized medication administration with bar coding technology. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Emmerechts A, Derde MP, Massart DL, et al. Database for computer-assisted processing of intravenous medication prescriptions. *Journal de Pharmacie Clinique* 1988;7(Suppl 2):261-5. Database: IPA.

Exclude - Unable to Retrieve Foreign

Enderlin GM, Summerfield MR, Burney MT. Use of a microcomputer in a pediatric hospital. *Am J Hosp Pharm* 1987;44(3):565-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Endoh A, Minato K, Komori M, et al. Quantitative comparison of human computer interaction for direct prescription entry systems. *Medinfo* 1995;8 Pt 2:1101-5. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Engelbrecht R, Hildebrand C. Telemedicine and diabetes. *Studies in Health Technology & Informatics* 1999;64:142-54. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Englebright JD, Franklin M. Managing a new medication administration process. *J Nurs Adm* 2005;35(9):410-3. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Enguidanos SM, Brumley RD. Risk of medication errors at hospital discharge and barriers to problem resolution. *Home Health Care Serv Q* 2005;24(1-2):123-35. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Ennis K, Maus R. Kokomo family care: automating the clinical practice. *Med Group Manage J* 2001;48(4):8-12. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Epinette WW. RXWriter and medication manager. *M D Computing* 1987;4(2):29-32. 2914124
Database: Inspec.
Exclude - No Outcomes of Interest

Epstein RH, Gratch DM, Grunwald Z. Development of a scheduled drug diversion surveillance system based on an analysis of atypical drug transactions. *Anesthesia & Analgesia* 2007;105(4):1053-60. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Erbele SM, Heck AM, Blankenship CS. Survey of computerized documentation system use in drug information centers. *Am J Health Syst Pharm* 2001;58(8):695-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Ertel GJ, Glenn DJ, Brown MD, et al. Utilization of a personal digital assistant by pharmacists at the point of care. *ASHP Annual Meeting* 2001;58. Database: IPA.
Exclude - Not a Primary Study

Ertle AR, Campbell EM, Hersh WR. Automated application of clinical practice guidelines for asthma management. *Proceedings/AMIA Annual Fall Symposium* 1996;552-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Ervin K, Culley C, Gross P, et al. Using a database management system to organize formulary decisions. *Ashp Midyear Clinical Meeting* 2004;39. Database: IPA.
Exclude - Not a Primary Study

Ervin KC, Skledar SJ, Hess MM. Designing and implementing a physician compliance profiling system for a drug use and disease state management program. *Ashp Midyear Clinical Meeting* 2001;36. Database: IPA.
Exclude - Not a Primary Study

Escobar A, Benitez A, Rivera B. Improving the quality of medication error reporting through failure mode and effects analysis (FMEA). *Ashp Midyear Clinical Meeting* 2005;40. Database: IPA.
Exclude - Not a Primary Study

Escoto KH, Hallock M, Wagner J, et al. Using variance analysis to detect hazards in a bar-code-assisted medication preparation process. *Joint Commission Journal on Quality & Safety* 2004;30(11):622-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Eskew A, Geisler M, O'Connor L, et al. Enhancing patient safety: clinician order entry with a pharmacy interface. *J Healthc Inf Manag* 2002;16(1):52-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Eslami S, Abu-Hanna A, de Keizer NF. Evaluation of outpatient computerized physician medication order entry systems: a systematic review. *J Am Med Inform Assoc* 2007;14(4):400-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Eslami S, de Keizer NF, Abu-Hanna A. The impact of computerized physician medication order entry in hospitalized patients--a systematic review. *Int J Med Inf* 2008;77(6):365-76. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Espinoza AD, Garcia-Vazquez J P, Rodriguez M D and others. Enhancing a Wearable help button to support the Medication Adherence of older adults. In Merida, Yucatan, Mexico: IEEE Computer Society; 2009. p.3-7. *Engineering Village Compendex and Inspec*.

Exclude - No Outcomes of Interest

Evans AR, Fitter M, Absolon PJ et al. Computerised system for the control, issue and audit of repeat medication. UKSC 120. UK: IBM United Kingdom Ltd., Winchester, UK; 1983 Aug. 2271180

Database: Inspec.

Exclude - No Outcomes of Interest

Evans RF, O'Reilly M. Medication safety in the operating room. *Seminars in Anesthesia, Perioperative Medicine and Pain* 2004;23(2):115-20. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Evans RS. The HELP system: a review of clinical applications in infectious diseases and antibiotic use. *MD Comput* 1991;8(5):282-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Evans S, Stemple C. Electronic health records and the value of health IT. *J Manag Care Pharm* 2008;14(6):S16-S18 Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Evans S, Woo P, Marconi K. Implementation of a remote pharmacist medication order entry service (RPMOES) for a small community hospital. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Evenson RC. Clinical judgment vs. multivariate formulas in assignment of psychotropic drugs. *J Clin Psychol* 1976;29(3): Database: PsycINFO.

Exclude - Not MMIT

Eye K, Hyland-Marciniak BJ, With C, et al. Portable physician profiling system: Application designed to measure changes in prescribing. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Falas T, Papadopoulos G, Stafylopatis A. A review of decision support systems in telecare. *J Med Syst* 2003;27(4):347-56. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fan S, Wen M, Hsu C and others. Health pal: A PDA phone that will take care of your health. In Montreal, QC, Canada: Institute of Electrical and Electronics Engineers Inc.; 2007. p.3703-8.20081311169831

Database: Compendex.

Exclude - Not MMIT

Fan W, Pei-Ran S, Bo-Han W. Using RFID to guard inpatient medication safety. In Worthington, OH, USA: ISRST; 2008. p.111-7. *Engineering Village Compendex and Inspec*.

Exclude - Unable to Retrieve

Fanberg H. The RFID revolution. *Mark Health Serv* 2004;24(3):43-4. Database: BSC.

Exclude - Not MMIT

Fanikos J, Curtis KA, Valez R, et al. Computerized restricted antibiotic program at a large teaching institution. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Fanikos J, Fiumara K, Baroletti S, et al. Impact of smart infusion technology on administration of anticoagulants (unfractionated Heparin, Argatroban, Lepirudin, and Bivalirudin). *Am J Cardiol* 2007;99(7):1002-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Farbstein K, Clough J. Improving medication safety across a multihospital system. *Jt Comm J Qual Improv* 2001;27(3):123-37. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Farrar K, Slee A, Hughes D. Use of structured electronic prescribing pathways to reduce medication errors. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Farrar K, Slee A, Yeats M. Hospital dispensing. On auto pilot. *Health Serv J* 2002;112(5826):26-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Farrar K, Slee A. Computerised prescribing and its impact on medication-error rates. *BJHC & IM* 2004;20(4):28-30. Database: CINAHL.

Exclude - No Outcomes of Interest

Farrar K, Caldwell NA, Robertson J, et al. Use of structured paediatric-prescribing screens to reduce the risk of medication errors in the care of children. *BJHC & IM* 2004;20(4):25-7.

Database: CINAHL.

Exclude - No Outcomes of Interest

Farrar KT, Slee AL, Hughes DK. Formulary management by physician order entry systems, fact or fiction. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Farrell AK, Caliendo G, Friedman T, et al. Impact of a computerized physician order entry driven algorithm on the prescribing and utilization of psychiatric drugs. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Farrell C. Why a VAT Tax Is Where It's At. *BusinessWeek Online* 2010;4

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=48547257&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Farrington JM, Alfano SL, Fitzsimmons RA. Guidelines for the use of ondansetron in the prevention and treatment of post-operative nausea and vomiting. *ASHP Annual Meeting* 1995;52: Database: IPA.

Exclude - Not a Primary Study

Farrington JM, Dembry LM. Development and implementation of guidelines for the monitoring of serum vancomycin concentrations. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Farrington P, Pugh S, Colville A, et al. A new method for active surveillance of adverse events from diphtheria/tetanus/pertussis and measles/mumps/rubella vaccines.[see comment]. *Lancet* 1995;345(8949):567-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Feifer RA, Nevins LM, McGuigan KA, et al. Mail-order prescriptions requiring clarification contact with the prescriber: prevalence, reasons, and implications. *J Manag Care Pharm* 2003;9(4):346-52. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Feinberg JL, Tobias DE, Cameron KA. MDS-Med Guide: Assessing medication effects using patient assessment data. *American Association of Colleges of Pharmacy Annual Meeting* 2004;101(Jul):49 Database: IPA.

Exclude - Not MMIT

Feldbaum J. Establishing order: for CIOs, creation of order sets is one of the first battles to be fought in the march toward CPOE. *Healthc Inform* 2009;26(8):51-2.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010392922&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1239&accno=2010392922
EBSCO CINAHL.

Exclude - Not a Primary Study

Feldman CM. Physician order entry: It really does make a difference. *Issues in Interdisciplinary Care* 2001;3(2):151-4. Database: CINAHL.

Exclude - Unable to Retrieve

Feldman JM. Do anesthesia information systems increase malpractice exposure? Results of a survey. *Anesth Analg* 2004;99(3):840-3. Database: Embase Sept 22-09.

Exclude - Not MMIT

Feldman PH, Murtaugh CM, Pezzin LE, et al. Just-in-time evidence-based e-mail “reminders” in home health care: impact on patient outcomes. *Health Serv Res* 2005;40(3):865-85. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Feldman PH, McDonald M, Rosati RJ, et al. Exploring the utility of automated drug alerts in home healthcare. *J Healthc Qual* 2006;28(1):29-40. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Felkey B. Electric connections. *American Druggist* 1996;213(Aug):43-4. Database: IPA.

Exclude - Not a Primary Study

Felkey BG. Introduction: Clinical workstation or how to reengineer pharmacy. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Felkey BG, Barker KN. Technology and automation in pharmaceutical care.[see comment]. *J Am Pharm Assoc (Wash)* 1996;NS36(5):309-14. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve

Felkey BG. Health system informatics.[see comment]. *Am J Health Syst Pharm* 1997;54(3):274-80. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Felkey BG, McAllister JC. Many faces of pharmacy informatics. *Computertalk for the Pharmacist* 1998;18(May-Jun):6-7. Database: IPA.

Exclude - Not a Primary Study

Felkey BG, Poikonen J. Technology and automation update. *J Manag Care Pharm* 1998;4(May-Jun):345-52. Database: IPA.

Exclude - Not MMIT

Felkey BG. Putting technology to work for you. *ASHP Annual Meeting* 2000;57: Database: IPA.

Exclude - Not a Primary Study

Felkey BG, Fox BI. Crossroads between pharmacy and technology. *California Pharmacist* 2004;48(4):16-9. Database: IPA.

Exclude - Not a Primary Study

Felkey BG, Fox BI. Pursuing patient portals. *Hosp Pharm* 2009;44(4):346-7. Database: IPA.

Exclude - Not MMIT

Felt-Lisk S, Johnson L, Fleming C, et al. Toward understanding EHR use in small physician practices. *Health Care Financ Rev* 2009;31(1):11-22. OVID EMBASE.

Exclude - Not a Primary Study

Fennie KP, Bova CA, Williams AB. Adjusting and censoring electronic monitoring device data. Implications for study outcomes. *Journal of Acquired Immune Deficiency Syndromes: JAIDS* 2006;43:Suppl-95 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fernando B, Savelyich BSP, Avery AJ, et al. Prescribing safety features of general practice computer systems: Evaluation using simulated test cases. *BMJ: British Medical Journal* 2004;328(7449):1171-2. Database: CINAHL.

Exclude - Not MMIT

Ferren AL. Gaining MD buy-in: physician order entry. *J Healthc Inf Manag* 2002;16(2):66-70. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ferris D, Menyah DK. Reducing the use of inappropriate medications in the hospitalized elderly. *Am J Health Syst Pharm* 2001;58(Sep 1):1588 Database: IPA.

Exclude - Not a Primary Study

Fetzer S. Research recap and reflection. *Nurs News (Meriden)* 2009;33(2):16
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010419757&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=673&accno=2010419757> EBSCO CINAHL.

Exclude - Not a Primary Study

Fidei GE. Evaluation of a pharmacy bar-code controlled drug management system in a community hospital. *Ashp Midyear Clinical Meeting* 1992;27. Database: IPA.

Exclude - Not a Primary Study

Field TS, Gurwitz J, Harrold L, et al. Strategies for detecting adverse drug events among older persons in the ambulatory setting. *J Am Med Inform Assoc* 2004;11(6):492-8.

Database: Embase Sept 22-09.

Exclude - Not MMIT

Fields M, Peterman J. Intravenous medication safety system averts high-risk medication errors and provides actionable data. *Nurs Adm Q* 2005;29(1):78-87. Database: CINAHL.

Exclude - Not MMIT

Fiery EM. One-voice wholesalers expand service to pharmacy. *American Druggist* 2004;190(Nov):52 Database: IPA.

Exclude - Not a Primary Study

Figar S, Waisman G, de Quiros FG, et al. Narrowing the gap in hypertension: effectiveness of a complex antihypertensive program in the elderly. *Disease Management* 2004;7(3):235-43. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Figge HL, Fox B, Tribble D. Electronic prescribing of controlled substances. *Am J Health Syst Pharm* 2009;66(14):1311-6. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Figge HL. Electronic prescribing in the ambulatory care setting. *Am J Health Syst Pharm* 2009;66(1):16-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Figge HL. Reducing medication errors using technological innovations. U S Pharmacist 2009;34(3):HS15-HS16 Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Fihn SD, McDonell MB, Vermes D, et al. A computerized intervention to improve timing of outpatient follow-up: a multicenter randomized trial in patients treated with warfarin.

National Consortium of Anticoagulation Clinics. J Gen Intern Med 1994;9(3):131-9.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Finch E, Mayne C. Thinking beyond CPOE to integrated IT strategy and management. J

Healthc Inf Manag 2004;18(1):24-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fine AM, Kalish LA, Forbes P, et al. Parent-driven technology for decision support in pediatric emergency care. Joint Commission Journal on Quality and Patient Safety

2009;35(6):307-15. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Finkelstein J, O'Connor G, Friedmann RH. Development and implementation of the home asthma telemonitoring (HAT) system to facilitate asthma self-care. Studies in Health

Technology & Informatics 2001;84(Pt:1):1-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Finley JM, Lewis RK, Schumock GT. Medication safety technology in community hospitals.

Ashp Summer Meeting 2004;60: Database: IPA.

Exclude - Not a Primary Study

Fireworker R B, Esposito V. Computerization of a hospital pharmacy. In Silver Spring, MD, USA: IEEE Comput. Soc. Press; 1984. p.246-50.2547060

Database: Inspec.

Exclude - Not a Primary Study

Firth G. Re-negotiating reproductive technologies: the 'public foetus' revisited. Feminist

Review 2009;(92):54-71. Scholar's Portal Sociological Abstracts.

Exclude - Not a Primary Study

Fischell R E. Microprocessor application to an artificial pancreas. In New York, NY, USA:

IEEE; 1980. p.15-21.1604619

Database: Inspec.

Exclude - Not MMIT

Fischell R E, Webster W G. The programmable implantable medication system (PIMS):

Design and applications. In New York, NY, USA: IEEE; 1983. p. 691.2320907

Database: Inspec.

Exclude - No Outcomes of Interest

fischer S, Stewart T, Mehta S, et al. Handheld computing in medicine. J Am Med Inform

Assoc 2003;10(2):139-49. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Fischer S, Stewart TE, Mehta S, et al. Handheld computing in medicine. *J Am Med Inform Assoc* 2003;10(2):139-49. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fiset P, Mathers L, Engstrom R, et al. Pharmacokinetics of computer-controlled alfentanil administration in children undergoing cardiac surgery. *Anesthesiology* 1995;83(5):944-55. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Fish CA, Kirking DM, Martin JB. Information systems for evaluating the quality of prescribing. *Ann Pharmacother* 1992;26(3):392-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fisher CM, Corrigan O, Henman M. A study of community pharmacy practice. 2. Prescription dispensing. *Journal of Social and Administrative Pharmacy* 1991;8(2):65-8. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

FitzHenry F. Measuring the quality of medication administration. In Washington, DC): 2005. p. 955. Grey Lit.

Exclude - Not a Primary Study

FitzHenry F, Doran JB, Matheny ME, et al. BCMA evaluation: finding significance in near misses. *AMIA 2008;Annual:Symposium* PMID:18999154 OVID MEDLINE.

Exclude - No Outcomes of Interest

Fitzmaurice DA, Hobbs FD, Murray ET, et al. Evaluation of computerized decision support for oral anticoagulation management based in primary care. *Br J Gen Pract* 1996;46(410):533-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Fitzmaurice DA, Hobbs FD, Delaney BC, et al. Review of computerized decision support systems for oral anticoagulation management. [Review] [15 refs]. *Br J Haematol* 1998;102(4):907-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fitzmaurice DA, Hobbs FD, Murray ET, et al. Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Arch Intern Med* 2000;160(15):2343-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Fitzpatrick R. Automated dispensing - Developing a business case to support investment. *Hospital Pharmacist* 2004;11(3):109-11. Database: IPA.

Exclude - Not a Primary Study

Fitzsimmons WE, Kaplan S, Carasiti ME, et al. Computer-assisted patient targeting and surveillance. *ASHP Annual Meeting* 1990;47: Database: IPA.

Exclude - Not a Primary Study

Flanagan ME, Patterson ES, Frankel RM, et al. Evaluation of a physician informatics tool to improve patient handoffs. *J Am Med Inform Assoc* 2009;16(4):509-15. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Flanders SJ, Juneja R, Roudebush CP, et al. Glycemic control and insulin safety: the impact of computerized intravenous insulin dosing. *Am J Med Qual* 2009;24(6):489-97. PMID:19666741 OVID MEDLINE.

Exclude - Not MMIT

Flebbe E, Jensen TB, Andersen PE. [Does electronic medicine prescription cause new types of errors?]. [Danish]. *Ugeskr Laeger* 2009;171(33):2260-4. PMID:19732502 OVID MEDLINE.

Exclude - Unable to Retrieve

Fleming DM. Practice based information systems for monitoring medicines. *Pharmaceutical Medicine* 1994;8(3-4):161-76. Database: IPA.

Exclude - No Outcomes of Interest

Flottorp S, Havelsrud K, Oxman A. Process evaluation of a cluster randomized trial of tailored interventions to implement guidelines in primary care--why is it so hard to change practice? *Fam Pract* 2003;20(3):333-9. Database: PsycINFO.

Exclude - Not a Primary Study

Flynn A, McGregory M. Pharmacy informatics task force update: Advancing medication-related clinical decision support. In Chicago, IL) SUBJECT(S) Identifier HIMSS; Healthcare information; Management systems Note(s) Held in e-store (RMS); Held on CD-ROM: 2009. p. 108. Grey Lit.

Exclude - No Outcomes of Interest

Flynn AJ. Smart infusion systems - How the addition of wireless capability improves smart pump management and utilization. *Ashp Midyear Clinical Meeting* 2007;42. Database: IPA.

Exclude - Not a Primary Study

Flynn AJ. Opportunity cost of pharmacists' nearly universal prospective order review. *Am J Health Syst Pharm* 2009;66(7):668-70. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Foisy MM, Tseng A. Development of an interactive computer-assisted program to manage medication therapy in HIV infected patients. *Drug Inf J* 1998;32(3):649-56. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Fonarow GC, Yancy CW, Albert NM, et al. Improving the use of evidence-based heart failure therapies in the outpatient setting: The IMPROVE HF performance improvement registry. *Am Heart J* 2007;154(1):12-38. Database: IPA.

Exclude - Not a Primary Study

Fong CJ, Elliott CH, Ulrick-Richmond BJ. Bar coding to the bedside: What it really means. *Ashp Midyear Clinical Meeting* 2003;38. Database: IPA.

Exclude - Not a Primary Study

Fong CJ, Elliot C, Engelman L, et al. Bar coding to the bedside: One year later-insights & unexpected benefits. *Ashp Midyear Clinical Meeting* 2004;39. Database: IPA.

Exclude - Not a Primary Study

Fonseca JA, Costa-Pereira A, Delgado L, et al. Asthma patients are willing to use mobile and web technologies to support self-management. *Allergy* 2006;61(3):389-90. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Font I, Poveda JL, Tordera M, et al. Pharmacy management based on standardization of processes and their decentralized implementation by clinical area pharmacists. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Font N, I, Climent C, Poveda Andres JL. Quality of drug treatment process through medication errors in a tertiary hospital. *Farmacia Hospitalaria* 2008;HOSP..(5):274-9. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Fontaine AL. Current requirements and emerging trends for labelling as a tool for communicating pharmacovigilance findings. *Drug Saf* 2004;27(8):579-89. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fontan J-E. Quality insurance of prescribing, dispensing and managing drugs in hospital pharmacy. *Journal de Pharmacie Clinique* 1995;14(4):280-8. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Food and Drug Administration H. Bar code label requirement for human drug products and biological products. Final rule. *Fed Regist* 2004;69(38):9119-71. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Food and Drug Administration H. Medical devices; general hospital and personal use devices; classification of remote medication management system. Final rule. *Fed Regist* 2007;72(202):59175-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fook V, Tee J, Yap K and others. Smart mote-based medical system for monitoring and handling medication among persons with dementia. In 2007; (Nara, Japan) SUBJECT(S) Identifier Ubiquitous computing; Home automation; Pervasive computing; ICOST Note(s) Includes bibliographical references and index.: 2007. p.54-62.Grey Lit.

Exclude - Not a Primary Study

Footo SO, Coleman JR. Medication administration: the implementation process of bar-coding for medication administration to enhance medication safety. *Nurs Econ* 2008;26(3):207-10. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Footo SO, Coleman JR. Success story. Medication administration: The implementation process of bar-coding for medication administration to enhance medication safety. *Nursing Economic\$* 2009;26(3):207-10. Database: CINAHL.

Exclude - Not MMIT

Foppe Van Mil JW, Tromp TF, McElnay J. Community pharmacy around the world. *International Pharmacy Journal* 2001;15(Jul):7-10. Database: IPA.
Exclude - Not MMIT

Forcinio H. Unit-of-use bar coding: Balancing the challenge of technological change with improved patient safety. *Pharmaceutical Technology North America* 2002;26(10):28-35. 2002447164132
Database: Compendex.
Exclude - Not a Primary Study

Forcinio H. Reaping the benefits of bar coding: Quick response comes to healthcare. *Pharmaceutical Technology* 1995;19(Jan):34 Database: IPA.
Exclude - Not a Primary Study

Forcinio H. Taking your medicine. *Managing Automation* 2005;20(6):46-8. 8721037
Database: Inspec.
Exclude - Not MMIT

Ford EW, McAlearney AS, Phillips MT, et al. Predicting computerized physician order entry system adoption in US hospitals: can the federal mandate be met? *Int J Med Inf* 2008;77(8):539-45. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Ford EW, Short JC. The impact of health system membership on patient safety initiatives. *Health Care Manage Rev* 2008;33(1):13-20. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Ford N. Why electronic prescribing? Is the technology ready yet? *Pharmacy Management* 1995;10(3):28-31. Database: IPA.
Exclude - Not MMIT

Ford N, Curtis C, Paul P. The use of electronic prescribing as part of a system to provide medicines management in secondary care. *BJHC & IM* 2001;17(8):26-9. Database: CINAHL.
Exclude - No Outcomes of Interest

Forni A, Skehan N, Hartman CA, et al. Evaluation of the impact of a tele-ICU pharmacist on the management of sedation in critically ill mechanically ventilated patients. *Ann Pharmacother* 2010;44(3):432-8. PMID:20164471 OVID MEDLINE.
Exclude - Not MMIT

Forni A, ChU.H.T., Fanikos J. Technology Utilization to Prevent Medication Errors. *Current Drug Safety* 2010;5(1):13-8. OVID EMBASE.
Exclude - No Outcomes of Interest

Forsstrom JJ, Gronroos P, Irjala K, et al. Linking patient medication data with laboratory information system. *Int J Biomed Comput* 1996;42(1-2):111-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Fort A, Narsinghani U, Bowyer F. Evaluating the safety and efficacy of Glucommander, a computer-based insulin infusion method, in management of diabetic ketoacidosis in children, and comparing its clinical performance with manually titrated insulin infusion. *J Pediatr Endocrinol* 2009;22(2):119-25. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Fortescue EB, Kaushal R, Landrigan CP, et al. Prioritizing strategies for preventing medication errors and adverse drug events in pediatric inpatients. *Pediatrics* 2003;111(4p1):722-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Fortier CR, Sura ME. Implementing BCMA: Assessing readiness, designing project plans, and managing change. *Ashp Summer Meeting 2009*;65: Database: IPA.

Exclude - Not a Primary Study

Fortuna RJ, Ross-Degnan D, Finkelstein J, et al. Clinician attitudes towards prescribing and implications for interventions in a multi-specialty group practice. *J Eval Clin Pract* 2008;14(6):969-73. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Foster J. Economic effectiveness of two different automated anesthesia medication dispensing devices at two different facilities. *Ashp Midyear Clinical Meeting 2006*;41: Database: IPA.

Exclude - Not a Primary Study

Foster RA, Antonelli PJ. Computerized physician-order entry: are we there yet? *Otolaryngol Clin North Am* 2002;35(6):1237-43. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fowles J, Weiner J, Chan K. Performance measures using electronic health records: Five case studies. *The Commonwealth Fund*; 2008.

http://www.commonwealthfund.org/publications/publications_show.htm?doc_id=685103
Grey Lit.

Exclude - Not a Primary Study

Fox BI. eMARs, smart pumps, & bar coding at the point of care: Is there evidence? *Ashp Midyear Clinical Meeting 2006*;41: Database: IPA.

Exclude - Not a Primary Study

Fox CH, Swanson A, Kahn LS, et al. Improving chronic kidney disease care in primary care practices: an upstate New York practice-based research network (UNYNET) study. *Journal of the American Board of Family Medicine: JABFM* 2008;21(6):522-30. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fox GN, Weidmann E, Diamond DE, et al. Hand-held electronic prescribing. *J Fam Pract* 2001;50(5):449-54. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fox J, Thomson R. Decision support and disease management: a logic engineering approach. *IEEE Transactions on Information Technology in Biomedicine* 1998;2(4):217-28. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fox N, Ward K, O'Rourke A. The birth of the e-clinic. Continuity or transformation in the UK governance of pharmaceutical consumption? *Soc Sci Med* 2005;61(7):1474-84. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Foy JL, Eastman RC, Nealon RC, et al. Automated therapeutic drug monitoring in an ambulatory care endocrine clinic. *Ann Pharmacother* 1992;26(5):675-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Foy JS, Eastman RC, Nealon RC, et al. Automated therapeutic drug monitoring in an ambulatory care endocrine clinic. *ASHP Annual Meeting* 1991;26(5):675-8. Database: IPA.

Exclude - Not a Primary Study

Fraizer C. E-prescribing in clinical practice. *J Med Pract Manage* 2004;20(3):148-51. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Frakes MA, High K, Stocking J. Transport nurse safety practices, perceptions, and experiences: the Air and Surface Transport Nurses Association survey. *Air Med J* 2009;28(5):250-5.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010420563&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=588&accno=2010420563 EBSCO CINAHL.

Exclude - Not a Primary Study

Frangou S, Sachpazidis I, Stassinakis A, et al. Telemonitoring of medication adherence in patients with schizophrenia. *Telemedicine and e-Health* 2005;11(6):675-83. Database: PsycINFO.

Exclude - Not MMIT

Frank G, Lawless ST, Steinberg TH. Improving physician communication through an automated, integrated sign-out system. *J Healthc Inf Manag* 2005;19(4):68-74. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Frank L, Galanos H, Penn S, et al. Using BPI and emerging technology to improve patient safety. *J Healthc Inf Manag* 2004;18(1):65-71. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Frank O, Kidd M. A guide to computer-generated prescriptions. *Aust Fam Physician* 1996;25(7):1162-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Franklin BD, O'Grady K. Dispensing errors in community pharmacy: Frequency, clinical significance and potential impact of authentication at the point of dispensing. *Int J Pharm Pract* 2007;15(4):273-81. Database: Embase Sept 22-09.

Exclude - Not MMIT

Franklin BD, O'Grady K, Voncina L, et al. An evaluation of two automated dispensing machines in UK hospital pharmacy. *Int J Pharm Pract* 2008;16(1):47-53. Database: IPA.

Exclude - Not MMIT

Franklin BD, Birch S, Savage I, et al. Methodological variability in detecting prescribing errors and consequences for the evaluation of interventions. *Pharmacoepidemiology & Drug Safety* 2009;18(11):992-9. PMID:19634116 OVID MEDLINE.

Exclude - Not MMIT

Franklin MJ, Sittig DF, Schmitz JL, et al. Modifiable templates facilitate customization of physician order entry. *Proceedings / AMIA* 1998;1998(1998):315-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fred T, Caplan C, Levy JM, et al. Clinician Feedback on Using Episode Groupers with Medicare Claims Data. *Health Care Financ Rev* 2009;31(1):51-61.
<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=48050717&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Fredriks DA, Coleman RW. Nomogram for dosing warfarin at steady state. *Clin Pharm* 1991;10(12):923-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Freire CC, Escobar Gimenes F.R., De Bortoli Cassiani S.H. Analysis of the computerized prescription in two clinics of a teaching hospital. *Medicina (Mex)* 2004;37(1-2):91-6. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Fretheim A, Oxman AD, Treweek S, et al. Rational Prescribing in Primary Care (RaPP-trial). A randomised trial of a tailored intervention to improve prescribing of antihypertensive and cholesterol-lowering drugs in general practice [ISRCTN48751230]. *BMC Health Serv Res* 2003;3(1):5 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Fretschner R, Bleicher W, Heininger A, et al. Patient data management systems in critical care. *J Am Soc Nephrol* 2001;12(Suppl 17):s83-s86 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Friedlin J, Dexter PR, Overhage JM. Details of a successful clinical decision support system. *AMIA* 2007;2007(2007):254-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Friedman C, Ash J, Tarczy-Hornoch P. Concordance between medication histories and outpatient electronic prescription claims in patients hospitalized with heart failure. In Washington, DC): 2005. p. 1109. Grey Lit.

Exclude - Not MMIT

Friedman LN, Halpern NA, Fackler JC. Implementing an electronic medical record. *Crit Care Clin* 2007;23(3):347-81. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Friedman MA, McCluskey CF. Observational assessment of clinical outcomes associated with the use of chemistry laboratory values in the Theratrac 2 system. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Friedman MA, Schueth A, Bell D. Interoperable electronic prescribing in the United States: A progress report. *Health Aff (Millwood)* 2009;28(2):393-403. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Froese P, Muller-Bohn T. Schleswig-Holstein health card. *Dtsch Apoth Ztg* 2004;145: Database: IPA.
Exclude - Not MMIT

Fu Q, Xue Z, Zhu J, et al. Anaesthesia record system on handheld computers--pilot experience and uses for quality control and clinical guidelines. *Computer Methods & Programs in Biomedicine* 2005;77(2):155-63. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Fuji KT, Galt K. Pharmacists and health information technology: Emerging issues in patient safety. *HEC Forum* 2008;20(3):259-75. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Fulda TR, Lyles A, Pugh MC, et al. Current status of prospective drug utilization review. *J Manag Care Pharm* 2004;10(5):433-41. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Fung EY, Fedorko L, Fisher J. Point of care medication bar-coding pilot in operating room. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Fung V, Ortiz E, Huang J, et al. Early experiences with e-Health services (1999-2002): Promise, reality, and implications. *Med Care* 2006;44(5):491-6. Database: PsycINFO.
Exclude - Not MMIT

Furukawa H, Maida C, Omori M, et al. Supply of injectable drugs for individual patients using the prescription entry system. Part 1. Evaluation of the prescription entry system of injectable drugs. *Japanese Journal of Hospital Pharmacy* 1996;22(6):615-27. Database: IPA.
Exclude - Unable to Retrieve Foreign

Furukawa H. Challenge for preventing medication errors-learn from errors: What is the most effective label display to prevent medication error for injectable drug? In Beijing, China: 2007. p.437-42. Grey Lit.
Exclude - No Outcomes of Interest

Gabrielline J, Lovly R. Computer assisted satellite pharmacy consultative service in a primary care clinic. *Pharm Times* 1985;55(Mar):146 Database: IPA.
Exclude - Not a Primary Study

Gabutti L, Lotscher N, Bianda J, et al. Would artificial neural networks implemented in clinical wards help nephrologists in predicting epoetin responsiveness? *BMC Nephrology* 2006;7(13):1-11. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gadzhanova S, Iankov II, Warren JR, et al. Developing high-specificity anti-hypertensive alerts by therapeutic state analysis of electronic prescribing records. *J Am Med Inform Assoc* 2007;14(1):100-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gaeta TJ, Fiorini M, Ender K, et al. Potential drug-drug interactions in elderly patients presenting with syncope. *J Emerg Med* 2002;22(2):159-62. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gaglio S, Ruggiero C, Spinelli G, et al. BREASTCAN: an expert system for postoperative breast cancer therapy. *Computers & Biomedical Research* 1986;19(5):445-61. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gaikwad R, Sketris I, Shepherd M, et al. Evaluation of accuracy of drug interaction alerts triggered by two electronic medical record systems in primary healthcare. *Health Informatics Journal* 2007;13(3):163-77. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gaillour F. Why do health systems flop with CPOE? *Physician Exec* 2004;30(2):28-9. Database: BSC.

Exclude - Not a Primary Study

Gainer A, Pancheri K, Zhang J. Improving the human computer interface design for a physician order entry system. *AMIA* 2003;847 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Galanter WL, Didomenico RJ, Polikaitis A. Preventing exacerbation of an ADE with automated decision support. *J Healthc Inf Manag* 2002;16(4):44-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Galanter WL, Hier DB, Jao C, et al. Computerized physician order entry of medications and clinical decision support can improve problem list documentation compliance. *Int J Med Inf* 2010;79(5):332-8. OVID EMBASE.

Exclude - Not MMIT

Galloway MJ, Foggin JJ, Dixon S. Introduction of computer assisted control of oral anticoagulation in general practice. *J Clin Pathol* 1995;48(12):1144-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Galusha C, Brown M, Kelly J. Bedside bar codes: Protecting patients and nurses. *Online Journal of Nursing Informatics* 2003;7(3):10-20. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Gamble KH. Cutting the cord. CIOs are leveraging wireless technologies to help nurses deliver patient care safely and efficiently. *Healthcare informatics : the business magazine for information and communication systems* 2009;26(9):30-3. OVID EMBASE.

Exclude - Not a Primary Study

Gamble KH. Is it registering? Patient portals, part II (see Financial Department for part I). *Healthc Inform* 2009;26(5):24-8. PMID:19514236 OVID MEDLINE.

Exclude - Not a Primary Study

Gamble KH. Identity crisis. The push to share data electronically--both inside and outside of the hospital walls--is forcing patient identification to the forefront. *Healthc Inform* 2010;27(1):22-6. PMID:20120889 OVID MEDLINE.

Exclude - Not a Primary Study

Ganapathy K and Ravindra A. mHealth: A potential tool for healthcare delivery in India. 2008. http://www.ehealth-connection.org/files/conf-materials/mHealth_A%20potential%20tool%20in%20India_0.pdf Grey Lit.

Exclude - Not a Primary Study

Gandhi TK, Weingart S.N., Seger A., et al. Does computerized prescribing help reduce medication errors and ADRs in the outpatient setting? *Formulary* 2001;36(8):613-4. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Gandhi TK, Weingart SN, Borus J, et al. Adverse drug events in ambulatory care. *N Engl J Med* 2003;348(16):1556-64. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gandhi TK, Weingart S, Seger A, et al. Outpatient prescribing errors and the impact of computerized prescribing. *J Gen Intern Med* 2005;20(9):837-41. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Gandhi TK, Poon EG, Sequist TD, et al. Primary care clinician attitudes towards ambulatory computerized physician order entry. *AMIA* 2005;961 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gannon K. How to sell physicians on order entry: Tips from M.D.s. *Hospital Pharmacist Report* 1996;10(Apr):14-5. Database: IPA.

Exclude - Not a Primary Study

Garber MC, Nau DP, Erickson SR, et al. The concordance of self-report with other measures of medication adherence: a summary of the literature. *Med Care* 2004;42(7):649-52.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Garcia-Alamino JM, Ward AM, Alonso-Coello P, et al. Self-monitoring and self-management of oral anticoagulation. *Cochrane Database Syst Rev* 2010;4: PMID:20393937 OVID MEDLINE.

Exclude - Not MMIT

Garde S, Baumgarten B, Basu O, et al. A meta-model of chemotherapy planning in the multi-hospital/multi-trial-center-environment of pediatric oncology. *Methods Inf Med* 2004;43(2):171-83. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gardetto NJ, Greaney K, Arai L, et al. Critical pathway for the management of acute heart failure at the veterans affairs san diego healthcare system: Transforming performance measures into cardiac care. *Critical Pathways in Cardiology* 2008;7(3):153-72. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Gardiner C, Williams K, Mackie IJ, et al. Can oral anticoagulation be managed using telemedicine and patient self-testing? A pilot study. *Clinical & Laboratory Haematology* 2006;28(2):122-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gardner RM, Ostler DV, Nelson BD, et al. The role of smart medical systems in the Space Station. *International Journal of Clinical Monitoring & Computing* 1989;6(2):91-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gardner RM. The HELP clinical decision-support system. *J Med Pract Manage* 1994;9(4):177-81. Database: Embase Sept 22-09.

Exclude - Not MMIT

Gardner RM, Pryor T, Warner H. The HELP hospital information system: Update 1998. *Int J Med Inf* 1999;54(3):169-82. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Gardner R, Clemmer T, Larsen K, et al. Computerized alert system use in clinical medicine. *Journal of Sound and Vibration* 1979;136-40. 1980050001471 Database: Compendex.

Exclude - No Outcomes of Interest

Garg AX, Adhikari NK, McDonald H, et al. Effects of computerized clinical decision support systems on practitioner performance and patient outcomes: a systematic review. *JAMA* 2005;293(10):1223-38. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Garich J, Cheung D, Sinclair D, et al. Implementation of a computerized physician order entry system at a 500 bed community hospital: case for pharmacist involvement. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Garjon Parra FJ. Prescribing drugs to patients receiving out-patient care. *Anales del Sistema Sanitario de Navarra* 2009;32(1):11-21. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Garnerin P, Goncerut J., Landau R., et al. Electronic anesthesia patient record: Does-it contribute to patient safety? *Med Hyg (Geneve)* 2004;62(2509):2526-30. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Garreau I, Derharoutunian C., Kiledjian E., et al. Clinical pharmacy in a geriatric unit: Impacts of clinical pharmacy interventions prior to medical order. *Pharmacie Hospitaliere Francaise* 1997;(SPEC. ISS.):46-7. Database: Embase Sept 22-09.

Exclude - Not MMIT

Garrett D. Tapping into the value of health data through secondary use. *Healthc Financ Manage* 2010;64(2):76-83. PMID:20178244 OVID MEDLINE.

Exclude - Not a Primary Study

Garrett LE, Jr., Hammond WE, Stead WW. The effects of computerized medical records on provider efficiency and quality of care. *Methods Inf Med* 1986;25(3):151-7. 2748427

Database: Inspec.

Exclude - Not MMIT

Garrett S K, Craig J B. Medication administration and the complexity of nursing workflow. In Chicago, IL, United states: Institute of Industrial Engineers; 2009. p. Delta.Engineering Village Compendex and Inspec.

Exclude - Not MMIT

Gaspar CM, Romero C, I, Caja M. Pharmaceutical care program in a traumatology hospital. [Spanish]. *Atencion Farmaceutica* 2009;11(5):301-8. OVID EMBASE.

Exclude - Unable to Retrieve

Gastelurrutia MA. Systems of dosing and administration of drugs and pharmaceutical care. *El Farmaceutico* 2001;28-37. Database: IPA.

Exclude - Unable to Retrieve Foreign

Gates DM, Swinkey NJ, Skow NA. Risks and rewards with automated dispensing technology. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Gaunt MJ. Electronic prescribing - Potential areas of weakness. *Pharm Times* 2009;75(10):30 OVID EMBASE.

Exclude - Not a Primary Study

Gawande AA, Bates DW. The use of information technology in improving medical performance. Part II. Physician-support tools. *Medgenmed [Computer File]: Medscape General Medicine* 2000;2(1):E13 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gay T. A case study on CPOE: A blueprint for a beginning. Massachusetts Technology Collaborative and New England Healthcare Institute; 2006.

http://web3.streamhoster.com/mtc/blueprint12_19_06.pdf Grey Lit.

Exclude - Not a Primary Study

Gbadamosi KS, DePew DD, Anderson G, et al. Collaborative improvement in the order and delivery process of intravenous infusion medications in the neonatal intensive care unit to decrease errors and utilize technology. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Gebhart F. Electronic Rx order entry comes with its own problems. Drug Topics 1999;143(22):42 Database: IPA.
Exclude - Not a Primary Study

Gebhart F. VA facility slashes drug errors bar-coding. Drug Topics 1999;143: Database: IPA.
Exclude - Not a Primary Study

Gebhart F. Hospitals - VHA and VA alike - Launch programs to cut errors. Drug Topics 2000;144(22):53 Database: IPA.
Exclude - Not a Primary Study

Gebhart F. California hospitals beat medication error deadline. Drug Topics 2003;147(23):HSE20 Database: IPA.
Exclude - Not a Primary Study

Gebhart F. VA study finds bar-coding brings new errors. Drug Topics 2003;147(4):HSE24 Database: IPA.
Exclude - Not a Primary Study

Gebhart F. New report compares merits of bedside bar-coding. Drug Topics 2004;148(22):HSE40 Database: IPA.
Exclude - Not a Primary Study

Gehlbach SH, Wilkinson WE, Hammond WE, et al. Improving drug prescribing in a primary care practice. Med Care 1984;22(Mar):193-201. Database: IPA.
Exclude - Not MMIT

Geier DA, Geier M. A review of the vaccine Adverse Event Reporting System database. Expert Opinion on Pharmacotherapy 2004;5(3):691-8. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Geiger G, Derman YD. Methodology for evaluating physician order entry (POE) implementations. J Eval Clin Pract 2003;9(4):401-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Geisler BP, Schuur JD, Pallin DJ. Estimates of electronic medical records in U.S. emergency departments. PLoS ONE 2010;5(2):e9274 OVID EMBASE.
Exclude - Not MMIT

Geiss E, Schwartz FW, Bruening R, et al. DOMINIG III. Centralized information system for general practitioners and out-patient medical services: Conception of realization. Electronics and electrical engineering, computer science 1976; Grey Lit.
Exclude - No Outcomes of Interest

Geissbuhler A, Miller RA. Clinical application of the UMLS in a computerized order entry and decision-support system. AMIA Annu Symp Proc 1998;Annual Symposium:320-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Geissbuhler A. Building man-man-machine synergies: experiences from the Vanderbilt and Geneva clinical information systems. *Int J Med Inf* 2003;69(2-3):127-33. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

George D, Austin-Bishop N. Error rates for computerized order entry by physicians versus nonphysicians. *Am J Health Syst Pharm* 2003;60(21):2250-2. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Georgiou A, Williamson M, Westbrook J, et al. The impact of computerised physician order entry systems on pathology services: A systematic review. *Int J Med Inf* 2007;76(7):514-29.
Exclude - Not a Primary Study

Gerard M, Das K, Trick WE, et al. Evolution of clinical decision support to increase influenza vaccination. *AMIA* 2007;962 Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Gestin-Boyer C, Rene C, Cupissol D, et al. [Prevention of cisplatin nephrotoxicity in cancer: adjusting the dosage with a computerized chrono-pump]. [French]. *C R Seances Soc Biol Fil* 1989;183(3):263-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Ghoreishi Nejad S, Martens R, Paranjape R. An agent-based diabetic patient simulation. In Berlin, Germany: Springer-Verlag; 2008. p.832-41.10262370
Database: Inspec.
Exclude - Not MMIT

Ghosh T, Norton M, Skiba D. Communication, coordination and knowledge sharing in the implementation of CPOE: impact on nursing practice. *AMIA* 2006;928 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Giannangelo K and Fenton S. SNOMED CT survey: An assessment of implementation in EMR/EHR applications. *Perspectives in Health Information Management*, 2008.
http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_038479.pdf Grey Lit.
Exclude - Not MMIT

Giannone G. Computer-supported weight-based drug infusion concentrations in the neonatal intensive care unit. *CIN COMPUT INFORM NURS* 2005;23(2):100-5. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Giannopoulos HT, Sterner-Allison J. Collaboration is the key. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Giblin JE, Lakes G. Home infusion therapy trial of a multitherapy remotely programmable ambulatory pump. *Ashp Midyear Clinical Meeting* 1992;27: Database: IPA.
Exclude - Not a Primary Study

Gibson JT. Are you making maximum use of bar codes? *American Druggist* 2004;190(Sep):33-4. Database: IPA.
Exclude - Not a Primary Study

Gibson M, Jenkins KN, Wilson R, et al. Multi-tasking in practice: coordinated activities in the computer supported doctor-patient consultation. *Int J Med Inf* 2005;74(6):425-36.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gibson M, Neil JK, Wilson R, et al. Verbal prescribing in general practice consultations. *Soc Sci Med* 2006;63(6):1684-98. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Gierl L, Steffen D, Ihracky D, et al. Methods, architecture, evaluation and usability of a case-based antibiotics advisor. *Computer Methods & Programs in Biomedicine* 2003;72(2):139-54. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gijssels-Wiersma DG, van den Bemt PM, Walenbergh-van Veen MC. Influence of computerised medication charts on medication errors in a hospital. *Drug Saf* 2005;28(12):1119-29. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gilabert-Perramon A, Lopez-Calahorra P, Escoda-Geli N, et al. Electronic prescription in Catalonia, Spain (Rec@t): a health tool. [Spanish]. *Med Clin (Barc)* 2010;134(SUPPL. 1):49-55. OVID EMBASE.

Exclude - No Outcomes of Interest

Gilby J. Computers in hospital pharmacy. *Pharmaceutical Journal* 1990;245(Sep 29):422-3.

Database: IPA.

Exclude - Not MMIT

Gillespie GM. Lost in translation: eliminate medical errors with clear communication. *CMA Today* 2009;42(2):13-7.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010470469&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2589&accno=2010470469

EBSCO CINAHL.

Exclude - Not a Primary Study

Gilmet GP, Mallon RP, Griffin BT, et al. The use of an integrated clinical laboratory and pharmacy diabetes database to provide physician performance feedback in an IPA-model HMO. *J Ambulatory Care Manage* 1998;21(1):12-23. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gilutz H, Novack L, Shvartzman P, et al. Computerized community cholesterol control (4C): meeting the challenge of secondary prevention. *Israel Medical Association Journal: Imaj* 2009;11(1):23-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ginsberg BH, Tan MH, Mazze R, et al. Staged diabetes management: computerizing a disease state management program. *J Med Syst* 1998;22(2):77-87. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Giokas D. Prince Edward Island implements province-wide drug information system: a small step for DIS; a giant leap for the pan-Canadian interoperable electronic health record. HEALTHC Q 2009;12(1):113-6.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010190221&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2877&accno=2010190221
EBSCO CINAHL.

Exclude - Not a Primary Study

Girouard D. Identifying medication-use system variances associated with computerized provider order entry. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Giurdanella P, Di Denia P. Does the electronic prescription reduce drugs errors? Comparison between electronic and manual prescription. Assistenza Infermieristica e Ricerca:Air 2007;26(2):92-8. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Glannelly BF. Using bar code technology with IV medication administration. ASHP Annual Meeting 1991;48: Database: IPA.

Exclude - Not a Primary Study

Glaser J, Kirby J. Evolution of the healthcare CIO. Healthcare financial management : journal of the Healthcare Financial Management Association 2009;63(11):38-41. OVID EMBASE.

Exclude - Not a Primary Study

Glassman PA, Simon B, Belperio P, et al. Improving recognition of drug interactions: benefits and barriers to using automated drug alerts. Med Care 2002;40(12):1161-71. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Glassman PA, Belperio P, Lanto A, et al. The utility of adding retrospective medication profiling to computerized provider order entry in an ambulatory care population. J Am Med Inform Assoc 2007;14(4):424-31. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Glessner MR, Pandak J. Online prospective drug utilization review in community practice: Clinical and economic impact. ASHP Annual Meeting 1992;49: Database: IPA.

Exclude - Not a Primary Study

Glowczewski J, Moskovitz R, Moran MT, et al. Computerized prescriber order entry (CPOE) net benefits realization one year after implementation in an urban teaching hospital. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Gluck ME. Is health information technology associated with patient safety in the United States? Findings Brief Health Care Financing & Organization 2009;12(3):1-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Glynn C, Kleinman RE, Smith S, et al. The evolution and implementation of a pediatric computerized order entry system: a case study. *J Healthc Inf Manag* 2004;18(2):64-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Godolphin W. Shared decision-making. *HEALTHC Q* 2009;12(Sp):e186-e190
PMID:19667767 OVID MEDLINE.
Exclude - Not a Primary Study

Goedert J. Rule oks E-scripts for controlled Rx. *Health Data Manag* 1922;18(5):20
PMID:20464897 OVID MEDLINE.
Exclude - Not a Primary Study

Gohil B, Gholamhosseini H, Harrison MJ, et al. Intelligent monitoring of critical pathological events during anesthesia. *Conference Proceedings: 2007*;4343-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Goldberg DE, Naylor MF, Larson AE, et al. Development of a mini computer program to identify medication orders requiring modification based on patient-specific renal function. *ASHP Annual Meeting 1990*;47(Jun): Database: IPA.
Exclude - Not a Primary Study

Goldberg HI, Ralston JD, Hirsch IB, et al. Using an Internet comanagement module to improve the quality of chronic disease care. *Joint Commission Journal on Quality & Safety* 2003;29(9):443-51. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Goldblum OM. Electronic prescribing: criteria for evaluating handheld prescribing systems and an evaluation of a new, handheld, wireless wide area network (WWAN) prescribing system. *Dermatol Online J* 2001;7(1):1 Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Goldblum OM. Practical applications of hand-held computers in dermatology. *Seminars in Cutaneous Medicine & Surgery* 2002;21(3):190-201. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Golden MS. An incident reporting system: documented at the point of service. *J Healthc Risk Manag* 1998;18(2):18-26. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Goldspiel BR, DeChristoforo R, Daniels CE. A continuous-improvement approach for reducing the number of chemotherapy-related medication errors. *Am J Health Syst Pharm* 2000;57:S4-S9 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Goldstein M. Using health information technology to improve hypertension management. *Current Hypertension Reports* 2008;10(3):201-7. Grey Lit.
Exclude - Not a Primary Study

Goldstein MK, Hoffman BB, Coleman RW, et al. Patient safety in guideline-based decision support for hypertension management: ATHENA DSS. *Proceedings / AMIA 2001*;9(6):214-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Goldstein MK, Coleman RW, Tu SW, et al. Translating research into practice: Organizational issues in implementing automated decision support for hypertension in three medical centers. *J Am Med Inform Assoc* 2004;11(5):368-76. Database: IPA.

Exclude - Not a Primary Study

Gomez EJ, Hernando Perez ME, Vering T, et al. The INCA system: a further step towards a telemedical artificial pancreas. *IEEE Transactions on Information Technology in Biomedicine* 2008;12(4):470-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gomez E, Hernando M, Garcia A, et al. Telemedicine as a tool for intensive management of diabetes: The DIABTel experience. *Comput Methods Programs Biomed* 2002;69(1):163-77.

Exclude - Not MMIT

Gondek K. New technologies. *Maryland Pharmacist* 1987;63(Jul):24-5. Database: IPA.

Exclude - Not MMIT

Gonzalez C, Torres F.P. Electronic medical records. A review and analysis of the current situation. *Diraya: Electronic medical records in Andalusia, Spain. Revista Espanola de Cardiologia Suplementos* 2007;7(C):37C-46C. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Goodell JG, Harry DJ, Vogt CP. Integration of an automated dispensing device into a computerized unit dose hospital pharmacy. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Goto S. Several practical approaches to the hospital information system. *IRYO - Japanese Journal of National Medical Services* 2005;58(11):615-20. Database: Embase Sept 22-09.

Exclude - Not MMIT

Gottlieb J. The long and winding road to computerized physician order entry. *Physician Exec* 2004;30(2):30-5. Database: BSC.

Exclude - Not a Primary Study

Gottlieb LK, Stone EM, Stone D, et al. Regulatory and policy barriers to effect clinical data exchange: Lessons learned from MedsInfo-ED. *Health Aff (Millwood)* 2005;24(5):1197-204. Database: CINAHL.

Exclude - Not a Primary Study

Goud R, Hasman A, Peek N. Development of a guideline-based decision support system with explanation facilities for outpatient therapy. *Comput Methods Programs Biomed* 2008;91(2):145-53. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Goud R, de Keizer NF, ter Riet G, et al. Effect of guideline based computerised decision support on decision making of multidisciplinary teams: cluster randomised trial in cardiac rehabilitation[published erratum appears in BMJ 2009 Jun 6;338:1370]. Br Med J 2009;338:1440
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010362275&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=694&accno=2010362275 EBSCO CINAHL.
Exclude - Not MMIT

Gouin-Thibault I, Levy C, Pautas E, et al. Improving anticoagulation control in hospitalized elderly patients on warfarin. J Am Geriatr Soc 2010;58(2):242-7. PMID:20374400 OVID MEDLINE.
Exclude - Not MMIT

Goundrey-Smith S. Pharmacy robots in UK hospitals: The benefits and implementation issues. Pharmaceutical Journal 2008;280(7502):599-602. Database: Embase Sept 22-09.
Exclude - Not MMIT

Gouveia WA. Computer applications in unit dose drug distribution systems. Am J Hosp Pharm 1984;41(10):2092-3. Database: IPA.
Exclude - Not a Primary Study

Gouveia WA. Managing pharmacy information systems. Am J Hosp Pharm 1993;50(1):113-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Gouveia WA. Transition to managed care. Part 1. Am J Health Syst Pharm 1998;55(3):229 Database: IPA.
Exclude - Not a Primary Study

Gouveia WA, Shane R, Clark T. Computerized prescriber order entry: Power, not panacea. Am J Health Syst Pharm 2003;60(18):1838 Database: IPA.
Exclude - Not a Primary Study

Gozdan MJ. Using technology to reduce medication errors. Nursing (Lond) 2009;39(6):57-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Grabowski BS. Pharmacy-based automated medication records: methods, application, and a survey of use. [Review] [8 refs]. Top Hosp Pharm Manage 1994;14(3):58-72. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Grabowski M, Filipiak KJ, Rudowski R, et al. Project of an expert system supporting risk stratification and therapeutic decision making in acute coronary syndromes. Pol J Pathol 2003;54(3):205-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Grad R, Pluye P, Granikov V, et al. Many family physicians will not manually update PDA software: an observational study... personal digital assistant. *Inform Prim Care* 2009;17(4):225-30.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010613951&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2542&accno=2010613951
 EBSCO CINAHL.
 Exclude - Not MMIT

Graham L, Tharp T. Development and implementation of an automated proactive approach toward improving pneumococcal vaccination rates in an in-patient acute care hospital setting. *Ashp Summer Meeting 2008*;64: Database: IPA.
 Exclude - Not a Primary Study

Graham TA, Bullard MJ, Kushniruk AW, et al. Assessing the sensibility of two clinical decision support systems. *J Med Syst* 2008;32(5):361-8. Database: Ovid MEDLINE(R).
 Exclude - Not MMIT

Grams RR, Zhang D, Yue B. A primary care application of an integrated computer-based pharmacy system. *J Med Syst* 1996;20(6):413-22. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Grandt D, Braun C, Hauser W. Frequency, relevance, causes of and strategies for prevention of medication errors. *Z Gerontol Geriatr* 2005;38(3):196-202. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Grandt D. Prevention of medication errors by electronic decision support. *Z Allgemeinmed* 2005;81(8):341-7. Database: Embase Sept 22-09.
 Exclude - Unable to Retrieve Foreign

Granlien MF, Hertzum M, Gudmundsen J. The gap between actual and mandated use of an electronic medication record three years after deployment. *Studies in Health Technology & Informatics* 2008;136:419-24. Grey Lit.
 Exclude - Not MMIT

Granlien M S, Hertzum M. Implementing new ways of working: Interventions and their effect on the use of an electronic medication record. In *Sanibel Island, FL, United states: Association for Computing Machinery*; 2009. p.321-30. *Engineering Village Compendex and Inspec*.
 Exclude - Not MMIT

Grant P. A new approach to diabetic control: fuzzy logic and insulin pump technology. *Medical Engineering & Physics* 2007;29(7):824-7. Database: Ovid MEDLINE(R).
 Exclude - Not MMIT

Grant RW, Wald JS, Poon EG, et al. Design and implementation of a web-based patient portal linked to an ambulatory care electronic health record: patient gateway for diabetes collaborative care. *Diabetes Technol Ther* 2006;8(5):576-86. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Grant RW, Meigs JB. Overcoming barriers to evidence-based diabetes care. *Current Diabetes Reviews* 2006;2(2):261-9. Database: Ovid MEDLINE(R).
 Exclude - Not MMIT

Gray DL, Ash SR, Jacobi J, et al. The training and use of an artificial neural network to monitor use of medication in treatment of complex patients. *J Clin Eng* 1991;16(4):331-6. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Gray MD, Felkey BG. Computerized prescriber order-entry systems: evaluation, selection, and implementation. *Am J Health Syst Pharm* 2004;61(2):190-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gray S P, Burkett D D. A pharmacy information system. In Arlington, VA, USA: Alliance for Engng. in Medicine & Biology; 1973. p. 321.617877

Database: Inspec.

Exclude - No Outcomes of Interest

Greatbatch D, Luff P, Health C, et al. Special issue on CSCW: III. Interpersonal communication and human-computer interaction: An examination of the use of computers in medical consultations. *Interacting with Computers* 1993;5(2):193-216. Database: PsycINFO.

Exclude - Not MMIT

Green BB, Ralston JD, Fishman PA, et al. Electronic communications and home blood pressure monitoring (e-BP) study: design, delivery, and evaluation framework.

Contemporary Clinical Trials 2008;29(3):376-95. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Green D. E-prescribing incentive: As described in the MIPPA legislation. Centers for Medicare & Medicaid Services; 2008.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=68574> Grey Lit.

Exclude - Not a Primary Study

Greenberg L. CPOE: How can clinical pharmacists help convince physicians that it's worthwhile? *Pharmacy Practice News* 2004;31(6):36-7. Database: IPA.

Exclude - Not a Primary Study

Greenberg MD, Ridgely MS, Bell DS. Electronic prescribing and HIPAA privacy regulation. *Inquiry* 2004;41(4):461-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Greene SK, Shi P, Dutta-Linn MM, et al. Accuracy of data on influenza vaccination status at four Vaccine Safety Datalink sites. *Am J Prev Med* 2009;37(6):552-5. PMID:19944924

OVID MEDLINE.

Exclude - Not MMIT

Greengold N. After the honeymoon: Tools to support evidence based clinical decision making. 2006. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=66886> Grey Lit.

Exclude - Not a Primary Study

Greer ML. RXPART: Prototype expert system for formulary decision making. *Ann Pharmacother* 1992;26(Feb):244-50. Database: IPA.

Exclude - Not a Primary Study

Greim JA, Shek C, Jones L, et al. Enterprise-wide drug-drug interaction alerting system. AMIA 2003;856 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Grimsno A. Electronic prescriptions--without side-effects? Tidsskr Nor Laegeforen 2006;126(13):1740-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Grindrod KA, Patel P, Martin JE. What interventions should pharmacists employ to impact health practitioners' prescribing practices? Ann Pharmacother 2006;40(9):1546-57.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gringras P, Santosh P, Baird G. Development of an Internet-based real-time system for monitoring pharmacological interventions in children with neurodevelopmental and neuropsychiatric disorders. Child Care Health Dev 2006;32(5):591-600. Database: PsycINFO.

Exclude - Not a Primary Study

Grissinger M, Globus NJ. Computerized order-entry systems and other technologies are supposed to make medication administration safer. But if poorly designed or used incorrectly, they can actually introduce errors. Keep your patients safe by avoiding the pitfalls we'll discuss here. Nursing (Lond) 2004;34(1):36-41. Database: IPA.

Exclude - Not a Primary Study

Groenroos P, Irjala K, Heiskanen J and others. Using computerized individual medication data to detect drug effects on clinical laboratory tests. In 222. Turku; Finland: 1995. p.31-6. Grey Lit.

Exclude - No Outcomes of Interest

Gronroos PE, Irjala KM, Selen GP, et al. Computerized monitoring of potentially interfering medication in thyroid function diagnostics. International Journal of Clinical Monitoring & Computing 1997;14(4):255-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Gronroos, P.E. Medication and laboratory: A study on computerized monitoring of drug-test and drug-drug interactions in hospital Turun Yliopisto (Finland) Editor. 1997. Grey Lit.

Exclude - Theses

Gross PA, Pujat D. Implementing practice guidelines for appropriate antimicrobial usage: a systematic review. Med Care 2001;39(8:Suppl:2):55-69. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gross PA, Bates DW. A pragmatic approach to implementing best practices for clinical decision support systems in computerized provider order entry systems. J Am Med Inform Assoc 2007;14(1):25-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gross TM, Kayne D, King A, et al. A bolus calculator is an effective means of controlling postprandial glycemia in patients on insulin pump therapy. Diabetes Technol Ther 2003;5(3):365-9. Database: CINAHL.

Exclude - Not MMIT

Grossman J and Cohen G. Despite regulatory changes, hospitals cautious in helping physicians purchase electronic medical records. 123. Center for Studying Health System Change; 2008. <http://www.hschange.org/CONTENT/1015/> Grey Lit.

Exclude - No Outcomes of Interest

Gruner A, Ljutow A, Schleinzner W, et al. Implementation of an electronic patient record. Experience in an interdisciplinary pain clinic. *Schmerz* 2008;22(1):24-33. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gruson D, Hilbert G., Vargas F., et al. Usual practices of antibiotic cycling. *Reanimation* 2002;11(3):200-8. Database: Embase Sept 22-09.

Exclude - Not MMIT

Gryskovich R. Putting safety first: West Virginia hospital uses point-of-care bar coding for documented improved safety in administering medications. *Health Manag Technol* 2002;23(5):38-9. 7323026

Database: Inspec.

Exclude - Not a Primary Study

Gryskovich RL, Wiegmann DK, Bain LD. Addressing medication errors: Which technological solutions are right for your institution? *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Guchelaar H, Kalmeijer M.D. Improved prescribing: Electronic prescribing in the hospital. *Pharm Weekbl* 2002;137(44):1558-61. Database: Embase Sept 22-09.

Exclude - Not MMIT

Guendelman S, Meade K, Benson M, et al. Improving asthma outcomes and self-management behaviors of inner-city children: A randomized trial of the Health Buddy interactive device and an asthma diary. *Archives of Pediatrics & Adolescent Medicine* 2002;156(2):114-20. Database: CINAHL.

Exclude - Not MMIT

Guerci A, Faccin N, Frenkiel N, et al. Realization of an assisted system for medical decision and follow-up in hematologic diseases treated with sequential chemotherapy. *Rev Med Interne* 1990;11(3):256-60. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Guerrero RM, Nickman NA, Jorgenson JA. Work activities before and after implementation of an automated dispensing system. *Am J Health Syst Pharm* 1996;53(5):548-54. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Guevremont C, Lefebvre P. Pharmaceutical care implementation with computerized interventions in a tertiary care teaching facility: Canadian approach. *ASHP Annual Meeting* 1997;54: Database: IPA.

Exclude - Not a Primary Study

Gugerty B, Wooldridge P, Coleman S and others. Development of the CISQ-MA to assess nursing attitudes towards electronic medication administration modules of healthcare information systems. In Bethesda, MD, USA: American Medical Informatics Assoc; 2002. p. 1035.8023610

Database: Inspec.

Exclude - Not a Primary Study

Guharoy R, Stalder B, Probst L, et al. Optimization of clinical services via effective use of wireless technology. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Guharoy R, Page N, Hirschman K, et al. Computerized physician order entry: Paradigm shift in pharmacy practice in a tertiary care teaching hospital. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.

Exclude - Not a Primary Study

Guigue L, Donadey T. Infections and computer-assisted therapeutic decisions. *Medecine et Maladies Infectieuses* 1999;29(12):758-67. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Gulliford SM, Schneider JK, Jorgenson JA. Using telemedicine technology for pharmaceutical services to ambulatory care patients. *Am J Health Syst Pharm* 1998;55(Jul 15):1512-5. Database: IPA.

Exclude - Not a Primary Study

Gumpper KE, Pedersen CA. Compare yourself with the ASHP 2007 informatics survey. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Gupta A, Crk I, Sarnikar S, et al. Drug effectiveness reporting and monitoring systems: Discussion and prototype development. *International Journal of Technology Management* 2009;47(1-3):174-90. 20091512030534

Database: Compendex.

Exclude - Not MMIT

Guptha SH, Mitra R, Puthrasingam S, et al. Application of evidence-based prescribing indicators in primary care: A pilot. *Qual Prim Care* 2003;11(2):107-16. Database: Embase Sept 22-09.

Exclude - Not MMIT

Gura KM, Chan L, McRae M, et al. Impact of bar code technology on a computerized automated TPN microcompounding system. *ASHP Annual Meeting* 1993;50: Database: IPA.

Exclude - Not a Primary Study

Gurtel A L, Gleser M A, Fallavollita A and others. PHAMIS pharmacy system. In New York, NY, USA: IEEE; 1979. p.885-91.1489925

Database: Inspec.

Exclude - Not MMIT

Gurwitch KD, Summerfield M, Calodney K. Implementation of a dose-checking module. *Ashp Midyear Clinical Meeting* 1989;24: Database: IPA.

Exclude - Not a Primary Study

Gustafsson LL, Widang K, Hoffmann M, et al. Computerized decision support in drug prescribing II: A national database to provide up-to-date and unbiased information. *Lakartidningen* 2003;100(15):1338-40. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Gustafsson LL, Widang K, Hoffmann M, et al. Computerized decision support in drug prescribing I: Better survey of patients' medications yields better quality of care. *Lakartidningen* 2003;100(15):1333-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hagland M. A model of innovation. Designing a medication-administration system for the smallest patients was a huge challenge. *Healthc Inform* 2009;26(4):28-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hagland M. For all the right reasons. Approaching CPOE from a patient safety and care quality perspective is the first critical step toward success. *Healthc Inform* 2009;26(9):40-4. PMID:19813574 OVID MEDLINE.

Exclude - Not a Primary Study

Hagland M. Administration Tech Trends 2010. Trend: data infrastructure. *Healthc Inform* 2010;27(2):12-4. PMID:20218062 OVID MEDLINE.

Exclude - Not a Primary Study

Hagland M. Clinical Tech Trends 2010. Tech Trend: CPOE. *Healthc Inform* 2010;27(2):16-8. PMID:20218063 OVID MEDLINE.

Exclude - Not a Primary Study

Hagland M. One-on-one with CHIME CIO of the Year Bill Spooner of Sharp HealthCare. *Healthc Inform* 2010;27(3):14-6.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010596036&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1239&accno=2010596036

EBSCO CINAHL.

Exclude - Not a Primary Study

Hagland M. Show us the evidence. *Healthc Inform* 2010;27(1):34-8. PMID:20120893 OVID MEDLINE.

Exclude - Not a Primary Study

Hajioff S. Computerized decision support systems: An overview. *Health Informatics Journal* 1998;4(1):23-8. Database: CINAHL.

Exclude - Not a Primary Study

Hakim A. E-prescribing and primary noncompliance: physician and patient experience. *Professional Case Management* 2010;15(2):62-7.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010621802&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=3340&accno=2010621802

EBSCO CINAHL.

Exclude - Not MMIT

Halamka J. Early experiences with E-prescribing. *J Healthc Inf Manag* 2006;20(2):12-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hallas J, Nissen A. Individualized drug utilization statistics. Analysing a population's drug use from the perspective of individual users. *Eur J Clin Pharmacol* 1994;47(4):367-72. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Halloran L. Phrases NPs hate to hear. Strategies for drive-you-crazy questions. *Adv Nurse Pract* 2009;17(3):69 PMID:19999439 OVID MEDLINE.
Exclude - Not a Primary Study

Halpern NA, Thompson RE, Greenstein RJ. A computerized intensive care unit order-writing protocol. *Ann Pharmacother* 1992;26(2):251-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hamann C, Poon E, Smith S, et al. Designing an electronic medication reconciliation system. *AMIA* 2005;976 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hameed S, Shabnam M. Integrated medication and emergency system. In /5; Kuala Lumpur): 2006. p.48-52. Grey Lit.
Exclude - No Outcomes of Interest

Hamilton RM. Valuable clinical decision support integrates with physician's workflow - Clinical decision support within a CPOE system can simplify the treatment-planning process. *Managed Healthcare Executive* 2003;13(12):40-1. Database: IPA.
Exclude - Not a Primary Study

Hamm M and Giang T. Focusing on food and fitness: State strategies for healthy communities and economic development. *National Conference of State Legislatures*; 2008. <http://www.ncsl.org/programs/health/webcastfeb08.htm#I> Grey Lit.
Exclude - Not MMIT

Hammersley VS, Morris CJ, Rodgers S, et al. Applying preventable drug-related morbidity indicators to the electronic patient record in UK primary care: methodological development. *J Clin Pharm Ther* 2006;31(3):223-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hammond KW, Snowden M, Risse SC, et al. An effective computer-based tardive dyskinesia monitoring system. *Am J Med Qual* 1995;10(3):133-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Hammond K, Snowden M, Adkins T, et al. Computerized TD reminders. *Hospital & Community Psychiatry* 1994;45(10):1043-4. Database: PsycINFO.
Exclude - Not a Primary Study

Hammond P, Harris AL, Das SK, et al. Safety and decision support in oncology. *Methods Inf Med* 1994;33(4):371-81. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hammond P, Modgil S, Wyatt JC. Safety and computer-aided design of chemotherapy plans. *Top Health Inf Manage* 2000;20(4):55-66. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hammond WE. The role of standards in electronic prescribing. *Health Aff (Millwood)* 2004;Suppl Web Exclusives(W4):325-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hammond W. A perspective on interoperability. 2008. http://www.ehealth-connection.org/files/conf-materials/Perspective%20on%20Interoperability_0.pdf Grey Lit.

Exclude - Not a Primary Study

Handler JA, Feied CF, Coonan K, et al. Computerized physician order entry and online decision support. *Acad Emerg Med* 2004;11(11):1135-41. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Handler SM, Altman RL, Perera S, et al. A systematic review of the performance characteristics of clinical event monitor signals used to detect adverse drug events in the hospital setting. *J Am Med Inform Assoc* 2007;14(4):451-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hansen RA, Dusetzina SB, Song L, et al. Depression affects adherence measurement but not the effectiveness of an adherence intervention in heart failure patients. *Journal of the American Pharmacists Association: JAPhA* 2009;49(6):760-8. PMID:19926556 OVID MEDLINE.

Exclude - Not a Primary Study

Hanson KK. Impact of barcode medication administration on medication safety: Importance of adherence and ongoing monitoring to sustain improvements. *Ashp Midyear Clinical Meeting* 2007;42. Database: IPA.

Exclude - Not a Primary Study

Hare GT, Reinhard SC, Brick JH, et al. Polypharmacy. Reducing adverse events among the elderly in NJ. Part 1. *N J Med* 1999;96(12):35-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hargrove F, Huffines SK. Results and impact of physician order entry on pharmacy processes. *Ashp Midyear Clinical Meeting* 1997;32. Database: IPA.

Exclude - Not a Primary Study

Hargrove FR, Huffines SK, Knight JR. Experiences and results of developing online order entry. *Ashp Midyear Clinical Meeting* 1998;33. Database: IPA.

Exclude - Not a Primary Study

Hargrove FR. Leveraging computerized provider order entry (CPOE) to meet pharmacy and therapeutic committee directives and patient care initiatives. *Ashp Midyear Clinical Meeting* 2006;41. Database: IPA.

Exclude - Not a Primary Study

Harmsen HH. Using barcodes in pharmacy. *Pharm Weekbl* 1993;128:477-82. Database: IPA.

Exclude - Unable to Retrieve Foreign

Harper P, Harper J. A proposed model for an internet-based computerised anticoagulant monitoring system. *Healthcare Review Online* 2005;9(4):1-9. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Harris CM. Prescribing in general practice: What's going on? *Pharmacy Management* 1991;7(2):15-7. Database: IPA.
Exclude - Not MMIT

Harris CM, Dajda R. The scale of repeat prescribing. *Br J Gen Pract* 1996;46(412):649-53. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Harrison J, Daly M. Leveraging health information technology to improve patient safety. *Public Administration & Management* 2009;14(1):218-37. Database: BSC.
Exclude - Not a Primary Study

Harrison MI, Koppel R, Bar-Lev S. Unintended consequences of information technologies in health care: An interactive sociotechnical analysis. *J Am Med Inform Assoc* 2007;14(5):542-9.
http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_802943_0_0_18/Harrison%20et%20al%20Unintended%20Conseq%20JAMIA.pdf Grey Lit.
Exclude - Not a Primary Study

Harriss C, Pringle M. Do general practice computer systems assist in medical audit? *Fam Pract* 1994;11(1):51-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Harry D, Howe R. Impact of an electronic medication administration record and bar code scanning system on the accuracy of medication administration in hospitalized patients. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
Exclude - Not a Primary Study

Hartley G. Scannable technology - Bar coded pharmaceuticals save lives. *New Zealand Pharmacy* 2006;26(2):17-9. Database: IPA.
Exclude - Not MMIT

Hartmann M, Specht M, Langkafel P. For the patients' good - Medication safety in the hospital. *Klinik Arzt* 2006;35(12):509-13. Database: Embase Sept 22-09.
Exclude - Not MMIT

Hartung DM, McGuinness MR. Evaluating the impact of computerized guidelines for ordering digoxin levels. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Harvey KJ, Vitry AI, Roughead E, et al. Pharmaceutical advertisements in prescribing software: an analysis. *Med J Aust* 2005;183(2):75-9. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Harvey KJ. The Pharmaceutical Benefits Scheme 2003-2004. *Australia and New Zealand Health Policy* 2005;2(1):2 Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Hasman A. Decision support systems and communication in medicine at the University of Limburg. In 11. Seattle, WA, USA: Publ by Alliance for Engineering in Medicine & Biology; 1989. p.1865-6.1990086071098
Database: Compendex.
Exclude - No Outcomes of Interest

Hasman A, Ament A, Arnou PC, et al. Inter-institutional information exchange in healthcare. Int J Biomed Comput 1992;31(1):5-16. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Hasman A, Smeets RP, Dupuits FM. Decision support for health professionals. Studies in Health Technology & Informatics 1998;51:227-30. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hasson M. Portable device appears reliable in evaluating glare of cataract patients. Ocular Surgery News 2009;27(14):9
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010352407&site=ehost-live>; EBSCO CINAHL.
Exclude - Not a Primary Study

Hasson M. Viability of ophthalmic electronic medical records hinges on interoperability. Ocular Surgery News 2009;27(14):1-4.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010352411&site=ehost-live>; EBSCO CINAHL.
Exclude - Not a Primary Study

Hasson M. Handheld corneal hydrometer provides accurate measurement of refractive index. Ocular Surgery News 2010;28(5):8-9.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010582157&site=ehost-live>; EBSCO CINAHL.
Exclude - Not a Primary Study

Hasson NK. Computerized physician order entry: Veterans Affairs Health Care System experience. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.
Exclude - Not a Primary Study

Hasspacher JB. Implementing a computer generated medication administration record. ASHP Annual Meeting 1991;48: Database: IPA.
Exclude - Not a Primary Study

Hatcher I, Sullivan M, Hutchinson J, et al. An intravenous medication safety system: preventing high-risk medication errors at the point of care. J Nurs Adm 2004;34(10):437-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Haugh R, Gearon CJ, Serb C, et al. Point and click. (If only it were that easy.). Hospitals & Health Networks 2002;76(1):36-50. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hausauer JH, Frey C, Barclay B, et al. Technician intervention: Quality assurance report. Ashp Midyear Clinical Meeting 1990;25: Database: IPA.
Exclude - Not a Primary Study

Havlovic DA. Including cost data in antimicrobial agent sensitivity reports. *Am J Hosp Pharm* 2001;46(6):1126 Database: IPA.

Exclude - Not MMIT

Hayashi T, Asakura S, Yamashita K, et al. Development of a patient medication instruction provision system which reflects the doctor's prescribing intentions. *Japanese Journal of Hospital Pharmacy* 1998;24(6):774-81. Database: IPA.

Exclude - No Outcomes of Interest

Hayney R. Electronic prescription for controlled substances - New prescribing option to benefit prescriber, pharmacist and patient. *Journal of the Pharmacy Society of Wisconsin* 2004;56 Database: IPA.

Exclude - No Outcomes of Interest

Hayward GL, Parnes AJ, Simon SR. Using health information technology to improve drug monitoring: a systematic review. *Pharmacoepidemiology & Drug Safety* 2009;18(12):1232-7. PMID:19725020 OVID MEDLINE.

Exclude - Not Primary Study

Hayward MK, Warren RD, Sykes J. Jackson Community Medical Record: a model for provider collaboration in a RHIO. *J Healthc Inf Manag* 2007;21(3):31-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hazebroucq G. Safety and traceability do exist for our customers. *S T P Pharma Pratiques* 2004;14(5):418-23. Database: Embase Sept 22-09.

Exclude - Not MMIT

Hazlehurst B, McMullen C, Gorman P, et al. How the ICU follows orders: care delivery as a complex activity system. *AMIA* 2003;284-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Healthcare Benchmarks and Quality Improvement. Wide-ranging collaborative drives improvement in patient safety: PDAs provided to staff for safer medication administration. *Healthc Benchmarks Qual Improv* 2010;17(1):1-5.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010540299&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2313&accno=2010540299 EBSCO CINAHL.

Exclude - Not a Primary Study

Heaton A. Public health, managed care, and pharmacy: Evolving trifecta. *J Manag Care Pharm* 2001;7(1):12-6. Database: IPA.

Exclude - Not a Primary Study

Hedge JA. Developing a bar code scanning interface to increase efficiency and accuracy. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Heffner JE, Brower K, Ellis R, et al. Using intranet-based order sets to standardize clinical care and prepare for computerized physician order entry. *Joint Commission Journal on Quality & Safety* 2004;30(7):366-76. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Heidenreich PA, Gholami P, Sahay A, et al. Clinical reminders attached to echocardiography reports of patients with reduced left ventricular ejection fraction increase use of beta-blockers: A randomized trial. *Circulation* 2007;115(22):2829-34. Database: Embase Sept 22-09.

Exclude - Not MMIT

Heijman JA, Koedijk FG, Van Der Velde E, et al. VIS-medication. Development and implementation of the medication module of a hospital ward information system using a pen-based computer. *Ziekenhuisfarmacie* 1998;14(4):199-204. Database: IPA.

Exclude - Unable to Retrieve Foreign

Heimly V. Consent-based access to core EHR information: the SUMO-project. *Studies in Health Technology & Informatics* 2008;136:431-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Heimly V, Berntsen KE. Consent-based access to core EHR information. Collaborative approaches in Norway. *Methods Inf Med* 2009;48(2):144-8. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Heindl B, Schmidt R, Schmid G, et al. A case-based consiliarius for therapy recommendation (ICONS): computer-based advice for calculated antibiotic therapy in intensive care medicine. *Computer Methods & Programs in Biomedicine* 1997;52(2):117-27. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Heininger A, Niemetz AH, Keim M, et al. Implementation of an interactive computer-assisted infection monitoring program at the bedside. *Infect Control Hosp Epidemiol* 1999;20(6):444-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Heinrichs W, Monk S, Eberle B. Automated anesthesia record systems. *Anaesthesist* 1997;46(7):574-82. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Heinze D, Feller P, McCorkle J et al. Computer-assisted auditing for high-volume medical coding. *Perspectives in Health Information Management, CAC Proceedings, 2006.* http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_032029.pdf Grey Lit.

Exclude - Not a Primary Study

Heller A. Patient safety technologies that won't break the bank. *Pharmacy Practice News* 1964;31(6):64 Database: IPA.

Exclude - Not a Primary Study

Helm GR, Fluno DA. Significance of order entry warning screens: Improving their effectiveness. *Ashp Midyear Clinical Meeting* 1999;34: Database: IPA.

Exclude - Not a Primary Study

Helmons PJ, Grouls RJ, Roos AN, et al. Using a clinical decision support system to determine the quality of antimicrobial dosing in intensive care patients with renal insufficiency. *Quality & Safety in Health Care* 2010;19(1):22-6.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010579264&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2345&accno=2010579264
EBSCO CINAHL.

Exclude - Not MMIT

Hemmerling TM. Automated anesthesia. *Current Opinion in Anaesthesiology* 2009;22(6):757-63. PMID:19812483 OVID MEDLINE.

Exclude - No Outcomes of Interest

Hemstreet BA, Stolpman N, Badesch DB, et al. Potassium and phosphorus repletion in hospitalized patients: implications for clinical practice and the potential use of healthcare information technology to improve prescribing and patient safety. *Current Medical Research & Opinion* 2006;22(12):2449-55. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Henderson J, Britt H, Miller G. Extent and utilisation of computerisation in Australian general practice. *Med J Aust* 2006;185(2):84-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Henderson J, Miller G, Britt H, et al. Effect of computerisation on Australian general practice: does it improve the quality of care? *Qual Prim Care* 2010;18(1):33-47. PMID:20359411 OVID MEDLINE.

Exclude - Not MMIT

Hengen KJ, Malone PM. Electronic prescription in pharmacy. *Ashp Midyear Clinical Meeting* 1997;32: Database: IPA.

Exclude - Not a Primary Study

Henneman PL, Fisher DL, Henneman EA, et al. Providers do not verify patient identity during computer order entry. *Acad Emerg Med* 2008;15(7):641-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Henricks W. Information system issues facing clinical laboratories serving complex integrated delivery systems. *J Healthc Inf Manag* 2000;14(3):55-67.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=60716> Grey Lit.

Exclude - Not a Primary Study

Hensing J, Dahlen D, Warden M, et al. Measuring the benefits of IT-enabled care transformation. *Healthc Financ Manage* 2008;62(2):74-80. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Herman KA. Utilizing a point of care documentation system for increased medication administration accuracy. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Hermes-DeSantis ER, Turck CJ. Combination therapy strategies for minimizing medication errors. *Hosp Pharm* 2006;41(Suppl. 1):S16-S22 Database: IPA.

Exclude - Not MMIT

Hernan MA, Cole SR. Invited commentary: Causal diagrams and measurement bias. *Am J Epidemiol* 2009;170(8):959-62. OVID EMBASE.
 Exclude - Not a Primary Study

Hernandez Martin J., Montero Hernandez M., Font Noguera I., et al. Evaluation of pharmaceutical interventions due to drug-related problems in hospitalized patients. *Atencion Farmaceutica* 2009;11(2):80-94. Database: Embase Sept 22-09.
 Exclude - Unable to Retrieve Foreign

Hershey C, Porter D, Breslau D, et al. Influence of simple computerized feedback on prescription charges in an ambulatory care: A randomized clinical trial. *Med Care* 1986;24(6):472-81. Exclude - Not MMIT

Hershey C, Goldberg H, Cohen D. The effect of computerized feedback coupled with a newsletter upon outpatient prescribing charges: A randomized controlled trial. *Med Care* 1988;26(1):88-94. Database: PsycINFO.
 Exclude - Not MMIT

Hess R, Bryce C, McTigue K, et al. The diabetes patient portal: Patient perspectives on structure and delivery. *Diabetes Spectrum* 2006;19(2):106-10. Database: Embase Sept 22-09.
 Exclude - Not MMIT

Hess V. Adjuvant!Online--an Internet-based decision tool for adjuvant chemotherapy in early breast cancer. *Ther Umsch* 2008;65(4):201-5. Database: Ovid MEDLINE(R).
 Exclude - Unable to Retrieve Foreign

Hickling K, Begg E, Moore M. A prospective randomised trial comparing individualised pharmacokinetic dosage prediction for aminoglycosides with prediction based on estimated creatinine clearance in critically ill patients. *Intensive Care Med* 1989;15(4):233-7. Exclude - Not MMIT

Hicks D. Scan to e-mail as a low-cost paperless orders transmission system. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
 Exclude - Not a Primary Study

Hidalgo Correias FJ, Bermejo Vicedo M.T., De Juana Velasco P., et al. Implementing an electronic medical prescription program in an INSALUD hospital. *Farmacia Hospitalaria* 2002;26(6):327-34. Database: Embase Sept 22-09.
 Exclude - Not MMIT

Hider P. Electronic prescribing: A critical appraisal of the literature. *NZHTA Report* 2002;5(2):1-113. Grey Lit.
 Exclude - Not a Primary Study

Hidle U. Implementing technology to improve medication safety in healthcare facilities: a literature review. *J N Y State Nurses Assoc* 2007;38(2):4-9. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Higgins DF. How wholesalers are using the computer to help you. *Pharm Times* 1974;40(10):50-3. Database: IPA.
 Exclude - Not a Primary Study

Higuchi K, Tsukamoto T, Nakano M, et al. Development and estimation of the management system for dispensing. Japanese Journal of Hospital Pharmacy 1992;18(6):584-93. Database: IPA.

Exclude - Not MMIT

Higuchi K, Tsukamoto T, Nakano M, et al. Estimation of the supporting functions for prescription making in an order entry system. Japanese Journal of Hospital Pharmacy 1993;19(6):555-66. Database: IPA.

Exclude - Unable to Retrieve Foreign

Higuchi N, Ichikawa N, Mine T, et al. Contribution to medical risk management of computerized prescription order entry systems - Improvement of master maintenance system using a commercially available order entry program. Japanese Journal of Pharmaceutical Health Care & Sciences 2004;30(6):382-8. Database: IPA.

Exclude - Not a Primary Study

Hill DA, Cacciatore M, Lamvu GM. Electronic prescribing influence on calcium supplementation: a randomized controlled trial. Am J Obstet Gynecol 2010;202(3):236-5. PMID:20044067 OVID MEDLINE.

Exclude - Not MMIT

Hilmas E, Sowan A, Gaffoor M, et al. Implementation and evaluation of a comprehensive system to deliver pediatric continuous infusion medications with standardized concentrations. Am J Health Syst Pharm 2010;67(1):58-69. PMID:20044370 OVID MEDLINE.

Exclude - No Outcomes of Interest

Hing ES, Burt CW, Woodwell DA. Electronic medical record use by office-based physicians and their practices: United States, 2006. Adv Data 2007;(393):1-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Hing E, Burt C, Woodwell D. Electronic medical record use by office-based physicians and their practices: United States, 2006. U.S. Department of Health and Human Services; 2007. http://library.ahima.org/xpedio/groups/public/documents/government/bok1_038413.pdf Grey Lit.

Exclude - Not MMIT

Hingl JF. Computerized physician order entry in a pediatric teaching institution: Design process, implementation and benefits. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Hingorani AD, Vallance P. Simple computer program for guiding management of cardiovascular risk factors and prescribing. Br Med J 1999;318(7176):101-5. Database: IPA.

Exclude - Not MMIT

Hirsch JD, Oen A, Robertson S, et al. Patient request for pharmacist counseling and satisfaction: Automated prescription delivery system versus regular pick-up counter. Journal of the American Pharmacists Association: JAPhA 2009;49(1):73-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Hitchens K. Hospitals controlling errors one year after tragedy. Hospital Pharmacist Report 2003;10(5):22-6. Database: IPA.

Exclude - Not a Primary Study

Ho G, Wheatley D, Scialfa C. Age differences in trust and reliance of a medication management system. Interacting with Computers 2005;17(6):690-710. Database: PsycINFO.

Exclude - Not a Primary Study

Ho L, Moh M, Walker Z and others. A prototype on RFID and sensor networks for elder healthcare: Progress report. In Philadelphia, PA, United states: Association for Computing Machinery; 2005. p.70-5.2005529621756

Database: Compendex.

Hobbs FD, Delaney BC, Carson A, et al. A prospective controlled trial of computerized decision support for lipid management in primary care. Fam Pract 1996;13(2):133-7.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Hoch I, Heymann AD, Kurman I, et al. Countrywide computer alerts to community physicians improve potassium testing in patients receiving diuretics. J Am Med Assoc 2003;10(6):541-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Hodari KT, Nanton JR, Carroll CL, et al. Adherence in dermatology: a review of the last 20 years. Journal of Dermatological Treatment 2006;17(3):136-42. Database: Ovid

MEDLINE(R).

Exclude - Not MMIT

Hodges N, Preston R. Practical consideration for implementation of new technology within a large health-system market. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Hoekstra M, Vogelzang M, Verbitskiy E, et al. Health technology assessment review: Computerized glucose regulation in the intensive care unit--how to create artificial control. Critical Care (London, England) 2009;13(5):223 PMID:19849827 OVID MEDLINE.

Exclude - Not a Primary Study

Hoffman LJ, Bulpitt AE, Lawton GC, et al. Implementation of an electronic medication administration record system for Point-of-Care Scanning: using bar coding and automated medication distribution technologies. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Hoffman MV, LaGory DN, Horn JP. Wireless, hand held device used by physicians to prescribe medication in an outpatient setting. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.

Exclude - Not a Primary Study

Hofmann UG, Michalski T, Tronnier V, et al. Personalized Pain Diary on a Handheld Device. IFMBE Processing 2009;25(5):133-6. Engineering Village Compendex and Inspec.

Exclude - No Outcomes of Interest

Hogan PA, Teller-Kook MB. Patient-oriented pharmacy practice model. Ashp Midyear Clinical Meeting 1994;29: Database: IPA.
Exclude - Not a Primary Study

Hohmann C, Radziwill R., Klotz J.M., et al. Implementation of a documentation system for drug-related problems in hospital (APS-Doc). Krankenhauspharmazie 2008;29(10):435-41. Database: Embase Sept 22-09.
Exclude - Unable to Retrieve

Hokanson JA, Keith MR, Guernsey BG, et al. Potential use of bar codes to implement automated dispensing quality assurance programs. Hosp Pharm 1985;20(5):327-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Hokanson JA, Richard PL, Doutre WH, et al. Key role of software in implementing computerbased information systems for the hospital pharmacy. Hosp Pharm 1988;18(3):113-6. Database: IPA.
Exclude - Unable to Retrieve

Holbrook A, Grootendorst P, Willison D, et al. Can current electronic systems meet drug safety and effectiveness requirements? AMIA 2005;335-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Holbrook A, Labiris R, Goldsmith CH, et al. Influence of decision aids on patient preferences for anticoagulant therapy: A randomized trial. Can Med Assoc J 2007;176(11):1583-7. Database: IPA.
Exclude - Not MMIT

Holbrook AM, Janjusevic V, Goldsmith CH, et al. A comprehensive appropriateness of prescribing questionnaire was validated by nominal consensus group. J Clin Epidemiol 2007;60(10):1022-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Holden RJ, Karsh B. A review of medical error reporting system design considerations and a proposed cross-level systems research framework. Hum Factors 2007;49(2):257-76. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Holder SD. Effectively preparing a pharmacy for a barcode medication verification project. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Hollowell LA, Perry TW, Alexander ML. Operating room controlled drug accountability incorporating pharmacy witnessed waste and electronic documentation. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.
Exclude - Not a Primary Study

Holmberg A, Brenninkmeijer R. Computerised protocol-driven prescribing. Disease Management & Health Outcomes 1998;3(3):105-14. Database: CINAHL.
Exclude - Not a Primary Study

Holtorf A-P, McAdam-Marx C, Schaaf D, et al. Systematic review on quality control for drug management programs: Is quality reported in the literature? BMC Health Serv Res 2009;9(38):1-11. OVID EMBASE.

Exclude - Not MMIT

Holzmueller CG, Pronovost P, Dickman F, et al. Creating the Web-based Intensive Care Unit Safety Reporting System. J Am Med Inform Assoc 2005;12(2):130-9. Database: Embase Sept 22-09.

Exclude - Not MMIT

Homme MB, Reynolds KK, Valdes R, Jr., et al. Dynamic pharmacogenetic models in anticoagulation therapy. Clin Lab Med 2008;28(4):539-52. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Honda Y, Yasunari K, Ogata M, et al. Dispensed drug information services for outpatients at Kumamoto University Hospital. Japanese Journal of Hospital Pharmacy 1998;24(5):576-83. Database: IPA.

Exclude - Not MMIT

Honeybourne C, Sutton S, Ward L. Knowledge in the palm of your hands: PDAs in the clinical setting. Health Information & Libraries Journal 2009;23(1):51-9. Database: CINAHL.

Exclude - Not MMIT

Hook J and Cusack C. Ambulatory computerized provider order entry: Findings from the AHRQ health IT portfolio. AHRQ; 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_890425_0_0_18/08-0063-EF.pdf Grey Lit.

Exclude - Not a Primary Study

Hook J, Pearlstein J, Samarth A et al. Using barcode medication administration to improve quality and safety: Findings from the AHRQ health IT portfolio. AHRQ; 2008.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_846906_0_0_18/09-0023-EF_bcma.pdf Grey Lit.

Exclude - Not a Primary Study

Hoover DA, Conklin CV. Successful implementation of bar code enabled point of care (BPOC): More than drug bar coding. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Hope C, Overhage J, Seger A, et al. A tiered approach is more cost effective than traditional pharmacist-based review for classifying computer-detected signals as adverse drug events. Journal of Biomedical Informatics 2003;36(1-2):92-8. Database: Embase Sept 22-09.

Exclude - Not MMIT

Hoppe-Tichy T. Bar code and computer-assisted ordering of medication. Krankenhauspharmazie 1993;14(6):298-9. Database: IPA.

Exclude - Unable to Retrieve

Hor CP, O'Donnell JM, Murphy AW, et al. General practitioners' attitudes and preparedness towards clinical decision support in e-prescribing (CDS-eP) adoption in the West of Ireland: a cross sectional study. BMC Med Inform Decis Mak 2010;10(1-3):8 Engineering Village Compendex and Inspec.

Exclude - No Outcomes of Interest

Horn J, Hansten P, Osborn J, et al. Clinical decision support systems: Customization of a drug interaction database to avoid alert fatigue. Ashp Summer Meeting 2009;65: Database: IPA.

Exclude - Not a Primary Study

Horn JP, Lesko A, Wissinger B. e-prescribing in a hospital clinic setting. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Horn JR. Computerized drug interaction screening: Pitfalls and future directions. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.

Exclude - Not a Primary Study

Horn MS. Does computerized ordering increase errors? JCOM 2005;12(5):239-40. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Horsky J, Kuperman GJ, Patel VL. Comprehensive analysis of a medication dosing error related to CPOE. J Am Med Inform Assoc 2005;12(4):377-82. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Horsky J, Kaufman DR, Patel VL. AMIA 2005;350-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Hripsak G, Kaushal R, Johnson K, et al. The United Hospital Fund meeting on evaluating health information exchange. Journal of Biomedical Informatics 2007;40(6 Suppl):S3-S10 http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PJ6GKF-1&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=4&_fmt=full&_orig=browse&_srch=doc-

info(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=414d9d9025e2ee1d6964d2632bb652d4 Grey Lit.

Exclude - Not a Primary Study

Hripsak G, Clayton PD, Jenders RA, et al. Design of a clinical event monitor. Computers & Biomedical Research 1996;29(3):194-221. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hsieh TC, Kuperman GJ, Jaggi T, et al. Characteristics and consequences of drug allergy alert overrides in a computerized physician order entry system. J Am Med Inform Assoc 2004;11(6):482-91. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Hu PJH, Wei C, Cheng T, et al. Predicting adequacy of vancomycin regimens: A learning-based classification approach to improving clinical decision making. *Decision Support Systems* 2007;43(4):1226-41. 9627801

Database: Inspec.

Exclude - Not MMIT

Huang HY, Lee TT. Applying bar code technology in nurses' medication administration. *Hu Li Tsa Chih - Journal of Nursing* 2009;56(2):70-4. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Hubbell PJ. The pharmacy as part of an integrated hospital information system: a focus on benefits. *Hosp Pharm* 1994;29(5):440-1. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hudgins J. Moving beyond vanilla. *Employee Benefit News* 2009;23(6):22-4.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=40537885&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Huertas Fernandez MJ, Baena-Canada JM, Martinez Bautista MJ, et al. Impact of computerised chemotherapy prescriptions on the prevention of medication errors. *Clinical & Translational Oncology: Official Publication of the Federation of Spanish Oncology Societies & of the National Cancer Institute of Mexico* 2006;8(11):821-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Huffines S. View from the other side: Changes in pharmacy practice after a decade of an integrated computerized provider order entry (CPOE) system. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Huffines SK, Sullivan TM. Pharmacy involvement in addressing substance abuse and drug diversion in a medical center. *ASHP Annual Meeting* 1996;53: Database: IPA.

Exclude - Not a Primary Study

Huffines SK, Hargrove FR, Knight JR. Strategies for successful implementation of medication online order entry system. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Hug BL, Witkowski DJ, Sox CM, et al. Adverse drug event rates in six community hospitals and the potential impact of computerized physician order entry for prevention. *J Gen Intern Med* 2010;25(1):31-8. OVID EMBASE.

Exclude - Not MMIT

Hughes D. When drugs don't work: Economic assessment of enhancing compliance with interventions supported by electronic monitoring devices. *Pharmacoeconomics* 2007;25(8):621-35. Database: PsycINFO.

Exclude - Not MMIT

Hughes DK, Farrar KT. Drug name confusion reduced by computerized prescribing. Ashp Midyear Clinical Meeting 1993;28: Database: IPA.
Exclude - Not a Primary Study

Hume M. Computer-aided drug selection can sharply cut adverse events. Qual Lett Healthc Lead 1999;11(3):10-2. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hummer J. Lean process in pharmacy renewal. AAACN Viewpoint 2010;32(1):1, 7-11. <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010547006&site=ehost-live>;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1228&accno=2010547006
EBSCO CINAHL.
Exclude - Not a Primary Study

Humphries TL, Delate T, Helling DK, et al. Impact of an automated dispensing system in outpatient pharmacies. Journal of the American Pharmacists Association: JAPhA 2008;48(6):774-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Hunt DL, Haynes RB, Hayward RS, et al. Automated direct-from-patient information collection for evidence-based diabetes care. Proceedings/AMIA Annual Fall Symposium 1997;1997:81-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Hunt DL, Haynes RB, Hanna SE, et al. Effects of computer-based clinical decision support systems on physician performance and patient outcomes: a systematic review. JAMA 1998;280(15):1339-46. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Hurdle JF, Weir CR, Roth B, et al. Critical gaps in the world's largest electronic medical record: Ad Hoc nursing narratives and invisible adverse drug events. AMIA 2003;309-12. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Hurley AC, Lancaster D, Hayes J, et al. The Medication Administration System--Nurses Assessment of Satisfaction (MAS-NAS) scale. Journal of Nursing Scholarship 2006;38(3):298-300. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Hurley D. Automated medication system-benefits for nursing. In Washington; DC: 1994. p. 1050. Grey Lit.
Exclude - Not a Primary Study

Hurley D. Lessons learned on the bar-coding trail. Pharmacy Practice News 2003;30(2):14-6. Database: IPA.
Exclude - Not a Primary Study

Hurley S, Dziukas L, McNeil J, et al. A randomized controlled clinical trial of pharmacokinetic theophylline dosing. Am Rev Respir Dis 1986;134(6):1219-24. Exclude - Not MMIT

Husch M, Sullivan C, Rooney D, et al. Insights from the sharp end of intravenous medication errors: implications for infusion pump technology. *Quality & Safety in Health Care* 2005;14(2):80-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Hussein G. MedStorm(TM): Innovative patient safety software for optimal therapeutics. *Clinical Research & Regulatory Affairs* 2003;20(3):327-30. Database: IPA.

Exclude - Not a Primary Study

Hussey TC. Physician order entry: Cure for medication errors or a problem in disguise? *Pharm Times* 2007;67(10):14 Database: IPA.

Exclude - Not a Primary Study

Husson M. Theriaque: Independent-drug database for good use of drugs by health practitioners. *Ann Pharm Fr* 2008;66(5-6):268-77. Database: Embase Sept 22-09.

Exclude - Not MMIT

Huston T. POC PT/INR testing: a better choice for patients and providers. *MLO Med Lab Obs* 2009;41(11):28 PMID:19960734 OVID MEDLINE.

Exclude - Not a Primary Study

Huston T. Trends in POCT for coagulation. *MLO Med Lab Obs* 2009;41(8):26
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010386048&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=246&accno=2010386048> EBSCO CINAHL.

Exclude - Not a Primary Study

Hwang HG, Chang IC, Hung WF, et al. The design and evaluation of clinical decision support systems in the area of pharmacokinetics. *Medical Informatics & the Internet in Medicine* 2004;29(3-4):239-51. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Hwang H, Chang I, Hung W, et al. The design and evaluation of clinical decision support systems in the area of pharmacokinetics. *Med Inform Internet Med* 2004;29(3-4):239-51. 8236878

Database: Inspec.

Exclude - No Outcomes of Interest

Hwang RW, Herndon JH. The business case for patient safety. *Clinical Orthopaedics & Related Research* 2007;457:21-34. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hynniman CE. Drug product distribution systems and departmental operations. *Am J Hosp Pharm* 1991;48(10 Suppl):S24-S35 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Hynniman CE. Medication dispensing in the future. *Pharm Times* 1992;58(Jun): Database: IPA.

Exclude - Not a Primary Study

Hypponen H, Salmivalli L, Nykanen P, et al. Testing a theoretical framework for interdisciplinary IT evaluation: The case of the Finnish Electronic Prescription. *International Journal of Healthcare Technology and Management* 2007;8(1-2):42-65. Database: Embase Sept 22-09.

Exclude - Not MMIT

I'Epercll K, Rudolf M., Pearson M., et al. General practitioner prescribing habits in asthma/COPD. *Asthma in General Practice* 1997;5(2):29-30. Database: Embase Sept 22-09.

Exclude - Not MMIT

Iizuka K, Hosokawa M, Hashimoto G, et al. Role of pharmacy practice in improving anti-cancer chemotherapy medication safety in a national university hospital in Japan. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Imperiali RD, Clapp MJ, Santoro JP. Evaluation and cost-justification of an automated medication dispensing system. *ASHP Annual Meeting* 1996;53: Database: IPA.

Exclude - Not a Primary Study

Ingersoll KS, Cohen J. The impact of medication regimen factors on adherence to chronic treatment: A review of literature. *J Behav Med* 2008;31(3):213-24. Database: CINAHL.

Exclude - Not a Primary Study

Isaac T, Weissman JS, Davis RB, et al. Overrides of medication alerts in ambulatory care. *Arch Intern Med* 2009;169(3):305-11. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Isaksen SF, Jonassen J, Malone DC, et al. Estimating risk factors for patients with potential drug-related problems using electronic pharmacy data. IMPROVE investigators. *Ann Pharmacother* 1999;33(4):406-12. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ishar R, Aydin C E. Predicting effective use of hospital computer systems. In Washington, DC, USA: IEEE Comput. Soc. Press; 1988. p.862-8.3372299

Database: Inspec.

Exclude - Not MMIT

Isherwood K. Computers and hospital pharmacy. *New Zealand Pharmacy* 1983;3(1):35-6. Database: IPA.

Exclude - No Outcomes of Interest

Ishimoto K, Ishihara M, Okada N, et al. Establishment and evaluation of a system for preventing mis-administration of powder using bar codes printed on drug envelopes. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2003;123(5):331-6. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Exclude - Unable to Retrieve Foreign

Ishizuka H, Waki Y, Horizuti M, et al. Development and evaluation of a new automated dispensing system. *Int J Biomed Comput* 1995;38(2):167-76. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Iteen SA, Ceppaglia J. Investigational drug information and order entry through use of a hospital-wide information system. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.
Exclude - Not a Primary Study

Iteen SA, Ceppaglia J. Investigational drug information through a hospitalwide computer system. *Am J Hosp Pharm* 1992;49(11):2746-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Iten A, Perrier A, Lovis C. Deployment of a computerized physician order entry: description of the process and challenges. *Revue Medicale Suisse* 2006;2(83):2344-6. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Ivey MF. Why consider a multiple hospital site medication cart fill? *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Ivey MF. Preparing for computerized physician order entry implementation at the health alliance of Greater Cincinnati. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
Exclude - Not a Primary Study

Iwata M, Nawata S, Takahashi Y, et al. The development of a new dispensing system for the appropriate use of injectable medicine - H [subscript] 2-receptor antagonist and proton pump inhibitor. *Japanese Journal of Pharmaceutical Health Care & Sciences* 2003;29(5):593-8.
Database: IPA.
Exclude - Not MMIT

Jacobs B. Electronic medical record, error detection, and error reduction: a pediatric critical care perspective. *Pediatric Critical Care Medicine* 2007;8(2:Suppl):Suppl-20 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Jacobs GS, Ernst TF, Sorrentino AC, et al. Computer assisted drug use evaluation: One hospital's approach. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.
Exclude - Not a Primary Study

Jacobs RA, Patchin GM, Savage LR. Impact on production and efficiency in a VA outpatient pharmacy setting following the implementation of an automated dispensing system, the Baker APS Pharmacy 2000 system with an Autoscript III robot. *ASHP Annual Meeting* 2000;57: Database: IPA.
Exclude - Not a Primary Study

Jain A, Atreja A, Harris C. Rapid notification of a drug recall using information technology. *JCOM* 2005;12(4):182-6. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

James CC. Look Doc, no wires. *Communications News* 2003;41(3):12-4. 7782452
Database: Inspec.
Exclude - No Outcomes of Interest

Janczak DR, Robinson E, Huenerberg K. Maximizing patient safety in a medical oncology practice: A journey through failure mode effects analysis to computerized physician order entry. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Janknegt R, Jacklin A, Kraemer I. Can a closed loop system add value above and beyond computerised physician order entry? *Ejhp Science* 2006;12(5):66-8. Database: IPA.

Exclude - Not a Primary Study

Jannett TC, Kay GN, Sheppard LC. Automated administration of lidocaine for the treatment of ventricular arrhythmias. *Med Prog Technol* 1990;16(1-2):53-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Janssen B, Menke R, Pourhassan F, et al. Guidelines based on decision support software. Quality management in neurological outpatient schizophrenia treatment. *Der Nervenarzt* 2006;77(5):S67-S75 Database: PsycINFO.

Exclude - Not a Primary Study

Jao C, Hier D, Galanter W. Automating the maintenance of problem list documentation using a clinical decision support system. *AMIA 2008;Annual:Symposium* PMID:18998927 OVID MEDLINE.

Exclude - No Outcomes of Interest

Jao CS, Hier DB. Overcoming limitations of data entry for the semi-automated detection of drug orphans in the EMR. *AMIA 2006;967* Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Jao C S, Hier D B, Galanter W L. Using clinical decision support to maintain medication and problem lists. A pilot study to yield higher patient safety. In Piscataway, NJ, USA: IEEE; 2008. p.739-43.10560314

Database: Inspec.

Exclude - No Outcomes of Interest

Jelliffe RW, Buell J, Kalaba R. Reduction of digitalis toxicity by computer-assisted glycoside dosage regimens. *Ann Intern Med* 1972;77(Dec):891-906. Database: IPA.

Exclude - Not MMIT

Jellinek SP, Cohen V, Likourezos A, et al. Analyzing a health-system's use of unfractionated heparin to ensure optimal anticoagulation. *J Pharm Technol* 2005;21(2):69-78. Database: Embase Sept 22-09.

Exclude - Not MMIT

Jenders RA. Standards in health information technology: promise and challenges. *AMIA 2007;1179-80*. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Jenkins G, Johnston PE, Patel NR, et al. The epidemiology of medication prescribing errors in the emergency department. *AMIA 2006;968* Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Jenrette DE, Mikell NG, Bromley HR. Evolving role of the ambulatory care clinical pharmacist: Integrating clinical and distributive functions. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Jenrette DE, Mikell NG, DeBruin A. Standardizing high alert intravenous infusions. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Jensen RJ, Anderson SA. CQI team approach to a point of use system implementation. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Jessee W. Healthcare's green shoots: reform, quality and HIT all moving forward, but there's still much to do. *Mod Healthc* 2009;39(41):22

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010441028&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010441028 EBSCO CINAHL.

Exclude - Not a Primary Study

Jha AK, Orav EJ, Ridgway AB, et al. Does the Leapfrog program help identify high-quality hospitals? *Jt Comm J Qual Patient Saf* 2008;34(6):318-25. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Jha A, Kuperman G, Teich J, et al. Identifying adverse drug events: Development of a computer-based monitor and comparison with chart review and stimulated voluntary report. *J Am Med Inform Assoc* 1998;5(3):305-14. Exclude - No Outcomes of Interest

Jha A, DesRoches C, Campbell E, et al. Use of electronic health records in U.S. hospitals. *The New England Journal of Medicine* 2009;360(16):1628-38. Database: PsycINFO.

Exclude - Not MMIT

Ji S, Matsumura Y, Kuwata S, et al. Creation of a master table for checking indication and contraindication of medicine from a knowledge base linked with a thesaurus. *J Med Syst* 2004;28(6):561-73. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Joanna Briggs Institute. Strategies to reduce medication errors with reference to older adults. *Nurs Stand* 2006;20(41):53-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

John G, Peter JV, Chacko B, et al. A computer-assisted recording, diagnosis and management of the medically ill system for use in the intensive care unit: A preliminary report. *Indian Journal of Critical Care Medicine* 2009;13(3):136-42. OVID EMBASE.

Exclude - No Outcomes of Interest

Johnson CL, Carlson RA, Tucker CL, et al. Using BCMA software to improve patient safety in Veterans Administration Medical Centers. *J Healthc Inf Manag* 2002;16(1):46-51.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Johnson C, Carlson R, Tucker C, et al. Using BCMA software to improve patient safety In veterans administration medical centers. J Healthc Inf Manag 2002;16(1):46-51.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=60723> Grey Lit.

Exclude - Not a Primary Study

Johnson KB, Davison C. Information technology: Its importance to child safety. Ambulatory Pediatrics 2004;4(1):64-72. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Johnson KB, FitzHenry F. Case report: activity diagrams for integrating electronic prescribing tools into clinical workflow. J Am Med Inform Assoc 2006;13(4):391-5.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Johnson K, Gadd C. Playing smallball: Approaches to evaluating pilot health information exchange systems. Journal of Biomedical Informatics 2007;40:S21-S26

[http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PJM9N4-](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PJM9N4-1&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=7&_fmt=full&_orig=browse&_srch=doc-)

[1&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=7&_fmt=full&_orig=browse&_srch=doc-](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PJM9N4-1&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=7&_fmt=full&_orig=browse&_srch=doc-)

[info\(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume\)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=1a841657f1e2958fb9b05ae42c4ae135](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PJM9N4-1&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=7&_fmt=full&_orig=browse&_srch=doc-info(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=1a841657f1e2958fb9b05ae42c4ae135) Grey Lit.

Exclude - Not a Primary Study

Johnson MT, Somani SM, Abramowitz PW, et al. Evaluation of an automated narcotic dispensing system. ASHP Annual Meeting 1991;48: Database: IPA.

Exclude - Not a Primary Study

Johnson N. Use of technology to improve drug therapy outcomes. Hosp Formul 2000;35(Jan):65-6. Database: IPA.

Exclude - No Outcomes of Interest

Johnson SC, Generali JA, Habershaw TA, et al. Clinical and economic impact of avoiding computer identified drug interactions. Ashp Midyear Clinical Meeting 1989;24: Database: IPA.

Exclude - Not a Primary Study

Johnson T, Currie G, Keill P, et al. NewYork-Presbyterian Hospital: translating innovation into practice. Jt Comm J Qual Patient Saf 2005;31(10):554-60. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Johnson VR, Hummel J, Kinninger T, et al. Immediate steps toward patient safety. Healthc Financ Manage 2004;58(2):56-61. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Johnston B, Wheeler L, Deuser J, et al. Outcomes of the Kaiser Permanente tele-home health research project. Arch Fam Med 2000;9(1):40-5. Grey Lit.

Exclude - Not MMIT

Johnston D, Pan E, Walker J et al. Patient safety in the physician's office: Assessing the value of ambulatory CPOE. California HealthCare Foundation; 2004.

<http://www.chcf.org/documents/ihealth/PatientSafetyInPhysiciansOfficeACPOE.pdf> Grey Lit.

Exclude - Not a Primary Study

Johnston D, Pan E, Walker J. The value of CPOE in ambulatory settings. J Healthc Inf Manag 2004;18(1):5-8. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=60710> Grey Lit.

Exclude - Not a Primary Study

Johnston G, Halpin E. Automated compliance: Reducing costs and maintaining quality. Pharmaceutical Technology Europe 2003;15(9):59-63. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Johnston ME, Langton KB, Haynes RB, et al. Effects of computer-based clinical decision support systems on clinician performance and patient outcome. A critical appraisal of research.[see comment]. Ann Intern Med 1994;120(2):135-42. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Jones CA, Clement L.T., Hanley-Lopez J., et al. The Breathmobile Program: Structure, implementation, and evolution of a large-scale, urban, pediatric asthma disease management program. Disease Management 2005;8(4):205-22. Database: Embase Sept 22-09.

Exclude - Not MMIT

Jones D, Groller S. Keep scanning: Tips for bar-coding compliance. Nephrology Nursing Journal 2008;35(2):176 Database: CINAHL.

Exclude - Not a Primary Study

Jones JL. Strategies for implementing CPOE in a children's hospital. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Jones JL. Implementing computerized prescriber order entry in a children's hospital. Am J Health Syst Pharm 2004;61(22):2425-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Jones K, Swensen N, Holley M, et al. Automation in a pediatric hospital: Implementation of a dispensing robot. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Jones MA, Wyld DC. A magic pill? The emergence of radio frequency identification (RFID) technology in the pharmaceutical supply chain. Journal of Pharmaceutical Marketing & Management 2005;17(1):17-33. Database: IPA.

Exclude - Not MMIT

Jones MJ. Internet-based prescription of sildenafil: a 2104-patient series. J Med Internet Res 2001;3(1):E2-Mar Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Jones MJ, Thomasson W. Establishing guidelines for internet-based prescribing. *South Med J* 2003;96(1):1-5. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Jones MS, Day PL, Raisch DW, et al. System validation: Case study using bar code technology. *Ashp Midyear Clinical Meeting* 1995;30: Database: IPA.

Exclude - Not a Primary Study

Jones MS, Raisch DW, Garnand D, et al. Use of a remote computerized system for study documentation in clinical trials. *Drug Inf J* 1998;32(4):1153-63. Database: IPA.

Exclude - Not a Primary Study

Jones RC, Dickson-Spillmann M, Mather MJ, et al. Accuracy of diagnostic registers and management of chronic obstructive pulmonary disease: the Devon primary care audit. *Respiratory Research* 2008;9:62 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Jones RT, Sullivan M, Barrett D. INRstar: computerised decision support software for anticoagulation management in primary care. *Inform Prim Care* 2005;13(3):215-21.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Jones S, Moss J. Computerized provider order entry: strategies for successful implementation. *J Nurs Adm* 2006;36(3):136-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Jonietz E. Paperless medicine. *Technol Rev* 2003;106(3):58-64. 2003217477127

Database: Compendex.

Exclude - Not a Primary Study

Jordan FM, Morales V, Kueh ET, et al. Interfacing a computerized physician order entry system to a pharmacy system. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Jordan S, Jones R, Sargeant MP. Adverse drug reactions: managing the risk. *J Nurs Manag* 2009;17(2):175-84. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Jorgenson JA, Leiker TK, Herzog CC. Combined medication-and-supply automated delivery system in an ambulatory setting. *Hosp Pharm* 2002;37(8):828-32. Database: IPA.

Exclude - No Outcomes of Interest

Joseph,L. Use of handheld technology (PDA) and reduction in medication errors: an evidence-based pilot project Fairleigh Dickinson University. 2009. EBSCO CINAHL.

Exclude - Not MMIT

Josephson DC. Automation's emerging role as a new quality assurance tool for the long-term care pharmacist. *Consultant Pharmacist* 1998;13(Sep):1028-32. Database: IPA.

Exclude - Not a Primary Study

Joshi MC, Joshi HS, Tariq K, et al. A prospective study of medication errors arising out of look-alike and sound-alike brand names confusion. *International Journal of Risk & Safety in Medicine* 2007;19(4):195-201. Database: IPA.

Exclude - Not MMIT

Jovanovic L, Peterson CM. Toward normoglycemia: studies in computer-assisted insulin delivery. *Diabetes Educ* 1987;13(3):302-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Jozefczyk KG. Of medication orders and deli sandwiches. *Am J Health Syst Pharm* 2001;58(Jan 1):78-9. Database: IPA.

Exclude - Not a Primary Study

Juneja R, Roudebush C, Kumar N, et al. Utilization of a computerized intravenous insulin infusion program to control blood glucose in the intensive care unit. *Diabetes Technol Ther* 2007;9(3):232-40. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Juneja R, Roudebush CP, Nasraway SA, et al. Computerized intensive insulin dosing can mitigate hypoglycemia and achieve tight glycemic control when glucose measurement is performed frequently and on time. *Critical Care (London, England)* 2009;13(5):R163 PMID:19822000 OVID MEDLINE.

Exclude - Not MMIT

Jungst J. Decreasing unit-based cabinet overrides by implementing after-hours pharmacist order entry in a non-24-hour pharmacy hospital. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Jungst JD, Spear HW, Weerasooriya JM. Improved compliance with Joint Commission on Accreditation of Healthcare Organizations pharmacy review standard after electronic medication administration record implementation. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Juvany RR, Sevilla SD, de la Pena Oliete MD, et al. Optimising the quality of the unit dose dispensing process through the implementation of the semi-automated Kardex system. *Farmacia Hospitalaria* 2007;HOSP..(1):38-42. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Jylha V, Saranto K. Electronic documentation in medication reconciliation - a challenge for health care professionals. *Appl Nurs Res* 2008;21(4):237-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kadobayashi M, Sato T, Matsui H, et al. Windows drug information system in prescription order entry system. *Japanese Journal of Hospital Pharmacy* 1995;21(1):86-92. Database: IPA.

Exclude - No Outcomes of Interest

Kaelber D, Bates D. Health information exchange and patient safety. *Journal of Biomedical Informatics* 2007;40:S40-S45
[http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PKXBSM-2&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=10&_fmt=full&_orig=browse&_srch=doc-info\(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume\)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=3cd55f777b69f508bbdc176ce79a97e8](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PKXBSM-2&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=10&_fmt=full&_orig=browse&_srch=doc-info(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=3cd55f777b69f508bbdc176ce79a97e8) Grey Lit.
Exclude - Not a Primary Study

Kaelber D, Shah S, Pan E et al. The value of personal health records. *Center for Information Technology Leadership*; 2008. http://www.citl.org/_pdf/CITL_PHR_Report.pdf Grey Lit.
Exclude - Not a Primary Study

Kahan NR, Blackman S, Kutz C, et al. A pharmacoepidemiological approach to investigating inappropriate physician prescribing in a managed care setting in Israel. *Am J Manag Care* 2005;11(2):89-90. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kaiser RH. Introduction of the electronic health card, electronic prescription, health professional card, and other telematic applications. *Onkologe* 2004;10(11):1247-9. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Kajbjer K. Procurement of prescriber support systems. *Studies in Health Technology & Informatics* 2008;136:729-34. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Kaldy J. How quality care rises from confusion and controversy. *Consultant Pharmacist* 2004;19(8):697-712. Database: Embase Sept 22-09.
Exclude - Not MMIT

Kalis DJ, Tranchida GJ, Schultz SD. Reduction of serious medication errors through computerized physician order entry. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Kalmeijer MD, Holtzer W, van Dongen R, et al. Implementation of a computerized physician medication order entry system at the Academic Medical Centre in Amsterdam. *Pharmacy World & Science* 2003;25(3):88-93. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Kalvelage SJ, Roberts M. Use of a robot and bar codes in a community hospital. *ASHP Annual Meeting* 1994;51: Database: IPA.
Exclude - Not a Primary Study

Kamal J, Rogers P, Saltz J, et al. Information warehouse as a tool to analyze Computerized Physician Order Entry order set utilization: opportunities for improvement. *AMIA* 2003;336-40. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kanaan S, Burke-Beebe S, Rippen H. Consumer control of electronic personal health information: What does it mean? Why is it important? U.S. Department of Health and Human Services; 2006.
http://library.ahima.org/xpedio/groups/public/documents/government/bok1_035684.pdf Grey Lit.

Exclude - Not MMIT

Kang KS, Lao P, Schiller L, et al. The influence of technology (CPOE) on the nature of pharmacist interventions. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Kanmaz TJ, Haupt BA, Peterson AM. Comparison of manual and bar-code systems for documenting pharmacists' interventions. *Am J Health Syst Pharm* 1997;54:1623-6. Database: IPA.

Exclude - Not MMIT

Kanstrup AM, Nohr C. Gaming against medical errors: methods and results from a design game on CPOE. *Studies in Health Technology & Informatics* 2009;148:188-96.

PMID:19745250 OVID MEDLINE.

Exclude - No Outcomes of Interest

Kanuho D, Sweeney EV, Shafer C, et al. Implementation of bedside medication verification and electronic medication administration record: Experience from two community hospitals. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Kaplan GG, bates D, McDonald D, et al. Inappropriate use of intravenous pantoprazole: Extent of the problem and successful solutions. *Clinical Gastroenterology and Hepatology* 2005;3(12):1207-14. Database: Embase Sept 22-09.

Exclude - Not MMIT

Kaplan SL, Donnelly AJ, Markway GG. Enhancement of a computer generated drug interaction reporting system to improve departmental clinical productivity measurement: descriptive report. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Karnon J, McIntosh A, Dean J, et al. A prospective hazard and improvement analytic approach to predicting the effectiveness of medication error interventions. *Safety Science* 2007;45(4):523-39. Database: Embase Sept 22-09.

Exclude - Not MMIT

Karow HS. Creating a culture of medication administration safety: laying the foundation for computerized provider order entry. *Jt Comm J Qual Improv* 2002;28(7):396-402. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Karsh B. Clinical practice improvement and redesign: How change in workflow can be supported by clinical decision support. *AHRQ*; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_874022_0_0_18/09-0054-EF.pdf Grey Lit.

Exclude - Not a Primary Study

Kasai T, Iwata M, Oosaki K. Computerized patient monitoring system. *Fujitsu Scientific and Technical Journal* 1974;10(3):67-82. 1974120006840

Database: Compendex.

Exclude - No Outcomes of Interest

Katsumura Y, Yasunaga H, Imamura T, et al. Quality evaluation of economic studies for medical safety management. *Nippon Koshu Eisei Zasshi - Japanese Journal of Public Health* 2007;54(7):447-53. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kattan M, Crain E, Steinbach S, et al. A randomized clinical trial of clinician feedback to improve quality of care for inner-city children with asthma. *Pediatrics* 2006;117(6):e1095-e1103 Exclude - Not MMIT

Katz N, Houle B, Fernandez KC, et al. Update on prescription monitoring in clinical practice: a survey of prescription monitoring program administrators. *PAIN MED* 2008;9(5):587-94. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kauffman RH, Boggs GL. Improving the quality of patient care with hospital-wide computerized drug reaction program. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Kaushal R. Using information technology to reduce rates of medication errors in pediatric hospitals. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Kaushal R, Barker KN, Bates DW. How can information technology improve patient safety and reduce medication errors in children's health care?[see comment. *Archives of Pediatrics & Adolescent Medicine* 2001;155(9):1002-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kaushal R, Bates D, Landrigan C, et al. Medication errors and adverse drug events in pediatric inpatients. *Journal of the American Medical Association* 2001;285(16):2114-20. Database: Embase Sept 22-09.

Exclude - Not MMIT

Kaushal R, Bates DW. Information technology and medication safety: what is the benefit? *Quality & Safety in Health Care* 2002;11(3):261-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kaushal R, Shojania KG, Bates DW. Effects of computerized physician order entry and clinical decision support systems on medication safety: a systematic review. *Arch Intern Med* 2003;163(12):1409-16. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kaushal R, Jaggi T, Walsh K, et al. Pediatric medication errors: What do we know? What gaps remain? *Ambulatory Pediatrics* 2004;4(1):73-81. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Kavanaugh DL, Minard PD. Comparative evaluation of manual and automated dispensing systems. *Ashp Midyear Clinical Meeting* 1991;26: Database: IPA.
Exclude - Not a Primary Study

Kawahara NE, Jordan FM. Beyond order entry: Use of a patient care computer system as a tool for physician education and promotion of cost-effective therapy. *Ashp Midyear Clinical Meeting* 1988;23: Database: IPA.
Exclude - Not a Primary Study

Kawai S, Ishimoto K, Kamiya A. A checking system for contraindications using a prescription, injection and disease name ordering system and its evaluation. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2001;121(11):807-15. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Kawano DF, Pereira LR, Ueta JM, et al. Medication misadventures: How to minimize them? *Revista Brasileira de Ciencias Farmaceuticas/Brazilian Journal of Pharmaceutical Sciences* 2006;42(4):487-95. Database: IPA.
Exclude - Not MMIT

Kawazoe Y, Ohe K. An ontology-based mediator of clinical information for decision support systems: a prototype of a clinical alert system for prescription. *Methods Inf Med* 2008;47(6):549-59. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kaye M. Mandating of electronic prescriptions for Medicare patients. *Online Journal of Nursing Informatics* 2008;12(2): Database: CINAHL.
Exclude - Not a Primary Study

Kearney N, Kidd L, Miller M, et al. Utilising handheld computers to monitor and support patients receiving chemotherapy: results of a UK-based feasibility study. *Support Care Cancer* 2006;14(7):742-52. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Keet R. Essential characteristics of an electronic prescription writer. [Review] [7 refs]. *J Healthc Inf Manag* 1999;13(3):53-61. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Kellett J. Decision support and the appropriate use of fibrinolysis in myocardial infarction. *Eff Clin Pract* 2001;4(1):1-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kelly JJ, Sweigard KW, Shields K, et al. John M. Eisenberg Patient Safety Awards. Safety, effectiveness, and efficiency: a Web-based virtual anticoagulation clinic. *Joint Commission Journal on Quality & Safety* 2003;29(12):646-51. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kelly K, Vaida AJ. A point about levothyroxine. *Pharm Times* 2000;70(4):60 Database: IPA.
Exclude - Not a Primary Study

Kelly KM, Boecler L, Dumitru D. Design and implementation of a neonatal computerized prescriber order entry (CPOE) system. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Kelly MW, Steffensmeier JJ, Ernst ME. Five year experience with a computerized order entry system: Perceptions and reality. *Ashp Summer Meeting* 2004;61: Database: IPA.

Exclude - Not a Primary Study

Kemmer AN. EAN-128 - Standard for supplementary coding. *Manufacturing Chemist* 2004;61(Mar):33 Database: IPA.

Exclude - Not a Primary Study

Kenagy J. Computerized provider order entry in multispecialty ambulatory care practices: A quantitative evaluation of information systems success. *Dissertation Abstracts International: Section B: The Sciences and Engineering* 2008;68(8-B): Database: PsycINFO.

Exclude - Theses

Kenny T, Wilson R, Purves IN. Prescribing patient information leaflets may be better than prescribing drugs. *Br Med J* 1998;317:80-1. Database: IPA.

Exclude - Not MMIT

Kent SS, Boecler LA. Implementation of a pharmacy computer system integrated with computerized physician order entry and an electronic medical record. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Kent SS. Planning for electronic medical Record/Computerized prescriber order entry implementation. *Ashp Summer Meeting* 2006;63: Database: IPA.

Exclude - Not a Primary Study

Keohane CA, Bane AD, Featherstone E, et al. Quantifying nursing workflow in medication administration. *J Nurs Adm* 2008;38(1):19-26. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kerkenbush NL, Lasome CE. The emerging role of electronic diaries in the management of diabetes mellitus. *AACN Clin Issues* 2003;14(3):371-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kern LM, Dhopeswarkar R, Barron Y, et al. Measuring the effects of health information technology on quality of care: a novel set of proposed metrics for electronic quality reporting. *Joint Commission Journal on Quality and Patient Safety* 2009;35(7):359-69. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kern L, Kaushal R. Health information technology and health information exchange in New York State: New initiatives in implementation and evaluation. *Journal of Biomedical Informatics* 2007;40:S17-S20
[http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PKXBSM-1&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=6&_fmt=full&_orig=browse&_srch=doc-info\(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume\)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=9c8805d4c03d6575db63de8e9f745056](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WHD-4PKXBSM-1&_user=822378&_coverDate=12%2F01%2F2007&_rdoc=6&_fmt=full&_orig=browse&_srch=doc-info(%23toc%236848%232007%23999599993.8998%23674051%23FLA%23display%23Volume)&_cdi=6848&_sort=d&_docanchor=&_ct=11&_acct=C000044540&_version=1&_urlVersion=0&_userid=822378&md5=9c8805d4c03d6575db63de8e9f745056) Grey Lit.
Exclude - Not a Primary Study

Kerr K. The Electronic Health Record in New Zealand - Part 1. *Healthcare Review Online* 2004;8(1): Database: Embase Sept 22-09.
Exclude - Not MMIT

Kerr RA. CPOE in a community hospital setting: A pharmacist's perspective. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Kerr RA. Case study: Barcode implementation success factors. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Kershaw B, White RH, Mungall D, et al. Computer-assisted dosing of heparin. Management with a pharmacy-based anticoagulation service.[see comment]. *Arch Intern Med* 1994;154(9):1005-11. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kester K, Baxter J, Freudenthal K. Errors associated with medications removed from automated dispensing machines usina override function. *Hosp Pharm* 2006;41(6):535-7. Database: IPA.
Exclude - Not MMIT

Kester KA, Rietsch S, Payne G. Remote order entry: Impact on automated dispensing machine overrides. *Ashp Summer Meeting* 2004;64: Database: IPA.
Exclude - Not a Primary Study

Kester L, Stoller J. Prevalence and Causes of Medication Errors: A Review. *Clinical Pulmonary Medicine* 2003;10(6):322-6. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Keyhani S, Hebert PL, Ross JS, et al. Electronic health record components and the quality of care. *Med Care* 2009;46(12):1267-72. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Keyser L, Pakyz A, Lee K. Evaluation of a standardized vancomycin nomogram in computerized physician order entry system. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Khajouei R, Jaspers MW. CPOE system design aspects and their qualitative effect on usability. *Studies in Health Technology & Informatics* 2008;136:309-14. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Khajouei R, Jaspers MW. The impact of CPOE medication systems' design aspects on usability, workflow and medication orders: A systematic review. *Methods Inf Med* 2010;49(1):3-19. PMID:19582333 OVID MEDLINE.

Exclude - Unable to Retrieve

Kheniene F, Bedouch P, Durand M, et al. Economic impact of an automated dispensing system in an intensive care unit. *Ann Fr Anesth Reanim* 2008;27(3):208-15. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Kho A, Henderson LE, Dressler DD, et al. Use of handheld computers in medical education. A systematic review. *J Gen Intern Med* 2006;21(5):531-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Khoury AT. Support of quality and business goals by an ambulatory automated medical record system in Kaiser Permanente of Ohio. *Eff Clin Pract* 1998;1(2):73-82. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Khovanov AV, Nechaev VI, Krylov VV, et al. An objective relation approach in the modeling of dispensary registration of patients with tuberculosis. *Problemy Tuberkuleza I Boleznij Legkih* 2007;tuberk.(4):10-3. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Khowaja K, Nizar R, Merchant R, et al. A systematic approach of tracking and reporting medication errors at a tertiary care university hospital, Karachi, Pakistan. *Therapeutics and Clinical Risk Management* 2008;4(4):673-9. Database: Embase Sept 22-09.

Exclude - Not MMIT

Kidd MR. The better medication management system. Implications for Australian general practice. *Aust Fam Physician* 2002;31(6):516-20. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kidder D, Bae J. Evaluation results from prospective drug utilization review: Medicaid demonstration. *Pharm Pract Manag Q* 1999;19(3):17-29. Database: IPA.

Exclude - Not MMIT

Kilbridge PM, Welebob EM, Classen DC. Development of the Leapfrog methodology for evaluating hospital implemented inpatient computerized physician order entry systems. *Quality & Safety in Health Care* 2006;15(2):81-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kile DM, Pasquarella MV, Durma MT, et al. Failure-mode and effects analysis - Morphine sulfate concentrate solution. *Ashp Midyear Clinical Meeting* 2003;38. Database: IPA.

Exclude - Not a Primary Study

Killelea K, Boecler L, Dumitru D, et al. Changes to physician orders made by pharmacists during verification in a computerized physician order entry (CPOE) system. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Kim C, Kim H, Nam J, et al. Internet diabetic patient management using a short messaging service automatically produced by a knowledge matrix system. Diabetes Care 2007;30(11):2857-8. Database: CINAHL.

Exclude - Not MMIT

Kim GR, Miller MR, Ardolino MA, et al. Capture and classification of problems during CPOE deployment in an academic pediatric center. AMIA 2007;414-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kim HC, Bae YH, Kim SW. Innovative ambulatory drug delivery system using an electrolytic hydrogel infusion pump. IEEE Trans Biomed Eng 1999;46(6):663-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kim HS, Kim NC, Ahn SH. Impact of a nurse short message service intervention for patients with diabetes. J Nurs Care Qual 2006;21(3):266-71. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kim HS, Cho H. Health level 7 development framework for medication administration. CIN COMPUT INFORM NURS 2009;27(5):307-17. PMID:19726925 OVID MEDLINE.

Exclude - No Outcomes of Interest

Kim J, Bates DW. Results of a survey on medical error reporting systems in Korean hospitals. Int J Med Inf 2006;75(2):148-55. 8878062

Database: Inspec.

Exclude - Not MMIT

Kim JT, Cho YH, Shin HT, et al. Study of the accuracy of prescription by using a computerized chemotherapy order system. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.

Exclude - Not a Primary Study

Kim JT, Kim SL, Baik JE, et al. TPN order system for the improvement of pharmacy affairs. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.

Exclude - Not a Primary Study

Kim J Y, Liao J, Lober W B. Applying social network analysis techniques to measure the efficacy of computerized medication records. In Bethesda, MD, USA: American Medical Informatics Assoc; 2002. p. 1063.8023622

Database: Inspec.

Exclude - Not a Primary Study

Kimmel K, Sensmeier J. A technological approach to enhancing patient safety. 2002.
<http://nyam.waldo.kohalibrary.com/cgi-bin/koha/opac-detail.pl?biblionumber=224691> saved
as NYAM Kimmel.pdf

Grey Lit.

Exclude - Not a Primary Study

Kimura M, Koyama T, Kaihara S, et al. Anticipator: A medical expert system implemented
by prolog/KR. Journal of Information Processing 1984;7(3):149-56. 2424505

Database: Inspec.

Exclude - Not a Primary Study

Kimura M, Shimizu K, Tsuchiya F, et al. Anticipator: an antibiotic medication consulting
system. Automedica 1986;7(1):33-41. 2760480

Database: Inspec.

Exclude - No Outcomes of Interest

Kincade K. Barcode scanners aren't just for groceries any more. Laser Focus World
2005;41(7):69-70. 8595530

Database: Inspec.

Exclude - Not MMIT

King A, Reed L. Reader's perspective. The safety and efficiency benefits of using CPOE
systems are overstated. Health Data Manag 2009;17(3):10 PMID:20349740 OVID
MEDLINE.

Exclude - Not a Primary Study

King MW, Pathan J, Kessler JM. Developmental steps in the creation of an online Adverse
Drug Event Program. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

King WJ, Le Saux N, Sampson M, et al. Effect of point of care information on inpatient
management of bronchiolitis. BMC Pediatrics 2007;7:4 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kini N, Savage B. CPOE primer. Physician Exec 2004;30(2):20-6. Database: Ovid
MEDLINE(R).

Exclude - Not MMIT

Kinnaird D, Cox T, Wilson JP. Unclaimed prescriptions in a clinic with computerized
prescriber order entry. Am J Health Syst Pharm 2003;60(14):1468-70. Database: Ovid
MEDLINE(R).

Exclude - Not MMIT

Kinninger T, Reeder L. The business case for medication safety. Healthc Financ Manage
2003;57(2):46-51. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kinsley M. Some pumps know best. Nursing Spectrum -- New York & New Jersey Edition
2009;18A(5):12-3. Database: CINAHL.

Exclude - Not a Primary Study

Kirby J, Barker B, Fernando DJ, et al. A prospective case control study of the benefits of electronic discharge summaries. *Journal of Telemedicine & Telecare* 2006;12:Suppl-1 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kirchheimer B. Electronic records on board: PeriBirth system reduces errors, increases safety on Lenox Hill's labor and delivery unit. *Nursing Spectrum -- New York & New Jersey Edition* 2009;21(21):18-9.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010451818&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1533&accno=2010451818
EBSCO CINAHL.
Exclude - Not a Primary Study

Kirkpatrick TK, Harrison D. An economical application of bar code technology to dispensing and departmental charges. *ASHP Annual Meeting* 1989;46: Database: IPA.
Exclude - Not a Primary Study

Kirschenbaum B. Will coding shortcuts of the past come back to haunt you? *Pharmacy Practice News* 2009;31(4):9 Database: IPA.
Exclude - Not a Primary Study

Kirschenbaum BE. Life lines: The impact of NOT getting started. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Kirschenbaum BE. Bar code technology: Progress in the last year. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Kishel JJ, Maguire M, Pankratz L, et al. Implementing an electronically based, nurse-driven pneumococcal vaccination protocol for inpatients. *Am J Health Syst Pharm* 2009;66(14):1304-8. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Klein EG, Santora JA, Pascale PM, et al. Automated dispensing: Comprehensive evaluation of accuracy, and time/cost impact with the Baxter ATC-212. *Ashp Midyear Clinical Meeting* 1993;28: Database: IPA.
Exclude - Not a Primary Study

Klein EG, Santora JA, Pascale PM, et al. Medication cart-filling time, accuracy, and cost with an automated dispensing system. *Am J Hosp Pharm* 1994;51(9):1193-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Klein M, Shneyder Y, Neiman S, et al. Multidisciplinary communication effectiveness flow sheet from emergency department area to medical/surgical floor. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Klein PR. Technology marathon: Lessons learned implementing a pharmacy information system, bar code automation and point of care bed side scanning and electronic medication documentation system within 120 days. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Kleinberg K. Leveraging the April 2006 FDA bar code regulation. Healthcare Information and Management Systems Society; 2006.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=66892> Grey Lit.

Exclude - Not a Primary Study

Klimek JS. MTM services standards improve patient safety. Health Manag Technol 2009;30(7):20-1.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010348957&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=707&accno=2010348957 EBSCO CINAHL.

Exclude - Not a Primary Study

Klinar I, Marusic M., Ljubic M., et al. SMS drug - How to improve patient compliance? Farmaceutski Glasnik 2006;62(1):10-5. Database: Embase Sept 22-09.

Exclude - Not MMIT

Kloss LL. Managing the unintended consequences of EHRs. Journal of the American Health Information Management Association 2009;80(8):19 OVID EMBASE.

Exclude - Not a Primary Study

Knab JH, Wallace MS, Wagner RL, et al. The use of a computer-based decision support system facilitates primary care physicians' management of chronic pain. Anesthesia & Analgesia 2001;93(3):712-20. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Knapp KK, Blalock SJ, Black BL. ASHP survey of ambulatory care responsibilities of pharmacists in managed care and integrated health systems--2001. Am J Health Syst Pharm 2001;58(22):2151-66. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Knaup P, Wiedemann T, Bachert A, et al. Efficiency and safety of chemotherapy plans for children: CATIPO--a nationwide approach. Artif Intell Med 2002;24(3):229-42. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Knaup P, Pilz J, Kaltschmidt J, et al. Standardized documentation of drug recommendations in discharge letters--a contribution to quality management in cooperative care. Methods Inf Med 2006;45(4):336-42. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Knight JR. Electronic prescribing - Features and functions for safe medication management. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.

Exclude - Not a Primary Study

Knudsen AK. From individual patient care to population-based error dynamics: New opportunities for hospital pharmacists in the management of automated health care information. *Ashp Midyear Clinical Meeting 2002*;37: Database: IPA.
Exclude - Not a Primary Study

Ko B, Pierce B, McLain MD, et al. Improving nursing home dental pre-op medication administration with a system analysis monitoring project by pharmacy, dental, and nursing. *Ashp Summer Meeting 2004*;61: Database: IPA.
Exclude - Not a Primary Study

Ko BJ. Project management tools for the 21st century: Process in the successful integration of a computer technology program Bar Code Medication Administration. *ASHP Annual Meeting 2001*;58: Database: IPA.
Exclude - Not a Primary Study

Kocakulah M.C, Upson J. Cost analysis of computerized physician order entry using value stream analysis: A case study. *Research in Healthcare Financial Management 2005*;10(1):13-25. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Kocur,J. Computer consultant for optimizing geriatric patient medication plan for multiple conditions: A medical expert system East Texas State UniversityEditor. 1995. Grey Lit.
Exclude - Theses

Koehler AM, Maibach H.I. Electronic monitoring in medication adherence measurement implications for dermatology. *American Journal of Clinical Dermatology 2001*;2(1):7-12. Database: Embase Sept 22-09.
Exclude - Not MMIT

Koepp TE, Evans SD, Nafts M, et al. Pharmacy participation in a point of care system - Lessons learned. *ASHP Annual Meeting 1996*;53: Database: IPA.
Exclude - Not a Primary Study

Koerner SD, Becker F. Advances in Navy pharmacy information technology: accessing Micromedex via the Composite Healthcare Computer System and local area networks. *Mil Med 1999*;164(7):481-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Kofoed K, Zalounina A, Andersen O, et al. Performance of the TREAT decision support system in an environment with a low prevalence of resistant pathogens. *J Antimicrob Chemother 2009*;63(2):400-4. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kohler F, Monchovet S, Patris A, et al. [Applications of a simple system of aid to medical decision-making with various patients in internal medicine: SELF]. [French]. *Rev Med Interne 1988*;9(2):196-207. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Kolb KW, Israel MK. Use of computerized databases for drug therapy quality management. *Top Hosp Pharm Manage 1991*;11(2):44-50. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Koopmanschap MA, Rutten FF. The drug budget silo mentality: the Dutch case. *Value in Health* 2003;6:Suppl-51 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kopec D, Kabir M, Reinharth D, et al. Human Errors in Medical Practice: Systematic Classification and Reduction with Automated Information Systems. *J Med Syst* 2003;27(4):297-313. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Kopec D, Shagas G, Kabir M, et al. Errors in medical practice: identification, classification and steps towards reduction. *Studies in Health Technology & Informatics* 2004;103:126-34. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Koplan K, Regan S, Goldszer R. A computerized aid to support smoking cessation treatment for hospital patients. *J Gen Intern Med* 2008;23(8): Database: PsycINFO.

Exclude - Not MMIT

Kopp BJ, Erstad BL. How to better detect and prevent adverse events and medication errors in your ICU. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Koppel R. Computerized physician order entry systems: the right prescription? *LDI Issue Brief* 2005;10(5):1-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Koppel R. What do we know about medication errors made via a CPOE system versus those made via handwritten orders? *Critical Care* 2005;9(5):427-8. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Koppel R. Defending computerized physician order entry from its supporters. *Am J Manag Care* 2006;12(7):369-70. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Koppel R, Leonard CE, Localio AR, et al. Identifying and quantifying medication errors: evaluation of rapidly discontinued medication orders submitted to a computerized physician order entry system. *J Am Med Inform Assoc* 2008;15(4):461-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Korn D. Bar codes in production control. *Manufacturing Chemist* 1984;55(Sep):38-9. Database: IPA.

Exclude - Not a Primary Study

Korpman RA. Medication processing: closing the provider-pharmacy gap.[see comment]. *Manag Care Interface* 1998;11(6):64-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kosty T, Dietz D. E-prescriptions: ready for prime time? Sorting it all out. *Computertalk for the Pharmacist* 2000;20(Sep-Oct):23-5. Database: IPA.

Exclude - Not a Primary Study

Kowiatek JG, Weber RJ, Skledar SJ, et al. Assessing and monitoring override medications in automated dispensing devices. *Jt Comm J Qual Patient Saf* 2006;32(6):309-17. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kozakiewicz JM. Utilization of a formal change process implementation of a CPOE system. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Kozakiewicz JM. Implementation of an electronic medication administration record. A critical step towards bedside verification. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Kozakiewicz JM, Benis LJ, Fisher SM, et al. Safe chemotherapy administration: using failure mode and effects analysis in computerized prescriber order entry. *Am J Health Syst Pharm* 2005;62(17):1813-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kozakiewicz JM, Wall K. Process improvement approach to tracking and trending computerized order entry medication variances using a formal error classification system. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Kozakiewicz JM, Seo TH, Early CK. Process improvement using clinical decision support for high-risk medications based on medication variance reporting. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Kozousek V. Glitazone use associated with diabetic macular EDEMA: Comment. *Evidence-Based Ophthalmology* 2010;11(1):44-5. OVID EMBASE.

Exclude - Not a Primary Study

Kozuki Y, Schepp KG. Visual-feedback therapy for antipsychotic medication adherence. *Int Clin Psychopharmacol* 2006;21(1):57-61. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Krahenbuhl-Melcher A, Krahenbuhl St. Hospital drug safety: Medication errors and adverse drug reactions. *Schweizerische Rundschau fur Medizin - Praxis* 2005;94(24-25):1031-8.

Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Kramer JS. Collaborative pharmacist and nurse before/after study to evaluate patient safety using electronically standardized admission and discharge medication reconciliation in a tertiary care hospital. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Kravet SJ, Knight A.M., Wright S.M. Ten lessons from implementing a computerized provider order entry system. *JCOM* 2007;14(2):105-9. Database: Embase Sept 22-09.

Exclude - Not MMIT

Kremen J, Blaha J, Kopecky P, et al. The treatment of hyperglycaemia in critically ill patients: comparison of standard protocol and computer algorithm. *Vnitr Lek* 2007;53(12):1269-73. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Krempa C, Olatubosun M, Zoppa LR. Streamlining the workflow process of a hospital pharmacy department implementation of a computerized physician order entry system. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Kremsdorf RD. CPOE in a community hospital setting - A physician's perspective. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Krishna S, Balas E.A., Boren S.A., et al. Patient acceptance of educational voice messages: A review of controlled clinical studies. *Methods Inf Med* 2002;41(5):360-9. Database: Embase Sept 22-09.
Exclude - Not MMIT

Krishnan S M, Vanicatte M, Lee I and others. Incorporating Interoperability Functionality for Increasing Patient Safety with PCA-Included Multiple Infusion Therapy. In Berlin, Germany: Springer Verlag; 2009. p.415-8. Engineering Village Compendex and Inspec.
Exclude - No Outcomes of Interest

Krizner K. Technology is the right prescription for minimizing medical errors. *Managed Healthcare Executive* 2002;12(9):44-5. Database: IPA.
Exclude - Not a Primary Study

Kroese WL, Avery AJ, Savelyich BS, et al. Assessing the accuracy of a computerized decision support system for digoxin dosing in primary care: an observational study. *J Clin Pharm Ther* 2005;30(3):279-83. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kromis LA, Durkin L, Ritter M. Adherence to continuous quality improvement principles lead to reduced harmful medication errors. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Kromis LA, Durkin L, Ritter M. Bedside bar code charting, smart pump dose mode programming and computer assisted physician order entry system utilization monitoring and quality improvement lead to reduced harmful medication errors. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Kubie A, James AH, Timms J, et al. Experience with a computer-assisted anticoagulant clinic. *Clinical & Laboratory Haematology* 1989;11(4):385-91. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Kubin CJ. Antimicrobial stewardship programs: Role in optimizing infectious disease outcomes. *Disease Management and Health Outcomes* 2008;16(6):403-10. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Kuehl AK, Chrischilles EA, Sorofman BA. System for exchanging information among pharmacists in different practice environments. *Am J Health Syst Pharm* 1998;55(10):1017-24. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kuiper SA, McCreadie SR, Mitchell JF, et al. Medication errors in inpatient pharmacy operations and technologies for improvement. *Am J Health Syst Pharm* 2007;64(9):955-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kun L G. The use of a personal computer for patient-condition-treatment in a CCU/ICU environment. In New York, NY, USA: IEEE; 1983. p. 715.2320912

Database: Inspec.

Exclude - No Outcomes of Interest

Kunac DL, Harrison-Woolrych M, Tatley M. Pharmacovigilance in New Zealand: The role of the New Zealand Pharmacovigilance Centre in facilitating safer medicines use. *N Z Med J* 2008;121(1283):76-89. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Kunhardt H. Evidence-based information technology: Concept for rational information processing in the health care system. 2006. Database: PsycINFO.

Exclude - Not MMIT

Kuo C, Li Y, Lee P and others. An interoperability infrastructure with portable prescription for improving patient safety - the framework of a national standard in Taiwan. In Piscataway, NJ, USA: IEEE; 2009. p.293-7.10799512

Database: Inspec.

Exclude - Not a Primary Study

Kuo GM, Phillips RL, Graham D, et al. Medication errors reported by US family physicians and their office staff. *Quality & Safety in Health Care* 2008;17(4):286-90. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kuper JJ. Accuracy of computerized monitoring of vancomycin restrictions. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Kuperman GJ, Bates DW, Teich JM, et al. A new knowledge structure for drug-drug interactions. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1994;836-40. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kuperman GJ, Cooley T, Tremblay J, et al. Decision support for medication use in an inpatient physician order entry application and a pharmacy application. *Studies in Health Technology & Informatics* 1998;52 Pt 1:467-71. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kuperman GJ, Teich JM, Tanasijevic MJ, et al. Improving response to critical laboratory results with automation: results of a randomized controlled trial. *J Am Med Inform Assoc* 1999;6(6):512-22. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kuperman GJ, Teich JM, Gandhi TK, et al. Patient safety and computerized medication ordering at Brigham and Women's Hospital. *Jt Comm J Qual Improv* 2001;27(10):509-21. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kuperman GJ, Gibson RF. Computer physician order entry: benefits, costs, and issues. *Ann Intern Med* 2003;139(1):31-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kuperman GJ, Marston E, Paterno M, et al. Creating an enterprise-wide allergy repository at Partners HealthCare System. *AMIA* 2003;376-80. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kuperman GJ, Gandhi TK, Bates DW. Effective drug-allergy checking: methodological and operational issues. *Journal of Biomedical Informatics* 2003;36(1-2):70-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kuperman GJ, Bobb A, Payne TH, et al. Medication-related clinical decision support in computerized provider order entry systems: a review. *J Am Med Inform Assoc* 2007;14(1):29-40. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kuritani Y, Shinke K, Kuramitsu A, et al. Development of dispensing support system to improve safety management - Electronic prescription saving system using PDA. *Japanese Journal of Pharmaceutical Health Care & Sciences* 2004;30(10):627-37. Database: IPA.

Exclude - Not a Primary Study

Kurtz MM, Baker E, Pearlson GD, et al. A virtual reality apartment as a measure of medication management skills in patients with schizophrenia: a pilot study. *Schizophr Bull* 2007;33(5):1162-70. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Kushniruk A, Borycki E, Kuwata S, et al. Predicting changes in workflow resulting from healthcare information systems: ensuring the safety of healthcare. *HEALTHC Q* 2006;9:Spec-8 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kushniruk AW, Borycki EM, Kuwata S, et al. Using a low-cost simulation approach for assessing the impact of a medication administration system on workflow. *Studies in Health Technology & Informatics* 2008;136:567-72. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kushniruk AW, Borycki EM, Anderson JG, et al. Preventing technology-induced errors in healthcare: the role of simulation. *Studies in Health Technology & Informatics* 2009;143:273-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kuwata S, Kushniruk A, Borycki E, et al. Using simulation methods to analyze and predict changes in workflow and potential problems in the use of a bar-coding medication order entry system. *AMIA 2006*;994 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Kwan JW. Update on utilization of infusion device technology. *Ashp Midyear Clinical Meeting 1993*;28: Database: IPA.

Exclude - Not a Primary Study

Kweekel DM, Van den Broek P.J., Teepe-Twiss I.M., et al. Project for surveillance of antimicrobial therapy advances rational prescriptions. A new system for effective use of antibiotics. *Pharm Weekbl 2003*;138(43):1498-504. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Kwok R, Dinh M, Dinh D, et al. Improving adherence to asthma clinical guidelines and discharge documentation from emergency departments: Implementation of a dynamic and integrated electronic decision support system. *EMA - Emergency Medicine Australasia* 2009;21(1):31-7. Database: Embase Sept 22-09.

Exclude - Not MMIT

Kwon H, Cho J, Kim H, et al. Development of web-based diabetic patient management system using short message service (SMS). *Diabetes Res Clin Pract* 2003;66(Suppl.):S133-S137 Database: Embase Sept 22-09.

Exclude - Not MMIT

Kynard LC, Meyser SR. Transition from the traditional pharmacy model toward pharmaceutical care using automation. *Ashp Midyear Clinical Meeting 2003*;38: Database: IPA.

Exclude - Not a Primary Study

Lacarelle B, Pisano P, Gauthier T, et al. Abbott PKS system: a new version for applied pharmacokinetics including Bayesian estimation. *Int J Biomed Comput* 1994;36(1-2):127-30. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lacasa C, Roure C, Martinez J, et al. Quality control of the drug administration of a general hospital. *Ashp Midyear Clinical Meeting 1995*;30: Database: IPA.

Exclude - Not a Primary Study

Lacasa C, Cot R, Roure C, et al. Medication errors in a general hospital. *European Hospital Pharmacy* 1998;4(2):35-40. Database: IPA.

Exclude - Unable to Retrieve Foreign

LaCourse D. Breakthrough design of hospital bar-coded medication process. In Atlanta, GA, United states: Institute of Industrial Engineers; 2005. 20064010153875 Database: Compendex.

Exclude - Unable to Retrieve

Ladd H, McCullough L, Revere D. Decision support for evidence-based public health practice and policy development in the global south. Invited Paper for the Rockefeller Foundation's Making the eHealth Connection: Global Partnerships, Local Solutions Meeting. Jul 2008. Bellagio, Italy., The Rockefeller Foundation; 2008 Jan 7. http://www.ehealth-connection.org/files/conf-materials/Decision%20Support%20for%20Public%20Health_0.pdf
Grey Lit.

Exclude - No Outcomes of Interest

Ladyzdynski P, Wojcicki J, Krzymien J, et al. Mobile telecare system for intensive insulin treatment and patient education. First applications for newly diagnosed type 1 diabetic patients. *Int J Artif Organs* 2006;29(11):1074-81. Database: Embase Sept 22-09.

Exclude - Not MMIT

Lagor,C. The role of information technology in a study on antithrombotic-related bleeding events The University of UtahEditor. 2005. Grey Lit.

Exclude - No Outcomes of Interest

Lahdenpera TS, Kyngas HA. Patients' views about information technology in the treatment of hypertension. *J Telemed Telecare* 2000;6(2):108-13. 6547879

Database: Inspec.

Exclude - Not MMIT

Lai C L, Chien S W, Chen S C and others. The application of RFID on inpatient medication administration to improve patient safety and reduce adverse drug events. In 2008; Rhodes, Greece): 2008. p.437-42.Grey Lit.

Exclude - No Outcomes of Interest

Lai C, Chien S, Chang L and others. Enhancing medication safety and healthcare for inpatients using RFID. In Piscataway, NJ, USA: Management of Engineering and Technology,Portland International Center; 2007. p.2783-90.9870749

Database: Inspec.

Exclude - Not a Primary Study

Lai JS, Yokoyama G. Impact of computerized prescriber order entry (CPOE) on clinical pharmacists: Challenges and trends in clinical pharmacy. *Ashp Summer Meeting* 2006;63:

Database: IPA.

Exclude - Not a Primary Study

Lai JT, Hou TW, Yeh CL, et al. Using Healthcare IC Cards to manage the drug doses of chronic disease patients. *Computers in Biology & Medicine* 2007;37(2):206-13. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Laicher A, Vesper A, Escher D, et al. Implementation of computerized information systems in the pharmaceutical technology department. *Pharmazeutische Industrie* 1994;56(4):385-9. Database: IPA.

Exclude - Not a Primary Study

Lake M. Finding relief in electronic prescribing. *Computertalk for the Pharmacist* 2004;24(6):13-4. Database: IPA.

Exclude - Not a Primary Study

Lallement A. Labeling and pharmaceutical safety. S T P Pharma Pratiques 1985;879-81.
Database: IPA.
Exclude - Not MMIT

Lam ZH, Hwang KS, Lee JY, et al. Active insulin infusion using optimal and derivative-weighted control. Medical Engineering & Physics 2002;24(10):663-72. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Lamb E. E-prescribing: Recommendations for Medicare draft standards issued. Pharmacy Today (Washington D C) 2004;10(11):24 Database: IPA.
Exclude - Not a Primary Study

Lamma E, Mello P, Nanetti A, et al. Artificial intelligence techniques for monitoring dangerous infections. IEEE Transactions on Information Technology in Biomedicine 2006;10(1):143-55. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Lamy JB, Duclos C, Bar-Hen A, et al. An iconic language for the graphical representation of medical concepts. BMC Med Inform Decis Mak 2008;8:16 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Lamy JB, Venot A, Bar-Hen A, et al. Design of a graphical and interactive interface for facilitating access to drug contraindications, cautions for use, interactions and adverse effects. BMC Med Inform Decis Mak 2008;8:21 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Lamy O, Burnand B. Clinical practice guidelines: Do they help improving medical practice? Med Hyg (Geneve) 2004;62(2506):2363-6. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Landis NT. Automated dispensing systems vary by function and location. Am J Hosp Pharm 1993;50(Nov):2242-8. Database: IPA.
Exclude - Not a Primary Study

Landon D and Mitchell G. Combat Emergency Medicine Expert System (CEMES): Project phase 2 report. 1998. Grey Lit.
Exclude - No Outcomes of Interest

Lang JR. Inexpensive microcomputer for intravenous admixture service applications. Am J Hosp Pharm 1983;40(3):406-8. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Langley J and Beasley C. Health information technology for improving quality of care in primary care settings. Institute for Healthcare Improvement; 2007.
http://library.ahima.org/xpedio/groups/public/documents/government/bok1_037314.pdf Grey Lit.
Exclude - Not a Primary Study

Lanoue E, Still C. Patient identification: Producing a better barcoded wristband. Patient Safety & Quality Healthcare 2008;12-6. <http://www.psqh.com/mayjun08/identification.html> Grey Lit.
Exclude - Not MMIT

Lanza V. Automatic record keeping in anaesthesia--a nine-year Italian experience. *International Journal of Clinical Monitoring & Computing* 1996;13(1):35-43. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lapane K, Cameron K, Feinberg J. Technology for improving medication monitoring in nursing homes. *AHRQ* 2005;4(2):401-13. Grey Lit.

Exclude - Not a Primary Study

Larizza M. A simple, live, cost-effective electronic tracking system for aseptic preparations: Improving communication and reducing disruptions. *Journal of Oncology Pharmacy Practice* 1991;14(2):91 Database: IPA.

Exclude - Not MMIT

Larsen MD, Nielsen LP, Jeffery L, et al. Medication errors on hospital admission. *Ugeskr Laeger* 2006;168(35):2887-90. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Larsen RA, Evans RS, Burke JP, et al. Improved perioperative antibiotic use and reduced surgical wound infections through use of computer decision analysis. *Infect Control Hosp Epidemiol* 1989;10(7):316-20. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Larson JR, Hisel TM, Joss JD, et al. Comparison of a computerized medical record to actual medication use. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Larson RL, Blake JP. Achieving order entry by physicians in a computerized medical record. *Hosp Pharm* 1988;23(Jun):551-3. Database: IPA.

Exclude - Not a Primary Study

Latham BD, Lehman RK. Medication reconciliation: From admission to discharge using electronically generated medication forms from a clinical information system. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Lau F. A clinical decision support system prototype for cardiovascular intensive care. *International Journal of Clinical Monitoring & Computing* 1994;11(3):157-69. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Laurie-Shaw B, Taylor W, Roach C. Focus on clinical best practices, patient safety and operational efficiency. *HEALTHC Q* 2004;10:Spec-6 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lavin M, Sierzega G, Pucklavage D, et al. Carts and care. Roll out safer medication delivery and smoother workflow with mobile technology. *Nurs Manag (Harrow)* 2009;41(7):16-8. Database: CINAHL.

Exclude - Not a Primary Study

Lavoie G, Tremblay L, Durant P, et al. Medicarte software developed for the Quebec microprocessor health card project. Medinfo 1995;8(Pt 2):1662 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lawrence D. Money matters. Part II: Hunkering down. Healthc Inform 2009;26(10):44-6. PMID:19883002 OVID MEDLINE.

Exclude - Not a Primary Study

Lawrence D. The field narrows. As CPOE takes center stage, many hospitals are finding that a pharmacy system from their core clinical vendor makes the most sense. Healthc Inform 2009;26(8):14-6. PMID:19722322 OVID MEDLINE.

Exclude - Not a Primary Study

Lawrence D. Steps forward on e-prescribing: as e-prescribing becomes more widespread, even hospital organizations without full EMR implementation are seeing gains in clinician workflow and patient safety. Healthc Inform 2010;27(5):24-6.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010655411&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1239&accno=2010655411
EBSCO CINAHL.

Exclude - Not a Primary Study

Lawrence KL, Kollef M. Antimicrobial stewardship in the Intensive care unit advances and obstacles. Am J Respir Crit Care Med 2009;179(6):434-8. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Layton D, Heeley E, Shakir S. Identification and evaluation of a possible signal of exacerbation of colitis during rofecoxib treatment, using Prescription-Event Monitoring Data. J Clin Pharm Ther 2004;29(2):171-81. Database: CINAHL.

Exclude - Not MMIT

Le Bouar Lacroux V., Lhopiteau K., Toledano N., et al. The infusion as an indicator of quality on the drug circuit. Journal de Pharmacie Clinique 2002;21(4):247-54. Database: Embase Sept 22-09.

Exclude - Not MMIT

Le Cosquer P. Help software trials: Conception and evaluation for the dispensation and management of drugs in clinical. Journal de Pharmacie Clinique 1997;15(SPEC. ISSUE):72-3. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Le Gonidec P, Diallo M, Djoussa-Kambou S, et al. Performances of an automated dispensing system combined with a computerized prescription order entry. Ann Pharm Fr 2008;67(2):84-90. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Le Jamtel H. Application of the CIP/39 code to cases. S T P Pharma Pratiques 1985;891-4. Database: IPA.

Exclude - Not MMIT

Le Normand Y, Milpied N, Kergueris MF, et al. Pharmacokinetic parameters of vancomycin for therapeutic regimens in neutropenic adult patients. *Int J Biomed Comput* 1994;36(1-2):121-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Le TT, Mark SM, Gumpper KF. Evaluation of a computerized provider order entry system in a pediatric hospital. *Ashp Summer Meeting* 2002;59: Database: IPA.

Exclude - Not a Primary Study

Leavitt M. Pilot testing of initial electronic prescribing standards: Cooperative agreements. 2003. <http://www.cms.hhs.gov/EPrescribing/Downloads/E-RxReporttoCongress.pdf> Grey Lit.

Exclude - Not a Primary Study

Leavitt M. Improving the medicare quality improvement organization program: Response to the institute of medicine study. 2006.

<http://www.himss.org/content/files/CMSQIOModernizationPlan8-31-06.pdf> Grey Lit.

Exclude - Not a Primary Study

Lebel D, Bussieres JF. Interface between a pharmacy system and a computerized physician order entry system (CPOES): A pilot project. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Lederer J, Best D. Reduction in anticoagulation-related adverse drug events using a trigger-based methodology. *Jt Comm J Qual Patient Saf* 2005;31(6):313-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lederman JR, Byers JA, Sincak CA. Implementation of an intravenous to oral medication therapy conversion program in a veterans affairs hospital utilizing an electronic chart. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Lederman R, Parkes C. Exploring errors in a medication process: An analysis of information delivery. In Maribor, Slovenia: Institute of Electrical and Electronics Engineers Computer Society; 2002. p.79-84.2002307033722

Database: Compendex.

Exclude - Not MMIT

Ledford A, Mackey BM, McCumber A, et al. Creation of an outpatient chemotherapy treatment record shared by pharmacy and nursing a chemotherapy infusion center to reduce the potential for medication errors and to improve efficiency. *Journal of Oncology Pharmacy Practice* 1996;14(2):96 Database: IPA.

Exclude - No Outcomes of Interest

Lee ET, Lee ME. Cybernetics and expert systems for machines: Interactions of medical expert systems. *Kybernetes* 1999;28(9):1081-3. 6447140

Database: Inspec.

Exclude - Not a Primary Study

Lee HR, Yoo SK, Jung SM, et al. A Web-based mobile asthma management system. *J Telemed Telecare* 2005;11(Suppl):56-9. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Lee J, Fulton G, Farr S and others. Online discharge notes increase utilization of computerized medication lists. In Bethesda, MD, USA: American Medical Informatics Assoc; 2002. p. 1077.8030448
Database: Inspec.
Exclude - Not MMIT

Lee RL. Use of portable data entry and bar code technology in hospital drug ordering. *Australian Journal of Hospital Pharmacy* 1990;20(Apr):186-8. Database: IPA.
Exclude - Unable to Retrieve

Lee WM, Retterer JA. Computerized physician order entry (CPOE) and the opportunity to improve pharmaceutical care. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Lee Y, Carroll C, Medhi D. A CORBA-based medical decision support system for medication error. In Bethesda, MD, USA: American Medical Informatics Assoc; 2002. p. 1078.8030449
Database: Inspec.
Exclude - Not a Primary Study

Leeman-Castillo B. Acceptability of a bilingual interactive computerized educational module in a poor, medically underserved patient population. *Journal of Health Communication* 2007;12(1):77-94. Database: PsycINFO.
Exclude - Not MMIT

Lefkowitz S, Cheiken H, Barnhart M, et al. Bare codes, applications in pharmacy practice: A 2-year retrospective report. *Ashp Midyear Clinical Meeting* 1989;24: Database: IPA.
Exclude - Not a Primary Study

Lefkowitz S, Cheiken H, Barnhart MR. Bar codes at the patients bedside: Experience with a point of care system. *ASHP Annual Meeting* 1990;47: Database: IPA.
Exclude - Not a Primary Study

Lefkowitz S, Cheiken H, Barnhart MR. A trial of the use of bar code technology to restructure a drug distribution and administration system. *Hosp Pharm* 1991;26(3):239-42. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Lehmann CU, Kim GR. Computerized provider order entry and patient safety. *Pediatr Clin North Am* 2006;53(6):1169-84. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Lehmann CU, Kim GR, Gujral R, et al. Decreasing errors in pediatric continuous intravenous infusions. *Pediatric Critical Care Medicine* 2006;7(3):225-30. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Lehmann ED, Deutsch T. Insulin dosage adjustment in diabetes. *J Biomed Eng* 1992;14(3):243-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Lehmann ED, Deutsch T. AIDA2: a Mk. II automated insulin dosage advisor. *J Biomed Eng* 1993;15(3):201-11. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lehmann ED, Deutsch T. Application of computers in diabetes care--a review. II. Computers for decision support and education. *Med Inform (Lond)* 1995;20(4):303-29. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lehmann ED, Deutsch T. Application of computers in diabetes care--a review. I. Computers for data collection and interpretation. *Med Inform (Lond)* 1995;20(4):281-302. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Leibovici L, Gitelman V, Yehezkelli Y, et al. Improving empirical antibiotic treatment: prospective, nonintervention testing of a decision support system. *J Intern Med* 1997;242(5):395-400. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Leirer VO, Morrow DG, Pariante GM, et al. Elders' nonadherence, its assessment, and computer assisted instruction for medication recall training. *J Am Geriatr Soc* 1988;36(10):877-84. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Leirer VO, Morrow DG, Tanke ED, et al. Elders' nonadherence: its assessment and medication reminding by voice mail. *Gerontologist* 1991;31(4):514-20. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lemke JM, Gardner DK, Pedersen CA, et al. Computerized prescriber order entry improves prescribing for patients in a neonatal intensive care unit. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Lenclen R. Computerized physician order entry softwares for pediatric units. *Arch Pediatr* 2005;12(6):918-20. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lenderink BW, Egberts TC. Closing the loop of the medication use process using electronic medication administration registration. *Pharmacy World & Science* 2004;26(4):185-90. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lenderink BW. Safe "closed loop" medication management: the need for barcoded unit doses. *European Journal of Hospital Pharmacy* 2004;10(6):140-3. Database: IPA.

Exclude - Unable to Retrieve Foreign

Lenert L, Sheiner L, Blaschke T. Improving drug dosing in hospitalized patients: automated modeling of pharmacokinetics for individualization of drug dosage regimens. *Comput Methods Programs Biomed* 1989;30(2-3):169-76. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lenert LA, Lurie J, Sheiner LB, et al. Advanced computer programs for drug dosing that combine pharmacokinetic and symbolic modeling of patients. *Comput Biomed Res* 1992;25(1):29-42. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lenert LA, Klostermann H, Coleman RW, et al. Practical computer-assisted dosing for aminoglycoside antibiotics. *Antimicrob Agents Chemother* 1992;36(6):1230-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lenert LA, Cher DJ. Use of meta-analytic results to facilitate shared decision making. *J Am Med Inform Assoc* 1999;6(5):412-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lenz T. Fewer stacks, fewer gaps: There's a lot to be gained - and saved - by streamlining paper-based workflows and processes. *Health Manag Technol* 2009;30(1):20-1. 10740048 Database: Inspec.

Exclude - Not a Primary Study

Leonard MS, Cimino M, Shaha S, et al. Risk reduction for adverse drug events through sequential implementation of patient safety initiatives in a children's hospital. *Pediatrics* 2006;118(4):e1124-e1129 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Leong TY, Kaiser K, Miksch S. Free and open source enabling technologies for patient-centric, guideline-based clinical decision support: a survey. *Yearbook of Medical Informatics* 2007;19:74-86. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Leslie T, Sagara L, Wallis K. Getting connected: The outlook for electronic prescribing in California. *Manatt Health Solutions*; 2008.

<http://www.chcf.org/topics/view.cfm?itemid=133793> Grey Lit.

Exclude - Not a Primary Study

Leu MG, Cheung M, Webster TR, et al. Centers speak up: the clinical context for health information technology in the ambulatory care setting. *J Gen Intern Med* 2008;23(4):372-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Levesque SM, O'Brien BS, Berard CM, et al. Improving pain management through a multidisciplinary team and use of information technology. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Levick D, O'Brien D. CPOE is much more than computers. *Physician Exec* 2003;29(6):48-52. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Levin RI, Koenig K, Corder M, et al. Risk stratification and prevention in chronic coronary artery disease: Use of a novel prognostic and computer-based clinical decision support system in a large primary managed-care group practice. *Disease Management* 2002;5(4):197-213. Database: Embase Sept 22-09.

Exclude - Not MMIT

Levison DL, Witkowski PL. Replacing cart exchange with automation frees pharmacists for expanded roles. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Levit F, Garside DB. Computer-assisted prescription writing. *Comput Biomed Res* 1977;10(5):501-10. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lew GB, Stephens R, Serrano VA. Medication error types with automated dispensing devices. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Lew M. The "Big bang": Lessons learned from implementing a bidirectional interface. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Lew P. Medication errors and quality improvement: University teaching hospital's experience. *California Journal of Hospital Pharmacy* 1993;5(May):9-11. Database: IPA.

Exclude - Not a Primary Study

Lewis C. Cost-effective, quality solutions to boost healthcare and bottom lines. *Caribbean Business* 2010;38(20):B8-B9

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=51302831&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Lewis PJ, Dornan T, Taylor D, et al. Prevalence, incidence and nature of prescribing errors in hospital inpatients: a systematic review. *Drug Saf* 2009;32(5):379-89. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lewis RK, Witte KW, Fruin DM, et al. Comparative analysis of controlled substance distribution systems. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Lewis RK, Carter B, Glover D, et al. Comprehensive services in an ambulatory care pharmacy. *Am J Health Syst Pharm* 1995;52(16):1793-7. Database: Embase Sept 22-09.

Exclude - Not MMIT

Li C, Xia J, Deng J, et al. A web-based quantitative signal detection system on adverse drug reaction in China. *Eur J Clin Pharmacol* 2009;65(7):729-41. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Liang H, Xue Y, Berger B. Web-based intervention support system for health promotion. *Decision Support Systems* 2006;42(1):435-49. Database: PsycINFO.

Exclude - Not MMIT

Liaw ST, Ugoni AM, Cairns C. Computer education. Don't forget the older GPs. Aust Fam Physician 2000;29(8):802-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Liaw ST, Pearce CM, Chondros P, et al. Doctors' perceptions and attitudes to prescribing within the Authority Prescribing System. Med J Aust 2003;178(5):203-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Liaw ST, Schattner P. Electronic decision support in general practice. What's the hold up? Aust Fam Physician 2003;32(11):941-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Libby D, Grove C, Adams M. Collaborative use of informatics among hospitals to benchmark medication use processes. Jt Comm J Qual Improv 1997;23(12):636-52. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Liberto RB, Merryfield DW, Jennings TS, et al. An integrated medication safety program in an integrated healthcare system. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Lin AC, Huang YC, Ivey MF, et al. The efficiency analysis of an automated prescription filling system in an outpatient pharmacy. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Lin AC, Huang YC, Panches G, et al. Effect of a robotic prescription-filling system on pharmacy staff activities and prescription-filling time. Am J Health Syst Pharm 2007;64(17):1832-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lin CA, Neafsey PJ, Anderson E. Advanced practice registered nurse usability testing of a tailored computer-mediated health communication program. CIN - Computers Informatics Nursing 2010;28(1):32-41. OVID EMBASE.

Exclude - Not MMIT

Lin CP, Nichol WP, Hoey P, et al. Approach for analysis of order check overrides in a computerized practitioner order entry system. AMIA 2005;1033 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lin CP, Payne TH, Nichol WP, et al. Evaluating clinical decision support systems: monitoring CPOE order check override rates in the Department of Veterans Affairs' Computerized Patient Record System. J Am Med Inform Assoc 2008;15(5):620-6. PMID:18579840 OVID MEDLINE.

Exclude - No Outcomes of Interest

Lin MF, Moyle W, Chang HJ, et al. Effect of an Interactive Computerized Psycho-education System on patients suffering from depression. J Clin Nurs 2008;17(5):667-76. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lin ND, Martins SB, Chan AS, et al. Identifying barriers to hypertension guideline adherence using clinician feedback at the point of care. *AMIA* 2006;494-8. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Lindblad B, Chilcott J, Rolls L. Mary Lanning Memorial Hospital: communication is key. *Joint Commission Journal on Quality and Safety* 2004;30(10):551-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Linder JA, Bates DW, Williams DH, et al. Acute infections in primary care: accuracy of electronic diagnoses and electronic antibiotic prescribing. *J Am Med Inform Assoc* 2006;13(1):61-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Linder JA, Chan JC, Bates DW. Appropriateness of antiviral prescribing for influenza in primary care: a retrospective analysis. *J Clin Pharm Ther* 2006;31(3):245-52. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Linder JA, Ma J, Bates DW, et al. Electronic health record use and the quality of ambulatory care in the United States. *Arch Intern Med* 2007;167(13):1400-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Linder JA, Rigotti NA, Schneider LI, et al. An electronic health record-based intervention to improve tobacco treatment in primary care: a cluster-randomized controlled trial. *Arch Intern Med* 2009;169(8):781-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Linkens DA, Hacisalihzade SS. Computer control systems and pharmacological drug administration: a survey. [Review] [78 refs]. *J Med Eng Technol* 1990;14(2):41-54. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Linnarsson R. Decision support for drug prescription integrated with computer-based patient records in primary care. *Med Inform (Lond)* 1993;18(2):131-42. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Linnarsson R, Nordgren K. A shared computer-based problem-oriented patient record for the primary care team. *Medinfo* 1995;8 Pt 2:1663 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lipton HL, Miller RH, Wimbush JJ. Electronic prescribing: ready for prime time? *J Healthc Inf Manag* 2003;17(4):72-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

List BA, Ballard JL, Langworthy KS, et al. Electronic health records in an outpatient breastfeeding medicine clinic. *J Hum Lact* 2008;24(1):58-68. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Liu C, Endoh A, Minato K, et al. Quantitative assessment of the effect of prescription order entry system in university hospital by entropy. *Japan Journal of Medical Informatics* 1998;18(2):149-55. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Liu JH, Venot A. Design of a computerized system for the retrospective analysis by the physician of his own drug prescriptions. *Studies in Health Technology & Informatics* 2001;84(Pt:2):2-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Liu J W S, Shih C S, Yeh H C and others. Point-of-care support for error-free medication process. In Piscataway, NJ, USA: IEEE; 2008. p.29-40.9904840

Database: Inspec.

Exclude - Not a Primary Study

Liu MY, Caliendo GC, Friedman TS, et al. Implementation of red cell factor guidelines utilizing a physician order entry system. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Ljungberg C, Kettis LA, Morlin C, et al. Primary care and hospital doctors experiences of a shared electronic patient medical record system: Transfer of information about patients drug therapies. *International Journal of Pharmacy Practice* 2010;18(SUPPL. 1):4 OVID EMBASE.

Exclude - No Outcomes of Interest

Ikemer D and rtz K. HITCh at HIMSS 2008: The state role in health information technology. *National Conference of State Legislatures*; 2008.

http://www.ncsl.org/print/health/forum/HIMSS_08.pdf Grey Lit.

Exclude - Not a Primary Study

Llabre MM, Weaver KE, Duran RE, et al. A measurement model of medication adherence to highly active antiretroviral therapy and its relation to viral load in HIV-positive adults. *AIDS Patient Care & Stds* 2006;20(10):701-11. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Llopis SP, Sanchez AA, Quintana VB. Integral computerization of health care for inpatients. Impact on primary care activities. *Farmacia Hospitalaria* 2003;HOSP..(4):231-9. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Lloyd TR. Automated dispensing and data collection system. *Pharmaceutical Technology* 1977;1(Dec):21-4. Database: IPA.

Exclude - Not a Primary Study

Lobach DF, Underwood HR. Computer-based decision support systems for implementing clinical practice guidelines. *Drug Benefit Trends* 1998;10(Oct):48-53. Database: IPA.

Exclude - Not a Primary Study

Lobach DF, Kawamoto K, Anstrom KJ, et al. Development, deployment and usability of a point-of-care decision support system for chronic disease management using the recently-approved HL7 decision support service standard. *Stud Health Technol Inform* 2007;129(Pt:2):2-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Locatelli F, Covic A, Macdougall IC, et al. Effect of computer-assisted European Best Practice Guideline implementation on adherence and target attainment: ORAMA results. *Journal of Nephrology* 2009;22(5):662-74. OVID EMBASE.

Exclude - Not MMIT

Lockwood W. Integrating e-prescribing into the workflow. *Computertalk for the Pharmacist* 2004;24(3):34-5. Database: IPA.

Exclude - Not a Primary Study

Lockwood W. Integration through interfaces. *Computertalk for the Pharmacist* 2004;24(1):36-8. Database: IPA.

Exclude - Not a Primary Study

Lockwood WA. In the e-prescribing world, which comes first? *America's Pharmacist* 2004;127(10):20 Database: IPA.

Exclude - Not a Primary Study

Lodder H, Wolf L, Vernee J. Sharing medication data using the InterCare architecture. In 1916; Hannover, Germany: 2000. p.647-51. Grey Lit.

Exclude - No Outcomes of Interest

Logan H, Baker K, Cowen H. Initial results of the use of a drug-interaction system. *Spec Care Dentist* 1988;8(6):252-5. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Logan M. AAMI president: Biomed to play pivotal role in evolving world of healthcare. *Biomed Instrum Technol* 2010;44(2):95-April OVID EMBASE.

Exclude - Not a Primary Study

Logan SS, Smith KS, Hellmuth DJ. Prevention of allergic reactions to medications utilizing quality assurance to improve reporting. *ASHP Annual Meeting* 1992;49: Database: IPA.

Exclude - Not a Primary Study

Long AJ, Chang P, Li YC, et al. The use of a CPOE log for the analysis of physicians' behavior when responding to drug-duplication reminders. *Int J Med Inf* 2008;77(8):499-506. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Longe KM. Bar coding. The inventory manager. *J Healthc Mater Manage* 1987;5(4):37-40. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Longhurst R. YouTube: a new space for birth? *Feminist Review* 2009;(93):46-63. Scholar's Portal Sociological Abstracts.

Exclude - Not a Primary Study

Lopez J. Automation in pharmacy. *California Journal of Hospital Pharmacy* 1991;3(Jul):5-6.
Database: IPA.
Exclude - Not a Primary Study

Lormand J. Bringing patient safety technology to the bedside: By instituting bar code technology from the pharmacy to the patient's bedside, a Louisiana health care system reduces its medication error rate. *Health Manag Technol* 2005;26(8):24-6. 8708126
Database: Inspec.
Exclude - Not a Primary Study

Lou AS, Betorz Latorre JJ, Garcia AC, et al. [Quality control of a computer system for repeat prescriptions]. [Spanish]. *Aten Primaria* 1991;8(3):195-6. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Louie C, Proffitt LS, Dong D. Successfully managing H2 blocker use with an automated medication system. *Hosp Formul* 2004;32: Database: IPA.
Exclude - No Outcomes of Interest

Louis MJ. Utilizing bar codes in hospital pharmacy. *Pharm Times* 1991;57: Database: IPA.
Exclude - Not a Primary Study

Low J. Computerized prescribing. *Australian Journal of Pharmacy* 1999;78(Feb):168
Database: IPA.
Exclude - Not MMIT

Low J. Challenges in using information technology to improve hospital healthcare delivery. *Australian Journal of Pharmacy* 2008;87(1033):28 Database: IPA.
Exclude - Not MMIT

Lowe DO, Pitre MM, DaSilva-Barbosa AP, et al. Computerized physician order entry interfaced to a pharmacy system and electronic medication administration record - A pilot experience. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Lowe DO, Pitre MM, DaSilva-Barbosa AP, et al. Development of a computerized physician order entry system interfaced to a pharmacy system and electronic medication administration record. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Lu CY, Ross-Degnan D, Soumerai SB, et al. Interventions designed to improve the quality and efficiency of medication use in managed care: a critical review of the literature - 2001-2007. *BMC Health Serv Res* 2008;8:75 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Luber AD, Holtzer C, Dong D, et al. Evaluation of the number and types of overrides via an automated dispensing system. *ASHP Annual Meeting* 1996;53: Database: IPA.
Exclude - Not a Primary Study

Lubiniecki M, Sobolik R. Strategies for sound-alike, look-alike drugs. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Lucas AJ. Improving medication safety in a neonatal intensive care unit. *Am J Health Syst Pharm* 2004;61(1):33-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lucas KS. Technician order entry certification: Providing pharmaceutical care without delaying medication access. *ASHP Annual Meeting* 1997;54: Database: IPA.

Exclude - Not a Primary Study

Lucas PJ, de Bruijn NC, Schurink K, et al. A probabilistic and decision-theoretic approach to the management of infectious disease at the ICU. *Artif Intell Med* 2000;19(3):251-79.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Luchins DJ, Klass D., Hanrahan P., et al. Computerized analysis of therapeutic drug monitoring practices in a state hospital system. *P and T* 2001;26(9):478-85. Database: Embase Sept 22-09.

Exclude - Not MMIT

Ludwig BC. Applying bar coding to confirm the 5-r's. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Lui K, Bryson SM, Irwin DB, et al. Evaluation of bayesian forecasting for individualized gentamicin dosage in infants weighing 1000 g or less. *Am J Dis Child* 1991;145(4):463-7.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lum GK, Todd S, Frater B, et al. The contribution of decentralized clinical pharmacy service programs to positive financial and clinical pharmacy outcomes, and improved staffing effectiveness. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Lumpkin J. Letter to Secretary Thompson: E-prescribing standards. 2004.

<http://www.ncvhs.hhs.gov/040902lt2.htm> Grey Lit.

Exclude - Not MMIT

Luna D, Otero V, Canosa D, et al. Analysis and redesign of a knowledge database for a drug-drug interactions alert system. *Stud Health Technol Inform* 2007;129(Pt:2):2-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Exclude - Not a Primary Study

Lunardi Filho WD, Macada AC, Lunardi G. [Support system for patient care planning and prescription for nursing care]. [Portuguese]. *Rev Bras Enferm* 1995;48(1):66-77. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Exclude - Unable to Retrieve Foreign

Lundell J, Hayes TL, Vurgun S, et al. Continuous activity monitoring and intelligent contextual prompting to improve medication adherence. *Conference Proceedings:*

2007;2007:6287-90. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Lundy SH, Anderson CL, Valentine ET. Just what the doctor ordered? CMS and DEA introduce new measures to facilitate e-prescribing. *Journal of Health and Life Sciences Law* 2009;2(4):79-114. PMID:19673178 OVID MEDLINE.

Exclude - Not a Primary Study

Lunin LF. Writing and processing drug information in the 80's. *Drug Inf J* 1981;15(Jan-Mar):20-4. Database: IPA.

Exclude - Not MMIT

Lunner K. Medicare reform 101 - What just happened to Medicare? *Pharmacy Today (Washington D C)* 2004;10(2):28 Database: IPA.

Exclude - Not a Primary Study

Luo J. Portable computing in psychiatry. *The Canadian Journal of Psychiatry / La Revue canadienne de psychiatrie* 2004;49(1):24-30. Database: PsycINFO.

Exclude - Not a Primary Study

Luo J. Computer physician order entry: To implement or not? *Primary Psychiatry* 2006;13(3): Database: PsycINFO.

Exclude - Not MMIT

Luo J. Computerized Medicine. *Primary Psychiatry* 2006;13(9): Database: PsycINFO.

Exclude - Not MMIT

Luo J. Electronic health information exchange: Key trends to watch. *Primary Psychiatry* 2006;13(5): Database: PsycINFO.

Exclude - Not MMIT

Luo J. Electronic Prescribing Systems with Computer Decision Support. *Primary Psychiatry* 2006;13(6): Database: PsycINFO.

Exclude - Not MMIT

Luo J. Radio Frequency Identification: Ready for Implementation Decision. *Primary Psychiatry* 2006;13(7): Database: PsycINFO.

Exclude - Not MMIT

Lustig L. Computerize chemotherapy prescribing. *Hospital Pharmacy Practice* 1999;9(Dec):386-7. Database: IPA.

Exclude - Not MMIT

Lux A. Cost-benefit analysis of a new health insurance card and electronic prescription in Germany. *J Telemed Telecare* 2002;8:Suppl-5 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lykowski G, Mahoney D. Computerized provider order entry improves workflow and outcomes. *Nurs Manag (Harrow)* 2004;35(2):40G-H. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lyman JA, Conaway M, Lowenhar S. Formulary access using a PDA-based drug reference tool: does it affect prescribing behavior? *AMIA 2008;Annual:Symposium* PMID:18998942 OVID MEDLINE.

Exclude - Not MMIT

Lynch O, McGrory J, Coyle E. Design of mobile phone applications for point of care result validation. In Anaheim, CA, USA: ACTA Press; 2006. p.41-5.9272515

Database: Inspec.

Exclude - No Outcomes of Interest

Lynx D. CPOE: Friend or foe? Keeposted 2002;28(2):4-5. Database: IPA.

Exclude - Not MMIT

Lyons A, Richardson S. Clinical decision support in critical care nursing. AACN Clin Issues 2003;14(3):295-301. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Lyons SS, Tripp-Reimer T, Sorofman BA, et al. Information technology for clinical guideline implementation: Perceptions of multidisciplinary stakeholders. J Am Med Inform Assoc 2005;12(1):64-71. Database: IPA.

Exclude - Not MMIT

Mabeck H. Implementation of an electronic medication system and disregarded power of the record. Stud Health Technol Inform 2008;136:443-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Mac-Kay MV, Sanchez BJ, Martinez-Lanao J, et al. Drug dosage in end-stage renal disease (ESRD) patients undergoing haemodialysis. A predictive study based on a microcomputer program. Clin Pharmacokinet 1993;25(3):243-57. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

MacDonald TM, Parkinson J, Davey PG, et al. PACT data for dispensed drugs linked to NHS numbers are available now. Br Med J 1998;316:1530-1. Database: IPA.

Exclude - Not MMIT

MacDougall C, Polk RE. Antimicrobial stewardship programs in health care systems. Clin Microbiol Rev 2005;18(4):638-56. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Macintyre CR, Kainer MA, Brown GV. A randomised, clinical trial comparing the effectiveness of hospital and community-based reminder systems for increasing uptake of influenza and pneumococcal vaccine in hospitalised patients aged 65 years and over. Gerontology 2003;49(1):33-40. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Exclude - Not MMIT

Mack EH, Wheeler DS, Embi PJ. Clinical decision support systems in the pediatric intensive care unit. Pediatric Critical Care Medicine 2009;10(1):23-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Mack TA. Decision support considerations in the development and implementation of an electronic medical record. Pharm Pract Manag Q 1998;18(1):21-34. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

MacKinnon NJ. Clinical decision support systems for pharmacy. ASHP Annual Meeting 1998;55: Database: IPA.

Exclude - Not a Primary Study

Mackowiak LR, Hayward SL. Issues of decision support in institutional pharmacy systems. Pharm Pract Manag Q 1998;18(1):35-45. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Maddox RR. Bar code medication administration: Lessons learned from three years of preparation. Hosp Pharm 2003;38(11):S24-S25 Database: IPA.

Exclude - Not a Primary Study

Maggenti G. An electronic device to help elderly people with their home medications. In 2007; Tours, France) SUBJECT(S) Identifier Ubiquitous computing; Home automation; Self-help devices for people with disabilities; Medical telematics; Smart homes; Health telematics; ICOST Note(s) Includes bibliographical references and index.: 2009. p.278-81. Grey Lit.

Exclude - Not a Primary Study

Magid SK, Pancoast PE, Fields T, et al. Employing clinical decision support to attain our strategic goal: the safe care of the surgical patient. J Healthc Inf Manag 2007;21(2):18-25.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Magrabi F, McDonnell G, Westbrook JI, et al. Using an accident model to design safe electronic medication management systems. Stud Health Technol Inform 2007;129(Pt:2):2-52. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Mahan CE, Spyropoulos AC. Venous thromboembolism prevention: A systematic review of methods to improve prophylaxis and decrease events in the hospitalized patient. Hosp Pract 2010;38(1):97-108. PMID:20469630 OVID MEDLINE.

Exclude - Unable to Retrieve

Mahanney LM. Use of technology in the VA health system - From bar codes to automated dispensing, Department of Veterans Affairs implements them all. Journal of the Pharmacy Society of Wisconsin 2004;9-12. Database: IPA.

Exclude - No Outcomes of Interest

Mahoney CD, Wyman CM, Marcoux MA, et al. Remote automated dispensing system reduces costs and improves inpatient hospice care. Ashp Midyear Clinical Meeting 1997;32: Database: IPA.

Exclude - Not a Primary Study

Mahrous H. Developing clinical wireless applications for health care professionals. American Association of Colleges of Pharmacy Annual Meeting 2000;102: Database: IPA.

Exclude - Not MMIT

Mahrous S, Dalbik M. E-prescribing: Using technology to prevent errors. Pharm Times 2006;72(11):102-5. Database: IPA.

Exclude - Not a Primary Study

Majkowski K, Bell D, Lapane K et al. Results and impact of electronic prescribing (e-Rx) use: 3rd Teleconference in a series of four on the medicare modernization act e-Rx pilot evaluation. AHRQ; 2007.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_807449_0_0_18/Nov2_eRxWebConference.pdf Grey Lit.

Exclude - Not a Primary Study

Makinen M, Forsstrom J, Aarimaa M, et al. A European survey on the possibilities and obstacles of electronic prescriptions in cross-border healthcare. *Telemedicine Journal and e-Health* 2006;12(4):484-9. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Makris M. Alternative models of delivery of anticoagulant services. [Review] [26 refs]. *Seminars in Thrombosis & Hemostasis* 1999;25(1):33-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Malachowski M, Jones CD. Evaluation of infusion pump errors and the potential impact of "smart" pump technology. *ASHP Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Malato LA, Kim S. End-user perceptions of a computerized medication system: is there resistance to change? *J Health Hum Serv Adm* 2004;27(1):34-55. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Manchester GW, Raia TJ, III, Scott J, et al. Primary Care Health Information System: a hybrid electronic-paper medical record system. *J Ambulatory Care Manage* 1992;15(3):13-29. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Mander T. Change imminent. *Pharmacy Practice News* 2004;29(1):4 Database: IPA.

Exclude - Not a Primary Study

Mangione CM, Gerzoff RB, Williamson DF, et al. The association between quality of care and the intensity of diabetes disease management programs. *Ann Intern Med* 2006;145(2):107-16. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Manser AJ. Computerized physician order entry: Strategic issues for Australian hospital pharmacists. *Australian Journal of Hospital Pharmacy* 1996;26(6):664-7. Database: IPA.

Exclude - No Outcomes of Interest

Mansfield BP. Review of existing automation developments. *ASHP Annual Meeting* 2000;57: Database: IPA.

Exclude - Not a Primary Study

Mansfield BP. Selecting and implementing technology to decrease errors. *ASHP Annual Meeting* 2000;57: Database: IPA.

Exclude - Not a Primary Study

Mansfield BP. Steps to design and implement a system to meet your institution needs: Case study. *ASHP Annual Meeting* 2000;57: Database: IPA.

Exclude - Not a Primary Study

Mansfield BP, Mansfield BP. Northwestern Medical Center - Experiences in the development, implementation and testing of an ideal medication management system. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Mansour H, Dilkhush D, Lannigan J, et al. The impact of a computerized potassium alert on adverse drug events and pharmacists' interventions. J Pharm Technol 2010;26(2):55-9. <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010636870&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=967&accno=2010636870 EBSCO CINAHL.

Exclude - Unable to Retrieve

Mansur J. A guide to the joint commission's medication management standards. Second. 2009. Grey Lit.

Exclude - Not MMIT

Manzo J, Tourville J. Design, implementation and management of automated pharmacy application and medication management standards for 69 facilities within a nationwide for-profit healthcare organization. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Marceau LD, Link C, Jamison RN, et al. Electronic diaries as a tool to improve pain management: is there any evidence? PAIN MED 2007;8:Suppl-9 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Marco F, Sedano C, Bermudez A, et al. A prospective controlled study of a computer-assisted acenocoumarol dosage program. Pathophysiology of Haemostasis and Thrombosis 2003;33(2):59-63. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Marcon L, Battersby BJ, Ruhmann A, et al. 'On-the-fly' optical encoding of combinatorial peptide libraries for profiling of protease specificity. Molecular Biosystems 2010;6(1):225-33. PMID:20024084 OVID MEDLINE.

Exclude - Not a Primary Study

Marcum ZA, Handler SM, Boyce R, et al. Medication misadventures in the elderly: A year in review. American Journal Geriatric Pharmacotherapy 2010;8(1):77-83. OVID EMBASE.

Exclude - No Outcomes of Interest

Margolis A, Flores F, Kierszenbaum M, et al. Warfarin 2.0--a computer program for warfarin management. Design and clinical use. Proceedings - the Annual Symposium on Computer Applications in Medical Care 1994;846-50. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mariani G, Manotti C, Dettori AG. A computerized regulation of dosage in oral anticoagulant therapy. Ric Clin Lab 1990;20(2):119-25. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Marietti C. Robots hooked on drugs. Robotic automation expands pharmacy services. Healthc Inform 1997;14(11):37-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Marini SD, Hasman A. Impact of BCMA on medication errors and patient safety: a summary. *Stud Health Technol Inform* 2009;146:439-44. PMID:19592882 OVID MEDLINE.

Exclude - Unable to Retrieve

Marini SD, Hasman A, Huijer HA-S. Information technology for medication administration: Assessing bedside readiness among nurses in Lebanon. *International Journal of Evidence-Based Healthcare* 2009;7(1):49-58. OVID PsychINFO.

Exclude - No Outcomes of Interest

Marino BL, Branowicki P, Bennett JA, et al. Evaluating process changes in a pediatric hospital medication system. *Outcomes Management* 2002;6(1):10-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Marino J. Technology to drive administration's government reform efforts. *Caribbean Business* 2009;37(44):20-3.

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&b=bth&AN=45443376&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

Markind JE, Alessandrini JM, Pugh MC. Adverse drug reaction reporting system. *ASHP Annual Meeting* 1991;48: Database: IPA.

Exclude - Not a Primary Study

Marple R N. Dispensing medication via computer. In *Physicians & computers* 3. 1986. p.28-9, 33.1987020023510

Database: Compendex.

Exclude - Unable to Retrieve

Marshall PD, Chin HL. The effects of an Electronic Medical Record on patient care: clinician attitudes in a large HMO. *Proceedings / AMIA 1998;Annual Symposium.*:150-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Marshall S. Automating medications management. In *Physician* 9. 1988. p.23-7.1992090283903

Database: Compendex.

Exclude - Not MMIT

Martens G, Zapf CL. A reference system in clinical anesthesia. *J Clin Monit* 1993;9(3):202-6. 1993111087983

Database: Compendex.

Exclude - Not a Primary Study

Martens JD, van der AA, Panis B, et al. Design and evaluation of a computer reminder system to improve prescribing behaviour of GPs. *Stud Health Technol Inform* 2006;124:617-23. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Martin-Baranera M, Planas I, Palau J, et al. IMASIS computer-based medical record project: dealing with the human factor. *Medinfo* 1995;8 Pt 1:333 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Martin E. Improving influenza vaccination rates in a pediatric asthma management program by utilization of an electronic medical record. *Clin Pediatr (Phila)* 2006;45(3):221-7.
Database: Embase Sept 22-09.
Exclude - Not MMIT

Martin JA. Radical redesign of the drug distribution system by implementing automated point of use technology results in positive outcomes. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.
Exclude - Not a Primary Study

Martin MK, Shuster KP, Palisano TG. Connecticut RxData project. *Proceedings / AMIA* 2002;494-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Martin P. Health industry bar code in Europe. *Med Device Technol* 1992;3(Aug-Sep):32-4.
Database: IPA.
Exclude - Not a Primary Study

Martin P, Haefeli WE, Martin-Facklam M. A drug database model as a central element for computer-supported dose adjustment within a CPOE system. *J Am Med Inform Assoc* 2004;11(5):427-32. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Martinez C, Herranz A, Barrueco N, et al. Technology advances in drug management in a general hospital. *Ashp Summer Meeting* 2005;62: Database: IPA.
Exclude - Not a Primary Study

Marucci C, Karoly MB, Begliomini R, et al. Implementation of new robotic dispensing technology into a system with an existing bar code point of care program. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Maruta E, Shimizu Y, Matsui K. Merit, demerit and its countermeasure that control drugs and medical materials in the pharmacy with the monistical control system. *Japanese Journal of Hospital Pharmacy* 1998;24(2):192-200. Database: IPA.
Exclude - No Outcomes of Interest

Marvin KC, Kuperman GJ. Supporting CPOE now and in the future. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
Exclude - Not a Primary Study

Marx K. Smoothing the path. *Healthc Inform* 2009;26(9):50-1.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010429784&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1239&accno=2010429784
EBSCO CINAHL.
Exclude - Not a Primary Study

Marxen TL, Nemeth J, Tran D. Improving adverse drug reaction reporting via technology and education. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.
Exclude - Not a Primary Study

Maslakowski CJ. Implementing a decision support tool to maximize drug therapy outcomes. Ashp Midyear Clinical Meeting 1996;31: Database: IPA.
Exclude - Not a Primary Study

Massaro T. Introducing Physician Order Entry at a Major Academic Medical Center: I. Impact on ORganizational Culture and Behavior. Acad Med 1993;68(1):20-4. Exclude - No Outcomes of Interest

Massaro T. Introducing physician order entry at a major academic medical center: II. Impact on medical education. Acad Med 1993;68(1):25-30. Exclude - Not a Primary Study

Massaut J, Reper P. Open source electronic health record and patient data management system for intensive care. Studies in Health Technology & Informatics 2008;141:139-45. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Mathieson S. E-prescribing. Rewriting the script. Health Serv J 2004;114(5935):suppl-8 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Matsumura Y, Yamaguchi T, Hasegawa H, et al. Alert system for inappropriate prescriptions relating to patients' clinical condition. Methods Inf Med 2009;48(6):566-73. PMID:19893857 OVID MEDLINE.
Exclude - Unable to Retrieve

Matsushita O, Higuchi K, Ishii N, et al. Behavioral approach to facilitate appropriate use of antibiotics. Rinsho Byori - Japanese Journal of Clinical Pathology 2008;56(11):994-1006. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Mattocks K, Lalime K, Tate JP, et al. The state of physician office-based health information technology in Connecticut: current use, barriers and future plans. Conn Med 2007;71(1):27-31. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Matulewicz AN, Mahoney CD, Metzger J, et al. Automating medication control in emergency department critical care rooms and operating room surgical suites. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Maurer C, Lecointre K, Cachin N, et al. Impact of medical prescription computerisation on the incidence of adverse drug effects. Rev Mal Respir 2003;20(3:Pt:1):t-63 Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Maviglia S. Preventing dispensing errors with bar coding. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Maviglia SM, Yoon CS, Bates DW, et al. KnowledgeLink: impact of context-sensitive information retrieval on clinicians' information needs. *J Am Med Inform Assoc* 2006;13(1):67-73. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Maviglia SM, Yoo JY, Franz C, et al. Cost-benefit analysis of a hospital pharmacy bar code solution. *Arch Intern Med* 2007;167(8):788-94. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Mawer GE. Computer assisted prescribing of drugs. [Review] [38 refs]. *Clin Pharmacokinet* 1976;1(1):67-78. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Maxwell DJ. Observations of pharmacy practice in HIV treatment centres in the United Kingdom. *Journal of Pharmacy Practice and Research* 2003;33(4):282-3. Database: IPA.
Exclude - Not MMIT

May EL. The case for bar coding: better information, better care--and better business. *Healthc Exec* 2003;18(5):8-13. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

May MW, Quattlebaum C, Parsons K. Rx Mobile: patient information at the point of care. *Am J Health Syst Pharm* 2006;63(5):456-60. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Mayfield,S. A multivariate analysis of the effects of CPOE on hospital quality and patient safety Medical University of South Carolina - College of Health ProfessionsEditor. 2008. Grey Lit.
Exclude - Theses

Mayhorn C. Perfecting the handheld computer for older adults: From cognitive theory to practical application. *International Journal of Cognitive Technology* 2007;12(1): Database: PsycINFO.
Exclude - Not MMIT

Mayhorn CB, Lanzolla VR, Wogalter MS, et al. Personal digital assistants (PDAs) as medication reminding tools: Exploring age differences in usability. *Gerontechnology* 2005;4(3):128-40. 9041501
Database: Inspec.
Exclude - Not MMIT

Mazze R, Langer O. Sakamoto,N.;1991. Self-monitoring and education: Computer technologies in diabetes. Database: PsycINFO.
Exclude - Not MMIT.

Mazzuca S, Vinicor F, Einterz R, et al. Effects of the clinical environment on physicians' response to postgraduate medical education. *American Educational Research Journal* 1990;27(3):473-88. Exclude - Not MMIT

McAlister N, Covvey H, Tong C, et al. Randomised controlled trial of computer assisted management of hypertension in primary care. *Br Med J* 1986;293:670-4. Exclude - Not MMIT

McBride JM, Clark TC, Larson CM, et al. Multidisciplinary process to ensure effective implementation of an advanced physician order entry system. Ashp Midyear Clinical Meeting 1999;34: Database: IPA.

Exclude - Not a Primary Study

McBride M. Rising the bar: An Illinois medical center decides “really good” isn’t good enough and sets out to lower an already low adverse drug event rate. Health Manag Technol 2005;26(12):32-3. 8805870

Database: Inspec.

Exclude - Not a Primary Study

McCarthy D and Mueller K. The New York City health and hospitals corporation: Transforming a public safety net delivery system to achieve higher performance. The Commonwealth Fund; 2008.

http://www.commonwealthfund.org/publications/publications_show.htm?doc_id=710848

Grey Lit.

Exclude - Not MMIT

McCartney PR. Using technology to promote perinatal patient safety. JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing 2006;35(3):424-31. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

McClelland K. Scripts on the wire. Chemist and Druggist 2004;253:22 Database: IPA.

Exclude - Not MMIT

McClure DL. Improving drug safety: Active surveillance systems should be paramount. Pharmaceutical Medicine 2009;23(3):127-30. OVID EMBASE.

Exclude - No Outcomes of Interest

McCluskey CF, Hammond RL. Implementation of a barcoded medication software system within an intravenous admixture area to track medication preparation and delivery, and improve workflow. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

McCormick EM. Allcare Medication Services offers consultation to long-term care facilities. Pharm Times 2004;55(Nov):36-7. Database: IPA.

Exclude - Not a Primary Study

McCowan C, Neville RG, Ricketts IW, et al. Lessons from a randomized controlled trial designed to evaluate computer decision support software to improve the management of asthma. Med Inform Internet Med 2001;26(3):191-201. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

McCoy C, Cunningham K, Afonso K, et al. Development and use of a restricted antimicrobial ordering process through a novel cPOE program. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

McCoy C. Evaluation of an computerized provider order entry (CPOE) indication based pharmacodynamic dosing model for meropenem. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

McCreadie SR, Stumpf JL, Benner TD. Building a better online formulary. Am J Health Syst Pharm 2002;59(19):1847-52. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

McCreadie SR, McGregory M, Siden R, et al. Development of a custom information system for an investigational drug service. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.
Exclude - Not a Primary Study

McCullough JS, Casey M, Moscovice I, et al. The effect of health information technology on quality in U.S. hospitals. Health Aff (Millwood) 2010;29(4):647-54.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010629265&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1759&accno=2010629265
EBSCO CINAHL.
Exclude - No Outcomes of Interest

McDaniel MR, DeJong DJ. Is the pharmacy department a sound investment? How one department documented the return on investment (ROI) to senior leadership. Ashp Midyear Clinical Meeting 1997;32: Database: IPA.
Exclude - Not a Primary Study

McDonald CJ, Tierney W.M. The medical gopher - A microcomputer system to help find, organize and decide about patient data. West J Med 1986;145(6):823-9. Database: Embase Sept 22-09.
Exclude - Not MMIT

McDonald CJ, Overhage JM, Tierney WM, et al. The Regenstrief Medical Record System: a quarter century experience. Int J Med Inf 1999;54(3):225-53. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

McDonald CJ. Computerization can create safety hazards: a bar-coding near miss. Ann Intern Med 2006;144(7):510-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

McDonald C, Hui S, Smith D, et al. Reminders to physicians from an introspective computer medical record. Ann Intern Med 1984;100(1):130-8. Exclude - Not MMIT

McDonald C, Hui S, Tierney W. Effects of computer reminders for influenza vaccination on morbidity during influenza epidemics. MD Comput 1992;9(5):304-12. Exclude - Not MMIT

McEvoy GK. Personal digital assistants (PDAS): Mobil access to information at the point of patient care. International Pharmaceutical Federation World Congress 2004;62:144 Database: IPA.
Exclude - Not MMIT

McEvoy M. The road to CPOE, an epoch journey. Keeposted 2004;30(2):10-1. Database: IPA.
Exclude - Not MMIT

McGarry, N. A study of perceived efficiency and perceived effectiveness when using healthcare informatics: A study at the District of Columbia Veterans Affairs Hospital The George Washington University Editor. 2006. Grey Lit.

Exclude - Not MMIT

McGee MK. Time for the industry to take its medicine. InformationWEEK 2005;(1056):120-8782001

Database: Inspec.

Exclude - Not a Primary Study

McGerald G, Dvorkin R, Levy D, et al. Prescriptions for schedule II opioids and benzodiazepines increase after the introduction of computer-generated prescriptions. Acad Emerg Med 2009;16(6):508-12. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

McGowan JJ, Richwine M. Electronic information access in support of clinical decision making: a comparative study of the impact on rural health care outcomes. Proceedings / AMIA 2000;565-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

McGregory ME, Burnham RC, Ginsberg-Evans LR, et al. Development of custom, barcode-enabled label printing software to improve the safety of medication repackaging. Ashp Midyear Clinical Meeting 2006;41. Database: IPA.

Exclude - Not a Primary Study

McKeough T. A Smart Sensor. Fast Company 2009;(141):66

<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=45425320&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.

Exclude - Not a Primary Study

McKinney M. Most improved winners follow varied paths. Hospitals & health networks / AHA 1940;84(1):38, 40. OVID EMBASE.

Exclude - Not a Primary Study

McKinney M. Decision support for docs. Hospitals & Health Networks 1944;83(12):42 PMID:20112759 OVID MEDLINE.

Exclude - Not a Primary Study

McKinney M. Plan before the panic. Hospitals & health networks / AHA 2009;83(11):35-6. OVID EMBASE.

Exclude - Not a Primary Study

McKinnon PS, Seaton T, Noirot LA, et al. Improving adherence to dyslipidemia medication guidelines in hospitalized diabetic patients using a technology-assisted pharmacist intervention. AMIA 2008;Annual:Symposium PMID:18999065 OVID MEDLINE.

Exclude - Not MMIT

McLaren S. Automated dispensing: Will new technology end drudgery or automate chaos? Can Pharm J 2004;128(Feb):16-9. Database: IPA.

Exclude - Not MMIT

McLaughlin CP. Just-in-time (JIT) for the patient. *Carolina Journal of Pharmacy* 1991;71(Sep):15-8. Database: IPA.
Exclude - Not MMIT

McLean AJ, Barned J, Ioannides-Demos L, et al. Computer-based prescription audit as a research, educational and management tool. *Aust Health Rev* 1984;7(4):260-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

McLean WM, MacKeigan LD. When does pharmaceutical care impact health outcomes? A comparison of community pharmacy-based studies of pharmaceutical care for patients with asthma. *Ann Pharmacother* 2005;39(4):625-31. Database: IPA.
Exclude - Not MMIT

McLeod SE, Lum E, Mitchell C. Value of medication reconciliation in reducing medication errors on admission to hospital. *Journal of Pharmacy Practice and Research* 2008;38(3):196-9. Database: Embase Sept 22-09.
Exclude - Not MMIT

McMichael J, Lieberman R, Doyle H, et al. An intelligent and cost-effective computer dosing system for individualizing FK506 therapy in transplantation and autoimmune disorders. *SO: Journal of clinical pharmacology* 1993;33(7):599-605.
<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/566/CN-00095566/frame.html> Database: Cochrane.
Exclude - Not MMIT

McMullin ST, Reichley RM, Kahn MG, et al. Automated system for identifying potential dosage problems at a large university hospital. *Am J Health Syst Pharm* 1997;54(5):545-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

McMullin ST. Standardization is necessary in the methods to assess the value of electronic prescribing systems. *J Manag Care Pharm* 2005;11(7):594-5. Database: IPA.
Exclude - Not MMIT

Mcneill BJ. Physician's clinic perspective on utilizing electronic prescribing. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

McNulty J, Donnelly E, Iorio K. Methodologies for sustaining barcode medication administration compliance. A multi-disciplinary approach. *J Healthc Inf Manag* 2009;23(4):30-3. PMID:19894484 OVID MEDLINE.
Exclude - No Outcomes of Interest

McNulty RM. CPOE chemotherapy ordering in a cancer hospital. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

McPhillips HA, Stille CJ, Smith D, et al. Potential medication dosing errors in outpatient pediatrics. *J Pediatr* 2005;147(6):761-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

McRoberts S. The use of bar code technology in medication administration. *Clin Nurse Spec* 2005;19(2):55-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Meadows M. Strategies to reduce medication errors. *FDA Consum* 2003;37(3):20-7.
Database: IPA.
Exclude - Not MMIT

Medina-Cuevas F, Navarrete-Navarro S, Avila-Figueroa C, et al. FARMAC: a program designed for monitoring the prescription of antimicrobials in hospitals. *Gac Med Mex* 2000;136(2):107-11. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Meglic M, Ivanovski M, Marusic A. Information technology tools to improve treatment of patients with depression: Focus on guidelines implementation. *Psychiatria Danubina* 2008;20(2): Database: PsycINFO.
Exclude - Unable to Retrieve Foreign

Mehta D, McCormack J, Fung P, et al. Target controlled infusion for kids: trials and simulations. *Conference Proceedings: 2008;2008:5818-21*. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Mehta R, Onatade R. Experience of inpatient electronic prescribing (EP) in UK hospitals: A perspective from pharmacy staff. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Meldrum D. Simple computerised repeat prescription control system. *British Medical Journal Clinical Research Ed* 1981;282(6280):1933-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Menachemi N, Burke D, Clawson A, et al. Information technologies in Florida's rural hospitals: does system affiliation matter? *J Rural Health* 2005;21(3):263-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Menachemi N, Brooks RG. Reviewing the benefits and costs of electronic health records and associated patient safety technologies. *J Med Syst* 2006;30(3):159-68. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Mendribil RG, Gravois L. Effective Pyxis inventory management utilizing Cardinal ASSIST reporting. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Menduno M. Software that plays hardball. Expert clinical systems fend off forgetfulness, mistakes, and fraud investigators. *Hosp Health Netw* 1998;72(10):44 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Menighan TE. The world of E-prescribing: Implications for pharmacy practice. *Computertalk for the Pharmacist* 2002;22(1):13-4. Database: IPA.
Exclude - Not a Primary Study

Mercer G. Industrywide standard does not exist, but latest printing technology lets barcode data fit on 2 mL drug vial. *U S Pharmacist* 1990;27(6):90 Database: IPA.

Exclude - Not MMIT

Meropol SB, Chan K, Chen Z, et al. Adverse events associated with prolonged antibiotic use. *Pharmacoepidemiology and Drug Safety* 2008;17(5):523-32. Database: Embase Sept 22-09.

Exclude - Not MMIT

Merry AF, Webster CS, Mathew DJ. A new, safety-oriented, integrated drug administration and automated anesthesia record system. *Anesthesia & Analgesia* 2003;93(2):385-90.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Merryfield DW, Degnan DD, Keller TE, et al. Electronic adverse drug event trigger measurement. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Merryfield DW, Steining KK, Gilley AE, et al. Accountability for clinical alert responses. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Mertz K. State legislatures and health information technology. In *National Conference of State Legislatures*; 2008. http://www.ncsl.org/print/health/forum/NCSL_HIMSS.pdf Grey Lit.

Exclude - Not a Primary Study

Metzen D, Capers W. Impact of computerized prescriber order entry (CPOE) on medication turnaround times. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Metzger J, Welebob E, Bates DW, et al. Mixed results in the safety performance of computerized physician order entry. *Health Aff (Millwood)* 2010;29(4):655-63.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010629266&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1759&accno=2010629266

EBSCO CINAHL.

Exclude - No Outcomes of Interest

Meyer GE, Smith JE, Brandell R. Application of bar codes to the hospital pharmacy unit dose medication dispensing system. *Ashp Midyear Clinical Meeting* 1989;24: Database: IPA.

Exclude - Not a Primary Study

Meyer GE, Brandell R, Smith JE, et al. Use of bar codes in inpatient drug distribution. *Am J Hosp Pharm* 1991;48(5):953-66. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Miasso AI, Oliveira RC, Silva AE, et al. Prescription errors in Brazilian hospitals: a multi-centre exploratory survey. *Cad Saude Publica* 2009;25(2):313-20. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Michener LM, Boecler L, Kent SS. Development and implementation of an adverse drug event alert system utilizing automated dispensing cabinets. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Mickle TR, Haymond J, Bess DT, et al. A pilot study of a clinical pharmacist supervised remote order entry program in a veterans administration medical center ambulatory care clinic. ASHP Annual Meeting 1990;47: Database: IPA.

Exclude - Not a Primary Study

Middleton B, Gandhi T, Bates D. The role of information technology in ambulatory care patient safety. 2002. <http://www.himss.org/ASP/ContentRedirector.asp?ContentID=60708> Grey Lit.

Exclude - Not a Primary Study

Middleton B. National committee on vital and health statistics (NCVHS): Statement of Blackford Middleton. Healthcare Information and Management Systems Society (HIMSS); 2006. http://www.himss.org/content/files/NCVHS_Testimony_Middleton.pdf Grey Lit.

Exclude - Not a Primary Study

Middleton B, Fleming M, Wiegand T et al. Best practices for community health information exchange. Center for Community Health Leadership; 2007.

http://www.misyscenter.com/NR/rdonlyres/6B8E9E8A-93BD-467D-A3BB-52B0E4DC6107/0/CCHL_BPG.pdf Grey Lit.

Exclude - Not a Primary Study

Milazzo CF. A pocket computer program for common calculations in clinical pediatrics. Comput Biol Med 1985;15(1):41-55. 2424486

Database: Inspec.

Exclude - Not a Primary Study

Militello K, Dolan C, Hansen CL, et al. Impact of pharmacist medication error prevention: Computerized mechanism to incorporate pharmacist interventions into MedMARx. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.

Exclude - Not a Primary Study

Millan J, Park S E, Kiefer S and others. TOPCARE: Implementation of a telematic homecare platform in cooperative health care provider networks. In Houston, TX, United states: Institute of Electrical and Electronics Engineers Inc.; 2002. p.1869-70.2003027310078 Database: Compendex.

Exclude - Not a Primary Study

Mille F, Degoulet P, Jaulent MC. Modeling and acquisition of drug-drug interaction knowledge. Stud Health Technol Inform 2007;129(Pt:2):2-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Miller A. Issues related to the development of a computer prescriber order entry (CPOE) system. Advances in Pharmacy 2002;1(1):28-45. Database: IPA.

Exclude - Not MMIT

Miller A. Incorporation of medication safety initiatives within a computerized prescriber order entry (CPOE) system. Ashp Summer Meeting 2004;61: Database: IPA.
Exclude - Not a Primary Study

Miller A. Patient safety initiatives in a computerized physician order entry system (HPS-P-354). International Pharmaceutical Federation World Congress 2004;62: Database: IPA.
Exclude - No Outcomes of Interest

Miller AS. Implementing physician order entry: Pharmacy's perspective. Ashp Midyear Clinical Meeting 1998;33: Database: IPA.
Exclude - Not a Primary Study

Miller AS. Multidisciplinary approach to implementing physician order entry: Pharmacy overview. Ashp Midyear Clinical Meeting 1999;34: Database: IPA.
Exclude - Not a Primary Study

Miller AS. Pharmacy issues: Drug nomenclature. Hosp Pharm 2000;35(Nov):1249-51.
Database: IPA.
Exclude - Not MMIT

Miller AS. Computerized prescriber order entry: Pharmacy issues: Clinical screenings and discharge prescriptions. Hosp Pharm 2001;36(12): Database: Embase Sept 22-09.
Exclude - Not MMIT

Miller AS. Pharmacy issues: Bidirectional order interface. Hosp Pharm 2001;36(Oct):1112-5.
Database: IPA.
Exclude - Not MMIT

Miller AS. Pharmacy issues: Unique IV drug ordering--epidural anesthesia. Hosp Pharm 2001;36(May):563-5. Database: IPA.
Exclude - Not MMIT

Miller AS. Pharmacy issues: Weight based dosing. Hosp Pharm 2001;36(Jul):790-4.
Database: IPA.
Exclude - Not MMIT

Miller AS. Review of patient safety outcomes from a PCOE system at an academic medical center. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.
Exclude - Not a Primary Study

Miller AS. Computerized prescriber order entry: The admission, transfer, and discharge process. Hosp Pharm 2002;37(1): Database: Embase Sept 22-09.
Exclude - Not MMIT

Miller AS. Computerized prescriber order entry. Hosp Pharm 2002;37(11):1218-21.
Database: IPA.
Exclude - Not MMIT

Miller AS. Nursing and pediatric issues in promoting safe medication practice within a CPOE system. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.
Exclude - Not a Primary Study

Miller AS. Nursing issues. Hosp Pharm 2002;37(3):321-2. Database: IPA.
Exclude - Not MMIT

Miller AS. Quality and operations improvement: Order set development and maintenance. Hosp Pharm 2002;37(7):765-8. Database: IPA.
Exclude - Not MMIT

Miller AS. Security issues. Hosp Pharm 2002;37(8):867-8. Database: IPA.
Exclude - Not MMIT

Miller AS. The implementation process (Part 3). Hosp Pharm 2002;37(12):1343-4. Database: IPA.
Exclude - Not MMIT

Miller AS. The implementation process (Part 1). Hosp Pharm 2002;37(10):1104-6. Database: IPA.
Exclude - Not MMIT

Miller AS. Computer prescriber order entry. Hosp Pharm 2003;38(8):794-8. Database: IPA.
Exclude - Not MMIT

Miller AS. Computerized prescriber order plan - Pharmacy work plan. Hosp Pharm 2003;38(10):981-4. Database: IPA.
Exclude - Not MMIT

Miller AS. Creating, maintaining and updating order sets in CPOE. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.
Exclude - Not a Primary Study

Miller AS. Downtime procedures (Part 2). Hosp Pharm 2003;38(7):694-7. Database: IPA.
Exclude - Not MMIT

Miller AS. Pharmacy issues: Chemotherapy. Hosp Pharm 2004;36(Sep):992-6. Database: IPA.
Exclude - Not MMIT

Miller AS. Pharmacy issues: Drug routes, dosage forms, and drug frequencies. Hosp Pharm 2004;35(Dec):1342-6. Database: IPA.
Exclude - Not MMIT

Miller AS. Pharmacy issues: Formulary changes and allergy checking. Hosp Pharm 2004;36(Nov):1209-10. Database: IPA.
Exclude - Not MMIT

Miller AS. The glory and the chaos of CPOE. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.
Exclude - Not a Primary Study

Miller BJ. MAR: Medication administration record corrections reduce medication errors through order entry education. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.
Exclude - Not a Primary Study

Miller DA, Zarowitz BJ, Petitta A, et al. Pharmacy technicians and computer technology to support clinical pharmacy services. Am J Hosp Pharm 1993;50(5):929-34. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Miller J and Eskew A. Computerized physician order entry. 2003.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=30507> Grey Lit.
Exclude - Not a Primary Study

Miller J. Bar code requirement would lead to adoption of efficient technology. *Managed Healthcare Executive* 2003;13(5):42-3. Database: IPA.
Exclude - Not a Primary Study

Miller PL, Frawley SJ. Trade-offs in producing patient-specific recommendations from a computer-based clinical guideline: a case study. *J Am Med Inform Assoc* 1995;2(4):238-42. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Miller R. Stop the carts. *Nurs Manag (Harrow)* 1999;30(1):44-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Miller RA, Gardner RM, Johnson KB, et al. Clinical decision support and electronic prescribing systems: a time for responsible thought and action. *J Am Med Inform Assoc* 2005;12(4):403-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Miller RS, Norris PR, Jenkins JM, et al. Systems initiatives reduce healthcare-associated infections: a study of 22,928 device days in a single trauma unit. *J Trauma* 2010;68(1):23-31. <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010528631&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=762&accno=2010528631 EBSCO CINAHL.
Exclude - Not MMIT

Miller R, Waiman L, Chen S, et al. The anatomy of decision support during inpatient care provider order entry (CPOE): Empirical observations from a decade of CPOE experience at Vanderbilt. *Journal of Biomedical Informatics* 2005;38:469-85. Exclude - Not a Primary Study

Miller WA. Computer oriented clinical hospital pharmacy services: A conceptual model. *J Clin Comput* 1974;3(Jan):276-87. Database: IPA.
Exclude - Not MMIT

Mills PD, Neily J, Mims E, et al. Improving the bar-coded medication administration system at the Department of Veterans Affairs. *Am J Health Syst Pharm* 2006;63(15):1442-7. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Mills PD, Neily J, Kinney LM, et al. Effective interventions and implementation strategies to reduce adverse drug events in the Veterans Affairs (VA) system. *Quality & Safety in Health Care* 2008;17(1):37-46. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Mills S, Blackburn C. Delivering quality. Rapid implementation of a centrally hosted, interoperable ambulatory EHR throughout all hospital markets of the Franciscan Missionaries of Our Lady Health System. *J Healthc Inf Manag* 2009;22(1):20-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Milstead JA. Leapfrog Group: a prince in disguise or just another frog? *Nurs Adm Q* 2002;26(4):16-25. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Milstein C, Venot A. Cost-related information to be provided by computerised drug-prescription systems to promote cost-effective prescribing. [Review] [18 refs]. *Pharmacoeconomics* 1997;12(2 Pt 1):130-9. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Mims E, Tucker C, Carlson R, et al. Quality-monitoring program for bar-code-assisted medication administration. *Am J Health Syst Pharm* 2009;66(12):1125-31. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Minard PD, Hanson DS, Bodner A, et al. Implementing changes in a private health care system. *ASHP Annual Meeting* 1999;56: Database: IPA.

Exclude - Not a Primary Study

Mir C, Gadri A, Zelger GL, et al. Impact of a computerized physician order entry system on compliance with prescription accuracy requirements. *Pharm World Sci* 2009;31(5):596-602. PMID:19554471 OVID MEDLINE.

Exclude - Not MMIT

Mishra S and Singh I. mHealth: A developing country perspective. 2008.

http://www.ehealth-connection.org/files/conf-materials/mHealth_%20A%20Developing%20Country%20Perspective_0.pdf Grey Lit.

Exclude - Not a Primary Study

Mitchell D. Early adopters and lemmings: Look before you leap into electronic records. *Physician Exec* 2005;31(4):18-22. Database: BSC.

Exclude - Not a Primary Study

Mitchell E, Sullivan F. A descriptive feast but an evaluative famine: systematic review of published articles on primary care computing during 1980-97. *BMJ* 2001;322(7281):279-82. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Mitra R, Marciello MA, Brain C, et al. Efficacy of computer-aided dosing of warfarin among patients in a rehabilitation hospital. *Am J Phys Med Rehabil* 2005;84(6):423-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mitrano EA, Clapp MD, Gordon Y, et al. Implementation of a bar code IV tracking system to reduce IV waste. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Mitrano EA, Clapp M, Gordon Y, et al. Implementation of a bar code IV tracking system to reduce IV waste. ASHP Annual Meeting 1997;54: Database: IPA.

Exclude - Not a Primary Study

Miyo K, Nittami YS, Kitagawa Y, et al. Development of case-based medication alerting and recommender system: a new approach to prevention for medication error. Studies in Health Technology & Informatics 2007;129(Pt:2):2-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mobach MP. The merits of a robot: a Dutch experience. Journal of Pharmacy & Pharmaceutical Sciences 2006;9(3):376-87. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Modai I, Sigler M. A computerized laboratory alerting system in a psychogeriatric unit. MD Comput 1998;15(2):95-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mohammad M, Kiffel C, Oliary J, et al. Physician evaluation of pharmacist interventions in an elderly ward. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Mohanalakshmi S, Jacob J. Design and implementation of an intelligent instrument for apnea patients. In Stevenage, UK: IET; 2007. p.562-6.10215173

Database: Inspec.

Exclude - Not a Primary Study

Mold A, Holden J, Miller DG. Pharmacy staff's perception of the effect of Electronic Prescribing (EP) on hospital pharmacy technician's future role development. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Molesa CS, Milad A. Conversion to low-dose acetaminophen containing combination analgesics to prevent acetaminophen toxicity. Ashp Summer Meeting 2007;64: Database: IPA.

Exclude - Not a Primary Study

Mollon B, Chong J, Jr., Holbrook AM, et al. Features predicting the success of computerized decision support for prescribing: a systematic review of randomized controlled trials. BMC Med Inform Decis Mak 2009;9:11 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Moniz B. Examining the unintended consequences of computerized provider order entry system implementation. Online Journal of Nursing Informatics 2009;13(1):1-12. Database: CINAHL.

Exclude - Not a Primary Study

Monroe S, Polk R. Antimicrobial use and bacterial resistance. Curr Opin Microbiol 2000;3(5):496-501. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Montani S, Bellazzi R, Quaglini S, et al. Meta-analysis of the effect of the use of computer-based systems on the metabolic control of patients with diabetes mellitus. *Diabetes Technol Ther* 2001;3(3):347-56. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Montelius E, Astrand B, Hovstadius B, et al. Individuals appreciate having their medication record on the web: a survey of attitudes to a national pharmacy register. *J Med Internet Res* 2008;10(4):e35 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Montgomery AA, Fahey T. A systematic review of the use of computers in the management of hypertension. *J Epidemiol Community Health* 1998;52(8):520-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Montgomery PA, Kan JM, Schaad AL. Systematic interventions to reduce enoxaparin doses in renal dysfunction and lower the rate of bleeding. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Montoya ID, Jano E. Online pharmacies: safety and regulatory considerations. *Int J Health Serv* 2007;37(2):279-89. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Moore JL, Hobbs RA, Mutnick BA, et al. Implementation of a quality assurance project to assess compliance with acute myocardial infarction oryx core measures and national guidelines. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Moore SM, Duffy E. Maintaining vigilance to promote best outcomes for hospitalized elders. *Crit Care Nurs Clin North Am* 2007;19(3):313-9. Database: CINAHL.

Exclude - Not a Primary Study

Moore TD, Dzierba SH, Miller A. Pharmacy computer system at the Ohio State University Hospitals. *Am J Hosp Pharm* 1984;41(Nov):2384-9. Database: IPA.

Exclude - Not MMIT

Morehead MD, Washington TG, Schneider R. Computer technology and the continuous improvements of the medication administration and distribution process. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Morelli J. Electronic Rx prescribing: Big business, promising future. *Drug Topics* 1997;141:19S-20S. Database: IPA.

Exclude - Not a Primary Study

Moreno P, Calleja MA, Orts A, et al. Improving pharmaceutical care through a new physician order entry computerized system. *Ashp Midyear Clinical Meeting* 1999;34: Database: IPA.

Exclude - Not a Primary Study

Moreno V, Farre M, Salva P. A program for oriented treatment of essential hypertension. *Comput Methods Programs Biomed* 1989;29(2):89-94. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Morera T, Gervasini G, Carrillo JA, et al. Early detection of drug interactions utilizing a computerized drug prescription handling system-focus on cerivastatin-gemfibrozil. *Eur J Clin Pharmacol* 2004;59(12):917-21. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Morgan B. Web portals enable electronic orders and results reporting. *MLO Med Lab Obs* 2009;41(12):26-7.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010512935&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=246&accno=2010512935 EBSCO CINAHL.
Exclude - Not a Primary Study

Morgan MW. In pursuit of a safe Canadian healthcare system. *Healthcarepapers* 2004;5(3):10-26. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Morgan MW. The VA advantage: the gold standard in clinical informatics. *Healthcarepapers* 2005;5(4):26-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Morice V, Seroussi B, Boisvieux JF. A real time control architecture for continuously managing patients in a care unit. *Methods Inf Med* 1995;34(5):475-88. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Morrell R, Wasilaukas B, Winslow R. Expert systems.[Review] [37 refs]. *Am J Hosp Pharm* 1994;51(16):2022-30. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Morrill TS. Automated pharmacy systems: Avoiding the pitfalls. *Hosp Mater Manage Q* 1982;4(Nov):79-85. Database: IPA.
Exclude - Not MMIT

Morris AH. Rational use of computerized protocols in the intensive care unit. *Critical Care (London, England)* 2001;5(5):249-54. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Morris AH. Decision support and safety of clinical environments. *Quality & Safety in Health Care* 2002;11(1):69-75. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Morris CJ, Cantrill JA, Avery AJ, et al. Preventing drug related morbidity: a process for facilitating changes in practice. *Quality & Safety in Health Care* 2006;15(2):116-21. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Morriss FH. Effectiveness of medication barcode scanning in a neonatal intensive care unit. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Morriss FH, Jr., Abramowitz PW, Carmen L, et al. "Nurses Don't Hate Change" -- survey of nurses in a neonatal intensive care unit regarding the implementation, use and effectiveness of a bar code medication administration system. *HEALTHC Q* 2009;12:Spec-40
PMID:19667791 OVID MEDLINE.

Exclude - Not a Primary Study

Morrissey J. An info-tech disconnect. Even as groups such as Leapfrog push IT as an answer to quality issues, doctors and executives say, 'not so fast'. *Mod Healthc* 2003;33(6):6-7.
Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Morrissey J. Too much too soon? JCAHO's proposed 2007 deadline for bedside bar-code technology has some providers scurrying, but vendors call the plan doable. *Mod Healthc* 2004;34(17):6-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Morrissey PE, Flynn ML, Lin S. Medication noncompliance and its implications in transplant recipients. *Drugs* 2007;67(10):1463-81. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Morsch M, Sheffer R, Glass S et al. The impact of physician quality measures on the coding process. *Perspectives in Health Information Management, CAC Proceedings*, 2008.
http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_040487.pdf Grey Lit.

Exclude - Not MMIT

Mosen D, Elliott CG, Egger MJ, et al. The effect of a computerized reminder system on the prevention of postoperative venous thromboembolism. *Chest* 2004;125(5):1635-41.
Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Moskovitz RM, Glowczewski JE, Moran MT, et al. Improving documentation of clinically significant interventions by staff pharmacists. *Ashp Midyear Clinical Meeting* 2005;40:
Database: IPA.

Exclude - Not a Primary Study

Motykie GD, Mokhtee D, Zebala LP, et al. The use of a Bayesian forecasting model in the management of warfarin therapy after total hip arthroplasty. *J Arthroplasty* 1999;14(8):988-93. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mount JC, Lewis K, Mason K, et al. Implementation of an automated patient-specific medication storage and management solution. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Mourtou E, Papathanasopoulos P, Pavlidis G. Implementing barcode technology in a Greek hospital: Experience and benefits. *Journal on Information Technology in Healthcare* 2007;5(2):83-96. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Moxey A, Robertson J, Newby D, et al. Computerized clinical decision support for prescribing: provision does not guarantee uptake. [Review] [78 refs]. J Am Med Inform Assoc 2010;17(1):25-33. PMID:20064798 OVID MEDLINE.

Exclude - Not a Primary Study

Mrozik W. Tracking and controlling UPS packages: A close parallel to drug delivery. Hosp Pharm 2003;38(11):S18-S19 Database: IPA.

Exclude - Not MMIT

Mukhopadhyay I, Lally F, Crome P. Appropriate prescribing in older people. Reviews in Clinical Gerontology 2002;17(2):139-51. Database: CINAHL.

Exclude - Not a Primary Study

Mull HJ, Nebeker JR. Informatics tools for the development of action-oriented triggers for outpatient adverse drug events. AMIA 2008;Annual:Symposium-9 PMID:18999297 OVID MEDLINE.

Exclude - Not MMIT

Mullaji A, Shetty GM. Computer-assisted TKA: greater precision, doubtful clinical efficacy: opposes. Orthopedics 2009;32(9):679

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010413373&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1212&accno=2010413373>

EBSCO CINAHL.

Exclude - Not a Primary Study

Muller R, Serogl M, Nauwerth U, et al. THEMPO: a knowledge-based system for therapy planning in pediatric oncology. Comput Biol Med 1997;27(3):177-200. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Muller RJ, Ernst TE. Implementation of a comprehensive quality assurance program in a major cancer center. Ashp Midyear Clinical Meeting 1991;26: Database: IPA.

Exclude - Not a Primary Study

Muller RJ, Ernst TE. Implementation of a comprehensive quality assurance program in a major cancer center. ASHP Annual Meeting 1991;48: Database: IPA.

Exclude - Not a Primary Study

Mullett CJ, Thomas JG, Smith CL, et al. Computerized antimicrobial decision support for hospitalized patients with a bloodstream infection. AMIA 2003;946 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mullett CJ, Thomas JG. Database-driven computerized antibiotic decision support: novel use of expert antibiotic susceptibility rules embedded in a pathogen-antibiotic logic matrix. AMIA 2003;480-3. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mullett CJ, Thomas JG, Smith CL, et al. Computerized antimicrobial decision support: an offline evaluation of a database-driven empiric antimicrobial guidance program in hospitalized patients with a bloodstream infection. *Int J Med Inf* 2004;73(5):455-60.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Mulligan M. System breakdown. 4 Med errors 250 scripts/day Solution: Pharmacist-driven initiatives. *Drug Topics* 2009;153(7):22-5. Database: IPA.

Exclude - Not a Primary Study

Mundy D, Chadwick DW. Electronic transmission of prescriptions: towards realising the dream. *International Journal of Electronic Healthcare* 2004;j.(1):112-25. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Mundy D, Chadwick D, Ball E. Some expectations and perceptions of electronic transfer of prescription systems. *BJHC & IM* 2004;21(2):34-6. Database: CINAHL.

Exclude - No Outcomes of Interest

Mundy DP, Chadwick DW. Security issues in the electronic transmission of prescriptions. *Med Inform Internet Med* 2003;28(4):253-77. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Mundy JP, Atkins CZ. Integration of clinical pharmacy services with handheld PDAs. *Ashp Midyear Clinical Meeting* 2001;36. Database: IPA.

Exclude - Not a Primary Study

Mungall D, Lord M, Cason S, et al. Developing and testing a system to improve the quality of heparin anticoagulation in patients with acute cardiac syndromes. *Am J Cardiol* 1998;82(5):574-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Munyer TO. Computer software technology in the 1990s. *Florida Pharmacy Today* 1991;55(Jan):6-8. Database: IPA.

Exclude - Not MMIT

Murchie CJ. Computer control of arterial blood pressure following cardiac surgery. *Intensive Care Nurs* 1987;3(1):3-7. Database: CINAHL.

Exclude - Not a Primary Study

Murff HJ. Medication errors in hospital care: Incidence and reduction strategies. *Journal of Pharmaceutical Finance* 2006;15(4):5-71. Database: CINAHL.

Exclude - No Outcomes of Interest

Murphy EM, Oxencis CJ, Klauck JA, et al. Medication reconciliation at an academic medical center: Implementation of a comprehensive program from admission to discharge. *Am J Health Syst Pharm* 2009;66(23):2126-31. OVID EMBASE.

Exclude - Not MMIT

Murphy JE, Malone DC, Olson BM, et al. Development of computerized alerts with management strategies for 25 serious drug-drug interactions. *Am J Health Syst Pharm* 2009;66(1):38-44. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Murray E. Managing anticoagulation. Practice Nurse 2009;33(7):21-2. Database: CINAHL.
Exclude - No Outcomes of Interest

Murray MA. Building a compliance team in pharmacy. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Murray MD, Young JM, Morrow DG, et al. Methodology of an ongoing, randomized, controlled trial to improve drug use for elderly patients with chronic heart failure. American Journal Geriatric Pharmacotherapy 2004;2(1):53-65. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Musser DB. The effect of a Technology-Linking Care (TLC) intervention on HIV/AIDS patients' therapeutic drug regimen adherence. University of Missouri - Saint Louis 2001; Ph D 2003; Database: CINAHL.
Exclude - Theses

Mutnick AH, Wong PK. Computerized prescriber order entry (CPOE): Does it hurt or help pharmaceutical care? Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Mutnick AH, Volles DF, Lyman JA. Piloting a pharmacy-based automated adverse drug event monitoring and prevention system. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Mutter M. One hospital's journey toward reducing medication errors. Joint Commission Journal on Quality & Safety 2003;29(6):279-88. Database: CINAHL.
Exclude - No Outcomes of Interest

Myers C, Lineen J. innovations in healthcare finance lessons from the 401(k) model. HFM 2008;62(10):38-42.
<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=34786700&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.
Exclude - Not a Primary Study

Myers TF, Venable HH, Hansen JA. Computer-enhanced neonatology practice evolution in an academic medical center. NICU Clinical Effectiveness Task Force. J Perinatol 1998;18(6 suppl 2):S38-S44 Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve

Mygatt PS. Too many numbers. Computertalk for the Pharmacist 1991;11(Nov-Dec):24-5. Database: IPA.
Exclude - Not a Primary Study

Myotoku M, Igarashi E, Kawaguchi S, et al. Survey of code printing on injections to prevent errors in dispensing injections. Japanese Journal of Pharmaceutical Health Care & Sciences 2003;29(5):661-4. Database: IPA.
Exclude - Not MMIT

N.F. ICT -- shortcut to better health and social services. African Business 2009;(358):46
<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=47000295&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.
Exclude - Not a Primary Study

Naditz A. Medication compliance-helping patients through technology: Modern “smart” pillboxes keep memory-short patients on their medical regimen. Telemedicine and e-Health 2008;14(9):875-80. 20085011773097
Database: Compendex.
Exclude - Not a Primary Study

Nadzam DM, Macklis RM. Promoting patient safety: is technology the solution? Jt Comm J Qual Improv 2001;27(8):430-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Nagle LM, Catford P. Toward a model of successful electronic health record adoption. HEALTHC Q 2008;11(3):84-91. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Naik P, Cuttica C. How to implement smart pump technology in a pediatric hospital setting: The good, the bad and the ugly. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.
Exclude - Not a Primary Study

Najera-Serna CC, Sansom H, Guidry L. Experience with the new McKesson HBOC Robot-Rx Cartless Delivery System. Ashp Midyear Clinical Meeting 1999;34: Database: IPA.
Exclude - Not a Primary Study

Nakajima K, Kurata Y, Takeda H. A web-based incident reporting system and multidisciplinary collaborative projects for patient safety in a Japanese hospital. Quality & Safety in Health Care 2005;14(2):123-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Nanji K, Cina J, Patel N, et al. Overcoming barriers to the implementation of a pharmacy bar code scanning system for medication dispensing: A case study. J Am Med Inform Assoc 2009;16(5):645-50. Exclude - Not MMIT

Naqvi,S. A semi-autonomous on-line chemotherapy prescription system Memorial University of Newfoundland (Canada)Editor. 2007. Grey Lit.
Exclude - Not a Primary Study

Nash JQ, Chandrakumar M, Farrington CP, et al. Feasibility study for identifying adverse events attributable to vaccination by record linkage. Epidemiology & Infection 1995;114(3):475-80. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Nash SB. Computerized physician order entry. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Nassaralla CL, Naessens JM, Hunt VL, et al. Medication reconciliation in ambulatory care: Attempts at improvement. Qual Safe Health Care 2009;18(5):402-7. OVID EMBASE.
Exclude - Not MMIT

Nazir T, Beatty PC. Anaesthetists' attitudes to monitoring instrument design options. *Br J Anaesth* 2000;85(5):781-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Nazzaro J. Looking at a computerized pharmacy in the military sector. *Computers Hosp* 1980;1(Sep-Oct):29-32. Database: IPA.

Exclude - Not MMIT

Nebeker JR, Hurdle JF, Hoffman J, et al. Developing a taxonomy for research in adverse drug events: potholes and signposts. *Proceedings / AMIA 2001*;493-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Negishi E, Hirai A, Yoshizaki N, et al. Electronic links between hospitals or clinics and pharmacies: structure and operation of WAKASHIO pharmacotherapy network. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2003;123(3):191-200. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Neiman S, Krishnan M, Lis A, et al. Improving the safety of using high-alert medications. *Ashp Midyear Clinical Meeting 2003*;38: Database: IPA.

Exclude - Not a Primary Study

Nejad S G, Paranjape R. An agent model of a diabetic patient. In Piscataway, NJ, USA: IEEE; 2007. p.214-8.9701791

Database: Inspec.

Exclude - Not MMIT

Nelson NC, Evans RS, Samore MH, et al. Detection and prevention of medication errors using real-time bedside nurse charting. *J Am Med Inform Assoc* 2005;12(4):390-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Nelson R. Physician direct entry: CPOE will save lives, but staff resistance and high cost are still barriers to acceptance. *Hospital Pharmacist Report* 2001;15(Aug):28 Database: IPA.

Exclude - Not a Primary Study

Nelson SJ, Zeng K, Bodenreider O. MyMedicationList: integrating personal medication records with resources. *AMIA 2008;Annual:Symposium* PMID:18998987 OVID MEDLINE.

Exclude - No Outcomes of Interest

Nemeth J, Krych R, Hiciano T, et al. Controlling clostridium difficile associated disease using a proactive pharmacy plan. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Nemytin I, Petrov VP, Zuev VK, et al. The use of artificial neuronal networks in the treatment of peptic ulcer. *Voen Med Zh* 1996;321(6):40-4. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Nesbitt DH, Whittenburg BD. Optimizing medication use in a cartless distribution system. *Ashp Midyear Clinical Meeting* 1999;34: Database: IPA.

Exclude - Not a Primary Study

Ness JE. Understanding and applying medication bar code technology: Pharmacy challenges. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Neuenschwander M. Review of current automated distribution systems. Ashp Midyear Clinical Meeting 1995;30: Database: IPA.

Exclude - Not a Primary Study

Neuenschwander M. Limiting or increasing opportunities for errors with dispensing automation. Hosp Pharm 1996;31(Sep):1102-6. Database: IPA.

Exclude - Not MMIT

Neuenschwander M. Review of current automated distribution systems. ASHP Annual Meeting 1996;53: Database: IPA.

Exclude - Not a Primary Study

Neuenschwander M, Cohen MR, Vaida AJ, et al. Practical guide to bar coding for patient medication safety. Am J Health Syst Pharm 2003;60(8):768-79. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Neuenschwander M. Rx for the pharmacy: Helps the medicine go down. Healthc Inform 2004;11(Dec):48-50. Database: IPA.

Exclude - Not a Primary Study

Neuenschwander M. Overview and update of automated dispensing technologies for inpatient and outpatient services. Ashp Midyear Clinical Meeting 1997;32: Database: IPA.

Exclude - Not a Primary Study

Neuman SL. Case presentation - New Jersey ePrescribing action coalition. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Newby DA, Robertson J. Computerised prescribing: assessing the impact on prescription repeats and on generic substitution of some commonly used antibiotics. Med J Aust 2010;192(4):192-5. PMID:20170455 OVID MEDLINE.

Exclude - Not MMIT

Newby LK, Harrington RA, Bhaskar MV, et al. An automated strategy for bedside aPTT determination and unfractionated heparin infusion adjustment in acute coronary syndromes: insights from PARAGON A. Journal of Thrombosis & Thrombolysis 2002;14(1):33-42. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Newell LM, Christensen D. Who's counting now? ROI for patient safety IT initiatives. J Healthc Inf Manag 2003;17(4):29-35. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ngalamou L, Campbell H. Diabetic information appliance. In Los Alamitos, CA, USA: IEEE Computer. Soc; 2002. p.3-6.7349234

Database: Inspec.

Exclude - Not MMIT

Nguyen PT, Howrie DL, Crowley KL, et al. Use of computer-generated alerts to identify pediatric patients at risk for nephrotoxicity. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Niazkhani Z, Pirnejad H, de Bont A, et al. Evaluating inter-professional work support by a computerized physician order entry (CPOE) system. *Studies in Health Technology & Informatics* 2008;136:321-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Niazkhani Z. Evaluating the impact of CPOE systems on medical workflow: a mixed method study. *Studies in Health Technology & Informatics* 2008;136:881-2. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Niazkhani Z, Pirnejad H, Berg M, et al. The impact of computerized provider order entry systems on inpatient clinical workflow: a literature review. *J Am Med Inform Assoc* 2009;16(4):539-49. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Niazkhani Z, Pirnejad H, van der SH, et al. Computerized provider order entry system - does it support the inter-professional medication process? Lessons from a Dutch academic hospital. *Methods Inf Med* 2010;49(1):20-7. PMID:19448890 OVID MEDLINE.
Exclude - Unable to Retrieve

Nibali SC. Prescribing with the International Common Denomination in paediatric primary care. A feasibility study. *Quaderni ACP* 2007;14(1): Database: Embase Sept 22-09.
Exclude - Unable to Retrieve Foreign

Nicholas PK, Agius CR. Toward safer IV medication administration: the narrow safety margins of many IV medications make this route particularly dangerous. *Am J Nurs* 2005;105(3 Suppl):25-30. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Nicholson JP, Mullins RA. Toward safer IV medication administration: the narrow safety margins of many IV medications make this route particularly dangerous. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Nicholson JP. Pharmacy involvement in a multi-disciplinary approach to improve medication safety in a community hospital. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Nicol NC. From danger to safety: A complete redesign of a health system medication management model. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Nielsen EW, Dybwik K. Drug interactions in an intensive care unit. *Tidsskr Nor Laegeforen* 2004;124(22):2907-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Nielsen S, Barratt MJ. Prescription drug misuse: is technology friend or foe? *Drug & Alcohol Review* 2009;28(1):81-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Nieuwlaat R, Barker L, Kim Y-K, et al. Underuse of evidence-based warfarin dosing methods for atrial fibrillation patients. *Thromb Res* 2010;125(4):e128-e131 OVID EMBASE.

Exclude - Not MMIT

Nihashi JI, Suzuki Y, Watanabe S, et al. Automatic dispensing checking system using two-dimensional barcode symbols. *Japanese Journal of Hospital Pharmacy* 1995;21(1):79-85.

Database: IPA.

Exclude - Not a Primary Study

Nihashi JI, Nishikawa M, Suzuki Y, et al. Problems and efficiencies of PC order entry system for prescriptions. *Japanese Journal of Hospital Pharmacy* 1997;23(1):75-81.

Database: IPA.

Exclude - Not a Primary Study

Niinimaki J, Forsstrom J. Approaches for certification of electronic prescription software. *Int J Med Inf* 1997;47(3):175-82. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Niinimaki J, Savolainen M, Forsstrom JJ. Methodology for security development of an electronic prescription system. *Proceedings / AMIA* 1998;1998(1998):245-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Nikitin AV. Methodological aspects of the optimal use of combined chemo- and immunotherapy. [Russian]Using telecommunications technology in patient care. *Antibiot Khimioter* 1990;35(10):35-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Nilsen JF. A computerized record system in primary health care. *Tidsskr Nor Laegeforen* 1982;102(25): Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Nix RJ, Hardesty C, Whittington C, et al. Using automation to replace the unit dose distribution system. *ASHP Annual Meeting* 1994;51: Database: IPA.

Exclude - Not a Primary Study

Noffsinger R, Chin S. Improving the delivery of care and reducing healthcare costs with the digitization of information. *J Healthc Inf Manag* 2000;14(2):23-30. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Nold EG, Williams TC. Bar codes and their potential applications in hospital pharmacy. *Am J Hosp Pharm* 1985;42(12):2722-32. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Nold EG. Preparing to implement an information system. *Am J Hosp Pharm* 1993;50(5):958-64. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Nold EG. Managing automation: Practical management skills for new supervisors. Ashp Midyear Clinical Meeting 1998;33: Database: IPA.

Exclude - Not a Primary Study

Nold EG. Health industry bar-code conference. Am J Hosp Pharm 2004;40(Oct):1698 Database: IPA.

Exclude - Not MMIT

Nooker J, Huffman M. Alternatives for decreasing telephone calls for obtaining first doses. Am J Hosp Pharm 1994;51:1576 Database: IPA.

Exclude - Not a Primary Study

North D. Controlling the costs of antibiotic resistance. Clin Ther 1993;15(Suppl A):3-11. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Novak SR, Pegelow ST, Kegley RC, et al. Look before you leap - The impact on pharmacy operations of implementing multiple technologies to decrease medication errors. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Novek J. Clinical or industrial pharmacy? Case studies of hospital pharmacy automation in Canada and France. Int J Health Serv 1998;28(3):445-65. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Novek J. Hospital pharmacy automation: collective mobility or collective control? Soc Sci Med 2000;51(4):491-503. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Nuckolls JG. Process improvement approach to the care of patients with type 2 diabetes. Providing physicians with tools to increase compliance and improve outcomes. Postgrad Med 2003;2003 May(Spec No):53-62. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Nyman K, Bergens A, Bjorin AS, et al. Feedback on prescribing profiles at a primary health center. Important element in quality assurance of drug prescription. Lakartidningen 2001;98(3):160-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

O'Brien M. Implementation of the EPIC electronic medical record/physician order-entry system. J Healthc Manag 2006;51(5):338-43. Database: BSC.

Exclude - No Outcomes of Interest

O'Connell KJ, Lum BL, Coleman RW. Computer-assisted implementation of the Joint Commission on Accreditation of Healthcare Organizations' (JCAHO) medication use indicator-1 (MU-1). Ashp Midyear Clinical Meeting 1995;30: Database: IPA.

Exclude - Not a Primary Study

O'Connor AB, Lang VJ, Quill TE. Eliminating analgesic meperidine use with a supported formulary restriction. Am J Med 2005;118(8):885-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

O'Donnell J. Parallax error is linked to AZT hospital mix-up. Drug Topics 2004;136: Database: IPA.
Exclude - Not a Primary Study

O'Malley C. Using telecommunications technology in patient care. Ashp Midyear Clinical Meeting 2000;35:45 Database: IPA.
Exclude - Not a Primary Study

O'Malley P. Computerized provider order entry and prescribing and the evidence for safe practice: update for the clinical nurse specialist. Clin Nurse Spec 2007;21(3):139-41. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

O'Malley P. Think bar-code medication administration eliminates adverse drug events? Think again! Clin Nurse Spec 2008;22(6):269-70. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

O'Neal BC, Worden JC, Couldry RJ. Telepharmacy and bar-code technology in an i.v. chemotherapy admixture area. Am J Health Syst Pharm 2009;66(13):1211-7. Database: CINAHL.
Exclude - Not a Primary Study

O'Reilly M, Talsma A, VanRiper S, et al. An anesthesia information system designed to provide physician-specific feedback improves timely administration of prophylactic antibiotics. Anesthesia & Analgesia 2006;103(4):908-12. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Oberlender R. Improving patient care by implementing a physician order-entry medication system. ASHP Annual Meeting 1995;52: Database: IPA.
Exclude - Not a Primary Study

Obez C, Barisic AM, Chatelier W, et al. The computer-assisted management programs for antibiotic therapies in connection with an application in geriatrics. Pathol Biol 2004;52(10):589-96. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Obsheatz MT. Bar coding and smart IV pumps - A seamless system for all medications. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.
Exclude - Not a Primary Study

Ocheltree GA, Nelson SP, Fisher WS. Automated dispensing stations: Bar code verification of stock refills. ASHP Annual Meeting 1997;54: Database: IPA.
Exclude - Not a Primary Study

Odwazny R, Hasler S, Abrams R, et al. Organizational and cultural changes for providing safe patient care. Qual Manag Health Care 2005;14(3):132-43. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Ogbuokiri ON, Handler S, Mark SM, et al. Analysis of medication administration errors intercepted by a bar-code medication administration system. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Ogletree N, Frinak S, Thaxton S, et al. Computerized Anemia Management Program (CAMP): An effective and appreciated approach to treating anemia. *Nephrology Nursing Journal* 2008;35(2):178 Database: CINAHL.
Exclude - Not MMIT

Ogura H, Sagara E, Yamamoto K, et al. Analysis of the online order entry process in an integrated hospital information system. *Comput Biol Med* 1985;15(6):381-93. Exclude - No Outcomes of Interest

Ogura H. Time duration in online order entry of prescription by physicians. *Yakuzaigaku* 1985;45(2): Database: Embase Sept 22-09.
Exclude - Unable to Retrieve Foreign

Oh SS. Ordering of continuous renal replacement therapy in a computerized provider order entry system. *Hosp Pharm* 2007;42(3):255-7. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Ohsfeldt RL, Ward MM, Schneider JE, et al. Implementation of hospital computerized physician order entry systems in a rural state: feasibility and financial impact. *J Am Med Inform Assoc* 2005;12(1):20-7. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Ohtsubo Y, Ishimito K, Tanioka M, et al. A checking system for injectable anticancer drugs using each patient's own data and its evaluation. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2002;122(6):389-97. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Oke TO. Primary health-care services with a functional ambulatory care clinical pharmacy in a low-income housing project clinic. *J Natl Med Assoc* 1994;86(6):465-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Oliver WW. Impact of e-medicine. *Georgia Pharmaceutical Journal* 2001;23(May):22-5. Database: IPA.
Exclude - Not MMIT

Olola CH, Rowan B, Narus S, et al. Enhancing continuity of care through an emergency medical card at Intermountain Healthcare: using continuity of Care Record (CCR) standard. *AMIA* 2007;2007 Oct 11(2007):1063 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Ong BK. Leveraging on information technology to enhance patient care: a doctor's perspective of implementation in a Singapore academic hospital. *Ann Acad Med Singapore* 2002;31(6):707-11. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Opasich C, Pinna GD, Sisti M, et al. An analysis of the decision process in the pharmacological treatment of a patient with chronic heart failure by means of a therapy management information system: the experience of the Montescano Heart Failure Unit. [Italian]. *G Ital Cardiol* 1998;28(11):1278-87. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Oppenkowski TP, Murray ET, Sandhar H, et al. External quality assessment for warfarin dosing using computerised decision support software. *J Clin Pathol* 2003;56(8):605-7.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ore R. What do legislators want to know about IT? National Conference of State Legislatures; 2007. http://www.ncsl.org/print/health/forum/Senator_Moore.pdf Grey Lit.

Exclude - Not a Primary Study

Oren E, Griffiths LP, Guglielmo BJ. Characteristics of antimicrobial overrides associated with automated dispensing machines. *Am J Health Syst Pharm* 2002;59(15):1445-8.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Oren E, Shaffer ER, Guglielmo BJ. Impact of emerging technologies on medication errors and adverse drug events. *Am J Health Syst Pharm* 2003;60(14):1447-58. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ornstein SM. Computer-generated physician and patient reminders: Tools to improve population adherence to selected preventive services. *J Fam Pract* 1991;32(1):82-90.

Database: Embase Sept 22-09.

Exclude - Not MMIT

Ortiz-Gomez JR, Monedero-Rodriguez P, Perez-Cajaraville JJ. Applications of informatics in anesthesiology: anesthesia graphics. *Rev Esp Anestesiol Reanim* 2002;49(3):141-9.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Osheroff JA, Pifer E, Teich J, et al. Improving medication use and outcomes with clinical decision support: A step-by-step guide. *Healthcare Information and Management Systems Society*; 2005. Grey Lit.

Exclude - Not a Primary Study

Osheroff JA, Teich JM, Middleton B, et al. A roadmap for national action on clinical decision support. *J Am Med Inform Assoc* 2007;14(2):141-5. Database: IPA.

Exclude - Not a Primary Study

Oszko MA. Determining i.v. infusion rates for critical care drugs using a microcomputer. *Hosp Pharm* 1989;24(6):465-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Otromptke J. New federal rules on e-prescribing and PQRI may provide relief for DPM's. *Podiatry Management* 2009;28(5):69-70. Database: CINAHL.

Exclude - Not MMIT

Ou NN, Armon JJ, Graner KK. Implementation of a computerized system to identify patients with heart failure not receiving reninangiotensin system inhibitor therapy: supporting pharmacist role in adherence to standard of care. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Overall JE, Faillace LA, Rhoades HM, et al. Computer-based monitoring of clinical care in a public psychiatric hospital unit. *Hospital & Community Psychiatry* 1987;38(4):381-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Overhage JM, Tierney WM, McDonald CJ. Design and implementation of the Indianapolis Network for Patient Care and Research. *Bull Med Libr Assoc* 1995;83(1):48-56. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Overhage JM, Middleton B, Miller RA, et al. Does national regulatory mandate of provider order entry portend greater benefit than risk for health care delivery? The 2001 ACMI debate. *The American College of Medical Informatics. J Am Med Inform Assoc* 2002;9(3):199-208. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Owens C, Zisser H, Jovanovic L, et al. Run-to-run control of blood glucose concentrations for people with Type 1 diabetes mellitus. *IEEE Trans Biomed Eng* 2006;53(6):996-1005. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ownby RL. Development of an interactive tailored information application to improve patient medication adherence. *AMIA 2005;2005:1069* Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Ozdas A, Miller RA. Care provider order entry (CPOE): a perspective on factors leading to success or to failure. *Yearbook of Medical Informatics* 2007;2007(1):128-37. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Pachler C, Plank J, Weinhandl H, et al. Tight glycaemic control by an automated algorithm with time-variant sampling in medical ICU patients. *Intensive Care Med* 2008;34(7):1224-30. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Padilla N. Current status of automated dispensing system. *Am J Hosp Pharm* 1994;51:704 Database: IPA.

Exclude - Not a Primary Study

Pain D, Fielden K, Shibl RA. Opinions on the use of clinical decision support systems for paediatric prescribing in a New Zealand hospital. *Logistics Information Management* 2003;16(3):201-6. 7676451

Database: Inspec.

Exclude - No Outcomes of Interest

Pakpahan R, Balas EA, Boren SA. Computable decision modules for patient safety in child health care. *Proceedings / AMIA 2002;2002:592-6*. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Palchuk MB, Seger DL, Alexeyev A, et al. Implementing renal impairment and geriatric decision support in ambulatory e-prescribing. AMIA 2005;2005:1071 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Palchuk MB, Seger DL, Recklet EG, et al. Weight-based pediatric prescribing in ambulatory setting. AMIA 2006;2006:1055 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Palchuk MB, Turchin A, Alexeyev A, et al. Reducing unintended consequences of e-prescribing on the path to nuanced prescriptions. AMIA 2008;2008 Nov 6:1079 PMID:18999103 OVID MEDLINE.

Exclude - No Outcomes of Interest

Palen L, Aalokke S. Of pill boxes and piano benches: "Home-made" methods for managing medication. In 2006; Banff, Alberta, Canada: 2006. p.79-88.Grey Lit.

Exclude - Not a Primary Study

Palmer MT, FitzGerald S. Integrating barcoding into medication dispensing processes in the pharmacy. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.

Exclude - Not a Primary Study

Paltiel O, Gordon L, Berg D, et al. Effect of a computerized alert on the management of hypokalemia in hospitalized patients. Arch Intern Med 2003;163(2):200-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Pandolfi N. Evolution and growth of the department of pharmacy at a university teaching hospital. Ashp Midyear Clinical Meeting 1992;27: Database: IPA.

Exclude - Not a Primary Study

Pankaskie M, Sullivan J. New players, new services: E-scripts revisited. J Am Pharm Assoc (Wash) 2004;40(4):566 Database: IPA.

Exclude - Not MMIT

Pantze R. EDP in the hospital pharmacy: Dose calculation in aminoglycoside therapy. Krankenhauspharmazie 1989;10(Jul):279-84. Database: IPA.

Exclude - Unable to Retrieve

Paoletti VC, Silverstein JH, Beach T, et al. Impact of a prescription action profile on residents' attitudes and perception of time management in a resident medicine clinic. Ashp Midyear Clinical Meeting 1998;33: Database: IPA.

Exclude - Not a Primary Study

Papke JA. Unclaimed prescriptions requisitioned through provider order entry. J Manag Care Pharm 1999;5(Nov-Dec):498-502. Database: IPA.

Exclude - Not MMIT

Papshev D, Peterson AM. Electronic prescribing in ambulatory practice: promises, pitfalls, and potential solutions. Am J Manag Care 2001;7(7):725-36. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Paradiso-Hardy F, Seto A, Ong S, et al. Use of a personal digital assistant in a pharmacy-directed warfarin dosing program. *Am J Health Syst Pharm* 2003;60(19):1943-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Pare G, Jaana M, Sicotte C. Exploring health information technology innovativeness and its antecedents in Canadian hospitals. *Methods Inf Med* 2010;49(1):28-36. PMID:20011805 OVID MEDLINE.

Exclude - Unable to Retrieve

Parente ST, McCullough JS. Health information technology and patient safety: evidence from panel data. *Health Aff (Millwood)* 2009;28(2):357-60.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010221233&site=ehost-live;Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1759&accno=2010221233>
EBSCO CINAHL.

Exclude - No Outcomes of Interest

Park W, Kim J, Chae Y, et al. Does the physician order-entry system increase the revenue of a general hospital? *Int J Med Inf* 2003;71(1):25-32. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Parker PJ. E-scribing: more realistic than ever. *Nurs Manag (Harrow)* 2002;33(4):50
Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Parker WP, Greco M. Computer-generated drug information notices for nurses. *Am J Hosp Pharm* 1985;42(5):1113-4. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Parl FF, O'Leary MF, Kaiser AB, et al. Implementation of a closed-loop reporting system for critical values and clinical communication in compliance with goals of the joint commission. *Clin Chem* 2010;56(3):417-23. OVID EMBASE.

Exclude - Not MMIT

Parnes JG, Barnes PB, Grice JM. Automation and information: Changing face of pharmacy. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Parrish F, Do N, Bouhaddou O and others. Implementation of RxNorm as a terminology mediation standard for exchanging pharmacy medication between federal agencies. In Washington, DC: 2006. p. 1057. Grey Lit.

Exclude - Not a Primary Study

Pasquarella MV, Kile DM, Young JM, et al. Patient focused pharmacy services - Implementation of a decentralized unit-based pharmacist program. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Patel SP, Eisenwine JE, Kromelis MR. Use of carousel technology to decrease medication errors and increase efficiency in an outpatient ambulatory pharmacy in a private, not-for-profit, university-affiliated pediatric hospital. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Patel UC, Davis L, Snead J. Implementation of surgery antimicrobial prophylaxis templates at a Veterans Affairs (VA) hospital. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Paterno MD, Cina JL, Goldhaber SZ, et al. Preventing DVT and PE in hospitalized patients: improving a successful electronic alert. *AMIA* 2006;2006:1058 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Patil SG, Gale TJ. Preliminary design of remotely used and monitored medication dispenser. *Conference Proceedings*: 2006;1:3616-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Patterson CB, Gorrell JJ, Spence CE, et al. Reconciling medications accurately and completely across the continuum. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Patterson ES. Addressing human factors in bar code medication administration systems. *Hosp Pharm* 2003;38(11):S16-S17 Database: IPA.

Exclude - No Outcomes of Interest

Patterson ES, Rogers ML, Render ML. Fifteen best practice recommendations for bar-code medication administration in the Veterans Health Administration. *Joint Commission Journal on Quality & Safety* 2004;30(7):355-65. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Patterson G, Gaspar C. Automated medication dispensing system impact on drug distribution and pharmacy expenditures. *Ashp Midyear Clinical Meeting* 1989;24: Database: IPA.

Exclude - Not a Primary Study

Paul M, Billebaud T, Kinoo J, et al. Use of informatics to validate the prescription of large spectrum antibiotics. *Journal de Pharmacie Clinique* 1988;7(Suppl 2):63-83. Database: IPA.

Exclude - Unable to Retrieve Foreign

Paul M, Nielsen AD, Goldberg E, et al. Prediction of specific pathogens in patients with sepsis: evaluation of TREAT, a computerized decision support system. *J Antimicrob Chemother* 2007;59(6):1204-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Payne TH. The transition to automated practitioner order entry in a teaching hospital: the VA Puget Sound experience. *Proceedings / AMIA* 1999;Annual Symposium.:589-93. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Payne TH. Commercial knowledge bases versus custom rule sets. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Payne TH. Order sets and protocols. Ashp Summer Meeting 2009;65: Database: IPA.
Exclude - Not a Primary Study

Payton J, Leder W, Hord E. Bar Code Medication Administration system improves patient safety. J Ark Med Soc 2007;104(4):84-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Pazzani MJ, See D, Schroeder E, et al. Application of an expert system in the management of HIV-infected patients. Journal of Acquired Immune Deficiency Syndromes & Human Retrovirology 1997;15(5):356-62. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Pearson LS. Use of bar coding technology. Med Device Technol 1994;5(3):42-3. Database: IPA.
Exclude - Not a Primary Study

Pearson SA, Soumerai S, Mah C, et al. Racial disparities in access after regulatory surveillance of benzodiazepines. Arch Intern Med 2006;166(5):572-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Pearson SA, Moxey A, Robertson J, et al. Do computerised clinical decision support systems for prescribing change practice? A systematic review of the literature (1990-2007). BMC Health Serv Res 2009;9:154 PMID:19715591 OVID MEDLINE.
Exclude - Not a Primary Study

Pearson WS, Bercovitz AR. Use of computerized medical records in home health and hospice agencies: United States, 2000. Vital & Health Statistics - Series 13: Data From the National Health Survey 2006;(161):1-14. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Peck C, Sheiner L, Halkin H and others. Practical application of computer-aided drug therapy. In 1974. p.321-3.1975120003468
Database: Compendex.
Exclude - No Outcomes of Interest

Pedersen CA, Schneider PJ, Santell JP. ASHP national survey of pharmacy practice in hospital settings: prescribing and transcribing--2001. Am J Health Syst Pharm 2001;58(23):2251-66. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Pedersen CA. ASHP national survey of pharmacy practice in acute care settings. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.
Exclude - Not a Primary Study

Pedersen CA, Schneider PJ, Scheckelhoff DJ. ASHP national survey of pharmacy practice in hospital settings: dispensing and administration--2002. Am J Health Syst Pharm 2003;60(1):52-68. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Pedersen CA, Schneider PJ, Scheckelhoff DJ. ASHP national survey of pharmacy practice in hospital settings: prescribing and transcribing--2004. *Am J Health Syst Pharm* 2005;62(4):378-90. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Pedersen CA, Schneider PJ, Scheckelhoff DJ. ASHP national survey of pharmacy practice in hospital settings: prescribing and transcribing--2007. *Am J Health Syst Pharm* 2008;65(9):827-43. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Pedersen CA, Schneider P, Scheckelhoff D. ASHP national survey of pharmacy practice in hospital settings: Dispensing and administration - 2008. *Am J Health Syst Pharm* 2009;66(10):926-46. Database: Embase Sept 22-09.

Exclude - Not MMIT

Pedersen CA, Schneider PJ, Scheckelhoff DJ. ASHP national survey of pharmacy practice in hospital settings: dispensing and administration--2008. *Am J Health Syst Pharm* 2009;66(10):926-46. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Peek N, Goud R, Abu-Hanna A. Application of statistical process control methods to monitor guideline adherence: a case study. *AMIA Annual Symposium Proceedings* 2008;581-5.

OVID EMBASE.

Exclude - Not MMIT

Peiris D, Murray J, Scully D, et al. Cardiovascular risk management at a Maori-led Primary Health Organisation--findings from a cross-sectional audit. *N Z Med J* 2008;121(1285):35-46. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Peiris DP, Joshi R, Webster RJ, et al. An electronic clinical decision support tool to assist primary care providers in cardiovascular disease risk management: development and mixed methods evaluation. *J Med Internet Res* 2009;11(4):e51 PMID:20018588 OVID MEDLINE.

Exclude - Not MMIT

Peithner G, Tschernuter U. Scanning in homeopathy. *Oaz Oesterreichische Apotheker Zeitung* 2003;57(8):376-8. Database: IPA.

Exclude - Not MMIT

Peitzman L. Automate to achieve meaningful use. *Health Manag Technol* 2009;30(11):25 <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010470331&site=ehost-live>; EBSCO CINAHL.

Exclude - Not a Primary Study

Pelayo S, Leroy N, Guerlinger S, et al. Cognitive analysis of physicians' medication ordering activity. *Studies in Health Technology & Informatics* 2005;116:929-34. Grey Lit.

Exclude - No Outcomes of Interest

Pelayo S, Bernonville S, Kolski C, et al. Applying a Human Factors Engineering approach to healthcare IT applications: example of a medication CPOE project. *Studies in Health Technology & Informatics* 2009;143:334-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Peleska J, Svejda D, Zvarova J. Computer supported decision making in therapy of arterial hypertension. *Int J Med Inf* 1997;45(1-2):25-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Peleska J, Zvara K, Vesely A, et al. Development of electronic form of the 1999 WHO/ISH hypertension guidelines. *Studies in Health Technology & Informatics* 2002;90:268-71.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Pellowe C. New catheterisation e-learning unit available to NHS staff. *Nurs Times* 2009;105(7):36

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010439028&site=ehost-live> EBSCO CINAHL.

Exclude - Not a Primary Study

Pelot J. Safety net's net architect. Interview by Elizabeth Gardner. *Health Data Manag* 2010;18(5):64 PMID:20464903 OVID MEDLINE.

Exclude - Not a Primary Study

Penfield S, Anderson K, Edmunds M et al. Towards health information liquidity: Realization of better, more efficient care from the free flow of health information. Booz, Allen, Hamilton; 2009.

http://www.boozallen.com/media/file/Toward_Health_Information_Liquidity.pdf Grey Lit.

Exclude - Not a Primary Study

Peng CC, Glassman PA, Castiglione B, et al. Retrospective drug utilization review: Incidence of clinically relevant potential drug-drug interactions in a large ambulatory population. *J Manag Care Pharm* 2003;9(6):513-22. Database: IPA.

Exclude - Not MMIT

Pennebaker G. Hospital patients get the green light. *Computertalk for the Pharmacist* 1987;7(May-Jun):8 Database: IPA.

Exclude - Not a Primary Study

Peoples MJ. Scanning makes more sense than ever. *Computertalk for the Pharmacist* 1998;18(Sep-Oct):34-5. Database: IPA.

Exclude - Not a Primary Study

Pepper GA, Towsley GL. Medication errors in nursing homes: Incidence and reduction strategies. *Journal of Pharmaceutical Finance, Economics & Policy* 2007;16(1):5-59.

Database: BSC.

Exclude - Not MMIT

Perea Falomir M, Bernaus Tosquella S. Online pharmaceutical care for the institutionalized elderly patient: Early experiences. [Spanish]. *Atencion Farmaceutica* 2010;12(1):50-4. OVID EMBASE.

Exclude - Unable to Retrieve

Pereira CC, Gomes FV, Cornelio RCAC, et al. Description and evaluation of the medication system of pharmacy service at an University hospital. *Acta Farmaceutica Bonaerense* 2009;28(1):91-6. Database: IPA.

Exclude - Unable to Retrieve Foreign

Perini VJ, Vermeulen LC, Jr. Comparison of automated medication-management systems. *Am J Hosp Pharm* 1994;51(15):1883-91. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Perkerson KA, Quercia RA, Goldman M, et al. A multidisciplinary initiative to increase identification and reporting of adverse drug reactions. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Perras C, Jacobs P, Boucher M et al. Technologies to reduce errors in dispensing and administration of medication in hospitals: Clinical and economic analyses. *Technology Report Number 121*. Ottawa: Canadian Agency for Drugs and Technologies in Health; 2009.
Exclude - Not a Primary Study

Perraut R. Micro-informatics and administration. A case of practical application in an overseas hospital. *Medicine et Armees* 1984;12(7):703-7. Database: Embase Sept 22-09.
Exclude - Not MMIT

Perrecone M. Electronic medication administration record system in an acute care hemodialysis unit. *Nephrology Nursing Journal* 2008;35(2):178 Database: CINAHL.
Exclude - Not a Primary Study

Perrecone M. Venous thromboembolism (VTE) prophylaxis: A new look at an old problem. *Crit Care Nurse* 2009;28(2):e50-e51 Database: CINAHL.
Exclude - Not MMIT

Perrin FV. Improving communication with your patients. *Drug Topics* 1988;May(2):48-56. Database: IPA.
Exclude - Not a Primary Study

Perrin FV. POS systems finally checking in. *Drug Topics* 1988;132:42-3. Database: IPA.
Exclude - Not a Primary Study

Perrin RA, Simpson N. RFID and bar codes--critical importance in enhancing safe patient care. *J Healthc Inf Manag* 2004;18(4):33-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Perrone E, Padula MS, Scarpa M, et al. Computerized medical records in monitoring hypertension. Longitudinal and horizontal evaluation of 25 general practitioners in a primary care setting of Modena, Italy. *Recenti Prog Med* 2009;100(1):4-8. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Perry L. Vendor launches bar-code suite to reduce med errors. *Drug Topics* 2002;146(4):B32 Database: IPA.
Exclude - Not a Primary Study

Perry L. Automated dispensing machines and robots are taking safety to new heights. *Drug Topics* 2008;152(8):32-3. Database: IPA.
Exclude - Not a Primary Study

Persell SD, Denecke-Dattalo TA, Dunham DP, et al. Patient-directed intervention versus clinician reminders alone to improve aspirin use in diabetes: a cluster randomized trial. *Jt Comm J Qual Patient Saf* 2008;34(2):98-105. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Persson M, Bohlin J, Eklund P. Development and maintenance of guideline-based decision support for pharmacological treatment of hypertension. *Computer Methods & Programs in Biomedicine* 2000;61(3):209-19. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Persson M, Mjorndal T, Carlberg B, et al. Evaluation of a computer-based decision support system for treatment of hypertension with drugs: retrospective, nonintervention testing of cost and guideline adherence. *J Intern Med* 2000;247(1):87-93. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Pestotnik SL. Medical informatics, decision support, and quality of care: Clinician's perspective. *ASHP Annual Meeting* 1998;55: Database: IPA.

Exclude - Not a Primary Study

Pestotnik SL. Preventing medication errors with health care technology. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Pestotnik SL. Expert clinical decision support systems to enhance antimicrobial stewardship programs: insights from the society of infectious diseases pharmacists. *Pharmacotherapy* 2005;25(8):1116-25. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Peters A, Kerner W. Analytic design and clinical application of an intelligent control system for pharmacotherapy with insulin--1 [German]. *Biomed Tech (Berl)* 1996;41(1-2):2-13. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Peters A, Kerner W. Analytic design and clinical application of an intelligent control system for pharmacotherapy with insulin--2. *Biomedizinische Technik (Berl)* 1996;41(3):42-53. <http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/822/CN-00124822/frame.html> Database: Cochrane.

Exclude - Unable to Retrieve Foreign

Petersen C. High tech toys offer pharmacy benefit management opportunity. *Managed Healthcare* 2000;10(3):39-40. Database: IPA.

Exclude - Not a Primary Study

Petersen D, Kato S. Computer program for drug dose tapering schedules: Cyclosporine and dexamethasone. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Petersen KG, Khalaf A, Jagle P, et al. A new software for initiating and optimising insulin treatment of out-patients. *Computer Methods & Programs in Biomedicine* 1990;32(3-4):325-31. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Peterson GM. Computer assisted drug therapy for epilepsy. *Drug Intelligence & Clinical Pharmacy* 1983;17(2):123-5. Database: IPA.

Exclude - Not a Primary Study

Peterson J F, Bates D W, Patel M and others. Gerios: Recommending drugs and dosing for elderly patients. In Bethesda, MD, USA: American Medical Informatics Assoc; 2002. p. 1127.8037214

Database: Inspec.

Exclude - No Outcomes of Interest

Peterson M, Byrom B, Dowlman N, et al. Optimizing clinical trial supply requirements: simulation of computer-controlled supply chain management. *Clinical Trials* 2004;1(4):399-412. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Petitta A, Miller DA, Zarowitz BJ, et al. A barcode-based clinical data capture system for the evaluation of drug usage. *ASHP Annual Meeting* 1990;47: Database: IPA.

Exclude - Not a Primary Study

Petropoulos CH, Petropoulos JB, Schoman-Finck LI. Impact of an interdisciplinary team on bar code medication administration. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Pette M, Eulitz M. The Multiple Sclerosis Documentation System MSDS. Discussion of a documentation standard for multiple sclerosis. *Nervenarzt* 2002;73(2):144-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Petersson M, Wihlborg J, Lovstrom R, et al. Systematizing medical alerts. *Studies in Health Technology & Informatics* 2008;136:753-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Pevnick JM, Asch SM, Adams JL, et al. Adoption and use of stand-alone electronic prescribing in a health plan-sponsored initiative. *Am J Manag Care* 2010;16(3):182-9.

PMID:20225913 OVID MEDLINE.

Exclude - Not MMIT

Pezzo MV, Pezzo SP. Physician evaluation after medical errors: Does having a computer decision aid help or hurt in hindsight? *Med Decis Making* 2006;26(1):48-56. Database:

Embase Sept 22-09.

Exclude - Not MMIT

Pham PA. Drug-drug interaction programs in clinical practice. *Clinical Pharmacology & Therapeutics* 2008;83(3):498-500. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Phillips IE, Nelsen C, Peterson J, et al. Improving aminoglycoside dosing through computerized clinical decision support and pharmacy therapeutic monitoring systems. *AMIA Annual Symposium Proceedings* 2008;1093 PMID:18999099 OVID MEDLINE.

Exclude - No Outcomes of Interest

Phillips JO, Massoud N. Case study II: Use of technological tools to optimize drug therapy. Ashp Midyear Clinical Meeting 1992;27: Database: IPA.
Exclude - Not a Primary Study

Phillips MT, Berner ES. Beating the system--pitfalls of bar code medication administration. J Healthc Inf Manag 2004;18(4):16-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Phillips W, Fleming D. Ethical concerns in the use of electronic medical records. Mo Med 2009;106(5):328-33. PMID:19902711 OVID MEDLINE.
Exclude - Not a Primary Study

Piacenti F. Development of an antiretroviral medication monitoring system for patients admitted with HIV: 2 Years later. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.
Exclude - Not a Primary Study

Piazza G, Goldhaber SZ. Computerized decision support for the cardiovascular clinician: applications for venous thromboembolism prevention and beyond. Circulation 2009;120(12):1133-7. PMID:19770412 OVID MEDLINE.
Exclude - No Outcomes of Interest

Pinckard AS. Employing software as a tool to alert pharmacists to patients requiring intervention. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.
Exclude - Not a Primary Study

Pires AR, Thigpen AR, Donnelly TM, et al. Use of an emergency medicine proprietary software package to optimize pharmaceutical care. ASHP Annual Meeting 1995;52: Database: IPA.
Exclude - Not a Primary Study

Pitre M, Ong K, Huh JH, et al. Thorough planning and full participation by pharmacists is key to MOE/MAR success. HEALTHC Q 2004;10(Sp):43-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Pizziferri L, Kittler AF, Volk LA, et al. Impact of an Electronic Health Record on oncologists' clinic time. AMIA Annual Symposium Proceedings 2005;1083 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Plagenz L. Impact of automation. Hospitals 1971;45:87-9. Database: IPA.
Exclude - Not MMIT

Plougmann S, Hejlesen OK, Cavan DA. DiasNet--a diabetes advisory system for communication and education via the internet. Int J Med Inf 2001;64(2-3):319-30. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Podichetty V, Penn D. The progressive roles of electronic medicine: benefits, concerns, and costs. Am J Med Sci 2004;328(2):94-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Poggio F. It's all about workflow. A misplaced focus on automating transactions will never bring about the healthcare system everyone seeks. *Healthc Inform* 2009;26(9):46-8. PMID:19813575 OVID MEDLINE.

Exclude - Not a Primary Study

Poggio F. Experts' corner. CPOE and the doc dilemma. *Healthc Inform* 2010;27(3):46-8. <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010596043&site=ehost-live>; EBSCO CINAHL.

Exclude - Not a Primary Study

Poikonen J. Quality of care, legal liability and pharmacy computer systems. *California Journal of Hospital Pharmacy* 1991;3(Jul):6-7. Database: IPA.

Exclude - Not a Primary Study

Poikonen J. A new term for transcribing. *Am J Health Syst Pharm* 2008;65(19):1801-2. <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010062417&site=ehost-live>; EBSCO CINAHL.

Exclude - Not a Primary Study

Poirier MF, Garreau M, Deniker P. Toward an automatic classification of psychotropic drugs, based on the computerized card-index of their characteristics. *Ann Med Psychol (Paris)* 1982;140(1):1-26. Database: PsycINFO.

Exclude - Not MMIT

Poirier TI. Drug interaction microcomputer software evaluation: SOAP. *Hosp Pharm* 1990;25(4):342-7. Database: Embase Sept 22-09.

Exclude - Not MMIT

Poirier TI, Giudici RA. Drug interaction microcomputer software evaluation: Rx triage. *Hosp Pharm* 1990;25(2):136-40. Database: Embase Sept 22-09.

Exclude - Not MMIT

Poirier TI, Giudici R. Evaluation of drug interaction microcomputer software: an updated comparison. *Hosp Pharm* 1995;30(10):888-94. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Poissant L, Pereira J, Tamblyn R, et al. The impact of electronic health records on time efficiency of physicians and nurses: A systematic review. *J Am Med Inform Assoc* 2005;12(5):505-16. Database: CINAHL.

Exclude - Not a Primary Study

Poller L, Wright D, Rowlands M. Prospective comparative study of computer programs used for management of warfarin. *J Clin Pathol* 1993;46(4):299-303. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Poller L, Shiach CR, MacCallum PK, et al. Multicentre randomised study of computerised anticoagulant dosage. European Concerted Action on Anticoagulation. *Lancet* 1998;352(9139):1505-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Poller L, Keown M, Ibrahim S, et al. A multicentre randomised clinical endpoint study of PARMA 5 computer-assisted oral anticoagulant dosage. *Br J Haematol* 2008;143(2):274-83. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Poller L, Keown M, Ibrahim S, et al. An international multicenter randomized study of computer-assisted oral anticoagulant dosage vs. medical staff dosage. *Journal of Thrombosis & Haemostasis* 2008;6(6):935-43. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Polydorou S, Gunderson EW, Levin FR. Training physicians to treat substance use disorders. *Current Psychiatry Reports* 2008;10(5):399-404. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Pons-Busom M, Aguas-Compaired M, Delas J, et al. Compliance with local guidelines for antibiotic prophylaxis in surgery. *Infect Control Hosp Epidemiol* 2004;25(4):308-12.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ponte CD. Managed care and the pharmacy profession revisited. *J Manag Care Pharm* 1999;5(2):78 Database: IPA.

Exclude - Not a Primary Study

Poon E. The use of bar code technology to improve medication safety: Reviewing the evidence. *Hosp Pharm* 2003;38(11):S5-S6 Database: IPA.

Exclude - Not a Primary Study

Poon EG, Cina JL, Churchill WW, et al. Effect of bar-code technology on the incidence of medication dispensing errors and potential adverse drug events in a hospital pharmacy.

AMIA Annual Symposium Proceedings 2005;1085 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Poon EG, Blumenfeld B, Hamann C, et al. Design and implementation of an application and associated services to support interdisciplinary medication reconciliation efforts at an integrated healthcare delivery network. *J Am Med Inform Assoc* 2006;13(6):581-92.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Poon EG, Cina JL, Churchill W, et al. Medication dispensing errors and potential adverse drug events before and after implementing bar code technology in the pharmacy. *Ann Intern Med* 2006;145(6):426-34. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Poon EG, Wright A, Simon SR, et al. Relationship between use of electronic health record features and health care quality: results of a statewide survey. *Med Care* 2010;48(3):203-9.

PMID:20125047 OVID MEDLINE.

Exclude - Not MMIT

Pope RA, Mattson CJ, Janousek JP, et al. A computer-based IV admixture system. *Methods Inf Med* 1982;21(2):65-9. 1895932

Database: Inspec.

Exclude - Not MMIT

Popescu M, Arthur G. OntoQuest: a physician decision support system based on ontological queries of the hospital database. AMIA Annual Symposium Proceedings 2006;639-43. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Popkin J, Kushniruk A, Borycki E, et al. The eFOSTr PROJECT: design, implementation and evaluation of a web-based Personal Health Record to support health professionals and families of children undergoing transplants. Studies in Health Technology & Informatics 2009;143:358-63. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Popovic T, Dojic L, Radonjic S, et al. System for the control of drug consumption of medical prescriptions of the self-interest health community of the city of Novi Sad and problems affecting this work and the results. Farmaceutski Glasnik 1990;46(4):138-42. Database: IPA.

Exclude - Not MMIT

Porter B. [Computerization in a community health service provider]. [Hebrew]. Harefuah 1999;137(7-8):274-7-351. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Porter BJ. Revising respiratory medication procedures to comply with regulatory standards and improve patient care. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Porter SC, Cai Z, Gribbons W, et al. The asthma kiosk: a patient-centered technology for collaborative decision support in the emergency department. J Am Med Inform Assoc 2004;11(6):458-67. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Porter SC, Kohane IS, Goldmann DA. Parents as partners in obtaining the medication history. J Am Med Inform Assoc 2005;12(3):299-305. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Porter SC, Forbes P, Feldman HA, et al. Impact of patient-centered decision support on quality of asthma care in the emergency department. Pediatrics 2006;117(1):e33-e42 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Porter SC, Kaushal R, Forbes PW, et al. Impact of a patient-centered technology on medication errors during pediatric emergency care. Ambulatory Pediatrics 2008;8(5):329-35. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Portwood JL, Bourret JA. Implementation of a day sheet bar-code system for controlled substances. Ashp Midyear Clinical Meeting 1991;26: Database: IPA.

Exclude - Not a Primary Study

Pour G, Bashyam P, Subramani J. MobiHealth: mobile agent-based system for mobile telemedicine. In Las Vegas, NV, USA: CSREA Press; 2007. p.459-64.10677483

Database: Inspec.

Exclude - No Outcomes of Interest

Pourrat X, Antier D, Crenn I, et al. A prescription and administration error of cisplatin: a case report. *Pharmacy World & Science* 2004;26(2):64-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Pous Serrano S, Dolz Lago JF, Estevan Estevan R, et al. Computerized physician order entry of pharmacotherapeutic protocols in a colorectal surgery unit. *Cirugia Espanola* 2006;80(4):195-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Powell V. Computerized physician outpatient prescription order entry: Department of Defense's version. *Hosp Pharm* 1997;32(1):65-74. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Powner D. Information technology: Centers for medicare & medicaid services needs to establish critical investment management capabilities. Report to the chairman, committee on finance, U.S. senate. 2005. <http://nyam.waldo.kohalibrary.com/cgi-bin/koha/opac-detail.pl?biblionumber=226627> saved as NYAM powner.pdf

Grey Lit.
Exclude - Not MMIT

Preece JF, Ashford JR, Hunt RG. Writing all prescriptions by computer. *J R Coll Gen Pract* 1984;34(269):655-7. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Prentice JP, Comer T. Evaluation of the UAB hospital automated medication management system: Medication availability for now and stat orders. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.
Exclude - Not a Primary Study

Previte JP. Information and communication system implementation in anesthesia. *Int Anesthesiol Clin* 2006;44(1):179-97. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Prgomet M, Georgiou A, Westbrook JI. The impact of mobile handheld technology on hospital physicians' work practices and patient care: a systematic review. [Review] [60 refs]. *J Am Med Inform Assoc* 2009;16(6):792-801. PMID:19717793 OVID MEDLINE.
Exclude - Not MMIT

Prier B, Siegel J, Hafford A. Automating pharmacy personnel: Using radio frequency and bar code technology to facilitate medication distribution. *Ashp Summer Meeting* 2007;64: Database: IPA.
Exclude - Not a Primary Study

Prieston ET, Rodgers PE. Kinetic dosing service enhanced by pharmacist order entry of drug level orders. *Ashp Midyear Clinical Meeting* 1994;29: Database: IPA.
Exclude - Not a Primary Study

Prieston ET, Moyer JL, Grant DK. Implementation of physician computerized order entry in a hospital based outpatient medical oncology clinic. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Prokosch H U, Hulse R K, Wall M and others. New decision support concepts for the pharmacy application in the HELP system. In Berlin, West Germany: Springer-Verlag; 1988. p.31-7.3495797

Database: Inspec.

Exclude - No Outcomes of Interest

Pronovost P, Hobson DB, Earsing K, et al. A practical tool to reduce medication errors during patient transfer from an intensive care unit. JCOM 2004;11(1):26-33. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Prot S, Bussieres JF, Lebel D. Role of computer aided prescriptions in health care facilities. Quebec Pharmacie 2004;51(8):651-5. Database: IPA.

Exclude - Unable to Retrieve Foreign

Protti D. Comparison of information technology in general practice in 10 countries. HEALTHC Q 2007;10(2):107-16. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Protti D, Bowden T, Johansen I. Adoption of information technology in primary care physician offices in New Zealand and Denmark, Part 4: Benefits comparisons. Inform Prim Care 2008;16(4):291-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Protti D, Bowden T, Johansen I. Adoption of information technology in primary care physician offices in New Zealand and Denmark, part 5: final comparisons. Inform Prim Care 2009;17(1):17-22. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Protti D, Johansen I, Perez-Torres F. Comparing the application of Health Information Technology in primary care in Denmark and Andalusia, Spain. Int J Med Inf 2009;78(4):270-83. Database: Embase Sept 22-09.

Exclude - Not MMIT

Puglisi RC. Computerization of the Veterans Administration pharmacy service. Mil Med 1981;146(2):124-6. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Puto K, Sohan U, Lopez-Samblas AM. Implementation of a potential and actual medication error reporting system. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Puustjarvi J, Puustjarvi L. The role of medicinal ontologies in querying and exchanging pharmaceutical information. International Journal of Electronic Healthcare 2009;5(1):1-13. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Qian Y, Ye X, Du W, et al. A computerized system for detecting signals due to drug-drug interactions in spontaneous reporting systems. Br J Clin Pharmacol 2010;69(1):67-73.

PMID:20078614 OVID MEDLINE.

Exclude - Not MMIT

Quan S, Zwarenstein MF, Dainty KN, et al. Legislative barriers to outpatient e-prescribing in a randomized trial. *HEALTHC Q* 2007;10(4):106-12. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Qureshi F. Minimising computer script errors. *Aust Fam Physician* 2001;30(4):350 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Rabert AS, Sebastian MM. The future is now: implementation of a tele-intensivist program. *J Nurs Adm* 2006;36(1):49-54. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Rabiller N. Automatic lists for dispensing of drugs and materials in functional units. *Pharmacie Hospitaliere Francaise* 1985;73:489-92. Database: Embase Sept 22-09.
Exclude - Not MMIT

Rabinowitz I, Tamir A. The SaM (Screening and Monitoring) approach to cardiovascular risk-reduction in primary care--cyclic monitoring and individual treatment of patients at cardiovascular risk using the electronic medical record. *European Journal of Cardiovascular Prevention & Rehabilitation* 2005;12(1):56-62. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Rabol LI, Anhoj J, Pedersen A, et al. Clinical decision support: Is the number of medication errors reduced? *Ugeskr Laeger* 2006;168(48):4179-84. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Raebel MA, Chester EA, Newsom EE, et al. Randomized trial to improve laboratory safety monitoring of ongoing drug therapy in ambulatory patients. *Pharmacotherapy* 2006;26(5):619-26. Exclude - Not MMIT

Rahimi B, Moberg A, Timpka T, et al. Implementing an integrated computerized patient record system: Towards an evidence-based information system implementation practice in healthcare. *AMIA Annual Symposium* 2008;616-20. PMID:18999062 OVID MEDLINE.
Exclude - Not MMIT

Rahimi B, Vimarlund V, Timpka T. Health information system implementation: A qualitative meta-analysis. *J Med Syst* 2009;33(5):359-68. OVID EMBASE.
Exclude - Not a Primary Study

Rahman Y, Knape T, Gargan M, et al. e-clinic: an electronic triage system for the management of type 2 Diabetes Mellitus. *Studies in Health Technology & Informatics* 2004;107(Pt:1):1-50. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Raible C. Unit-dose-system. *Krankenhauspharmazie* 2008;29(6): Database: Embase Sept 22-09.
Exclude - Not MMIT

Raine T. The case for coding. *Packaging News* 2006;39-41. 20070110344563 Database: Compendex.
Exclude - Not MMIT

Ralston JD, Carrell D, Reid R, et al. Patient web services integrated with a shared medical record: patient use and satisfaction. *J Am Med Inform Assoc* 2007;14(6):798-806. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ram D, Peretz B. Administering local anaesthesia to paediatric dental patients -- current status and prospects for the future. *Int J Paediatr Dent* 2002;12(2):80-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ramos Gonzalez, J. A design and implementation of an electronic physician orders entry system University of Puerto Rico, Mayaguez (Puerto Rico) Editor. 2006. Grey Lit.

Exclude - No Outcomes of Interest

Ramsay J, Hans-Joachim P, Thull B, et al. The evaluation of an information system for intensive care. *Behaviour & Information Technology* 1997;16(1):17-24. Database: PsycINFO.

Exclude - Not MMIT

Rapp SM. Eight-point framework designed to yield safer health care amidst greater EHR use. *Orthopedics Today* 2009;29(11):24

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010475193&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1950&accno=2010475193

EBSCO CINAHL.

Exclude - Not a Primary Study

Rask K, Culler S, Scott T, et al. Adopting National Quality Forum medication safe practices: Progress and barriers to hospital implementation. *Journal of Hospital Medicine (Online)* 2007;2(4):212-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rastegar DA, Knight AM, Monolakis JS. Antiretroviral medication errors among hospitalized patients with HIV infection. *Clin Infect Dis* 2006;43(7):933-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rastoul S, Portier E, Bonan B, et al. Quality assurance in a satellite pharmacy in a pediatric oncology and hematology department. *Journal de Pharmacie Clinique* 1996;15(3):202-15.

Database: IPA.

Exclude - Unable to Retrieve Foreign

Raths D. Is the bar still too high? *Healthc Inform* 2009;26(9):38

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010429777&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=1239&accno=2010429777

EBSCO CINAHL.

Exclude - Not a Primary Study

Ravnan MC, Ravnan SL, Manzo BA. Physician attitudes toward computerized medication order entry. *Ashp Midyear Clinical Meeting* 2001;36. Database: IPA.

Exclude - Not a Primary Study

Ray MD, Aldrich LT, Lew PJ. Experience with an automated point-of-use unit-dose drug distribution system. *Hosp Pharm* 1995;30(1):18-3. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Raybardhan S, Balen RM, Partovi N, et al. Documenting drug-related problems with personal digital assistants in a multisite health system. *Am J Health Syst Pharm* 2005;62(17):1782-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Raymoure SH, Kunz K, Hall K, et al. Effect of a pharmacy generated medication administration record on medication errors. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Reading LM. Implementation of a computerized prescriber order entry system: A work in progress at a cardiac/pulmonary center. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Real ML, Banahan BF, III, Brown TR, et al. Computerized services in hospital pharmacy departments. *Am J Hosp Pharm* 1993;50(1):121-3. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Reardon G, Pleil AM, Zaenger AF. Evaluation of a rule-based program to describe antibiotic utilization for otitis media among three medical plans. *J Manag Care Pharm* 1999;5:232-40. Database: IPA.
Exclude - Not MMIT

Rebidas D, Smith ST, Denomme P. Redesigning medication distribution systems in the OR. *AORN J* 1999;69(1):184-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Reckmann MH, Westbrook JI, Koh Y, et al. Does computerized provider order entry reduce prescribing errors for hospital inpatients? A systematic review. *J Am Med Inform Assoc* 2009;16(5):613-23. PMID:19567798 OVID MEDLINE.
Exclude - Not a Primary Study

Reed DM, Hepler C, Helling D. Antibiotic use review in ambulatory care using computer assisted medical record audit. *Am J Hosp Pharm* 1982;39(2):280-4. Database: Embase Sept 22-09.
Exclude - Not MMIT

Reed MC, Grossman JM. Limited information technology for patient care in physician offices. *Issue Brief/Center for Studying Health System Change* 2004;(89):1-6. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Reed MJ, Fothergill J. The Livingston Paediatric Dose Calculator. *Emergency Medicine Journal* 2007;24(8):567-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Reese R. EMRs help to boost productivity. 2002.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=63036> Grey Lit.
 Exclude - Not a Primary Study

Refsum C, Kumarapeli P, Gunaratne A, et al. Measuring the impact of different brands of computer systems on the clinical consultation: a pilot study. *Inform Prim Care* 2008;16(2):119-27. Database: Ovid MEDLINE(R).
 Exclude - Not MMIT

Regan D and Stevens C. Improved outpatient medication management using online physician prescribing. Healthcare Information and Management Systems Society; 2002.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=62722> Grey Lit.
 Exclude - Not a Primary Study

Reich PR. Computerized prescription writing revisited. *Manag Care Interface* 2004;14(Jan):12 Database: IPA.
 Exclude - No Outcomes of Interest

Reichert JC, Glasgow M, Narus SP, et al. Using LOINC to link an EMR to the pertinent paragraph in a structured reference knowledge base. *Proceedings / AMIA 2002*;652-6. Database: Ovid MEDLINE(R).
 Exclude - Not MMIT

Reina-Tosina J, Roa L, Caceres J, et al. New approaches toward the fully digital integrated management of a burn unit. *IEEE Trans Biomed Eng* 2002;49(12):1470-6. Database: Embase Sept 22-09.
 Exclude - Not a Primary Study

Reinhardt ER. Technical paradigms for realizing ubiquitous care. *Studies in Health Technology & Informatics* 2008;134:129-34. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Reisner LA, Dong D, Guglielmo BJ, et al. Pro-active pharmacy engagement in medication reconciliation at a university hospital. *Ashp Summer Meeting 2007*;64(JUN):P29D Database: IPA.
 Exclude - Not a Primary Study

Relihan E, Silke B, O'Grady F. Internally-developed electronic reporting system for medication errors. *Ir Med J* 2009;102(7):223-4. OVID EMBASE.
 Exclude - No Outcomes of Interest

Remane Y, Vogel J, Hensel G. SAP based chemotherapy compounding tool - Boon or bane? *Krankenhauspharmazie* 2008;29(10):442-7. Database: Embase Sept 22-09.
 Exclude - Not a Primary Study

Rempher KJ, Lasome CE, Lasome TJ. Leveraging palm technology in the advanced practice nursing environment. *AACN Clin Issues* 2003;14(3):363-70. Database: Ovid MEDLINE(R).
 Exclude - Not MMIT

Rems M, Bohanec M, Urh B, et al. Implementation problems of decision support system for nosocomial infection. *Studies in Health Technology & Informatics* 1997;43 Pt A:358-62. Database: Ovid MEDLINE(R).
 Exclude - Not a Primary Study

Renshaw TM, Inman NS, Ly D. Computer assisted physician order entry within a neonatal intensive care unit. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Resetar E, Reichley RM, Noirot LA, et al. Customizing a commercial rule base for detecting drug-drug interactions. *AMIA 2005*;1094 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Resetar E, Reichley RM, Noirot LA, et al. Strategies for reducing nuisance alerts in a dose checking application. *AMIA 2005*;624-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Resetar E, Reichley RM, Noirot LA, et al. Implementing daily dosing rules using a commercial rule base. *AMIA 2006*;1073 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Reuther LO, List SB, Christensen HR. Application of the medicine profile in an ambulatory setting. *Ugeskr Laeger* 2008;170(33):2427-32. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Reynolds G, Boyer D, Mackey K, et al. Alerting strategies in computerized physician order entry: a novel use of a dashboard-style analytics tool in a children's hospital. *AMIA Annual Symposium 2008*;1108 PMID:18999063 OVID MEDLINE.

Exclude - No Outcomes of Interest

Reynolds K. Tech update. From paper to computerized entry: Creating a culture shift. *Nurs Manag (Harrow)* 2009;39(2):51-3. Database: CINAHL.

Exclude - Not MMIT

Reynolds MS, Kirkwood CF, Ostrosky JD, et al. Effect of an educational computer screen on direct physician order entry of anti-anaerobic drugs. *Ashp Midyear Clinical Meeting* 1988;23: Database: IPA.

Exclude - Not a Primary Study

Rhea S. Going mobile: wireless devices and technology bring surge in advanced applications for health monitoring and treatment, but legal and privacy issues remain. *Mod Healthc* 2010;40(18):28-30.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010646753&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010646753 EBSCO CINAHL.

Exclude - Not a Primary Study

Ribbons R. A difficult pill to swallow: Inpatient e-prescribing, rural and metropolitan experiences compared. *Journal on Information Technology in Healthcare* 2005;3(2):

Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Ribbons R, Dall V, Webb R. The better medication management system (BMMS). In 2002; Melbourne, Australia): 2002. Grey Lit (BMMS). Source: In: Ribbons, R. M.; Dall, V.; Webb, R.; National health informatics conference; Improving quality by lowering barriers M. Ribbons; p. 105; Brunswick East, Vic.;; Health Informatics Society of Australia Ltd.,; [2002?].

Exclude - No Outcomes of Interest

Ricciardelli M. Assuring cost effective ondansetron dosing: Two tiered approach. ASHP Annual Meeting 1995;52: Database: IPA.

Exclude - Not a Primary Study

Ricciardi TN, Masarie FE, Middleton B. Clinical benchmarking enabled by the digital health record. Studies in Health Technology & Informatics 2001;84(Pt:1):1-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rich DS. Compliance chatroom. Reconciling automated dispensing with prior pharmacist review. Joint Commission Benchmark 2002;4(2):3 Database: CINAHL.

Exclude - Not MMIT

Richardson RF, Caldwell JS, Hartz JL, et al. Evaluation of an automated dispensing machine in a large government hospital. Ashp Midyear Clinical Meeting 1991;26: Database: IPA.

Exclude - Not a Primary Study

Richesson RL, Malloy J, Paulus K, et al. An automated standardized system for managing adverse events in clinical research networks. Drug Saf 2008;31(10):807-22. Database: Embase Sept 22-09.

Exclude - Not MMIT

Riddell B. Information technology and the clinical nurse specialist. "Smart" infusion pumps and the clinical nurse specialist. Clinical Nurse Specialist: The Journal for Advanced Nursing Practice 2006;20(3):117-8. Database: CINAHL.

Exclude - Not a Primary Study

Riddle BM, Vinson BE. Cost benefit analysis of a computerized controlled substance distribution system. Ashp Midyear Clinical Meeting 1989;24: Database: IPA.

Exclude - Not a Primary Study

Ridinger MHT. The electronic prescription conundrum: Why "e-Rx" isn't so "e-Z". Clin Pharmacol Ther 2007;81(1):13-5. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Ried LD. Economic impact of a college of pharmacy, antiepileptic drug substitutions, and computerized prescriber order entry. Journal of the American Pharmacists Association: JAPhA 2009;49(6):838 PMID:19926568 OVID MEDLINE.

Exclude - Not MMIT

Riekert KA, Rand C. Electronic monitoring of medication adherence: When is high-tech best? Journal of Clinical Psychology in Medical Settings 2002;9(1):25-34. Database: Embase Sept 22-09.

Exclude - Not MMIT

Riley K. Long term care and automated pharmacy systems: Work in progress. Consultant Pharmacist 1997;12(Nov):1283-4. Database: IPA.

Exclude - Not MMIT

Rind D M, Safran C. The development and evaluation of computer-generated alerts in an inpatient setting. In Amsterdam, Netherlands: North-Holland; 1992. p.249-54.4373159

Database: Inspec.

Exclude - Unable to Retrieve

Rind D, Kim J, Sturges E and others. SeniorMed: Connecting patients to their medication records. In Washington, DC: 1999. p. 1147. Grey Lit.

Exclude - Not MMIT

Rinfret S, Lussier MT, Peirce A, et al. The impact of a multidisciplinary information technology-supported program on blood pressure control in primary care. Circulation 2009;Cardiovascular quality and outcomes. 2(3):170-7. OVID EMBASE.

Exclude - Not MMIT

Ringold DJ, Santell JP, Schneider PJ, et al. ASHP national survey of pharmacy practice in acute care settings: Prescribing and transcribing--1998. Am J Health Syst Pharm 1999;56:142-57. Database: IPA.

Exclude - Not MMIT

Ringold DJ, Santell JP, Schneider PJ. ASHP national survey of pharmacy practice in acute care settings: dispensing and administration--1999. Am J Health Syst Pharm 2000;57(19):1759-75. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Riou C, Pouliquen B, Le Beux P. A computer-assisted drug prescription system: the model and its implementation in the ATM knowledge base. Methods Inf Med 1999;38(1):25-30.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Risselada AJ. A task for experts. To a better oral anticoagulation therapy in the hospital. Pharm Weekbl 2000;135(7):235-41. Database: Embase Sept 22-09.

Exclude - Not MMIT

Rivers KJ, Dailey RE. Choosing a home infusion pharmacy software system. Infusion 1997;3(Feb):21-30. Database: IPA.

Exclude - Not MMIT

Roark DC. Bar codes & drug administration: Can new technology reduce the number of medication errors? Am J Nurs 2004;104(1):63-6. Database: CINAHL.

Exclude - Not a Primary Study

Roark DC. Bar codes and drug administration. Am J Nurs 2004;104(1):63-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Roark D, Miguel K. Bar coding's replacement? Nurs Manag (Harrow) 2006;37(2):29-31. Database: BSC.

Exclude - Not a Primary Study

Robbins GB. Transitioning from paper to virtual: Trials and tribulations of converting to an electronic medication administration record (eMAR) prior to computerized physician order entry (CPOE) in a community hospital. *Ashp Midyear Clinical Meeting 2006*;41: Database: IPA.

Exclude - Not a Primary Study

Robbins MF, Steiner C, Ratzman K, et al. Characteristics of a patient-safety-compliant computerized prescriber order entry (CPOE) database. *Ashp Midyear Clinical Meeting 2006*;41: Database: IPA.

Exclude - Not a Primary Study

Robbins MF, Steiner C, Ratzman K, et al. Enhancement of a computerized prescriber order entry (CPOE) database: Inclusion of renal and hepatic dosage adjustment information. *Ashp Midyear Clinical Meeting 2006*;41: Database: IPA.

Exclude - Not a Primary Study

Roberts GW, Farmer CJ, Cheney PC, et al. Clinical decision support implemented with academic detailing improves prescribing of key renally cleared drugs in the hospital setting. *J Am Med Inform Assoc* 2010;17(3):308-12.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010646540&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=2362&accno=2010646540 EBSCO CINAHL.

Exclude - Not MMIT

Roberts YD, Job ML, Jerris RC. Cost impact of a computer-assisted antibiotic monitoring program in a community hospital. *ASHP Annual Meeting 1992*;49: Database: IPA.

Exclude - Not a Primary Study

Robertson J, Henry D, Dobbins T, et al. Prescribing patterns in general practice. A comparison of two data sources. *Aust Fam Physician* 1999;28(11):1186-90. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Robertson J, Walkom E, Pearson SA, et al. The impact of pharmacy computerised clinical decision support on prescribing, clinical and patient outcomes: A systematic review of the literature. *International Journal of Pharmacy Practice* 2010;18(2):69-87. PMID:20441116 OVID MEDLINE.

Exclude - Not a Primary Study

Robertson L, Smith M, Tannenbaum D, et al. Pilot implementation of an online disease management system for depression in Australia. *J Telemed Telecare* 2005;11(Suppl. 2):108-11. 8766834

Database: Inspec.

Exclude - Not a Primary Study

Robeznieks A, Lubell J. 'Meaningful use' at last: regs aim to boost quality, safety and efficiency. *Mod Healthc* 2010;40(1):4

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010519353&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010519353 EBSCO CINAHL.

Exclude - Not a Primary Study

Robeznieks A. More than meaningful use: HIMSS attendees urged to focus on patient care. *Mod Healthc* 2010;40(10):30-1.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010584258&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010584258 EBSCO CINAHL.

Exclude - Not a Primary Study

Robinson BR, Johnson D. For emergency medical service helicopter pilots, all wind is local. *Air Med J* 256;28(5):232-6. PMID:19748447 OVID MEDLINE.

Exclude - Not a Primary Study

Robinson D, Bass G, Anderson-Harper H, et al. Impact of the meditrol automated medication system on hospital pharmacy cost and quality of care. *Ashp Midyear Clinical Meeting* 1989;24:119-26. Database: IPA.

Exclude - Not a Primary Study

Robinson DL, Heigham M, Clark J. Using Failure Mode and Effects Analysis for safe administration of chemotherapy to hospitalized children with cancer. *Jt Comm J Qual Patient Saf* 2006;32(3):161-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Robinson G. PRODIGY: A support tool for prescribers. *Practice Nurse* 2005;27(9):20-1. Database: CINAHL.

Exclude - Not MMIT

Rode D. AHIMA's advocacy priorities for 2009. *Journal of the American Health Information Management Association* 2009;80(1):16-8. OVID EMBASE.

Exclude - Not a Primary Study

Rode D. Reassessing privacy and security compliance. *Journal of the American Health Information Management Association* 2009;80(10):20-2. OVID EMBASE.

Exclude - Not a Primary Study

Rodman J, Jelliffe R, Tuey D, et al. Clinical studies with computer-assisted initial lidocaine therapy. *Arch Intern Med* 1984;144(4):703-9. Exclude - Not MMIT

Rodriguez N J, Borges J A, Soler Y and others. PDA vs. laptop: A comparison of two versions of a nursing documentation application. In Los Alamitos, CA, USA: IEEE; 2003.

p.201-6.7763192

Database: Inspec.

Exclude - Not MMIT

Rodway A, Fleetwood JA, Laker MF, et al. Standardised monitoring of patients on long-term medication in primary care. *Br J Gen Pract* 2002;52(Suppl):S37-S39 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Roemer LK, Richardson SJ, Rocha RA, et al. Exploratory case method to determine the frequency of redundant orders within manually consolidated order lists. *Int J Med Inf* 2004;73(7-8):639-45. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Roger France FH. The advancement, progress and challenges in medical informatics in Belgium. *Bull Mem Acad R Med Belg* 2007;162(1-2):129-36. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Roggow DJ, Solie CJ, Tracy MF, et al. Clinical nurse specialist leadership in computerized provider order entry design. *Clin Nurse Spec* 2005;19(4):209-14. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Roig RJ, Sanchez DS, Oliete MD, et al. Optimising the quality of the unit dose dispensing process through the implementation of the semi-automated Kardex(R) system. *Farmacia Hospitalaria* 2007;31(1):38-42. Database: IPA.

Exclude - Not MMIT

Roitman J, Kalra S. Interventions to improve adherence to lipid lowering medication. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2007;27(2):61-118. Database: CINAHL.

Exclude - Not a Primary Study

Roland MO, Zander LI, Evans M, et al. Evaluation of a computer assisted repeat prescribing programme in a general practice. *British Medical Journal Clinical Research Ed* 1985;291(6493):456-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Roman-Vinas R. Factors of success in the implementation of the technologies of the information and the communication in the health systems. The human factor. [Spanish]. *Med Clin (Barc)* 2010;134(SUPPL. 1):39-44. OVID EMBASE.

Exclude - No Outcomes of Interest

Romera IC. The growth of outpatient units. *EJHP Practice* 2009;15(2):50 OVID EMBASE.

Exclude - Not a Primary Study

Romero AV, Malone DC. Accuracy of adverse-drug-event reports collected using an automated dispensing system. *Am J Health Syst Pharm* 2005;62(13):1375-80. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rommers MK, Teepe-Twiss IM, Guchelaar HJ. Preventing adverse drug events in hospital practice: an overview. *Pharmacoepidemiology & Drug Safety* 2007;16(10):1129-35.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ros JJW, Cramers B, Prins J, et al. The contribution of informatics to medication safety: Business process redesign in the hospital. *Pharm Weekbl* 2002;137(44):1570-4. Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Rosati K. Using electronic health information for pharmacovigilance: the promise and the pitfalls. *Journal of Health & Life Sciences Law* 2009;2(4):171-239. PMID:19673181 OVID MEDLINE.

Exclude - Not a Primary Study

Roscoe JC, Goransson E, Slancar M, et al. Multidisciplinary approach to ensure safety in the prescribing and administration of chemotherapy. *Journal of Oncology Pharmacy Practice* 2000;6(2):60-3. Database: IPA.

Exclude - Not a Primary Study

Rose JL. Improved and expanded pharmacy care in rural Alaska through telepharmacy and alternative methods demonstration project. *Int J Circumpolar Health* 2007;Suppl 1(66):14-22. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rose PE, Fitzmaurice D. New approaches to the delivery of anticoagulant services. *Blood Rev* 1998;12(2):84-90. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rosenbaum BP, Patel SG, Guyer DL, et al. The pharmaceutical management system at Shade Tree Family Clinic: A medical student-run free clinic's experience. *Informatics for health & social care* 2009;33(3):151-7. Database: CINAHL.

Exclude - Not a Primary Study

Rosenberg JM, Fuentes RJ, Kirschenbaum HL, et al. Comparative content evaluation of the de Haen and DRUGDEX drug information systems. *Am J Hosp Pharm* 1983;40(Oct):1664-9. Database: IPA.

Exclude - Not MMIT

Rosenberg SN, Shnaiden TL, Wegh AA, et al. Supporting the patient's role in guideline compliance: A controlled study. *Am J Manag Care* 2008;14(11):737-44. Database: CINAHL.

Exclude - Not MMIT

Rosenbloom ST, Grande J, Geissbuhler A, et al. Experience in implementing inpatient clinical note capture via a provider order entry system. *J Am Med Inform Assoc* 2004;11(4):310-5. Database: CINAHL.

Exclude - Not MMIT

Rosenbloom S. Approaches to evaluating electronic prescribing. *J Am Med Inform Assoc* 2006;13(4):399-401.

<http://search.ebscohost.com/login.aspx?direct=true&db=lxh&AN=21635005&site=ehost-live>
Database: LISTA.

Exclude - Not a Primary Study

Rosenfeld S, Mendelson D. Health information technology policy: Legislative and regulatory progress in 2003, and prospects for the future. 2004. <http://nyam.waldo.kohalibrary.com/cgi-bin/koha/opac-detail.pl?biblionumber=225547> Grey Lit.

Exclude - Not a Primary Study

Rosenkranz K O, Reichertz P L. DAVID-a dialog system for acquisition and validation of information on drugs. In Amsterdam, Netherlands: North-Holland; 1974. p.883-8.1006961 Database: Inspec.

Exclude - Not MMIT

Rosenstein AH. Measuring the benefits of clinical decision support: return on investment. *Pharm Pract Manag Q* 2000;20(1):28-43. Database: IPA.

Exclude - Not a Primary Study

Ross J. Collaboration -- integrating nursing, pharmacy and information technology into a barcode medication administration system implementation. CARING Newsletter 2008;23(1):1-8. Database: CINAHL.

Exclude - Not a Primary Study

Ross S. Results of a survey of an online physician community regarding use of electronic medical records in office practices. J Med Pract Manage 2009;24(4):254-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rossi L, Margola L, Manzelli V, et al. wHospital: a web-based application with digital signature for drugs dispensing management. Conference proceedings : Annual International Conference of the IEEE Engineering in Medicine and Biology Society IEEE Engineering in Medicine and Biology Society Conference 2006;Suppl:6793-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rossi RA, Every NR. A computerized intervention to decrease the use of calcium channel blockers in hypertension. J Gen Intern Med 1997;12(11):672-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rossos PG, Abrams H, Wu R, et al. Active physician participation key to smooth MOE/MAR rollout. HEALTHC Q 2006;10(Sp):58-64. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rostom A, O'Connor A, Tugwell P, et al. A randomized trial of a computerized versus an audio-booklet decision aid for women considering post-menopausal hormone replacement therapy. Patient Education & Counseling 2002;46(1):67-74. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Roten I, Marty S, Beney J. Electronic screening of medical records to detect inpatients at risk of drug-related problems. Pharmacy World & Science 2010;32(1):103-7. PMID:20012362 OVID MEDLINE.

Exclude - Not MMIT

Roth KB, Palmer R, Linck C, et al. Implementation of Pyxis profiling in a large teaching institution. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.

Exclude - Not a Primary Study

Rothschild J. Computerized physician order entry in the critical care and general inpatient setting: a narrative review. J Crit Care 2004;19(4):271-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rothschild JM, Lee TH, Bae T, et al. Clinician use of palmtop drug reference guide. J Am Med Inform Assoc 2002;9(3):223-9. Database: CINAHL.

Exclude - Not MMIT

Rothschild JM, Keohane CA, Thompson S, et al. Intelligent intravenous infusion pumps to improve medication administration safety. AMIA 2003;992 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rothschild JM, Keohane CA, Cook EF, et al. A controlled trial of smart infusion pumps to improve medication safety in critically ill patients. *Crit Care Med* 2005;33(3):533-40.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rothschild JM, Fang E, Gottschall J, et al. Use and perceived benefits of handheld PDA clinical reference applications. *AMIA Annual Symposium proceedings* 2005;2005:1099

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rotich JK, Hannan TJ, Smith FE, et al. Installing and implementing a computer-based patient record system in sub-Saharan Africa: the Mosoriot Medical Record System. *J Am Med Inform Assoc* 2003;10(4):295-303. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rotman BL, Sullivan AN, McDonald T, et al. A randomized evaluation of a computer-based physician's workstation: design considerations and baseline results. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1995;693-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rough S. Improving the medication administration process - The impact of point-of-care bar code medication scanning technology. *Journal of the Pharmacy Society of Wisconsin* 2004;(May-Jun):4 Database: IPA.

Exclude - No Outcomes of Interest

Rough SS. Development of a physician order entry project plan to maximize medication use safety. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Rousseau N, McColl E, Newton J, et al. Practice based, longitudinal, qualitative interview study of computerised evidence based guidelines in primary care. *Br Med J* 2003;326(7384):314-21. Companion 6796. Exclude - Not MMIT

Roux RK, Pedersen CA, Schneider PJ, et al. Evaluation of medication rules within a physician order entry system on appropriate prescribing of low molecular weight heparins (LMWH). *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Roycroft R. Does IT 'cut the mustard' in primary care? *Inform Prim Care* 2004;12(2):97-102. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Rozak CT, Ronshagen JP. Effect of automation on dispensing errors. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Rozich JD, Haraden CR, Resar RK. Adverse drug event trigger tool: A practical methodology for measuring medication related harm. *Quality & Safety in Health Care* 2003;12(3):194-200. Database: CINAHL.

Exclude - Not MMIT

Rozonkiewiecs M, Walters J. Robots in the Rx: Automating medication distribution. *Healthc Inform* 1995;12(12):20-6. Database: IPA.

Exclude - Not a Primary Study

Rubio FM, Aldaz FR, Garcia GC, et al. Computer-aided electronic prescribing in Spanish hospitals. *Farmacia Hospitalaria* 2005;29(4):236-40. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Rudman WJ, Bailey JH, Hope C, et al. Impact of a web-based reporting system on the collection of medication error occurrence data. *Advances in Patient Safety* 2005;3: Implementation Issues: Grey Lit.

Exclude - Not MMIT

Ruffin M. Information technology. Part 2: Interactive media enhances medicine. *Physician Exec* 1996;22(8):46-8. Database: BSC.

Exclude - Not a Primary Study

Ruixo JJ, Romero G, Vargas MI, et al. Pharmacoeconomic impact of the implementation of an automated dispensing system in the emergency department of a university hospital. *Farmacia Hospitalaria* 2000;24(6):390-7. Database: IPA.

Exclude - Unable to Retrieve Foreign

Ruiz-Pena JL, Duque P, Izquierdo G. Optimization of treatment with interferon beta in multiple sclerosis. Usefulness of automatic system application criteria. *BMC Neurology* 2008;8(3): Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ruland CM. A survey about the usefulness of computerized systems to support illness management in clinical practice. *Int J Med Inf* 2004;73(11-12):797-805. Database: Embase Sept 22-09.

Exclude - Not MMIT

Rulon V. The right information at the right time: Improving the quality of health information improves quality of care. *Journal of the American Health Information Management Association* 2009;80(8):10 OVID EMBASE.

Exclude - Not a Primary Study

Runy LA. Your ICD-10 planning starts now! *Hospitals & health networks / AHA* 2009;83(12):33-40. OVID EMBASE.

Exclude - Not a Primary Study

Ruotsalainen P, Iivari AK, Doupi P. Finland's strategy and implementation of citizens' access to health information. *Studies in Health Technology & Informatics* 2008;137:379-85. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Rupp MT. E-prescribing: The value proposition. *America's Pharmacist* 2005;127(4):23-6. Database: IPA.

Exclude - Not a Primary Study

Rupp MT. Maximizing the effectiveness of e-prescribing between physicians and community pharmacies. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Russ A L, Saleem J J, McManus M S and others. Computerized Medication Alerts and Prescriber Mental Models: Observing Routine Patient Care. In 11. San Antonio, TX, United states: Human Factors and Ergonomics Society Annual Meeting Proceedings; 2009. p.655-9. Engineering Village Compendex and Inspec.

Exclude - Unable to Retrieve

Russell J. How technology solutions can impact nursing retention. *Nurs Econ* 2008;26(3):188-90. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Russo MF, Lam PK, Goldberg MJ. Pyxis installation in a large teaching institution. *Ashp Summer Meeting* 2002;59. Database: IPA.

Exclude - Not a Primary Study

Ryan F, Byrne S, O'Shea S. Managing oral anticoagulation therapy: improving clinical outcomes. A review. *J Clin Pharm Ther* 2008;33(6):581-90. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ryan F, Byrne S, O'Shea S. Randomized controlled trial of supervised patient self-testing of warfarin therapy using an internet-based expert system. *Journal of Thrombosis and Haemostasis* 2009;7(8):1284-90. Database: Embase Sept 22-09.

Exclude - Not MMIT

Ryan MJ. Australian electronic medication management policy and systems. *Journal of Pharmacy Practice and Research* 2007;37(1):49-55. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Ryan ML, Rinke R, de Leon RF. Selecting a pharmacy computer system for the future. *Pharm Pract Manag Q* 1995;15(3):1-14. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Rybak MJ. Antimicrobial stewardship. *Pharmacotherapy* 2007;27(10 Pt 2):131S-5S. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Saager L, Collins GL, Burnside B, et al. A randomized study in diabetic patients undergoing cardiac surgery comparing computer-guided glucose management with a standard sliding scale protocol. *Journal of Cardiothoracic & Vascular Anesthesia* 2008;22(3):377-82. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Exclude - Not MMIT

Saathoff A. Human factors considerations relevant to CPOE implementations. *J Healthc Inf Manag* 2005;19(3):71-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Saeed M, Lieu C, Raber G, et al. MIMIC II: a massive temporal ICU patient database to support research in intelligent patient monitoring. *Computers in Cardiology* 2002;29:641-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Safian SC. EHR triggers: a force for improved care--and revenue. *Healthc Financ Manage* 2009;63(9):108-14. PMID:19743656 OVID MEDLINE.

Exclude - Not a Primary Study

Safren SA, Hendriksen ES, Desousa N, et al. Use of an on-line pager system to increase adherence to antiretroviral medications. *AIDS Care* 2003;15(6):787-93. Database: PsycINFO.

Exclude - Not MMIT

Sager RB, Woodward C. Restricted medication management program: A university health-system's approach for improvement. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Saginur M, Graham ID, Forster AJ, et al. The uptake of technologies designed to influence medication safety in Canadian hospitals. *J Eval Clin Pract* 2008;14(1):27-35. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Saginur, M. Technologies to improve medication safety in hospitals: A study of their effectiveness and use in Canada University of Ottawa (Canada) Editor. 2005. Grey Lit.

Exclude - Not a Primary Study

Sakai Y, Orii T, Liu Z, et al. Physicians' evaluations of the prescription order entry system in university hospitals. *Japanese Journal of Hospital Pharmacy* 1998;24(4):409-15. Database: IPA.

Exclude - Unable to Retrieve Foreign

Sakowski J, Leonard T, Colburn S, et al. Using a bar-coded medication administration system to prevent medication errors in a community hospital network. *Am J Health Syst Pharm* 2005;62(24):2619-25. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sakowski J, Newman JM, Dozier K. Severity of medication administration errors detected by a bar-code medication administration system. *Am J Health Syst Pharm* 2008;65(17):1661-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Salasidis R, Padjen AL, Fleischer D. Patient management in the ICU: the PDB System. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1991;990-2. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Saleem JJ, Patterson ES, Militello L, et al. Impact of clinical reminder redesign on learnability, efficiency, usability, and workload for ambulatory clinic nurses. *J Am Med Inform Assoc* 2007;14(5):632-40. Database: IPA.

Exclude - Not MMIT

Saliba A. Reduction of insulin errors in a community hospital. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Salmivalli L, Hilmola OP. Business pluralism of electronic prescriptions: state of development in Europe and the USA. *International Journal of Electronic Healthcare* 2006;2(2):132-48. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Salzer LB. Basics of bar coding in health care. Consultant Pharmacist 2004;2(Mar-Apr):125-6. Database: IPA.

Exclude - Not MMIT

Samarth A and Grant E. Using health IT to determine medication adherence: Findings from the AHRQ health IT portfolio. AHRQ Publication No. 10-0010-EF. AHRQ; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_909099_0_0_18/10-0010-EF.pdf Grey Lit.

Exclude - Not a Primary Study

Samore MH, Bateman K, Alder SC, et al. Clinical decision support and appropriateness of antimicrobial prescribing: a randomized trial. JAMA 2005;294(18):2305-14. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Sanborn M, Sheehan VC, Cohen T. Clinical transformation: Formulary standardization in a large healthcare system. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Sanborn MD, Krueger JL. Survey of automation in hospital pharmacy practice. Am J Health Syst Pharm 1995;52(1):97-8. Database: IPA.

Exclude - Not a Primary Study

Sanchez-Cosio R, Mutnick AH, Abramowitz PW, et al. Study of continuity of pharmaceutical care between a tertiary hospital and local pharmacies. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.

Exclude - Not a Primary Study

Sandberg WS, Sandberg EH, Seim AR, et al. Real-time checking of electronic anesthesia records for documentation errors and automatically text messaging clinicians improves quality of documentation. Anesthesia & Analgesia 2008;106(1):192-201. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Sanders K H, Jr., Radford W E. The computer in a programmable implantable medication system (PIMS). In New York, NY, USA: IEEE; 1982. p.682-5.2031120

Database: Inspec.

Exclude - Not a Primary Study

Sandhoff BG, Nies LK, Olson KL, et al. Clinical pharmacy cardiac risk service for managing patients with coronary artery disease in a health maintenance organization. Am J Health Syst Pharm 2007;64(1):77-84. Database: Embase Sept 22-09.

Exclude - Not MMIT

Sandhuja S, Girouard DL. Failure mode and effects analysis (FMEA) of a computerized unfractionated heparin (UFH) nomogram. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Sani S, Shane R, Geller R, et al. Implementation of a quality assurance program to ensure the safety and effectiveness of a centralized pharmacy automation system. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Santell JP. ASHP national survey of hospital-based pharmaceutical services--1994. Am J Health Syst Pharm 1995;52(11):1179-98. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Santell JP, Cousins DD. Error risks associated with computer order entry. Drug Topics 2004;147: Database: IPA.

Exclude - Not a Primary Study

Santucci RA. Editorial Comment. Urology 2010;75(5):1185-92. OVID EMBASE.

Exclude - Not a Primary Study

Sanz EJ. Methodological approaches to ADR detection in out-patient children. Bratisl Lek Listy 1991;92(12):597-602. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sanz F, Loza M, Ahlgrimm E and others. Telematics in community pharmacies to support responsible self-medication. The TESEMED project. In 1996; (Copenhagen): 1996. p.141-8. Grey Lit.

Exclude - Not MMIT

Sanz F, Silveira C, Mircheva J, et al. Telematics applications to support the role of the community pharmacists as self-medication advisors. TESEMED Consortium. Stud Health Technol Inform 1999;68:764-7. Exclude - Not MMIT

Sarachek N S, Kreithen D E. Comparative speed and accuracy of microcomputer ICU calculations. In Bethesda, MD, USA: American Assoc. Medical Systems and Informatics; 1983. p.313-4.2188344

Database: Inspec.

Exclude - Not MMIT

Saranto K, Ensio A, Jokinen T. Patient medication: How is it documented? Stud Health Technol Inform 2006;122:738-41. Exclude - Not MMIT

Saraoglu HM, Sanli S. A fuzzy logic-based decision support system on anesthetic depth control for helping anesthetists in surgeries. J Med Syst 2007;31(6):511-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sarasin F, Schifferli JA. Immuno-allergic interstitial nephritis: steroids yes or no? [French]. Schweizerische Medizinische Wochenschrift 1990;120(49):1858-61. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Sarasohn-Kahn J and Holt M. The prescription infrastructure: Are we ready for eprescribing? California HealthCare Foundation; 2006.

<http://www.chcf.org/topics/view.cfm?itemID=118337> Grey Lit.

Exclude - Not a Primary Study

Sardinha C. Electronic prescribing: Next revolution in pharmacy? *J Manag Care Pharm* 2004;4(1):35-9. Database: IPA.

Exclude - No Outcomes of Interest

Sary A, Johnsen J A. The development and impact of a drug exception reporting system in a psychiatric facility. In New York, NY, USA: IEEE; 1980. p.1752-5.1644532

Database: Inspec.

Exclude - Not MMIT

Sassi G, Striano B, Merlo UA. A reporting system for the assessment of chemotherapy toxicity. *Journal of Oncology Pharmacy Practice* 2005;11(2):63-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Satchu S, Hua Y, Ryl A, et al. Development and implementation of an electronic viewable medication profile. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Sato N. Drug control system by bar code. *Pharmaceuticals Monthly* 1988;30(May):957-60. Database: IPA.

Exclude - Not a Primary Study

Saubermann LA. Implementation of a computerized provider order entry (POE) system greatly improves allergy documentation in a pharmacy computer system. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Saull-McCaig S, Pacheco R, Kozak P, et al. Implementing MOE/MAR: balancing project management with change management. *HEALTHC Q* 2004;10(Sp):27-38. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Savage SW, Schneider PJ, Pedersen CA. Utility of an online medication-error-reporting system. *Am J Health Syst Pharm* 2005;62(21):2265-70. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Saya FG, Shane R. A stepwise approach to the evaluation and selection of a hospital pharmacy information system. *Pharm Pract Manag Q* 1995;15(3):15-22. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Scalise D. CPOE (computerized physician order entry). An executive's guide. *Hospitals & Health Networks* 2002;76(6):41-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Scalise D. The case for e-prescribing. *Hospitals & Health Networks* 2007;81(2):45-51. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Scanlon M. Computer physician order entry and the real world: we're only humans. *Joint Commission Journal on Quality & Safety* 2004;30(6):342-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Scarafile PD, Campbell BD, Kilroy JE, et al. Computer-assisted concurrent antibiotic review in a community hospital. *Am J Hosp Pharm* 1985;42(2):313-5. Database: IPA.
Exclude - Not a Primary Study

Scavuzzo J, Gamba N. Bridging the gap: the Virtual Chemotherapy Unit. *J Pediatr Oncol Nurs* 2004;21(1):27-32. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Schade CP, Sullivan FM, de Lusignan S, et al. e-Prescribing, efficiency, quality: lessons from the computerization of UK family practice. *J Am Med Inform Assoc* 2006;13(5):470-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Schadow G. Structured product labeling improves detection of drug-intolerance issues. *J Am Med Inform Assoc* 2009;16(2):211-9. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Schafer T, Rothe N, Kim E, et al. Introduction of an electronic charting software with computerized physician order entry and medication safety features - Experience at the University Hospital Freiburg, Germany. [German]. *Krankenhauspharmazie* 2009;30(12):569-74. OVID EMBASE.
Exclude - Unable to Retrieve

Schecke T, Langen M, Popp HJ, et al. Knowledge-based decision support for patient monitoring in cardioanesthesia. *Int J Clin Monit Comput* 1992;9(1):1-11. 4220021 Database: Inspec.
Exclude - Not a Primary Study

Scheckelhoff DJ, Mark SM, Gumpfer KF. Development of a medication safety strategic plan for a hospital or health system. *ASHP Annual Meeting* 2001;58: Database: IPA.
Exclude - Not a Primary Study

Scheckelhoff DJ. ASHP's approach to advocating for bar code unit dose medication systems. *Hosp Pharm* 2003;38(11):S9-S10 Database: IPA.
Exclude - Not MMIT

Schedlbauer A, Prasad V, Mulvaney C, et al. What evidence supports the use of computerized alerts and prompts to improve clinicians' prescribing behavior? *J Am Med Inform Assoc* 2009;16(4):531-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Scheepers-Hoeks AM, Grouls RJ, Neef C, et al. Strategy for implementation and first results of advanced clinical decision support in hospital pharmacy practice. *Studies in Health Technology & Informatics* 2009;148:142-8. PMID:19745244 OVID MEDLINE.
Exclude - No Outcomes of Interest

Scherr D, Zweiker R, Kollmann A, et al. Mobile phone-based surveillance of cardiac patients at home. *Journal of Telemedicine & Telecare* 2006;12(5):255-61. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Schetz M, Dasta J, Goldstein S, et al. Drug-induced acute kidney injury. *Current Opinion in Critical Care* 2005;11(6):555-65. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schiff G, Bates DW. Electronic point-of-care prescribing: Writing the 'script'. *Disease Management and Health Outcomes* 2000;7(6):297-304. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Schiff GD, Rucker TD. Computerized prescribing: building the electronic infrastructure for better medication usage. *JAMA* 1998;279(13):1024-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schiff GD. Computerized prescriber order entry: models and hurdles. *Am J Health Syst Pharm* 2002;59(15):1456-60. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schiff GD, Klass D, Peterson J, et al. Linking laboratory and pharmacy: opportunities for reducing errors and improving care. *Arch Intern Med* 2003;163(8):893-900. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schissler K, Triplett J, Ortega J, et al. Medication reconciliation within a computerized medical record. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Schlesinger JM, Blumenfeld B, Broverman C. Component architecture in HIS: a drug order entry case study. *Studies in Health Technology & Informatics* 1997;45:54-60. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schloerb PR. Electronic parenteral and enteral nutrition. *Journal of Parenteral & Enteral Nutrition* 2000;24(1):23-9. Database: IPA.

Exclude - Not MMIT

Schmiddek JM, Weeks WB. What do we know about financial returns on investments in patient safety? A literature review. *Jt Comm J Qual Patient Saf* 2005;31(12):690-9 (10). Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schmidt NJ. Effective change to the distribution process, which decreased medication turnaround time. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Schmidt S, Grimm A. Health service research of telemedicine applications. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2009;52(3):270-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schmidt W, Alberici G, Cook L. Development of a computerized system for inventory of controlled drug substances. *NIDA Res Monogr* 1988;81:163-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schmitt D, Allenet B, Brudieu E, et al. Acceptability of computerized prescription writing in the hospital. Survey of 44 drug-prescribing physicians. *Presse Med* 2001;30(30):1478-81. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Schneck LH. E-prescribing can be new tool in quality-care arsenal. *Mgma Connexion/Medical Group Management Association* 2001;6(1):32-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schneider AJ, Murray WB, Mentzer SC, et al. "Helper:" A critical events prompter for unexpected emergencies. *J Clin Monit* 1995;11(6):358-64. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schneider BL, Gulseth MP, Cusick MA, et al. Computer program to assist pharmacy management of an inpatient warfarin dosing service. *Am J Health Syst Pharm* 2005;62(22):2393-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schneider PJ. Using technology to enhance measurement of drug-use safety. *Am J Health Syst Pharm* 2002;59(23):2330-2. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schneider PJ. Framing the opportunity for bar code medication administration. *Hosp Pharm* 2003;38(11):S3-S4 Database: IPA.

Exclude - Not MMIT

Schneider R, LaBatt EP. Implementation of a computerized physician order entry system. *Ashp Midyear Clinical Meeting* 1994;29: Database: IPA.

Exclude - Not a Primary Study

Schneider R. Comparison of implementation of a physician order entry system in the inpatient and outpatient setting. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Schneider R, Carlson R. Implementation of a closed loop verification program: Reporting problematic bar codes in a bar code medication administration system from the reporting facility to manufacturer notification. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Schneider R, Bagby J, Carlson R. Bar-code medication administration: a systems perspective. *Am J Health Syst Pharm* 2008;65(23):2216-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schneider R, Bagby J, Carlson R. Informatics interchange. Bar-code medication administration: a systems perspective. *Am J Health Syst Pharm* 2008;65(23):2216-9. <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010122091&site=ehost-live>; EBSCO CINAHL.

Exclude - Not a Primary Study

Schneider S, Gustavson B, Li T et al. Personal health information management and the design of consumer health information technology: Secondary analysis of data from the medical expenditure panel survey. AHRQ; 2009.

http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_1248_907541_0_0_18/09-0090-EF.pdf Grey Lit.

Exclude - Not a Primary Study

Schnider TW, Gaeta R, Brose W, et al. Derivation and cross-validation of pharmacokinetic parameters for computer-controlled infusion of lidocaine in pain therapy. *Anesthesiology* 1996;84(5):1043-50. Database: Embase Sept 22-09.

Exclude - Not MMIT

Schnipper JL, Linder JA, Palchuk MB, et al. "Smart Forms" in an Electronic Medical Record: Documentation-based Clinical Decision Support to Improve Disease Management. *J Am Med Inform Assoc* 2008;15(4):513-23. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Schnipper JL, Gandhi TK, Wald JS, et al. Design and implementation of a web-based patient portal linked to an electronic health record designed to improve medication safety: the Patient Gateway medications module. *Inform Prim Care* 2008;16(2):147-55 (9). Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schnurrer JU, Frolich JC. Incidence and prevention of lethal undesirable drug effects. *Internist* 2003;44(7):889-95. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schock P. Less paper, less fuss, better patient care. *Health Manag Technol* 2007;28(2):62-3. 9332633

Database: Inspec.

Exclude - Not a Primary Study

Schoenman JA, Keeler J, Moiduddin A et al. Roadmap for the adoption of health information technology in rural communities. Bethesda, MD: Federal Office of Rural Health Policy; 2006. http://www.norc.org/NR/rdonlyres/6A09114C-1B4D-4834-A942-8D6E0EDB799B/0/HIT_Paper_Final.pdf Grey Lit.

Exclude - Not a Primary Study

Schreier G, Hayn D, Kastner P, et al. A mobile-phone based teledermatology system to support self-management of patients suffering from psoriasis. *Conference Proceedings*: 2008;2008:5338-41. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schrezenmeir J, Achterberg H, Bergeler J, et al. Controlled study on the use of hand-held insulin dosage computers enabling conversion to and optimizing of meal-related insulin therapy regimens. *Life Support Syst* 1985;3(Suppl 1):561-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schroeder CG, Pierpaoli PG. Direct order entry by physicians in a computerized hospital information system. *Am J Hosp Pharm* 1986;43(2):355-9. Database: IPA.

Exclude - Not a Primary Study

Schuenke C. Computerized Physician Order Entry (CPOE) Technology success requires increased pharmacist involvement. *Journal of the Pharmacy Society of Wisconsin* 2009;(March/April):29-30. Database: IPA.

Exclude - Not MMIT

Schuetz D. If you build it, will they come? e-Prescribing's "Field of dreams". *Computertalk for the Pharmacist* 1938;22(5):38 Database: IPA.

Exclude - Not a Primary Study

Schug SH. Health Telematics - Current Developments and Consequences for Hospitals and Care-providers. *Klinikarzt* 2003;32(11):391-7. Database: Embase Sept 22-09.

Exclude - Not MMIT

Schug SH, Redders M. Health telematics projects in the perspective of the German federal states. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2005;48(6):649-56. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schulmeister L. Ten simple strategies to prevent chemotherapy errors. *Clin J Oncol Nurs* 2005;9(2):201-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schulmeister L. Look-alike, sound-alike oncology medications. *Clin J Oncol Nurs* 2006;10(1):35-41. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Schultz EN, Temkin LA, Mahoney CD. Use of bar code technology to improve inventory management. *Am J Hosp Pharm* 1990;47(7):1592-4. Database: IPA.

Exclude - Not MMIT

Schulz G, Beyer J, Hohleweg F, et al. Diabetes self-adjustment by a computerized program--first experiences in inpatient and outpatient treatment. *Life Support Syst* 1985;3(Suppl 1):578-82. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schumock GT, Nair VP, Finley JM, et al. Penetration of medication safety technology in community hospitals. *J Med Syst* 2003;27(6):531-41. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schwartz G. Automated efficiencies implemented at Rush Medical Center, Chicago. *U S Pharmacist* 2003;28(10):106-7. Database: IPA.

Exclude - Not MMIT

Schwartz JI, Fink JL, III. Legal issues associated with pharmacokinetic software. *Am J Hosp Pharm* 1989;46(1):120-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Schwartz JR, Erush SC. Evaluation of documentation of patient height, weight, and allergy information at a university medical center. *Ashp Summer Meeting* 2003;60: Database: IPA.

Exclude - Not a Primary Study

Sciamanna CN, Gifford DR, Smith RJ. Design and acceptability of patient-oriented computerized diabetes care reminders for use at the point of care. *Med Inform Internet Med* 2004;29(2):157-68. Database: Embase Sept 22-09.

Exclude - Not MMIT

Scimeca PG, Weinblatt M. A PC database to facilitate treatment of pediatric hematology/oncology patients. *Pediatr Hematol Oncol* 1995;12(3):259-69. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Sciar DA, VanDerLinde LP, Skaer TL. Strategic data-base management for DUR and ADR reporting. *Ashp Midyear Clinical Meeting* 1989;24: Database: IPA.

Exclude - Not a Primary Study

Sciar DA, VanDerLinde LP, Skaer TL. Medical information for DUR & ADR reporting: Multi-year experience. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.

Exclude - Not a Primary Study

Scott-Cawiezell J, Madsen RW, Pepper GA, et al. Medication safety teams' guided implementation of electronic medication administration records in five nursing homes. *Jt Comm J Qual Patient Saf* 2009;35(1):29-35. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Scott A, Coote W. Whither Divisions of General Practice? An empirical and policy analysis of the impact of Divisions within the Australian health care system. *Med J Aust* 2007;187(2):68-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Scott IA, Denaro CP, Bennett CJ, et al. Achieving better in-hospital and after-hospital care of patients with acute cardiac disease. *Med J Aust* 2004;180(Suppl 10):S83-S88 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Scott JS, Phillips LL. Early implementation experiences with an orders interface from a clinical information system to a pharmacy system: Challenges of blending two workflows. *ASHP Annual Meeting* 1998;55: Database: IPA.

Exclude - Not a Primary Study

Scott JS, Fitzgerald K, Sampson E, et al. Impact of robotic technology on the accuracy and quality of a centralized cart fill process. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Scotto M, Abraham T, Chiu R, et al. Implementation of a bar-code IV infusion pump program. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Seals AB, Duffy VG. Toward development of a computer-based methodology for evaluating and reducing medication administration errors. *Ergonomics* 2005;48(9):1151-68. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Searles B, Nasrallah F, Graham S, et al. Electronic data management for the Hemochron Jr. Signature coagulation analyzer. *J Extra Corpor Technol* 2002;34(3):182-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Section of Pharmacy Informatics and Technology ASoH-SP. ASHP Statement on bar-code-enabled medication administration technology. *Am J Health Syst Pharm* 2009;66:588-90.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Seden K, Back D, Khoo S. Antiretroviral drug interactions: often unrecognized, frequently unavoidable, sometimes unmanageable. *J Antimicrob Chemother* 2009;64(1):5-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Seelinger PM. Financial/workload evaluation for an automated controlled substance dispensing system. *Ashp Midyear Clinical Meeting* 1992;27: Database: IPA.

Exclude - Not a Primary Study

Segarra J, DeStefano JJ, Davis RH. Streamlining outpatient prescription dispensing utilizing prescriber order entry. *Ashp Midyear Clinical Meeting* 1991;26: Database: IPA.

Exclude - Not a Primary Study

Segars, Solomon DK, Gourley DR, et al. Implementation of a commercially available clinical decision support system to decrease adverse drug events: an exploratory descriptive analysis. *Ashp Summer Meeting* 2003;60: Database: IPA.

Exclude - Not a Primary Study

Seger AC, Gandhi TK, Hope C, et al. Development of a computerized adverse drug event (ADE) monitor in the outpatient setting. *Advances in Patient Safety: From Research to Implementation* 2005;2: Concepts and Methodology: Grey Lit.

Exclude - Not a Primary Study

Seger AC, Jha AK, Bates DW. Adverse drug event detection in a community hospital utilizing computerised medication and laboratory data. *Drug Saf* 2007;30(9):817-24.

Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Seger DL. Computerized POE: Changing roles for the clinical pharmacist. *J Am Pharm Assoc (Wash)* 1999;39(5):710 Database: IPA.

Exclude - No Outcomes of Interest

Seger DL, Gandhi TK, Leape LL, et al. Pilot study to improve medication prescribing in outpatients. *Ashp Midyear Clinical Meeting* 1999;34: Database: IPA.

Exclude - Not a Primary Study

Selam JL. Implantable insulin pumps: a major piece of computerized diabetes therapy. *Hormone & Metabolic Research Supplement* 1990;24:144-54. Database: Ovid MEDLINE(R).

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sellman JS, Decarolis D, Schullo-Feulner A, et al. Information resources used in antimicrobial prescribing. *J Am Med Inform Assoc* 2004;11(4):281-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Sellors C, Levine M, Sellors J, et al. Computer networking to enhance pharmacist-physician communication: A pilot demonstration project in community settings. *Can Pharm J* 2004;137(8):26-30. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Sengstack PP, Gugerty B. CPOE systems: success factors and implementation issues. *J Healthc Inf Manag* 2004;18(1):36-45. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Senholzi C, Fricker MP. Improving the quality of care: A regional medication-safety effort. *Pharmacol Toxicol* 2002;27(7):341-4. Database: IPA.

Exclude - Not MMIT

Sensmeier J. Statement of the healthcare information and management systems society. *Healthcare Information and Management Systems Society*; 2002.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=29278> Grey Lit.

Exclude - Not a Primary Study

Sensmeier J. The latest? A shift from operations to outcomes. *Nurs Manag (Harrow)* 2009;2-9. Database: BSC.

Exclude - Not MMIT

Sensmeier J, Horowitz JK. Advance care delivery through technology. *Nurs Manag (Harrow)* 2003;34:2-6. Database: BSC.

Exclude - Not a Primary Study

Sensmeier J. Survey says: care, communication enhanced by IT. Nurses report ups and downs of current systems. *Nurs Manag (Harrow)* 2006;Suppl 2(4):6 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Senst BL. Steps in multidisciplinary implementation of `smart pump` technology to optimize patient safety. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Serb C. Small and rural IT challenges. *Hospitals & Health Networks*; 2006.

http://www.hhnmag.com/hhnmag/jsp/articledisplay.jsp?dcrpath=HHNMAG/PubsNewsArticle/data/2006August/0608HHN_FEA_RuralIT_1&domain=HHNMAG Grey Lit.

Exclude - Not a Primary Study

Serna J, Sanchez M, Pintor R, et al. Imputation of consumption to patient in three different clinical units with automated dispensing systems. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Seroussi B, Bouaud J, Dreau H, et al. ASTI: a guideline-based drug-ordering system for primary care. *Studies in Health Technology & Informatics* 2001;84(Part 1):528-32. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Seroussi B, Bouaud J, Chatellier G. Modeling patient-specific therapeutic strategy in the guideline-based management of a chronic disease. *Studies in Health Technology & Informatics* 2003;95:537-42. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Seroussi B, Bouaud J. Reminder-based or on-demand decision support systems: a preliminary study in primary care with the management of hypertension. *Studies in Health Technology & Informatics* 2004;101:142-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Serpa MD, Ting J, Lee MP, et al. Computerized ambulatory care pharmacy information system for direct order entry by prescribers. *Am J Hosp Pharm* 1990;47(2):361-3. Database: IPA.

Exclude - Not a Primary Study

Seto A, Huh J, Chau M. Development of an electronic pharmacy patient profiling system in the era of computerized physician order entry. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Shabot MM, LoBue M, Chen J. Wireless clinical alerts for physiologic, laboratory and medication data. *Proceedings / AMIA 2000*;789-93. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shabot M M, LoBue M, Chen J. Wireless clinical alerts for critical medication, laboratory and physiologic data. In Maui, USA: IEEE; 2000. p. 111.2000225139512

Database: Compendex.

Exclude - Not a Primary Study

Shachak A, Hadas-Dayagi M, Ziv A, et al. Primary care physicians' use of an electronic medical record system: a cognitive task analysis. *J Gen Intern Med* 2009;24(3):341-8.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Shaeffer GK, Reiss KJ, Wood DP. Steps in implementing technology. *ASHP Annual Meeting* 1996;53: Database: IPA.

Exclude - Not a Primary Study

Shah A. Automated dispensing - practical tips on managing implementation. *Hospital Pharmacist* 2004;11(5):198-200. Database: IPA.

Exclude - Not a Primary Study

Shalo S. The cyber six: ePharma's hot growth spots. *Pharmaceutical Executive* 2002;22(1):72-4. Database: IPA.

Exclude - Not MMIT

Shamliyan TA, Duval S, Du J, et al. Just what the doctor ordered. Review of the evidence of the impact of computerized physician order entry system on medication errors. *Health Serv Res* 2008;43(1 Part 1):32-53. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shane R. CPOE: the science and the art. *Am J Health Syst Pharm* 2003;60:1273-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shannon T, Ratchford A, Southward D, et al. The development of a computerised equipment and drug calculator for use in resuscitation. *Emergency Medicine Journal* 2002;19(3):215-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Shapiro J, Bessette M, Levine SR, et al. HandiStroke: A handheld tool for the emergent evaluation of acute stroke patients. *Acad Emerg Med* 2003;10(12):1325-8. Database: CINAHL.

Exclude - Not a Primary Study

Shapiro RS. Legal bases for the control of analgesic drugs. *J Pain Symptom Manage* 1994;9(3):153-9. Database: Sociological Abstracts.

Exclude - Not MMIT

Shaw NT, Mador RL, Ho S, et al. Understanding the impact on intensive care staff workflow due to the introduction of a critical care information system: a mixed methods research methodology. *Studies in Health Technology & Informatics* 2009;143:186-91. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shea HE, Preston C, Hudson S. Electronic patient diaries in a clinical trial - The holistic approach. *Drug Inf J* 2004;38(3):225-38. Database: Embase Sept 22-09.

Exclude - Not MMIT

Shea S, DuMouchel W, Bahamonde L. A Meta-analysis of 16 randomized controlled trials to evaluate computer-based clinical reminder systems for preventive care in the ambulatory setting. *J Am Med Inform Assoc* 1996;3(6):399-409. Exclude - Not a Primary Study

Shea SB, Brown FA, Flowers WP, et al. Automated medication dispensing strategy. *Ashp Midyear Clinical Meeting* 2003;38. Database: IPA.

Exclude - Not a Primary Study

Shebl NA, Franklin BD, Barber N. Clinical decision support systems and antibiotic use. *Pharmacy World & Science* 2007;29(4):342-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sheehan B, Stetson P, Wilcox A, et al. Sociotechnical analysis of a neonatal ICU in the context of CPOE. *AMIA Annual Symposium* 2008; PMID:18999093 OVID MEDLINE.

Exclude - No Outcomes of Interest

Sheehan B, Chused A, Graham PL, III, et al. Frequency and types of alerts for antibiotic prescribing in a neonatal ICU. *Studies in Health Technology & Informatics* 2009;146:521-5. PMID:19592898 OVID MEDLINE.

Exclude - Not MMIT

Sheehan VC, Sanborn MD, Dillon T. Pharmacist response to advanced decision support medication alerts. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Sheiner LB, Halkin H, Peck C, et al. Improved computer-assisted digoxin therapy. A method using feedback of measured serum digoxin concentration. Ann Intern Med 1975;82(5):619-27. Database: IPA.

Exclude - No Outcomes of Interest

Shek C, Sittig DF, Kuperman GJ, et al. Pharmacists' satisfaction with a physician order entry (POE) system. Ashp Midyear Clinical Meeting 1999;34: Database: IPA.

Exclude - Not a Primary Study

Sheppard L C, Kouchoukos N T, Kirklin J W. Surgical intensive care process control system. In New York, NY, USA: IEEE; 1969. p.113-7.114054

Database: Inspec.

Exclude - Not MMIT

Sherr J, Tamborlane WV. Past, present, and future of insulin pump therapy: A better shot at diabetes control. Mt Sinai J Med 2008;75:352-61. Database: Embase Sept 22-09.

Exclude - Not MMIT

Shieh JS, Fu M, Huang SJ, et al. Comparison of the applicability of rule-based and self-organizing fuzzy logic controllers for sedation control of intracranial pressure pattern in a neurosurgical intensive care unit. IEEE Trans Biomed Eng 2006;53(8):1700-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shieh S C, Lin C C, Yang T F and others. Using RFID technology on clinic's pharmacy operation management and development of intelligent medicine dispensing cabinet. In Singapore, Singapore: Inst. of Elec. and Elec. Eng. Computer Society; 2008. p.2006-9.20091311989136

Database: Compendex.

Exclude - Not a Primary Study

Shiffman RN. Towards effective implementation of a pediatric asthma guideline: integration of decision support and clinical workflow support. Proceedings - the Annual Symposium on Computer Applications in Medical Care 1994;797-801. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shiffman RN, Liaw Y, Brandt CA, et al. Computer-based guideline implementation systems: A systematic review of functionality and effectiveness. J Am Med Inform Assoc 1999;6(2):104-14. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Shimada K, Takada H, Mitsuyama S, et al. Drug-recommendation system for patients with infectious diseases. AMIA Annual Symposium Proceedings 2005;1112 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shindo A, Matsuda A, Tani S, et al. Construction of a safety management system for drug use by using an RFID tag. *Studies in Health Technology & Informatics* 2006;122:770 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shirley KL. Analysis of medication administration time and the effects of an automated distribution system. *Ashp Midyear Clinical Meeting* 1998;33: Database: IPA.

Exclude - Not a Primary Study

Shklovskiy-Kordi N, Zingerman B, Rivkind N and others. Multimedia case history as a tool for management of patients and clinical trials. In New York, NY, United states: Institute of Electrical and Electronics Engineers Computer Society; 2003. p. 269.2003347605196 Database: Compendex.

Exclude - Not a Primary Study

Shojania KG. Safe medication prescribing and monitoring in the outpatient setting. *Can Med Assoc J* 2006;174(9):1257-8. Database: PsycINFO.

Exclude - Not a Primary Study

Shojania KG, Jennings A, Mayhew A, et al. The effects of on-screen, point of care computer reminders on processes and outcomes of care. (Review). *The Cochrane Database of Systematic Reviews* 2009;(3):CD001096 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Shojania KG, Jennings A, Mayhew A, et al. Effect of point-of-care computer reminders on physician behaviour: A systematic review. *Can Med Assoc J* 2010;182(5):E216-E225 PMID:20212028 OVID MEDLINE.

Exclude - Not a Primary Study

Short D, Frischer M, Bashford J. Barriers to the adoption of computerised decision support systems in general practice consultations: A qualitative study of GPs' perspectives. *Int J Med Inf* 2004;73(4):357-62. Exclude - Not MMIT

Shortliffe EH. CPOE and the facilitation of medication errors. *Journal of Biomedical Informatics* 2005;38(4):257-8. 2005339301072

Database: Compendex.

Exclude - Not a Primary Study

Shudai M, Fujii K, Kuroda J, et al. Construction and evaluation of new anti-cancer drug prescription support system to promote proper use of anticancer drugs. *Gan to Kagaku Ryoho [Japanese Journal of Cancer & Chemotherapy]* 2008;35(10):1717-20. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Shulman LN, Miller R, Ambinder E, et al. Principles of safe practice using an oncology EHR system for chemotherapy ordering, preparation, and administration, part 2 of 2. *Journal of Oncology Practice* 2008;4(5):254-7. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Shulman LN. Principles of safe practice using an oncology EHR system for chemotherapy ordering, preparation, and administration, part 1 of 2. *Journal of Oncology Practice* 2008;4(4):203-6. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Shulman R, Jani Y. Comparison of supplementary prescribers' and doctors' compliance with guidelines for drug dosing in haemofiltration on an intensive care unit. *Pharmaceutical Journal* 2005;274(7346):492-3. Database: Embase Sept 22-09.
Exclude - Not MMIT

Shulman R, Finney S, O'Sullivan C, et al. Tight glycaemic control: A prospective observational study of a computerised decision-supported intensive insulin therapy protocol. *Critical Care* 2007;11(4):R75 Database: Embase Sept 22-09.
Exclude - Not MMIT

Shultz KH. Medication reconciliation: An electronic solution to improve patient safety. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Shultz WE, Post C. 1993 Barcoding good manufacturing practices. *Wholesale Drugs Magazine* 1993;45(Jun):60-1. Database: IPA.
Exclude - Not a Primary Study

Shumway JM, Jackowitz AI, Abate MA. Analysis of physicians', pharmacists', and nurses' attitudes toward the use of computers to access drug information. *Methods Inf Med* 1990;29(2):99-103. Database: IPA.
Exclude - Not MMIT

Shuttleworth T. Computer-generated medication profile review as replacement for automatic stop orders. *Hosp Pharm* 1996;31(3):214-8. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Shuttleworth TA, Jorgenson JA. Automating drug distribution to save cognitive services at an academic medical center. *ASHP Annual Meeting* 1996;53: Database: IPA.
Exclude - Not a Primary Study

Shuttleworth TA, Ruelle S. Detecting medication errors with automated distribution. *Ashp Midyear Clinical Meeting* 1996;December 1996:P-217R Database: IPA.
Exclude - Not a Primary Study

Shuttleworth TA, Jorgenson JA. Implementing a strategic plan to automate drug distribution. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.
Exclude - Not a Primary Study

Sibley C. Treating the common code: Security printing for drugs. *Manufacturing Chemist* 1965;49(Sep):65 Database: IPA.
Exclude - Not a Primary Study

Siegel C, Alexander M. Acceptance and impact of the computer in clinical decisions. *Hospital & Community Psychiatry* 1984;35(8):773-5. Database: PsycINFO.
Exclude - Not a Primary Study

Siegel C, Laska E, fischer S. Clinical review: Computerization and implementation. Journal of Psychiatric Treatment & Evaluation 1983;5(4):321-4. Database: PsycINFO.
Exclude - Not a Primary Study

Siegel C, Alexander MJ, Dlugacz YD, et al. Evaluation of a computerized drug review system: impact, attitudes, and interactions. Computers & Biomedical Research 1984;17(5):419-35. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Siek K A, Khan D U, Ross S E. A usability inspection of medication management in three personal health applications. In Berlin, Germany: Springer Verlag; 2009. p.129-38. Engineering Village Compendex and Inspec.
Exclude - Not MMIT

Siek K A, Khan D, Ross S. A usability inspection of medication management in three personal health applications. In 2001; San Diego, CA: 2009. p.129-38. Grey Lit Source: In: Kurosu, Masaaki; Human centered design: first international conference, HCD 2009; p. 129-138; Berlin; Springer; c2009 Series: LECTURE NOTES IN COMPUTER SCIENCE Number: 2009; NUMB 5619 ISSN: 0302-9743.
Exclude - Not MMIT

Siepmann JP, Bachman JW. HTN-APT: computer aid in hypertension management. J Fam Pract 1987;24(3):313-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Sierra P. Potential and real drug interactions in critically ill patients. Rev Esp Anesthesiol Reanim 1998;44(10):383-7. Database: Embase Sept 22-09.
Exclude - Not MMIT

Sievers MA, Bokhart GH, Hazelton SM. Network implementation of pharmacy automation. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.
Exclude - Not a Primary Study

Sikorskii A, Given C, Given B, et al. Symptom management for cancer patients: A trial comparing two multimodal interventions. J Pain Symptom Manage 2007;34(3):253-64. Database: PsycINFO.
Exclude - Not MMIT

Sikri S, Sansgiry SS, Sanborn MD, et al. Effect of a remote order scanning system on processing medication orders. Am J Health Syst Pharm 2006;63(15):1438-41. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Silone JE. Avoiding common pitfalls in electronic health record implementation. Ocular Surgery News 2009;27(21):68-9.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010465038&site=ehost-live> EBSCO CINAHL.
Exclude - Not a Primary Study

Silva AEBC, Miasso A, Oliveira R, et al. The process of drug dispensing and distribution at four Brazilian hospitals: A multicenter descriptive study. *Latin American Journal of Pharmacy* 2008;27(3):446-53. Database: Embase Sept 22-09.

Exclude - Not MMIT

Silva MM, Mendonca T, Esteves S. Personalized neuromuscular blockade through control: clinical and technical evaluation. *IEEE Conference Proceedings* 2008;2008:5826-9.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Silverman JB, Stapinski C, Bates DW, et al. Adverse drug event prevention program (ADEPP): Putting the literature into action. *Ashp Midyear Clinical Meeting* 2001;36:

Database: IPA.

Exclude - Not a Primary Study

Silverman JB, Faria CE, Hartman CA, et al. The implementation of barcode technology in an investigational drug service. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Silverman P, Fanikos J, DePeiro D, et al. Computer medication order entry intervention to control physician prescribing. *Ashp Midyear Clinical Meeting* 1997;32: Database: IPA.

Exclude - Not a Primary Study

Silvern DA, Tsunekage T, Scannell S, et al. Assisted drug risk management using computer-controlled infusion pumps and a programmable bedside monitor. *J Clin Eng* 1989;14(5):381-9. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Silverstein S. Barriers to computerized prescribing. *JAMA* 1998;280(6):516-7. Database: IPA.

Exclude - Not a Primary Study

Silvester BV, Carr SJ. A shared electronic health record: lessons from the coalface. *Med J Aust* 2009;190(Suppl 11):S113-S116 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Simborg DW, Derewicz HJ. A highly automated hospital medication system. Five years' experience and evaluation. *Ann Intern Med* 1975;83(3):342-6. Database: Ovid

MEDLINE(R).

Exclude - Not MMIT

Simborg DW. Medication prescribing on a university medical service-the incidence of drug combinations with potential adverse interactions. *Johns Hopkins Med J* 1976;139(1):23-6.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Siminoff LA. A decision aid to assist in adjuvant therapy choices for breast cancer.

Psychooncology 2006;15(11):1001-13. Database: Embase Sept 22-09.

Exclude - Not MMIT

Simko RJ, Stanek EJ. Treatment patterns for heart failure in a primary care environment. *Am J Manag Care* 1997;3(11):1779-80. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Simmer TF, Steffen WM. Impact of a pharmacy and nursing service survey after a one month trial with the Pyxis Medstation. *Ashp Midyear Clinical Meeting* 1993;28: Database: IPA.

Exclude - Not a Primary Study

Simon SR, Chan KA, Soumerai SB, et al. Potentially inappropriate medication use by elderly persons in U.S. Health Maintenance Organizations, 2000-2001. *J Am Geriatr Soc* 2005;53(2):227-32. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Simon SR, Smith DH, Feldstein AC, et al. Computerized prescribing alerts and group academic detailing to reduce the use of potentially inappropriate medications in older people. *J Am Geriatr Soc* 2006;54(6):963-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Simon SR, McCarthy ML, Kaushal R, et al. Electronic health records: which practices have them, and how are clinicians using them? *J Eval Clin Pract* 2008;14(1):43-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Simon SR, Soran CS, Kaushal R, et al. Physicians' use of key functions in electronic health records from 2005 to 2007: a statewide survey. *J Am Med Inform Assoc* 2009;16(4):465-70. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Simonaitis L, Belsito A, Overhage JM. Enhancing an ePrescribing system by adding medication histories and formularies: the Regenstrief Medication Hub. *AMIA 2008;Annual Symposium Proceedings* 2008:677-81. PMID:18999153 OVID MEDLINE.

Exclude - No Outcomes of Interest

Simoni-Wastila L, Tompkins C. Balancing diversion control and medical necessity: the case of prescription drugs with abuse potential. *Substance Use & Misuse* 2001;36(9-10):1275-96. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Simonian AI. Building a better drug database. *Am J Health Syst Pharm* 2004;61(4):397-400. Database: IPA.

Exclude - Not a Primary Study

Simonian AI. Medication order communication using fax and document-imaging technologies. *Am J Health Syst Pharm* 2008;65(6):570-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Simonian MM. Electronic prescribing systems in pediatrics: The rationale and functionality requirements. *Pediatrics* 2007;119(6):e1413-e1422 Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Simonsmeier LM. Pharmacy fails to warn aspirin-sensitive patient. *Pharm Times* 2004;71(1):100 Database: IPA.

Exclude - Not a Primary Study

Simonson JK, Anzisi L, Hill AA, et al. Screening, counseling and monitoring drug therapies including those outside the comfort zone. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Simpson N. Bar coding for patient safety. HIMSS report. Healthcare Information and Management Systems Society; 2001.

<http://www.himss.org/asp/ContentRedirector.asp?ContentID=29276> Grey Lit.

Exclude - Not a Primary Study

Simpson N. Exploring the impact of the FDA proposed rule for bar code labeling. Report. Healthcare Information and Management Systems Society; 2003.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=31671> Grey Lit.

Exclude - Not a Primary Study

Simpson RL. Information technology. Nurses -- yes, nurses -- improve physician order entry. Nurs Manag (Harrow) 2006;31(9):22-3. Database: CINAHL.

Exclude - Not a Primary Study

Simpson WR. Bedside barcode verification at point of care. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Sims N. IV infusion pumps: The missing component in bar code medication administration systems. Hosp Pharm 2003;38(11):S26-S27 Database: IPA.

Exclude - Not MMIT

Sinclair-Pingel J, Wright L, Grisso AG, et al. Implementation of standardized concentrations for continuous infusions in a pediatric hospital. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Singer MV, Haft R, Barlam T, et al. Vancomycin control measures at a tertiary-care hospital: impact of interventions on volume and patterns of use. Infect Control Hosp Epidemiol 1998;19(4):248-53. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Sinnett MJ. Computerized physician order entry: Montefiore experience. Ashp Midyear Clinical Meeting 2001;36: Database: IPA.

Exclude - Not a Primary Study

Sinnett MJ, Cusick DD. Pharmacy services in a fully implemented CPOE organization? Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Sintchenko V, Coiera E, Iredell JR, et al. Comparative impact of guidelines, clinical data, and decision support on prescribing decisions: an interactive web experiment with simulated cases. J Am Med Inform Assoc 2004;11(1):71-7. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Sintchenko V, Magrabi F, Tipper S. Are we measuring the right end-points? Variables that affect the impact of computerised decision support on patient outcomes: a systematic review. *Medical Informatics & the Internet in Medicine* 2007;32(3):225-40. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sintchenko V, Coiera E, Gilbert GL. Decision support systems for antibiotic prescribing. *Current Opinion in Infectious Diseases* 2008;21(6):573-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sipkoff M. New dispensing, bar-coding tools aim for enhanced safety. *Drug Topics* 2003;147: Database: IPA.

Exclude - Not a Primary Study

Sipkoff M. Automated dispensing system offers savings, greater safety. *Drug Topics* 2004;148: Database: IPA.

Exclude - Not a Primary Study

Sipkoff M. Exclusive survey shows retail R.Ph.s have less to spend than their hospital peers but are happier with what they have. *Drug Topics* 2004;148(16):37-8. Database: IPA.

Exclude - Not a Primary Study

Sipkoff M. Will e-prescribing increase generic utilization? Experts believe so. *Drug Topics* 2010;10S Database: IPA.

Exclude - Not a Primary Study

Sirajuddin AM, Osheroff JA, Sittig DF, et al. Implementation pearls from a new guidebook on improving medication use and outcomes with clinical decision support. Effective CDS is essential for addressing healthcare performance improvement imperatives. *J Healthc Inf Manag* 2009;23(4):38-45. PMID:19894486 OVID MEDLINE.

Exclude - No Outcomes of Interest

Siska MH. Pharmacy Informatics and risk management: Minimizing negative events and consequences. *Ashp Summer Meeting* 2009;65: Database: IPA.

Exclude - Not a Primary Study

Sittig DF, Stead WW. Computer-based physician order entry: the state of the art. [Review]. *J Am Med Inform Assoc* 1994;1(2):108-23. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sittig DF, Campbell E, Guappone K, et al. Recommendations for monitoring and evaluation of in-patient Computer-based Provider Order Entry systems: results of a Delphi survey. *AMIA* 2007;671-5. Background. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Sittig DF, Wright A, Osheroff JA, et al. Grand challenges in clinical decision support. *Journal of Biomedical Informatics* 2008;41(2):387-92. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Siwicki B. Electronic prescriptions: just what the doctor ordered. *Health Data Manag* 1995;3(10):62-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sjoberg B, Backstrom T, Arvidsson LB, et al. Design and implementation of a point-of-care computerized system for drug therapy in Stockholm metropolitan health region--Bridging the gap between knowledge and practice. *Int J Med Inf* 2007;76(7):497-506. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Skinner R, Caldwell J, Vitale P. Computerized screening for appropriate dosing of renally eliminated medications. *Proceedings - the Annual Symposium on Computer Applications in Medical Care* 1994;971 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Skledar S, Guttendorf S, Kowiatek J, et al. Standardizing infusion concentrations for safety: Process and results. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Skylis IM, Shoup LK, Woroniecki CL. Evaluation of a computerized pharmacy patient charge system which uses bar code technology. *Ashp Midyear Clinical Meeting* 1988;23: Database: IPA.

Exclude - Not a Primary Study

Sleath BL, Facchinetti NJ. Impact of computerization on New England hospital pharmacy personnel. *Am J Hosp Pharm* 1990;47(9):2070-2. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Slee A, Farrar K, Hughes D. Introducing automation reduces medication errors. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Slee A, Farrar K, Hughes D. Implementing an automated dispensing system. *Pharmaceutical Journal* 2002;268(7191):437-8. Database: IPA.

Exclude - Not MMIT

Sloot PMA. A Grid-based HIV expert system. *J Clin Monit Comput* 2005;19(4-5):263-78. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Smeets R, Talmon J, Meinardi H, et al. Validating a decision support system for anti-epileptic drug treatment. Part II: adjusting anti-epileptic drug treatment. *Int J Med Inf* 1999;55(3):199-209. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Smeets R, Talmon J, Meinardi H, et al. Validating a decision support system for anti-epileptic drug treatment. Part I: initiating anti-epileptic drug treatment. *Int J Med Inf* 1999;55(3):189-98. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Smestad NR. Bar-coding outpatient prescriptions and inpatient vials for accuracy=safier care. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Smith DS, Park JY, Musen MA. Therapy planning as constraint satisfaction: a computer-based antiretroviral therapy advisor for the management of HIV. AMIA Proceedings 1998;Annual Symposium:627-31. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Smith JE, Meyer GE. Organizational approach to implementing bar code technology in a university hospital. Am J Hosp Pharm 1987;44(Mar):572-3. Database: IPA.

Exclude - Not MMIT

Smith M, Dang D, Lee J. E-prescribing: Clinical implications for patients with diabetes. Journal of Diabetes Science & Technology 2009;3(5):1215-8. PMID:20144439 OVID MEDLINE.

Exclude - No Outcomes of Interest

Smith MW, Joseph GJ. Pharmacy data in the VA health care system. MED CARE RES REV 2003;60(Suppl 3):123S Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Smith S, Tang TB, Terry JG, et al. Development of a miniaturised drug delivery system with wireless power transfer and communication. IET Nanobiotechnology 2007;1(5):80-6.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Smith SA, Shah ND, Bryant SC, et al. Chronic care model and shared care in diabetes: randomized trial of an electronic decision support system. Mayo Clin Proc 2008;83(7):747-57. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Smith TM. Optimizing the use of electronic medication administration records in a rural hospital. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Smithline N, Christenson E. Physicians and the Internet: understanding where we are and where we are going. J Ambulatory Care Manage 2001;24(4):39-53. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Smits H. Network medicine: Epilepsy and pharmacy. Pharm Weekbl 1983;118(14): Database: Embase Sept 22-09.

Exclude - Not MMIT

Smorra SR, Migaki G, Stoner S. Implementation and evaluation of an internet based medication formulary. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Snoff CA. Project HITCH June meeting: Health information technology and the safety net. U.S. Department of Health and Human Services; 2007.

<http://www.ncsl.org/print/health/forum/Cheryl.pdf> Grey Lit.

Exclude - Not a Primary Study

Snyder-Halpern R, Wagner MC. Planning for implementation of a vendor-based clinical information system. Case study. *Comput Nurs* 2000;18(1):9-12. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Snyder R, Weston M, Fields W, et al. Computerized provider order entry system field research: The impact of contextual factors on study implementation. *Int J Med Inf* 2006;75(10-11):730-40. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Snyder RA, Fields WL. Measuring hospital readiness for information technology (IT) innovation: A multisite study of the Organizational Information Technology Innovation Readiness Scale. *J Nurs Meas* 2006;14(1):45-55. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Snyder RA, Abarca J, Meza JL, et al. Reliability evaluation of the adapted national coordinating council medication error reporting and prevention (NCC MERP) index. *Pharmacoepidemiology & Drug Safety* 2007;16(9):1006-13. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Sodorff MM, Galusha CT. Impact of point-of-care barcode technology on medication error prevention. *Advances in Pharmacy* 2004;2(2):168-74. Database: IPA.

Exclude - Not a Primary Study

Sojer R, Burkle T, Criegee-Rieck M, et al. Knowledge modelling and knowledge representation in hospital information systems to improve drug safety. *Journal on Information Technology in Healthcare* 2005;4(1):165-70. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Solet DJ, Norvell JM, Rutan GH, et al. Lost in translation: challenges and opportunities in physician-to-physician communication during patient handoffs. *Acad Med* 2005;80(12):1094-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Solomon DH, Shmerling R, Schur P, et al. A computer based intervention to reduce unnecessary serologic testing. *J Rheumatol* 1999;26(12):2578-84. Database: Embase Sept 22-09.

Exclude - Not MMIT

Solomon I, Maharshak N, Chechik G, et al. Applying an artificial neural network to warfarin maintenance dose prediction. *Israel Medical Association Journal (IMAJ)* 2004;6(12):732-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Solomon MR. Information technology to support self-management in chronic care: A systematic review. *Disease Management and Health Outcomes* 2008;16(6):391-401. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Solovy A, Hoppszallern S, Brown SB. 100 Most Wired: being most wired in a down economy. *Hospitals & Health Networks* 2009;83(7):28-37. PMID:19708612 OVID MEDLINE.

Exclude - Not a Primary Study

Soman S, Zasuwa G, Yee J. Automation, decision support, and expert systems in nephrology. *Advances in Chronic Kidney Disease* 2008;15(1):42-55. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Somerville KT, Phillips K. The MedPlanner: An Internet-based dynamic medication education tool. *Advances in Pharmacy* 2003;1(2):112-20. Database: IPA.

Exclude - Not MMIT

Song L, Chui WC, Lau CP, et al. A 3-year study of medication incidents in an acute general hospital. *J Clin Pharm Ther* 2008;33(2):109-14. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Sood SP. Telepharmacy and ePharmacy: Siamese or discrete? *International Journal of Healthcare Technology and Management* 2008;9(5-6): Database: Embase Sept 22-09.

Exclude - Not MMIT

Soper J, Chan GT, Skinner JR, et al. Management of oral anticoagulation in a population of children with cardiac disease using a computerised system to support decision-making. *Cardiol Young* 2006;16(3):256-60. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Soran CS, Simon SR, Jenter CA, et al. Do electronic health records create more errors than they prevent? *AMIA 2008;Annual:Symposium* PMID:18998877 OVID MEDLINE.

Exclude - Not MMIT

Sorrell,S. Enhancement of adherence to tricyclic antidepressants by computerized supervision. *Illinois Institute of Technology* Editor. 1983. Grey Lit.

Exclude - No Outcomes of Interest

Souder DE, Gouveia WA, Sheretz D, et al. Computer-assisted intravenous admixture system. *Am J Hosp Pharm* 1973;30(Nov):1015-20. Database: IPA.

Exclude - Not a Primary Study

Soulliard D, Hong M, Saubermann L. Development of a pharmacy-managed medication dictionary in a newly implemented computerized prescriber order-entry system. *Am J Health Syst Pharm* 2004;61(6):617-22. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Soulliard DE, Hong MK, Sauberman LA. Development of a pharmacy managed medication dictionary in a newly implemented computerized provider order entry system. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Soumerai SB, McLaughlin TJ, Avorn J. Quality assurance for drug prescribing. [Review]. *Qual Assur Health Care* 1990;2(1):37-58. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sovie MJ, Caba JM. EMAR & barcoding: the good, the bad, & the reality. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Sowan AK, Mohamed G, Soeken K, et al. A comparison of medication administrations errors using CPOE orders vs. handwritten orders for pediatric continuous drug infusions. AMIA 2006;1105 Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Sowan AK. The effect of a computerized physician order entry system on managing continuous infusion medications at a pediatric intensive care unit. University of Maryland 2009;Ph.D thesis: Database: CINAHL.

Exclude - Theses

Spader C. Cool tools: cutting-edge gadgets sharpen nurses' efficiency. Nursing Spectrum -- DC, Maryland & Virginia Edition 2009;19(11):20-1.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010398314&site=ehost-live> EBSCO CINAHL.

Exclude - Not a Primary Study

Spahni S, Lovis C, Ackermann M, et al. Securing chemotherapies: fabrication, prescription, administration and complete traceability. Studies in Health Technology & Informatics 2007;129(Pt:2):2-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Spaniel F, Vohlidka P, Hrdlicka J, et al. ITAREPS: Information technology aided relapse prevention programme in schizophrenia. Schizophr Res 2008;98(1-3):312-7. Database: PsycINFO.

Exclude - Not MMIT

Spann B. Enhanced real time reporting of adverse drug reactions. ASHP Annual Meeting 2001;58: Database: IPA.

Exclude - Not a Primary Study

Spaulding T J, Furukawa M F, Raghu T S. Performance impacts of medication management systems: Process matters. In Koloa, Kauai, HI, United states: IEEE Computer Society; 2010. Engineering Village Compendex and Inspec.

Exclude - No Outcomes of Interest

Speizer RD, Guharoy R, Rodgers L, et al. Re-engineering of a medication distribution system in the emergency department in a tertiary care institution via implementation of an automated dispensing system. Ashp Midyear Clinical Meeting 2000;35: Database: IPA.

Exclude - Not a Primary Study

Spencer MD, Stein GC, Cantwell KM. Hot topics from the government affairs division. Ashp Summer Meeting 2005;62: Database: IPA.

Exclude - Not a Primary Study

Sperzel WD, Broverman CA, Kapusnik-Uner JE, et al. The need for a concept-based medication vocabulary as an enabling infrastructure in health informatics. AMIA Proceedings 1998;Annual Symposium:865-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Spetz J, Keane D. Evaluating success. Strategies and challenges for understanding IT implementation in a rural hospital. *J Healthc Inf Manag* 2009;23(1):62-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Spetz J, Keane D. Information technology implementation in a rural hospital: A cautionary tale. *J Healthc Manag* 2009;54(5):337-47. OVID EMBASE.

Exclude - Not MMIT

Sphiris N, Nanas S, Paraskevopoulou H, et al. A decision support software package for medical treatment of I.C.U. patients. *Studies in Health Technology & Informatics* 1997;43 Pt B:911-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Spiro RF. Electronic-prescribing: Will long-term care be ready? *Consultant Pharmacist* 2006;20(8): Database: Embase Sept 22-09.

Exclude - Not MMIT

Spiro RF. Electronic prescribing in the long term care setting. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Spiro RF. Electronic prescribing in long-term care: an overview of five pilot projects. *Consultant Pharmacist* 2008;23(1):16-26. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Spitzer K, Thie A, Kunze K. Spontaneous subarachnoid haemorrhage: expert system for appraisal of the prognosis and computer-supported decision for therapy. *J Neurol* 1988;235(6):335-42. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Spooner SA, Council on Clinical Information Technology AAoP. Special requirements of electronic health record systems in pediatrics. *Pediatrics* 2007;119(3):631-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Spurr C. Case study: provider order entry. *ASHP Annual Meeting* 1997;54: Database: IPA.

Exclude - Not a Primary Study

Spurr C. CPOE - Implications for nursing. *Ashp Summer Meeting* 2002;59: Database: IPA.

Exclude - Not a Primary Study

Squires M, Biesiada D, Fanizza R, et al. New approaches to improving patient safety: strategy, technology and funding. *HEALTHC Q* 2005;8(3):120-2. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Sreih A. Spironolactone: The missing drug in the treatment of patients hospitalized with congestive heart failure. *JCOM* 2004;11(9): Database: Embase Sept 22-09.

Exclude - Not MMIT

Stablein D, Welebob E, Johnson E, et al. Understanding hospital readiness for computerized physician order entry. *Joint Commission Journal on Quality & Safety* 2003;29(7):336-44. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Stapinski CD, Silverman JB, Churchill WW. The clinical utility of an integrated adverse drug event computer alert system. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Starkey C, Michaelis J, de Lusignan S. Computerised systematic secondary prevention in ischaemic heart disease: a study in one practice. *Public Health* 2000;114(3):169-75. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Starnes HM. Use of technology in a high volume ambulatory pharmacy. *Ashp Midyear Clinical Meeting* 1995;30: Database: IPA.
Exclude - Not a Primary Study

Staroselsky M, Volk LA, Tsurikova R, et al. An effort to improve electronic health record medication list accuracy between visits: patients' and physicians' response. *Int J Med Inf* 2008;77(3):153-60. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Steckel SD, Nillni MH, Prince CA, et al. Challenges of building a pharmacy computer system at a cancer research center. *Ashp Midyear Clinical Meeting* 1994;29: Database: IPA.
Exclude - Not a Primary Study

Stefanacci R. The long and wired road. *Pharmacy Practice* 1998;19(7):25 Database: IPA.
Exclude - Not MMIT

Stefanchik M. Point-of-care information systems: Prioritizing bedside applications. *Comput Healthc* 1987;8(4):42-7. 2889757
Database: Inspec.
Exclude - Not MMIT

Stein GC. Current federal regulatory trends. *Ashp Summer Meeting* 2002;59: Database: IPA.
Exclude - Not a Primary Study

Stein JD, Chui CD. Improving ambulatory pharmacy services through automation. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.
Exclude - Not a Primary Study

Steinbrook R. The (slowly) vanishing prescription pad. *N Engl J Med* 2008;359(2):115-7. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Steinschaden T, Petersson G, Astrand B. Physicians' attitudes towards eprescribing: A comparative web survey in Austria and Sweden. *Inform Prim Care* 2009;17(4):241-8. PMID:20359402 OVID MEDLINE.
Exclude - Unable to Retrieve

Stephenson T. The National Patient Safety Agency. Arch Dis Child 2005;90(3):226-8. Database: CINAHL.
Exclude - Not a Primary Study

Stergachis A, Keene D, Somani S. Informatics for medicines management systems in resource-limited settings. 2008. http://ehealth-connection.org.libaccess.lib.mcmaster.ca/files/conf-materials/Informatics%20for%20Medicines%20Mngmt%20Systems_0.pdf Grey Lit.
Exclude - Not a Primary Study

Sterling J. Recent publications on medications and pharmacy. Hosp Pharm 2007;42(8):768-73. Database: IPA.
Exclude - Not MMIT

Sterling J. Recent publications on medications and pharmacy. Hosp Pharm 2008;43(1):66-71. Database: IPA.
Exclude - Not MMIT

Sterns AA. Curriculum design and program to train older adults to use personal digital assistants. Gerontologist 2005;45(6):828-34. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Sterns A, Mayhorn C. Persuasive pillboxes: Improving medication adherence with personal digital assistants. In Eindhoven, Netherlands: Springer Verlag; 2006. p.195-8.20062910005554
Database: Compendex.
Exclude - No Outcomes of Interest

Stettheimer T. Too steep to climb: proposed meaningful-use regs ask too much, too soon of providers. Mod Healthc 2010;40(4):25
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010541386&site=ehost-live> EBSCO CINAHL.
Exclude - Not a Primary Study

Steurbaut K, Van Hoecke S, Colpaert K, et al. Use of web services for computerized medical decision support, including infection control and antibiotic management, in the intensive care unit. Journal of Telemedicine & Telecare 2010;16(1):25-9. PMID:20086264 OVID MEDLINE.
Exclude - No Outcomes of Interest

Stevens BP. Kids' healthcare gets technology boost. Pharmacy Practice News 2003;30(1):42-3. Database: IPA.
Exclude - Not a Primary Study

Stevens BP. Best practices in CPOE rack up impressive gains in safety. Pharmacy Practice News 2009;30(3):9 Database: IPA.
Exclude - Not a Primary Study

Stevens R. Pharmacy aspects of the Exmouth Care card trial. Pharmaceutical Journal 1991;246:359-61. Database: IPA.
Exclude - Not MMIT

Stevens RG. Computer technology: Patient retained medical records. *Pharmaceutical Journal* 1984;232:570-3. Database: IPA.

Exclude - Not MMIT

Stevens R G, Crabble A M. Computerised patient retained records: A working system. In Weybridge, Surrey, UK: *British Journal Healthcare Comput*; 1985. p.250-61.2605411

Database: Inspec.

Exclude - No Outcomes of Interest

Stevenson JE, Nilsson GC, Petersson GI, et al. Nurses' experience of using electronic patient records in everyday practice in acute/inpatient ward settings: a literature review. *Health Informatics Journal* 2010;16(1):63-72.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010632373&site=ehost-live> EBSCO CINAHL.

Exclude - Not MMIT

Stewart KJ. Respiratory care in the computer age. *Respir Care* 2004;49(4):361-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Stiefel FC, Huyse FJ, Sollner W, et al. Operationalizing integrated care on a clinical level: the INTERMED project. *Med Clin North Am* 2006;90(4):713-58. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Exclude - Not a Primary Study

Stiening KK, Trimble JM, Merryfield DW. Computerized clinical alerts - Reducing the noise. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Stiltner RL, Ploetz PA, Kraus CK. Implementation and evaluation of an automated drug distribution system as a component of the university community clinic pharmacy service models. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.

Exclude - Not a Primary Study

Stockley IH. Computerized automatic warnings about drug interactions are now available. *Br Med J* 1998;314:303 Database: IPA.

Exclude - Not MMIT

Stone BL, Boehme S, Mundorff MB, et al. Hospital admission medication reconciliation in medically complex children: An observational study. *Arch Dis Child* 2010;95(4):250-5.

OVID EMBASE.

Exclude - Not MMIT

Stoner NS, Tanfield CJ, Talbot DC. Computerized prescribing of chemotherapy reduces errors. *Br Med J* 1996;312:707 Database: IPA.

Exclude - Not a Primary Study

Stoop AP, Bal R, Berg M. OZIS and the politics of safety: using ICT to create a regionally accessible patient medication record. *Int J Med Inf* 2007;76:Suppl-35 Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Exclude - No Outcomes of Interest

Storz E. S. Implementation of electronically submit reports of adverse events to the Boards of Health in Germany. *Pharmazeutische Industrie* 2007;69(4): Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Stout C. *Current Advances in Inpatient Psychiatric Care*, Greenwood Press;1993. Use of computers in the psychiatric setting. Database: PsycINFO.
Exclude - Not MMIT.

Strack T, Bergeler J, Beyer J, et al. Computer assisted conventional insulin therapy. *Life Support Syst* 1985;3 Suppl 1:568-72. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Straight M. One strategy to reduce medication errors: the effect of an online continuing education module on nurses' use of the Lexi-Comp feature of the Pyxis MedStation 2000. *CIN COMPUT INFORM NURS* 2008;26(1):23-30. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Strain JJ, Caliendo G, Himelein C. Using computer databases to predict and avoid drug-drug interactions in the cancer patient requiring psychotropics. *Psychooncology* 1998;7(4):321-32. Database: CINAHL.
Exclude - Not MMIT

Strange M. Why should the National Programme for IT include the electronic transmission of prescriptions? *BJHC & IM* 2004;21(2):30-2. Database: CINAHL.
Exclude - Not MMIT

Stratton TP, Worley MM, Schmidt M, et al. Implementing after-hours pharmacy coverage for critical access hospitals in northeast Minnesota. *Am J Health Syst Pharm* 2008;65(18):1727-34. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Stroecker S. Polished automation tools allow patient safety to shine. *Nurs Manag (Harrow)* 2003;34(12):34-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Stuart GW, Laraia MT, Ornstein SM, et al. An interactive voice response system to enhance antidepressant medication compliance. *Top Health Inf Manage* 2003;24(1):15-20. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Stuhlmiller B. Databases can help lighten formulary complexities. *Managed Healthcare* 1999;9(Jul):46-7. Database: IPA.
Exclude - Not a Primary Study

Stumpf PG. Practical solutions to improve safety in the obstetrics/gynecology office setting and in the operating room. *Obstetrics & Gynecology Clinics of North America* 2008;35(1):19-35. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Sturgeon J. Do-it-yourself bar-coding vs. outsourcing. *Pharmacy Practice News* 2005;29(12):5 Database: IPA.
Exclude - Not a Primary Study

Sturgeon JL, Williams M, van Servellen G. Efficacy of green tea in the prevention of cancers. *Nursing and Health Sciences* 2009;11(4):436-46. OVID EMBASE.

Exclude - Not a Primary Study

Sturm C, Boehmer R, Monch V, et al. Implementation of an automated antibiotic utilization review system in a German university hospital. *Ashp Midyear Clinical Meeting* 1995;30:

Database: IPA.

Exclude - Not a Primary Study

Sturzlinger H, Hiebinger C, Pertl D, et al. Computerized physician order entry effectiveness and efficiency of electronic medication ordering with decision support systems. Cologne: German Agency for Health Technology Assessment at the German Institute for Medical Documentation and Information (DAHTA DIMDI) 2000; Grey Lit -english summary saved for system, full-text german version in grey lit folder.

Exclude - Not MMIT

Stuurman-Bieze AG, van den Berg PB, Tromp TF, et al. Computer-assisted medication review for asthmatic patients as a basis for intervention. Constructing and validating an algorithmic computer instrument in pharmacy practice. *Pharmacy World & Science* 2004;26(5):289-96. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Stuurman-Bieze AG, de Boer WO, Kokenberg ME, et al. Complex pharmaceutical care intervention in pulmonary care: part A. The process and pharmacists' professional satisfaction. *Pharmacy World & Science* 2005;27(5):376-84. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Suave LS, Jadalla AA. Integrated heart failure telemonitoring system for homecare. *Cardiology Journal* 2010;17(2):200-4. OVID EMBASE.

Exclude - No Outcomes of Interest

Sue CA. The glory and chaos of selecting and implementing automated dispensing technologies. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Sugiyama T, Takada M, Shibayama T, et al. Computerized checking system for powder drug dispensing utilizing data from a prescription order entry system. *Japanese Journal of Hospital Pharmacy* 1998;24(2):179-85. Database: IPA.

Exclude - Not a Primary Study

Sugiyama T, Shibayama T, Takagi N, et al. Computerized provision system for the instruction of medication to patients associated with a prescription order entry system. *Japanese Journal of Hospital Pharmacy* 1998;24(3):292-300. Database: IPA.

Exclude - Not a Primary Study

Exclude - Not a Primary Study

Sugiyama T, Shibayama T, Takano Y, et al. Development of prescription checking system: Subsystem in dispensing support system. *Japanese Journal of Hospital Pharmacy* 1999;25(1):88-97. Database: IPA.

Exclude - Not a Primary Study

Sugiyama T, Niwa T, Takagi N, et al. Investigating the usefulness of a prescription checking system in risk management. Japanese Journal of Pharmaceutical Health Care & Sciences 2003;29(1):73-6. Database: IPA.

Exclude - Not MMIT

Sullivan JE, Buchino JJ. Medication errors in pediatrics--the octopus evading defeat. J Surg Oncol 2004;88(3):182-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sullivan M. Integration of a decision support system in a large academic medical center. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.

Exclude - Not a Primary Study

Summerhayes M. Oncology pharmacy at the M.D. Anderson Cancer Center. Hospital Pharmacist 2000;7(Feb):49-51. Database: IPA.

Exclude - Not a Primary Study

Summers R. A methodology for the design, implementation and evaluation of intelligent systems with an application to critical care medicine. Biological and Medical Sciences 1992; Grey Lit.

Exclude - Not MMIT

Sun CC, Chang P. Automatic appropriateness-evaluation and consultation-suggestion of antibiotics usage via mining of previous prescription data in hospital information system. AMIA 2005;1125 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Sun PR, Wang BH, Wu F. A new method to guard inpatient medication safety by the implementation of RFID. J Med Syst 2008;32(4):327-32. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Sung TY, Hung FH, Chiu HW. Implementation of an integrated drug information system for inpatients to reduce medication errors in administering stage. Conference Proceedings: 2008;2008:743-6. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Sunol R, Vallejo P, Groene O, et al. Implementation of patient safety strategies in European hospitals. Quality & Safety in Health Care 2009;18:Suppl-61 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Suplido S, Sommers S. Impact of computerized physician order entry prescribing on medication errors in the outpatient setting. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Sura ME. Measuring impact: Data collection, analysis, monitoring and response. Ashp Summer Meeting 2009;65: Database: IPA.

Exclude - Not a Primary Study

Susca P. Hospital puts drugs behind bars. ID Systems 1989;9(7): 1990010101565 Database: Compendex.

Exclude - Not MMIT

Susman E. E-prescribing improves the bottom line in pharmacy. *Pharmacy Practice News* 1970;31(6):70 Database: IPA.
Exclude - Not a Primary Study

Susman E. Bar codes working their way into hospitals to combat errors. *Pharmacy Practice News* 2004;31(7):23 Database: IPA.
Exclude - Not a Primary Study

Sutter TL, Wellman GS, Mott DA, et al. Evaluation of access discrepancies associated with an automated storage and distribution cabinet. *Ashp Midyear Clinical Meeting* 1996;31: Database: IPA.
Exclude - Not a Primary Study

Suttman H. Computer-controlled infusion. *Anaesthetist* 1988;37(8): Database: Embase Sept 22-09.
Exclude - Not MMIT

Sutton DR, Taylor P, Earle K. Evaluation of PROforma as a language for implementing medical guidelines in a practical context. *BMC Med Inform Decis Mak* 2006;6:20 Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Svoboda E. Health Care Takes the Stage. *Fast Company* 2010;(145):48-50.
<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=49228531&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.
Exclude - Not a Primary Study

Swanson D. Automated dispensing - an overview of the types of systems available. *Hospital Pharmacist* 2004;11(2):66-8. Database: IPA.
Exclude - Not a Primary Study

Swanson TA, Blair P, Madigan L. Reduction in medication errors through redesign of the medication use system. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
Exclude - Not a Primary Study

Sward K, Orme J, Jr., Sorenson D, et al. Reasons for declining computerized insulin protocol recommendations: application of a framework. *Journal of Biomedical Informatics* 2008;41(3):488-97. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Sweeney J. Electronic patient medical records in primary care: Changes to GP terms of service. *Humanities, psychology and social science* 2002; Grey Lit.
Exclude - No Outcomes of Interest

Sweet BV, Tamer HR, Siden R, et al. Improving investigational drug service operations through development of an innovative computer system. *Am J Health Syst Pharm* 2008;65(10):969-73. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Sweidan M, Reeve JF, Brien JA, et al. Quality of drug interaction alerts in prescribing and dispensing software. *Med J Aust* 2009;190(5):251-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Syed S, Salah K, Saleh M. Medication reconciliation: A process of germination in healthcare. Ashp Summer Meeting 2004;64: Database: IPA.

Exclude - Not a Primary Study

Szabo J. Electronic health records. Health system hopes to build physician loyalty with EHR program. Hospitals & health networks / AHA 2010;84(1):15 OVID EMBASE.

Exclude - Not a Primary Study

Szandzik EG. Bar-code technology and hospital pharmacy practice. J Pharm Technol 1988;4(Sep-Oct):182-7. Database: IPA.

Exclude - Unable to Retrieve

Szeinbach S, Seoane-Vazquez E, Parekh A, et al. Dispensing errors in community pharmacy: perceived influence of sociotechnical factors. Int J Qual Health Care 2007;19(4):203-9.

Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Szeinbach SL, Taylor TH, Gillenwater EL. Automated dispensing technologies: Effect on managed care. J Manag Care Pharm 1995;1(Sep-Oct):121-7. Database: IPA.

Exclude - Not a Primary Study

Szilagy PG, Rodewald LE, Savageau J, et al. Improving influenza vaccination rates in children with asthma: A test of a computerized reminder system and an analysis of factors predicting vaccination compliance. Pediatrics 1992;90(6):871-5. Database: CINAHL.

Exclude - Not MMIT

Tabor RA, Kolar GR, Martinez BE. Decentralization of pharmacy services. Ashp Midyear Clinical Meeting 1990;25: Database: IPA.

Exclude - Not a Primary Study

Tackley R. Integrating anaesthesia and intensive care into the National Care Record. Br J Anaesth 2006;97(1):69-76. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Taft LM, Evans RS, Shyu CR, et al. Countering imbalanced datasets to improve adverse drug event predictive models in labor and delivery. Journal of Biomedical Informatics 2009;42(2):356-64. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Tahil FA. Clinical informatics. BluefishRx prescription writer. J PSYCHIATR PRACT 2005;11(1):58-61. Database: CINAHL.

Exclude - Not MMIT

Tait DI. New developments and future direction of prison pharmacy. Hospital Pharmacist 1999;6(Apr):100-3. Database: IPA.

Exclude - Not MMIT

Takada M, Honda T, Nakata I, et al. Benefits and risks of electronic patient records on the works of the Pharmaceutical Department in the National Hospital Organization, Kyoto Medical Center. IRYO - Japanese Journal of National Medical Services 2005;59(5):255-61. Database: Embase Sept 22-09.

Exclude - Not MMIT

Takahashi E, Yoshida K, Izuno T, et al. Protocol care for hypertension supported by an expert system. *Medinfo 1995*;8 Pt 2:954 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Takahashi Y, Goto H, Saito T. Health Care System using Face Robot and Tele-operation via Internet. In Singapore, Singapore: Nanyang Technological University; 2002. p.1025-30.2004208164395

Database: Compendex.

Exclude - Not a Primary Study

Takayama K, Seino T, Sugiura M, et al. Development of an inspection-supporting system using drug images for unit dose packages. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2001;121(11):821-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Talbert DA, Roush E, Velazco L, et al. An incremental pharmacy informatics model for use in a rural hospital. *AMIA 2005*;1131 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tallis RC, Edmond ED, O'Halloran A. Computer system to prevent prescription of contraindicated drugs. *British Journal of Pharmaceutical Practice* 1984;6(Jul):223-8.

Database: IPA.

Exclude - Unable to Retrieve

Tallis RC. Medicines for the elderly. Part 2. Computer assisted prescribing. *Pharmaceutical Journal* 2004;235: Database: IPA.

Exclude - Unable to Retrieve

Tamai IY, Rubenstein LZ, Josephson KR, et al. Impact of computerized drug profiles and consulting pharmacist on outpatient prescribing patterns: Clinical trial. *Drug Intelligence & Clinical Pharmacy* 1987;21(Nov):890-5. Database: IPA.

Exclude - Not MMIT

Tamblyn R. Improving patient safety through computerized drug management: the devil is in the details. *Healthcare Papers* 1982;5(3):52-68. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tamblyn RM, Jacques A, Laprise R, et al. The Office of the Future Project: the integration of new technology into office practice. Academic detailing through the super highway. Quebec Research Group on Medication Use in the Elderly. *Clinical Performance & Quality Health Care* 1997;5(2):104-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tamuz M, Harrison MI. Improving patient safety in hospitals: Contributions of high-reliability theory and normal accident theory. *Health Serv Res* 2006;41(4:Pt:2):t-76

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tan HH, Soon T, Chan RK. Quality improvement: experience of a sexually transmitted infection clinic in Singapore. *International Journal of STD & AIDS* 2008;19(12):800-4.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tan K, Dear PR, Newell SJ. Clinical decision support systems for neonatal care. *Cochrane Database Syst Rev* 2005;(2):CD004211 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Tan KC, Khor EF, Cai J, et al. Automating the drug scheduling of cancer chemotherapy via evolutionary computation. *Artif Intell Med* 2002;25(2):169-85. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Tan RS, Isaacks S. Computerized records and quality of care. *Annals of Long-Term Care* 1999;7(9):348-53. Database: Embase Sept 22-09.
Exclude - Not MMIT

Tang PC, Young CY. Active Guidelines: integrating Web-based guidelines with computer-based patient records. *AMIA Proceedings* 2000;843-7. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Tang PC, Black W, Buchanan J, et al. PAMF Online: integrating EHealth with an electronic medical record system. *AMIA* 2003;644-8. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Tanner TB. ScriptWriter. A relational database to manage outpatient medical treatment. *The Annual Symposium on Computer Applications in Medical Care* 1994;Proceedings:1034 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Tant D. Having practice pharmacists is not only way of reducing prescribing costs. *Br Med J* 1999;318:872 Database: IPA.
Exclude - Not MMIT

Tauman AV, Virata MD, Boyce J. Implementation of a criteria-based anti-infective approval system through computerized physician order entry (CPOE). *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.
Exclude - Not a Primary Study

Tavella A. Safety in cytostatic compounding: The impact of automation. *EJHP Practice* 2009;15(3):58-9. OVID EMBASE.
Exclude - Not a Primary Study

Tavernier A, Coussaert E, d'Hollander A, et al. Model-based pharmacokinetic regulation in computer-assisted anesthesia an interactive system: CARIN. *Acta Anaesthesiol Belg* 1987;38(1):63-8. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Taxis K, Dean B, Barber N. Hospital drug distribution systems in the UK and Germany - A study of medication errors. *Pharmacy World & Science* 1999;21(1):25-31. Exclude - Not MMIT

Taylor B, Dinh M, Kwok R, et al. Electronic interface for emergency department management of asthma: A randomized control trial of clinician performance. *EMA - Emergency Medicine Australasia* 2008;20(1):38-44. Database: Embase Sept 22-09.
Exclude - Not MMIT

Taylor R, Manzo J, Sinnett M. Quantifying value for physician order-entry systems: a balance of cost and quality. *Healthc Financ Manage* 2002;56(7):44-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Taylor SD, Birdwell SW, Schneider DN, et al. Development and validation of criteria to identify patients requiring clinical pharmacist intervention. *ASHP Annual Meeting* 1992;49: Database: IPA.

Exclude - Not a Primary Study

Teagarden JR, Nagle B, Aubert R, et al. Dispensing error rate in a highly automated mail-service pharmacy practice. *Pharmacotherapy* 2005;25(11 I):1629-35. Database: Embase Sept 22-09.

Exclude - Not MMIT

Teall AM. The power of the PDA. *Adv Nurse Pract* 2009;17(8):43-5. PMID:20014663 OVID MEDLINE.

Exclude - Not a Primary Study

Teich J. Electronic prescribing in the ambulatory care environment: Promise, progress, barriers, solutions. *eHealth Initiative*; 2004.
http://ehr.medigent.com.libaccess.lib.mcmaster.ca/assets/collaborate/2004/03/30/NCVHS_eRx_testimony_Mar03.pdf Grey Lit.

Exclude - Not a Primary Study

Teich JM, Spurr CD, Schmiz JL, et al. Enhancement of clinician workflow with computer order entry. *The Annual Symposium on Computer Applications in Medical Care* 1995;Proceedings:459-63. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Teich JM, Petronzio AM, Gerner JR, et al. An information system to promote intravenous-to-oral medication conversion. *AMIA Proceedings* 1999;Annual Symposium:415-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Teich JM, Glaser JP, Beckley RF, et al. The Brigham integrated computing system (BICS): advanced clinical systems in an academic hospital environment. *Int J Med Inf* 1999;54(3):197-208. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Teich JM, Osheroff JA, Pifer EA, et al. Clinical decision support in electronic prescribing: recommendations and an action plan: report of the joint clinical decision support workgroup. *J Am Med Inform Assoc* 2005;12(4):365-76. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Teich J and Marchibroda J. Electronic prescribing: Toward maximum value and rapid adoption. Executive summary. *eHealth Initiative*; 2004.
[http://ehr.medigent.com/assets/collaborate/2004/04/14/eHealth Initiative Electronic Prescribing Report 04.14.04 Executive Summary.pdf](http://ehr.medigent.com/assets/collaborate/2004/04/14/eHealth_Initiative_Electronic_Prescribing_Report_04.14.04_Executive_Summary.pdf) Grey Lit.

Exclude - Not a Primary Study

Teich J, Osheroff J, Pifer E et al. Clinical decision support for electronic prescribing: Recommendations and an action plan. 2005.
<http://www.amia.org/mbrcenter/pubs/docs/cdswhitepaperforhhs-final2005-03-08.pdf> Grey Lit.

Exclude - Not a Primary Study

Telling K. Electronic prescriptions in Sweden: the success story continues. BJHC & IM 2005;22(10):22-4. 8818280
Database: Inspec.

Exclude - Not a Primary Study

Telling K. Healthcare in Sweden. Electronic prescriptions in Sweden: The success story continues. BJHC & IM 2005;22(10):22-4. Database: CINAHL.

Exclude - Not a Primary Study

Teltsch D, Pinelle D, Winslade N, et al. Decision support for community-based empirical antibiotic prescribing. AMIA 2005;1133 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Temkin LA, Welch DW, Mahoney CD. Cost savings from computerization and addition of bar coding capability to a pharmacy purchasing and inventory management system. Ashp Midyear Clinical Meeting 1988;23: Database: IPA.

Exclude - Not a Primary Study

Tempelhof MW. Personal digital assistants: A review of current and potential utilization among medical residents. Teaching & Learning in Medicine 2009;21(2):100-4. Database: CINAHL.

Exclude - Not a Primary Study

Temple J, Ludwig B. Implementation and evaluation of carousel dispensing technology in a university medical center pharmacy. Am J Health Syst Pharm 2010;67(10):821-9.
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010664325&site=ehost-live>; EBSCO CINAHL.

Exclude - Not MMIT

ten Berg MJ, Huisman A, van den Bemt PM, et al. Linking laboratory and medication data: new opportunities for pharmacoepidemiological research. Clinical Chemistry & Laboratory Medicine 2007;45(1):13-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

ter Wee RJ, van der KE, Brenninkmeijer RF, et al. Development of an electronic prescription processing option: an aid for general practice. Br J Gen Pract 1991;41(345):151-4. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tessier A, Bedouch P, Allenet B, et al. Computerized prescriber order entry associated with pharmacist participation in physician round in a French teaching hospital: assessment of pharmacist's interventions. Ashp Summer Meeting 2004;62: Database: IPA.

Exclude - Not a Primary Study

Thall I. RFID and healthcare applications: What comes after bar code? *Hosp Pharm* 2003;38(11):S28-S29 Database: IPA.
Exclude - Not MMIT

Thielke TS. Automation support of patient-focused care. *Top Hosp Pharm Manage* 1994;14(1):53-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Thielke TS. Impact of bar code medication administration. *Hosp Pharm* 2003;38(11):S22-S23 Database: IPA.
Exclude - Unable to Retrieve

Thomas A. Medical perspective on PCOE and safe medication practice. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Thomas AN, Marchant AE, Ogden MC, et al. Implementation of a tight glycaemic control protocol using a web-based insulin dose calculator. *Anaesthesia* 2005;60(11):1093-100. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Thomas AN. Computerised decision support to achieve tight glycaemic control in critical illness. *British Journal of Intensive Care* 2009;19(1):29-35. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Thomas H, Lewis G, Watson M, et al. Computerised Patient-specific Guidelines for Management of Common Mental Disorders in Primary Care: a Randomised Controlled Trial. *Br J Gen Pract* 2004;54:832-7. Exclude - Not MMIT

Thomas MP. Practical pharmacology: Medication errors. *Clin Pediatr (Phila)* 2003;42(4): Database: Embase Sept 22-09.
Exclude - Not MMIT

Thomas N. The role of pharmacoeconomics in disease management. A pharmaceutical benefit management company perspective. *Pharmacoeconomics* 1996;9 Suppl 1:9-15. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Thomas N, Fifer SK. The combination of collaborative drug therapy management and E-prescribing. *Manag Care Interface* 2003;16(1):38-42. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Thomas R, Mitocaru C, Hanlon J. Evaluating E&M coding accuracy of GoCode as compared to internal medicine physicians and auditors. *Perspectives in Health Information Management, CAC Proceedings*, 2008.
http://library.ahima.org/xpedio/groups/public/documents/ahima/bok1_040464.pdf Grey Lit.
Exclude - Not MMIT

Thomas SM, Davis DC. The characteristics of personal order sets in a computerized physician order entry system at a community hospital. *AMIA* 2003;1031 Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Thomas WH. Medicament prescribing, administration and monitoring system. Australian Journal of Hospital Pharmacy 1976;6(4):145-8. Database: IPA.

Exclude - Unable to Retrieve

Thomasset KB, Faris R. Survey of pharmacy services provision in the emergency department. Am J Health Syst Pharm 2003;60(15):1561-4. Database: IPA.

Exclude - Not MMIT

Thompson DA, Duling L, Holzmueller C, et al. Computerized physician order entry, a factor in medication errors: Descriptive analysis of events in the Intensive Care Unit Safety Reporting System. JCOM 2005;12(8):407-12. Database: Embase Sept 22-09.

Exclude - Not MMIT

Thompson K. Assessing bedside barcoding readiness. Ashp Midyear Clinical Meeting 2002;37: Database: IPA.

Exclude - Not a Primary Study

Thomson P, Dowding D, Swanson V, et al. A computerised guidance tree (decision aid) for hypertension, based on decision analysis: development and preliminary evaluation. European Journal of Cardiovascular Nursing 2006;5(2):146-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Thomson R, Robinson A, Greenaway J, et al. Development and description of a decision analysis based decision support tool for stroke prevention in atrial fibrillation. Quality & Safety in Health Care 2002;11(1):25-31. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Thomson RG, Eccles MP, Steen IN, et al. A patient decision aid to support shared decision-making on anti-thrombotic treatment of patients with atrial fibrillation: randomised controlled trial. Quality & Safety in Health Care 2007;16(3):216-23. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Thorne A, Williamson S, Jellison T, et al. Implementation of home-based medication order entry at a community hospital. Am J Health Syst Pharm 2009;66(21):1939-42.

PMID:19850789 OVID MEDLINE.

Exclude - No Outcomes of Interest

Thornton JP, Schumock G, Kanafotska C. Development of a computerized pharmacy therapeutic recommendation tracking program. Top Hosp Pharm Manage 1992;11(4):44-51. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Thornton PD, Simon S, Mathew TH. Towards safer drug prescribing, dispensing and administration in hospitals. J Qual Clin Pract 1999;19(1):41-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Thull B, Rau G, Hanrath P. Clinical decision support: From data to clinical information. In Amsterdam, Neth: IEEE; 1996. p.2162-4.1998194115272

Database: Compendex.

Exclude - Not a Primary Study

Thull B, Janssens U, Rau G, et al. Approach to computer-based medication planning and coordination support in intensive care units. *Technol Health Care* 1997;5(3):219-33. 5711045 Database: Inspec.

Exclude - No Outcomes of Interest

Thursky K. Use of computerized decision support systems to improve antibiotic prescribing. *Expert Review of Antiinfective Therapy* 2006;4(3):491-507. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Thursky KA, Mahemoff M. User-centered design techniques for a computerised antibiotic decision support system in an intensive care unit. *Int J Med Inf* 2007;76(10):760-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tierney JA, Roefaro J, Creamer K, et al. Prevention of potential adverse drug events in a computerized physician ordering system. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Tierney M, McLurg D, Macmillan C. Transferring medication order entry from pharmacists to pharmacy technicians. *Can J Hosp Pharm* 1999;52(4):240-3. Database: IPA.

Exclude - Not MMIT

Tierney WM, Overhage JM, Takesue BY, et al. Computerizing guidelines to improve care and patient outcomes: the example of heart failure. *J Am Med Inform Assoc* 1995;2(5):316-22. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tilson L, Sabra K. Pharmacists' role in the introduction of a clinical information system to an intensive care unit (ICU). *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Tilva SB, Keefer LA, Summerfield MR, et al. A novel process for verification of dispensing accuracy of premixed IV solutions using bi-dimensional bar code technology. *Ashp Summer Meeting* 2002;59: Database: IPA.

Exclude - Not a Primary Study

Tjaden P. A comment on White and Du Mont's 'Visualizing sexual assault: An exploration of the use of optical technologies in the medico-legal contact'. *Soc Sci Med* 2009;68(1):9-11. Scholar's Portal Sociological Abstracts.

Exclude - Not a Primary Study

Tohara D, Miyake H, Shimaoka A, et al. The construction of a checking system for drug-contraindicant disease in order entry system. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

Tokunaga J, Kubota M, Tsuruta S, et al. Support system with a composition analysis for prescription order of injection agents. *Japanese Journal of Pharmaceutical Health Care & Sciences* 2003;29(6):720-4. Database: IPA.

Exclude - Not MMIT

Tolman C, Richardson D, Bartlett C, et al. Structured conversion from thrice weekly to weekly erythropoietic regimens using a computerized decision-support system: a randomized clinical study. *J Am Soc Nephrol* 2005;16(5):1463-70. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Tomasi E, Facchini LA, Maia MD. Health information technology in primary health care in developing countries: a literature review. *Bull World Health Organ* 2004;82(11):867-74.
Database: IPA.
Exclude - Not a Primary Study

Tomita A. Effects of novel infection control system using electronic medical records of Showa University Northern Yokohama Hospital. *Journal of the Showa Medical Association* 2006;65(6): Database: Embase Sept 22-09.
Exclude - Not MMIT

Tooher R, Middleton P, Pham C, et al. A systematic review of strategies to improve prophylaxis for venous thromboembolism in hospitals. *Ann Surg* 2005;241(3):397-415.
Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Torchio M, Molino G, Cavanna A, et al. PEPTY: a knowledge-based program for assisting medical reasoning in peptic diseases. *Computer Methods & Programs in Biomedicine* 1989;28(4):249-56. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Torniainen K. In my opinion: Hospital pharmacists are needed in the development of data systems in Finland. *EJHP Practice* 2009;15(1):11 OVID EMBASE.
Exclude - Not a Primary Study

Toschlog EA, Newton C, Allen N, et al. Morbidity reduction in critically ill trauma patients through use of a computerized insulin infusion protocol: a preliminary study. *Journal of Trauma-Injury Infection & Critical Care* 2007;62(6):1370-5. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Tourville J. How technology is helping children's medical center of Dallas reach zero-error tolerance. *U S Pharmacist* 2003;28(6):80-9. Database: IPA.
Exclude - Not MMIT

Tourville JF, Hetey SK, Eisenwine JE, et al. Role of automation in closing the loop on the medication use process and the redesign of the practice environment at a university affiliated, tertiary care, pediatric teaching hospital. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.
Exclude - Not a Primary Study

Tourville JF, Eisenwine JE, Kromelis MR, et al. Development and testing of a two-way wireless, server-based infusion device, used to decrease medication errors and improve real time documentation. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.
Exclude - Not a Primary Study

Toussi M, Ebrahimi V, Le Toumelin P, et al. An automated method for analyzing adherence to therapeutic guidelines: application in diabetes. *Studies in Health Technology & Informatics* 2008;136:339-44. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Townsend P. Developing a database to manage use of unlicensed medicines. *Hospital Pharmacist* 2006;13(8):299-300. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Tran K, Levy-Santoro M, Kohn B, et al. Medication safety initiative: Monitoring small volume potassium chloride orders. *Ashp Midyear Clinical Meeting* 2007;42. Database: IPA.

Exclude - Not a Primary Study

Travaglia J, Westbrook M, Braithwaite J. Implementation of a patient safety incident management system as viewed by doctors, nurses and allied health professionals. *Health (N Y)* 2009;no. 3(pp. 277-296). Database: Sociological Abstracts.

Exclude - Not MMIT

Treweek S. A new quality improvement study every day? Using QTools to build quality improvement projects around primary care electronic medical record systems. *Studies in Health Technology & Informatics* 2004;107(Pt:2):2-9. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Trice A. Refine prescribing systems. *Hospital Pharmacy Practice* 2004;8:301-3. Database: IPA.

Exclude - Not MMIT

Triller DM, Donnelly J, Ruge J. Travel-related savings through a rural, clinic-based automated drug dispensing system. *J Community Health* 2005;30(6):467-76. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Trindade GR, Anderson LA, Van Pelt RA. Using standard order protocols and pre-printed order forms to enhance pharmacy efficiency. *ASHP Annual Meeting* 1993; Database: IPA.

Exclude - Not a Primary Study

Trivedi MH, Claassen CA, Grannemann BD, et al. Assessing physicians' use of treatment algorithms: Project IMPACTS study design and rationale. *Contemporary Clinical Trials* 2007;28(2):192-212. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Trivedi MH, Daly EJ. Measurement-based care for refractory depression: a clinical decision support model for clinical research and practice. *DRUG ALCOHOL DEPENDENCE* 2007;88:Suppl-71 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Trivedi M, Daly E. Measurement-based care for refractory depression: A clinical decision support model for clinical research and practice. *Drug Alcohol Depend* 2007;88(Suppl 2):S61-S71 Database: PsycINFO.

Exclude - Not a Primary Study

Troester S. Drive nursing activities to the bedside with a closed-loop system. *Nurs Manag (Harrow)* 2006;37(12):18 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Troiano D, Morrison J, Federico F, et al. Safely automating the medication use process. Not as easy as it looks. *J Healthc Inf Manag* 2009;23(4):17-23. PMID:19894482 OVID MEDLINE.

Exclude - No Outcomes of Interest

Trotter A. Computerized physician order entry system in pediatric inpatients: Prevention of medication errors and adverse drug events. *Monatsschrift fur Kinderheilkunde* 2009;157(2): Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Tsai P H, Chuang Y T, Chou T S and others. iNuC: An intelligent mobile nursing cart. In Tianjin, China: IEEE Computer Society; 2009. Engineering Village Compendex and Inspec.

Exclude - No Outcomes of Interest

Tseng SH, Shyong-Jiang DD, Hoi HS, et al. Effect of free treatment and surveillance on HIV-infected persons who have tuberculosis, Taiwan, 1993-2006. *Emerg Infect Dis* 2009;15(2):332-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Tsuchiya F. Medication errors caused by order entry system and prevention measures. In Beijing, China: Springer-Verlag; 2007. p.535-43.9610422

Database: Inspec.

Exclude - Not a Primary Study

Tucker A. NHS ETP pilots: Pharmacy2U consortium's solution goes live. *BJHC & IM* 2002;19(4):30-2. Database: CINAHL.

Exclude - Not MMIT

Tucker WM. When less is more: reducing the incidence of antipsychotic polypharmacy. *J PSYCHIATR PRACT* 2007;13(3):202-4. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Tudiver F. Primary care providers' perceptions of home diabetes telemedicine care in the IDEATel project. *J Rural Health* 2007;23(1): Database: Embase Sept 22-09.

Exclude - Not MMIT

Tudor RS, Hovorka R, Cavan DA, et al. DIAS-NIDDM--a model-based decision support system for insulin dose adjustment in insulin-treated subjects with NIDDM. *Computer Methods & Programs in Biomedicine* 1998;56(2):175-91. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Tulloch J, Evans B. Evaluation of the accuracy of the Saskatchewan Health Pharmaceutical Information Program for determining a patient's medication use immediately before admission. *Can J Hosp Pharm* 2009;62(1):21-7. Database: Embase Sept 22-09.

Exclude - Not MMIT

Tully M. The impact of information technology on the performance of clinical pharmacy services. *J Clin Pharm Ther* 2000;25(4):243-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Tully MP, Seston M, Cantrill J. Motivators and barriers to the implementation of pharmacist-run prescription monitoring and review services in two settings. *Int J Pharm Pract* 2000;8(3):188-97. Database: Embase Sept 22-09.

Exclude - Not MMIT

Tung Y, Duffy L, Gyamfi J, et al. Improvements in immunization compliance using a computerized tracking system for inner city clinics. *Clin Pediatr (Phila)* 2003;42(7):603-11. Database: Embase Sept 22-09.

Exclude - Not MMIT

Turchin A, Gandhi TK, Coley CM, et al. The use of electronic medication reconciliation to establish the predictors of validity of computerized medication records. *Studies in Health Technology & Informatics* 2007;129(Pt:2):2-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Turchin A, Hamann C, Schnipper JL, et al. Evaluation of an inpatient computerized medication reconciliation system. *J Am Med Inform Assoc* 2008;15(4):449-52. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Turisco F and Rhoads J. Equipped for efficiency: Improving nursing care through technology. California HealthCare Foundation; 2008.

<http://www.chcf.org/topics/view.cfm?itemID=133816> Grey Lit.

Exclude - Not a Primary Study

Ubach C, Bate A, Ryan M, et al. Using discrete choice experiments to evaluate alternative electronic prescribing systems. *Int J Pharm Pract* 2002;10(3):191-200. List. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Uddin Z, Bear RA. Public-private partnerships in the Canadian environment: options for hospital pharmacies. *Healthc Manage Forum* 1997;10(4):45-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ukens C. Canadian mail-service R.Ph.s counseling face-to-face. *Drug Topics* 1994;138:39 Database: IPA.

Exclude - Not a Primary Study

Ukens C. RoboR.Ph.: Is community pharmacy ready for automated dispensing? *Drug Topics* 1996;140(8):60-6. Database: IPA.

Exclude - Not a Primary Study

Ukens C. Are you ready? *Drug Topics* 2000;144:34-6. Database: IPA.

Exclude - Not a Primary Study

Ukens C. DEA proposes rule to verify e-scripts of controlled drugs. *Drug Topics* 2001;145:60 Database: IPA.

Exclude - Not a Primary Study

Ukens C. Pediatric groups hail new med safety guidelines. *Drug Topics* 2002;146(14): Database: IPA.

Exclude - Not a Primary Study

Ukens C. Firm offers chains automated Rx reminders. Drug Topics 2004;148(17):56
Database: IPA.
Exclude - Not a Primary Study

Ukens C. Half of independents able to use e-prescribing. Drug Topics 2004;148(17):58
Database: IPA.
Exclude - Not a Primary Study

Ukens C. NABP targets patient safety with new hire. Drug Topics 2004;148(8):34 Database:
IPA.
Exclude - Not a Primary Study

Ukens C. Young entrepreneurs roll out Rx refill technology. Drug Topics 2004;148(3):104
Database: IPA.
Exclude - Not a Primary Study

Underhill JG, Thomas MW. Use of an electronic bulletin board and e-mail system to
facilitate a concurrent drug use evaluation program. Ashp Midyear Clinical Meeting 1995;30:
Database: IPA.
Exclude - Not a Primary Study

Underwood MA. Prime time for pharmacists--telepharmacy is here. Pharm Times
2004;63:22 Database: IPA.
Exclude - Not a Primary Study

Upperman JS, Staley P, Friend K, et al. The introduction of computerized physician order
entry and change management in a tertiary pediatric hospital. Pediatrics 2005;116(5):e634-
e642 Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Urbanski AP. Extended possibilities of pharmaceuticals delivery to patients using
dematerialized prescriptions. Studies in Health Technology & Informatics 2004;103:28-35.
Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Urquhart C, Currell R, Grant M, et al. Nursing record systems: effects on nursing practice
and healthcare outcomes. Cochrane Database Syst Rev2009;Issue 1:
<http://www.mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD002099/frame.html>
Grey Lit.
Exclude - Not MMIT

Urquhart C, Currell R, Grant M, et al. Nursing record systems: Effects on nursing practice
and healthcare outcomes. Cochrane Database Syst Rev2009;(4): Grey Lit.
Exclude - Not a Primary Study

Vadher B, Patterson DL, Leaning M. Evaluation of a decision support system for initiation
and control of oral anticoagulation in a randomised trial. BMJ 1997;314(7089):1252-6.
Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Vadher BD. Validation of an algorithm for oval anticoagulant dosing and appointment
scheduling. Clin Lab Haematol 1995;17(4): Database: Embase Sept 22-09.
Exclude - Not MMIT

Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. *Clinical & Laboratory Haematology* 1997;19(3):203-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Vaida A, Kercher L. Practical tools for medication safety in acute care. *Journal of the American Pharmacists Association: JAPhA* 2003;43(5:Suppl 1):S48-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Vaida AJ. Observations on the proposed FDA bar coding rule. *Hosp Pharm* 2003;38(11):S7-S8 Database: IPA.

Exclude - Not MMIT

Vaidya V, Sowan AK, Mills ME, et al. Evaluating the safety and efficiency of a CPOE system for continuous medication infusions in a pediatric ICU. *AMIA* 2006;1128 Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Valenzuela Gamez JC, Lopez Gonzalez AM, Pedraza Cezon LA, et al. A project for the implementation of a unit-dose drug dispensation system in an intensive care unit. *Farmacia Hospitalaria* 2005;HOSP.(5):318-22. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Vallee-Smejda S, Hahn M, Aubin N, et al. Recording practices and satisfaction of hemophiliac patients using two different data entry Systems. *CIN - Computers Informatics Nursing* 2009;27(6):372-8. OVID EMBASE.

Exclude - Not MMIT

Valley SL. Hard and soft options: New technology for pharmacists. *New Jersey Journal of Pharmacy* 1990;1:1-7. Database: IPA.

Exclude - No Outcomes of Interest

Van Cleave J, Leslie LK. Approaching ADHD as a chronic condition: implications for long-term adherence. *Journal of Psychosocial Nursing & Mental Health Services* 2008;46(8):28-37. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Van de Castle B, Kim J, Pedreira MLG, et al. Information technology and patient safety in nursing practice: an international perspective. *Int J Med Inf* 2004;73(7-8):607-14. 8164061 Database: Inspec.

Exclude - Not MMIT

van den Bemt PM, Egberts TC, de Jong-van den Berg LT, et al. Drug-related problems in hospitalised patients. *Drug Saf* 2000;22(4):321-33. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

van der Sijs H, Aarts J, Vulto A, et al. Overriding of drug safety alerts in computerized physician order entry. *J Am Med Inform Assoc* 2006;13(2):138-47. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

van der Sijs H, Lammers L, van den Tweel A, et al. Time-dependent drug-drug interaction alerts in care provider order entry: software may inhibit medication error reductions. *J Am Med Inform Assoc* 2009;16(6):864-8. PMID:19717806 OVID MEDLINE.

Exclude - Not MMIT

van der Sijs H, Bouamar R, van Gelder T, et al. Functionality test for drug safety alerting in computerized physician order entry systems. *Int J Med Inf* 2010;79(4):243-51. OVID EMBASE.

Exclude - Not MMIT

van der Sijs H, van Gelder T, Vulto A, et al. Understanding handling of drug safety alerts: a simulation study. *Int J Med Inf* 2010;79(5):361-9. OVID EMBASE.

Exclude - Not MMIT

van der Westerlaken MM, Zoer J, Flier-van Vuuren M. Using bar codes in medicine distribution and administration. *Ejhp Science* 2006;12(5):60-2. Database: IPA.

Exclude - Not a Primary Study

van Gruting CW, de Gier JJ. Medication assistance: the development of drug surveillance and drug information in The Netherlands. *Ann Pharmacother* 1992;26(7-8):1008-12. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

van Hyfte D, de Vries Robbe P, Tjandra-Maga T, et al. Towards a more rational use of psychoactive substances in clinical practice. *Pharmacopsychiatry* 2001;34(1):13-8. Database: PsycINFO.

Exclude - Not a Primary Study

van Mil JW, Dudok van Heel MC, Boersma M, et al. Interventions and documentation for drug-related problems in Dutch community pharmacies. *Am J Health Syst Pharm* 2001;58(15):1428-31. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

van Mil JW. Pharmaceutical care in community pharmacy: practice and research in the Netherlands. *Ann Pharmacother* 2005;39(10):1720-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

van Onzenoort HA, van de Plas A, Kessels AG, et al. Disappointing results of bar-code verification. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

van Roon EN, Langendijk PN, Flikweert S, et al. Weighing interactions and actions. Structured assessment of drug-drug interactions. *Pharm Weekbl* 2003;138(24):850-5. Database: IPA.

Exclude - Not MMIT

van Roon EN, Flikweert S, le Comte M, et al. Clinical relevance of drug-drug interactions : a structured assessment procedure. *Drug Saf* 2005;28(12):1131-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

van Rosse F, Maat B, Rademaker CM, et al. The effect of computerized physician order entry on medication prescription errors and clinical outcome in pediatric and intensive care: a systematic review. *Pediatrics* 2009;123(4):1184-90. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Van Schaik P, Flynn D, Van Wersch A, et al. The acceptance of a computerised decision-support system in primary care: A preliminary investigation. *Behaviour & Information Technology* 2004;23(5):321-6. Database: BSC.

Exclude - Not MMIT

van Tilburg CM, Leistikow IP, Rademaker CM, et al. Health Care Failure Mode and Effect Analysis: a useful proactive risk analysis in a pediatric oncology ward. *Quality & Safety in Health Care* 2006;15(1):58-63. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Van Wijk BLG, Klungel O, Heerdink E, et al. Initial non-compliance with antihypertensive monotherapy is followed by complete discontinuation of antihypertensive therapy. *Pharmacoepidemiology and Drug Safety* 2006;15(8):587-93. Database: Embase Sept 22-09.

Exclude - Not MMIT

Van Wyk JT, van Wijk MA. Assessment of the possibility to classify patients according to cholesterol guideline screening criteria using routinely recorded electronic patient record data. *Studies in Health Technology & Informatics* 2002;93:39-46. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Van D, V, McRae A, Wiseman M, et al. Successful introduction of Tallman letters to reduce medication selection errors in a hospital network. *Journal of Pharmacy Practice and Research* 2008;38(4):263-6. Database: Embase Sept 22-09.

Exclude - Not MMIT

Vander Linde JB, Nordberg M, West N, et al. Improving anticoagulation management with a multidisciplinary anticoagulation team. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Vander Linde JB, Sypula DL. Medication order verification at a remote hospital pharmacy. *Ashp Midyear Clinical Meeting* 2002;37: Database: IPA.

Exclude - Not a Primary Study

Vander Linde JB, Cook C, Manthey C. Improving medication safety with a health system medication safety committee. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.

Exclude - Not a Primary Study

VanDerLinde LP, Scott TS. Concurrent A.D.R./D.U.E. monitoring in a tertiary care hospital via computerized therapeutic risk assessment (TRAC) algorithms. *ASHP Annual Meeting* 1991;48: Database: IPA.

Exclude - Not a Primary Study

VanDerLinde LP, Phillips JO. Problem-oriented pharmacotherapy: role of information technology. *Ashp Midyear Clinical Meeting* 1992;27: Database: IPA.

Exclude - Not a Primary Study

Varveris DA, Morton NS. Target controlled infusion of propofol for induction and maintenance of anaesthesia using the paedfusor: an open pilot study. *Paediatr Anaesth* 2002;12(7):589-93. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Veach M. AHRQ Study shows need for thorough testing of CPOE. *Healthc Financ Manage* 2005;59(5):25-6. Database: BSC.
Exclude - Not MMIT

Vecchione A. Automation giants upgrade dispensing products. *Hospital Pharmacist Report* 1995;9:45 Database: IPA.
Exclude - Not a Primary Study

Vecchione A. Link or sink: Hospitals take action to integrate IS. *Hospital Pharmacist Report* 1995;9:1-10. Database: IPA.
Exclude - Not a Primary Study

Vecchione A. Bar coding: new future for some old technology. *Hospital Pharmacist Report* 1999;13:45 Database: IPA.
Exclude - Not a Primary Study

Vecchione A. Tracking med errors. *Hospital Pharmacist Report* 1999;13:30-2. Database: IPA.
Exclude - Not a Primary Study

Vecchione A. New VA initiatives serve to enhance patient safety. *Hospital Pharmacist Report* 2000;14:54-5. Database: IPA.
Exclude - Not a Primary Study

Velotta CL. Usability of computers for medication order entry by home healthcare nurses. *University of Kentucky* 2003; Ph D 2003; Database: CINAHL.
Exclude - Theses

Veltri MA, Glenn DJ, Gladwell T. Development and implementation of a pharmacy based, automatic dose standardization program for pediatrics. *Ashp Midyear Clinical Meeting* 1997;32: Database: IPA.
Exclude - Not a Primary Study

Venkatraman R, Durai R. Errors in medicine administration: how can they be minimised? *Journal of Perioperative Practice* 2008;18(6):249-53. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Venot A. Electronic prescribing for the elderly: will it improve medication usage?. [Review]. *Drugs & Aging* 1999;15(2):77-80. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Ver-Bruggen S. New diagnosis. *Brand* 2005;4(3):10-7. 2005299215133
Database: Compendex.
Exclude - Not MMIT

Vera-Donoso CD, Llopis B, Oliver F, et al. Selective chemoprophylaxis guided by multifactorial analysis in superficial bladder cancer. *Eur Urol* 1990;17(3):219-22. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Verstappen SM, Jacobs JW, van der Veen MJ, et al. Intensive treatment with methotrexate in early rheumatoid arthritis: aiming for remission. *Computer Assisted Management in Early Rheumatoid Arthritis (CAMERA, an open-label strategy trial)*. *Ann Rheum Dis* 2007;66(11):1443-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Verzija JM. TOP elaborates complex medication schedules. Drug compliance and HIV patients. *Pharm Weekbl* 1999;134(12): Database: Embase Sept 22-09.

Exclude - Not MMIT

Vess J. Implementation of a computer assisted treatment planning and outcome evaluation system in a forensic psychiatric hospital. *Psychiatric Rehabilitation Journal* 2001;25(2):124-32. Database: PsycINFO.

Exclude - Not a Primary Study

Vigoda MM, Lubarsky DA. Failure to recognize loss of incoming data in an anesthesia record-keeping system may have increased medical liability. *Anesthesia & Analgesia* 2006;102(6):1798-802. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Vigoda MM, Gencorelli FJ, Lubarsky DA. Discrepancies in medication entries between anesthetic and pharmacy records using electronic databases. *Anesthesia & Analgesia* 2007;105(4):1061-5. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Vilella A. The role of mobile phones in improving vaccination rates in travelers. *Prev Med* 2004;38(4): Database: Embase Sept 22-09.

Exclude - Not MMIT

Villani A, Malfatto G, Della RF, et al. Disease management for heart failure patients: role of wireless technologies for telemedicine. The ICAROS project. *G Ital Cardiol* 2007;(Rome) (2):107-14. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Vincent WR, Martin CA, Winstead PS, et al. Effects of a pharmacist-to-dose computerized request on promptness of antimicrobial therapy. *J Am Med Inform Assoc* 2009;16(1):47-53. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Virk P, Bates DW, Halamka J, et al. Analyzing transaction workflows in an ePrescribing system. *AMIA* 2006;1129 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Vishwanath A, Brodsky L, Shaha S, et al. Patterns and changes in prescriber attitudes toward PDA prescription-assistive technology. *Int J Med Inf* 2009;78(5):330-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Visscher S, Lucas PJ, Schurink CA, et al. Modelling treatment effects in a clinical Bayesian network using Boolean threshold functions. *Artif Intell Med* 2009;46(3):251-66. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Visser J. [A quarter of a century of computer-assisted anticoagulant treatment]. [Dutch]. *Ned Tijdschr Geneesk* 1997;141(1):55-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Vitello SJ, Kellick KA, Pudhorodsky TG, et al. Implementation and promotion of a computerized provider order entry program at a Department of Veterans Affairs medical center. *Ashp Midyear Clinical Meeting* 1993;28: Database: IPA.

Exclude - Not a Primary Study

Vivian JC. E-signatures and pharmacy practice. *U S Pharmacist* 2001;26:50-3. Database: IPA.

Exclude - Not MMIT

Vivian JC. Rogue internet pharmacies. *U S Pharmacist* 2009;34(1): Database: Embase Sept 22-09.

Exclude - Not MMIT

Vlachogiannis JG. Expert medication system using a C language integrated production system. *Engineering Intelligent Systems for Electrical Engineering and Communications* 2006;14(3):137-45. 9279218

Database: Inspec.

Exclude - No Outcomes of Interest

Vlahavas I, Pomportsis A, Sakellariou I. IDIS-KS: an intelligent drug information system as a knowledge server. *Studies in Health Technology & Informatics* 1997;43 Pt A:368-72.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Vlahovic-Palcevski V, Bergman U. Quality of prescribing for the elderly in Croatia-computerized pharmacy data can be used to screen for potentially inappropriate prescribing. *Eur J Clin Pharmacol* 2004;60(3):217-20. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Vogelzang M, Zijlstra F, Nijsten MW. Design and implementation of GRIP: a computerized glucose control system at a surgical intensive care unit. *BMC Med Inform Decis Mak* 2005;5:38 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Vogelzang M, Loeff BG, Regtien JG, et al. Computer-assisted glucose control in critically ill patients. *Intensive Care Med* 2008;34(8):1421-7. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Voit DT, McDonnell A, Delgado-Flores JO, et al. Evaluation of hydroxyurea prescribing in a large academic teaching hospital affiliated with a major cancer center. *Ashp Midyear Clinical Meeting* 2007;42: Database: IPA.

Exclude - Not a Primary Study

Vollmer W, Kirshner M, Peters D, et al. Use and impact of an automated telephone outreach system for asthma in a managed care setting. *Am J Manag Care* 2006;12(12):725-33.

Exclude - Not MMIT

Vora S, Verber L, Potts S, et al. Effect of a novel birth intervention and reminder-recall on on-time immunization compliance in high-risk children. *Human Vaccines* 2009;5(6):395-402. PMID:19029825 OVID MEDLINE.

Exclude - Not MMIT

Voytovich RM, Outten RL. An aminoglycoside monitoring service in a community hospital using a microcomputer. *Hosp Pharm* 1987;22(3):267-72. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Vrabel RB, Mackowiak L. Automation in pharmacy: two institutions' experiences with novel distribution systems. *Hosp Formul* 1995;30(2):106-13. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Vrijens B, Belmans A, Matthys K, et al. Effect of intervention through a pharmaceutical care program on patient adherence with prescribed once-daily atorvastatin.

Pharmacoepidemiology & Drug Safety 2006;15(2):115-21. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Vrontos J, McAlearney AS, Schneider PJ, et al. The effect of smart pumps on patient safety: a qualitative study. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Vrublevskii OP, Stolbovoi AV. [Choice of antibacterial therapy with the use of a computerized consultation system]. [Russian]. *Anesteziol Reanimatol* 1989;(1):17-9.

Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Vu S, Boyd RA, Jones AT. Quality assurance of automatic substitution program involving angiotensin II receptor Mockers: Focus on medication reconciliation. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Wadhwa S, Saxena A, Wadhwa B. Hospital information management system: an evolutionary knowledge management perspective. *International Journal of Electronic Healthcare* 2007;j.(2):232-60. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wagner MM, Hogan WR. The accuracy of medication data in an outpatient electronic medical record. *J Am Med Inform Assoc* 1996;3(3):234-44. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Waitman LR, Pearson D, Hargrove FR, et al. Enhancing Computerized Provider Order Entry (CPOE) for neonatal intensive care. *AMIA* 2003;1078 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Waitman L, Pearson D, Hargrove F, et al. Enhancing computerized provider order entry (CPOE) for neonatal intensive care. *Proceedings of the AMIA Symposium* 2003;1078

<http://www.amia.org/pubs/proceedings/symposia/2003/462.pdf> Grey Lit.

Exclude - Not a Primary Study

Wakefield DS, Brokel J, Ward MM, et al. An exploratory study measuring verbal order content and context. *Qual Safe Health Care* 2009;18(3):169-73. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Walk SU, Fritz S, Pruszydlo M, et al. Pharmaceutical counselling of drug switches at the interface between primary and tertiary care. *Krankenhauspharmazie* 2009;30(5):218-21. Database: Embase Sept 22-09.

Exclude - Not MMIT

Wallasch T-M. Application examples of an IT-solution for the managed healthcare of patients with chronic headache. *Nervenheilkunde* 2009;28(6): Database: Embase Sept 22-09.

Exclude - Not MMIT

Waller A, Franklin V, Pagliari C, et al. Participatory design of a text message scheduling system to support young people with diabetes. *Health Informatics Journal* 2006;12(4):307-21. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Walsh K, Antony J. Improving patient safety and quality: what are the challenges and gaps in introducing an integrated electronic adverse incident and recording system within health care industry? *Int J Health Care Qual Assur* 2007;20(2-3):107-15. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Walsh P. Prescription and medication fulfilment come of age. *M D Computing* 2000;17(3):45-8. 6667176

Database: Inspec.

Exclude - Not MMIT

Walton R, Ilic Z. Knowledge engineering for drug prescribing guidelines. *Studies in Health Technology & Informatics* 1995;16:75-85. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Walton R, Dovey S, Harvey E, et al. Computer support for determining drug dose: systematic review and meta-analysis. *BMJ* 1999;318(7189):984-90. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Walton R, Dovey S, Harvey E, et al. Information in practice. Computer support for determining drug dose: Systematic review and meta-analysis. *BMJ: British Medical Journal* 1999;318(7189):984-90. Database: CINAHL.

Exclude - Not a Primary Study

Walton RT, Gierl C, Yudkin P, et al. Evaluation of computer support for prescribing (CAPSULE) using simulated cases. *BMJ* 1997;315(7111):791-5. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Walton RT, Gierl C, Yudkin P, et al. Information in practice. Evaluation of computer support for prescribing (CAPSULE) using simulated cases. *BMJ: British Medical Journal* 1997;315(7111):791-5. Database: CINAHL.

Exclude - No Outcomes of Interest

Walton RT, Harvey E, Dovey S, et al. Computerised advice on drug dosage to improve prescribing practice. *Cochrane Database Syst Rev*2001;(1):CD002894 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Wan D. Magic Medicine Cabinet: A situated portal for consumer healthcare. *Lecture Notes in Computer Science* 1999;1707:352-5. 6604232

Database: Inspec.

Exclude - Not a Primary Study

Wang F, Li Z. Development of a clinical pharmacy automatic medicine dispensing and prescription information processing system. In Singapore, Singapore: Nanyang Technol. Univ; 1994. p.852-5.5358817

Database: Inspec.

Exclude - No Outcomes of Interest

Wang JK, Shabot MM, Duncan RG, et al. A clinical rules taxonomy for the implementation of a computerized physician order entry (CPOE) system. *AMIA Proceedings 2002* 2002;860-3. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Wang JK, Herzog NS, Kaushal R, et al. Prevention of pediatric medication errors by hospital pharmacists and the potential benefit of computerized physician order entry. *Pediatrics* 2007;119(1):e77-e85 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wang M, Zao J K, Tsai P H and others. Wedjat: a mobile phone based medicine in-take reminder and monitor. In Piscataway, NJ, USA: IEEE; 2009. p.423-30.10839683

Database: Inspec.

Exclude - Not a Primary Study

Wang WL, Lin CH. Study of two-dimensional barcode prescription system for pharmacists' activities of NHI contracted pharmacy. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2008;128(1):123-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wang X, Hripcsak G, Markatou M, et al. Active computerized pharmacovigilance using natural language processing, statistics, and electronic health records: a feasibility study. *J Am Med Inform Assoc* 2009;16(3):328-37. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ward-McKnight R, West DS. Electronic prescribing reduces errors and improves patient safety. *J Ark Med Soc* 2007;104(3):60-1. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Warmuth M, Moellentin D, Aziz S, et al. The use of a decision support software in pharmacy computer system to proactively prevent adverse drug events. *Ashp Summer Meeting* 2003;60: Database: IPA.

Exclude - Not a Primary Study

Warner A, Menachemi N, Brooks R. Information technologies relevant to pharmacy practice in hospitals: Results of a statewide survey. *Hosp Pharm* 2005;40(3):233-9. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Warner H, Jr., Blue SR, Sorenson D, et al. New computer-based tools for empiric antibiotic decision support. *AMIA Annual Fall Symposium 1997 1997*;Proceedings:238-42. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Warren J, Gadzhanova S, Stanek J, et al. Understanding caseload and practice through analysis of therapeutic state transitions. *AMIA 2005*;784-8. Database: Ovid MEDLINE(R).

Exclude - No Outcomes of Interest

Warren J, Gaikwad R, Mabotuwana T, et al. Utilising practice management system data for quality improvement in use of blood pressure lowering medications in general practice. *N Z Med J* 2008;121(1285): Database: Embase Sept 22-09.

Exclude - Not MMIT

Washington L, Washington T. Implementation of a computerized non-formulary request form at a veterans affairs medical center. *Ashp Midyear Clinical Meeting 2002*;37: Database: IPA.

Exclude - Not a Primary Study

Washington L, Katsh E, Sondheimer N. Dispute resolution: Planning for disputed information in EHRs and PHRS. *Journal of the American Health Information Management Association 2009*;80(11):26-30. OVID EMBASE.

Exclude - Not a Primary Study

Wassenberg MWM. Hypertension management in primary care: Standard care and attitude towards a disease management model. *Neth J Med* 2004;62(10):375-82. Database: Embase Sept 22-09.

Exclude - Not MMIT

Watanabe M, Sugiura M, Seino T, et al. The construction and evaluation of the preventing method for the input mischoice in a prescription order entry system--usefulness of a three-character input and a warning screen display system. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan 2002*;122(10):841-7. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

Waxlax TJ. Evaluation of the appropriate use of computerized physician order entry (CPOE) system for medication orders in an inpatient setting. *Ashp Midyear Clinical Meeting 2007*;42: Database: IPA.

Exclude - Not a Primary Study

Weant KA, Cook AM, Bradley V, et al. Pharmacy resident experience in implementation of computerized prescriber order entry (CPOE). *Ashp Midyear Clinical Meeting 2005*;40: Database: IPA.

Exclude - Not a Primary Study

Weant KA, Cook AM, Armitstead JA. Medication-error reporting and pharmacy resident experience during implementation of computerized prescriber order entry. *Am J Health Syst Pharm* 2007;64(5):526-30. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Weaver R. Computers and medical knowledge: The diffusion of decision support technology. 1991. Grey Lit.

Exclude - No Outcomes of Interest

Webber D, Kotok A. Collaborative e-business and ebXML: A new approach and a standard to make it happen. The real payoff for e-business exists in business-to-business commerce, and as experience from other industries shows, collaboration is key. *Managed Healthcare* 2004;12(1):46 Database: IPA.

Exclude - Not a Primary Study

Weber-Jahnke J. Design of decoupled clinical decision support for service-oriented architectures. *International Journal of Software Engineering & Knowledge Engineering* 2009;19(2):159-83. Database: BSC.

Exclude - Not MMIT

Weber JN, Hatai JK, Carlson T, et al. Surgery/anesthesiology medication perpetual inventory system. *Ashp Midyear Clinical Meeting* 1994;29: Database: IPA.

Exclude - Not a Primary Study

Weber RJ. Core competencies in hospital pharmacy - Medication order review. *Hosp Pharm* 2006;41(3): Database: Embase Sept 22-09.

Exclude - Not MMIT

Weber RJ. Establishing a patient-centered philosophy of pharmacy practice. *Hosp Pharm* 2006;41(1): Database: Embase Sept 22-09.

Exclude - Not MMIT

Weber RJ. Implementing a bar-code medication administration system. *Hosp Pharm* 2008;43(12):1016-23. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Weber RJ. Director's forum. *Hosp Pharm* 2009;44(3): Database: Embase Sept 22-09.

Exclude - Not MMIT

Weber V, White A, McIlvried R. An electronic medical record (EMR)-based intervention to reduce polypharmacy and falls in an ambulatory rural elderly population. *J Gen Intern Med* 2008;23(4):399-404. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Weber V, Bloom F, Pierdon S, et al. Employing the electronic health record to improve diabetes care: a multifaceted intervention in an integrated delivery system. *J Gen Intern Med* 2008;23(4):379-82. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Webster C, Copenhaver J. Structured data entry in a workflow-enabled electronic patient record. *J Med Pract Manage* 2001;17(3):157-61. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Webster NR, Cohen AT. Closed-loop administration of atracurium. Steady-state neuromuscular blockade during surgery using a computer controlled closed-loop atracurium infusion. *Anaesthesia* 1987;42(10):1085-91. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wechsler J. HHS task force: Congress to focus efforts on detecting and reducing medical errors. *Hosp Formul* 2001;36:472-3. Database: IPA.

Exclude - Not a Primary Study

Wechsler J. Drug development gets budgetary boost as regulatory updates start to emerge. *Hosp Formul* 2002;37(2):102-3. Database: IPA.

Exclude - Not a Primary Study

Wechsler J. Drug safety in the limelight. *Pharmaceutical Technology Europe* 2003;15(5):24-32. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Wechsler J. Drug safety in the limelight. *Pharmaceutical Technology* 2003;27(5):24 Database: IPA.

Exclude - Not a Primary Study

Wechsler J. Regulating the Internet. *Pharmaceutical Executive* 2003;19: Database: IPA.

Exclude - Not MMIT

Wechsler J. Electronic information-systems: New systems promote drug development and patient safety. *Pharmaceutical Technology* 2004;28(2): Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Wechsler J. FDA leaders stress science and compliance: Stiffer enforcement of quality standards aims to restore public confidence in agency actions. *BioPharm International* 2009;22(10):16-20. OVID EMBASE.

Exclude - Not a Primary Study

Wechsler J. FDA seeks more timely drug information electronically: The new sentinel system aims to expand access to data on medical product safety and patient effects. *BioPharm International* 2010;23(3):18-9. OVID EMBASE.

Exclude - Not a Primary Study

Weckman HN, Janzen SK. The critical nature of early nursing involvement for introducing new technologies. *Online Journal of Issues in Nursing* 2009;14(2):1-11.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010504667&site=ehost-live> EBSCO CINAHL.

Exclude - No Outcomes of Interest

Weeks G, Stanley L, Vinson M. Automation of the medication history process: A case report. *Hosp Pharm* 2005;40(12):1057-61. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Weeks WB, Bagian JP. Developing a culture of safety in the Veterans Health Administration. *Eff Clin Pract* 2000;3(6):270-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Weerasooriya JM, Jungst JD. Effects of electronic medication administration record implementation on medication error reporting rates. Ashp Midyear Clinical Meeting 2005;40: Database: IPA.

Exclude - Not a Primary Study

Weerasooriya JM, Jungst JD. Effects of off-site pharmacist computerized order entry (COE) on after-hours entry into the pharmacy department. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Wei C, Haque S, Li Z. Design of a Web-based antibiotics management system. In Athens, GA, USA: CSREA Press - Univ. Georgia; 2000. p.129-34.7168854

Database: Inspec.

Exclude - No Outcomes of Interest

Wei X. The method of discovering the irrational use of drugs based on the electronic prescriptions. Pharmaceutical Care and Research 2009;9(2): Database: Embase Sept 22-09.

Exclude - Not MMIT

Weibel KJ, Nolly R, Apple A, et al. Evaluation of using failure mode and effect analysis (FMEA) to analyze i.v. robot technology prior to implementation. Ashp Summer Meeting 2004;61: Database: IPA.

Exclude - Not a Primary Study

Weimar C. Going all-digital is easier said than done. Physician Exec 2009;35(2):20-2.

Database: BSC.

Exclude - Not a Primary Study

Weiner B. A structured care plan for coronary artery bypass surgery: The pharmacist's perspective. P and T 1994;19(10): Database: Embase Sept 22-09.

Exclude - Not MMIT

Weiner M, Callahan CM, Tierney WM, et al. Using information technology to improve the health care of older adults. Ann Intern Med 2003;139(5:Part 2):430-6. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Weingart SN, Cleary A, Seger A, et al. Medication reconciliation in ambulatory oncology. Jt Comm J Qual Patient Saf 2007;33(12):750-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Weingarten S. A measurable difference: Clinical decision support improves care and safety. Healthcare Information and Management Systems Society; 2007.

<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=66792> Grey Lit.

Exclude - Not a Primary Study

Weingarten S. Healthcare information technology and improving the quality of patient care. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Weinhold FE, Osborne J, Renes P, et al. Tracking medication dispensed from a hospital pharmacy. Am J Health Syst Pharm 1999;56:1190 Database: IPA.

Exclude - Not a Primary Study

Weinstock M. Defining meaningful use. H&HN: Hospitals & Health Networks 2010;84(2):
<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010572973&site=ehost-live> EBSCO CINAHL.

Exclude - Not a Primary Study

Weir C, Hoffman J, Nebeker JR, et al. Nurse's role in tracking adverse drug events: the impact of provider order entry. Nurs Adm Q 2005;29(1):39-44. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Weir R. Automated dispensing: A role in the realm of modern Medicines Management? Irish Pharmacy Journal 2003;81(7):300-4. Database: IPA.

Exclude - Not MMIT

Weiss AM. Buying prescription drugs on the internet: promises and pitfalls. Cleve Clin J Med 2006;73(3):282-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Weiss J, Krebs S, Hoffmann C, et al. Survey of adverse drug reactions on a pediatric ward: a strategy for early and detailed detection. Pediatrics 2002;110:254-7. Database: CINAHL.

Exclude - Not MMIT

Weissman A M. Prospective, concurrent and retrospective Drug Use Review: Use of hospital-based minicomputers and distributed data processing. In New York, NY, USA: IEEE; 1982. p.351-4.2015374

Database: Inspec.

Exclude - Not a Primary Study

Weitkamp JH, Ozdas A, LaFleur B, et al. Fluconazole prophylaxis for prevention of invasive fungal infections in targeted highest risk preterm infants limits drug exposure. J Perinatol 2008;28(6):405-11. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wellman GS, Wilson GN, Govern VL, et al. Decision making analysis for programmable multichannel infusion systems. Ashp Midyear Clinical Meeting 1990;25: Database: IPA.

Exclude - Not a Primary Study

Wells S. J. Online management of cardiovascular risk in New Zealand with PREDICT- Getting evidence to the "moment of care". Healthcare Review Online 2005;9(1): Database: Embase Sept 22-09.

Exclude - Not MMIT

Wells LA. Management case study. Ashp Midyear Clinical Meeting 2004;39: Database: IPA.

Exclude - Not a Primary Study

Went K. Safer intensive care prescribing: Engaging users in the implementation of an electronic prescribing system. British Journal of Healthcare Computing and Information Management 2009; Grey Lit.

Exclude - Not a Primary Study

Wenzer HS, Bottger U, Boye N. A socio-technical study of an ubiquitous CPOE-system in local use. *Studies in Health Technology & Informatics* 2006;124:326-32. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wess ML, Schauer DP, Johnston JA, et al. Application of a decision support tool for anticoagulation in patients with non-valvular atrial fibrillation. *J Gen Intern Med* 2008;23(4):411-7. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wess P, Gutzler H. EDP in the hospital pharmacy. Infection and resistance data: Collection, storing and evaluation. *Krankenhauspharmazie* 1990;11:185-8. Database: IPA.

Exclude - No Outcomes of Interest

West D, Szeinbach SL. Pharmacy computers: Vendors' perspective. *Computertalk for the Pharmacist* 1997;17:22-5. Database: IPA.

Exclude - Not a Primary Study

West D, Hastings J, Earley A. Economic justification of the use of dispensing technologies in independent community pharmacies. *American Journal of Clinical Dermatology* 2001;123:17-25. Database: IPA.

Exclude - Not MMIT

West DW, Levine S, Magram G, et al. Pediatric medication order error rates related to the mode of order transmission. *Archives of Pediatrics & Adolescent Medicine* 1994;148(12):1322-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Westbrook JI, Braithwaite J, Georgiou A, et al. Multimethod evaluation of information and communication technologies in health in the context of wicked problems and sociotechnical theory. *J Am Med Inform Assoc* 2007;14(6):746-55. Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Westerlund T. The quality of self-care counselling by pharmacy practitioners, supported by IT-based clinical guidelines. *Pharm World Sci* 2007;29(2):67-72. Database: Embase Sept 22-09.

Exclude - Not MMIT

Westfall JM, Fernald DH, Staton EW, et al. Applied strategies for improving patient safety: a comprehensive process to improve care in rural and frontier communities. *J Rural Health* 2004;20(4):355-62. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Westphal JF, Farinotti R. Computerized physician order entry, a tool for added safety in the hospital. *Presse Med* 2003;32(24):1138-46. Database: Ovid MEDLINE(R).

Exclude - Unable to Retrieve Foreign

White RH, Hong R, Venook AP, et al. Initiation of warfarin therapy: comparison of physician dosing with computer-assisted dosing. *J Gen Intern Med* 1987;2(3):141-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

White RH, Mungall D. Outpatient management of warfarin therapy: comparison of computer-predicted dosage adjustment to skilled professional care. *Ther Drug Monit* 1991;13(1):46-50. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Whiting SO, Gale A. Computerized physician order entry usage in North America: the doctor is in. *HEALTHC Q* 2008;11(3):94-7. Database: Ovid MEDLINE(R).
Exclude - No Outcomes of Interest

Whitman G, Cowell V, Parris K, et al. Prophylactic antibiotic use: hardwiring of physician behavior, not education, leads to compliance. *J Am Coll Surg* 2008;207(1):88-94. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Whittemore K. Controlled substances and electronic prescribing: What you need to know. *Computertalk for the Pharmacist* 2003;23(6):12-3. Database: IPA.
Exclude - Not a Primary Study

Whittemore K. E-prescribing from the physician's perspective. *America's Pharmacist* 2004;127(8):20 Database: IPA.
Exclude - Not a Primary Study

Whittemore K. E-renewals will give pharmacists more control. *America's Pharmacist* 2004;127(9):22 Database: IPA.
Exclude - Not a Primary Study

Whittemore K. Electronic prescription transmission. *National Association of Retail Druggists Journal* 2004;117:18-20. Database: IPA.
Exclude - No Outcomes of Interest

Whittemore K. The case for electronic prescribing. *Journal of the Pharmacy Society of Wisconsin* 2004; Database: IPA.
Exclude - Not MMIT

Whittemore K. Working with state pharmacy boards to achieve safety and efficiency. *Computertalk for the Pharmacist* 2004;24(3):32-3. Database: IPA.
Exclude - Not a Primary Study

Whittemore K. The case for electronic prescribing. *Nebraska Mortar & Pestle* 2005;68(2):26-7. Database: IPA.
Exclude - Not MMIT

Wickham V, Miedema F, Gamedinger K, et al. Bar-coded patient ID: review an organizational approach to vendor selection. *Nurs Manag (Harrow)* 2006;37(12):22-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wiedemann T, Knap P, Bachert A, et al. Computer-aided documentation and therapy planning in pediatric oncology. *Studies in Health Technology & Informatics* 1998;52 Pt 2:1306-9. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wiener F, Groth T, Mortimer O, et al. A knowledge-based information system for monitoring drug levels. *Computer Methods & Programs in Biomedicine* 1989;29(2):115-28. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Will EJ, Richardson D, Tolman C, et al. Development and exploitation of a clinical decision support system for the management of renal anaemia. *Nephrology Dialysis Transplantation* 2007;22(Suppl 4):iv31-iv36 Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Willcourt RJ, Pager D, Wendel J, et al. Induction of labor with pulsatile oxytocin by a computer-controlled pump. *Am J Obstet Gynecol* 1994;170(2):603-8. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Willems J, Raymakers A, Daneels F, et al. Computer applications in a hospital pharmacy. *Methods Inf Med* 1975;14(4):194-8. 847481

Database: Inspec.

Exclude - No Outcomes of Interest

Williams AG. Insulin algorithms in the self-management of insulin-dependent diabetes: the interactive 'Apple Juice' program. *Med Inform (Lond)* 1996;21(4):327-44. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Williams CK, Maddox RR. Implementation of an i.v. medication safety system. *Am J Health Syst Pharm* 2005;62(5):530-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Williams CP. Implementation of a telepharmacy program within a health system. *Ashp Midyear Clinical Meeting* 2003;38: Database: IPA.

Exclude - Not a Primary Study

Williams CT, Lee VF, Marsalis LD. Implementation of an automated medication/supply distribution system. *Hosp Mater Manage Q* 1994;16(2):76-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Williams CT. Inside a closed-loop medication strategy. *Nurs Manag (Harrow)* 2004;3(Suppl 5):8-9. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Williams F. The role of the electronic medical record (EMR) in care delivery development in developing countries: A systematic review. *Inform Prim Care* 2008;16(2):139-45. Database: Embase Sept 22-09.

Exclude - Not MMIT

Williams Y, Apple A, Greene W, et al. Minimizing heparin risk: Evaluation of the impact of an automated "clinical rule". *Ashp Summer Meeting* 2004;61: Database: IPA.

Exclude - Not a Primary Study

Williams Y, Apple A, Nolly R, et al. Improving enoxaparin prescribing: Evaluation of the impact of an automated "clinical rule". *Ashp Summer Meeting* 2005;62: Database: IPA.

Exclude - Not a Primary Study

Williamson JE. Budget- and eco-savvy hospitals boost reprocessing compliance. *Healthcare Purchasing News* 2010;34(3):36-43.
<http://libaccess.mcmaster.ca/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=48633313&site=ehost-live&scope=site EBSCO-BusinessSourceComplete>.
Exclude - Not a Primary Study

Williamson S. Management of oral anticancer therapies. *Pharmaceutical Journal* 2008;281(4 October 2008):399-402. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Willson W. Electronic prescribing within the context of the NHS Care Records Service: An opportunity for new computing? *British Journal of Healthcare Computing* 2002;21(2):38-40. Database: CINAHL.
Exclude - No Outcomes of Interest

Wilson GN, Moore KA. Multiple-order intravenous drug management system in a medical/surgical intensive care unit: Opportunities and challenges for pharmacy practice. *Ashp Midyear Clinical Meeting* 1990;25: Database: IPA.
Exclude - Not a Primary Study

Wilson GN, Moore KA, Wellman GS, et al. Patient specific bar codes for IV drug therapy in an ICU. *Ashp Midyear Clinical Meeting* 1991;26: Database: IPA.
Exclude - Not a Primary Study

Wilson J. The perfect scan - 100% Bar code medication administration. *Healthcare Information and Management Systems Society*; 2007. Grey Lit.
Exclude - Not a Primary Study

Wilson J. The perfect scan - 100% Bar code medication administration. *Healthcare Information and Management Systems Society*; 2007.
<http://www.himss.org/ASP/ContentRedirector.asp?ContentID=66799> Grey Lit.
Exclude - Not MMIT

Wilson J, Anderson MA. Casting electronic safety nets across care continuums. *Nurs Manag (Harrow)* 2004;35(Suppl 5):4-7. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wilson JR, Ceccanti J, Frederick C, et al. Computer modelling and microcostin methodologies in preliminary feasibility studies of automated dispensing systems. *Ashp Midyear Clinical Meeting* 1988;23: Database: IPA.
Exclude - Not a Primary Study

Wilson K, Sullivan M. Preventing medication errors with smart infusion technology. *Am J Health Syst Pharm* 2004;61(2):177-83. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wilson KH. Implementation and evaluation of a fluoroquinolone formulary change in a large community hospital. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
Exclude - Not a Primary Study

Wilson L. A systematic approach: new study looks at what drives top performance in clinical quality, efficiency at systems. *Mod Healthc* 2009;39(32):26-8.

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010382170&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010382170 EBSCO CINAHL.

Exclude - Not a Primary Study

Windle J, Van Milligan G, Duffy S, et al. Rapid deployment of physician order entry using web-based, disease-specific order sets. *AMIA 2003;Annual Symposium Proceedings 2003*:1079 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Windle J, Van Milligan G, Duffy S, et al. Web-based physician order entry: an open source solution with broad physician involvement. *AMIA 2003;Annual Symposium Proceedings 2003*:724-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Wingfield J, Foster C. Consent and confidentiality: Legal implications of electronic transmission of prescriptions. *Pharmaceutical Journal* 2002;269(7214):328-31. Database: IPA.

Exclude - Not MMIT

Winkelman WJ, Leonard KJ, Rossos PG. Patient-perceived usefulness of online electronic medical records: Employing grounded theory in the development of information and communication technologies for use by patients living with chronic illness. *J Am Med Inform Assoc* 2005;12(3):306-14. Database: IPA.

Exclude - Not MMIT

Winsor SJ, Hartman CH, Silverman JB. Efficacy of order entry warfarin-drug interaction alert systems based on physician responses and patient outcomes in general medicine patients. *Ashp Midyear Clinical Meeting* 2001;36: Database: IPA.

Exclude - Not a Primary Study

Winterbottom LM, Fong AM, Benkstein KL, et al. Impact of a clinical pharmacy consult service on guideline adherence and management of gabapentin for neuropathic pain. *J Manag Care Pharm* 2006;12(1):61-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Winters J. Microcomputers for improvement of quality of stock formulations in public pharmacies. *Pharm Weekbl* 1983;118: Database: Embase Sept 22-09.

Exclude - Unable to Retrieve Foreign

Wise J, Operario D. Use of electronic reminder devices to improve adherence to antiretroviral therapy: A systematic review. *Aids Patient Care STDS* 2008;22(6):495-504. Database: PsycINFO.

Exclude - Not MMIT

Witteman CLM, Kunst H. SelectCare--in aid of psychotherapists' treatment decisions. *Computers in Human Behavior* 1999;15(1999):143-59. Database: PsycINFO.

Exclude - Not MMIT

Witter J, Jatko M, Eisert S and others. Computerized clinical decision support systems have the potential to detect drug-lab interactions in outpatient clinics. In Proceedings 2002. Bethesda, MD, USA: American Medical Informatics Assoc; 2002. p. 1202.8037236
Database: Inspec.

Exclude - Not MMIT

Wittmer PD. Reducing pharmacist medication order profiling errors. Ashp Summer Meeting 2004;62: Database: IPA.

Exclude - Not a Primary Study

Wittwer WK. Nursing involvement, buy-in and workflow considerations: Overcoming the hurdles. Ashp Midyear Clinical Meeting 2003;38: Database: IPA.

Exclude - Not a Primary Study

Wogen SE. Improving the efficiency of the prescription process and promoting plan adherence. Drug Benefit Trends 2003;15(9):35-40. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Wolf ZR. Pursuing safe medication use and the promise of technology. MedSurg Nursing: official journal of the Academy of Medical-Surgical Nurses 2007;16(2):92-100. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Womer RB, Tracy E, Soo-Hoo W, et al. Multidisciplinary systems approach to chemotherapy safety: rebuilding processes and holding the gains. J Clin Oncol 2002;20(24):4705-12.

Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Wong B, Calvey J, Yeung S, et al. Two-year follow-up of the computerized physician order entry system compared to the order turn-around time of the paper-based system. Ashp Midyear Clinical Meeting 2007;42: Database: IPA.

Exclude - Not a Primary Study

Exclude - Not a Primary Study

Wong BJ. An approach to preventing methotrexate prescribing errors in rheumatoid arthritis. Hosp Pharm 1993;28(11):1081-2. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Exclude - Not a Primary Study

Wong BJ, Vogenberg FR. Systematic prevention of methotrexate prescribing errors in rheumatoid arthritis patients. Ashp Midyear Clinical Meeting 1995;30: Database: IPA.

Exclude - Not a Primary Study

Wong BJ, Rancourt MD, Clark ST. Enhancing pharmacy clinical services through the use of an automated drug dispensing system. Ashp Midyear Clinical Meeting 1997;32: Database: IPA.

Exclude - Not a Primary Study

Wong BJ, Chandler-Toufieli DM, Sullivan DJ, et al. Enhancing the renal dosing adjustment process using automation. Ashp Midyear Clinical Meeting 1999;34: Database: IPA.

Exclude - Not a Primary Study

Exclude - Not a Primary Study

Wong BJ, Yeung SK, Swinney KW, et al. Comparison of order turn-around time in computerized physician order entry (CPOE) to order turn-around time in the paper-based system. *Ashp Midyear Clinical Meeting 2006*;41: Database: IPA.
Exclude - Not a Primary Study

Wong C, Geiger G, Derman Y D and others. Redesigning the medication ordering, dispensing, and administration process in an acute care academic health sciences centre. In *Piscataway, NJ, USA: IEEE; 2003. p.1894-902.8045513*
Database: Inspec.
Exclude - No Outcomes of Interest

Wong D, DeBusk M, Sankrithi U. Implementation of an ambulatory medication management application in a pediatric emergency department. *Ashp Summer Meeting 2004*;64: Database: IPA.
Exclude - Not a Primary Study

Wong ICK, Wong LYL, Cranswick NE. Minimising medication errors in children. *Arch Dis Child 2009*;94:161-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wong PK, Fecher M, Hill GW. Implementation of one way order entry interface between a pharmacy and a nursing computer systems. *Ashp Midyear Clinical Meeting 1995*;30: Database: IPA.
Exclude - Not a Primary Study

Wood JR, Laffel LMB. Technology and intensive management in youth with type 1 diabetes: state of the art. *Current Diabetes Reports 2007*;7(2):104-13. Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Wood L, Martinez C. The general practice research database: role in pharmacovigilance. *Drug Saf 2004*;27(12):871-81. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Woodall SC. Remote order entry and video verification: Reducing after-hours medication errors in a rural hospital. *Joint Commission Journal on Quality & Safety 2004*;30(8):442-7. Database: CINAHL.
Exclude - Not MMIT

Woodall SC. Telepharmacy. *Ashp Midyear Clinical Meeting 2006*;41: Database: IPA.
Exclude - Not a Primary Study

Woodruff AE, Hunt CA. Involvement in medical informatics may enable pharmacists to expand their consultation potential and improve the quality of healthcare. *Ann Pharmacother 1992*;26(1):100-4. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Woosley RL. Drug labeling revisions: Guaranteed to fail? *JAMA 2000*;284(23):3047-9. Database: IPA.
Exclude - Not a Primary Study

Work M. Improving medication safety with a wireless mobile barcode system in a community hospital. *Patient Safety & Quality Healthcare* 2005;May/June 2005: <http://www.psqh.com/mayjun05/casestudy.html>.
Exclude - Not a Primary Study

Wright A, Sittig DF. SANDS: a service-oriented architecture for clinical decision support in a National Health Information Network. *Journal of Biomedical Informatics* 2008;41(6):962-81. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wright AA, Katz IT. Bar coding for patient safety. *N Engl J Med* 2005;353(4):329-31. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wright L, Hargrove FR, Waitman LR, et al. Computerized decision support for intravenous fluid management in pediatric patients. *Hosp Pharm* 2004;39(10):996-1001. Database: Embase Sept 22-09.
Exclude - Not a Primary Study

Wright L, Waitman LR, Grisso A. Computerized provider order entry. Implementation of pediatric-specific decision support in a CPOE system. *Hosp Pharm* 2004;39(4):381-5. Database: IPA.
Exclude - Not a Primary Study

Wright L, Hargrove FR, Grisso AG. Improving communication and safety with CPOE decision support tools. *Hosp Pharm* 2004;39(11):1106-17. Database: IPA.
Exclude - Not a Primary Study

Wright L, Hargrove FR. Use of decision support in a computerized prescriber order entry system to prevent medication errors associated with ordering of potassium chloride in a pediatric critical care unit. *Ashp Midyear Clinical Meeting* 2004;39: Database: IPA.
Exclude - Not a Primary Study

Wright L, Feldott CC, Hargrove FR. Designing decision support for insulin ordering in a computerized provider order entry system. *Hosp Pharm* 2007;42(2):158-61. Database: Embase Sept 22-09.
Exclude - No Outcomes of Interest

Wright R, Handler SM, Ruby C, et al. Update on drug-related problems in the elderly. *American Journal Geriatric Pharmacotherapy* 2006;4(1):78-84. Database: Embase Sept 22-09.
Exclude - Not MMIT

Wrobel JP. Are we ready for the Better Medication Management System? *Med J Aust* 2003;178(9):448-50. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Wu A, Snyder C, Huang I, et al. A randomized trial of the impact of a programmable medication reminder device on quality of life in patients with HIV. *Aids Patient Care STDS* 2006;20(11):773-81. Database: PsycINFO.
Exclude - Not MMIT

Wu H, Ozok A. Medication adherence among the elderly and technology aids: Results from an online survey study. In A.A.Ozok and P.Zaphiris (Eds.): Online Communities, editor. 5621. 2003; San Diego, CA) SUBJECT(S) Identifier Information technology; Computer networks; Online social networks Note(s) Includes bibliographical references and index.: 2009. p.719-27.Grey Lit.

Exclude - Not MMIT

Wu JH, Shen WS, Lin LM, et al. Testing the technology acceptance model for evaluating healthcare professionals' intention to use an adverse event reporting system. *Int J Qual Health Care* 2008;20(2):123-9. Database: Embase Sept 22-09.

Exclude - Not MMIT

Wu R, Peters W, Morgan MW. The next generation of clinical decision support: linking evidence to best practice. *J Healthc Inf Manag* 2002;16(4):50-5. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Wu RC, Abrams H, Baker M, et al. Implementation of a computerized physician order entry system of medications at the University Health Network--physicians' perspectives on the critical issues. *HEALTHC Q* 2006;9(1):106-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Wu,S.j. Impact of information quality on the use and effectiveness of computerized clinical reminders Purdue Univeristy. 2010. OVID PsychINFO.

Exclude - Not MMIT

Wu Y. Internet-based mode of real-time drug supervision and administration. *Chinese Journal of New Drugs* 2008;17(22): Database: Embase Sept 22-09.

Exclude - Not MMIT

Wurster M, Doran T. Anticoagulation management: a new approach. *Disease Management* 2006;9(4):201-9. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Wurster MW. Computerized patient management system improves compliance, efficiency and revenue in an anticoagulation clinic. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Wyatt J, Walton R. Computer based prescribing. *Br Med J* 1995;311(7014):1181-2. Database: IPA.

Exclude - Not a Primary Study

Xie M, Johnson K. Applying human factors research to alert-fatigue in e-prescribing. *AMIA* 2007;1161 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Yamada H, Nishimura S, Shimamori Y, et al. The development and operation of a package inserts service system for electronic medical records. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2003;123(3):201-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Yamada H, Makino E, Niinuma Y, et al. Construction and evaluation of a cancer chemotherapy regimen database using an electronic medical chart network. *Yakugaku Zasshi - Journal of the Pharmaceutical Society of Japan* 2005;125(7):567-77. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Yamamoto K, Orii T, Kotaki H, et al. Development of a computerized accounting system for therapeutic drug monitoring [Japanese]. *Journal of the Nippon Hospital Pharmacists Association* 1998;24(5):567-75. Database: IPA.

Exclude - Not MMIT

Yamamoto LG, Wiebe RA. Pediatric and adult emergency management assistance using computerized guidelines. *Am J Emerg Med* 1989;7(1):91-6. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Yamashiro H, Toi M. Update of evidence in chemotherapy for breast cancer. *International Journal of Clinical Oncology* 2008;13(1):3-7. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Yan Q, Hunt CA. Preventing adverse drug events (ADEs): The role of computer information systems. *Drug Inf J* 2000;34(4):1247-60. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Yang J, Nugroho AS, Yamauchi K, et al. Efficacy of interferon treatment for chronic hepatitis C predicted by feature subset selection and support vector machine. *J Med Syst* 2007;31(2):117-23. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Yang WH, Cai WM, Xu B, et al. Practice on effectively running pharmacy intravenous admixture services. *Pharmaceutical Care and Research* 2006;6(3):218-21. Database: Embase Sept 22-09.

Exclude - Not MMIT

Yang Y, Han X, Bao F, et al. A smart-card-enabled privacy preserving E-prescription system. *IEEE Transactions on Information Technology in Biomedicine* 2004;8(1):47-58. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Yang ZW, Hu JH, Quan SC. Implementation of barcode technology in pharmacy intravenous admixture services. *Pharmaceutical Care and Research* 2005;5(Part 3):213-5. Database: Embase Sept 22-09.

Exclude - Not MMIT

Yang Z, Hu JH, Wang Z, et al. Design and application of drug dispensing software with on-line drug information. *Pharmaceutical Care and Research* 2006;6(4): Database: Embase Sept 22-09.

Exclude - No Outcomes of Interest

Yap KY, Chan A, Chui WK. Improving pharmaceutical care in oncology by pharmacoinformatics: the evolving role of informatics and the internet for drug therapy. [Review] [100 refs]. LANCET ONCOL 2009;10(10):1011-9. PMID:19796753 OVID MEDLINE.

Exclude - No Outcomes of Interest

Yarnall KS, Michener JL, Hammond WE. The medical record: a comprehensive computer system for the family physician. J Am Board Fam Pract 1994;7(4):324-34. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Yates C. Implementing a bar-coded bedside medication administration system. Crit Care Nurs Q 2007;30(2):189-95. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ye X, Fu Z, Wang H, et al. A computerized system for signal detection in spontaneous reporting system of Shanghai China. Pharmacoepidemiology & Drug Safety 2009;18(2):154-8. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Yee W, Ross D. Use of a system-wide electronic event reporting system to improve medication safety. Ashp Midyear Clinical Meeting 2006;41: Database: IPA.

Exclude - Not a Primary Study

Yeh H C, Hsiu P C, Shih C S and others. APAMAT. a prescription algebra for medication authoring tool. In Piscataway, NJ, USA: IEEE; 2006. p.4284-91.9772284 Database: Inspec.

Exclude - Not a Primary Study

Yeh H C, Hsiu P C, Shih C S and others. APAMAT: A prescription algebra for medication authoring tool. In Taipei, Taiwan: Institute of Electrical and Electronics Engineers Inc.; 2006. p.4284-91.20073510787141

Database: Compendex.

Exclude - Not a Primary Study

Yeh Y T, Liu C T, Wu S J and others. Assessment of user satisfaction with an internet-based integrated patient education system for diabetes management. In Singapore, Singapore: Inst. of Elec. and Elec. Eng. Computer Society; 2008. p.85-9.20084111624098

Database: Compendex.

Exclude - Not MMIT

Yen K, Shane EL, Pawar SS, et al. Time motion study in a pediatric emergency department before and after computer physician order entry. Ann Emerg Med 2009;53(4):462-8.

Database: CINAHL.

Exclude - Not MMIT

Ying A. Mobile physician order entry. J Healthc Inf Manag 2003;17(1):58-63. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Ying H, Lin F, MacArthur RD, et al. A fuzzy discrete event system approach to determining optimal HIV/AIDS treatment regimens. *IEEE Transactions on Information Technology in Biomedicine* 2006;10(4):663-76. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Ying H, Lin F, MacArthur RD, et al. A self-learning fuzzy discrete event system for HIV/AIDS treatment regimen selection. *IEEE Transactions on Systems, Man, & Cybernetics, Part B: Cybernetics* 2007;37(4):966-79. Database: Ovid MEDLINE(R).

Exclude - Not MMIT

Young J, Stevenson KB. Real-time surveillance and decision support: Optimizing infection control and antimicrobial choices at the point of care. *Am J Infect Control* 2008;36(3 Suppl.):S67-S74 Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Youngs C. Refining dose range checking and drug interactions to improve medication safety. *Ashp Midyear Clinical Meeting* 2000;35: Database: IPA.

Exclude - Not a Primary Study

Yourman L, Concato J, Agostini JV. Use of computer decision support interventions to improve medication prescribing in older adults: a systematic review. *American Journal Geriatric Pharmacotherapy* 2008;6(2):119-29. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Youssef A, McCoy C, Cunningham K, et al. Development and implementation of a customized CPOE model for enoxaparin ordering. *Ashp Midyear Clinical Meeting* 2005;40: Database: IPA.

Exclude - Not a Primary Study

Yu FB, Jr., Leising S, Turner S. Development approach to an enterprise-wide medication reconciliation tool in a free-standing pediatric hospital with commercial best-of-breed systems. *AMIA* 2007;1164 Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Yu JC, Ho M. Development of antibiotic quick orders within a computerized physician order entry (CPOE) system to improve the timeliness and duration of antimicrobial prophylaxis for surgical patients. *Ashp Midyear Clinical Meeting* 2006;41: Database: IPA.

Exclude - Not a Primary Study

Yu VL, Fagan LM, Hannigan J, et al. Antimicrobial selection by a computer: Blinded evaluation by infectious diseases experts. *Journal of the American Medical Association* 1979;242(12):1279-82. Database: IPA.

Exclude - Not MMIT

Yuce Y. MAMAS: Mobile asthma monitoring and assessment system. *Journal on Information Technology in Healthcare* 2007;5(4):229-38. Database: Embase Sept 22-09.

Exclude - Not a Primary Study

Zai AH, Grant RW, Estey G, et al. Lessons from implementing a combined workflow-informatics system for diabetes management. *J Am Med Inform Assoc* 2008;15(4):524-33. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Zakhour H. Computers cut costs in the lab. Health Serv J 1988;98(5086):150-1. 3102521
Database: Inspec.
Exclude - Not a Primary Study

Zallen BG. Actual practice with Interpractice Systems: first experiences. HMO Pract
1993;7(2):61-6. Database: Ovid MEDLINE(R).
Exclude - Not a Primary Study

Zalman D, Odeh M, Oliven A. Physicians' assessment of computerized prescribing.
Harefuah 2000;138(6):434-40, 519. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Zarn D. MOE/MAR project management: A (well-informed) bird's eye view. HEALTHC Q
2006;10:Spec No:39 Database: CINAHL.
Exclude - Not a Primary Study

Zarowitz BJ, Petitta A, Mlynarek M, et al. Cimetidine evaluation in critically ill medical and
surgical patients. Ashp Midyear Clinical Meeting 1991;26: Database: IPA.
Exclude - Not a Primary Study

Zarowitz BJ. Utilizing automation and technology to meet the challenges of pharmacy
practice. ASHP Annual Meeting 1994;51: Database: IPA.
Exclude - Not a Primary Study

Zeisset A. ICD-10-CM enhancements: A look at the features that will improve coding
accuracy. Journal of the American Health Information Management Association
2009;80(2):55-8. OVID EMBASE.
Exclude - Not a Primary Study

Zeng K, Bodenreider O, Nelson SJ. Design and implementation of a personal medication
record-MyMedicationList. AMIA 2008;Annual:Symposium-8 PMID:18998962 OVID
MEDLINE.
Exclude - No Outcomes of Interest

Zeng R-J. Factors influencing clinical intervention of prescription automatic screening
system. Pharmaceutical Care and Research 2004;4(3): Database: Embase Sept 22-09.
Exclude - Unable to Retrieve

Zheng K, Padman R, Johnson MP, et al. An interface-driven analysis of user interactions
with an electronic health records system. J Am Med Inform Assoc 2009;16(2):228-37.
Database: Ovid MEDLINE(R).
Exclude - Not MMIT

Zhou Q, Zhu LL, Yan XF, et al. Drug utilization of clarithromycin for gastrointestinal
disease treatment. World Journal of Gastroenterology 2008;14(39):6065-71. Database: Ovid
MEDLINE(R).
Exclude - Not MMIT

Ziegenhagen DJ, Frye C, Kottmair S. Patient-oriented healthcare programs. Concepts and
practical experience in the field of chronic heart failure. Z Arztl Fortbild Qualitatssich
2005;99(3):209-15. Database: Ovid MEDLINE(R).
Exclude - Unable to Retrieve Foreign

Zigmond J. H1N1 under surveillance: feds, consumers getting plenty of assistance in tracking pandemic flu. *Mod Healthc* 2009;39(50):33

<http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=2010504843&site=ehost-live>; Publisher URL: www.cinahl.com/cgi-bin/refsvc?jid=772&accno=2010504843 EBSCO CINAHL.

Exclude - Not a Primary Study

Zillich AJ. Antimicrobial use control measures to prevent and control antimicrobial resistance in US hospitals. *Infect Control Hosp Epidemiol* 2006;27(10):1088-95. Database: Embase Sept 22-09.

Exclude - Not MMIT

Zimmerman CR, Chaffee BW, Lazarou J, et al. Maintaining the enterprisewide continuity and interoperability of patient allergy data. *Am J Health Syst Pharm* 2009;66(7):671-9. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Zito JM. Psychotropic practice patterns for youth: A 10-year perspective. *Arch Pediatr Adolesc Med* 2003;157(1):17-25. Database: Embase Sept 22-09.

Exclude - Not MMIT

Zoka R, Hyams P, Broderson R, et al. Problems concerning documentation of infusion orders and medication administration in a physician order entry computer system at intensive care units. *Ashp Midyear Clinical Meeting* 1991;26. Database: IPA.

Exclude - Not a Primary Study

Zolnierz M. Hospital pharmacy-based, computer-generated Tikosyn (Dofetilide) physician order protocol. *Hosp Pharm* 2003;38(7):659-61. Database: IPA.

Exclude - Not MMIT

Zytkowski ME. Nursing informatics: the key to unlocking contemporary nursing practice. *AACN Clin Issues* 2003;14(3):271-81. Database: Ovid MEDLINE(R).

Exclude - Not a Primary Study

Appendix F. Glossary of Terms

Adverse drug event (ADE). Harm caused by the use of a drug. ADEs also include adverse drug reactions, which is harm directly cause by a drug at the normal doses. ADEs can also be classified as preventable or not.

Source: Nebeker JR, Barach P, Samore MH. Clarifying adverse drug events: A clinician's guide to terminology, documentation, and Reporting. *Annals of Internal Medicine*. 2004;140:795-801.

Adverse event. An adverse event is a specific undesirable medical occurrence. It can be either a new undesirable medical problem or worsening of an existing health or medical problem.

Source: <http://www.gsk-clinicalstudyregister.com/glossary.jsp>.

Bar Code Medication Administration (BCMA). BCMA is a barcode system consisting of a barcode reader, a portable computer with wireless connection, a computer server, and software. Patients and medications are barcoded and both barcodes must match before the medication is administered. Often BCMA systems also record medication events and timing. Source: http://en.wikipedia.org/wiki/Bar_Code_Medication_Administration

Clinical Decision-Support System (CDSS). Computer tools or applications to assist in clinical decisions by providing evidence-based knowledge in the context of patient specific data. CDSSs for this report must be capable of integrating patient specific information from an existing system and external evidence to provide an alert or reminder to the clinician about actions to be or not be taken.

Source: Health IT.hhs.gov glossary

Clinical Outcomes. For this report we defined clinical outcomes liberally as any clinical morbidity, mortality, adverse event or clinical surrogate such as improved LDL cholesterol, asthma symptoms or quality of life, as the primary outcome of the study. They are also defined in this report as those things that happen to, and are important to patients in the study or real life situations.

Computerized Provider Order Entry (CPOE). A computer application that allows a provider's orders for diagnostic and treatment services (such as medications, laboratory, and other tests) to be entered and transferred electronically. During ordering or monitoring the CPOE system can compare the order against standards for dosing, checks for allergies or interactions with other medications and warns the physician about potential problems including duplication. Most CPOE systems are integrated into other existing health IT.

Source: Health IT.hhs.gov glossary

Cost-Benefit Analysis. Cost-Benefit Analysis (CBA) requires programme consequences to be valued in monetary units, thus enabling the analyst to make a direct comparison of the programme's incremental cost with its incremental consequences in commensurate units of measurements. CBA compares discounted future streams of incremental programme benefits with incremental programme costs; the difference between these two streams being the net

social benefits of the programme. In simple terms, the goal of analysis is to identify whether a programme's benefits exceed its costs a positive net social benefit indicating that programme is worthwhile.

Source: Drummond MF, Methods for the Economic Evaluation of Health Care Programmes, Ch-7, 3rd Edition, 2005, Oxford University Press

Cost-Effectiveness Analysis. Cost-Effective Analysis (CEA) is one form of full economic evaluation where both the costs and consequences of health programmes or treatments are examined. In CEA, the incremental cost of a programme from a particular viewpoint is compared to the incremental health effects of the programme, where the health effects are measured in natural units related to the objective of the programme. The results are usually expressed as a cost per unit of effect.

Source: Drummond MF, Methods for the Economic Evaluation of Health Care Programmes, Ch-5, 6, 3rd Edition, 2005, Oxford University Press.

Cost study. The cost study designation is a broad umbrella term used for all studies that include costs. More formal costs studies include cost-benefit, cost-utility, cost effectiveness analyses.

Cost-Utility Analysis. Cost-Utility Analysis (CUA) is one form of evaluation where both that focuses particular attention on the quality of the health outcome produced or forgone by health programmes or treatments. In CUA, the incremental cost of a programme from a particular viewpoint is compared to the incremental health improvement attributable to the programme, where health improvement is measured in quality-adjusted life-years (QALYs) gained, or possibly some variant, like disability adjusted life-years (DALYs) gained. The results are expressed per QALY gained.

Source: Drummond MF, Methods for the Economic Evaluation of Health Care Programmes, Ch-6, 3rd Edition, 2005, Oxford University Press.

e-Prescribing. A type of computer technology that clinicians use handheld or personal computer devices to review drug and formulary coverage and to transmit prescriptions to a printer or to a local pharmacy and often store this information. e-Prescribing software can be integrated into existing clinical information systems to allow physician access to patient specific information to screen for drug interactions and allergies. e-Prescribing systems are less complex than CPOE systems that allow ordering of drugs. For this report we use author-derived designations to differentiate between e-Prescribing and CPOE systems that are used to order medications.

Source: Health IT.hhs.gov glossary.

Electronic Data Interchange (EDI). Refers to the exchange of routine business transactions from one computer to another in a standard format, using standard communications protocols. This report concentrates on EDI in the communication between clinicians and pharmacists to perfect the order or prescription.

Source: Centre for Medicare and Medicaid Services, HHS

Electronic Health Record (EHR). An electronic record of health-related information on an individual that conforms to nationally recognized interoperability standards and that can be created, managed, and consulted by authorized clinicians and staff across more than one health care organization. An EHR system is usually broader than an EMR system. EMRs have traditionally been hospital based. For this report we use whatever designation the authors provide in their studies.

Source: Health IT.hhs.gov report page 15

Electronic Medical Record (EMR). An electronic record of health-related information on an individual that can be created, gathered, managed, and consulted by authorized clinicians and staff within one health care organization. These EMR systems are often hospital based and often not connected with information on the patient available outside the hospital system.

Source: Health IT.hhs.gov report page 15

Electronic Medication Administration Record (eMAR). Electronic medication administration record systems are hospital based, point-of-care systems that usually incorporate BCMA capabilities to make the administration of medications safer for patients by reducing error rates and allowing nurses to more efficiently manage medication tasks. eMAR systems record all medication administrative events including time of administration and integrate with pharmacy information systems.

Source: fgraham blog post

Health Information Technology (health IT). Health IT is the application of information processing involving both computer hardware and software that deals with the storage, retrieval, sharing, and use of health care information, data, and knowledge for communication and decision making.

Source: Health IT.hhs.gov glossary

Intermediate Outcomes. For this evidence report intermediate outcomes were defined as satisfaction with system, usability, knowledge, skills, and attitude, and other related issues.

Major Endpoint. Also known as the primary outcome. The major endpoint is the main outcome that researchers determine to be the most important of any of the measures taken during planning and implementation of a study. Most studies have one to two major endpoints and multiple endpoints. Study size calculations are based on the major endpoint.

Medication Errors. Any error that occurs during the medication management process (prescribing, order communication, dispensing, administering, and monitoring). These can be potential errors--ones that are identified and addressed before the patient receives the medication or actual errors. The actual errors are ones that occur when the patient receives the wrong medication, the wrong dose or form, or at the wrong time. Medication errors can also be preventable and non-preventable. We used author identified statements of our classification of medication errors in this report.

Source: Ferner RE, Aronson JK. Clarification of terms used in medication errors: Definitions and classifications. Drug Safety. 2006 Nov;(11):1011-22.

Medication Management. Medication management is a continuum that covers all aspects of prescription medication. Medication management includes the five phases of the medication process (prescribing and ordering, order communication, dispensing, administering, and monitoring). Bell and colleagues in their seminal work on describing and modeling medication management outline the phases as being prescribe, transmit, dispense, administer and monitor¹ For this report, based on input from our TEP, to have greater clarity of what is occurring in the transmit phase, especially the active involvement by the pharmacist, we refer to the transmission of the order or prescription and the bi-directional communication between prescriber and pharmacy staff as “order communication”.

Medication management can also include procurement, storage, reconciliation, and reporting involved in the assessment of patients for the need for drugs through to optimal care and monitoring after the drugs are prescribed. For this report we also included issues related to education or training in the use of health IT in medication management.

Medication Management through Health Information Technology (MMIT). MMIT systems are electronic systems that (1) collect, process, or exchange health information about patients and formal caregivers, (2) are integrated with existing health IT such as EHR or EMR systems, and (3) provide advice or suggestions to either the health care provider or the patients and their families on issues related to medication management.

Source: <http://www.ahrq.gov/clinic/tp/medmgttp.htm>.

Medication Monitoring. The process of assessing a patient’s response to a medication and documenting its outcomes based on physical findings, history, laboratory testing, or a combination of any of these.

Source: Handler SM, Nace DA, Studenski SA, et al. Medication error reporting in long term care. *Am J Geriatr Pharmacother* 2004;2(3):190-6.

Medication Monitoring Errors. Errors due to inadequate laboratory evaluation of drug therapies or a delayed or failed response to signs or symptoms of drug toxicity or laboratory evidence of toxicity.

Source: Fillit H, Rockwood K, Woodhouse L. Brocklehurst’s Textbook of Geriatric Medicine and Gerontology. 7th. Churchill Livingstone; 2010. Geriatric Pharmacotherapy and Polypharmacy.

Medication Reconciliation. A formal process of identifying the most complete and accurate list of medications a patient is taking and using that list to provide correct medications for the patient anywhere within the health care system.

Source: <http://www.wicheckpoint.org/DefinitionOfTerms.aspx>

Patient Safety. Freedom from accidental injuries during the course of receiving medical care.

Source: <http://www.bvs.org.ar/pdf/seguridadpaciente.pdf>

Personal Health Record (PHR). An electronic record of health-related information on an individual that is maintained by the person themselves. The PHR can conform to nationally

recognized interoperability standards. Data may be stand alone and entered only by patients and their caregivers or be fully integrated with EHRs and other health IT systems.

Source: Health IT.hhs.gov

Pharmacy Information System. An application that provides complete support for the pharmacy (hospital, community based or other pharmacies) from an operational, clinical and management perspective, helping to optimize patient safety, streamline workflow and reduce operational costs.

Source: <http://www.himssanalytics.org/docs/Definitions-By-Term.pdf>

Pragmatic Trial. Pragmatic trials are designed to find out about how effective a treatment actually is in routine, everyday practice. Pragmatic trials answer questions about the overall effectiveness of an intervention, and cannot study the contributions of its different components. Pragmatic trials are used to test an overall ‘package’ of care, including the contribution of the therapeutic relationship, patients’ expectations, and any specific therapy that is used. Generally a pragmatic trial would compare the effect of this package of care with another treatment, not with a placebo. Pragmatic trials are used with the aim of providing the evidence that will help policy makers, practitioners or patients make choices between two interventions. They help define the best use of limited resources.

Source: <http://www.frtcm.org/Pragmatic%20trials%20CTM%202004%2012%20136-40.pdf>

Primary Outcome: *See Major Endpoint.*

Process Changes: Also known as Process Outcomes. These are study outcomes related to how the care process happens. For example, time to perform tasks, workflow changes, improved efficiencies, modifications of prescriptions, and errors in prescriptions are considered to be process changes or outcomes for studies of MMIT.

Qualitative Research. Qualitative research seeks out the ‘why’, not the ‘how’ of its topic through the analysis of unstructured information—things like interview transcripts, open ended survey responses, emails, notes, feedback forms, photos and videos. It doesn’t just rely on statistics or numbers, which are the domain of quantitative researchers. Qualitative research is used to gain insight into people’s attitudes, behaviors, value systems, concerns, motivations, aspirations, culture or lifestyles. It’s used to inform business decisions, policy formation, communication and research. Focus groups, in-depth interviews, content analysis, ethnography, evaluation and semiotics are among the many formal approaches that are used, but qualitative research also involves the analysis of any unstructured material, including customer feedback forms, reports or media clips.

Source: <http://www.qsrinternational.com/what-is-qualitative-research.aspx>.

Signs. Evidence of disease ascertained by the clinician using direct observation or tools such as a stethoscope or blood pressure monitor. These signs are used to diagnosis a disease or disorder or monitor the progress of a healthcare issue.

Sustainability. The ability of a health service to provide ongoing access to appropriate quality care in a cost-effective and health-effective manner.

Source: Humphreys JS, Wakerman J, Wells R. What do we mean by sustainable rural health services? Implications for rural health research. *Aust J Rural Health* 2006;14(1):33-5.

Symptoms. Symptoms are patient reported issues (e.g., pain, fatigue, or depression) that the clinician considers along with signs to ascertain a disease or disorder or monitor disease progression.

Tall Man letters. Use of capital letters in look-alike drug names to help guarantee differentiation, For example, Novo**LOG** and Novo**LIN**, and Huma**LOG** and Humu**LIN**, helped differentiate these products.

Usability. Usability is a measure of how learnable, efficient, memorable, error free, and satisfactory a computer system or program is. Standard methods are available that measure the usability of a system and provide strategies to improve its usability aspects. A system that has high usability will be used and used efficiently.

Source: Nielsen J. *Usability Engineering*. Academic Press. San Diego, CA. 1993

Use. A simple measure or count of how often a system or application is used.

Source: Nielsen J. *Usability Engineering*. Academic Press. San Diego, CA. 1993

Usefulness. Usefulness is a soft measure of whether the system or application meets its stated goals.

Source: Nielsen J. *Usability Engineering*. Academic Press. San Diego, CA. 1993

Value proposition. Broadly speaking, ‘value proposition’ refers to the benefits one receives by adopting a particular product, approach, or technology, as compared to what you currently have, or what some other competitive offering would provide. In monetary terms, the value proposition is what the customer gets for his/her money/time. It can also be regarded as differences in performance and/or cost between two different alternatives, such as response speed, product or service quality, and the relative performance in terms of satisfaction or preference. Search terms: ‘return on investment,’ ‘cost benefit,’ ‘relative value,’ ‘relative performance,’ etc.

Source: Dr. Norm Archer, McMaster University, July 2009.

Value of health IT. Clinical, organizational, financial or other benefits derived from the adoption, utilization, and diffusion of health IT less the costs of achieving these benefits (<http://grants.nih.gov/grants/guide/rfa-files/RFA-HS-04-012.html>).