## "Restoring America's Wetlands"—References

Birdsey, R.A. (1996) Regional Estimates of Timber Volume and Forest Carbon for Fully Stocked Timberland, Average Management After Final Clearcut Harvest. In Forests and Global Change: Volume 2, Forest Management Opportunities for Mitigating Carbon Emissions, eds. R.N. Sampson and D. Hair, American Forests, Washington, DC.

Bishop, A., M. Vrtiska. May 2008.Effects of the Wetlands Reserve Program on Waterfowl Carrying Capacity in the Rainwater Basin Region of South-Central Nebraska. 51p.

Faulkner, S., M. Baldwin, W. Barrow, H. Waddle, B. Keeland, S. Walls, D. James, T. Moorman. June 2010. Final Report: Regional Estimates of Ecological Services Derived from U.S. Department of Agriculture Conservation Programs in the Mississippi Alluvial Valley.

Gleason, R.A., Laubhan, M.K., and Euliss, N.H., Jr., eds., 2008, Ecosystem services derived from wetland conservation practices in the United States Prairie Pothole Region with an emphasis on the U.S. Department of Agriculture Conservation Reserve and Wetlands Reserve Programs: U.S. Geological Professional Paper 1745, 58 p.

King, S.L., D. Twedt, R. Wilson. The Role of the Wetland Reserve Program in Conservation Efforts in the Mississippi River Alluvial Valley. 2006. V34. No 4. P 914-920

Carbon Sequestration: <u>http://www.epa.gov/sequestration/rates.html</u> Wetlands: <u>http://water.epa.gov/type/wetlands/toc.cfm</u>