

**Table 96. Select Family Planning and Medical Service Use by Women, 15 to 44 Years of Age: 2002**

[61,561 represents 61,561,000. Based on the National Survey of Family Growth, see Appendix III]

Characteristic	Number (1,000)	Percent using—							
		Family planning services			Medical services				
		At least one family planning service	Birth control method	Birth control checkup or test	At least one medical service	Pregnancy test	Pap smear	Pelvic exam	Counseling/test/treatment for sexually transmitted diseases
<b>All women</b> <sup>1</sup>	<b>61,561</b>	<b>41.7</b>	<b>33.9</b>	<b>23.6</b>	<b>69.1</b>	<b>19.7</b>	<b>64.4</b>	<b>59.7</b>	<b>12.6</b>
15 to 19 years old	9,834	39.9	31.1	22.0	40.6	18.3	34.6	27.0	15.2
15 to 17 years old	5,819	31.8	22.2	15.8	28.0	11.4	23.2	17.9	11.1
18 to 19 years old	4,016	51.6	43.9	31.0	58.9	28.2	51.2	40.2	21.1
20 to 24 years old	9,840	63.3	54.0	35.7	75.7	31.5	69.7	60.6	22.3
25 to 29 years old	9,249	55.4	46.3	30.2	75.9	30.2	70.7	66.0	16.6
30 to 34 years old	10,272	47.0	39.1	27.2	78.1	22.2	72.7	69.7	12.2
35 to 39 years old	10,853	30.5	23.9	18.6	71.5	13.6	68.3	65.9	6.9
40 to 44 years old	11,512	19.5	14.0	10.8	71.8	5.9	69.4	67.0	4.4
Currently married	28,327	39.5	31.5	21.3	77.2	21.1	73.1	69.8	8.1
Currently cohabiting	5,570	50.4	43.2	30.2	77.2	31.0	72.2	64.7	20.3
Never married, not cohabiting	21,568	44.4	36.4	25.4	55.8	16.5	50.7	43.9	15.9
Formerly married, not cohabiting	6,096	34.5	28.0	22.0	71.0	14.1	66.0	64.0	14.4
White only, Non-Hispanic	39,498	43.1	36.4	25.4	70.1	17.3	66.1	63.2	11.8
Black only, Non-Hispanic	8,250	39.4	30.6	21.8	74.4	23.7	69.1	58.1	15.7
Hispanic <sup>2</sup>	9,107	39.7	28.9	20.6	63.4	24.3	57.1	48.5	12.5

<sup>1</sup> Includes other races not shown separately. <sup>2</sup> Persons of Hispanic origin may be of any race.

Source: U.S. National Center for Health Statistics, special tabulations from the *Advance Data*, Number 350, "Use of Contraception and Use of Family Planning Services in the United States: 1982–2002," December 2004.

**Table 97. Current Contraceptive Use by Women by Age, Race, Hispanic Origin, Marital, and Cohabitation Status: 2006 to 2008**

[61,864 represents 61,864,000. Based on the Continuous National Survey of Family Growth conducted from July 2006 through December 2008. Contraceptive use reported by women 15–44 years of age during heterosexual vaginal intercourse. Women using more than one method of contraception are classified by most effective method reported]

Contraceptive status and method	Unit	2006–2008							
		Race/ethnicity				Marital and cohabitation status			
		All women <sup>1</sup>	White only, Non-Hispanic	Black only, Non-Hispanic	Hispanic <sup>2</sup>	Never married, not cohabiting	Currently married	Formerly married, not cohabiting	Currently cohabiting
<b>All women</b>	<b>(1,000)</b>	<b>61,864</b>	<b>37,660</b>	<b>8,452</b>	<b>10,377</b>	<b>22,847</b>	<b>27,006</b>	<b>5,190</b>	<b>6,821</b>
PERCENT DISTRIBUTION									
Using contraception (contraceptors) <sup>3</sup>	Percent	61.8	64.7	54.5	58.5	39.3	78.6	60.6	71.2
Female sterilization	Percent	16.7	14.9	21.8	19.6	4.5	23.6	35.3	16.3
Male sterilization	Percent	6.1	8.3	1.1	3.4	0.3	12.7	2.3	2.2
Pill	Percent	17.3	21.2	11.4	11.4	18.1	16.3	11.4	23.2
Implant, Lunelle™	Percent	0.7	0.5	0.6	1.5	0.6	0.7	(S)	1.0
3-month injectable (Depo-Provera™)	Percent	2.0	1.4	4.1	2.6	2.2	1.4	2.6	3.1
Intrauterine device (IUD)	Percent	1.5	1.6	1.7	1.2	1.5	1.0	0.8	3.7
Diaphragm	Percent	3.4	3.3	2.8	4.8	1.1	5.3	2.1	4.7
Condom	Percent	10.0	9.5	8.8	9.4	9.1	11.7	4.1	10.2
Periodic abstinence-calendar rhythm	Percent	0.5	0.5	(S)	0.6	0.2	1.0	(S)	0.4
Periodic abstinence-natural family planning	Percent	0.1	(S)	(S)	(S)	(S)	0.2	(S)	(S)
Withdrawal	Percent	3.2	3.3	2.1	3.0	1.5	4.5	1.4	5.3
Other methods <sup>4</sup>	Percent	0.3	0.3	(S)	0.5	0.2	0.3	(S)	(S)
Not using contraception	Percent	38.2	35.3	45.5	41.5	60.7	21.4	39.4	28.8
Surgically sterile-female (noncontraceptive)	Percent	0.4	0.2	0.4	0.7	0.4	0.3	1.0	(S)
Nonsurgically sterile-female or male <sup>4</sup>	Percent	1.7	1.6	1.8	1.8	2.1	1.0	2.7	2.2
Pregnant or postpartum	Percent	5.4	4.9	5.7	8.3	2.6	7.2	2.6	10.5
Seeking pregnancy	Percent	4.1	3.5	4.4	6.2	1.3	6.4	0.8	7.1
Other nonuse:									
Never had intercourse or no intercourse in									
3 months before interview	Percent	19.2	18.3	22.6	18.8	45.6	0.9	21.1	1.8
Had intercourse in 3 months before interview	Percent	7.3	6.7	10.6	5.8	8.7	5.5	11.3	6.9

S Figure does not meet publication standards. <sup>1</sup> Includes other races not shown separately. <sup>2</sup> Persons of Hispanic origin may be any race. <sup>3</sup> Includes diaphragm, emergency contraception, female condom or vaginal pouch, foam, cervical cap, Today™ sponge, suppository or insert, jelly or cream, and other methods. <sup>4</sup> Persons sterile from illness, accident, or congenital conditions.

Source: U.S. National Center for Health Statistics, *National Survey of Family Growth*, "Use of Contraception in the United States: 1982–2008," Series 23, No. 29, May 2010. See also <<http://cdc.gov/nchs/nsfg/>>.