

Table 182. Reported AIDS Cases for Adults and Adolescents by Transmission Category and Sex: 2007

[Provisional. For cases reported in the year shown. Includes Puerto Rico, Virgin Islands, Guam, and U.S. Pacific Islands. Acquired immunodeficiency syndrome (AIDS) is a specific group of diseases or conditions which are indicative of severe immunosuppression related to infection with the human immunodeficiency virus (HIV). Data are subject to retrospective changes and may differ from those data in Table 180. For information on HIV death rates, go to Table 126]

Transmission category	2007			Cumulative through 2007 ¹		
	Total	Male	Female	Total	Male	Female
Persons 13 years old and over, total	38,297	28,320	9,977	1,021,242	820,037	201,205
Male-to-male sexual contact	14,383	14,383	(X)	445,645	445,645	(X)
Injection drug use	4,736	3,103	1,633	235,842	166,251	69,591
Male-to-male sexual contact and injection drug use	1,514	1,514	(X)	67,797	67,797	(X)
Hemophilia/coagulation disorder	46	37	9	5,567	5,212	355
High-risk heterosexual contact	7,504	2,791	4,713	142,852	52,623	90,229
Sex with injection drug user	985	281	704	38,766	11,941	26,825
Sex with bisexual male	233	(X)	233	5,415	(X)	5,415
Sex with person with hemophilia	14	4	10	603	90	513
Sex with HIV-infected person, transfusion recipient	56	31	25	1,403	584	819
Sex with HIV-infected person, risk factor not specified	6,216	2,475	3,741	96,665	40,008	56,657
Receipt of blood transfusion, blood components, or tissue	109	50	59	9,315	5,181	4,134
Other/risk not reported or identified	10,005	6,442	3,563	114,224	77,328	36,896

X Not applicable. ¹ Includes persons with a diagnosis of AIDS, reported from the beginning of the epidemic through 2007. Cumulative total includes persons with characteristics unknown. The first AIDS cases were reported in the United States in June 1981.

Source: U.S. Centers for Disease Control and Prevention, Atlanta, GA, *HIV/AIDS Surveillance Report, 2007*, Volume 19, <<http://www.cdc.gov/hiv/surveillance/resources/reports/2007report/>> (revised February 2009).

Table 183. Estimated Numbers of Persons Living with Acquired Immunodeficiency Syndrome (AIDS) by Year, Age, and Selected Characteristics: 2000 to 2007

[Includes Puerto Rico, Virgin Islands, Guam, and U.S. Pacific Islands. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting. See the Commentary section in the report]

Age and characteristic	2000	2004	2005	2006	2007
Total ^{1, 2}	322,242	411,089	432,029	451,837	470,902
AGE AS OF END OF YEAR					
Less than 13 years old	2,916	1,797	1,485	1,193	929
13 and 14 years old	518	802	801	757	680
15 to 24 years old	5,086	7,013	7,700	8,184	8,821
25 to 34 years old	55,678	47,368	45,623	44,829	44,729
35 to 44 years old	145,524	164,185	163,885	161,009	155,702
45 to 54 years old	85,268	136,084	149,317	162,562	175,604
55 to 64 years old	21,701	42,954	50,485	58,464	67,101
65 years old and over	5,550	10,886	12,733	14,839	17,336
RACE/ETHNICITY					
White, not Hispanic	119,172	143,338	149,107	154,770	160,010
Black, not Hispanic	132,104	174,479	183,930	192,793	201,404
Asian/Pacific Islander	2,498	3,834	4,191	4,564	5,011
American Indian/Alaska Native	1,073	1,462	1,563	1,632	1,698
Hispanic ³	64,833	84,427	89,415	94,100	98,726
TRANSMISSION CATEGORY					
MALE ADULT/ADOLESCENT					
Males 13 years old and over, total	250,315	314,618	329,884	344,370	358,332
Male-to-male sexual contact	143,390	186,036	196,868	207,551	218,136
Injection drug use	58,238	64,651	65,716	66,509	67,121
Male-to-male sexual contact and injection drug use	24,968	28,510	29,248	29,771	30,196
High-risk heterosexual contact ⁴	20,839	32,153	34,671	37,065	39,353
Other ⁵	2,880	3,267	3,381	3,474	3,527
FEMALE ADULT/ADOLESCENT					
Females 13 years old and over, total	68,015	92,443	98,101	103,447	108,576
Injection drug use	28,212	32,824	33,684	34,314	34,845
Heterosexual contact ⁴	38,131	57,368	62,037	66,626	71,088
Other ⁵	1,671	2,250	2,380	2,506	2,643
CHILD (LESS THAN 13 YEARS OLD AT DIAGNOSIS)					
Total	3,912	4,028	4,044	4,021	3,992
Perinatal	3,709	3,828	3,841	3,822	3,800
Other ⁶	202	201	203	198	192

¹ Includes persons of unknown or multiple race and of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total. ² Persons who reported multiple racial categories or whose race was unknown are included in the total numbers. ³ Hispanics can be of any race. ⁴ Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. ⁵ Includes hemophilia, blood transfusion, perinatal and risk not reported or not identified. ⁶ Includes hemophilia, blood transfusion, and risk not reported or not identified.

Source: U.S. Centers for Disease Control and Prevention, Atlanta, GA, *HIV/AIDS Surveillance Report, 2008*, Volume 20, <<http://www.cdc.gov/hiv/surveillance/resources/reports/2008report/>>.