## Table 194. Selected Respiratory Diseases Among Persons 18 Years of Age and Over by Selected Characteristics: 2008

[In thousands (225,227 represents 225,227,000). Respondents were asked in two separate questions if they had ever been told by a doctor or other health professional that they had emphysema or asthma. Respondents who had been told they had asthma were asked if they still had asthma. Respondents were asked in three separate questions if they had been told by a doctor or other health professional in the past 12 months that they had hay fever, sinusitis, or bronchitis. Based on the National Health Interview Survey, a sample survey of the civilian noninstitutionalized population; see Appendix [II]

		Selected respiratory conditions <sup>1</sup>						
Selected characteristic	Total	Emphy-	Asthma		Hay		Chronic	
	persons	sema	Ever	Still	fever	Sinusitis	bronchitis	
Total <sup>2</sup>	225,227	3,789	28,260	16,380	18,022	30,621	9,832	
SEX								
Male Female	108,755 116,472	1,769 2,020	11,705 16,555	6,136 10,244	7,463 10,559	10,080 20,541	3,121 6,711	
AGE	, i	,						
18 to 44 years 45 to 64 years 65 to 74 years 75 years and over	110,615 77,335 19,869 17,409	222 1,573 1,124 870	14,888 9,265 2,376 1,731	7,949 5,768 1,548 1,116	7,379 7,823 1,537 1,283	12,850 12,072 3,362 2,337	3,486 4,251 1,172 922	
RACE								
Race alone: <sup>3</sup> White Black or African American American Indian or Alaska Native Asian Native Hawaiian or other Pacific Islander Two or more races <sup>5</sup>	222,430 182,651 26,765 2,178 10,501 334 2,798	3,708 3,443 217 (B) (B) - 4 81	27,538 22,650 3,627 263 966 (B) 722	15,881 13,032 2,163 148 506 (B) 499	17,709 15,130 1,699 4 132 723 (B) 313	30,237 25,593 3,645 227 740 (B) 384	9,649 8,442 1,020 <sup>4</sup> 52 <sup>4</sup> 124 (B) 183	
HISPANIC ORIGIN AND RACE 6 Hispanic or Latino	30,583	191	2,950	1,639	1,554	2,693	631	
Mexican or Mexican American	19,089 194,645 154,483 26,051	4 93 3,598 3,258 214	1,546 25,310 19,951 3,525	825 14,741 11,542 2,107	872 16,468 13,647 1,666	1,526 27,928 23,112 3,582	304 9,201 7,834 1,011	

<sup>–</sup> Represents zero. B Figure too small to meet statistical standards for reliability of a derived figure. ¹ A person may be represented in more than one column. ² Total includes other races not shown separately. ³ Refers to persons who indicated only a single race group. ⁴ Figure does not meet standard of reliability or precision. ⁵ Refers to all persons who indicated more than one race group. ⁵ Persons of Hispanic or Latino origin may be any race or combination of races.

Source: National Center for Health Statistics, Summary Health Statistics for the U.S. Population: National Health Interview Survey, 2008, Vital and Health Statistics, Series 10, Number 242, 2009. See also <a href="http://www.cdc.gov/nchs/data/series/sr-10/sr-10">http://www.cdc.gov/nchs/data/series/sr-10/sr-10</a> 242.pdf>.

## Table 195. Persons 18 Years of Age and Over With Migraines and Pains in the Neck, Lower Back, Face, or Jaw by Selected Characteristics: 2008

[In thousands (225,227 represents 225,227,000). Based on National Health Interview Survey, a sample survey of the civilian noninstitutionalized population, Appendix III]

Selected characteristic		Migraine and pain 1					
		Migraine or		Pain in	Pain in		
	Total	severe	Pain in	lower	face or		
	persons	headache 2	neck 3	back 3	jaw 3		
Total 4	225,227	30,262	31,447	61,719	9,753		
SEX							
Male	108,755	8,886	12,385	27,342	3,051		
Female	116,472	21,376	19,062	34,377	6,701		
AGE							
18 to 44 years	110,615	18,352	14,017	27,136	5,092		
45 to 64 years	77,335	10,260	12,744	22,778	3,523		
65 to 74 years	19,869	1,121	2,618	6,359	649		
75 years and over	17,409	529	2,068	5,447	489		
RACE							
Race alone 5	222,430	29,671	30,791	60,637	9,569		
White	182,651	24,921	26,709	51,792	8,399		
Black or African American	26,765	3,524	2,789	6,272	904		
American Indian or Alaska Native	2,178	342	351	663	<sup>6</sup> 63		
Asian	10,501	844	929	1,822	<sup>6</sup> 196		
Native Hawaiian or other Pacific Islander	334	(B)	(B)	<sup>6</sup> 88	(B)		
Two or more races 7	2,798	591	655	1,082	183		
HISPANIC ORIGIN AND RACE 8							
Hispanic or Latino	30,583	4,050	4,082	7,453	1,033		
Mexican or Mexican American.	19,089	2,501	2,255	4,167	637		
Not Hispanic or Latino	194,645	26,212	27,365	54,266	8,719		
White, single race	154,483 26.051	21,309 3,319	23,025 2,668	45,183 6,032	7,494 853		
Diack of Affical Afficial, Single face	20,031	3,319	۷,000	0,032	633		

B Figure to small to meet statistical standards for reliability of a derived figure. ¹ A person may be represented in more than one column. ² Respondents were asked, "During the past 3 months, did you have a severe headache or migraine?" Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor aches or pains. ³ Respondents were asked, "During the past 3 months, did you have a neck pain; or low back pain; or facial ache or pain in the jaw muscles or the joint in front of the ear?" Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor aches or pains. ¹ Total includes other races not shown separately. ⁵ Refers to persons who indicated only a single race group. ⁶ Figure does not meet standard of reliability or precision. ⁷ Refers to all persons who indicated more than one race group. ⁶ Persons of Hispanic or Latino origin may be any race or combination of races.

Source: National Center for Health Statistics, Summary Health Statistics for the U.S. Population: National Health Interview Survey, 2008, Vital and Health Statistics, Series 10, Number 242, 2009. See also <a href="http://www.cdc.gov/nchs/data/series/sr\_10/sr10\_242.pdf">http://www.cdc.gov/nchs/data/series/sr\_10/sr10\_242.pdf</a>.