Table 194. Persons 18 Years of Age and Over With Migraines and Pains in the Neck. Lower Back. Face. or law by Selected Characteristics: 2007 In thousands (223.181 represents 223.181.000). Based on National Health Interview Survey, a sample survey of the civilian noninstitutionalized populaion, Appendix III1

Migraines and pains 1

a l					
Selected characteristic	Total persons		Pain in neck ³	Pains in lower back ³	Pains in face or jaw 3
Total ⁴	223,181	27,364	29,019	57,070	9,062
SEX Male Female	107,750 115,431		11,665 17,354	25,334 31,736	2,581 6,481
AGE 18 to 44 years old. 45 to 64 years old. 65 to 74 years old. 75 years and over.	76,136 19,258	9,277 1,075	11,833 12,073 2,833 2,280	24,555 21,860 5,650 5,005	4,649 3,455 607 350
RACE Race alone ⁵	220,175	26,792	28,485	56,016	8,825

65 to 74 years old	19,258	1,075	2,833	5,650	607
75 years and over	16,897	585	2,280	5,005	350
RACE					
Race alone ⁵	220,175	26,792	28,485	56,016	8,825
White	180,815	22,241	24,727	47,523	7,532
Black or African American	26,366	3,010	2,473	6,006	814
American Indian or Alaska Native	2,222	463	336	543	⁶ 192
Asian	10,437	1,049	897	1,799	264
Native Hawaiian or other Pacific Islander	335	(B)	(B)	⁶ 146	(B)

16,897	585	2,280	5,005	350
220,175	26,792	28,485	56,016	8,825
180,815	22,241	24,727	47,523	7,532
26,366	3,010	2,473	6,006	814 6192
2,222	463	336	543	⁶ 192
10,437	1,049	897	1,799	264
335	(B)	(B)	⁶ 146	(B)
3,006	572	534	1,053	237
,			,	
	180,815 26,366 2,222 10,437 335	220,175 26,792 180,815 22,241 26,366 3,010 2,222 463 10,437 1,049 335 (B)	220,175 26,792 28,485 180,815 22,241 24,727 26,366 3,010 2,473 2,222 463 336 10,437 1,049 897 335 (B) (B)	220,175 26,792 28,485 56,016 180,815 22,241 24,727 47,523 26,366 3,010 2,473 6,006 2,222 463 336 543 10,437 1,049 897 1,799 335 (B) (B) 6146

White	180,815	22,241	24,727	47,523	7,53
Black or African American	26,366	3,010	2,473	6,006	814
American Indian or Alaska Native	2,222	463	336	543	⁶ 19
Asian	10,437	1,049	897	1,799	26
Native Hawaiian or other Pacific Islander	335	(B)	(B)	⁶ 146	(B
Two or more races 7	3,006	572	534	1,053	23
HISPANIC ORIGIN AND RACE 8					
Hispanic or Latino	29.857	3.462	3.321	6.851	97
Mexican or Mexican American	18,309	1,928	1,925	3,955	510
Not Hispanic or Latino	193,324	23,902	25,698	50,218	8.09

70 16 92

White, single race....... 153,359 19,132 21,729 41,463 6.711 25.574 2.933 2.408 5,666 759 Black or African American, single race B Figure to small to meet statistical standards for reliability of a derived figure.

¹ A person may be represented in more than Respondents were asked. "During the past 3 months, did you have a severe headache or migraine?"

Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor

³ Respondents were asked. "During the past 3 months, did you have a neck pain; or low back pain; or facial ache or pain in the jaw muscles or the joint in front of the ear?" Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor aches or pains. ⁴ Total includes other races not shown separately.

Refers to persons who indicated only a single race group.

8 Refers to persons who indicated more than one race group.

9 Persons of Hispanic or Latino origin may be any race or combination. of races

Source: U.S. National Center for Health Statistics. Vital and Health Statistics, Series 10, Number 240, Summary Health Statistics for U.S. Adults: National Health Interview Survey, 2007; http://www.cdc.gov/nchs/data/series/sr 10/sr10 240.pdf>